RESOURCE ARTICLE

Prox1 dynamically regulates downstream targets and chromatin accessibility

during venous to lymphatic endothelial cell transdifferentiation in the embryo

Lin Grimm[1,2,3,*], Elizabeth Mason[1,2*], Stefanie Dudczig[1,2], Jan Kazenwadel [4,5], Tyrone Chen[1,2], Oliver

Yu[1,2], Neil I. Bower[3], Scott Paterson[1,2,3], Kazuhide Okuda[1,2], Maria Rondon Galeano[1,2], Sakurako

Kobayashi[1,2], Anne Senabouth[3,4], Anne K. Lagendijk[3], Joseph Powell[3,6,7,8], Kelly A. Smith[9], Natasha L.

Harvey[4,5], Katarzyna Koltowska[10] and Benjamin M. Hogan[1,2,3,9 #].

[1] Organogenesis and Cancer Program, Peter MacCallum Cancer Centre, Melbourne, VIC 3000, Australia.

[2] Sir Peter MacCallum Department of Oncology, University of Melbourne, Melbourne, VIC 3000, Australia.

[3] Division of Genomics of Development and Disease, Institute for Molecular Bioscience, The University of

Queensland, St Lucia, Brisbane, Queensland 4072, Australia.

[4] Centre for Cancer Biology, University of South Australia and SA Pathology, Adelaide, South Australia, 5001,

Australia.

[5] SA Pathology, Adelaide, South Australia, 5001, Australia.

[6] Garvan Institute of Medical Research, 384 Victoria Street, Darlinghurst, Sydney, 2010, Australia.

[7] School of Medical Sciences, 18 High Street, University of New South Wales, Kensington, Sydney, Australia.

[8] Garvan-Weizmann Centre for Cellular Genomics, Garvan Institute of Medical Research, 384 Victoria Street,

Darlinghurst, Sydney, 2010, Australia.

[9] Department of Anatomy and Physiology, University of Melbourne, Melbourne, VIC 3000, Australia.

[10] Department of Immunology, Genetics and Pathology, Uppsala University, Uppsala, Sweden.

 # Author for Correspondence:

Professor Ben Hogan

Organogenesis and Cancer Program,

Peter MacCallum Cancer Centre,

Melbourne, VIC 3000, Australia

E-mail: ben.hogan@petermac.org

41 **Keywords:** Lymphatics, Lymphangiogenesis, Prox1, Single cell, RNA Seq, ATAC seq.

42 **Summary**: 149 words

43 **Main Text**: 4378 words

During development, lymphatic vasculature forms as a second, distinct vascular network derived from blood vessels. Transdifferentiation of venous into lymphatic endothelial cells (LECs) is the first step in this process. Specification, transdifferentiation and maintenance of LEC fate requires the transcription factor Prox1 but downstream mechanisms and markers remain to be described in full. We present a single cell transcriptomic atlas of lymphangiogenesis in zebrafish revealing new markers and hallmarks of LEC differentiation. We further profile single cell transcriptomic and chromatin accessibility changes in Prox1 mutants. This work reveals dynamic, multifaceted roles for Prox1 that include blocking blood vascular and hematopoietic fate and up-regulating essential factors for lymphatic development. We reveal that Prox1 controls chromatin state and evidence for Prox1-independent regulation of early LEC differentiation. This work thus provides new insights and a detailed resource for further understanding an enigmatic developmental process that is central in pathological lymphangiogenesis in disease.

Introduction

Lymphatic vasculature plays crucial physiological roles that include the drainage of interstitial fluids, trafficking of immune cells and drainage of dietary lipids. The formation of new lymphatic vessels from pre-existing vessels (lymphangiogenesis) occurs in both development and disease. Signalling through Vegfr3 (Flt4) can be triggered by Vegfc or Vegfd and drives lymphangiogenesis in settings as diverse as development, cancer metastasis, inflammation and ocular disease [1]. In the embryo, lymphangiogenesis begins when the first LEC progenitors depart the cardinal veins (CVs) from E9.5 in mice and 32 hours post fertilisation (hpf) in zebrafish [2]. Vegfc-Flt4 signalling drives LEC progenitor sprouting but also up-regulates Prox1 expression in both zebrafish and mice [3-7]. The transcription factor Prox1 acts as the master regulator of LEC fate [8] and is exclusively expressed in developing LECs in early embryonic vasculature. Loss of Prox1 (Prox1a and Prox1b in zebrafish) leads to a loss of developing lymphatic vessels [4,9]. Following departure from the CV, LEC progenitors go on to colonise embryonic tissues and organs and remodel to form functional lymphatic vessels [10]. While at later stages in mammals there are contributions of LECs from non-venous origins, early embryonic lymphanigogenesis occurs chiefly from the CVs [11-17]. In mice, both the earliest stages of LEC progenitor sprouting from CVs and maintenance of LEC fate in stable lymphatics are dependent on the function of Prox1 [9,18].

Despite over two decades of study of this enigmatic developmental process, the

transcriptomic changes that occur as embryonic venous endothelial cells (VECs)

transdifferentiate into LEC progenitors, and further differentiate into mature lymphatics,

have not been transcriptionally profiled *in vivo*. In the absence of Prox1 in conditional

knockout mice, LECs have been shown to lose the expression of some LEC markers and to gain

expression of some blood vascular endothelial cell (BEC) markers[18]. Prox1 is known to

autoregulate its own expression and to also regulate Flt4 expression in a positive regulatory

loop during early development[7]. Yet how Prox1 controls the transcriptome during LEC

specification, differentiation and maintenance has not been described. This is in part because

of the technical challenge of accessing early mouse endothelial cells in wildtype and mutant

embryos, a problem that is not limiting when using the zebrafish embryo.

As recent studies have demonstrated highly conserved expression and function of Prox1

homologues in zebrafish[4,6,19,20], we here took advantage of the accessibility of the zebrafish

embryo to examine developmental lymphangiogenesis using single cell transcriptomics. We

further analysed zebrafish zygotic *prox1a* mutants with single cell RNA sequencing and single

cell ATAC sequencing, as well as maternal-zygotic double *prox1a/prox1b* mutant (null Prox1

mutants) vasculature with single cell RNA sequencing. Overall, we reveal that embryonic

lymphangiogenesis is a highly conserved process involving dynamic and discrete Prox1-

dependent and independent mechanisms. We provide a resource of new markers of VEC-LEC

transdifferentiation, differentiating and mature cell types and we demonstrate the utility of

this resource to identify a new Prox1-dependent regulator of lymphangiogenesis, the

transmembrane protein Tspan18.

Results

A single cell RNA-seq atlas of embryonic lymphangiogenesis

Zebrafish secondary angiogenesis occurs when Prox1-positive LECs and VECs both sprout

from the cardinal vein (CV) in the trunk or the head in a progressive process between ~32 and

~48hpf. In the trunk, sprouting LECs migrate dorsally and invest the horizontal myoseptum,

where they form a transient pool of parachordal LECs (PLs) from approximately 48hpf[21,22]. In

craniofacial regions of the embryo, LECs sprout from the CVs at several locations[23,24]. After

this, LECs throughout the embryo proliferate and migrate extensively (between ~56hpf-
80hpf) to colonise new regions and tissues[24-27]. In the trunk, LECs anastomose to form the
first lymphatic vessels at around 4 days post fertilisation; forming the thoracic duct (TD),
dorsal longitudinal lymphatic vessels (DLLVs) and intersegmental lymphatic vessels (ISLVs). In
craniofacial regions, they assemble from disparate sources into lateral (LFL), medial (MFL),
otolithic lymphatic vessels (OLV) and lymphatic branchial arches (LBA), as well as forming a
lymphatic loop (LL) in the head that will later give rise to a unique mural LEC population in the
brain (muLECs, also known as FGPs or brain LECs)[28-30]. By 5dpf, major lymphatics in the
craniofacial and trunk regions of the embryo are functional and drain dyes and fluids
deposited in the peripheral tissues[22,31].
119
To profile stages of development spanning key steps in lymphatic differentiation, we selected:
40hpf, when specification and initial sprouting of LECs are occurring; 3dpf, when immature
LECs are migrating through the embryo; 4dpf when LECs are assembling into vessels; and 5dpf
when lymphatics are functional and maturing[32] (**Fig. 1a**). We used transgenic zebrafish strains
that specifically labelled embryonic vasculature to allow for fluorescence activated cell sorting
(FACS, full details in methods). We dissociated whole transgenic embryos and FAC sorted
endothelial cells, for profiling on the 10X Chromium scRNA-seq platform. We sequenced
35,634 cells across 3 runs, then merged and normalised the data[33,34], filtering low-quality
libraries (**Extended data Fig 1a**). To define the cellular identity of each cluster, we
systematically evaluated the expression of known markers summarised in **Extended data**
Table 1a. We identified 9,771 lymphatic and venous endothelial cells that comprise our single
cell atlas of lymphangiogenesis (**Fig 1b**, **Extended data Fig 1b-c**).
132
This dataset is displayed[35] as a UMAP in **Fig 1b** coloured according to developmental stage
and cell phenotype, with venous (VEC) to lymphatic (LEC) trajectory of differentiation
confirmed by RNA-velocity analysis[36]. Comparison of cells at 40hpf with later stages revealed
that the earlier populations are transcriptionally distinct from VECs and LECs at 3, 4 and 5dpf.
Notably, the 40hpf "VEC preLEC" cluster contained both *prox1a+* and *prox1a-* cells forming a
single population rather than discrete clusters. This indicates that LECs are not
transcriptionally distinct from VECs at 40hpf, suggesting early *prox1a+* cells are specified but
not yet differentiated LECs (**Fig 1c**)[4]. We identified 3 main classes of LECs at 3, 4 and 5dpf:

muLECs (1511 cells) marked by expression of *osr2*[28], canonical LECs marked by expression of

cdh6 (2669 cells), and a smaller sub-population of LECs expressing low levels of *prox1a*

(LEC_low_prox1a, 310 cells) (**Fig 1c**, **Extended data Fig 1d-e**).   To define markers of

differentiating canonical LECs at each developmental stage, we applied differential

expression (DE) analysis (**Extended data Table 1b,**). This analysis not only captured known

LEC markers including *prox1a, angpt2a, tbx1* and *cldn11b*, but also uncovered new genes

commonly expressed in canonical LECs across all developmental stages including *grp156,*

hapln3, cdh6 and *tspan18a* (**Fig 1d, Extended data Fig 1f**).   We extended this approach and

evaluated global differences between all canonical LECs and all VECs (n=1240 genes,

Extended data table 1c). GO analysis[37] confirmed the association of biological processes

known to be associated with lymphatics with genes upregulated in LEC (n=752 LEC gene set),

these terms included "lymphangiogenesis", "glycolytic process", "lymph vessel development"

and "ameboidal-type cell migration" (**Fig 1e**, **Extended data Table 1d**).

To provide further confidence in the specificity of the new LEC markers defined by this scRNA-

seq resource, we used DE analysis to defined cluster specific gene expression. We noted that

this analysis suggested higher expression of the well-known marker *lyve1b* in muLECs than in

LECs, which we validated by examining expression levels of *lyve1b* using the established

transgenic line *Tg(lyve1b:DsRed2)* (heat map in **Fig 2a**). We further identified expression of

the known kidney epithelial solute transporter *slc7a7a* as uniquely expressed in muLEC cells

and *fabp11a* as a canonical LEC and VEC marker excluded from muLECs. We generated both

new *slc7a7a*–Citrine and a *fabp11a-Citrine* BAC transgenic strains, which confirmed the

restricted muLEC and vascular expression patterns and further validated our scSeq atlas of

lymphatic differentiation (**Fig 2b-c**).   Overall, this atlas therefore identifies n=752 specific

markers of lymphangiogenesis, the majority of which are new markers, representing new

candidate regulators and spanning four developmental stages.

Prox1 maintains lymphatic endothelial cell identity by repressing blood vascular fate and

promoting lymphatic vascular fate at the level of the transcriptome

Prox1 is both necessary and sufficient to drive VEC to LEC trans-differentiation in mammals

and this is proposed to occur by Prox1 simultaneously initiating LEC fate and repressing the

VEC fate program[8,18]. Prox1 expression is also necessary for the maintenance of LEC identity

during development[18,38,39]. Despite the role for Prox1 being well-studied in mouse models,

the transcriptomic program controlled by Prox1 *in vivo* has never been profiled. Zebrafish

prox1a zygotic and maternal zygotic mutants have been previously described[4,20]. The zygotic

mutants retain maternal deposition of *prox1a* in the oocyte, sufficient to drive normal PL

formation, LEC migration to the horizontal myoseptum (HM) and initial assembly of

lymphatics by 4dpf[4,19], however mutants have a reduction in total LEC numbers throughout

the face and trunk[4]. When the maternal contribution of *prox1a* is removed, lymphatic

development is more severely impaired[4]. We hypothesised that the zygotic mutants which

form lymphatic vessels, likely have abnormal vessel identity in the absence of zygotic Prox1.

Thus, we first addressed the question of how Prox1 regulates lymphatic differentiation by

taking advantage of this unique model. We applied scRNA-seq to endothelial cells FAC sorted

from *Zprox1a^{-/-}* mutants[20] and WT sibling zebrafish at 4dpf, during lymphatic vessel assembly

(n=8075 cells, **Extended data Fig 2a**, **Extended data Table 2a**). Cluster analysis revealed 3

populations of LECs (n=2068) and a single population of VECs (n=1051) comprising both

mutant and wild type cells, and perhaps most striking, a single population of mutant cells

(n=484) transcriptionally similar to VECs, marked by the expression of *aqp1a.1* (**Fig 3a-b**,

Extended data Figs 2b-e). All three LEC clusters (LEC, LEC_S1, LEC_S2) showed graded

expression of lymphatic markers *prox1a*, *cldn11b*, *cdh6* and *angpt2a*, such that "LEC" most

closely resembles canonical LECs from our WT atlas. LEC_S1 and S2 represent alternative LEC

sub-types with different proliferative potential based on S-phase occupancy and cell cycle

marker *mki67* (**Fig 3b**, **Extended data Fig 2e**). The expression of *prox1a* was lowest in the

more proliferative LEC_S1 cells, and highest in the less proliferative LEC_S2 (**Fig 3c**).   RNA-

velocity analysis suggested a trajectory between the mutant cluster and the LEC cluster.

Taken together, these findings suggest that the mutant specific cluster sits on a trajectory

between LEC and VEC fate, representing cells either failing to fully differentiate from VEC to

LEC or LECs undergoing dedifferentiation.

To survey how Prox1 maintains normal LEC differentiation, we performed DE analysis

comparing the mutant cluster with WT LEC (**Fig 3f**, **Extended data Table 2b**). Overall, a much

larger set of genes were downregulated than upregulated in the mutant cluster (n=1034 vs

n=294) with almost half of the most downregulated genes (AvLogFC) highly enriched for LEC

markers (eg. *tbx1*, *cdh6*, *cldn11b*). Of the upregulated genes, almost 75% were VEC markers

(eg. *sox7*, *kdrl*, *cdh5*) and again this was enriched in the most highly upregulated genes (**Fig**

3f, Extended data Fig 2f). Comparing the change in gene expression between wildtype LEC

cluster and VEC cluster, with the change between wildtype LEC and mutant cluster, we found

striking concordance suggesting the mutant cluster is shifted along a LEC fate to VEC fate

trajectory (**Fig 3g**). Overall, there is a simultaneous loss of lymphatic fate and re-acquisition

of blood vascular gene expression, consistent with work in mouse and with Prox1 function

being highly conserved between vertebrates. To validate these observations, we used

confocal imaging of Z*prox1a*$^{-/-}$ mutants and observed upregulation of blood vascular markers

cdh5 (**Fig 3h-i, Extended data Fig 3a,c**) and *kdrl* (**Fig 3j-k, Extended data Fig 3d-g**) in *lyve1b*-

positive lymphatics vessels. We saw a coincident reduction in the expression of *lyve1b* in

these vessels (**Extended data Fig3b**). Interestingly, the difference in *kdrl* levels (relative

intensity) in lymphatics between mutants and wildtype was more significant at 5dpf than

4dpf, suggesting that progressive dedifferentiation may be occurring in the mutant (**Fig 3k**).

This single cell profile thus demonstrates that Prox1 alone is sufficient to maintain LEC fate

and identifies the gene expression maintained by Prox1 in the differentiating lymphatic

vasculature.

Prox1 mutant cells that fail to differentiate, display a unique, immature chromatin

signature

At the level of gene expression, Prox1 function is essential for VEC to LEC differentiation,

however it is unknown if Prox1 controls chromatin accessibility during this process. Thus, we

next profiled Z*prox1a*$^{-/-}$ mutants at 4dpf during vessel assembly using single nuclei (sn) ATAC-

seq (n=3,731 nuclei, **Extended data Fig 4a, Extended data Table 3a**). Cluster analysis (**Fig 4a**)

and overall accessibility of key markers (**Fig 4b**) identified similar populations to the scRNA-

seq profiling: canonical LEC (LEC_01 n=114 nuclei, LEC_02 n=213 nuclei) and VEC (VEC_01

n=157 nuclei) clusters, and a small but discrete population comprised almost entirely of

mutant cells (Mutant Cluster n=47 nuclei). Notably, we found that there was almost no

contribution of mutant cells to the LEC clusters (**Fig 4a** lower bar plot), indicating a loss of fate

when cell identity is determined at the level of chromatin state **(Extended data Fig 4b-e)**. To

validate the utility of this dataset, we assessed concordance between gene expression and

chromatin accessibility in WT LECs and WT VECs (**Fig 4c**). DA (differential accessibility) analysis

revealed the more accessible regions in the LEC clusters were associated with LEC genes

identified in our atlas, and the less accessible regions with VEC genes (**Fig 4c, Extended data**

table 3b). These regions were enriched for motifs of lymphatic-specific transcription factors

including Mafa, Mafb, and the Tbx family **(Extended data table 3c)**. Several markers of LEC

and VEC identity demonstrated overall concordance between chromatin accessibility (gene

score) and gene expression (**Figure 4d**), including DA regions in *cdh6* (**Fig 4d and e**) and *cdh5*

(**Fig 4d and f**). This was consistent with observations in the scRNA-seq data, providing further

evidence that the Prox1a deficient cells assembling into fragmented lymphatic vessels are not

differentiated LECs.

Given that cells in the mutant specific cluster lacking Prox1 demonstrate loss of lymphatic

gene expression and acquisition of blood vascular gene expression, we expected chromatin

accessibility to change in an equally coordinated manner. However, analysis of accessibility

at individual genes (gene score) for the Mutant Cluster revealed changes inconsistent with a

simple fate shift (**Fig 5a**-**b, Extended data Fig 4f**). Notably, at the level of individual peaks we

found that 1726 peaks displayed a striking increase in accessibility compared to WT LEC and

1794 peaks an increase compared with WT VEC (**Fig 5b**, **Extended data tables 4a-b**). Of these,

431 were common peaks identifying more accessible chromatin regions in the mutant cluster

than in either WT LEC or WT VEC (**Fig 5b**). This suggests that some regions of chromatin open

up more than usual during LEC or VEC differentiation in the absence of Prox1. Most genes

with increased accessibility did not show a concurrent increase in gene expression, and we

observed no pattern or trend in expression overall (**Fig 5c**). These more open regions were

enriched for motifs of early acting transcription factors involved in embryonic vasculogenesis

and haematopoiesis, including Erg, Etv2, Ets1, Fli1 and Spi1/Pu.1 (**Fig 5d, Extended data table**

4c). We take this to suggest that in the absence of Prox1, transcription factors that normally

drive earlier transitions during vasculogenesis and haematopoiesis activate a unique

developmental chromatin signature. This demonstrates that Prox1 coordinates the correct

accessibility of the chromatin while maintaining LEC fate in endothelial cells. Furthermore, we

note that a large number of the genes with increased chromatin accessibility are known

regulators of lymphangiogenesis and so this increased accessibility may be a sensitive way to
identify key regulatory genes (**Extended data table 4c**).

Prox1 is primarily a negative regulator of blood and blood vascular fate during the
specification of LEC fate and VEC-LEC transdifferentiation

While the above data identifies the role of Prox1 at stages when it is maintaining LEC identity
in assembling LECs, Prox1 is essential for the very earliest decision made when a VEC becomes
specified to develop into a LEC[9]. To ask how Prox1 controls the earliest stages of venous to
lymphatic transdifferentiation, we first needed to generate a complete Prox1 loss of function
model in zebrafish. We generated *prox1a*$^{+/-}$, *prox1b*$^{+/-}$ double heterozygous animals (lacking
both Prox1 homologues) and then used germline transplantation approaches to produce
animals carrying a double mutant germline (**Extended data Fig 5a-b**). Genetic crossing of
these animals generated embryos that were maternal zygotic (MZ) *MZprox1a*$^{-/-}$, *MZprox1b*$^{-/-}$
mutant zebrafish completely lacking *prox1a/b* transcript expression or maternal deposition
(**Fig 6a-b; Extended data Fig 5c**). A quantitative phenotypic analysis revealed double MZ
mutants show a severe reduction of facial lymphatics and a near complete loss of lymphatic
vessels in the trunk by 4dpf (**Fig 6d-e, Extended data Fig 5f-h**) [4]. Furthermore, *MZprox1a*$^{-/-}$,
MZprox1b$^{-/-}$ mutants initially show a delay in the formation of PLs as these cells emigrate the
CV to invest the HM, but despite this delay, these cells eventually seed the HM before failing
to undertake any further migration (**Fig 6c; Extended data Fig 5i-m**). We saw evidence for
genetic interaction between *prox1a* and *prox1b*, suggesting that both can contribute in
zebrafish lymphatic development (**Extended data Fig 5**).

To test whether LECs require Prox1a in a cell autonomous manner, we performed embryonic
transplantation to generate chimeric embryos, transplanting *MZprox1a*$^{-/-}$ mutant cells into
wildtype hosts (**Extended data Fig 6a**). MZ*prox1a*$^{-/-}$ mutant cells failed to form lymphatic
vessels in otherwise wildtype hosts and we noted mutant PLs stuck in the HM by 5dpf,
confirming that Prox1a functions cell autonomously during zebrafish lymphangiogenesis
(**Extended data Fig 6b-d**). Taken together, the *MZprox1a*$^{-/-}$, *MZprox1b*$^{-/-}$ mutant phenotype is
more severe than any previously described zebrafish Prox1 mutants and Prox1 is necessary
cell autonomously for zebrafish lymphatic development.

Next, we profiled the endothelium of *MZprox1a^-/-^*, *MZprox1b^-/-^* mutant zebrafish at 40hpf

(when cells are both being specified to LEC fate and sprouting from the CVs) using single cell

RNA seq as described above (**Extended data table 5a**). Using cluster analysis and key marker

expression, we identified populations of cells that included VECs and also included *prox1a-*

positive cells. Based on expression of markers associated with EC sprouting (*mki67, pcna*), we

defined two populations of secondary sprouts of venous origin (LEC_VEC_01 n=713 cells,

LEC_VEC_02 n=677 cells) and a single population of cells representing the cardinal vein (PCV

n=812 cells; **Fig 6f-i**). Consistent with our observations in **Fig 1,** cells expressing *prox1a* at

40hpf were not transcriptionally distinct from sprouting VECs and failed to form a "lymphatic

progenitor" cluster, suggesting they are specified and express *prox1a* but not yet

differentiated. RNA velocity analysis suggested that these three clusters remained closely

related, consistent with little differentiation between these populations at this stage. To

identify the earliest transcriptional changes controlled by Prox1 in lymphatic development,

we used DE to evaluate the global differences between *MZprox1a^-/-^*, *MZprox1b^-/-^* and WT in

the LEC_VEC_01 and LEC_VEC_02 clusters of sprouting cells (**Fig 6j, Extended data table 5b**).

We found a significant upregulation of a large set of genes in the absence of Prox1 that were

identified as enriched for blood vascular and haematopoietic genes by GO-term and marker

analysis (**Fig 6k,** n=1,027 genes, **Extended data table 5c**), but we saw little evidence of

positively regulated genes downstream of Prox1 (just n=27 genes downregulated in

*MZprox1a^-/-^*, *MZprox1b^-/-^*). This demonstrates that in the earliest stages of LEC specification

Prox1 is not actively driving a fate program but rather is acting primarily to down-regulate

blood vascular and hematopoietic fate. This profile identifies these earliest Prox1-dependent

changes.

Sprouting Prox1-expressing cells show changes in expression that are not controlled by

Prox1

During their departure from the CV, zebrafish LECs show differences from sprouting VECs in

their proliferation and morphology [40,41]. While Prox1 specifies and maintains LEC fate, these

early changes could potentially be regulated by Prox1-independent mechanisms that have

yet to be described. To test this, we first examined the 27 genes that we identified as

positively regulated in response to Prox1 function (down in mutants; **Fig 6l**). We found that

none of these genes were known regulators of lymphangiogenesis. Next, we compared

wildtype *prox1a*-expressing and non-expressing cells in the 40hpf secondary sprout clusters

using DE analysis (**Fig 6m,o; Extended data table 5d**). We found that n=71 genes were more

highly expressed in the early specified Prox1+ LECs than in sprouting VECs, however there was

essentially no overlap with the genes that were Prox1-dependent (**Fig 6n**). Despite this, the

n=71 genes were clearly enriched for known regulators of early lymphatic vascular

development. This included the Yap target *ccn2a,* as well as *s1pr1, nrp2b, rasa1b, mafbb* and

cldn11a, all up-regulated in LECs at this early stage and previously linked to lymphatic

development [42-47] (**Fig 6o**). Together, we take this as evidence that Prox1-independent

mechanisms exist in early specified LECs and that they control at least several essential

regulators and markers of lymphangiogenesis.

Tspan18 homologues are Prox1 targets essential for embryonic lymphangiogenesis

While we find that Prox1 homologues function primarily to negatively regulate gene

expression at the earliest stages of lymphangiogenesis, there were still 27 genes positively

regulated by Prox1 at 40hpf (**Fig 7a**). One of these genes, *tspan18b* encodes a four-

transmembrane domain Tetraspanin protein. Tetraspanins are a highly evolutionarily

conserved family of proteins involved in the organisation of membrane microdomains and

have been shown to be able to influence cell adhesion, signalling and behaviour [48]. No

Tetraspanins have been specifically linked to the process of developmental

lymphangiogenesis. We noted that both *tspan18b* and *tspan18a* were down-regulated in

MZprox1a$^{-/-}$, MZprox1b$^{-/-}$ mutants, although *tspan18a* was expressed at much lower levels

than *tspan18b* at this stage (**Fig 7a**). In our 4dpf *Zprox1a$^{-/-}$* mutant dataset, both paralogues

were down-regulated in the mutant cluster compared with mature LECs (**Fig 7b**). Both genes

displayed expression in LECs in our time course atlas that was enriched in lymphatic vessels

compared with muLECs and *tspan18a* was highly enriched in LECs compared with VECs at 3,

4 and 5dpf (**Fig 7c**). In our sn-ATAC seq dataset at 4dpf, we identified LEC specific peaks of

chromatin accessibility that were increased in the absence of Prox1 for *tspan18b* but Prox1-

dependent for *tspan18a* (**Fig 7d**). This suggests that Prox1 regulation of Tspan18 paralogues

is likely to be direct in the earliest stages of VEC-LEC transdifferentiation, during specification

of the lymphatic lineage. To test if this may be correct and to probe evolutionary conservation

of this relationship, we performed ChIP seq for PROX1 in human LECs. We found that PROX1

binds directly to intragenic sites and the TSS at the *TSPAN18* locus in human LECs (**Fig 7e**).

Finally, we tested if *tspan18a* and *tspan18b* are functionally relevant targets of Prox1 in
zebrafish lymphatic development. We used a CRISPR F0 knockout approach[49] and confirmed
CRISPR targeting by PCR. We carefully quantified LEC numbers in the trunk of
tspan18a/tspan18b crispants at 5dpf and found a loss of lymphatic vessels and strong
reduction in LEC numbers but no change in the number of VECs in venous ISVs, which form
concomitant with LECs during secondary sprouting (**Fig 7f**). The reduction in LECs in this
setting was not a phenocopy of the more severe phenotype in *MZprox1a*[−/−], *MZprox1b*[−/−]
mutants, suggesting that the *tspan18* paralogues likely perform only a part of the job
necessary for normal lymphatic development downstream of Prox1 (**Fig 7f**). Overall, the
example of *tspan18a* and *tspan18b* demonstrates the utility of this resource of singe cell RNA
and snATAC seq data for the discovery of highly conserved targets of Prox1 and of functionally
relevant regulators of lymphatic development.

Discussion
In this study, we present a single cell RNA seq analysis of four key stages of embryonic
lymphangiogenesis in a vertebrate embryo, revealing new markers and potential regulators
of LEC differentiation. We profile the endothelial cells of *Zprox1a* mutants, which form
lymphatic structures, but these "lymphatics" dedifferentiate. This identifies the
transcriptional code maintained by Prox1 in order to maintain LEC identity and the highly
conserved nature of Prox1 function comparing zebrafish with mice. We further probed the
Zprox1a mutant model using snATAC seq and found that while in mutants the transcriptome
reverts to a blood vascular fate in the absence of Prox1, the chromatin displays ectopically
open regions and peaks that suggest a more immature state. In particular, in the absence of
Prox1 at 4dpf, a large number of opened chromatin regions are enriched for transcription
factor motifs for earlier acting vasculogenic and haematopoietic transcription factors. This
suggests that regulators of earlier vascular fate become more active and increase chromatin
accessibility in the absence of Prox1. This role of Prox1 has not been previously appreciated
and this may suggest its combinatorial function with other developmental transcription
factors that are dysfunctional in the absence of Prox1.

Additional biological insights from this study come from our analysis of double maternal
zygotic *prox1a, prox1b* mutants. These mutants are presumed "null" mutants for Prox1
orthologues and they revealed (through scRNA seq) that during its earliest role in vascular
development, in LEC fate specification and VEC-LEC transdifferentiation, Prox1 functions
primarily to negatively regulate blood and blood vascular fate. While this may represent a
downstream program rather than direct repression of gene expression by Prox1 orthologues,
it is notable that Prox1 and *Drosophila* Prospero have been reported to be able to function as
repressors in a context dependent manner [50-54]. We also found that there is significant change
in early sprouting *prox1a*-positive LECs that is not regulated by Prox1-function. While this may
seem counter-intuitive, many of the genes increased in these cells independent of Prox1
function are already implicated in lymphatic development and we take this to suggest that
mechanisms regulated independently of Prox1 homologues exist in the earliest sprouting
LECs. The regulators of these factors at this stage remain to be described. Finally, by further
characterisation of Prox1 regulation of Tspan18 homologues (a and b), we identify that Prox1
controls lymphangiogenesis through a previously undescribed factor that is likely to act at the
LEC membrane. The Tspan18 CRISPR knockout phenotype is more subtle than *MZprox1a$^{-/-}$*,
MZprox1b$^{-/-}$ mutants, suggesting that multiple downstream targets are likely at play and
orchestrate lymphatic development in response to Prox1 function.
409
Altogether this study describes the process of developmental venous to lymphatic
transdifferentiation, early LEC differentiation and LEC maintenance in detail, *in vivo*. We
describe the role of Prox1 during this process, revealing a conserved and dynamic regulatory
process with unprecedented resolution. This resource will help to understand
lymphangiogenesis in contexts beyond the embryo, such as in pathological
lymphangiogenesis in metastasis, inflammation and tissue repair. Finally, we suggest that
combining zebrafish mutant models, with their inherent ease of cellular accessibility, and the
resolution afforded by single cell profiling, will be an approach that yields powerful new
insights in many areas of developmental biology in the future.
419

Acknowledgements

This project was supported in part by NHMRC Ideas grant 2004300 and ARC Discovery project
DP180102846. B.M.H. was supported by NHMRC Fellowship 1155221. We thank Angelika

Christ and the IMB Sequence Facility (University of Queensland), Tim Semple and Peter Mac

Genomics Facility, the Research Computing Facility at Peter Mac, FACS facilities at UQ and

Petermac and Isaac Virshup for assistance.

Author contributions

L.G. and E.M. performed, analysed experiments and co-wrote manuscript. B.M.H, K.K and

N.L.H conceptualised experiments, analysed data and co-wrote manuscript. S.D, T.C, O.Y,

N.I.B, S.P, K.O, A.S and A.L. performed and analysed experiments. J.P, and K.A.S, provided key

unpublished reagents and analysed data.

Conflicts of interest

The authors declare no conflict of interests.

Online Methods

Zebrafish husbandry

Zebrafish work was conducted in compliance with animal ethics committees at the Peter

MacCallum Cancer Centre, the University of Melbourne and the University of Queensland.

Published transgenic lines used were: *Tg(fli1a:nEGFP)^{y7}* [55]*; Tg(-5.2lyve1b:DsRed)^{nz101}* [24]*;* and

Tg(kdrl:Has.HRAS-mCherry)^{s916} [21]*.* Published mutant lines used were *prox1a^{i278}* [20]*, prox1b* [4,19]*.*

Tg(lyve1b:Venus)^{uq51bh} was generated as previously described[28] but here using an

independent genomic integration with the same construct.

Generation of maternal zygotic mutants

Germline replacement was performed using embryonic transplantation as previously

described [4,56]. Maternal zygotic (MZ) *prox1a* mutant embryos were made by crossing germline

replaced *prox1a$^{i278-/-}$;prox1b$^{sa0035+/-}$* females with *prox1a$^{i278+/-}$;prox1b$^{sa0035+/-}$* males [4,56].

Genotyping of individual embryos during transplantation and phenotypic analysis was

performed as previously described [4].

Transgenesis, genome-editing and genotyping

All microinjections were performed as previously described [57]. CRISPR genome editing for

tspan18a/b was performed as previously described [58], and all genotyping confirmed using

455 PCR [49]. *slc7a7a^BAC^:slc7a7a-Citrine^uom10^* and *fabp11a^BAC^:fabp11a-Citrine^uom10^* recombineering

456 was performed as previously described [59].

457 Guide RNA primers previously described [49]:

458   *tspan18a*-T1-5'-CCCAGATTCCCACCCCTAGA-3' (product size 290bp)

459   *tspan18a*-T2-5'-GGCGGGATGCTGTTCCTACT-3' (product size 270bp)

460   *tspan18b*-T1-5'-TGAAGCCTCCCCGAGCCCCT-3' (product size 285bp)

461   *tspan18b*-T2-5'-TGTGGACCCCACAGGATTTA-3' (product size 110bp)

462 Primers for PCR genotyping:

463   *tspan18a*-T1-geno-forward:5'-AGTGCTGTGGGAAAATGAGC-3'

464   *tspan18a*-T1-geno-reverse:5'-AAATCCCATACAAACAATAGAAGG-3'

465   *tspan18a*-T2-geno-forward:5'-AGTGCTGTGGGAAAATGAGC-3'

466   *tspan18a*-T2-geno-reverse:5'-AAATCCCATACAAACAATAGAAGG-3'

467   *tspan18b*-T1-geno-forward:5'-AGTGCTTTAGTATGCATGTTTCTGT-3'

468   *tspan18b*-T1-geno-reverse:5'-TGCTAGATCAATCGTGGAATGG-3'

469   *tspan18b*-T2-geno-forward:5'-GTGGCTCGTTCCTCCTCG-3'

470   *tspan18b*-T2-geno-reverse:5'-CAGGATGATGTAGACGCCCG-3'

471 Primers for BAC recombineering:

472   *slc7a7a*-BAC-Citrine-forward:5'-

473   AACTGCTTTAGACAGTGTTTTTTGGTACCATCCCATATATTTAAAAAACAGCCACCATGGTGA

474   GCAAGGGCGAGGAG-3'

475   *slc7a7a*-BAC-Citrine-reverse:5'-

476   TTCGACACCTCAGGGGATGCCTCTTCTGCAGGCGTAGGGCTGTAGGACGCTCAGAAGAACT

477   CGTCAAGAAGGCG-3'

478   *fabp11a*-BAC-Citrine-forward:5'-

479   TTACAGCTGTTGCGAGATTGAAAAGTAGAGGAGCATCATTATTCGGGAAAGCCACCATGGT

480   GAGCAAGGGCGAGGAG-3'

481   *fabp11a*-BAC-Citrine-reverse:5'-

482   TCAAAGTTGTCGCTGGTGGTCATTTTCCACGTTCCTACGAATTTGTCAACTCAGAAGAACTC

483   GTCAAGAAGGCG-3'

485 **Imaging and quantification**

486 Imaging was conducted at the Centre for Advanced Histology and Microscopy (Peter

MacCallum Cancer Centre). Imaging of live samples was performed using a Zeiss LSM 710 FCS

confocal microscope, a Zeiss LSM 780 FCS confocal microscope, or an Olympus FV3000

confocal microscope. Mounting and imaging were performed as previously described [60].

In Figs. 2b, 6a-c, and 7f, quantification of vascular phenotypes was performed as previously

described [60,61]. In Figs 3h-k, *cdh5:Venus* or *kdrl:GFP* intensity in the TD was measured using

Imaris software (Bitplane) and normalised to fluorescence intensity of the DA for *cdh5* and to

the fluorescence intensity PCV for *kdrl* in the same embryos.

Fluorescence activated cell sorting

FACS was performed at the Peter MacCallum Cancer Centre and the University of

Queensland. For **Fig 1**, we isolated cells using the following transgenic lines: 40hpf,

Tg(fli1a:nEGFP)^{y7}; 3,4 and 5dpf, *Tg(-5.2lyve1b:Venus)^{uq47bh}, Tg(kdrl:Has.HRAS-mCherry)^{s916}*

For **Fig 3 and Fig 4**, we isolated cells using the following transgenic lines: 4dpf,

Tg(fli1a:nEGFP)^{y7}, Tg(-5.2lyve1b:DsRed)^{nz101}. For **Fig 6**, we isolated cells using the following

transgenic lines: 40hpf, *Tg(fli1a:nEGFP)^{y7}* .To dissociate embryos and obtain single cell

suspensions, we followed published protocols [62]. Briefly, at the desired developmental stage

we deyolked embryos by pipetting up and down and rinsing in calcium free ringers solution,

we centrifuged at 2000rpm for 5' at 4°C and remove supernatant and dissociated the cells by

incubating in liberase [2.5mg/mL] (Cat #5401119001 Sigma-Aldrich) diluted at a 1:35 ratio in

DPBS at 28.5°C for approximately 5', homogenizing the samples during and after the

incubation. To stop the reaction, we added CaCl2 to a final concentration of 1-2mM and FBS

to a final concentration of 5-10%. We centrifuged at 2000rpm for 5' at 4°C. discarded the

supernatant, in order to be able to asses live vs. dead cells, we re-suspend the cells solution

in Zombie Violet TM Viability Dye (Cat# 423113, BioLegend) and incubated for 20' at RT softly

rocking, we rinsed the cells by centrifuging and resuspending in DPBS/EDTA, for ATAC-seq

experiments samples were resuspended in 2%BSA/PBS. Suspension was filtered through a

strainer and taken to the FAC sorting facility. In the Flow Cytometry facility, we used the BD

FACS Aria Fusion sorter (BD Biosciences), we based the selection for the desired population

on FSC and SCC, alive cells were selected based on the Zombie Violet profile and double

positive cells for the desired transgenics were targeted according to the expression profiles

of single cells. Double positive cells were sorted in 300uL 100% FBS in a cold block and taken

immediately to the sequencing facility.

scRNA-seq library preparation

Library preparation and sequencing was performed at the Institute for Molecular Bioscience

Sequencing Facility (University of Queensland) or Peter Mac Genomics Facility. Single cell

suspensions were sorted by FACS, spun down to concentrate and a cell count was performed

to determine post-sort viability and cell concentration. Single cell suspension was partitioned

and barcoded using the 10X Genomics Chromium Controller (10X Genomics) and the Single

Cell 3' Library and Gel Bead Kit (V2 10X Genomics PN-120237; V3.1; 10X Genomics; PN-

1000123). The cells were loaded onto the Chromium Single Cell Chip A (10X Genomics; PN-

120236), B (10X Genomics; PN-1000073 or PN-1000074) or G (10X Genomics; PN-1000120)

to target 10,000 cells. GEM generation and barcoding, cDNA amplification, and library

construction was performed according to the 10X Genomics Chromium User Guide. The

resulting single cell transcriptome libraries contained unique sample indices for each sample.

The libraries were quantified on the Agilent BioAnalyzer 2100 using the High Sensitivity DNA

Kit (Agilent, 5067-4626). Libraries were pooled in equimolar ratios, and the pool was

quantified by qPCR using the KAPA Library Quantification Kit - Illumina/Universal (KAPA

Biosystems, KK4824) in combination with the Life Technologies Viia 7 real time PCR

instrument. After the initial sequencing run, libraries were re-pooled according to estimated

captured cells as determined using the Cell Ranger software (10X Genomics).

Sequencing of scRNA-seq libraries

At the IMB(UQ) genomics facility, denatured libraries were loaded onto an Illumina NextSeq-

500 and sequenced using a 150-cycle High-Output Kit as follows: 26bp (Read1), 8bp (i7 index),

98bp (Read2). Read1 supplies the cell barcode and UMI, i7 the sample index, and Read2 the

3' sequence of the transcript. At the Peter Mac Molecular Genomics facility, single cell

transcriptome libraries were sequenced on an Illumina NovaSeq 6000 using S4 300-cycle

chemistry. Read1 supplies the cell barcode and UMI, i7 the sample index, and Read2 the 3'

sequence of the transcript. Sequencing read lengths were trimmed to 28bp (Read1), 8bp (i7

index), 91bp (Read2), ensuring compatibility with the 10X Genomics analysis software, Cell

Ranger.

scRNA-seq data processing and analysis

All code and documentation associated with this analysis is publicly available under an open source software license at:

https://atlassian.petermac.org.au/bitbucket/users/tyrone.chen/repos/hogan_lab/browse/2022_Grimm_Mason_et_al_PROX1/

Relevant functions are in italics for reference. Where necessary fastq files were made using *Cell Ranger* [63] (version 3.1.0 or 3.0.2) *mkfastq*. Sequencing QC was assessed using *FastQC* 0.11.6 and *MultiQC* viewer for aggregated reports. *Cell Ranger count* and *aggr* were used to generate aggregated count files mapped to GRCz11 (Ensembl 101), without depth normalisation. Doublets were identified from the filtered aggregated count files using *Scrublet* [64] in *Python* version 3.6 and filtered from subsequent analyses. For the $MZprox1^{-/-}$ mutant and $Zprox1^{-/-}$ mutant datasets filtered aggregated count files were processed, sc-transform normalised, filtered and clustered (louvain) using *Seurat* version 2.0 [65] and 3.0 [34] respectively for *R statistical software* version 4.0.2. QC was evaluated before and after normalisation using plot functions in *Seurat* and *scater* 1.20.1 [66], and all thresholds and settings are described in scripting. Cluster solutions were evaluated using *ClusTree* [67]. Datasets used in the atlas of lymphangiogenesis were processed, filtered, merged and log-normalised using *Seurat* version 3.0 [34], with QC and settings as above. Merged data was clustered and normalised using *CSS simspec* [33] and clustering and cluster evaluation performed on this object only, as described above. For all scRNA-seq datasets cluster phenotype was determined using the expression of key markers (Supplementary Table), with the aid of *CellXGene* visualisation software [68]. All downstream analysis and plotting were performed using *Seurat* version 3.0 [34] with default settings. All gene ontology analyses were performed using Panther.db [37] (Biological Process Complete).

Preparation of single nuclei for snATAC-seq

Single cell suspensions were sorted by FACS and prepared for nuclei isolation as previously described by 10x Genomics Demonstrated Protocol for Single Cell ATAC Sequencing (CG000169 - Rev D). Cell suspensions were pelleted (300 x g for 5 minutes) and rinsed with PBS + 0.04% BSA. Cells were resuspended in 95uL of freshly prepared lysis buffer (10mM Tris-HCl pH 7.4, 10 mM NaCl, 3 mM MgCl2, 0.1% Tween-20, 0.1% NP40 Substitute, 0.01%

Digitonin, and 1% BSA) and incubated on ice for 1 minute. 100uL of chilled wash buffer

(10 mM Tris-HCl, pH 7.4, 10 mM NaCl, 3 mM MgCl2, 0.1% Tween-20, 1% BSA) was used to

neutralise the reaction, before the nuclei were pelleted (500 x g for 5 minutes) and

resuspended again in 7uL of 1x Nuclei Buffer (10X Genomics Cat# PN-2000153/2000207).

Presence of healthy and intact nuclei was assessed by visual inspection on a brightfield

microscope using Trypan Blue staining (Thermo Fisher Cat# T10282) and Countess Cell

Counting Chamber Slides (Thermo Fisher Cat# C10228).

snATAC-seq library preparation and sequencing

Single nuclei suspensions were resuspended at approximately 5000 nuclei per μL before

undergoing tagmentation for 60min at 37°C. After tagmentation nuclei were partitioned and

barcoded using the 10X Genomics Chromium Controller (10X Genomics) and the Single Cell

ATAC Reagent Kit (V1.1; 10X Genomics; PN-1000176). Tagmented nuclei were loaded onto

the Chromium Single Cell Chip H (10X Genomics; PN-1000162), GEM generation, barcoding

and library construction was performed according to the 10X Genomics Chromium User

Guide. The resulting single cell ATAC libraries contained unique sample indices for each

sample. The libraries were quantified on the Agilent BioAnalyzer 2100 using the High

Sensitivity DNA Kit (Agilent, 5067-4626) and pooled in equimolar ratios. Sequencing was

performed on an Illumina NextSeq 500 using a 150-cycle High-Output Kit as follows: 50bp

(Read1), 8bp (i7 index), 16bp (i5 index), 50bp (Read2) achieving a read depth of 25,000 read

pairs per nucleus.

snATAC-seq processing and analysis

FASTQ files generated from sequencing were used as inputs to 10X Genomics *Cell Ranger*

ATAC 2.0.0. *cellranger-atac count* was used to generate count files mapped to GRCz11

(ENSEMBL 101), without depth normalisation. Resulting fragment files were read into *ArchR*

1.0.1 for *R statistical software 4.0.5* as a tile matrix with 500-bp bins. All remaining steps in

the ATAC-Seq analysis were performed within *ArchR 1.0.1*. QC filtering was performed, and

only high-quality cells with a TSS enrichment score greater than 4 and greater than 1,000

unique nuclear fragments were retained. Doublets were predicted using *addDoubletScores*

and filtered using *filterDoublets.* Data normalization and dimensionality reduction were

performed using iterative Latent Semantic Indexing (LSI) and Uniform Manifold

Approximation and Projection (UMAP) embeddings were used for visualisation in reduced dimension space. Separate from the *ArchR 1.0.1* package, cluster solutions were independently evaluated using *clustree 0.4.3*. A Gene Score Matrix that stores predicted gene expression was then generated based on the accessibility of regulatory elements in the vicinity of the gene. We used gene scores of endothelial markers for cluster annotation and subsetting. Differentially accessible genes were identified by differential testing using *getMarkerFeatures*. Local chromatin accessibility of the marker genes was visualised using the *plotBrowserTrack*.

ChIP-seq library preparation and analysis

For ChIP, antibodies used were goat anti-Prox1 (R&D Systems, AF2727), rabbit anti-H3K4Me1 (Cell Signaling, #5326), rabbit anti-H3K27Me3 (Cell Signaling, #9733), rabbit anti-H3K27Ac (Cell Signaling, #8173), rabbit anti-H3K4Me3 (Cell Signaling, #9751). hLEC were harvested and processed for ChIP as previously described[69] using a truCHIP™ Low Cell Chromatin Shearing Kit with SDS Shearing Buffer (Covaris). Briefly, 10 million cells/ml were cross-linked using 1% formaldehyde for 5 minutes, neutralised with glycine, lysed and nuclei washed. For transcription factor ChIP, chromatin from 3 million cells was sheared using a 130µl microtube in a Covaris sonicator at the recommended settings for 8 minutes. 5µg antibody was used to immunoprecipitate sheared DNA. For histone mark ChIP, chromatin from 1 million cells was sheared and 1µg antibody was used. Following washing and reversal of crosslinks, DNA was purified using a Qiagen MinElute PCR Purification Kit. ChIP-Seq library preparation and sequencing was carried out at the ACRF Cancer Genomics Facility, Centre for Cancer Biology, Adelaide. ChIP-Seq libraries were prepared according to Illumina's TruSeq Sample Prep Guide Revision A with some minor modifications as previously described[69]. Briefly, DNA was end-repaired, followed by adenylation of the 3' ends. Next, Illumina indexing adapters were ligated to the DNA. 18 cycles of PCR were performed to enrich for successfully adapter ligated molecules. Ligation products were purified with a 2% Pippin Prep gel (Sage Science), selecting a size range of 250 – 300 base pairs. The size distribution and yield of the purified libraries were determined using an Agilent Bioanalyzer High Sensitivity chip and Qubit dsDNA HS assay (Invitrogen), respectively. Libraries were pooled in equimolar ratios and sequenced in one lane of a HiSeq 2500 short read flowcell (1 x 50bp). Reads were mapped to the UCSC hg19 genome with BWA[70] version 0.7.9a-r786, allowing at most 3 alignments. Read depth was

646   calculated via *genomeCoverageBed* from BEDTools[71], using the scale parameter to normalise

647   to reads-per-million. Read depth was plotted relative to annotated transcription start sites

648   via custom Python scripts utilising Matplotlib.

Figure 1: A single cell RNAseq developmental atlas of lymphangiogenesis in zebrafish

a. Schematic representation of four key stages of zebrafish lymphatic development in
head and trunk: 40hpf encompasses both specification (Prox1-induction) and
sprouting; 3dpf migration of LECs; 4dpf assembly of lymphatic vessels; 5dpf
maturation functional lymphatics.

b. UMAP visualization of 9771 cells filtered for VEC and LEC populations (n=6 samples;
see **Extended data table 1a** for cluster identification and **Extended data fig 1** for whole
dataset) coloured according to developmental stage (top), predicted cell phenotype
(middle), and RNA-velocity (bottom).

c. UMAP visualization of key marker gene expression. Colour scale represents log-
normalised expression. LEC markers: *prox1a, cdh6*. BEC marker: *cdh5*. LEC and VEC
markers: *mrc1a, flt4*. AEC marker: *flt1*. muLEC markers: *osr2, hexa*.

d. Dot plot of top genes commonly marking LECs at 3, 4 and 5dpf, with expression
displayed in VECs (all stages), VEC_preLEC (40hpf) and LECs at 3, 4 and 5dpf (see also
Extended data table 1b, **Extended data figure 1f**). Colour scale represents average
log-normalised expression and point size represents percentage of cells expressing
gene.

e. Bar plot summarizing GO BP analyses of genes DE between 3, 4 and 5dpf LEC and VEC
populations (**Extended data table 1c-d**). Y-axis represents enriched BP term, x-axis
represents the -log10(*adjusted p value*), bars are coloured and ordered according to
fold enrichment of the GO term in LEC. GO terms enriched in LEC are coloured purple,
terms enriched in VEC are coloured red.

Hpf, Hours post-fertilisation.dpf, Days post fertilisation. BEC, Blood vascular endothelial cell.
VEC, Venous endothelial cell. LEC, Lymphatic endothelial cell. muLEC, mural Lymphatic
endothelial cell (a.k.a FGP or brain LEC).

Figure 2: Transgenic marker strains confirm cluster identity and identify vessel specific markers

    a. UMAP of *lyve1b* expression predicts higher expression in muLECs than LEC populations (left), confirmed by a heat map of a *Tg(lyve1b:DsRed2)* zebrafish larvae at 6dpf showing high *lyve1b* expression in craniofacial (middle) and moderate expression in trunk lymphatic vessels (right).

    b. UMAP of *slc7a7a* expression predicts restricted expression in muLEC populations (left), confirmed in lateral confocal projections showing co-expression of *lyve1b* (magenta) and a new BAC transgenic strain for *slc7a7* (green) in the zebrafish head at 5dpf (middle) which is not expressed in veins and lymphatic vessels of the trunk (right).

    c. UMAP of *fabp11a* expression predicts expression in LEC and VEC but not muLEC populations (left), confirmed in lateral confocal projections showing co-expression of *lyve1b* and a new BAC transgenic strain for *fabp11a* marking venous and lymphatic vessels in the trunk (right) without showing expression in the vasculature of the head (middle) at 5dpf.

Lateral confocal images, anterior to the left. muLECs, mural lymphatic endothelial cells; facial LV, facial lymphatic vessels; vISV, venous intersegmental vessel; PCV, posterior cardinal vein; TD, thoracic duct; ISLV, intersomitic lymphatic vessel. Scale bars, 80 μm for head (middle) in (a) and 50 μm for trunk images (right).

Figure 3: Single cell RNA seq analysis reveals a fate shift from LEC to VEC in the absence of

Prox1 in zygotic *prox1a* mutants

    a. UMAP visualization of 8075 cells filtered for VEC and LEC subpopulations (n=2

        samples; see **Extended data table 2a** for cluster annotation, **Extended data Fig3** for

        full dataset) coloured according to predicted cell phenotype.

    b. UMAP visualization of marker gene expression. Colour scale represents SCT-

        normalised expression. LEC markers: *prox1a, cldn11b, cdh6, angpt2a*. LEC and VEC

        markers: *flt4, mrc1a, lyve1b*. BEC markers: *cdh5, kdrl*. Mutant cluster markers: *frem1b,*

        aqp1a.1.

    c. UMAP visualization with cells coloured according to genotype (*Zprox1a^-/-* or WT)

        identifies a mutant specific cluster and RNA velocity analysis suggests a trajectory

        between the mutant cluster and the LEC cluster, likely indicative of a fate shift.

    d. Stacked bar plot representing the genotype composition of cells in a given phenotypic

        cluster.

e. Dot plot of marker expression across defined clusters (indicated on Y-axis). Colour
scale represents average SCT-normalised expression and point size represents
percentage of cells expressing gene.

f. Bar plot of average log fold change in gene expression comparing WT LECs with the
Mutant Cluster. Y-axis represents average log fold change, x-axis represents
differentially expressed genes (Wilcoxin Rank Sum *adjusted p value* < 0.05) and bars
are coloured according to status as a LEC (purple) or VEC (red) marker in **Fig. 1**. Pie
charts (right) indicate the LEC and VEC marker composition of genes up-regulated
(n=365) and down regulated (n=1922) in the Mutant Cluster, demonstrating a fate
shift.

g. Concordance in the fate shift between Mutant Cluster and WT LEC with the WT LEC
and VEC trajectory. Each point represents significant DE genes (n=2287) between
Mutant Cluster and LEC WT, coloured according to LEC or VEC marker status. X-axis
represents average log fold change relative to the Mutant Cluster, Y-axis represents
average log fold change relative to LEC WT.

h. Lateral confocal images of *cdh5* (green) and *lyve1b* (magenta) expression in the
developing trunk in WT and *Zprox1a* mutants at 4dpf.

i. Quantification of *cdh5* intensity in the thoracic duct in WT and mutants (relative to
expression in the DA).

j. Lateral confocal images of *kdrl* (green) and *lyve1b* (magenta) expression at 4dpf (left)
and 5dpf (right).

k. Quantification of *kdrl* intensity in the thoracic duct in WT and mutants (relative to
expression in the PCV) at 4 and 5dpf.

WT, wildtype. Z, zygotic. TD, thoracic duct, DA, Dorsal Aorta. PCV posterior cardinal vein. ISLV,
intersomitic lymphatic vessel. DLLV, Dorsal longitudinal lymphatic vessel. PL, parachordal LEC.
Scale bars, 50 μm. *, The error bars represent mean ± s.e.m.; *p=0.01 and **p=0.0094 from
an unpaired, two-sided t-test.

Figure 4: Single nuclei ATAC seq reveals chromatin accessibility changes between venous

and lymphatic endothelial cells

    a. UMAP visualization of snATAC-seq from *Zprox1a⁻ᐟ⁻* mutant and WT endothelial cells at

        4dpf (n=3731 nuclei), coloured by cell phenotype (top) and genotype (middle) (see

Extended data Fig4 for full dataset). Bar plot summarising the composition of wildtype
and mutants in each cluster. A mutant specific cluster is identified, similar to **Fig 3**.

b. UMAPs visualization indicating accessibility of key marker genes (gene score) that
confirm predicted cluster phenotypes.

c. Bar plot of log2 fold change for the differentially accessible peaks (n=4185 peaks,
Wilcoxon Rank Sum, *FDR* < 0.05) between WT LECs and WT VECs. LEC/VEC markers
identified in scRNA-seq are coloured in red/purple respectively and demonstrate a
strong correlation between chromatin state and the VEC to LEC fate shift. 228/1561
more accessible peaks in the LEC cluster were associated with LEC markers and
398/2624 more accessible peaks in VEC cluster associated with VEC markers. Pie
charts (right) summarising the proportion of LEC/VEC markers in the differentially
accessible peaks.

d. Dot plots of scRNA-seq (top) and snATAC-seq accessibility (bottom) data summarising
the behaviour of key LEC (*prox1a, cdh6, tbx1*) and VEC (*cdh5, flt1, gata6*) markers in
WT LECs, WT VECs and the Mutant Cluster at 4dpf. LEC genes are upregulated (scRNA-
seq) and chromatin is more permissive (snATAC-seq) in WT LEC, and VEC genes are
upregulated and more permissive in WT VEC. The size of the dots represents the
proportion of cells or nuclei, and colour represents either SCT-normalised expression
or Gene Score of Accessibility.

e. Genome accessibility track of LEC marker *cdh6* (left) and VEC marker *cdh5* (right)
indicating these genes are more accessible in WT LEC and WT VEC cell phenotypes
respectively. Red bars represent peaks in the reproducible peak set from snATAC-seq.
DAP between WT LECs and WT VECs (Wilcoxon Rank Sum, *FDR* < 0.05) were
highlighted in blue.

DAP, differentially accessible peaks. FDR, false discovery rate. snATAC-seq, single nuclei
ATAC-seq.

Figure 5: Zygotic *prox1a* mutants display a unique chromatin accessibility state in fate-

shifted endothelial cells

a. Heatmaps of accessibility (gene score) for all genes (n=32 020 genes) in WT LECs, WT

VECs and Mutant Cluster showing that mutant cluster cells display a unique chromatin

state at many genes.

b. Schematic illustrates DAP analyses between Mutant Cluster vs. LEC WT and Mutant
Cluster vs. VEC WT. Venn diagram indicates all individual peaks that were found to be
increased in accessibility in the Mutant Cluster vs LEC or VEC with n=431 DAP
commonly more open than in both LEC WT and VEC WT (FDR < 0.05, log2 fold change
> 1.5).
c. Dot plot summarizing the scRNA-seq expression level of n=13 genes with DAP more
open in the Mutant Cluster at 4dpf, demonstrating that accessibility changes for these
genes did not correlate with changes in transcription. The size of the dot represents
the proportion of cells that express the markers in the cluster, and colour represents
SCT-normalised expression.
d. Top 10 enriched motifs (HOMER analysis, adjusted p value < 0.05) in the n=431 peaks
that are more open in the Mutant Cluster than WT LEC and VEC. Percentages on the
right-hand side represent the proportion of regions enriched for the motif in the
n=431 peak set, and the whole genome (background).
e. Genome accessibility track of *prox1a*. Red bars represent peaks in the reproducible
peak set from snATAC-seq.
f. Genome accessibility track of *tbx1*. Red bars represent peaks in the reproducible peak
set from snATAC-seq
g. Genome accessibility track of *kdr*. Red bars represent peaks in the reproducible peak
set from snATAC-seq
804
805 DAP, differentially accessible peaks. FDR, false discovery rate. snATAC-seq, single nuclei
806 ATAC-seq.
807

Figure 6: The role of Prox1 in LEC specification and sprouting revealed by single cell profiling

of maternal zygotic *prox1a;prox1b* double mutants

a. Lateral confocal images of zebrafish heads at 4dpf showing endothelial cell nuclei
(green) and venous and lymphatic vessels (white) in control (upper) and *MZprox1a$^{-/-}$*
;MZprox1b$^{-/-}$ embryos (lower). Facial lymphatics are absent or shorter or missing
(asterisk) in the mutants (asterisk). Scale bar, 80 μm.

b. Lateral confocal images of zebrafish trunks at 4dpf in control (upper) and *MZprox1a$^{-/-}$*
;MZprox1b$^{-/-}$ mutants (lower), showing absent lymphatic vessels (asterisk) but
retained PLs (arrows in lower). Scale bars, 50 μm.

c. Lateral confocal images of PLs in the horizontal myoseptum at 2, 3 and 4dpf. PLs form
later in *MZprox1a$^{-/-}$;MZprox1b$^{-/-}$* mutants and accumulate in the horizontal
myoseptum while the PLs of control embryos emigrate the HM by 4dpf. Scale bars,
50 μm.

d. Quantification of lymphatic endothelial cell (LEC) number from 4dpf heads control:
n=19, *MZprox1a$^{-/-}$;MZprox1b$^{-/-}$* mutants: n=18) and trunks (control: n=29, *MZprox1a$^{-/-}$*
;MZprox1b$^{-/-}$ mutants:n=25). Error bars represent mean ± s.e.m.; **** p < 0.0001,
from an unpaired, two-sided t-test.

e. Quantification of number of PLs per somite in trunks at 2dpf in controls (n=41) and
MZprox1a$^{-/-}$;MZprox1b$^{-/-}$ mutants (n=38), at 3dpf in controls (n=11) and *MZprox1a$^{-/-}$*
;MZprox1b$^{-/-}$ mutants (n=10) and at 4dpf in controls (n=29) and *MZprox1a$^{-/-}$*
;MZprox1b$^{-/-}$ mutants (n=25). Error bars represent mean ± s.e.m.; **** p < 0.0001, *
p < 0.05, from an unpaired, two-sided t-test.

f. UMAP visualization of filtered scRNA-seq data showing 2202 endothelial cells (n=4
samples; **Extended data table 6a**) coloured according to predicted cell phenotype
(top), genotype (middle), and RNA-velocity (bottom).

g. UMAP visualization of key marker gene expression. Colour scale represents SCT-
normalised expression.

h. Stacked bar plot representing the genotype composition of cells in each cluster.

i. Dot plot of key marker expression. Colour scale represents average SCT-normalised
expression and point size represents percentage of cells expressing gene.

j. Bar plot indicating average log fold change of n=1054 significantly different genes
between the *MZprox1a/b$^{-/-}$* and WT cells in LEC_VEC_01 and LEC_VEC_02 clusters at
40hpf (Wilcoxin Rank Sum *adjusted p value* < 0.05).

k. Bar plot summarizing GO BP analyses of n=1027 genes upregulated in *MZprox1a/b^-/-*.

Y-axis represents enriched BP term, x-axis represents the -log10(*adjusted p value*),

bars are coloured according to fold enrichment.

l. Dot plot of n=27 genes downregulated in 40hpf *MZprox1a/b^-/-* and WT cells in

LEC_VEC_01 and LEC_VEC_02 clusters.

m. Schematic outlining the selection of n=109 *prox1a* expressing cells from n=915 *prox1a*

negative counterparts in the 40hpf WT sample (whole dataset in **Fig1**). DEG analysis

revealed n=71 genes upregulated in *prox1a* positive cells (Wilcoxin Rank Sum *adjusted*

p value < 0.05).

n. Venn diagram illustrating genes upregulated in *prox1a* positive (vs negative) WT cells

at 40hpf, and downregulated in *MZprox1a/b^-/-* mutant cells at 40hpf.

o. Dot plot of n=71 genes upregulated in WT *prox1a* positive (vs negative) cells at 40hpf.

Colour scale represents average SCT-normalised expression and point size represents

percentage of cells expressing gene. Bold labels indicate known lymphatic genes.

LL, lymphatic loop. OLV, otholitic lymphatic vessel. LFL, lateral facial lymphatic vessel. MFL,

medial facial lymphatic vessel. LBA, lymphatic branchial arches. DLLV, dorsal longitudinal

lymphatic vessel. PL, parachordal LEC. ISLV, intersomitic lymphatic vessel. TD, thoracic duct.

DA, dorsal aorta. PCV, posterior cardinal vein.

Figure 7: Identification of Tspan18 paralogues as functionally essential downstream targets of Prox1 in lymphangiogenesis

   a. Dot plot of expression of *tspan18b* and *tspan18a* in WT and *MZprox1a/b$^{-/-}$* mutant cells at 40hpf. The size of the dot represents the proportion of cells that express the markers in the cluster, and colour represents SCT-normalised expression.

   b. Dot plot of expression of *tspan18b* and *tspan18a* in Wt and *Zprox1a$^{-/-}$* mutant cells at 4dpf. The size of the dot represents the proportion of cells that express the markers in the cluster, and colour represents SCT-normalised expression.

   c. UMAP visualization of *tspan18b* and *tspan18a* expression in the single cell atlas of lymphangiogenesis. Colour scale represents log-normalised expression.

   d. Genome accessibility track of *tspan18b* (left) and *tspan18a* (right). Red bars represent peaks in the reproducible peak set from snATAC-seq. Differentially accessible peaks between Mutant Cluster and WT LECs (*p value* < 0.05) were highlighted in blue

   e. ChIP-seq profile of Prox1 binding together with H3K27Me3, H3K4Me1, H3K27AC, and H3K4Me3 profiles at the TSPAN18 locus (hg19) in human LECs.

   f. Lateral confocal images of zebrafish trunks depicting mature lymphatic vessels in the uninjected control which are missing in tspan18a;b CRISPR injected embryos at 5dpf (asterisks) identifying a functional requirement for tspan18 paralogues in lymphangiogenesis. Note that the formation of intersegmental vessels is also severely impaired in tspan18a;b crispants (arrowheads).. Scale bars, 50 µm. Quantification of LEC numbers across 4 somites and VEC numbers per vISV at 5dpf. Error bars represent mean ± s.e.m.; ** p = 0.0012, ns = not significant from an unpaired, two-sided t-test.

DLLV, dorsal longitudinal lymphatic vessel. ISLV, intersomitic lymphatic vessel. TD, thoracic duct. PCV, posterior cardinal vein. vISV, venous intersegmental vessel.

   EXTENDED DATA FIGURES AND FIGURE LEGENDS

Extended data Fig 1: Full dataset and analysis for data in Fig 1.

a. UMAP visualization of n=35634 endothelial cells sequenced for the single cell atlas of

lymphangiogenesis (n=6 samples; see also **Extended data table 1a** for markers used

to identify clusters) coloured according to developmental stage (top left), cell cycle

phase (top right), predicted cell phenotype (bottom left), and cluster solution (RNA-

snn-res.1).

b. Clustree analysis demonstrating the relationship between different cluster resolutions

for the n=9771 venous and lymphatic endothelial cells in Figure 1a. Resolution 0.4

(boxed) was used for all downstream analyses.

c. UMAP visualization of cluster resolution 0.4 (as appears in **Fig 1**).

d. Stacked bar plot of resolution 0.4 cluster composition cell cycle phase.

e. Stacked bar plot of resolution 0.4 cluster composition developmental stage.

f.   Venn diagram of significant genes commonly upregulated in LEC compared to VEC at

903   3, 4 and 5dpf.

Extended data Fig 2: Full dataset and analysis for data in Fig 3.

a. UMAP visualization of n=8075 endothelial cells sequenced comparing lymphangiogenesis in WT and *Zprox1a* mutants at 4dpf (n=2 samples; see also **Extended data table 2a** for markers used to identify clusters) coloured according to developmental stage (top left), cell cycle phase (top right), predicted cell phenotype (bottom left), and cluster solution (SCT-snn-res.8)

b. Clustree analysis demonstrating the relationship between different cluster resolutions
for the n=3063 venous and lymphatic endothelial cells in **Figure 3a**. Resolution 0.2
(boxed) was used for all downstream analyses.

c. UMAP visualization of cluster resolution 0.2

d. Stacked bar plot of resolution 0.2 cluster composition genotype

e. Stacked bar plot of resolution 0.2 cluster composition cell cycle phase

f. Venn diagram indicates the overlap between genes significantly upregulated in VEC
compared to LEC at each developmental stage of the single cell atlas, and those
upregulated in the *Zprox1a$^{-/-}$* Mutant Cluster at 4dpf compared to WT LECs. N=85
genes (coloured red) are uniquely associated with the Mutant Cluster and WT VEC at
4dpf, compared to N=4 genes in WT VEC at 3 and 5dpf.

Extended data Fig 3: Additional phenotypic analysis of blood vascular markers in *Zprox1a⁻/⁻* mutants associated with Fig 3

    a. Lateral confocal images of zebrafish trunks at 4dpf showing *cdh5* expression (green) and *lyve1b* expression in lymphatic vessels (magenta) in control (upper) and *Zprox1a* mutant embryos (lower). Transgenic lines are indicated. Scale bar, 50 μm.

    b. Quantification of *lyve1b* expression intensity in the TD relative to the PCV in sibling and *Zprox1a* mutant embryos at 4dpf.

c. Quantification of *cdh5* expression intensity in the TD relative to the DA in sibling and
Zprox1a mutant embryos at 4dpf.

d. Lateral confocal images of zebrafish trunks at 4dpf showing *kdrl* expression (green)
and *lyve1b* expression in lymphatic vessels (magenta) in control (upper) and *Zprox1a*
mutants (lower). Transgenic lines are indicated. Scale bars, 50 μm.

e. Quantification of *kdrl* expression intensity in the TD relative to the DA in sibling and
Zprox1a mutant embryos at 4dpf and 5dpf (upper).

f. Lateral confocal images of zebrafish trunks at 5dpf showing *kdrl* expression (green)
and *lyve1b* expression in lymphatic vessels (magenta) in control (upper) and *Zprox1a*
mutants (lower). Transgenic lines are indicated. Scale bars, 50 μm.

g. Quantification of the number of TD segments displaying *kdrl* expression in *Zprox1a*
mutant embryos at 4dpf and 5dpf (lower).

DLLV, dorsal longitudinal lymphatic vessel. ISLV, intersomitic lymphatic vessel. TD, thoracic
duct. DA, Dorsal aorta. PCV, posterior cardinal vein. vISV, venous intersegmental vessel. PL,
Parachoral lymphatic endothelial cells.

Extended data Fig 4: Additional data and analysis for data in Fig 4.

a. UMAP visualization of n=3731 single nuclei ATAC-seq data comparing lymphangiogenesis in WT and *Zprox1a* mutants at 4dpf. Nuclei coloured according to cluster (res-0.8).

b. UMAP visualisation with nuclei coloured according to predicted phenotype based on
accessibility changes at the genes defined in **Extended data table 1c.**

c. UMAP visualisation with nuclei coloured according to genotype.

d. Stacked bar plot indicating the genotype composition of nuclei captured per cell type.

e. Density plot indicating the majority of snATAC-seq peaks fall around the TSS. Y-axis
represents distance from TSS (bp) and y-axis represents normalised Tn5 insertion
profile.

f. Heatmaps of accessibility (gene score) for DEG (**Fig 3f**) upregulated (left, n=300
mapped genes) and downregulated (right, n=1052 mapped genes) in the Mutant
Cluster compared to WT LECs. Colour indicates row-wise Z-score, each row represents
a differentially expressed gene in scRNA-seq data, columns are snATAC-seq clusters.
Boxed genes downregulated in Mutant display more permissive chromatin in Mutant
than WT LEC, showing that mutant cluster cells display a unique chromatin state at
many genes.

Extended data Fig 5: **Additional phenotypic analysis of *MZprox1a^{-/-};MZprox1b^{-/-}* mutants**

associated with Fig 6

a. Schematic explanation of the germline replacement transplantation method used to

generate *MZprox1a^{-/-};MZprox1b^{-/-}* mutants.

b. Overall morphology of control and *MZprox1a^{-/-};MZprox1b^{-/-}* mutants at 5dpf. Arrows

indicate oedema around eyes, heart and intestine. Scale bars 500 μm.

c-d. Lateral confocal images showing vascular defects in control and mutant in the craniofacial

(**c**) and trunk (**d**) regions of the larvae at 4dpf. Genotypes and transgenic labels are indicated.

MZprox1a-/- and *MZprox1a+/-, MZprox1b-/-* animals both showed a strong loss of lymphatic

vessels in craniofacial and trunk regions. The *MZprox1a-/-, MZprox1a-/-* mutants showed a

robust loss of craniofacial vessels equivalent to the other mutant phenotypes but a more

severe loss of trunk lymphatics and an accumulation of PLs in the HM.

e. Genotypes that are shown in the quantification. Colour codes are used to identify

individual embryonic genotypes in the dot plots showing quantification in **f-h**.

f. Quantification of the number of craniofacial (head) LECs within facial lymphatic

vessels showing evidence for contributions from *prox1a* and *prox1b*.

g. Quantification of the number of mural LECs (muLECs) showing a major role for *prox1a*

and a role for *prox1b* in muLEC development.

h. Quantification of the number of trunk LECs within vessels showing a dominant role for

prox1a in trunk lymphangiogenesis.

i. Lateral confocal images showing defects in the formation of PLs in mutants upon loss

of *prox1a* compared with controls in the trunk. Genotypes are indicated.

j. Colour coded list of analysed genotypes abbreviated in (**k-m**). Each embryo has a

defined genotype for *prox1b* which is represented in colour code as displayed in j.

k. The number of PLs formed (per somite) at 2dpf in genotypes indicated. Decreasing

gene dosage for *prox1a* and *prox1b* progressively reduced the initial seeding of the

HM by PLs.

l. The number of cells in sprouts departing the PCV is unchanged in mutants showing

that sprouting occurs normally.

m. The number of cells in vISVs is unchanged in mutants indicating no effect on the

venous endothelium.

Z, zygotic; MZ, maternal and zygotic; HM, horizontal myoseptum; LL, lymphatic loop; OLV,

otholitic lymphatic vessel; LFL, lateral facial lymphatic vessel; MFL, medial facial lymphatic

vessel; LBA, lymphatic branchial arches; DLLV, dorsal longitudinal lymphatic vessel; PL,

parachordal LEC; ISLV, intersomitic lymphatic vessel; TD, thoracic duct; DA, dorsal aorta; PCV,

posterior cardinal vein.

1005

1006

Extended data Fig 6: **Embryonic transplantation data demonstrating endothelial cell autonomy of *prox1a* maternal zygotic mutant phenotype associated with Fig 6.**

    a. Schematic showing the endothelial cell transplantation approach used to test cell autonomy for the vascular phenotypes seen in *MZprox1a* mutants. Cell identities were traced following transplantation in unlabelled host embryos using expression of the pan -endothelial marker *fli1a* and venous/lymphatic marker *lyve1b* (indicated).

b. Examples of 5dpf wildtype hosts engrafted with wildtype donor cells (upper) that
contribute to arteries (aISVs, DA), veins (PCV, vISVs) and lymphatic vessels (TD, ISLVs)
and examples of wildtype embryos engrafted with *MZprox1a* mutant cells
contributing arteries (aISVs, DA) and veins (vISV, PCV) but not lymphatic vessels. Note
the PL accumulation in the horizontal myoseptum..

c. The number of embryos imaged and graft sizes scored post-tranplantation. As the
contribution to the different EC types is affected by the graft size, we analysed grafts
of comparable sizes and position from wildtype and mutant embryos. Grafts were
categorised as large, medium or small. Small (spanning 1-2 somites), medium
(spanning 2-3 somites) and large grafts (spanning 4-5 somites). The number of ECs
contributing to each EC type was scored at single cell resolution and the percentage
of grafts with a contribution to each lineage calculated accordingly. The numbers
within the different graft groups indicate the number of analysed embryos within that
group.

d. The percentage of embryos with vascular grafts that display contributions to arteries,
veins and lymphatics. *MZprox1a*$^{-/-}$ mutant cells contribute to arteries and veins but
showed a near complete loss of contribution to lymphatics. The size of the grafts did
not influence the outcome. *Z1b** indicates a mixed prox1b background. p < 0.0001
from an unpaired, two-sided t-test.

DA, dorsal aorta. PCV, posterior cardinal vein. PL, parachordal LEC. TD, thoracic duct. aISV,
arterial intersegmental vessel. vISV, venous intersegmental vessel. ISLV, intersegmental
lymphatic vessel.

1035

References

Oliver, G., Kipnis, J., Randolph, G. J. & Harvey, N. L. The Lymphatic Vasculature in the 21(st) Century: Novel Functional Roles in Homeostasis and Disease. *Cell* **182**, 270-296, doi:10.1016/j.cell.2020.06.039 (2020).

Koltowska, K., Betterman, K. L., Harvey, N. L. & Hogan, B. M. Getting out and about: the emergence and morphogenesis of the vertebrate lymphatic vasculature. *Development* **140**, 1857-1870, doi:10.1242/dev.089565 (2013).

Baek, S. *et al.* The Alternative Splicing Regulator Nova2 Constrains Vascular Erk Signaling to Limit Specification of the Lymphatic Lineage. *Dev Cell* **49**, 279-292 e275, doi:10.1016/j.devcel.2019.03.017 (2019).

Koltowska, K. *et al.* Vegfc Regulates Bipotential Precursor Division and Prox1 Expression to Promote Lymphatic Identity in Zebrafish. *Cell Rep* **13**, 1828-1841, doi:10.1016/j.celrep.2015.10.055 (2015).

Deng, Y., Atri, D., Eichmann, A. & Simons, M. Endothelial ERK signaling controls lymphatic fate specification. *The Journal of clinical investigation* **123**, 1202-1215, doi:10.1172/JCI63034 (2013).

Shin, M. *et al.* Vegfc acts through ERK to induce sprouting and differentiation of trunk lymphatic progenitors. *Development* **143**, 3785-3795, doi:10.1242/dev.137901 (2016).

Srinivasan, R. S. *et al.* The Prox1-Vegfr3 feedback loop maintains the identity and the number of lymphatic endothelial cell progenitors. *Genes Dev* **28**, 2175-2187, doi:10.1101/gad.216226.113 (2014).

Oliver, G. & Srinivasan, R. S. Endothelial cell plasticity: how to become and remain a lymphatic endothelial cell. *Development* **137**, 363-372, doi:10.1242/dev.035360 (2010).

Wigle, J. T. & Oliver, G. Prox1 function is required for the development of the murine lymphatic system. *Cell* **98**, 769-778, doi:10.1016/s0092-8674(00)81511-1 (1999).

Hagerling, R. *et al.* A novel multistep mechanism for initial lymphangiogenesis in mouse embryos based on ultramicroscopy. *The EMBO journal* **32**, 629-644, doi:10.1038/emboj.2012.340 (2013).

Klotz, L. *et al.* Cardiac lymphatics are heterogeneous in origin and respond to injury. *Nature* **522**, 62-67, doi:10.1038/nature14483 (2015).

Martinez-Corral, I. *et al.* Nonvenous origin of dermal lymphatic vasculature. *Circ Res* **116**, 1649-1654, doi:10.1161/CIRCRESAHA.116.306170 (2015).

Stanczuk, L. *et al.* cKit Lineage Hemogenic Endothelium-Derived Cells Contribute to Mesenteric Lymphatic Vessels. *Cell Rep* **10**, 1708-1721, doi:10.1016/j.celrep.2015.02.026 (2015).

Srinivasan, R. S. *et al.* Lineage tracing demonstrates the venous origin of the mammalian lymphatic vasculature. *Genes Dev* **21**, 2422-2432, doi:10.1101/gad.1588407 (2007).

Srinivasan, R. S. *et al.* The nuclear hormone receptor Coup-TFII is required for the initiation and early maintenance of Prox1 expression in lymphatic endothelial cells. *Genes Dev* **24**, 696-707, doi:10.1101/gad.1859310 (2010).

Yang, Y. *et al.* Lymphatic endothelial progenitors bud from the cardinal vein and intersomitic vessels in mammalian embryos. *Blood* **120**, 2340-2348, doi:10.1182/blood-2012-05-428607 (2012).

17 Stone, O. A. & Stainier, D. Y. R. Paraxial Mesoderm Is the Major Source of
Lymphatic Endothelium. *Dev Cell* **50**, 247-255 e243,
doi:10.1016/j.devcel.2019.04.034 (2019).
18 Johnson, N. C. *et al.* Lymphatic endothelial cell identity is reversible and its
maintenance requires Prox1 activity. *Genes Dev* **22**, 3282-3291,
doi:10.1101/gad.1727208 (2008).
19 Tao, S. *et al.* Zebrafish prox1b mutants develop a lymphatic vasculature, and prox1b
does not specifically mark lymphatic endothelial cells. *PLoS One* **6**, e28934,
doi:10.1371/journal.pone.0028934 (2011).
20 van Impel, A. *et al.* Divergence of zebrafish and mouse lymphatic cell fate
specification pathways. *Development* **141**, 1228-1238, doi:10.1242/dev.105031
(2014).
21 Hogan, B. M. *et al.* Ccbe1 is required for embryonic lymphangiogenesis and venous
sprouting. *Nat Genet* **41**, 396-398, doi:10.1038/ng.321 (2009).
22 Yaniv, K. *et al.* Live imaging of lymphatic development in the zebrafish. *Nat Med* **12**,
711-716, doi:10.1038/nm1427 (2006).
23 Eng, T. C. *et al.* Zebrafish facial lymphatics develop through sequential addition of
venous and non-venous progenitors. *EMBO Rep* **20**, doi:10.15252/embr.201847079
(2019).
24 Okuda, K. S. *et al.* lyve1 expression reveals novel lymphatic vessels and new
mechanisms for lymphatic vessel development in zebrafish. *Development* **139**, 2381-
2391, doi:10.1242/dev.077701 (2012).
25 Bussmann, J. *et al.* Arteries provide essential guidance cues for lymphatic endothelial
cells in the zebrafish trunk. *Development* **137**, 2653-2657, doi:10.1242/dev.048207
(2010).
26 Cha, Y. R. *et al.* Chemokine signaling directs trunk lymphatic network formation
along the preexisting blood vasculature. *Dev Cell* **22**, 824-836,
doi:10.1016/j.devcel.2012.01.011 (2012).
27 Jung, H. M. *et al.* Development of the larval lymphatic system in zebrafish.
Development **144**, 2070-2081, doi:10.1242/dev.145755 (2017).
28 Bower, N. I. *et al.* Mural lymphatic endothelial cells regulate meningeal angiogenesis
in the zebrafish. *Nat Neurosci* **20**, 774-783, doi:10.1038/nn.4558 (2017).
29 van Lessen, M. *et al.* Intracellular uptake of macromolecules by brain lymphatic
endothelial cells during zebrafish embryonic development. *Elife* **6**,
doi:10.7554/eLife.25932 (2017).
30 Venero Galanternik, M. *et al.* A novel perivascular cell population in the zebrafish
brain. *Elife* **6**, doi:10.7554/eLife.24369 (2017).
31 Kuchler, A. M. *et al.* Development of the zebrafish lymphatic system requires
VEGFC signaling. *Curr Biol* **16**, 1244-1248, doi:10.1016/j.cub.2006.05.026 (2006).
32 Hogan, B. M. & Schulte-Merker, S. How to Plumb a Pisces: Understanding Vascular
Development and Disease Using Zebrafish Embryos. *Dev Cell* **42**, 567-583,
doi:10.1016/j.devcel.2017.08.015 (2017).
33 He, Z., Brazovskaja, A., Ebert, S., Camp, J. G. & Treutlein, B. CSS: cluster similarity
spectrum integration of single-cell genomics data. *Genome Biol* **21**, 224,
doi:10.1186/s13059-020-02147-4 (2020).
34 Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902
e1821, doi:10.1016/j.cell.2019.05.031 (2019).
35 Becht, E. *et al.* Dimensionality reduction for visualizing single-cell data using UMAP.
Nat Biotechnol, doi:10.1038/nbt.4314 (2018).

36 La Manno, G. *et al.* RNA velocity of single cells. *Nature* **560**, 494-498,
doi:10.1038/s41586-018-0414-6 (2018).
37 Thomas, P. D. *et al.* Applications for protein sequence-function evolution data:
mRNA/protein expression analysis and coding SNP scoring tools. *Nucleic Acids Res*
34, W645-650, doi:10.1093/nar/gkl229 (2006).
38 Wigle, J. T. *et al.* An essential role for Prox1 in the induction of the lymphatic
endothelial cell phenotype. *EMBO J* **21**, 1505-1513, doi:10.1093/emboj/21.7.1505
(2002).
39 Kang, J. *et al.* An exquisite cross-control mechanism among endothelial cell fate
regulators directs the plasticity and heterogeneity of lymphatic endothelial cells.
Blood **116**, 140-150, doi:10.1182/blood-2009-11-252270 (2010).
40 Jerafi-Vider, A. *et al.* VEGFC/FLT4-induced cell-cycle arrest mediates sprouting and
differentiation of venous and lymphatic endothelial cells. *Cell Rep* **35**, 109255,
doi:10.1016/j.celrep.2021.109255 (2021).
41 Koltowska, K. *et al.* The RNA helicase Ddx21 controls Vegfc-driven developmental
lymphangiogenesis by balancing endothelial cell ribosome biogenesis and p53
function. *Nat Cell Biol* **23**, 1136-1147, doi:10.1038/s41556-021-00784-w (2021).
42 Burrows, P. E. *et al.* Lymphatic abnormalities are associated with RASA1 gene
mutations in mouse and man. *Proc Natl Acad Sci U S A* **110**, 8621-8626,
doi:10.1073/pnas.1222722110 (2013).
43 Frye, M. *et al.* EphrinB2-EphB4 signalling provides Rho-mediated homeostatic
control of lymphatic endothelial cell junction integrity. *Elife* **9**,
doi:10.7554/eLife.57732 (2020).
44 Geng, X. *et al.* S1PR1 regulates the quiescence of lymphatic vessels by inhibiting
laminar shear stress-dependent VEGF-C signaling. *JCI Insight* **5**,
doi:10.1172/jci.insight.137652 (2020).
45 Grimm, L. *et al.* Yap1 promotes sprouting and proliferation of lymphatic progenitors
downstream of Vegfc in the zebrafish trunk. *Elife* **8**, doi:10.7554/eLife.42881 (2019).
46 Koltowska, K. *et al.* mafba is a downstream transcriptional effector of Vegfc
signaling essential for embryonic lymphangiogenesis in zebrafish. *Genes Dev* **29**,
1618-1630, doi:10.1101/gad.263210.115 (2015).
47 Yuan, L. *et al.* Abnormal lymphatic vessel development in neuropilin 2 mutant mice.
Development **129**, 4797-4806 (2002).
48 Termini, C. M. & Gillette, J. M. Tetraspanins Function as Regulators of Cellular
Signaling. *Front Cell Dev Biol* **5**, 34, doi:10.3389/fcell.2017.00034 (2017).
49 Wu, R. S. *et al.* A Rapid Method for Directed Gene Knockout for Screening in G0
Zebrafish. *Dev Cell* **46**, 112-125 e114, doi:10.1016/j.devcel.2018.06.003 (2018).
50 Armour, S. M. *et al.* An HDAC3-PROX1 corepressor module acts on HNF4alpha to
control hepatic triglycerides. *Nat Commun* **8**, 549, doi:10.1038/s41467-017-00772-5
(2017).
51 Liu, X. *et al.* Mitotic Implantation of the Transcription Factor Prospero via Phase
Separation Drives Terminal Neuronal Differentiation. *Dev Cell* **52**, 277-293 e278,
doi:10.1016/j.devcel.2019.11.019 (2020).
52 Qin, J. *et al.* Prospero-related homeobox (Prox1) is a corepressor of human liver
receptor homolog-1 and suppresses the transcription of the cholesterol 7-alpha-
hydroxylase gene. *Mol Endocrinol* **18**, 2424-2439, doi:10.1210/me.2004-0009 (2004).
53 Song, K. H., Li, T. & Chiang, J. Y. A Prospero-related homeodomain protein is a
novel co-regulator of hepatocyte nuclear factor 4alpha that regulates the cholesterol
7alpha-hydroxylase gene. *J Biol Chem* **281**, 10081-10088,
doi:10.1074/jbc.M513420200 (2006).

54   Takeda, Y. & Jetten, A. M. Prospero-related homeobox 1 (Prox1) functions as a novel
1183   modulator of retinoic acid-related orphan receptors alpha- and gamma-mediated
1184   transactivation. *Nucleic Acids Res* **41**, 6992-7008, doi:10.1093/nar/gkt447 (2013).
1185   55   Lawson, N. D. & Weinstein, B. M. In vivo imaging of embryonic vascular
1186   development using transgenic zebrafish. *Dev Biol* **248**, 307-318,
1187   doi:10.1006/dbio.2002.0711 (2002).
1188   56   Ciruna, B. *et al.* Production of maternal-zygotic mutant zebrafish by germ-line
1189   replacement. *Proc Natl Acad Sci U S A* **99**, 14919-14924,
1190   doi:10.1073/pnas.222459999 (2002).
1191   57   Westerfield, M. *The Zebrafish Book. A Guide for the Laboratory use of Zebrafish
1192   (Danio rerio)*. 4 edn, (University of Oregorn Press, 2000).
1193   58   Gagnon, J. A. *et al.* Efficient mutagenesis by Cas9 protein-mediated oligonucleotide
1194   insertion and large-scale assessment of single-guide RNAs. *PLoS One* **9**, e98186,
1195   doi:10.1371/journal.pone.0098186 (2014).
1196   59   Bussmann, J. & Schulte-Merker, S. Rapid BAC selection for tol2-mediated
1197   transgenesis in zebrafish. *Development* **138**, 4327-4332, doi:10.1242/dev.068080
1198   (2011).
1199   60   Okuda, K. S., Baek, S. & Hogan, B. M. Visualization and Tools for Analysis of
1200   Zebrafish Lymphatic Development. *Methods Mol Biol* **1846**, 55-70, doi:10.1007/978-
1201   1-4939-8712-2_4 (2018).
1202   61   Bower, N. I. *et al.* Vegfd modulates both angiogenesis and lymphangiogenesis during
1203   zebrafish embryonic development. *Development* **144**, 507-518,
1204   doi:10.1242/dev.146969 (2017).
1205   62   Kartopawiro, J. *et al.* Arap3 is dysregulated in a mouse model of hypotrichosis-
1206   lymphedema-telangiectasia and regulates lymphatic vascular development. *Hum Mol
1207   Genet* **23**, 1286-1297, doi:10.1093/hmg/ddt518 (2014).
1208   63   Zheng, G. X. *et al.* Massively parallel digital transcriptional profiling of single cells.
1209   *Nat Commun* **8**, 14049, doi:10.1038/ncomms14049 (2017).
1210   64   Wolock, S. L., Lopez, R. & Klein, A. M. Scrublet: Computational Identification of
1211   Cell Doublets in Single-Cell Transcriptomic Data. *Cell Syst* **8**, 281-291 e289,
1212   doi:10.1016/j.cels.2018.11.005 (2019).
1213   65   Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell
1214   transcriptomic data across different conditions, technologies, and species. *Nat
1215   Biotechnol* **36**, 411-420, doi:10.1038/nbt.4096 (2018).
1216   66   McCarthy, D. J., Campbell, K. R., Lun, A. T. & Wills, Q. F. Scater: pre-processing,
1217   quality control, normalization and visualization of single-cell RNA-seq data in R.
1218   *Bioinformatics* **33**, 1179-1186, doi:10.1093/bioinformatics/btw777 (2017).
1219   67   Zappia, L. & Oshlack, A. Clustering trees: a visualization for evaluating clusterings at
1220   multiple resolutions. *Gigascience* **7**, doi:10.1093/gigascience/giy083 (2018).
1221   68   Kejie Li, Z. O., Dongdong Lin, Michael Mingueneau, Will Chen, David Sexton,
1222   Baohong Zhang. cellxgene VIP unleashes full power of interactive visualization,
1223   plotting and analysis of scRNA-seq data in the scale of millions of cells. *bioRxiv
1224   **2020.08.28.270652**, doi:https://doi.org/10.1101/2020.08.28.270652 (2020).
1225   69   Kazenwadel, J. *et al.* GATA2 is required for lymphatic vessel valve development and
1226   maintenance. *J Clin Invest* **125**, 2979-2994, doi:10.1172/JCI78888 (2015).
1227   70   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler
1228   transform. *Bioinformatics* **25**, 1754-1760, doi:10.1093/bioinformatics/btp324 (2009).
1229   71   Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing
1230   genomic features. *Bioinformatics* **26**, 841-842, doi:10.1093/bioinformatics/btq033
1231   (2010).

1232

1233

1234