Antisense oligonucleotides targeting exon 11 are able to partially rescue the Neurofibromatosis Type 2 phenotype *in vitro.*

N.Catasús[1], I.Rosas[1,2], S.Bonache[1,2], A.Negro[1,2], A.Plana[3], H.Salvador[4], E.Serra[5], I.Blanco[1,2] and E.Castellanos[1,2] on behalf of the NF2 Spanish National Reference Centre HUGTP-ICO-IGTP

1) Clinical Genomics Research Group, Germans Trias i Pujol Research Institute (IGTP); Can Ruti Campus, Badalona, Barcelona, 08916; Spain

2) Clinical Genomics Unit, Germans Trias i Pujol University Hospital (HUGTiP), Can Ruti Campus, Badalona, Barcelona, 08916; Spain

3) Dermatology Department, Germans Trias i Pujol University Hospital (HUGTiP), Can Ruti Campus, Badalona, Barcelona, 08916; Spain

4) Pediatric Oncology Unit, Hospital Sant Joan de Déu, Esplugues, Barcelona, Spain

5) Hereditary Cancer Group, Germans Trias i Pujol Research Institute (IGTP-PMPPC); Can Ruti Campus, Badalona, Barcelona, 08916; Spain

Correspondence should be addressed to Dr. Elisabeth Castellanos (ecastellanos@igtp.cat)

Corresponding author:

Dr. Elisabeth Castellanos

Clinical Genomics Unit - Genetics Service

Laboratori Clinic Metropolitana Nord (LCMN)

Hospital Universitari Germans Trias i Pujol

Crta. Canyet, s/n. Badalona, Barcelona, Spain

Email: ecastellanos@igtp.cat

Abstract

Neurofibromatosis type 2 (NF2) is an autosomal dominant condition characterized by the development of multiple tumours of the nervous system caused by loss of function variants in the *NF2* gene. The clinical presentation of the disease is variable and related to the type of the inherited constitutional mutation. Here, we evaluated the use of antisense oligonucleotides, specifically phosphorodiamidate morpholino oligomers (PMOs), to reduce the severity of the effects of *NF2* truncating variants by converting them to milder hypomorphic forms *in vitro*. Our results showed that the PMOs designed for variants located at +/- 13 from the intron-exon boundary region interfered with the correct splicing signalling of both *NF2* wild-type and mutated alleles. On the other hand, we were able to specifically induce the skipping of exons 4, 8 and 11 of both the mutated and the WT allele maintaining the *NF2* gene reading frame at cDNA level. Only the skipping of exon 11 produced a hypomorphic Merlin (Merlin-e11) able to partially rescue the phenotype observed in primary fibroblast cultures from NF2 patients harbouring a germline loss of function variant. This encouraging result is an *in vitro* proof of concept of the use of antisense therapy to develop personalized therapies for NF2.

INTRODUCTION

Neurofibromatosis type 2 (NF2) (OMIM 101000) is an autosomal dominant (AD) disease caused by loss-of-function (LOF) variants in the *NF2* tumour suppressor gene (22q12.2) [1,2]. The reported incidence is between 1 in 28,000-40,000 [3] and the hallmark of the disease is the presence of bilateral vestibular schwannomas (BVS) which clinically present with hearing loss, tinnitus and imbalance [4]. Common features of NF2 include cranial, spinal, peripheral nerve, and intradermal schwannomas; cranial and spinal meningiomas; and intrinsic central nervous system (CNS) tumours, usually spinal ependymomas [5,6]. Despite their benign nature, tumours are associated with high morbidity due to their multiplicity and anatomical location, which cause pain and nerve dysfunction leading to a reduced quality of life and life expectancy [7].

There are limited therapies available for these patients and radiotherapy and surgery for tumour resection are the classical treatments [8]. Currently, medical therapy is also an option since Bevacizumab, an anti-angiogenic monoclonal antibody, has been demonstrated to be an effective treatment for vestibular schwannomas (VS) inducing the reduction of tumour growth and an improvement in hearing in adult NF2 patients [6,9]. However, Bevacizumab does not seem to be equally effective in all NF2 patients and, in addition, it has no effect on other types of tumours present due to the disease, such as meningiomas [10]. Efforts have been made to develop other therapies for NF2-related tumours, such as Losartan (antihypertensive), Lapatinib (tyrosine kinase inhibitor) or AR-42 (Histone deacetylase (HDAC) inhibitor) [11–14], but there still remains a need to identify clinically relevant therapeutic targets to treat NF2 patients.

The *NF2* gene consists of 17 exons [15,16] with a wide mutational spectrum, 85-90% are germline point mutations and 10-15% are large deletions distributed throughout the entire gene [17,18]. *NF2* encodes for Merlin (Moesin-Ezrin-Radixin-Like protein), a 595aa scaffold protein involved in membrane protein organization, regulation of cell-cell adhesion and with a regulatory role in the cytoskeleton architecture [19,20]. Merlin is comprised of three structural

regions: a N-terminal FERM domain which is subdivided into three globular domains (F1, F2 and F3), an α-helical coiled-coil domain and a C-terminal hydrophilic tail [21].

The genotype-phenotype correlation of NF2 has long been studied [22–30]. For NF2, it is well reported that nonsense variants and large deletions, which account for around 30% of cases, are associated with severe manifestations of the disease, whereas missense, or in-frame variants present a milder form of NF2. Besides, about 25% of the point mutations affect the correct splicing of the *NF2* gene, and this mechanism is associated with a variable severity of the disease [27,29,31]. The well-known association between the genetic variant type and disease severity may constitute an opportunity to develop personalized gene therapies based on the specific variant of the *NF2* gene.

Antisense gene therapy has been successfully used to regulate gene expression in some pathologies. Most of the target diseases are inherited in an autosomal recessive manner, such as Spinal Muscular Atrophy (SMA) [32], homozygous familial hypercholesterolemia (HoFH) [33], or Duchenne muscular dystrophy (DMD) [34] the latter two have been approved by the FDA and are amongst others currently in the advanced stages of clinical trials, [35,36]. Furthermore, there are also examples for AD diseases, such as Retinitis pigmentosa (RP) [37], amyotrophic lateral sclerosis [38] or Huntington's disease [39]. In all cases, antisense oligonucleotides (ASOs) have been used. These are single-stranded modified DNA-like molecules of 15 to 30 nucleotides that act at RNA level by specifically correcting or changing the expression of the target protein to a less deleterious form [40,41]. ASOs have been therapeutically used to reduce protein production whether via RNase H cleavage or mRNA-ASO complex to block mRNA-ribosome interaction. It is also possible to modulate the splice signalling and among the different types of ASOs, Phosphorodiamidate Morpholino Oligomers (PMOs) have highly desirable molecular properties that make them suitable for this last approach. In PMOs the deoxyribose sugar element is replaced by a six-member morpholino ring, and the phosphodiester inter-subunit bonds are

replaced by phosphorodiamidate linkages; all of which produce a structure that provides high-targeting specificity, stability, low toxicity and resistance to nucleases, assuring the maintenance of a good long-term activity within the cell [42].

In this study we assessed two different approaches to test *in vitro* antisense therapy for NF2: the first consisted of modulating and correcting the aberrant splicing of the *NF2* gene originating from different point-mutations located near the canonical splice site; the second consisted of forcing the skipping of those exons that carry a frameshift or a nonsense variant, while preserving the reading frame. Thus, taking into consideration the NF2 genotype-phenotype association, nonsense variants that generate a more severe phenotype could be altered at RNA level to generate less deleterious protein forms.

RESULTS

ANTISENSE THERAPY FOR *NF2* SPLICING VARIANTS

The experimental approach for *NF2* splicing variants consisted of designing specific PMOs for each variant to be tested in order to mask the aberrant splice signalling triggered by the pathogenic variant and promote the spliceosome to properly recognize the correct splice site without altering the wild type (WT) allele (Figure 1). Two different PMOs were designed for each variant. The first design was a 25-mer variant-specific facilitated by Gene Tools according to their guidelines for intron and exon binding subunits and self-complementary moieties. We made a second design using splice site predictors to study the variant surrounding sequence (acceptor splicing sites, exonic splicing enhancers (ESEs) or exonic splicing silencers (ESSs)) (Human Splice Finder) in order to avoid the induction of possible new splicing signals due to the effect of the PMO. In addition, the design avoided masking the canonical splicing GT-AG sites and branch points (YNYYRAY) when possible (Table 1, Figure 2A).

We tested the PMOs' efficacy in primary fibroblasts from NF2 patients harbouring heterozygous germline variants located near the canonical splice site of exons 3, 8 and 15 (Table S1) from the Spanish National Reference Centre (CSUR) on Phakomatoses. A dose-response study was performed to test the effect on splicing of different concentrations of PMOs after 24h of treatment and the efficiency was assessed at cDNA level by RT-PCR. The effect of PMOs on the splicing of *NF2* was analyzed by electrophoresis and confirmed by Sanger sequencing, revealing a maximum effect at 20μM. A time course was carried out treating fibroblasts with 20μM PMO during 24h, 48h and 72h. PMO treatment in Patient_Spl_1 and Patient_Spl_2 increased the skipping of exons 15 and 8 respectively in both of the designs, in contrast to the expected effect of blocking only the germline variant and restoring the NF2 correct splicing without altering the transcription of the WT allele (Figure 2B, B.I and B.II). Similarly, the treatment of PMOs of patients Patient_Spl_3 and Patient_Spl_4, both with nearby variants near the canonical splice site of exon 2, revealed that the variant-specific PMO caused the skipping of both exons 2 and 3 of the NF2 gene instead of correcting the aberrant NF2 splicing due to germline variants (Figure 2B, B.III and B.IV). The same effect was observed with the two different designs and the effect was confirmed by Sanger sequencing (Figure S1). Fibroblasts from healthy donors (FibCtrl) were also treated with each designed variant-specific PMO and they showed the same effect as fibroblasts derived from patients. The effect of the treatment on Merlin was assayed by Western Blot and no improvement in Merlin levels was detected after the treatment at the studied time points (Figure S2).

ANTISENSE THERAPY FOR *NF2* TRUNCATING VARIANTS

Differently from the previous approach, this aimed to test if PMOs could be used to specifically force the skipping of in-frame exons carrying a frameshift or nonsense variant, thus maintaining the *NF2* gene reading frame, and test if the generated Merlin protein could act as an hypomorphic form, still preserving partial function and to some extent able to rescue an *in vitro* phenotype. In this way, taking into account the genotype-phenotype association in which

nonsense or truncating variants are related to more severe phenotypes than in-frame variants, those that generate a more severe phenotype could be modified to less deleterious forms (Figure 3), assessing if the effect of the exon skipping on both alleles was more beneficial than a truncating variant in heterozygosis. A pair of PMOs were designed to be complementary to both 5'- and 3'- intron-exon boundaries to induce skipping of exons 4, 8 or 11 of the *NF2* gene (Exon-Specific PMOs) (Table 2).

Fibroblasts from healthy donors were treated with a pair of PMOs designed to target the acceptor and the donor sites of each exon. A dose response study confirmed the efficiency of the treatment showing full skipping of exons 4 and 8 at 20μM (Figure S3 A and B). For exon 11, two different designs were tested (Table 2; PMO_ES11_v1 and PMO_ES11_v2), while the first (PMO_ES11_v1) failed to induce skipping (data not shown), the second design (PMO_ES11_v2) showed a maximum effect when inducing the skipping of exon 11 at 40μM, and the cells showed an expression of more than 50% of the exon-less form (Figure S3.C). As for the splicing variants, a time course experiment was performed to test the effect at distinct time points (24h, 48h and 72h). In all cases, no differences were observed over time, indicating that the maximum effect of the treatment at RNA level was reached at 24h and maintained at 72h (Figure S3).

We analyzed whether the skipping of both the in-frame *NF2* exon harbouring a truncating variant and the wild-type *NF2* allele, was less deleterious than harbouring a non-functional allele in primary NF2 patients' derived fibroblasts. First, we studied the capacity of restoring Merlin levels *in vitro* after the PMO treatment (Figure 4). Unexpectedly, Merlin western blot showed progressive loss of protein signal when targeting exons 4 (Merlin-e4) or 8 (Merlin-e8), studied in two and five samples respectively at the 24h, 48h and 72h time points (Figure 4, A and B). In order to elucidate if the induced skipping of exons 4 or 8 caused a reduction of Merlin levels; or if there was a loss in the detection of Merlin by the antibody used, possibly due to a misfolded protein; the exon-less NF2 cDNAs were cloned to a N-terminal FLAG® Tag Expression Vector and transfected to HeLa cells. Flag-Merlin Western Blot showed that there was little or no expression

of Merlin-e4 or Merlin-e8, indicating that these new Merlin forms could not be synthesized or were degraded shortly after synthesis (Figure S4).

The skipping of exon 11 generated a shorter form of Merlin, indicating that a potential hypomorphic Merlin could be resulting from the skipping of this exon (Merlin-e11) (Figure 4C).

E11 PMO treatment of *NF2*+/- fibroblasts partially rescued the phenotype *in vitro*

Functionality of Merlin-e11 was studied in primary fibroblasts harbouring a heterozygous truncating variant in exon 11: one from an adult patient (Patient_ES11_1) and another from a paediatric patient (Patient_ES11_2) (Table S1). Given the complexity of determining differences in severity in the phenotype from studying the signalling pathways in which Merlin is involved [29], physiological read outs were assessed. Due to the role of Merlin in cytoskeletal organization [38,39], Merlin-e11's functional activity was first tested through actin immunofluorescence. Phalloidin staining revealed an improvement of the organizational capacity of the cytoskeleton after PMO treatment, although the phenotype observed in both samples was slightly different: patient_ES11_1 showed prominent membrane ruffles that decreased in response to treatment, (Figure 5A) while in patient_ES11_2 we observed an improvement in the cell-cell contact organization (Figure 5B). Specifically, paediatric derived NF2+/- fibroblasts showed an improvement in cell culture, as it seemed that the treatment and expression of Merlin-e11 allowed fibroblasts to recover part of the cell-cell contact inhibition and grow as a monolayer instead of growing overlapped.

In addition, proliferation rates were tested after 72h of treatment by flow cytometry. patient_ES11_1 showed a statistically significant depletion of proliferation when treated with PMOs (65.6%, $p<0.05$) (Figure 6A) while patient_ES11_2 showed a less pronounced decrease (25.1%) (Figure 6B). Considering both samples, primary fibroblasts with a truncating variant in exon 11 showed a 51% reduction in proliferation when compared with the vehicle (Endo-Porter,

p<0.01) and 39.7% compared to treatment control (p<0.01), on average (Figure 6C, Figure S5). PMO treatment did not induce significant differences in cell viability (Figure S6).

DISCUSSION

Neurofibromatosis type 2 (NF2) is a multisystem genetic disorder for which the development of effective therapeutic options with no adverse consequences is needed. The clinical presentation of the disease is variable and related to the type of germline variant inherited in the *NF2* gene [8]. In this context, the development of personalized therapies based on the type of *NF2* variant constitutes an opportunity. In this study, we tested the use of PMOs as antisense therapy to reduce the severity of the effects of *NF2* truncating and splicing variants by changing them to milder hypomorphic forms *in vitro* by two different approaches.

The first approach targeted the splicing variants with the aim of concealing the erroneous splice signal caused by the pathogenic variant and restoring levels of Merlin, in a similar way to the approach that our group used for a *NF2* deep intronic variant [43]. In this study it has not been possible to design PMOs that permit correction of the splicing signalling and furthermore, it was not possible to achieve specificity to target the mutated allele as the same effect was observed in both patient-derived fibroblasts and control fibroblasts. We hypothesize that when the splicing variant is within +/- 13 nucleotides of the acceptor or donor splice site, the PMO could allosterically interfere with the branch point and the U2 snRNA incorporation into the spliceosome [44]. The spatiotemporal localization of the two alleles during splicing could prevent the PMO from specifically targeting the mutated allele and thus it would interfere in the cis-regulation of RNA elements of both alleles equally, resulting in an increased pathogenic effect [45]. Since the approach requires the PMO to be specific for the allele harbouring the pathogenic variant, there is little room for variation in the PMO design.

As far as we know, there are no studies in the literature that have been able to repair the LOF variant effect and recover gene function through the use of ASOs for a splicing mutation located in the canonical splice regions in genes causing autosomal dominant diseases. It has only been possible through CRISPR/Cas technology by the induction of single nucleotide substitutions [46,47]. Allele-specific approaches have been successfully tested for intronic variants through splice modulation ASOs that mask the erroneous signalling and given their intronic location and distance to splicing regulatory regions, proper protein synthesis can be recovered [48–52].

Furthermore, the approach for the truncating variants tried to benefit from the fact that these variants are associated with more severe phenotypes in comparison to in-frame variants in *NF2* [28,30,53]. Thus, we proposed inducing skipping of the exon carrying the variant with the hypothesis of generating potential homozygous hypomorphic forms of Merlin, still aiming to ameliorate the pathogenicity of the heterozygous truncating variant *in vitro*. The skipping of exons 4 or 8 harbouring *NF2* LOF variants was not able to generate an exon-less Merlin, because Merlin-e4 and Merlin-48 were degraded. Exons 4 and 8 of the *NF2* gene code for the subdomains B and C respectively of the FERM domain, where the majority of reported pathogenic variants occur; and which has the most potential for anti-mitogenic signalling and tumour suppressor function [17,21]. The observed results could be due to the fact that the lack of an exon coding for the FERM domain would cause an unbearable aberrant folding of the protein and its synthesis would be prevented or it would be forcibly degraded. In addition, the FERM domain is responsible for the intramolecular interaction with the C-terminal domain, which results in Merlin being in a closed and active conformation [54–56], thus, if this folding fails, it is possible that the resulting protein would not be able to execute the suppressor activity.

For autosomal recessive diseases, such as SMA or DMD [57,58], an autosomal dominant with a dominant negative effect, Frontotemporal dementia with parkinsonism-17 (FTDP-17) or RP, allele specific ASOs have been used to induce degradation of the disease-causing allele via RNAse

H activity and for this purpose, the native DNA backbone is not modified [44,59–61]. Recent evidence suggests that Merlin could act in a negative dominant manner through its self-dimerization through FERM-FERM interactions [62], so the use of RNAse H dependent ASOs could be a possible antisense approach for mutations located in the FERM domain, in both in-frame and out of frame exons. In addition, in contrast to the approach tested in this study, we hypothesize that the degradation of the mutant transcript could prevent interference with the spliceosome during the splicing of the wild-type allele, and therefore, the use of RNAse H dependent ASOs should be also taken into consideration for patients harbouring splicing variants in the *NF2* gene. In the light of our findings, the degradation of the *NF2* mutated allele could bring an improvement in the *in vitro* phenotype and open the door for future gene therapy approaches.

When targeting exon 11 of the *NF2* gene, an exon-less Merlin was expressed (Merlin-e11). Merlin, as a member of the ERM protein family, is involved in the link between actin cytoskeleton and adherent junctions with a relevant role in dynamic cytoskeleton remodelling, such as in membrane ruffling or in the formation of actin micro-spikes. Consistent with other studies which found that Merlin loss leads to a dramatic increase of membrane ruffles [63], here we showed a decrease in membrane ruffle formation in response to the expression of Merlin-e11 that could be associated to a partial rescue of the phenotype. Additionally, we showed an improvement in the cytoskeleton organization after treatment, which could be due to a recovery of the Merlin function as a scaffold protein regulating the linkage between membrane proteins and the cortical cytoskeleton [52,64,65]. Consistent with the role as a tumour suppressor protein, Merlin-deficient cells loose contact inhibition of proliferation [54,66,67] and although much remains to be discovered about the adhesive signalling pathways involved, in this study we showed that the induction of Merlin-e11 by PMO treatment contributed to the recovery of cell-cell contact inhibition of proliferation in fibroblasts derived from NF2 patients harbouring truncating mutations in exon 11.

This work is a proof of concept of the use antisense therapy *in vitro* as a personalized therapy for NF2 patients harbouring truncating variants at exon 11 of the *NF2* gene. Further studies will be needed to elucidate whether there are other mechanisms or designs to apply ASOs as a therapeutic approach for this disease, as in our experience it has not been possible to recover Merlin levels for variants located at in-frame exons 4 or 8, or to correct splicing signals *in vitro* in the four variants located near canonical splice sites tested, although other splicing variants should be tested to confirm these results.

On the other hand, our approach has shown encouraging results when targeting exon 11, although further characterization assays need to be performed to better characterize the function of Merlin-e11 before considering it for further *in vivo* or pre-clinical assays.

MATERIAL AND METHODS

All procedures performed were in accordance with the ethical standards of the IGTP Institutional Review Board, which approved this study and with the 1964 Helsinki declaration and its later amendments.

Samples: samples from 14 NF2 patients from the Spanish Reference Centre were included in the study (Table S1) after providing written informed consent. Samples included in the study were those harbouring a splicing variant or a truncating variant in an in-frame exon of NF2.

Genetic NF2 test: Genetic testing was performed using the customized I2HCP panel [66] in blood or tissue when available. To detect splicing variants, patients were studied both at RNA and DNA levels. Variants were validated by Sanger sequencing and analyzed using CLC-workbench (Qiagen, Hilden, Germany) in all primary cultures established.

Antisense specificity and efficacy: 25-mer specific Phosphorodiamidate Morpholino Oligomers (PMOs) for four patients with splicing variants, and a pair of PMOs to induce exon skipping of exons 4, 8, and 11 of the *NF2* gene were designed, synthesized and purified by Gene Tools (Philomath, OR, USA). Endo-Porter (Gene Tools) was used as vehicle to deliver PMOs into cells, according to the manufacturer's instructions. Patient primary fibroblasts carrying *NF2* germline pathogenic variants were treated with the specific single or paired PMOs. A dose-response and a time course study were performed to set-up treatment conditions and the effect of PMOs was assessed by RT-PCR and Sanger sequencing. Levels of Merlin after PMO treatment were analyzed by Western Blot. For all experiments, the correct effect of the treatment at RNA level was confirmed prior to Western Blot analysis.

Analysis of Merlin-e11 phenotype recovery: an actin immunocytochemistry assay was performed to study the cytoskeletal organization after PMO treatment using Phalloidin (Molecular Probes, Invitrogen). Click-iT™ EdU Alexa Fluor™ 488 Flow Cytometry Assay Kit (Molecular Probes, Invitrogen) was used, according to the manufacturer's protocol, to determine cell proliferation. Cell viability was assessed using RealTime-Glo™ MT Cell Viability Assay (Promega) following the manufacturer's instructions.

NF2 **Cloning and expression:** *NF2* gene was cloned using the Gateway® Gene Cloning system (Invitrogen) after treatment to induce the exon-skipping to a FLAG® Tag Expression Vector. The expression vector was transfected into HeLa cells (Lipofectamine LTX with Plus Reagent, Invitrogen) according to the manufacturer's instructions. Flag expression was assessed by Western Blot (Monoclonal ANTI-FLAG® M2 antibody, Sigma-Aldrich).

Statistics: Statistical analysis was carried out using GraphPad Prism software v7. The Shapiro–Wilk test was used to test normality and for multiple group comparisons, a two-tailed unpaired t test was performed. Statistical significance is indicated by *p<0.05, **p<0.01, and ***p<0.001.

For more information, see Supplementary Materials and Methods.

ACKNOWLEDGEMENTS

We thank the HGTP Clinical Services and staff for their collaboration in generating and collecting patient samples and data and the Hereditary Cancer Group at the IGTP for their help in improving this work. We thank the IGTP Flow Cytometry core facility and its staff for their contribution and technical support. We would like to acknowledge the constant support of the different NF lay associations: Asociación de Afectados de Neurofibromatosis, Chromo22 and ACNefi. This work has been supported by: The Children's Tumor Foundation (CTF-2019-05-005), the Spanish Association of NF Affected (AANF), Fundación Proyecto Neurofibromatosis, the Catalan NF Association (AcNeFi), the Ministry Science and Innovation (PI20/00215); Fundació La Marató de TV3 (126/C/2020) and the Government of Catalonia (2017 SGR 496).

AUTHOR CONTRIBUTORS

E.C. conceived the study and wrote the manuscript that was revised, corrected, and improved by all the authors. N.C. performed most of the experimental work, analyzed the data, generated the figures for the paper and contributed to writing the manuscript. I.R, S.B. and A.N. performed experimental work. A.P. and H.S. contributed with sample collection. E.S. and I.B. provided scientific input. All authors approved the final version of the manuscript.

REFERENCES

Evans DGR, Huson SM, Donnai D, *et al.* A genetic study of type 2 neurofibromatosis in the United Kingdom . II . Guidelines for genetic counselling. 1992;:847–52.

Evans DG, Howard E, Giblin C, *et al.* Birth incidence and prevalence of tumor-prone syndromes: Estimates from a UK family genetic register service. *American Journal of Medical Genetics, Part A* 2010;**152**:327–32. doi:10.1002/ajmg.a.33139

Evans DG, Bowers NL, Tobi S, *et al.* Schwannomatosis: a genetic and epidemiological study. *Journal of Neurology, Neurosurgery & Psychiatry* 2018;**89**:1215–9. doi:10.1136/jnnp-2018-318538

Lloyd SKW, Evans DGR. *Neurofibromatosis type 2 (NF2): diagnosis and management*. 1st ed. Elsevier B.V. 2013. doi:10.1016/B978-0-444-52902-2.00054-0

Evans DGR. Orphanet Journal of Rare Diseases. 2009;**2**:1–11. doi:10.1186/1750-1172-4-16

Forde C, King AT, Rutherford SA, *et al.* Disease course of neurofibromatosis type 2: A 30-year follow-up study of 353 patients seen at a single institution. *Neuro-Oncology* 2021;**23**:1113–24. doi:10.1093/neuonc/noaa284

Lloyd SKW, Evans DGR. *Neurofibromatosis type 2 (NF2): diagnosis and management.* 1st ed. Elsevier B.V. 2013. doi:10.1016/B978-0-444-52902-2.00054-0

Lloyd SKW, Evans DGR. *Neurofibromatosis type 2 (NF2): diagnosis and management*. 1st ed. Elsevier B.V. 2013. doi:10.1016/B978-0-444-52902-2.00054-0

Plotkin SR, Duda DG, Muzikansky A, *et al.* Multicenter, Prospective, Phase II and Biomarker Study of High-Dose Bevacizumab as Induction Therapy in Patients With Neurofibromatosis Type 2 and Progressive Vestibular Schwannoma. *Journal of Clinical Oncology* 2019;:JCO.19.01367. doi:10.1200/jco.19.01367

Nunes FP, Merker VL, Jennings D, *et al.* Bevacizumab Treatment for Meningiomas in NF2: A Retrospective Analysis of 15 Patients. *PLoS ONE* 2013;**8**. doi:10.1371/journal.pone.0059941

Wu L, Vasilijic S, Sun Y, *et al.* Losartan prevents tumor-induced hearing loss and augments radiation efficacy in NF2 schwannoma rodent models. 2021. https://www.science.org

Karajannis MA, Legault G, Hagiwara M, *et al.* Phase II trial of lapatinib in adult and pediatric patients with neurofibromatosis type 2 and progressive vestibular schwannomas. *Neuro-Oncology* 2012;**14**:1163–70. doi:10.1093/neuonc/nos146

Osorio DS, Hu J, Mitchell C, *et al.* Effect of lapatinib on meningioma growth in adults with neurofibromatosis type 2. *Journal of Neuro-Oncology* 2018;**139**:749–55. doi:10.1007/s11060-018-2922-5

Welling DB, Collier KA, Burns SS, *et al.* Early phase clinical studies of AR-42, a histone deacetylase inhibitor, for neurofibromatosis type 2-associated vestibular schwannomas and meningiomas. *Laryngoscope Investigative Otolaryngology* 2021;**6**:1008–19. doi:10.1002/lio2.643

Schmucker B, Tang Y, Kressel M. Novel alternatively spliced isoforms of the neurofibromatosis type 2 tumor suppressor are targeted to the nucleus and cytoplasmic granules. *Human Molecular Genetics* 1999;**8**:1561–70. doi:10.1093/hmg/8.8.1561

Su F, Zhou Z, Su W, *et al.* A novel alternative splicing isoform of NF2 identified in human schwann cells. *Oncology Letters* 2016;**12**:977–82. doi:10.3892/ol.2016.4685

Ahronowitz I, Xin W, Kiely R, *et al.* Mutational Spectrum of the NF2 Gene : A Meta-Analysis of 12 Years of Research and Diagnostic Laboratory Findings. 2007;**28**. doi:10.1002/humu

Halliday D, Parry A, Evans DG. Neurofibromatosis type 2 and related disorders. *Current Opinion in Oncology* 2019;**31**:562–7. doi:10.1097/CCO.0000000000000579

Asthagiri AR, Parry DM, Butman JA, *et al.* Neurofibromatosis type 2. *Lancet* 2009;**373**:1974–86. doi:10.1016/S0140-6736(09)60259-2

Chiasson-MacKenzie C, Morris ZS, Baca Q, *et al.* NF2/Merlin mediates contact-dependent inhibition of EGFR mobility and internalization via cortical actomyosin. *Journal of Cell Biology* 2015;**211**:391–405. doi:10.1083/jcb.201503081

Cooper J, Giancotti FG. Molecular insights into NF2/Merlin tumor suppressor function. *FEBS Letters* 2014;**588**:2743–52. doi:10.1016/j.febslet.2014.04.001

Evans DGR, Trueman L, Wallace A, *et al.* Genotype / phenotype correlations in type 2 neurofibromatosis (NF2): evidence for more severe disease associated with truncating mutations. 1998;:450–5.

Heineman TE, Evans DGR, Campagne F, *et al.* In silico analysis of NF2 gene missense mutations in neurofibromatosis type 2: From genotype to phenotype. *Otology and Neurotology* 2015;**36**:908–14. doi:10.1097/MAO.0000000000000639

Selvanathan SK, Shenton A, Ferner R, *et al.* Further genotype - phenotype correlations in neurofibromatosis 2. *Clinical Genetics* 2010;**77**:163–70. doi:10.1111/j.1399-0004.2009.01315.x

Halliday D, Emmanouil B, Pretorius P, *et al.* Genetic severity score predicts clinical phenotype in NF2. *Journal of Medical Genetics* 2017;**54**:657–64. doi:10.1136/jmedgenet-2017-104519

Emmanouil B, Houston R, May A, *et al.* Progression of hearing loss in neurofibromatosis type 2 according to genetic severity. *Laryngoscope* 2019;**129**:974–80. doi:10.1002/lary.27586

Halliday D, Emmanouil B, Vassallo G, *et al.* Trends in phenotype in the English paediatric neurofibromatosis type 2 cohort stratified by genetic severity. *Clinical Genetics* 2019;**96**:151–62. doi:10.1111/cge.13551

Baser ME, Kuramoto L, Joe H, *et al.* Genotype-Phenotype Correlations for Nervous System Tumors in Neurofibromatosis 2: A Population-Based Study. *The American Journal of Human Genetics* 2004;**75**:231–9. doi:10.1086/422700

Catasús N, Garcia B, Galván-Femenía I, *et al.* Revisiting the UK genetic severity score for NF2: A proposal for the addition of a functional genetic component. *Journal of Medical Genetics* 2021;:1–9. doi:10.1136/jmedgenet-2020-107548

Ruttledge MH, Andermann a a, Phelan CM, *et al.* Type of mutation in the neurofibromatosis type 2 gene (NF2) frequently determines severity of disease. *American Journal of Human Genetics* 1996;**59**:331–42.

Halliday D, Emmanouil B, Pretorius P, *et al.* Genetic Severity Score predicts clinical phenotype in NF2. 2017;:657–64. doi:10.1136/jmedgenet-2017-104519

Hua Y, Vickers TA, Baker BF, *et al.* Enhancement of SMN2 exon 7 inclusion by antisense oligonucleotides targeting the exon. *PLoS Biology* 2007;**5**:729–44. doi:10.1371/journal.pbio.0050073

Crooke RM, Graham MJ, Lemonidis KM, *et al.* An apolipoprotein B antisense oligonucleotide lowers LDL cholesterol in hyperlipidemic mice without causing hepatic steatosis. *Journal of Lipid Research* 2005;**46**:872–84. doi:10.1194/jlr.M400492-JLR200

McClorey G, Moulton HM, Iversen PL, *et al.* Antisense oligonucleotide-induced exon skipping restores dystrophin expression in vitro in a canine model of DMD. *Gene Therapy* 2006;**13**:1373–81. doi:10.1038/sj.gt.3302800

Stein CA, Castanotto D. FDA-Approved Oligonucleotide Therapies in 2017. *Molecular Therapy* 2017;**25**:1069–75. doi:10.1016/j.ymthe.2017.03.023

Xiong H, Veedu RN, Diermeier SD. Recent advances in oligonucleotide therapeutics in oncology. *International Journal of Molecular Sciences* 2021;**22**. doi:10.3390/ijms22073295

Dominant NREA. Antisense Oligonucleotide-Based Downregulation of. ;**3**:1–12.

Miller T, Cudkowicz M, Shaw PJ, *et al.* Phase 1–2 Trial of Antisense Oligonucleotide Tofersen for SOD1 ALS . *New England Journal of Medicine* 2020;**383**:109–19. doi:10.1056/nejmoa2003715

Tabrizi SJ, Leavitt BR, Landwehrmeyer GB, *et al.* Targeting Huntingtin Expression in Patients with Huntington's Disease. *New England Journal of Medicine* 2019;**380**:2307–16. doi:10.1056/nejmoa1900907

Gleave ME, Monia BP. Antisense therapy for cancer. *Nature Reviews Cancer* 2005;**5**:468–79. doi:10.1038/nrc1631

Levin AA. Treating Disease at the RNA Level with Oligonucleotides. *New England Journal of Medicine* 2019;**380**:57–70. doi:10.1056/nejmra1705346

Sazani P, Kang SH, Maier MA, *et al.* Nuclear antisense effects of neutral, anionic and cationic oligonucleotide analogs. *Nucleic acids research* 2001;**29**:3965–74. doi:10.1093/nar/29.19.3965

Castellanos E, Rosas I, Solanes A, *et al.* In vitro antisense therapeutics for a deep intronic mutation causing Neurofibromatosis type 2. *European Journal of Human Genetics* 2013;**21**:769–73. doi:10.1038/ejhg.2012.261

Tang Z, Zhao J, Pearson ZJ, *et al.* Rna-targeting splicing modifiers: Drug development and screening assays. *Molecules* 2021;**26**. doi:10.3390/molecules26082263

Maass PG, Barutcu AR, Shechner DM, *et al.* Spatiotemporal allele organization by allele-specific CRISPR live-cell imaging (SNP-CLING). *Nature Structural and Molecular Biology* 2018;**25**:176–84. doi:10.1038/s41594-017-0015-3

György B, Nist-Lund C, Pan B, *et al.* Allele-specific gene editing prevents deafness in a model of dominant progressive hearing loss. *Nature Medicine* 2019;**25**:1123–30. doi:10.1038/s41591-019-0500-9

Kocher T, Peking P, Klausegger A, *et al.* Cut and Paste: Efficient Homology-Directed Repair of a Dominant Negative KRT14 Mutation via CRISPR/Cas9 Nickases. *Molecular Therapy* 2017;**25**:2585–98. doi:10.1016/j.ymthe.2017.08.015

Tomkiewicz TZ, Suárez-Herrera N, Cremers FPM, *et al.* Antisense oligonucleotide-based rescue of aberrant splicing defects caused by 15 pathogenic variants in abca4. *International Journal of Molecular Sciences* 2021;**22**. doi:10.3390/ijms22094621

Aguti S, Bolduc V, Ala P, *et al.* Exon-Skipping Oligonucleotides Restore Functional Collagen VI by Correcting a Common COL6A1 Mutation in Ullrich CMD. *Molecular Therapy - Nucleic Acids* 2020;**21**:205–16. doi:10.1016/j.omtn.2020.05.029

Panagiotopoulos AL, Karguth N, Pavlou M, *et al.* Antisense Oligonucleotide- and CRISPR-Cas9-Mediated Rescue of mRNA Splicing for a Deep Intronic CLRN1 Mutation. *Molecular Therapy - Nucleic Acids* 2020;**21**:1050–61. doi:10.1016/j.omtn.2020.07.036

Pros E, Fernández-Rodríguez J, Canet B, *et al.* Antisense therapeutics for neurofibromatosis type 1 caused by deep intronic mutations. *Human Mutation* 2009;**30**:454–62. doi:10.1002/humu.20933

Castellanos E, Carrato C, Rosas I, *et al.* In vitro antisense therapeutics for a deep intronic mutation causing Neurofibromatosis type 2. 2013;:769–73. doi:10.1038/ejhg.2012.261

Kluwe L, MacCollin M, Tatagiba M, *et al.* Phenotypic variability associated with 14 splice-site mutations in the NF2 gene. *American Journal of Medical Genetics* 1998;**77**:228–33. doi:10.1002/(SICI)1096-8628(19980518)77:3<228::AID-AJMG8>3.0.CO;2-L

Morrison H, Sherman LS, Legg J, *et al.* The NF2 tumor suppressor gene product, merlin, mediates contact inhibition of growth through interactions with CD44. *Genes and Development* 2001;**15**:968–80. doi:10.1101/gad.189601

Scoles DR. The merlin interacting proteins reveal multiple targets for NF2 therapy. 2008;**1785**:32–54. doi:10.1016/j.bbcan.2007.10.001

Li Q, Nance MR, Kulikauskas R, *et al.* Self-masking in an Intact ERM-merlin Protein: An Active Role for the Central α-Helical Domain. *Journal of Molecular Biology* 2007;**365**:1446–59. doi:10.1016/j.jmb.2006.10.075

Passini MA, Bu J, Richards AM, *et al.* Antisense oligonucleotides delivered to the mouse CNS ameliorate symptoms of severe spinal muscular atrophy. *Science Translational Medicine* 2011;**3**. doi:10.1126/scitranslmed.3001777

Wheeler TM, Leger AJ, Pandey SK, *et al.* Targeting nuclear RNA for in vivo correction of myotonic dystrophy. *Nature* 2012;**488**:111–7. doi:10.1038/nature11362

Panza F, Lozupone M, Seripa D, *et al.* Development of disease-modifying drugs for frontotemporal dementia spectrum disorders. *Nature Reviews Neurology* 2020;**16**:213–28. doi:10.1038/s41582-020-0330-x

Murray SF, Jazayeri A, Matthes MT, *et al.* Allele-specific inhibition of rhodopsin with an antisense oligonucleotide slows photoreceptor cell degeneration. *Investigative Ophthalmology and Visual Science* 2015;**56**:6362–75. doi:10.1167/iovs.15-16400

DeVos SL, Miller RL, Schoch KM, *et al.* Tau reduction prevents neuronal loss and reverses pathological tau deposition and seeding in mice with tauopathy. *Science Translational Medicine* 2017;**9**. doi:10.1126/scitranslmed.aag0481

Hennigan RF, Thomson CS, Ratner N. Merlin Tumor Suppressor Function is Regulated by PIP 2 -Mediated Dimerization . 2021;:1–24.

The Nf2 Tumor Suppressor, Merlin, Functions in Rac-Dependent Signaling. http://www.developmental.

Chiasson-Mackenzie C, Morris ZS, Liu CH, *et al.* Merlin/ERM proteins regulate growth factor-induced macropinocytosis and receptor recycling by organizing the plasma membrane:Cytoskeleton interface. *Genes and Development* 2018;**32**:1201–14. doi:10.1101/gad.317354.118

McClatchey AI, Fehon RG. Merlin and the ERM proteins - regulators of receptor distribution and signaling at the cell cortex. Trends in Cell Biology. 2009;**19**:198–206. doi:10.1016/j.tcb.2009.02.006

Johnson KC, Kissil JL, Fry JL, *et al.* Cellular transformation by a FERM domain mutant of the Nf2 tumor suppressor gene. *Oncogene* 2002;**21**:5990–7. doi:10.1038/sj.onc.1205693

Lallemand D, Curto M, Saotome I, *et al.* NF2 deficiency promotes tumorigenesis and metastasis by destabilizing adherens junctions. *Genes and Development* 2003;**17**:1090–100. doi:10.1101/gad.1054603

Castellanos E, Gel B, Rosas I, *et al.* A comprehensive custom panel design for routine hereditary cancer testing: Preserving control, improving diagnostics and revealing a complex variation landscape. *Scientific Reports* 2017;**7**. doi:10.1038/srep39348

FIGURE LEGENDS

Figure 1. Experimental approach to test antisense therapy for splicing *NF2* variants. In blue, expected outcome after PMO treatment. **(A)** Experimental design for splicing variants that induce exon skipping and result in a truncating protein. **(B)** Experimental design for variants inducing the inclusion of intronic regions as coding sequences, resulting in alterations in the gene reading frame that finally generate a truncating protein. In both approaches, the PMO is expected to mask the aberrant splicing signal and rescue the NF2 phenotype *in vitro*.

Figure 2. (A) Design of the PMO specific for each variant. For the second design (v2), Canonical Splice Sites (CSS), Branch Points (BR), Acceptor and Donor Sites (AS and DS), Exonic Splicing Enhancer sequences (ESE) and Exonic Splicing Silencer (ESS) sequences indicated were taken into account and excluded, when possible, for the sequence design in order to avoid masking the correct splicing signals. The exonic part of the sequence is shown in a gray box and the unlabeled part constitute intronic regions. Tested genetic variants are highlighted in red. **(B) The use of variant-specific PMO did not allow to correct the aberrant splicing signaling.** The effect of the PMOs is shown at cDNA level through dose response and a time course experiments. As, the same effect was observed in both of the PMO designs (v1 and v2), only results from the second design (v2) are shown. Merlins' forms according to the expected molecular weight are indicated next to the bands in the agarose gel. PMO_Ctrl was used as an experiment control. Fib_Ctrl stands for fibroblasts without NF2 mutation, ES for Exon Skipping and UT for Untreated.

Figure 3. Experimental approach to test antisense therapy for truncating *NF2* variants. In blue, expected outcome after PMO treatment**.** PMOs are designed at the donor and acceptor sites of an in-frame exon to induce skipping of exons harboring truncating variants. The generated in-frame deletion is expected to ameliorate the phenotype *in vitro*.

Figure 4. Merlin levels after inducing skipping of in-frame exons harboring truncating variants. Merlin Western Blot after exon-specific PMOs treatment showed a decrease in Merlin total

levels and no presence of exon-less, shorter forms of Merlin-e4 (A) and Merlin-e8 (B). Merlin-e11 was detected and a slight increase was determined in total Merlin levels (C). Increases or decreases in Merlin levels are considered with respect to the untreated sample. Fibroblast control, with no NF2 mutation, are used as experimental control. UT stands for Untreated. NPE stands for Normalized Protein Expression. 0µM are samples treated with Endo-Porter.

Figure 5. E11 PMO treatment of *NF2*$^{+/-}$ fibroblasts induced improvement in actin cytoskeleton organization. Actin immunocytochemistry analysis revealed cytoskeletal abnormalities in patient's derived fibroblasts. Patient_ES11_1 (A) showed a decrease in membrane ruffle formation and Patient_ES11_2 (B) responded to the treatment showing improved culture organization. DAPI was used to stain cell nuclei and Phalloidin is shown in red. Scale bar: 75µm.

Figure 6. Expression of Merlin-e11 could allow a reduction in the proliferation capacity of patient's fibroblasts. Quantification of EdU-positive cells (percentage over DAPI-positive nuclei). A significant decrease was detected in the proliferation rate of treated fibroblasts when compared to Endo-Porter for Patient_ES11_1 (A). Patient_ES11_2 (B) showed the same tendency. (C) Representation of the mean values of both patients. Mean±SD is represented in the bar, representing three independent experiments. *$p < 0.05$, **$p < 0.01$ denotes statistical significance using unpaired t-test. See also Figure S5.

TABLES

Table 1. NF2 splicing loss-of-function PMO design						
Patient's code	c.	g.	p.	Variant Effect	PMO_Design 1	PMO_Design 2
Patient_Spl_Ctrl (Experimental Control) (PMO_Ctrl)	c.1446_1477ins	g.74409T>A	p.Pro482Profs*39	Insertion of a cryptic exon of 167pb between exons 13 and 14	CATCCCTCAAATCTCTTACCGTTCT*	
Patient_Spl_1	c.1736A>G	g.83045A>G	p.Lys525Asnfs*19	ES of exon 15	**Spl1_PMO_v1**: AGAAAGAGACCCTGGGTACCTTTTT	**Spl1_PMO_v2**: CCTTTTAATGGTATTGTGCTTGCTG
Patient_Spl_2	c.810+1dupG	g.62785dupG	p.Phe271_295del	ES of exon 8	**Spl2_PMO_v1**: GCAGTACACACATGTCCTACCCTCC	**Spl2_PMO_v2**: CCCTCCTTGTCACTGTACGAGATGT
Patient_Spl_3	c.241-9A>G	g.35526A>G	p.Val81Phefs*44	New acceptor site. Inclusion of 8 intronic nucleotides as coding sequence	**Spl3_PMO_v1**: AGTACCTGCAGAACTGCAGAGCAAA	**Spl3_PMO_v2**: GCAGAACTGCAGAGCAAAAGACAAA
Patient_Spl_4	c.241-13T>A	g.35522T>A	p.Val81Phefs*44	New acceptor site. Inclusion of 10 intronic nucleotides as coding sequence	**Spl4_PMO_v1**: AGTACCTGCAGAATTGCTGAGCAAA	**Spl4_PMO_v2**: GCAGAATTGCTGAGCAAAAGACAAA

*Exon Skipping (ES). * Published in Castellanos et al. 2013*

Table 2. Exon-specific PMOs design to target NF2 truncating variants

Exon	Target *NF2* variant	Target region	PMOs designed	PMO_Design 1	PMO_Design 2
4 (n=2)	Any truncating at exon 4	Exon 4_5'	PMO Pair	**PMO_5'ES4**: ACCTGAAAGGAGCAACAAGGGAGAC	
		Exon 4_3'		**PMO_3'ES4**: TTTCTTCTTTGAGCCTACCTTGGCC	
8 (n=5)	Any truncating at exon 8	Exon 8_5'	PMO Pair	**PMO_5'ES8**: TTTATTCTGTGGATCCAATAAGAAC	
		Exon 8_3'		**PMO_3'ES8**: TGCAGTACACACATGTCCTACCTCC	
11 (n=2)	Any truncating at exon 11	Exon 11_5'	PMO Pair	**PMO_5'ES11_v1**: CGCTCCATCTGCGAGGGGTGAAGAA	**PMO_5'ES11_v2**: CGCTCCATCTGCGAGGGGTGAAGAA
		Exon 11_3'		**PMO_3'ES11_v1**: CCTCAGAAATCACCAGTGCTTCGTT	**PMO_3'ES11_v2**: CCTCAGAAATCACCAGTGCTTCGTT

n indicates the number of patients derived fibroblasts in which PMOs is tested

A. Pathogenic variants inducing exon skipping

Exon skipping

Partially defective protein

Normal protein
Rescue *NF2* phenotype

B. Pathogenic variants inducing inclusion of intronic regions as coding sequences

No protein synthesis
Patient's severe phenotype

Normal protein
Rescue *NF2* phenotype

A

Patient Spl1

Spl1_PMO_v1
Spl1_PMO_v2
GGCAGCAGCAAGCACAATACCATTAAAAGGGTACCCAGGGTCTCTTTCTTTG
Exon 15
c.1736A>G
CSS
BP
AS
AS
AS
ESE
ESE
ESS

Patient Spl2

Spl2_PMO_v1
Spl2_PMO_v2
AAACATCTCGTACAGTGACAAGGAG GGTAGGACATGTGTGTACTGCAGATGGGT
Exon 8
c.810+1dupG
CSS
AS   AS
AS
ESE   ESE
ESS   ESS

Patients Spl3 and Spl4

c.241-9A>G
Spl3_PMO_v1
Spl3_PMO_v2
TAGTGTGTTTGTCTTTTGCTCTGCAGTTCTGCAG GTACTGGATCATGAT
Exon 3
CSS
c.241-13T>A
Spl4_PMO_v2
Spl4_PMO_v1
DS   DS
DS
ESE   ESE
ESS   ESS

B

B.I. Patient Spl1

Dose Response

0µM   5µM   10µM   20µM

Exons 14, 15, 17 (Isoform I, WT)
Heteroduplex (14,15,17 and 14,17)

Exons 14, 16 (Isoform II, ES15)
Heteroduplex (14,6 and 14, 17)
Exons 14, 17 (Isoform I, ES15)

Time Course (20 µM)

0h   24h   48h   72h   FibCtrl Treated

B.II. Patient Spl2

Dose Response

0 µM   5µM   10µM   20µM

WT
Exon Skipping 8

Time Course (20 µM)

0h   24h   48h   72h   FibCtrl Treated

B.III. Patient Spl3

Dose Response

0µM   5µM   10µM   20µM

WT
Heteroduplex WT and ES
of exons 2 and 3
ES of exons 2 and 3

Time Course (20 µM)

0h   24h   48h   72h   FibCtrl Treated

B.IV. Patient Spl4

Dose Response

0µ   5µM   10µM   20µM

WT
Heteroduplex WT and ES
of exons 2 and 3
ES of exons 2 and 3

Time Course (20µM)

0h   24h   48h   72h   FibCtrl Treated

No protein synthesis
Patient's severe phenotype

PMO
treatment

In frame exon skipping

Partially defective protein
Patient's mild phenotype

A. Truncating mutation in exon 4: inducing skipping of exon 4 (ES4)

UT | Endo-Porter 24h | 20uM 24h | Endo-Porter 48h | 20uM 48h | Endo-Porter 72h | 20uM 72h | PMO Ctrl | Fib_Ctrl

B. Truncating mutation in exon 8: inducing skipping of exon 8 (ES8)

UT | Endo-Porter 24h | 20uM 24h | Endo-Porter 48h | 20uM 48h | Endo-Porter 72h | 20uM 72h | PMO Ctrl | Fib_Ctrl

C. Truncating mutation in exon 11: inducing skipping of exon 11 (ES11)

UT | Endo-Porter 24h | 40uM 24h | Endo-Porter 48h | 40uM 48h | Endo-Porter 72h | 40uM 72h | PMO Ctrl | Fib_Ctrl

A. Patient_ES11_1

B. Patient_ES11_2

C. Merlin-e11