Antisense oligonucleotides targeting exon 11 are able to partially rescue the
Neurofibromatosis Type 2 phenotype *in vitro*.

N.Catasús[1], I.Rosas[1,2], S.Bonache[1,2], A.Negro[1,2], M.Torres-Martin[1,2], A.Plana[3],
H.Salvador[4], E.Serra[5], I.Blanco[1,2] and E.Castellanos[1,2] on behalf of the NF2 Spanish
National Reference Centre HUGTP-ICO-IGTP

1) Clinical Genomics Research Group, Germans Trias i Pujol Research Institute (IGTP);
Can Ruti Campus, Badalona, Barcelona, 08916; Spain

2) Clinical Genetics Department, Germans Trias i Pujol University Hospital (HUGTiP),
Can Ruti Campus, Badalona, Barcelona, 08916; Spain

3) Dermatology Department, Germans Trias i Pujol University Hospital (HUGTiP), Can
Ruti Campus, Badalona, Barcelona, 08916; Spain

4) Pediatric Oncology Unit, Hospital Sant Joan de Déu, Esplugues, Barcelona, Spain

5) Hereditary Cancer Group, Germans Trias i Pujol Research Institute (IGTP-PMPPC);
Can Ruti Campus, Badalona, Barcelona, 08916; Spain

Correspondence should be addressed to Dr. Elisabeth Castellanos
(ecastellanos@igtp.cat)

Corresponding author:

Dr. Elisabeth Castellanos

Clinical Genomics Unit - Genetics Service

Laboratori Clinic Metropolitana Nord (LCMN)

Hospital Universitari Germans Trias i Pujol

Crta. Canyet, s/n. Badalona, Barcelona, Spain

Email: ecastellanos@igtp.cat

Abstract

Neurofibromatosis type 2 (NF2) is an autosomal dominant condition caused by loss of

function variants in the *NF2* gene, which codes for the protein Merlin, and characterized

by the development of multiple tumours of the nervous system. The clinical

presentation of the disease is variable and related to the type of the inherited germline

variant. Here, we tested if PMOs could be used to correct the splice signalling caused by

variants at +/- 13 within the intron-exon boundary region. Here we show that the PMOs

designed for these variants do not constitute a therapeutic approach. Furthermore, we

evaluated the use of phosphorodiamidate morpholino oligomers (PMOs) to reduce the

severity of the effects of *NF2* truncating variants with the aim of generating milder

hypomorphic isoforms *in vitro* through the induction of the in-frame deletion of the

exon-carrying variant. We were able to specifically induce the skipping of exons 4, 8

and 11 maintaining the *NF2* gene reading frame at cDNA level. Only the skipping of

exon 11 produced a hypomorphic Merlin (Merlin-e11), able to partially rescue the

observed phenotype in primary fibroblast cultures from NF2 patients, being

encouraging for the treatment of patients harbouring truncating variants located in exon

11.

INTRODUCTION

Neurofibromatosis type 2 (NF2) (OMIM 101000) is an autosomal dominant (AD)

disease caused by loss-of-function (LOF) variants in the *NF2* tumour suppressor gene

(22q12.2) [1,2]. The reported incidence is between 1 in 28,000-40,000[3] and the hallmark

of the disease is the presence of bilateral vestibular schwannomas (BVS) which

clinically present with hearing loss, tinnitus and imbalance[4]. Common features of NF2

include cranial, spinal, peripheral nerve, and intradermal schwannomas; cranial and

spinal meningiomas; and intrinsic central nervous system (CNS) tumours, usually spinal ependymomas[5,6]. Despite their benign nature, tumours are associated with high morbidity due to their multiplicity and anatomical location, which cause pain and nerve dysfunction leading to a reduced quality of life and life expectancy[4].

There are limited therapies available for these patients and radiotherapy and surgery for tumour resection are the classical treatments[4]. Currently, medical therapy is also an option since Bevacizumab, a monoclonal antibody targeting the Vascular Endothelial Growth Factor (VEGF), has been demonstrated to be an effective treatment for vestibular schwannomas (VS) inducing the reduction of tumour growth and an improvement in hearing in adult NF2 patients[6,7]. However, Bevacizumab does not seem to be equally effective in all NF2 patients and, in addition, it has no effect on other types of tumours arising from the disease , such as meningiomas[8]. Efforts have been made to develop other therapies for NF2-related tumours, such as Losartan (antihypertensive), Brigatinib and Lapatinib (tyrosine kinase inhibitors) or AR-42 (Histone deacetylase (HDAC) inhibitor)[9–13], but there still remains a need to identify clinically relevant therapeutic targets to treat NF2 patients.

The *NF2* gene consists of 17 exons[14,15] with a wide mutational spectrum. It is estimated that 85-90% of the pathogenic variants in *NF2* are germline point mutations and 10-15% are large deletions, distributed throughout the entire gene[16,17]. *NF2* encodes for Merlin (Moesin-Ezrin-Radixin-Like protein), a 595aa scaffold protein involved in membrane protein organization, regulation of cell-cell adhesion and with a regulatory role in the cytoskeleton architecture[18,19]. Merlin is comprised of three structural regions: a N-terminal FERM domain which is subdivided into three globular domains (F1, F2 and F3), an α-helical coiled-coil domain and a C-terminal hydrophilic tail[20].

The genotype-phenotype correlation of NF2 has long been studied[21–29]. For NF2, it is

well reported that truncating variants and large deletions, which account for around 30%

of cases, are associated with severe manifestations of the disease, whereas missense or

in-frame variants present milder forms of NF2. Data reveal that individuals with the

latter variants show a late disease onset and are diagnosed later compared to individuals

with truncating mutations, as well as, a reduced risk of developing peripheral nerve

tumours, spinal tumours and meningiomas. Thus, being these genetics variants

associated with milder disease burden[21,23,27,30,31]. Besides, about 25% of the point

mutations affect the correct splicing of the *NF2* gene, and this mechanism is associated

with a variable severity of the disease[26,28,30]. NF2 is also estimated to have a high

presence of mosaicism, reported to affect at least 30% and up to 50% of *de novo*

patients, which often challenges the molecular diagnosis of the disease[32] NF2

mosaicism is associated to mild clinical presentations, although significant variability

has been reported[28,32]. Thus, the well-known association between the germline genetic

variant type and disease severity may constitute an opportunity to develop personalized

gene therapies based on the specific variant of the *NF2* gene.

Antisense gene therapy has been successfully used to regulate gene expression in some

pathologies. Most of the target diseases are inherited in an autosomal recessive manner,

such as Spinal Muscular Atrophy (SMA)[33], homozygous familial hypercholesterolemia

(HoFH)[34], or Duchenne muscular dystrophy (DMD)[35] the latter two have been approved

by the FDA and are amongst others currently in the advanced stages of clinical

trials[36,37]. Furthermore, there are also examples for AD diseases, such as Retinitis

pigmentosa (RP)[38], amyotrophic lateral sclerosis[39] or Huntington's disease[40]. In all

cases, antisense oligonucleotides (ASOs) have been used. These are single-stranded

molecules that act at RNA level by specifically correcting or modifying the expression

of the target protein to a less deleterious form[41,42]. ASOs have been therapeutically used

to reduce protein production whether via RNase H cleavage or mRNA-ASO complex to

block mRNA-ribosome interaction. It is also possible to modulate the splice signalling

and among the different types of ASOs, Phosphorodiamidate Morpholino Oligomers

(PMOs) have highly desirable molecular properties that make them suitable for this last

approach. PMOs are DNA-analogue molecules that comprise a nucleic acid base, a

methylenemorpholino ring and phosphodiester inter-subunit bonds are replaced by

phosphorodiamidate linkages, all of which produce a structure that provides high-

targeting specificity, stability, low toxicity and resistance to nucleases, assuring the

maintenance of a good long-term activity within the cell[43].

In this study we propose an approach to test *in vitro* antisense therapy for NF2. This

consists of forcing the skipping of exons harbouring a frameshift or nonsense variant

while preserving the gene reading frame, being potentially applicable to 9 of the 15 *NF2*

exons in which truncating variants have been reported. Thus, taking into consideration

the NF2 genotype-phenotype association, frameshift or nonsense variants that generate

a more severe phenotype could be altered at RNA level and potentially generate less

deleterious protein forms, resulting in a mitigation of phenotype severity. In addition,

we tested the feasibility of modulating and correcting the aberrant splicing of the *NF2*

gene originating from different point-mutations located near canonical splice sites.

RESULTS

Antisense therapy strategies for *NF2*

The first approach of this study aimed to test if PMOs could be used to specifically

force the skipping of in-frame exons carrying a truncating variant, thus maintaining the

NF2 gene reading frame, and test if the generated Merlin protein could act as an

hypomorphic form, still preserving partial function and to some extent, able to rescue an

in vitro phenotype. In this way, taking into account that nonsense or frameshift variants

are associated to more severe phenotypes than in-frame variants, based on genotype-

phenotype correlations, these truncating variants could be modified to less deleterious

forms (Figure 1A), with the need to assess whether the effect of the exon skipping on

both alleles was more beneficial than harbouring a truncating variant in heterozygosis.

A second strategy was evaluated for *NF2* splicing variants. This experimental approach

consisted of designing specific PMOs for a particular splicing variant in order to mask

the aberrant splice signalling triggered by the pathogenic variant and promote the

spliceosome to properly recognize the correct splice site without altering the wild type

(WT) allele. This could be used to target splicing variants that induce the skipping of an

exon (Figure 1B) or for those that induce inclusion of intronic regions as coding

sequences, as previously tested[44] (Figure 1C).

Evaluating the use of PMOs for truncating variants affecting the *NF2* gene

We first investigated *NF2* exons that could be appropriate candidates for

generating an hypomorphic Merlin isoform after forcing its skipping, while preserving

the gene reading frame. The evaluation of the *NF2* transcript (NM_00268.3) identified 9

out of 15 exons that could be skipped while maintaining the translational reading frame.

In addition, for the remaining out-of-frame exons it would be possible to induce

skipping of two consecutive exons (6&7, 12&13 and 14&15) (Figure S1). All these

twelve combinations were prioritized to be *in silico* analysed. We also assessed exon

length, since longer exons encode for a more significant proportion of the protein and

thus, could cause greater impact on its structure and stability and, as a consequence,

functionality. Most *NF2* exons were in the range of normality without any exon overly

long (>500bp). We indicated all unique nonsense and frameshift variants described in

the publicly available dataset LOVD (Leiden Open Variation Database) and also all

truncating and splicing variants identified in our cohort. As previously described, no

hotspots were detected and pathogenic variants were spread over the whole gene[16,17],

although several were described in the FERM domain[16,20] in LOVD database. Next, we

annotated Merlin protein post-translational modifications (PMTs) and the theoretical

effect of skipping for each exon (or consecutive exons) in the resulting Merlin protein

by *PredictProtein*[45,46]. Predicted features are shown in Figure S1. Seven candidate

exons showed a high conservation score (>6.5), three were predicted to induce a high

change of surface area, while five out twelve exons or combination of them showed

several phosphorylation sites, which could be relevant for Merlin regulation.

Considering all the different *in silico* indicators, exons 5 and 11 might be the most

suitable candidates since they showed less conservation score, did not present any

phosphorylation site and the predicted impact over the protein structure would be

relatively low. Nonetheless, besides pairs of exons 12&13 and 14&15, the rest of the in-

frame exons presented *in silico* metrics relatively similar among them, and therefore,

there was no strong evidence for discard any of the exons for further *in vitro* analysis. In

the light of this, we tested the effect on Merlin after forcing exon-skipping on all the

available patient's primary fibroblasts with a truncating variant, corresponding to exons

4, 8 and 11.

A pair of PMOs were designed to be complementary to both 5'- and 3'- intron-exon

boundaries to induce skipping of exons 4, 8 or 11 of the *NF2* gene (Exon-Specific

PMOs) and the induction of the exon skipping was first assessed in fibroblasts from

healthy donors (NF2$^{(+/+)}$) (Table 1). These were delivered to primary fibroblasts from

healthy donors (NF2$^{(+/+)}$) through an endocytosis-mediated process with the use of the

Endo-Porter. The effect of the treatment was considered in relation to the cells treated

with Endo-Porter to determine exclusively the effect of the PMO treatment. Dose

response and time course studies confirmed the efficiency of the PMO treatment at

RNA level, showing full skipping of exons 4 and 8 at 20µM at 24h, 48h and 72h

(Figure 2A and Figure S2). For exon 11, two different designs were tested (Table 1;

PMO_ES11_v1 and PMO_ES11_v2): while the first (PMO_ES11_v1) failed to induce

skipping (data not shown), for the second design (PMO_ES11_v2) the maximum

skipping effect observed was at the highest tested dose (40µM), achieving more than

50% of the exon-less form after 72h of treatment (Figure 2A and Figure S2).

Next, we analysed whether the skipping of both the *NF2* alleles, the one

harbouring a truncating variant and the WT allele, was less deleterious than harbouring

a non-functional allele. First, we studied the capacity of restoring Merlin levels *in vitro*

after the PMO treatment. Merlin western blot showed progressive loss of protein signal

when targeting exons 4 (Merlin-e4) or 8 (Merlin-e8), studied at 24h, 48h and 72h time

points in two and five patient samples, respectively (Figure 2B). In order to verify that

the induced skipping of exons 4 or 8 caused a reduction of Merlin levels rather than a

lack of immunoreactivity; the exon-less *NF2* cDNAs were cloned to a N-terminal

FLAG® Tag Expression Vector and transfected to HeLa cells. Flag-Merlin Western

Blot showed that there was little or no expression of Merlin-e4 or Merlin-e8, indicating

that these new Merlin forms could not be synthesized or were degraded shortly after

synthesis (Figure S3).

Conversely, skipping of exon 11 generated a shorter isoform of Merlin (Merlin-

e11), and quantification of these results determine a slight increase in the total levels of

Merlin (WT Merlin and Merlin-e11) (Figure 2B and Figure S4). These results indicated

that a potential hypomorphic Merlin-e11 could result from the skipping of this exon.

E11 PMO treatment of *NF2*$^{(+/-)}$ fibroblasts partially rescued the phenotype *in vitro*

Functionality of Merlin-e11 was studied in primary fibroblasts harbouring a

heterozygous truncating variant in exon 11: one from an adult patient (Patient_ES11_1)

and another from a paediatric patient (Patient_ES11_2) (Table S1). Given the

complexity of determining differences in severity in the phenotype from studying the

signalling pathways in which Merlin is involved [29], physiological read outs were

assessed in patients derived fibroblasts (*NF2*$^{(+/-)}$). Due to the role of Merlin in

cytoskeletal organization [38,39], Merlin-e11's functional activity was first tested

through actin immunofluorescence. Phalloidin staining revealed an altered phenotype in

both primary fibroblast samples, but it was slightly different in each patient tested:

patient_ES11_1 showed prominent membrane ruffles, while patient_ES11_2 showed

alterations in cytoskeletal abnormalities and a tendency to growth as aggregates, when

compared to primary *NF2*$^{((+/+))}$ fibroblasts (Figure 3), and quite similar to the phenotype

observed in other primary *NF2*$^{(+/-)}$ fibroblasts (Figure S5). After PMO treatment an

improvement of the organizational capacity of the cytoskeleton was observed,

membrane ruffles were decreased in patient_ES11_1, and in patient_ES11_2 we

observed an improvement in the cell-cell contact organization (Figure 3). Specifically,

paediatric derived NF2$^{(+/-)}$ fibroblasts recovered part of the cell-cell contact inhibition

and recover the capacity of growth as monolayer after the induction of Merlin-e11.

In addition, proliferation rates were tested after 72h of treatment by flow cytometry.

Patient_ES11_1 showed a statistically significant depletion of proliferation when treated

with PMOs (65.6%, $p<0.05$) (Figure 4A) while patient_ES11_2 showed a less

pronounced decrease (25.1%) (Figure 4B). Considering both samples, primary

fibroblasts with a truncating variant in exon 11 and expressing Merlin-e11 by PMO

treatment showed a 51% reduction in proliferation when compared with the vehicle

(Endo-Porter, p<0.01) and 39.7% compared to treatment control (PMO with no effect

over the *NF2* gene) (p<0.01), on average (Figure 4C, Figure S6). In order to determine

absence of potential toxic effects, a viability test was performed, revealing that PMO

treatment did not induce significant differences in cell viability (Figure S7).

Antisense therapy for *NF2* splicing variants

This additional strategy intended to use PMOs, rather than to induce skipping, to mask

specifically splicing variants and thus recover the correct splice signal. Two different

PMOs were designed for each variant. The first design was a 25-mer variant-specific

facilitated by *Gene Tools* according to their guidelines for intron and exon binding

subunits and self-complementary moieties (data not shown). We made a second design

using splice site predictors to study the variant surrounding sequence (acceptor splicing

sites, exonic splicing enhancers (ESEs) or exonic splicing silencers (ESSs)) (based on

Human Splice Finder) in order to avoid the induction of possible new splicing signals

due to the effect of the PMO. In addition, the design avoided masking the canonical

splicing GT-AG sites and branch points (YNYYRAY) when possible (Table 2, Figure

5A). We tested the PMOs' efficacy in primary fibroblasts from NF2 patients harbouring

heterozygous germline variants (NF2$^{(+/-)}$) located near the canonical splice site of exons

3, 8 and 15 (Table S1). A dose-response study was performed to test the effect on

splicing of different concentrations of PMOs after 24h of treatment, and the efficiency

was assessed at cDNA level by RT-PCR. The effect of PMOs on the splicing of *NF2*

was analysed by electrophoresis and confirmed by Sanger sequencing, in which the

maximum observed effect was at the highest tested dose (20µM) (Figure 5B: Dose

Response; Figure S8). A time course was carried out treating fibroblasts with 20µM

PMO during 24h, 48h and 72h. PMO treatment in Patient_Spl_1 and Patient_Spl_2

increased the skipping of exons 15 and 8 respectively, in contrast to the desired effect of

inducing the inclusion of skipped exon without altering the transcription of the WT

allele (Figure 5B: Time Course, B.I and B.II; Figure S8). Similarly, the treatment of

PMOs of patients Patient_Spl_3 and Patient_Spl_4, both with nearby variants near the

canonical splice site of exon 2, revealed that the variant-specific PMO caused the

skipping of both exons 2 and 3 of the *NF2* gene instead of correcting the aberrant *NF2*

splicing due to germline variants (Figure 5B: Time Course, B.III and B.IV; Figure S8).

Their effect was confirmed by Sanger sequencing (Figure S9). $NF2^{(+/+)}$ fibroblasts from

healthy donors were also treated with each designed variant-specific PMO and they

showed the same effect as $NF2^{(+/-)}$ fibroblasts derived from patients (Figure 5B and

Figure S8). The effect of the treatment on Merlin was assayed by Western Blot and no

improvement in Merlin levels was detected at the studied time points (Figure S10).

DISCUSSION

Neurofibromatosis type 2 (NF2) is a multisystem genetic disorder for which the

development of effective therapeutic options with no adverse consequences is needed.

The clinical presentation of the disease is variable and related to the type of germline

variant inherited in the *NF2* gene[4]. In this context, the development of personalized

therapies based on the type of *NF2* variant constitutes an opportunity. In this study, we

tested the use of PMOs as antisense therapy to reduce the severity of the effects of *NF2*

germline truncating and splicing variants by changing them to milder hypomorphic

forms *in vitro* by two different approaches.

In the in-frame exon skipping approach, we targeted nonsense and frameshift variants

trying to benefit from the fact that these variants are associated with more severe

phenotypes in comparison to in-frame germinal variants in *NF2*[27,29,47]. Thus, we

proposed inducing skipping of the exon carrying the truncating variant with the

hypothesis of generating potential homozygous hypomorphic forms of Merlin, to

ameliorate the pathogenicity of the heterozygous truncating variant *in vitro*. We did not

consider mosaic patients in this approach since their phenotype could depend on the

type of variant and the percentage of affected cells[32].

Firstly, we evaluated the theoretical effect of skipping any in-frame exon, and although

in silico studies did not show any outstanding candidate, the predicted effect on exons 5

and 11 was encouraging. Therefore, we decided to test all exons for which samples were

available. The skipping of exons 4 or 8 harbouring *NF2* LOF variants was not able to

generate an exon-less Merlin, because Merlin-e4 and Merlin-e8 were degraded. Exons 4

and 8 of the *NF2* gene code for the subdomains B and C respectively of the FERM

domain, where the majority of reported pathogenic variants occur[16,20]. The observed

results could be due to the fact that the lack of an exon coding for the FERM domain

would cause an unbearable aberrant folding of the protein and its synthesis would be

prevented or it would be forcibly degraded, although these effects were not predicted *in*

silico. In addition, the FERM domain is responsible for the intramolecular interaction

with the C-terminal domain, which results in Merlin being in a closed and active

conformation[48–50] to induce the most potential for anti-mitogenic signalling. Thus, if this

folding fails, it is possible that the resulting protein would not be able to execute the

tumour suppressor activity.

When targeting exon 11 of the *NF2* gene, an exon-less Merlin was expressed (Merlin-

e11). Merlin, as a member of the ERM protein family, is involved in the link between

actin cytoskeleton and adherent junctions with a relevant role in dynamic cytoskeleton

remodelling, such as in membrane ruffling or in the formation of actin micro-spikes[51–

53]. Consistent with other studies which described that Merlin loss leads to a dramatic

increase of membrane ruffles and actin cytoskeletal disoganization[44,54], we found that

$NF2^{(+/-)}$ primary fibroblasts tested in this study also showed actin cytoskeleton

abnormalities. In addition, here we observed a decrease in membrane ruffle formation in

response to the expression of Merlin-e11 that could be associated to a partial rescue of

the phenotype and an improvement in the cytoskeleton organization after treatment,

which could be due to a recovery of the Merlin function as a scaffold protein regulating

the linkage between membrane proteins and the cortical cytoskeleton[44,54,55]. Consistent

with the role as a tumour suppressor protein, Merlin-deficient cells loose contact

inhibition[48,52,56] and although much remains to be discovered about the adhesive

signalling pathways involved, in this study we showed that the induction of Merlin-e11

by PMO treatment contributed to the recovery of cell-cell contact inhibition of

proliferation in fibroblasts $NF2^{(+/-)}$ derived from NF2 patients harbouring truncating

mutations in exon 11. The tested dose in this study are higher than FDA approved

PMOs for other inherited diseases[36,57,58]. Thus, redesigns and modifications of the tested

antisense molecule could be performed to improve its efficacy and reduce the required

dose, in addition to further pharmacodynamics and pharmacokinetics studies to

determine the appropriate range doses and the associated toxicity and effectiveness.

Furthermore, treatment effect should be tested in different cell types, especially in

schwannoma and meningioma originating cells, as well as *in vivo* studies to determine if

the treatment could prevent tumour formation or could reduce tumour burden, when

administered systemically before considering the applicability of these molecules in

patients

Thus, this study constitutes a proof of concept of a potential medical approach for these

variants, although there this is still much to be elucidated about the effect of the PMOs

treatment and its therapeutically approach. Further data is required to assess whether

this approach could delay the appearance of tumors, cause patients a reduced tumor

burden or slower their growth.

A second strategy was directed to splicing variants with the aim of concealing the

erroneous splice signal caused by the pathogenic variant and restoring levels of Merlin,

in a similar way to the approach that our group used for a *NF2* deep intronic variant[44] or

to induce exon retention by blocking an homozygous single nucleotide variant located 6

nucleotides within exon 7 of *SMN2*[59]. In this study, it has not been possible to design

PMOs that permit specifically the correction of aberrant splicing signalling and

furthermore, it was not possible to achieve enough specificity to target the mutated

allele as the same effect was observed in both patient-derived fibroblasts and control

fibroblasts. Based on our results, we hypothesize that when the splicing variant is within

+/- 13 nucleotides of the acceptor or donor splice site, the PMO could allosterically

interfere with the branch point and the U2 snRNA incorporation into the spliceosome[60].

The spatiotemporal localization of the two alleles during splicing could prevent the

PMO from specifically targeting the mutated allele and thus it would interfere in the cis-

regulation of RNA elements of both alleles equally, resulting in an increased pathogenic

effect[61]. Since the approach requires the PMO to be specific for the allele harbouring the

pathogenic variant, there is little room for variation in the PMO design for the variants

tested in this study.

As far as we know, there are no studies in the literature that have been able to repair the

LOF variant effect and induce exon retention through the use of ASOs for a splicing

variants located in the canonical splice regions in genes causing autosomal dominant

diseases. The most described effect of these molecules targeting intron/exon boundary is exon skipping. It has only been possible through CRISPR/Cas technology by the induction of single nucleotide substitutions[62,63] Allele-specific approaches have been successfully tested for intronic variants through splice modulation ASOs that mask the erroneous signalling and given their intronic location and distance to splicing regulatory regions, proper protein synthesis can be recovered[44,64–67]. Few works have applied ESSENCE (exon-specific splicing enhancement by small chimeric effectors) method to rescue disease-associated exon skipping and modulate alternative splicing[33,68,69]. Furthermore, for autosomal recessive diseases, such as SMA or DMD[70,71], or for autosomal dominant diseases with a dominant negative effect, such as Frontotemporal dementia with parkinsonism-17 (FTDP-17) or RP, allele specific ASOs have been used to induce degradation of the disease-causing allele via RNAse H activity and for this purpose, the native DNA backbone is not modified[60,72–74]. Some evidences suggest that Merlin could act in a negative dominant manner through its self-dimerization through FERM-FERM interactions[75–77], but this seems to be dependent on patient's variant[78]. In the light of our findings, and if Merlin dominant effect is confirmed, the degradation of the *NF2* mutated allele by the use of RNAse H dependent ASOs could be a possible antisense approach for nonsense, frameshift and splicing pathogenic variants located in the FERM domain, in both in-frame and out-of-frame exons, and then bringing an improvement in the *in vitro* phenotype and opening the door for future gene therapy approaches.

In addition, for those truncating variants that do not exert a dominant-negative effect, the generation of hypomorphic Merlin variants could be beneficial for the patient, and therefore, more variants and hypomorphic Merlin forms should be analysed to understand the contribution of this effect over patient phenotype.

This work is a proof of concept of the use antisense therapy *in vitro* as a personalized

therapy for NF2 patients harbouring germline truncating variants at exon 11 of the *NF2*

gene. Our approach has shown encouraging results when targeting exon 11, although

further characterization assays need to be performed to better characterize the function

of Merlin-e11 before considering it for further *in vivo* or pre-clinical assays.

In addition, further studies will be needed to elucidate whether there are other

mechanisms or designs to apply ASOs as a therapeutic approach for this disease, as in

our experience it has not been possible to recover Merlin levels for variants located at

in-frame exons 4 or 8, or to correct splicing signals *in vitro* in the four variants located

near canonical splice sites tested, although other splicing variants should be assessed to

confirm these results.

MATERIAL AND METHODS

All procedures performed were in accordance with the ethical standards of the IGTP

Institutional Review Board, which approved this study and with the 1964 Helsinki

declaration and its later amendments.

Samples: samples from 14 NF2 patients from the Spanish National Reference Centre

(CSUR) on Phakomatoses were included in the study (Table S1) after providing written

informed consent. Samples included in the study were those harbouring a splicing

variant or a truncating variant in an in-frame exon of NF2. **Controls**: Fibroblasts

$NF2^{(+/+)}$ stand as non-affected cells for experimental control (WT), with no mutation in

the *NF2* gene, obtained from healthy donors.

Genetic *NF2* test: Genetic testing was performed using the customized I2HCP panel[79]

in blood or tissue when available. To detect splicing variants, patients were studied both

at RNA and DNA levels. Variants were validated by Sanger sequencing and analyzed

using CLC-workbench (Qiagen, Hilden, Germany) in all primary cultures established

using (NM_000268.3; NG_009057.1; LRG_511) as reference sequences.

***In silico NF2* analysis:**NF2 exons were evaluated using NM_000268.3 and

NG_009057.1 reference sequences. *NF2* pathogenic variants described in Leiden Open

Variation Database [LOVD] in addition with our genetic variants database identified in

our NF2 cohort[28]. All *in silico* analysis of exons (and combinations of them) that when

skipped, an in-frame Merlin protein is theoretically maintained were based on the

analysis performed by Leier et al 2021[80]. Post Translation Modifications (PTM) were

retrieved from Phosfosite[81]. Only phosphorylation, acetylation and ubiquitylation were

considered as PTMs. Analysis of *in silico* protein modifications by the exon skipping

was performed using the web-server version of PredictProtein

(https://www.PredictProtein.org)[45,46].

Antisense specificity and efficacy: 25-mer specific Phosphorodiamidate Morpholino

Oligomers (PMOs) for four patients with splicing variants, and a pair of PMOs to

induce exon skipping of exons 4, 8, and 11 of the *NF2* gene were designed

(NM_000268.3; NG_009057.1; LRG_511), synthesized and purified by Gene Tools

(Philomath, OR, USA). Endo-Porter (Gene Tools) was used as vehicle to deliver PMOs

into cells at 6μM (manufacturer recommended dose), according to the manufacturer's

instructions. Patient primary fibroblasts carrying *NF2* germline pathogenic variants

were treated with the specific single or paired PMOs. A dose-response and a time

course study were performed to set-up treatment conditions and the effect of PMOs was

assessed by RT-PCR and Sanger sequencing. Levels of Merlin after PMO treatment

were analyzed by Western Blot. For all experiments, the correct effect of the treatment

at RNA level was confirmed prior to Western Blot analysis.

Analysis of Merlin-e11 phenotype recovery: an actin immunocytochemistry assay was performed to study the cytoskeletal organization after PMO treatment using Phalloidin (Molecular Probes, Invitrogen). Click-iT™ EdU Alexa Fluor™ 488 Flow Cytometry Assay Kit (Molecular Probes, Invitrogen) was used, according to the manufacturer's protocol, to determine cell proliferation. Cell viability was assessed using RealTime-Glo™ MT Cell Viability Assay (Promega) following the manufacturer's instructions.

NF2 **Cloning and expression:** *NF2* gene was cloned using the Gateway® Gene Cloning system (Invitrogen) after treatment to induce the exon-skipping to a FLAG® Tag Expression Vector. The expression vector was transfected into HeLa cells (Lipofectamine LTX with Plus Reagent, Invitrogen) according to the manufacturer's instructions. Flag expression was assessed by Western Blot (Monoclonal ANTI-FLAG® M2 antibody, Sigma-Aldrich).

Statistics: Statistical analysis was carried out using GraphPad Prism software v7. The Shapiro–Wilk test was used to test normality and for multiple group comparisons, a two-tailed unpaired t test was performed. Statistical significance is indicated by $*p<0.05$, $**p<0.01$, and $***p<0.001$. For more information, see Supplementary Materials and Methods.

ACKNOWLEDGEMENTS

We thank the HGTP Clinical Services and staff for their collaboration in generating and collecting patient samples and data and the Hereditary Cancer Group at the IGTP for their help in improving this work. We thank the IGTP Flow Cytometry core facility and its staff for their contribution and technical support. We would like to acknowledge the constant support of the different NF lay associations: Asociación de Afectados de

Neurofibromatosis, Chromo22 and ACNefi. This study has been funded by Instituto de Salud Carlos III through the project PI20/00215 (Co-funded by European Regional Development Fund "A way to make Europe"), Fundació La Marató de TV3 (126/C/2020), the Children's Tumor Foundation (CTF-2019-05-005), the Spanish Association of NF Affected (AANF), Fundación Proyecto Neurofibromatosis, the Catalan NF Association (AcNeFi), and the Government of Catalonia (2017 SGR 496).

AUTHOR CONTRIBUTORS

E.C. conceived the study and wrote the manuscript that was revised, corrected, and improved by all the authors. N.C. performed most of the experimental work, analyzed the data, generated the figures for the paper and contributed to writing the manuscript. I.R, S.B., and A.N. performed experimental work. M.T-M. performed bioinformatics analysis. A.P. and H.S. contributed with sample collection. E.S. and I.B. provided scientific input. All authors approved the final version of the manuscript.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflicts of interest.

KEY WORDS

Neurofibromatosis type 2, antisense therapy, antisense oligomers, genotype-phenotype correlations, Merlin

DATA AVAILABILITY STATEMENT

The authors confirm that the data supporting the findings of this study are available within the article [and/or] its supplementary materials.

REFERENCES

1. Evans, D. G. R., Huson, S. M., Donnai, D., Neary, W., Blair, V., Newton, V. & Strachan, T. A genetic study of type 2 neurofibromatosis in the United Kingdom . II . Guidelines for genetic counselling. 847–852 (1992).

2. Evans, D. G., Howard, E., Giblin, C., Clancy, T., Spencer, H., Huson, S. M. & Lalloo, F. Birth incidence and prevalence of tumor-prone syndromes: Estimates from a UK family genetic register service. *Am. J. Med. Genet. Part A* **152**, 327–332 (2010).

3. Evans, D. G., Bowers, N. L., Tobi, S., Hartley, C., Wallace, A. J., King, A. T., Lloyd, S. K. W., Rutherford, S. A., Hammerbeck-Ward, C., Pathmanaban, O. N., *et al.* Schwannomatosis: A genetic and epidemiological study. *J. Neurol. Neurosurg. Psychiatry* (2018) doi:10.1136/jnnp-2018-318538.

4. Lloyd, S. K. W. & Evans, D. G. R. *Neurofibromatosis type 2 (NF2): diagnosis and management. Peripheral Nerve Disorders* vol. 115 (Elsevier B.V., 2013).

5. Evans, D. G. R. Neurofibromatosis type 2 (NF2): A clinical and molecular review. *Orphanet J. Rare Dis.* **4**, (2009).

6. Forde, C., King, A. T., Rutherford, S. A., Hammerbeck-Ward, C., Lloyd, S. K., Freeman, S. R., Pathmanaban, O. N., Stapleton, E., Thomas, O. M., Laitt, R. D., *et al.* Disease course of neurofibromatosis type 2: A 30-year follow-up study of 353 patients seen at a single institution. *Neuro. Oncol.* **23**, 1113–1124 (2021).

7. Plotkin, S. R., Duda, D. G., Muzikansky, A., Allen, J., Blakeley, J., Rosser, T., Campian, J. L., Wade Clapp, D., Fisher, M. J., Tonsgard, J., *et al.* Multicenter, prospective, phase II and biomarker study of high-dose bevacizumab as induction therapy in patients with neurofibromatosis type 2 and progressive vestibular schwannoma. *J. Clin. Oncol.* **37**, 3446–3454 (2019).

8. Nunes, F. P., Merker, V. L., Jennings, D., Caruso, P. A., di Tomaso, E., Muzikansky, A., Barker, F. G., Stemmer-Rachamimov, A. & Plotkin, S. R. Bevacizumab Treatment for Meningiomas in NF2: A Retrospective Analysis of 15 Patients. *PLoS One* **8**, (2013).

9.   Wu, L., Vasilijic, S., Sun, Y., Chen, J., Landegger, L. D., Zhang, Y., Zhou, W.,
Ren, J., Early, S., Yin, Z., *et al. Losartan prevents tumor-induced hearing loss*
and augments radiation efficacy in NF2 schwannoma rodent models. *Sci. Transl.*
Med vol. 13 https://www.science.org (2021).

10.   Chang, L. S., Oblinger, J. L., Smith, A. E., Ferrer, M., Angus, S. P., Hawley, E.,
Petrilli, A. M., Beauchamp, R. L., Riecken, L. B., Erdin, S., *et al.* Brigatinib
causes tumor shrinkage in both NF2-deficient meningioma and schwannoma
through inhibition of multiple tyrosine kinases but not ALK. *PLoS One* **16**,
(2021).

11.   Karajannis, M. A., Legault, G., Hagiwara, M., Ballas, M. S., Brown, K.,
Nusbaum, A. O., Hochman, T., Goldberg, J. D., Koch, K. M., Golfinos, J. G., *et*
al. Phase II trial of lapatinib in adult and pediatric patients with
neurofibromatosis type 2 and progressive vestibular schwannomas. *Neuro.*
Oncol. **14**, 1163–1170 (2012).

12.   Osorio, D. S., Hu, J., Mitchell, C., Allen, J. C., Stanek, J., Hagiwara, M. &
Karajannis, M. A. Effect of lapatinib on meningioma growth in adults with
neurofibromatosis type 2. *J. Neurooncol.* **139**, 749–755 (2018).

13.   Welling, D. B., Collier, K. A., Burns, S. S., Oblinger, J. L., Shu, E., Miles-
Markley, B. A., Hofmeister, C. C., Makary, M. S., Slone, H. W., Blakeley, J. O.,
et al. Early phase clinical studies of AR-42, a histone deacetylase inhibitor, for
neurofibromatosis type 2-associated vestibular schwannomas and meningiomas.
Laryngoscope Investig. Otolaryngol. **6**, 1008–1019 (2021).

14.   Schmucker, B., Tang, Y. & Kressel, M. Novel alternatively spliced isoforms of
the neurofibromatosis type 2 tumor suppressor are targeted to the nucleus and
cytoplasmic granules. *Hum. Mol. Genet.* **8**, 1561–1570 (1999).

15.   Su, F., Zhou, Z., Su, W., Wang, Z. & Wu, Q. A novel alternative splicing isoform
of NF2 identified in human schwann cells. *Oncol. Lett.* **12**, 977–982 (2016).

16.   Ahronowitz, I., Xin, W., Kiely, R., Sims, K., Maccollin, M. & Nunes, F. P.
Mutational Spectrum of the NF2 Gene: A Meta-Analysis of 12 Years of
Research and Diagnostic Laboratory Findings. **28**, (2007).

17.   Halliday, D., Parry, A. & Evans, D. G. Neurofibromatosis type 2 and related
disorders. *Curr. Opin. Oncol.* **31**, 562–567 (2019).

18.   Asthagiri, A. R., Parry, D. M., Butman, J. A., Kim, H. J., Tsilou, E. T., Zhuang,
Z. & Lonser, R. R. Neurofibromatosis type 2. *Lancet* **373**, 1974–86 (2009).

19.   Chiasson-MacKenzie, C., Morris, Z. S., Baca, Q., Morris, B., Coker, J. K.,
Mirchev, R., Jensen, A. E., Carey, T., Stott, S. L., Golan, D. E., *et al.* NF2/Merlin
mediates contact-dependent inhibition of EGFR mobility and internalization via
cortical actomyosin. *J. Cell Biol.* **211**, 391–405 (2015).

20.   Cooper, J. & Giancotti, F. G. Molecular insights into NF2/Merlin tumor
suppressor function. *FEBS Lett.* **588**, 2743–2752 (2014).

21.   Evans, D. G. R., Trueman, L., Wallace, A., Collins, S. & Strachan, T. Genotype /
phenotype correlations in type 2 neurofibromatosis (NF2): evidence for more
severe disease associated with truncating mutations. 450–455 (1998).

22.   Heineman, T. E., Evans, D. G. R., Campagne, F. & Selesnick, S. H. In silico
analysis of NF2 gene missense mutations in neurofibromatosis type 2: From
genotype to phenotype. *Otol. Neurotol.* **36**, 908–914 (2015).

23.   Sk, S., Shenton, A., Ferner, R., Aj, W. & Sm, H. Further genotype – phenotype
correlations in neurofibromatosis 2. 163–170 (2010) doi:10.1111/j.1399-
0004.2009.01315.x.

24.   Halliday, D., Emmanouil, B., Pretorius, P., Mackeith, S., Painter, S., Tomkins,
H., Evans, D. G. & Parry, A. Genetic Severity Score predicts clinical phenotype
in NF2. 657–664 (2017) doi:10.1136/jmedgenet-2017-104519.

25.   Emmanouil, B., Houston, R., May, A., Ramsden, J. D., Hanemann, C. O.,
Halliday, D., Parry, A. & Mackeith, S. Progression of hearing loss in
neurofibromatosis type 2 according to genetic severity. *Laryngoscope* **129**, 974–
980 (2019).

26.   Halliday, D., Emmanouil, B., Vassallo, G., Lascelles, K., Nicholson, J.,
Chandratre, S., Anand, G., Wasik, M., Pretorius, P., Evans, D. G., *et al.* Trends in

phenotype in the English paediatric neurofibromatosis type 2 cohort stratified by
genetic severity. *Clin. Genet.* **96**, 151–162 (2019).

27. Baser, M. E., Kuramoto, L., Joe, H., Friedman, J. M., Wallace, A. J., Gillespie, J.
E., Ramsden, R. T. & Evans, D. G. R. Genotype-Phenotype Correlations for
Nervous System Tumors in Neurofibromatosis 2: A Population-Based Study. *Am.*
J. Hum. Genet. **75**, 231–239 (2004).

28. Catasús, N., Garcia, B., Galván-Femenía, I., Plana, A., Negro, A., Rosas, I., Ros,
A., Amilibia, E., Becerra, J. L., Hostalot, C., *et al.* Revisiting the UK genetic
severity score for NF2: A proposal for the addition of a functional genetic
component. *J. Med. Genet.* 1–9 (2021) doi:10.1136/jmedgenet-2020-107548.

29. Ruttledge, M. H., Andermann, a a, Phelan, C. M., Claudio, J. O., Han, F. Y.,
Chretien, N., Rangaratnam, S., MacCollin, M., Short, P., Parry, D., *et al.* Type of
mutation in the neurofibromatosis type 2 gene (NF2) frequently determines
severity of disease. *Am. J. Hum. Genet.* **59**, 331–42 (1996).

30. Halliday, D., Emmanouil, B., Pretorius, P., MacKeith, S., Painter, S., Tomkins,
H., Evans, D. G. & Parry, A. Genetic severity score predicts clinical phenotype in
NF2. *J. Med. Genet.* **54**, 657–664 (2017).

31. Teranishi, Y., Miyawaki, S., Nakatomi, H., Ohara, K., Hongo, H., Dofuku, S.,
Okano, A., Takayanagi, S., Ota, T., Yoshimura, J., *et al.* Early prediction of
functional prognosis in neurofibromatosis type 2 patients based on genotype–
phenotype correlation with targeted deep sequencing. *Sci. Rep.* **12**, 1–10 (2022).

32. Gareth Evans, D., Hartley, C. L., Smith, P. T., King, A. T., Bowers, N. L., Tobi,
S., Wallace, A. J., Perry, M., Anup, R., W Lloyd, S. K., *et al.* Incidence of
mosaicism in 1055 de novo NF2 cases: much higher than previous estimates with
high utility of next-generation sequencing. (2020) doi:10.1038/s41436.

33. Hua, Y., Vickers, T. A., Baker, B. F., Bennett, C. F. & Krainer, A. R.
Enhancement of SMN2 exon 7 inclusion by antisense oligonucleotides targeting
the exon. *PLoS Biol.* **5**, 729–744 (2007).

34. Crooke, R. M., Graham, M. J., Lemonidis, K. M., Whipple, C. P., Koo, S. &

Perera, R. J. An apolipoprotein B antisense oligonucleotide lowers LDL
cholesterol in hyperlipidemic mice without causing hepatic steatosis. *J. Lipid*
Res. **46**, 872–884 (2005).

35. McClorey, G., Moulton, H. M., Iversen, P. L., Fletcher, S. & Wilton, S. D.
Antisense oligonucleotide-induced exon skipping restores dystrophin expression
in vitro in a canine model of DMD. *Gene Ther.* **13**, 1373–1381 (2006).

36. Stein, C. A. & Castanotto, D. FDA-Approved Oligonucleotide Therapies in 2017.
Mol. Ther. **25**, 1069–1075 (2017).

37. Xiong, H., Veedu, R. N. & Diermeier, S. D. Recent advances in oligonucleotide
therapeutics in oncology. *Int. J. Mol. Sci.* **22**, (2021).

38. Dominant, N. R. E. A. Antisense Oligonucleotide-Based Downregulation of. **3**,
1–12.

39. Miller, T., Cudkowicz, M., Shaw, P. J., Andersen, P. M., Atassi, N., Bucelli, R.
C., Genge, A., Glass, J., Ladha, S., Ludolph, A. L., *et al.* Phase 1–2 Trial of
Antisense Oligonucleotide Tofersen for SOD1 ALS . *N. Engl. J. Med.* **383**, 109–
119 (2020).

40. Tabrizi, S. J., Leavitt, B. R., Landwehrmeyer, G. B., Wild, E. J., Saft, C., Barker,
R. A., Blair, N. F., Craufurd, D., Priller, J., Rickards, H., *et al.* Targeting
Huntingtin Expression in Patients with Huntington's Disease. *N. Engl. J. Med.*
380, 2307–2316 (2019).

41. Gleave, M. E. & Monia, B. P. Antisense therapy for cancer. *Nat. Rev. Cancer* **5**,
468–479 (2005).

42. Levin, A. A. Treating Disease at the RNA Level with Oligonucleotides. *N. Engl.*
J. Med. **380**, 57–70 (2019).

43. Sazani, P., Kang, S. H., Maier, M. A., Wei, C., Dillman, J., Summerton, J.,
Manoharan, M. & Kole, R. Nuclear antisense effects of neutral, anionic and
cationic oligonucleotide analogs. *Nucleic Acids Res.* **29**, 3965–74 (2001).

44. Castellanos, E., Rosas, I., Solanes, A., Bielsa, I., Lázaro, C., Carrato, C.,

Hostalot, C., Prades, P., Roca-Ribas, F., Blanco, I., *et al.* In vitro antisense
therapeutics for a deep intronic mutation causing Neurofibromatosis type 2. *Eur.*
J. Hum. Genet. **21**, 769–773 (2013).

45.   Yachdav, G., Kloppmann, E., Kajan, L., Hecht, M., Goldberg, T., Hamp, T.,
Hönigschmid, P., Schafferhans, A., Roos, M., Bernhofer, M., *et al.* PredictProtein
- An open resource for online prediction of protein structural and functional
features. *Nucleic Acids Res.* **42**, 337–343 (2014).

46.   Bernhofer, M., Dallago, C., Karl, T., Satagopam, V., Heinzinger, M., Littmann,
M., Olenyi, T., Qiu, J., Schütze, K., Yachdav, G., *et al.* PredictProtein -
Predicting protein structure and function for 29 years. *Nucleic Acids Res.* **49**,
W535–W540 (2021).

47.   Kluwe, L., MacCollin, M., Tatagiba, M., Thomas, S., Hazim, W., Haase, W. &
Mautner, V. F. Phenotypic variability associated with 14 splice-site mutations in
the NF2 gene. *Am. J. Med. Genet.* **77**, 228–233 (1998).

48.   Morrison, H., Sherman, L. S., Legg, J., Banine, F., Isacke, C., Haipek, C. A.,
Gutmann, D. H., Ponta, H. & Herrlich, P. The NF2 tumor suppressor gene
product, merlin, mediates contact inhibition of growth through interactions with
CD44. *Genes Dev.* **15**, 968–980 (2001).

49.   Scoles, D. R. The merlin interacting proteins reveal multiple targets for NF2
therapy. **1785**, 32–54 (2008).

50.   Li, Q., Nance, M. R., Kulikauskas, R., Nyberg, K., Fehon, R., Karplus, P. A.,
Bretscher, A. & Tesmer, J. J. G. Self-masking in an Intact ERM-merlin Protein:
An Active Role for the Central α-Helical Domain. *J. Mol. Biol.* **365**, 1446–1459
(2007).

51.   Pelton, P. D., Sherman, L. S., Rizvi, T. A., Marchionni, M. A., Wood, P.,
Friedman, R. A. & Ratner, N. Ruffling membrane, stress fiber, cell spreading and
proliferation abnormalities in human Schwannoma cells. *Oncogene* **17**, 2195–
2209 (1998).

52.   Lallemand, D., Curto, M., Saotome, I., Giovannini, M. & McClatchey, A. I. NF2

deficiency promotes tumorigenesis and metastasis by destabilizing adherens
junctions. *Genes Dev.* **17**, 1090–1100 (2003).

53.   Stamenkovic, I. & Yu, Q. Merlin, a "Magic" Linker Between
the Extracellular Cues and Intracellular Signaling Pathways that Regulate Cell
Motility, Proliferation, and Survival. *Curr. Protein Pept. Sci.* **11**, 471–484
(2010).

54.   Chiasson-Mackenzie, C., Morris, Z. S., Liu, C. H., Bradford, W. B., Koorman, T.
& McClatchey, A. I. Merlin/ERM proteins regulate growth factor-induced
macropinocytosis and receptor recycling by organizing the plasma
membrane:Cytoskeleton interface. *Genes Dev.* **32**, 1201–1214 (2018).

55.   McClatchey, A. I. & Fehon, R. G. Merlin and the ERM proteins - regulators of
receptor distribution and signaling at the cell cortex. *Trends in Cell Biology* vol.
19 198–206 at https://doi.org/10.1016/j.tcb.2009.02.006 (2009).

56.   Johnson, K. C., Kissil, J. L., Fry, J. L. & Jacks, T. Cellular transformation by a
FERM domain mutant of the Nf2 tumor suppressor gene. *Oncogene* **21**, 5990–
5997 (2002).

57.   Popplewell, L. J., Adkin, C., Arechavala-Gomeza, V., Aartsma-Rus, A., de
Winter, C. L., Wilton, S. D., Morgan, J. E., Muntoni, F., Graham, I. R. &
Dickson, G. Comparative analysis of antisense oligonucleotide sequences
targeting exon 53 of the human DMD gene: Implications for future clinical trials.
Neuromuscul. Disord. **20**, 102–110 (2010).

58.   Dhuri, K., Bechtold, C., Quijano, E., Pham, H., Gupta, A., Vikram, A. & Bahal,
R. Antisense oligonucleotides: An emerging area in drug discovery and
development. *Journal of Clinical Medicine* vol. 9 1–24 at
https://doi.org/10.3390/JCM9062004 (2020).

59.   Madocsai, C., Lim, S. R., Geib, T., Lam, B. J. & Hertel, K. J. Correction of
SMN2 Pre-mRNA splicing by antisense U7 small nuclear RNAs. *Mol. Ther.* **12**,
1013–1022 (2005).

60.   Tang, Z., Zhao, J., Pearson, Z. J., Boskovic, Z. V. & Wang, J. Rna-targeting

splicing modifiers: Drug development and screening assays. *Molecules* **26**, (2021).

61. Maass, P. G., Barutcu, A. R., Shechner, D. M., Weiner, C. L., Melé, M. & Rinn, J. L. Spatiotemporal allele organization by allele-specific CRISPR live-cell imaging (SNP-CLING). *Nat. Struct. Mol. Biol.* **25**, 176–184 (2018).

62. György, B., Nist-Lund, C., Pan, B., Asai, Y., Karavitaki, K. D., Kleinstiver, B. P., Garcia, S. P., Zaborowski, M. P., Solanes, P., Spataro, S., *et al.* Allele-specific gene editing prevents deafness in a model of dominant progressive hearing loss. *Nat. Med.* **25**, 1123–1130 (2019).

63. Kocher, T., Peking, P., Klausegger, A., Murauer, E. M., Hofbauer, J. P., Wally, V., Lettner, T., Hainzl, S., Ablinger, M., Bauer, J. W., *et al.* Cut and Paste: Efficient Homology-Directed Repair of a Dominant Negative KRT14 Mutation via CRISPR/Cas9 Nickases. *Mol. Ther.* **25**, 2585–2598 (2017).

64. Tomkiewicz, T. Z., Suárez-Herrera, N., Cremers, F. P. M., Collin, R. W. J. & Garanto, A. Antisense oligonucleotide-based rescue of aberrant splicing defects caused by 15 pathogenic variants in abca4. *Int. J. Mol. Sci.* **22**, (2021).

65. Aguti, S., Bolduc, V., Ala, P., Turmaine, M., Bönnemann, C. G., Muntoni, F. & Zhou, H. Exon-Skipping Oligonucleotides Restore Functional Collagen VI by Correcting a Common COL6A1 Mutation in Ullrich CMD. *Mol. Ther. - Nucleic Acids* **21**, 205–216 (2020).

66. Panagiotopoulos, A. L., Karguth, N., Pavlou, M., Böhm, S., Gasparoni, G., Walter, J., Graf, A., Blum, H., Biel, M., Riedmayr, L. M., *et al.* Antisense Oligonucleotide- and CRISPR-Cas9-Mediated Rescue of mRNA Splicing for a Deep Intronic CLRN1 Mutation. *Mol. Ther. - Nucleic Acids* **21**, 1050–1061 (2020).

67. Pros, E., Fernández-Rodríguez, J., Canet, B., Benito, L., Sánchez, A., Benavides, A., Ramos, F. J., López-Ariztegui, M. A., Capellá, G., Blanco, I., *et al.* Antisense therapeutics for neurofibromatosis type 1 caused by deep intronic mutations. *Hum. Mutat.* **30**, 454–462 (2009).

68. Cartegni, L. & Krainer, A. R. Correction of disease-associated exon skipping by
synthetic exon-specific activators. *Nat. Struct. Biol.* **10**, 120–125 (2003).

69. Wilusz, J. E., Devanney, S. C. & Caputi, M. Chimeric peptide nucleic acid
compounds modulate splicing of the bcl-x gene in vitro and in vivo. *Nucleic*
Acids Res. **33**, 6547–6554 (2005).

70. Passini, M. A., Bu, J., Richards, A. M., Kinnecom, C., Sardi, S. P., Stanek, L. M.,
Hua, Y., Rigo, F., Matson, J., Hung, G., *et al.* Antisense oligonucleotides
delivered to the mouse CNS ameliorate symptoms of severe spinal muscular
atrophy. *Sci. Transl. Med.* **3**, (2011).

71. Wheeler, T. M., Leger, A. J., Pandey, S. K., Mac Leod, A. R., Wheeler, T. M.,
Cheng, S. H., Wentworth, B. M., Bennett, C. F. & Thornton, C. A. Targeting
nuclear RNA for in vivo correction of myotonic dystrophy. *Nature* **488**, 111–117
(2012).

72. Panza, F., Lozupone, M., Seripa, D., Daniele, A., Watling, M., Giannelli, G. &
Imbimbo, B. P. Development of disease-modifying drugs for frontotemporal
dementia spectrum disorders. *Nat. Rev. Neurol.* **16**, 213–228 (2020).

73. Murray, S. F., Jazayeri, A., Matthes, M. T., Yasumura, D., Yang, H., Peralta, R.,
Watt, A., Freier, S., Hung, G., Adamson, P. S., *et al.* Allele-specific inhibition of
rhodopsin with an antisense oligonucleotide slows photoreceptor cell
degeneration. *Investig. Ophthalmol. Vis. Sci.* **56**, 6362–6375 (2015).

74. DeVos, S. L., Miller, R. L., Schoch, K. M., Holmes, B. B., Kebodeaux, C. S.,
Wegener, A. J., Chen, G., Shen, T., Tran, H., Nichols, B., *et al.* Tau reduction
prevents neuronal loss and reverses pathological tau deposition and seeding in
mice with tauopathy. *Sci. Transl. Med.* **9**, (2017).

75. Truong, K., Ahmad, I., Jason Clark, J., Seline, A., Bertroche, T., Mostaert, B.,
Van Daele, D. J. & Hansen, M. R. Nf2 Mutation in Schwann Cells Delays
Functional Neural Recovery Following Injury. *Neuroscience* **374**, 205–213
(2018).

76. Cockburn, K., Biechele, S., Garner, J. & Rossant, J. The hippo pathway member

nf2 is required for inner cell mass specification. *Curr. Biol.* **23**, 1195–1201
(2013).

77. Hennigan, R. F., Thomson, C. S. & Ratner, N. Merlin Tumor Suppressor
Function is Regulated by PIP2-Mediated Dimerization.
doi:10.1101/2021.11.11.468247.

78. Bachir, S., Shah, S., Shapiro, S., Koehler, A., Mahammedi, A., Samy, R. N.,
Zuccarello, M., Schorry, E. & Sengupta, S. Neurofibromatosis type 2 (NF2) and
the implications for vestibular schwannoma and meningioma pathogenesis.
International Journal of Molecular Sciences vol. 22 1–12 at
https://doi.org/10.3390/ijms22020690 (2021).

79. Castellanos, E., Gel, B., Rosas, I., Tornero, E., Santín, S., Pluvinet, R., Velasco,
J., Sumoy, L., Valle, J., Perucho, M., *et al.* A comprehensive custom panel design
for routine hereditary cancer testing: preserving control , improving diagnostics
and revealing a complex variation landscape. *Nat. Publ. Gr.* 1–12 (2017)
doi:10.1038/srep39348.

80. Leier, A., Moore, M., Liu, H., Daniel, M., Hyde, A. M., Messiaen, L., Korf, B.
R., Selvakumaran, J., Ciszewski, L., Lambert, L., *et al.* Targeted exon skipping
of NF1 exon 17 as a therapeutic for neurofibromatosis type I. *Mol. Ther. Nucleic*
Acid **28**, 261–278 (2022).

81. Hornbeck, P. V, Zhang, B., Murray, B., Kornhauser, J. M., Latham, V. & Psp, P.
R. PhosphoSitePlus , 2014: mutations , PTMs and recalibrations. **43**, 512–520
(2015).

 FIGURE LEGENDS

Figure 1. Experimental approaches to test antisense therapy for *NF2* variants. (A)

Truncating variants. In blue, expected outcome after PMO treatment**.** PMOs are

designed at the donor and acceptor sites of an in-frame exon to induce skipping of exons

harbouring truncating variants. The generated in-frame deletion is expected to

ameliorate the phenotype *in vitro*. (B) **Experimental approach to test antisense**

therapy for splicing *NF2* variants that induce exon skipping and result in a truncating

protein and (C) that **induce the inclusion of intronic regions as coding sequences**,

resulting in alterations in the gene reading frame that finally generate a truncating

protein. In blue, expected outcome after PMO treatment. In both approaches, the PMO

is expected to mask the aberrant splicing signal and rescue the NF2 phenotype *in vitro*.

Figure 2. *NF2* RNA transcript and **Merlin levels after inducing skipping of in-frame**

exons harbouring truncating variants. (A) *NF2* RNA transcripts after inducing

skipping of exons 4, 8 and 11 of the *NF2* gene showed that we were able to generate

these exon-less forms at RNA level. The different lanes correspond to the following

conditions: (1) Untreated 24h; (2) Treated (20µM for ES4 and ES8, 40 µM for ES11)

24h; (3) Untreated 48h; (4) Treated 48h; (5) Untreated 72h; (6) Treated 72h. (B) Merlin

Western Blot after exon-specific PMOs treatment showed a decrease in Merlin total

levels and no presence of exon-less, shorter forms of Merlin-e4 and Merlin-e8. Merlin-

e11 was detected and a slight increase was determined in total Merlin levels. Graphs

show total protein quantification from three independent experiments in fibroblasts

from one patient. Increases or decreases in Merlin levels are considered in respect to

the untreated sample.

NF2$^{(+/+)}$ fibroblasts stand for control fibroblasts from healthy donors. When indicated

"untreated" stands for patient's fibroblasts (NF2$^{(+/-)}$) without PMO treatment, "Endo-

porter" for patient's fibroblasts (NF2$^{(+/-)}$) treated with PMO's vehicle, "treated" for

patient's fibroblasts (NF2$^{(+/-)}$) treated with a pair of PMOs for each exon, and "PMO

Ctrl" for patient's fibroblasts (NF2$^{(+/-)}$) treated with a control PMO that does not

interfere with *NF2*. ES stands for Exon Skipping and NPE for Normalized Protein

Expression.

Figure 3. E11 PMO treatment of *NF2*$^{(+/-)}$ fibroblasts induced improvement in actin

cytoskeleton organization. Actin immunocytochemistry analysis revealed cytoskeletal

abnormalities in patient's derived fibroblasts. Patient_ES11_1 showed a decrease in

membrane ruffle formation and Patient_ES11_2 responded to the treatment showing

improved culture organization. DAPI was used to stain cell nuclei and Phalloidin is

shown in red. *NF2*($^{(+/+)}$) fibroblasts stand for control fibroblasts from healthy donors.

When indicated "untreated" stands for patient's fibroblasts (NF2$^{(+/-)}$) without PMO

treatment, "Endo-porter" for patient's fibroblasts (NF2$^{(+/-)}$) treated with PMO's vehicle

and "treated" for patient's fibroblasts (NF2$^{(+/-)}$) treated with ES11 PMO. Scale bar:

75µm.

Figure 4. Expression of Merlin-e11 was able to reduce the proliferation capacity of

patient fibroblasts. Quantification of EdU-positive cells (percentage over DAPI-

positive nuclei). A significant decrease was detected in the proliferation rate of treated

fibroblasts when compared to Endo-Porter for Patient_ES11_1 (A). Patient_ES11_2 (B)

showed the same tendency. (C) Representation of the mean values of both patients.

Mean±SD is represented in the bar, representing three independent experiments.

*p<0.05, **p<0.01 denotes statistical significance using unpaired t-test. See also Figure

S5. In the figure, "Endo-porter" stands for patient's fibroblasts (NF2$^{(+/-)}$) treated with

PMO's vehicle, "treated" for patient's fibroblasts (NF2$^{(+/-)}$) treated with specific PMO

and "PMO Ctrl" for patient's fibroblasts (NF2$^{(+/-)}$) treated with a control PMO that does

not interfere with *NF2*.

Figure 5. (A) Design of the PMO specific for each variant. For the second design

(v2), Canonical Splice Sites (CSS), Branch Points (BR), Acceptor and Donor Sites (AS

and DS), Exonic Splicing Enhancer sequences (ESE) and Exonic Splicing Silencer

(ESS) sequences indicated were taken into account and excluded, when possible, for the

sequence design in order to avoid masking the correct splicing signals. The exonic part

of the sequence is shown in a grey box and the unlabelled part constitute intronic

regions. Tested genetic variants are highlighted in red. **(B) The use of variant-specific**

PMO did not allow correction of the aberrant splicing signalling. The effect of the

PMOs v2 is shown at cDNA level through dose response and a time course experiments

for each patient (Spl_1, Spl_2, Spl_3 and Spl_4). *NF2* transcripts according to the

expected molecular weight are indicated next to the bands in the agarose gel. *NF2*($^{(+/+)}$)

fibroblasts stand for control fibroblasts from healthy donors and *NF2*($^{(+/+)}$) fibroblasts

treated indicate that have been treated with the specific PMO of each patient at 20µM.

When indicated "untreated" stands for patient's fibroblasts (NF2$^{(+/-)}$) without PMO

treatment, "Endo-porter" for patient's fibroblasts (NF2$^{(+/-)}$) treated with PMO's vehicle

and "treated" for patient's fibroblasts (NF2$^{(+/-)}$) treated with specific PMO. ES stands for

Exon Skipping and WT for Wild Type.

TABLES

Table 1. Exon-specific PMOs design to target *NF2* truncating variants

Exon	Target *NF2* variant	Target region	PMOs designed	PMO_Design 1	PMO_Design 2
4 (n=2)	Any truncating at exon 4	Exon 4_5'	PMO Pair	**PMO_5'ES4**: ACCTGAAAGGAGCAACAAGGGAGAC	
		Exon 4_3'		**PMO_3'ES4**: TTTCTTCTTTGAGCCTACCTTGGCC	
8 (n=5)	Any truncating at exon 8	Exon 8_5'	PMO Pair	**PMO_5'ES8**: TTTATTCTGTGGATCCAATAAGAAC	
		Exon 8_3'		**PMO_3'ES8**: TGCAGTACACACATGTCCTACCTCC	
11 (n=2)	Any truncating at exon 11	Exon 11_5'	PMO Pair	**PMO_5'ES11_v1**: AGGCGCTGCCGCTCCATCTGCGAGG	**PMO_5'ES11_v2**: CGCTCCATCTGCGAGGGGTGAAGAA
		Exon 11_3'		**PMO_3'ES11_v1**: CCAGCCCCTCAGAAATCACCAGTGC	**PMO_3'ES11_v2**: CCTCAGAAATCACCAGTGCTTCGTT

n indicates the number of patients derived fibroblasts in which PMOs is tested

Table 2. NF2 splicing loss-of-function PMO design

Patient's code	c.	g.	p.	Variant Effect	PMO_Design
Patient_Spl_Ctrl (PMO_Ctrl)	c-1447-240T>A	g.74408☐T>A	p.Pro482Profs*39	Insertion of a cryptic exon of 167pb between exons 13 and 14	CATCCCTCAAATCTCTTACCGTTCT*
Patient_Spl_1	c.1736A>G	g.83045A>G	p.Lys525Asnfs*19	ES of exon 15	**Spl1_PMO_v2:** CCTTTTAATGGTATTGTGCTTGCTG
Patient_Spl_2	c.810+1dupG	g.62785dupG	p.Phe271_295del	ES of exon 8	**Spl2_PMO_v2:** CCCTCCTTGTCACTGTACGAGATGT
Patient_Spl_3	c.241-9A>G	g.35526A>G	p.Val81Phefs*44	New acceptor site exon 3. Inclusion of 8 intronic nucleotides as coding sequence	**Spl3_PMO_v2:** GCAGAACTGCAGAGCAAAAGACAAA
Patient_Spl_4	c.241-13T>A	g.35522T>A	p.Val81Phefs*44	New acceptor site exon 3. Inclusion of 10 intronic nucleotides as coding sequence	**Spl4_PMO_v2:** GCAGAATTGCTGAGCAAAAGACAAA

*Exon Skipping (ES). * Published in Castellanos et al. 2013*

A. Truncating variants

No protein synthesis
Patient's severe phenotype

In frame exon skipping

Partially defective protein
Patient's mild phenotype

B. Splicing variants inducing exon skipping

Exon skipping

Partially defective protein

Normal protein
Rescue *NF2* phenotype

C. Splicing variants inducing inclusion of intronic regions as coding sequences

No protein synthesis
Patient's severe phenotype

Normal protein
Rescue *NF2* phenotype

A. *NF2* RNA transcript after inducing ES

ES4 *NF2* transcript

ES8 *NF2* transcript

ES11 *NF2* transcript

B. Merlin levels after ES of in-frame exons harbouring truncating variants

Merlin-e4

Merlin-e8

Merlin-e11

A. Patient_ES11_1

B. Patient_ES11_2

C. Merlin-e11

A

Patient Spl1

Patient Spl2

Patients Spl3 and Spl4

B

B.I. Patient Spl1

Dose Response

Endo-Porter 5µM 10µM 20µM NF2(+/+) fibroblasts

Time Course (20µM)

Untreated Treated 24h Treated 48h Treated 72h NF2(+/+) fibroblasts treated

→ Exons 14, 15, 17 (Isoform I, WT)
→ Heteroduplex (14, 15, 17 and 14, 17)
→ Exons 14, 16 (Isoform II, ES15)
→ Heteroduplex (14, 6 amd 14, 17)
→ Exons 14, 17 (Isoform I, ES15)

B.II. Patient Spl2

Dose Response

Endo-Porter 5µM 10µM 20µM NF2(+/+) fibroblasts

Time Course (20µM)

Untreated Treated 24h Treated 48h Treated 72h NF2(+/+) fibroblasts treated

→ WT
→ Exon Skipping 8

B.III. Patient Spl3

Dose Response

Endo-Porter 5µM 10µM 20µM NF2(+/+) fibroblasts

Time Course (20µM)

Untreated Treated 24h Treated 48h Treated 72h NF2(+/+) fibroblasts treated

→ WT
→ Heteroduplex WT and ES of exons 2 and 3
→ ES of exons 2 and 3

B.IV. Patient Spl4

Dose Response

Endo-Porter 5µM 10µM 20µM NF2(+/+) fibroblasts

Time Course (20µM)

Untreated Treated 24h Treated 48h Treated 72h NF2(+/+) fibroblasts treated

→ WT
→ Heteroduplex WT and ES of exons 2 and 3
→ ES of exons 2 and 3