Phosphorylation by casein kinase 2 ensures ER-phagy receptor TEX264 binding to ATG8 proteins

Haruka Chino[1,] Akinori Yamasaki[2,6], Koji L Ode[3], Hiroki R Ueda[3,4], Nobuo N Noda[2,5*], Noboru Mizushima[1]*

[1] Department of Biochemistry and Molecular Biology, and [3] Department of Systems Pharmacology, Graduate School of Medicine, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

[2] Institute of Microbial Chemistry (BIKAKEN), 3-14-23 Kamiosaki, Shinagawa-ku, Tokyo 141-0021, Japan

[4] Laboratory for Synthetic Biology, RIKEN Center for Biosystems Dynamics Research, 1-3 Yamadaoka, Suita, Osaka 565-0871, Japan

[5] Division of Biological Molecular Mechanisms, Institute for Genetic Medicine, Hokkaido University, Kita 15, Nishi 7, Sapporo 060-0815, Japan

[6] Present address: Cell Biology Center, Institute of Innovative Research, Tokyo Institute of Technology, 4259 Nagatsuta-cho, Midori-ku, Yokohama, Kanagawa, 226-8503, Japan

*Correspondence

Noboru Mizushima, Department of Biochemistry and Molecular Biology, Graduate School of Medicine, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan; Tel: +81-3-5841-3440; Fax: +81-3-3815-1490; E-mail: nmizu@m.u-tokyo.ac.jp

Nobuo N Noda, Division of Biological Molecular Mechanisms, Institute for Genetic Medicine, Hokkaido University, Kita 15, Nishi 7, Sapporo 060-0815, Japan; Tel: +81-11-706-5069; Fax: +81-11-706-7826; E-mail: nn@igm.hokudai.ac.jp

Running title   Phosphorylation of TEX264 by CK2

Abstract (<175 words)

Selective autophagy cargos are recruited to autophagosomes primarily by interacting with autophagosomal ATG8 family proteins via the LC3-interacting region (LIR). The upstream sequence of most LIRs contains negatively charged residues such as Asp, Glu, and phosphorylated Ser and Thr. However, the significance of LIR phosphorylation (compared with having acidic amino acids) and the structural basis of phosphorylated LIR–ATG8 binding are not entirely understood. Here, we show that the serine residues upstream of the core LIR of the endoplasmic reticulum (ER)-phagy receptor TEX264 are phosphorylated by casein kinase 2, which is critical for its interaction with ATG8s, autophagosomal localization, and ER-phagy. Structural analysis showed that phosphorylation of these serine residues increased binding affinity by producing multiple hydrogen bonds with ATG8s that cannot be mimicked by acidic residues. This binding mode is different from those of other ER-phagy receptors that utilize a downstream helix, which is absent from TEX264, to increase affinity. These results suggest that phosphorylation of the LIR is critically important for strong LIR–ATG8 interactions, even in the absence of auxiliary interactions.

Keywords: crystal structure/ ER-phagy/ LC3-interacting region/ phosphorylation/ selective autophagy

Introduction

Macroautophagy (hereafter, autophagy) is an intracellular degradation system through which cytoplasmic components are degraded in lysosomes. During autophagy, a portion of the cytoplasm is sequestered by autophagosomes, which then fuse with lysosomes to degrade their internal material (Nakatogawa, 2020; Søreng *et al*, 2018). Autophagy degrades intracellular materials not only randomly but also selectively (Gatica *et al*, 2018; Johansen & Lamark, 2020). In selective autophagy, certain soluble proteins, protein aggregates, organelles (e.g., mitochondria, endoplasmic reticulum (ER), and lysosomes), and intracellular pathogens are selectively recognized and degraded (Anding & Baehrecke, 2017; Chino & Mizushima, 2020; Gomes & Dikic, 2014; Palikaras *et al*, 2018). Selective autophagy is important for maintaining cellular homeostasis and is associated with human diseases (Dikic & Elazar, 2018; Hübner & Dikic, 2020; Mizushima & Levine, 2020). Most of the selective substrates are recognized by autophagosomal ATG8 family proteins directly or indirectly via selective autophagy receptors (Kirkin & Rogov, 2019). ATG8 proteins are ubiquitin-like proteins that can be covalently conjugated to phosphatidylethanolamine in the autophagic membrane (Mizushima, 2020). ATG8 proteins are classified into two subfamilies in mammals, namely, the LC3 (including LC3A, LC3B, and LC3C) and GABARAP (including GABARAP, GABARAPL1, and GABARAPL2) subfamilies.

Interactions with ATG8 proteins involve the LC3-interacting region (LIR) in selective substrates (Johansen & Lamark, 2020). The canonical LIR motif consists of the consensus sequence Θ_0-X_1-X_2-Γ_3 (called the core LIR), with Θ representing an aromatic residue (W/F/Y), Γ an aliphatic residue (L/V/I), and X any amino acid. The side chains of the aromatic and aliphatic residues of the LIR motif interact with two hydrophobic pockets on the surface of ATG8s to stabilize an intermolecular β-sheet formed by the LIR motif and β-strand 2 within ATG8s (Suzuki *et al*, 2014). In addition to these canonical interactions, other interactions sometimes critically enhance binding. For example, negatively charged residues are often found at the X_{-3}, X_{-2}, X_{-1}, X_1, and X_2 positions. Acidic residues such as Asp and Glu and phosphorylated Ser and Thr at X_{-3}, X_{-2}, and

X_{-1} can form electrostatic interactions with Lys46 and Lys48 of Atg8, whereas

acidic residues at X_1 and X_2 form electrostatic interactions with Arg67 and Arg28

of Atg8 in yeasts (Noda *et al*, 2010). Phosphorylation of Ser and Thr residues at

these positions could serve as a molecular switch. Upon bacterial infection,

microbe-derived lipopolysaccharides induce phosphorylation of S177 (X_{-1}) of

optineurin by TANK binding kinase 1 (TBK1) and increase the affinity to ATG8s,

promoting selective autophagy (Rogov *et al*, 2013; Wild *et al*, 2011). In addition,

the mitophagy receptors BNIP3 and NIX (also known as BNIP3L) are also

phosphorylated to function (Rogov *et al*, 2017; Zhu *et al*, 2013). These

publications indicated the formation of additional intermolecular hydrogen bonds

between Ser in positions X_{-1} and X_{-2} and Arg10, Arg11, Lys51 in

LC3/GABARAPs. Besides these, some LIRs possess an α-helix downstream of

the core LIR, which markedly increases their affinity with ATG8s through the

additional intermolecular hydrogen bonds between electronegative residues

after the core LIR and α-helix 3 of ATG8s (R69 and R70 in LC3B) (Johansen &

Lamark, 2020; Li *et al*, 2018; Mochida *et al*, 2020; Sakurai *et al*, 2017). However,

the structural basis by which these residues surrounding the core LIR enhance

binding to ATG8s is not completely understood because these previous reports

utilized nuclear magnetic resonance analysis or crystallography using

phosphomimicking peptides or single-chain constructs.

In this study, we show that the serine residues upstream of the core LIR

of TEX264, a receptor for autophagic degradation of the endoplasmic reticulum

(ER-phagy) (An *et al*, 2019; Chino *et al*, 2019), are phosphorylated by casein

kinase 2 (CK2). Phosphorylation of the LIR was required for the interaction of

TEX264 with ATG8s, autophagosomal localization, and ER-phagy receptor

function. According to the results of crystallographic analysis, we propose that

the phosphorylation of residues upstream of the core LIR is critically important to

produce strong LIR–ATG8 interactions, even in the absence of auxiliary

interactions, by producing multiple hydrogen bonds in addition to ionic

interactions.

Results

Ser266, Ser269, Ser271, and Ser272 upstream of the core LIR in TEX264 are phosphorylated

TEX264 contains multiple highly conserved serine residues upstream of the core LIR (Fig 1A). An anti-phosphoserine antibody reacted to immunopurified FLAG-tagged TEX264 (TEX264–FLAG), suggesting that the serine residues of TEX264 were phosphorylated (Fig 1B). To determine the exact phosphorylation sites, we performed tandem mass spectrometry (MS/MS). To detect a peptide containing the LIR sequence in MS/MS analysis, we generated a TEX264^{G280K} mutant such that the otherwise too long trypsinized peptides around the LIR in wild-type (WT) TEX264 (SEHSYSESGASGSSFEELDLEGEGPLGESR) could be digested by trypsin at the introduced Lys residue. MS/MS analysis revealed multiple phosphorylations in various combinations around the LIR sequence. Among them, Ser266, Ser269, Ser271, and Ser272 were identified as being phosphorylated (Fig 1A, asterisks, Fig EV1A)

Ser271 and Ser272 are essential for ATG8 binding and autophagosomal localization of TEX264

To investigate the importance of the phosphorylation of Ser266, Ser269, Ser271, and Ser272 in TEX264, we replaced them with alanine and tested the interaction of TEX264 with ATG8s. Among mammalian ATG8s, TEX264 is known to interact with LC3A, LC3B, GABARAP, and GABARAPL1 (An *et al.*, 2019; Chino *et al.*, 2019). Alanine-substitution mutants of either or any combination of the serine residues decreased the level of serine phosphorylation (Fig 1C). Unexpectedly, mutation of the core LIR (LIR4A, FEEL/AAAA) also impaired serine phosphorylation (discussed later). As with the core LIR mutant, single or multiple serine residue mutants including Ser271 and Ser272 abolished the interaction with LC3A/B and GABARAP, although TEX264^{S269A} still weakly bound to GABARAP (Fig 1C). Alanine substitution of Ser266 did not affect the interaction with LC3A/B and GABARAP (Fig 1C). C-terminal GFP-tagged TEX264

(TEX264-GFP) was present in the ER and formed punctate structures with LC3

during starvation (Fig 1D and E). By contrast, although the TEX264^{S269A},

TEX264^{S271A}, and TEX264^{S272A} mutants still localized to the ER, their

accumulation at LC3 structures was significantly suppressed (Fig 1D and E).

Consistent with the immunoprecipitation data, TEX264^{S266A} formed punctate

structures and localized to autophagosomes like wild-type TEX264 (Fig EV1B

and C). These results suggest that the serine residues, particularly Ser271 and

Ser272, upstream of the core LIR are required for ATG8 binding and

autophagosomal localization of TEX264.

To determine whether these serine residues are important for the

ER-phagy receptor function of TEX264, we monitored ER-phagy activity using a

doxycycline-inducible ER-phagy reporter with an N-terminal ER signal sequence

followed by tandem monomeric RFP and GFP and the ER retention sequence

KDEL (Chino *et al.*, 2019). Upon delivery of this reporter to lysosomes, GFP, but

not RFP, is degraded, producing an RFP fragment. When expressed in

TEX264-knockout (KO) cells, WT TEX264-FLAG recovered ER-phagy activity,

but the TEX264^{LIR4A}, TEX264$^{S271A, S272A}$, and TEX264$^{S269A, S271A, S272A}$ mutants did

not (Fig 1F and G). Thus, Ser269, Ser271 and Ser272 are required for the

ER-phagy receptor function of TEX264.

Phosphorylated TEX264 localizes to autophagosomes

To further characterize the phosphorylation of TEX264, we generated an

antibody against the peptide GASGpSpSFEEL (a.a. 267–276 of TEX264)

containing phosphorylated Ser271 and Ser272 (pTEX264), because

phosphorylation of Ser271 and Ser272 is particularly important for its interaction

with ATG8s (Fig1C). The anti-pTEX264 antibody could detect endogenous and

exogenous TEX264, but not the exogenous TEX264^{S271,272A} mutant (Fig EV1D).

Phosphatase treatment completely abolished the pTEX264 signals (Fig EV1E),

validating that the anti-pTEX264 antibody specifically recognizes

phosphorylated TEX264.

Although this antibody barely detected endogenous TEX264 by

immunofluorescence microscopy, it could detect stably overexpressed

TEX264-GFP (Fig 2A). The GFP signal of TEX264–GFP was observed as

reticular and punctate patterns, whereas the pTEX264 signal mainly

accumulated at autophagosomes under starvation conditions. The number of

pTEX264-positive puncta increased following the treatment with the lysosome

inhibitor bafilomycin A_1, indicating that pTEX264 is efficiently degraded in

lysosomes (Fig 2A and B). We determined whether endogenous pTEX264

localized to autophagosomes by a membrane floatation assay. In a stepwise

OptiPrep gradient, autophagosomes and autolysosomes containing LC3-II and

LAMP1 were collected in fractions 3 and 5, where endogenous pTEX264 was

cofractionated (Fig 2C). The ER marker SEC61B was not enriched in these

fractions. These data suggest that phosphorylated TEX264 preferentially

localizes to autophagosomes under autophagy-induction conditions such as

starvation and mTORC1 inhibition.

194   Consistently, the level of phosphorylated TEX264 was decreased under

starvations, which was restored by lysosome inhibition, and increased in

autophagy-deficient ATG3-KO and FIP200-KO cells (Fig 2D and E). However,

the ratio of phosphorylated TEX264 to total TEX264 was not changed

significantly under these conditions (Fig 2D and F). These data suggest that,

while phosphorylated TEX264 is efficiently recruited to autophagosomes, there

is an equilibrium between the phosphorylated and dephosphorylated forms.

CK2 is important for starvation-induced ER-phagy through TEX264

phosphorylation

Ser271 and Ser272 of TEX264 fulfill the criteria for a consensus motif

recognized by CK2, namely, S-X-X-E/D/Yp/Sp (Meggio *et al*, 1994; Pinna, 2002;

Rekha & Srinivasan, 2003). Thus, we tested the requirement of CK2 activity for

TEX264 phosphorylation by using CX4945, a specific inhibitor of CK2. CX4945

suppressed TEX264 phosphorylation, whereas a CK1 inhibitor (D4476) did not

(Fig 3A). Also in an invitro system, recombinant TEX264-FLAG was

phosphorylated by CK2, which was inhibited by the CK2 inhibitor (Fig 3B). These

Since the two glutamate residues (Glu274, Glu275) inside the core LIR

are part of the CK2 consensus motif, they may be required for the first priming

event, which triggers processive phosphorylation of further upstream serine

residues (Zhou *et al*, 2017). To test this hypothesis, we analyzed the

phosphorylation status of an alanine substitution mutant of TEX264

(TEX264^{E274A,E275A}). We used the anti-phosphoserine antibody instead of the

anti-pTEX264 antibody because it could not detect the TEX264^{E274A,E275A} mutant

regardless of its phosphorylation status. The TEX264^{E274A,E275A} mutant as well as

the TEX264$^{S271A, S272A}$ and TEX264^{S269A,S271A,S272A} mutants showed significantly

reduced levels of serine phosphorylation (Fig 3D and E). This also explained

why the core LIR4A mutant significantly decreased the level of serine

phosphorylation (Fig 1C). These results suggest that the two glutamates inside

the core LIR are the priming site for CK2-mediated TEX264 phosphorylation.

To determine the significance of the CK2-mediated phosphorylation of

TEX264, we analyzed the effect of the CK2 inhibitor CX4945 on the

autophagosomal localization of TEX264. CK2 inhibition significantly impaired the

formation of TEX264 puncta (Fig 3F and G), suggesting that CK2-mediated

phosphorylation is important for the autophagosomal localization of TEX264.

Next, we measured ER-phagy activity using the ER-phagy reporter and found

that the CK2 inhibitor significantly suppressed the production of the RFP

fragment under starvation conditions (Fig 3H and I). The CK2 inhibitor did not

inhibit starvation-induced bulk autophagy judged using the GFP-LC3-RFP

reporter (Fig EV2) (Kaizuka *et al*, 2016). These data suggest that CK2 activity is

important for starvation-induced ER-phagy through TEX264 phosphorylation

(Fig 3J).

Phosphorylation of the TEX264 LIR increases its affinity to ATG8s

Next, we determined whether the phosphorylation of the LIR in TEX264

promoted binding to ATG8s by an isothermal titration calorimetry (ITC) assay

(Fig 4A, Fig EV3A and Table EV1). The TEX264 peptide (residues 269–278) with

S272 phosphorylation (p-S272) showed enhanced affinity to LC3B and

GABARAP (37- and 23-fold, respectively). Moreover, multiple phosphorylations

(p-S271, p-S272 and p-S269, p-S271, p-S272) further strengthened the affinity

of TEX264 to LC3B and GABARAP, resulting in sub-μM Kd values (0.25–0.52

μM). S266 phosphorylation (p-S266) only slightly enhanced the affinity to LC3B

and GABARAP (1.8- and 6.5-fold, respectively) (Fig EV3B). The

phosphomimetic TEX264$^{S271D, S272D}$ (2SD-LIR), TEX264$^{S269D, S271D, S272D}$

(3SD-LIR), TEX264$^{S271, 272E}$ (2SE-LIR), and TEX264$^{S269E, S271E, S272E}$ (3SE-LIR)

peptides also demonstrated increased binding to LC3B and GABARAP

compared with WT TEX264 and TEX264$^{S271A, S272A}$ peptides; however, the Kd

value was within the order of μM (the minimum was 5.0 and 2.1 μM for LC3B

and GABARAP, respectively). Consistently, compared with WT TEX264,

TEX264$^{S269D, S271D, S272D}$ and TEX264$^{S269E, S271E, S272E}$ showed a significant

reduction in their affinity to LC3A, LC3A/B (the antibody reacts to both LC3A and

LC3B), and GABARAP (Fig 4B), accumulation at autophagosomes during

starvation (Fig 4C and D), and ability to promote ER-phagy (Fig 4E and F).

These data suggest that multiple phosphorylations of the upstream region of the

core LIR sufficiently strengthen the affinity of TEX264 to ATG8s to mediate

ER-phagy, which cannot be complemented by Asp or Glu mutations.

Crystallographic analysis reveals canonical interactions between

GABARAP and TEX264 LIR

To elucidate the structural basis of ATG8 binding by TEX264 LIR and its

enhancement by phosphorylation, we determined the crystal structure of

GABARAP complexed with phosphorylated TEX264 LIR (residues 269–278 with

p-S271,p-S272; referred to as LIR$^{p-S271,p-S272}$) and that of GABARAP fused to

TEX264 LIR with the S272D mutation (residues 271–281, referred to as

LIR^{S272D}) at a resolution of 3.0 and 2.0 Å, respectively (Fig 5A and B, Table EV2,

Protein Data Bank [PDB] code 7VEC and 7VED, respectively). The asymmetric

unit contained 12 and 2 copies of GABARAP for LIR$^{p-S271,p-S272}$ and LIR^{S272D},

respectively, which formed a complex with each LIR in a similar manner (Fig

EV4A–D). In both structures, the TEX264 LIR formed an intermolecular β-sheet

with β-strand 2 and bound to the hydrophobic pockets 1 and 2 of GABARAP

using the side-chains of Phe273 at position Θ_0 and Leu276 at position Γ_3,

respectively, as with canonical LIRs (Fig 5A and B). Moreover, Glu274 at position

X_1 and Glu275 at position X_2 and/or Asp277 at position X_4 formed electrostatic

interactions with Arg67 and Arg28 of GABARAP, respectively. In the case of

LIR$^{p-S271,p-S272}$, Leu278 at position X_5 formed hydrophobic interactions with

α-helix 3 of GABARAP (Fig 5A). These auxiliary interactions within the core LIR

and its downstream region are also observed for other LIRs to enhance their

binding affinity to ATG8s. These observations suggest that the manner by which

the TEX264 LIR interacts with ATG8s is similar to that of other canonical LIRs.

Phosphorylation at the X_{-2} position in TEX264 LIR produces multiple

hydrogen bonds for strong interactions with GABARAP

Although the conformation and interaction manner of the core LIR motif with

GABARAP are essentially similar between the LIR$^{p-S271,p-S272}$ and LIR^{S272D}

structures, those of the upstream region of the core LIR are totally distinct

between them (Fig 5C and D). In the LIR^{S272D} structure, the main-chain of LIR

was bent at Asp272 (position X_{-1}) at a right angle and the upstream region was

located away from α-helix 1 of GABARAP. The side-chain of Asp272 formed a

hydrogen bond with that of GABARAP Lys48. On the other hand, in the

LIR$^{p-S271,p-S272}$ structure, the upstream region had a stretched β-conformation

continuous with the core LIR motif and interacted with α-helix 1 of GABARAP.

The phosphate group of p-Ser272 (position X_{-1}) formed a hydrogen bond with

the side-chain of GABARAP Lys46, while that of p-Ser271 (position X_{-2}) formed

three hydrogen bonds with the side-chain of Tyr5 and His9 at α-helix 1 and

Lys48 (whose side-chain is locked by Glu17) of GABARAP (Fig 5E). We

confirmed the contribution of Tyr5, His9, and Lys48 to the high affinity with

TEX264 LIR p-Ser271,p-Ser272 by ITC. The Kd values of GABARAP K48A and

These

results suggest that the 4-fold enhancement of the affinity of TEX264 with

GABARAP by phosphorylation at Ser271 (p-Ser272 vs. p-Ser271,p-Ser272 in

Fig 4A), which enables sub-μM affinity, is presumably established by these

multiple hydrogen bonds.

We then wondered whether the formation of multiple hydrogen bonds of

p-Ser at position X_{-2} is specific to TEX264 or conserved among other LIRs. In

contrast to the strict conservation of Lys48 among all six ATG8 paralogs, His9

was conserved only among the GABARAP family and Tyr5 was conserved in

GABARAP/GABARAPL1. We inspected all of the structures of

GABARAP/GABARAPL1 deposited in the PDB and found that the crystal

structure of GABARAPL1 complexed with the phosphorylated LIR from the Golgi

protein short coiled-coil protein (SCOC) (PDB 7AA7; (Wirth *et al*, 2021)) showed

a strikingly similar interaction: the phosphate group of p-Ser at position X_{-2} of

SCOC LIR formed three hydrogen bonds with the side-chain of His9, Lys48, and

either Tyr5 or Arg47 of GABARAPL1 (Fig 5F) (Wirth *et al.*, 2021). The authors

revealed that Ser phosphorylation at position X_{-2} of the long SCOC LIR peptide

(containing four consecutive Glu residues at positions X_{-7}–X_{-4}) marginally

enhanced binding affinity (2.4- and 2.6-fold for GABARAP and GABARAPL1,

respectively), whereas phosphorylation at position X_{-1} did not. Moreover, they

also showed that Ser phosphorylation at position X_{-2} of the short SCOC LIR

peptide (lacking four Glu residues at positions X_{-7}–X_{-4}) enhanced binding affinity

dramatically (67- and 275-fold with GABARAP and GABARAPL1, respectively)

(Wirth *et al.*, 2021). Our ITC data showed that two or three Asp/Glu mutations at

positions X_{-3}–X_{-1} or single phosphorylation at position X_{-1} of TEX264 LIR

enhanced its binding affinity with GABARAP to a similar extent (approximately

20-fold), which was further enhanced (4-fold) by phosphorylation at position X_{-2}

(Fig 4A). These observations suggest that the enhancement of binding affinity by

phosphorylation at the upstream region of the core LIR is achieved by two

distinct mechanisms: the first endows negative charges to the LIR for

non-specific interactions with positively-charged ATG8s (Noda *et al.*, 2010), and

the second creates specific interactions with ATG8s, as observed for triple

hydrogen bonding between p-Ser at X_{-2} and Tyr5, His9, and Lys48 of

GABARAP/GABARAPL1. A phosphomimetic Asp or Glu mutation at the

upstream region would be able to account for only the former mechanism since

the chemical structure of the carboxyl side-chain of Asp/Glu cannot form similar

specific hydrogen bonds with those observed for the phosphate group.

Consistent with this idea, there is no example of triple hydrogen bonding of Asp

or Glu to Tyr5, His9, and Lys48 of GABARAP/GABARAPL1 among the dozens

of LIR-bound structures deposited in the PDB (Fig EV5). Thus, phosphorylation

but not Asp/Glu incorporation at the upstream region of the core LIR is essential

for TEX264 to generate ATG8-binding affinity strong enough (sub-μM Kd values)

to initiate ER-phagy.

Discussion

In this study, we showed that CK2-mediated phosphorylation of TEX264 is required for its ER-phagy receptor function, and the phosphorylated serine residues upstream of the core LIR are important for generating high affinity to ATG8s that cannot be mimicked by the substitution of acidic residues.

In many studies, phosphorylated Ser and Thr residues are substituted with acidic residues to mimic their phosphorylation status (Thorsness & Koshland, 1987). However, this approach does not always work. For example, inhibition of yeast Ste5, a MAPK cascade scaffold protein, requires multiple phosphorylations, but the replacement of Ser/Thr with Glu cannot mimic inhibition (Strickfaden *et al*, 2007). This may be because phosphorylated Ser and Thr are, nominally, charged twice as much as Glu and Asp at physiological pH (Cooper *et al*, 1983) and could, therefore, strengthen electrostatic interactions. However, it cannot fully explain the differences in binding. Our structural analysis suggests that the phosphorylated residue at position X_{-2} in TEX264 LIR could form three hydrogen bonds in an ideal configuration with GABARAP. This hydrogen-bonding pattern is quite similar to the recently reported binding mediated by p-Ser at position X_{-2} of SCOC LIR bound to GABARAPL1 (Wirth *et al.*, 2021). In contrast, the interaction of the phosphorylated residue at position X_{-1} is weaker and less specific: p-Ser/p-Thr at position X_{-1} of TEX264 and SCOC LIRs formed only one and no hydrogen bond with GABARAP and GABARAPL1, respectively. These observations suggest that GABARAP and GABARAPL1 conserve the specific binding site (composed of Tyr5, His9, and Lys48) for the phosphorylated residue at position X_{-2} of LIRs, which plays a critical role in increasing binding affinity. Whether such a specific binding pocket is conserved among other members of ATG8s needs to be addressed in the future. LC3 family proteins have conserved Arg and Lys residues (Arg11 and Lys51 in the case of LC3B) that correspond to His9 and Lys48 of GABARAP, respectively, which might form hydrogen bonds and/or salt-bridges with p-S271 of TEX264 to enhance binding affinity. Collectively, we propose that phosphorylation of the LIR is important for not only spatiotemporal

regulation but also strong interactions with ATG8s, which are accomplished by

the formation of specific hydrogen bonds that cannot be mimicked by acidic

residues, in addition to the previously described ionic interactions (Noda *et al.*,

2010).

Some selective autophagy receptors have only acidic amino acids

upstream of the core LIR. p62 (also known as SQSTM1), one of the most

well-studied autophagy receptors, has a "DDD" sequence followed by the core

LIR sequence "WTHL." The Kd value of p62–LC3 interactions is 1.5 µM

(Rozenknop *et al*, 2011), which is comparable with those of the interactions

between TEX264 2D/2E-LIR and LC3B and between 3D/3E-LIR and LC3B. In

contrast, other ER-phagy receptors such as FAM134B and SEC62 show high

affinity to GABARAPs with Kd values of 0.11–0.30 µM and 0.19–0.57 µM, and to

LC3s with Kd values of 0.21–1.11 µM and 0.92–5.18 µM, respectively (Mochida

et al., 2020). These two receptors possess acidic LIRs with a C-terminal short

helix, which is used in various LIRs to enhance ATG8-binding affinity to sub-µM

or even smaller Kd values through interactions with α-helix 3. Although TEX264

LIR does not possess a C-terminal helix, multiple phosphorylated serine

residues can instead generate high affinity (sub-µM Kd values) to ATG8s via

specific interactions, which enable this receptor to initiate ER-phagy, which

appears to require higher affinity with ATG8s than other types of autophagy.

Our data show that the ratio of phosphorylated TEX264 to total TEX264

is not changed under ER-phagy-inducing conditions. This may be because

phosphorylated TEX264 is efficiently degraded and/or there is an equilibrium

between the phosphorylated and dephosphorylated forms. To maintain the ratio,

it is assumed that TEX264 phosphorylation is upregulated by starvation. It is also

possible that phosphorylation could act as a switch as TEX264 has more than

one function. Besides ER-phagy, TEX264 is required for DNA quality control at

the inner nuclear membrane (Fielden *et al*, 2020). It is not known whether this

nuclear function of TEX264 also requires phosphorylation. If not,

phosphorylation and dephosphorylation could switch between these two

different functions. This may be similar to the case of AMPK-mediated

phosphorylation of the Beclin 1 LIR in response to nutrient starvation, which

underlies the differential regulation of the pro- and non-autophagic functions of

the VPS34 complex (Kim *et al*, 2013).

Materials and Methods

Cell culture

HeLa cells, HEK293T cells, and mouse embryonic fibroblasts (MEFs) were

authenticated by RIKEN. They were cultured in Dulbecco's modified Eagle's

medium (DMEM) (D6546; Sigma-Aldrich) supplemented with 10% fetal bovine

serum (S1820-500; Biowest) and 2 mM L-glutamine (25030-081; Gibco) in a 5%

CO_2 incubator. For starvation treatment, cells were washed twice with

phosphate-buffered saline (PBS) and incubated in amino acid-free DMEM

(048-33575; Wako Pure Chemical Industries) without serum. FIP200-KO HeLa

cells (Tsuboyama *et al*, 2016) and ATG3-KO HeLa cells (Sou *et al*, 2008) have

been described previously. For bafilomycin A_1 treatment, cells were cultured with

100 nM bafilomycin A_1 (B1793; Sigma-Aldrich). For CK2 inhibitor treatment, cells

were cultured with 20 µM CX4945 (S2248; Selleck). WT HeLa cells and

TEX264-KO HeLa cells stably expressing the ER-phagy reporter

pCW57.1-CMV-ssRFPGFP-KDEL (Chino *et al.*, 2019) were cultured with 0.5

µg/mL doxycycline (D3447; Sigma-Aldrich) for 24 h. Thereafter, cells were

washed with PBS twice and incubated in DMEM or amino acid-free DMEM

without serum for 9 h.

Plasmids

cDNA encoding human *TEX264* (Chino *et al.*, 2019) was inserted into pMRX-IN

(Morita *et al*, 2018) or pMRX-IB (Morita *et al.*, 2018) (these plasmids were

generated from pMXs (Kitamura *et al*, 2003)). DNAs encoding enhanced GFP

and 3×FLAG were also used for tagging. Amino acid substitution constructs

were prepared by PCR-mediated site-directed mutagenesis.

Antibodies and reagents

For immunoblotting, mouse monoclonal anti-HSP90 (610419; BD Biosciences),

anti-RFP (M208-3; MBL), and anti-FLAG (F4042; Sigma) antibodies, rabbit

polyclonal anti-TEX264 (NBP1-89866; Novus), anti-p62 (PM045; MBL),

anti-LC3A (4599s; Cell Signaling), anti-LC3B (4108s; Cell Signaling),

anti-GABARAP (13733s; Cell Signaling), and anti-RFP (a gift from Dr. T. Endo,

Kyoto Sangyo University) antibodies, and an anti-phosphoserine antibody

(ab9332; Abcam) were used as primary antibodies. Anti-mouse (111-035-003;

Jackson ImmunoResearch Laboratories, Inc.) and anti-rabbit (111-035-144;

Jackson ImmunoResearch Laboratories, Inc.) HRP-conjugated IgGs were used

as secondary antibodies. For immunostaining, mouse monoclonal anti-LC3

(M152-3; MBL) was used as a primary antibody. Alexa Fluor 568-conjugated

anti-mouse IgG (A-11004; Thermo Fisher Scientific) and Alexa Fluor

660-conjugated anti-rabbit IgG (A-21074; Molecular Probes) antibodies were

used as secondary antibodies. A rabbit polyclonal anti-pTEX264 antibody was

generated (by ABclonal) against the peptide GASGpSpSFEEL (a.a. 267–276 of

TEX264) containing phosphorylated Ser271 and Ser272 and was used for

immunoblotting and immunofluorescence microscopy. For transient transfection,

Fugene HD (VPE2311; Promega) was used according to the manufacturer's

instructions.

Preparation of retroviruses

To prepare retroviruses, HEK293T cells were transiently transfected with a

retroviral vector together with pCG-VSV-G and pCG-gag-pol (gifts from Dr. T.

Yasui, National Institutes of Biomedical Innovation, Health and Nutrition) using

Fugene HD. After cells were cultured for 2 days, the supernatant was collected

and passed through a 0.45-µm syringe filter unit (SLHV033RB; EMD Millipore).

Generation of stable cell lines

Cells were cultured with retrovirus and 8 µg/mL polybrene (H9268;

Sigma-Aldrich), and stable transformants were selected with blasticidin

(022-18713; Wako Pure Chemical Industries) or geneticin (10131; Thermo

Fisher Scientific).

Immunoprecipitation and immunoblotting

Cell lysates were prepared in a lysis buffer (50 mM Tris-HCl, pH 7.5, 150 mM

NaCl, 1 mM EDTA, 1% Triton X-100, phosphatase inhibitor cocktail [07574-61;

Nacalai Tesque], and protease inhibitor cocktail [EDTA-free]; [03969-34; Nacalai

Tesque]). After centrifugation at 17,700 × g for 10 min, the supernatants were

subjected to immunoprecipitation using anti-FLAG M2 affinity gel (A2220;

Sigma-Aldrich). Precipitated immunocomplexes were washed three times in

washing buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 1 mM EDTA, and 1%

Triton X-100) and boiled in sample buffer (46.7 mM Tris-HCl, pH 6.8, 5% glycerol,

1.67% sodium dodecyl sulfate, 1.55% dithiothreitol, and 0.02% bromophenol

blue). The samples were subsequently separated by sodium dodecyl

sulfate-polyacrylamide gel electrophoresis and transferred to Immobilon-P

polyvinylidene difluoride membranes (IPVH00010; EMD Millipore).

Immunoblotting analysis was performed with the indicated antibodies.

Super-Signal West Pico Chemiluminescent Substrate (1856135; Thermo Fisher

Scientific) or Immobilon Western Chemiluminescent HRP Substrate (P90715;

EMD Millipore) was used to visualize the signals, which were detected on a

Fusion System Solo 7S (M&S Instruments, Inc.). Contrast and brightness

adjustments and quantification were performed using Fiji software (ImageJ;

National Institutes of Health (Schindelin et al., 2012) and Photoshop CS6

(Adobe).

Immunocytochemistry

Cells grown on coverslips were washed with PBS and fixed with 4%

paraformaldehyde in PBS for 10 min at room temperature or 100% methanol for

15 min at −30°C. Fixed cells were permeabilized with 50 µg/mL digitonin (D141;

Sigma-Aldrich) in PBS for 5 min, blocked with 3% bovine serum albumin in PBS

for 15 min, and incubated with primary antibodies for 1 h. After washing three

times with PBS, cells were incubated with Alexa Fluor 568/660-conjugated goat

anti-mouse or anti-rabbit IgG secondary antibodies for 1 h. Fluorescence

microscopy was performed on a confocal laser microscope (FV3000; Olympus)

with a PLANAPO 60×1.4 NA oil-immersion objective lens and captured with

FluoView software (Olympus). The number of punctate structures and the

ER-autophagosome ratio were determined using Fiji software (ImageJ).

Cell fractionation

Cells from four 10-cm dishes cultured with 250 nM Torin1 (4247; Tocris

Bioscience) and 100 nM bafilomycin A_1 (B1793; Sigma-Aldrich) for 16 h were

harvested and washed twice with ice-cold PBS. The cell pellets were collected

after centrifugation at 600 × g for 3 min and resuspended in 2 ml ice-cold

homogenization buffer (250 mM sucrose, 20 mM Hepes-KOH, pH 7.4, 1 mM

EDTA, and complete EDTA-free protease inhibitor). Cells were disrupted by

hydrodynamic shearing (27-gauge needle). The homogenized cells were

centrifuged twice at 3,000 × g for 8 min to remove cell debris and undisrupted

cells. The supernatant was diluted with an equal volume of 60% OptiPrep

(1114542; CosmoBio) in homogenization buffer. Discontinuous OptiPrep

gradients were generated in SW41 tubes (344059; Beckman Coulter) by

overlaying the following OptiPrep solutions in homogenization buffer: 2.4 ml of

the diluted supernatant in 30% OptiPrep, 1.8 ml in 20%, 2 ml in 15%, 2 ml in

10%, 2.0 ml in 5%, and 2.0 ml in 0%. The gradients were centrifuged at 150,200

× g in SW41 Ti rotors (Beckman Coulter) for 3 h, and then 14 fractions (0.8 ml

each) were collected from the top. Proteins in each fraction were isolated by

precipitation using trichloroacetic acid. The final pellet was suspended in sample

buffer and heated at 55°C for 10min.

In vitro Phosphorylation Assay

Recombinant C-terminally FLAG-Tagged TEX264 was produced by

PURExpress in vitro protein synthesis kit (E6800; New England Biolabs).

Phosphoproteomic profiling of TEX264

HeLa cells stably expressing TEX264^{G280K}-FLAG were cultured with amino acid-free DMEM (048-33575; Wako Pure Chemical Industries) without serum for 1 h and lysed with lysis buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 1 mM EDTA, 1% NP-40, phosphatase inhibitor cocktail [07574-61; Nacalai Tesque], and protease inhibitor cocktail [EDTA-free]; [03969-34; Nacalai Tesque]). After centrifugation at 17,700 × g for 10 min, the supernatants were incubated with anti-FLAG M2 magnetic beads for 3 h at 4°C with gentle rotation. The eluted proteins were enzymatically digested according to a phase-transfer surfactant protocol (Rappsilber *et al*, 2007). Then, 50-μL eluted samples were mixed with 85 μL phase-transfer surfactant buffer. The samples were reduced with 10 mM dithiothreitol at room temperature for 30 min and alkylated with 50 mM of 2-iodoacetamide (804744; Sigma-Aldrich) at room temperature for 30 min. Next, the samples were diluted 5-fold by adding 50 mM NH_4HCO_3 solution followed by digestion with 1 μg Lysyl Endopeptidase (LysC; 121-05063; Wako Pure Chemical Industries) at 37°C for 4 h. The samples were further digested with 1 μg trypsin at 37°C for 8 h. An equal volume of ethyl acetate acidified with 0.5% TFA was added to the digested samples. After centrifugation at 10,000 × g for 10 min twice at room temperature, the aqueous phase containing peptides was collected and dried using an evaporator. The dried peptides were solubilized in 100 μL of 2% acetonitrile and 0.1% TFA. Phosphorylated peptides were enriched using a Titansphere Phos-TiO MP Kit (5010-21281; GL Science) according to the user's manual. Eluted phosphopeptides were desalted through SDB and GC tips (GL Science). The desalted eluted peptides were then dried using an evaporator.

Liquid chromatography (LC)-MS/MS analysis

The dried samples were dissolved in 2% acetonitrile and 0.1% TFA and loaded onto the LC-MS system with a Q-Exactive MS instrument (Thermo Fisher Scientific) equipped with a nano HPLC system (Advance UHPLC; Bruker Daltonics) and an HTC-PAL autosampler (CTC Analytics) with a trap column (0.3 × 5 mm, L-column, ODS; Chemicals Evaluation and Research Institute). The samples were separated by a gradient using mobile phases A (0.1% formic acid/H_2O) and B (0.1% formic acid and 100% acetonitrile) at a flow rate of 300 nL/min (4% to 32% B for 190 min, 32% to 95% B for 1 min, 95% B for 2 min, 95% to 4% B for 1 min, and 4% B for 6 min) with a home-made capillary column (length of 200 mm and inner diameter of 100 μm) packed with 2-μm C18 resin (L-column2; Chemicals Evaluation and Research Institute). Then, the eluted peptides were electrosprayed (1.8–2.3 kV) and introduced into the MS equipment. Data were obtained using the positive ion mode of data-dependent MS/MS (ddMS^2) acquisition. Full MS spectra were obtained with a scan range of 350–1800 *m/z* with 70,000 FWHM resolution at 200 *m/z*. MS^2 spectra were obtained with 17,500 FWHM resolution at 200 *m/z*. For ddMS^2 acquisition, the 10 highest precursor ions (excluding isotopes of a cluster) above an intensity threshold of 1.7e4 with a charge state from 2+ to 5+ were selected at the 4.0 *m/z* isolation window. A 20-s dynamic exclusion was applied. The obtained raw data were subjected to a database search (UniProt, reviewed mouse database as of September 13, 2018) with Sequest HT algorithm running on Proteome Discoverer 2.2 (Thermo Fisher Scientific). The database was modified to include the TEX264^{G280K} sequence. The parameters for the database search were as follows: peptide cleavage was set to trypsin; missed cleavage sites were allowed to be up to two residues; peptide lengths were set to 6–144 amino acids; and mass tolerances were set to 10 ppm for precursor ions and 0.02 Da for fragment ions. For modification conditions, carbamidomethylation at cysteine was set as a fixed modification. Oxidation at methionine and phosphorylation at serine, threonine, and tyrosine were set as variable modifications. The maximum

number of equal modifications per peptide was set to six. A significance
threshold of $p < 0.05$ was applied.

Purification of proteins and peptides

For crystallization and ITC measurement, synthesized TEX264 LIR peptides
were purchased from GenScript. Human GABARAP (residues 1–116) and
human LC3B (residues 1–120) were prepared as described previously (Mochida
et al., 2020). Human TEX264 (residues 271–281 with mutation of S272D) fused
to the N-terminus of human GABARAP with F3S/V4T mutations was inserted
downstream of the human rhinovirus 3C protease recognition site of the
pGEX6P-1 vector (Cytiva) and expressed in *Escherichia coli* BL21 (DE3). After
the bacteria were cultured at 37°C until $OD_{600} = 1.0$, they were supplemented
with 100 μM isopropyl-β-D-thiogalactopyranoside and further incubated at 16°C
overnight. Cells were collected by centrifugation, resuspended with PBS
containing 5 mM EDTA, and sonicated for 10 min. After centrifugation, the
recovered supernatants were subjected to GST-accept resins (Nacalai Tesque).
The resins were washed three times with PBS and the bound proteins were
eluted with 10 mM glutathione and 50 mM Tris, pH 8.0. The eluates were
incubated with human rhinovirus 3C protease overnight at 4°C and purified by
ion-exchange chromatography using a HiTrap DEAE FF column (Cytiva) with
buffers A (20 mM Tris, pH 8.0) and B (20 mM Tris, pH 8.0 and 1 M NaCl) and
size-exclusion chromatography using HiLoad 26/60 Superdex 200 prep grade
(Cytiva) with 20 mM HEPES, pH 6.8 and 150 mM NaCl. The purified proteins in
20 mM HEPES, pH 6.8 and 150 mM NaCl were concentrated by Centriprep
(Millipore) and kept at -80°C until use. Peptides were subjected to size-exclusion
chromatography using Superdex peptide 10/300 GL (Cytiva) with 20 mM HEPES,
pH 6.8 and 150 mM NaCl and kept at -80°C until use.

Crystallization and diffraction data collection

All crystallizations were performed through the sitting-drop vapor-diffusion
method at 20°C by mixing protein and reservoir solutions in a 1:1 volume ratio.

For crystallization of TEX264 LIR^{S272D}-GABARAP fusion, the protein was mixed

with the reservoir solution consisting of 0.2 M potassium sodium tartrate, 0.1 M

sodium citrate, pH 5.6, and 2.0 M ammonium sulfate. For crystallization of

GABARAP complexed with TEX264 LIR$^{p\text{-}S271,272}$, the protein and peptide were

mixed in a 1:1.2 molar ratio and then mixed with the reservoir solution consisting

of 30% PEG4000, 0.2 M ammonium acetate, and 0.1 M bis-Tris, pH 5.6. The

crystals of TEX264 LIR^{S272D}-GABARAP fusion were soaked in the reservoir

solution supplemented with 30.7% glycerol and frozen in liquid nitrogen. The

flash-cooled crystals were kept in a stream of nitrogen gas at -178°C during data

collection. Diffraction data collection was performed using an EIGER X4M

detector at beamline BL-1A (KEK) with a wavelength of 1.1000 Å. The diffraction

data were indexed, integrated, and scaled using X-ray Detector Software

(Kabsch, 2010). The crystals of the GABARAP-TEX264 LIR$^{p\text{-}S271,272}$ complex

were soaked in the reservoir solution supplemented with 5% glycerol and

flash-frozen in a stream of nitrogen gas and kept at -173°C during data collection.

Diffraction data collection was performed using an in-house XtaLAB

SynergyCustom (Rigaku) that was equipped with a HyPix-6000 detector and

MicroMax-007 HF X-ray generator with a wavelength of 1.54178 Å. The

diffraction data were indexed and integrated using CrysAlisPro ver. 1.171.40.14a

(Rigaku) and merged and scaled using Aimless in the CCP4 program package

(Winn *et al*, 2011).

Structure determination

The structure of the TEX264 LIR^{S272D}-GABARAP fusion protein was determined

by the molecular replacement method using the Phenix program (Liebschner *et*

al, 2019). The crystal structure of GABARAP (PDBID: 1GNU) was used as a

search model. Crystallographic refinement was performed using the Phenix and

COOT programs (Emsley *et al*, 2010; Liebschner *et al.*, 2019). The structure of

the GABARAP-TEX264 LIR$^{p\text{-}S271,272}$ complex was determined by molecular

replacement using MrBUMP (Keegan & Winn, 2008). The crystal structure of

GABARAP complexed with AnkB LIR (PDB 5YIR) was used as a search model.

Crystallographic refinement was performed using REFMAC5 (Murshudov *et al*,

2011), Phenix, and COOT programs. All structural images in the manuscript

were prepared with PyMOL (The PyMOL Molecular Graphics System, Version

2.0 Schrödinger; LLC.).

ITC

The peptides used in ITC are listed in Fig 4A. ITC experiments were performed

using a Microcal iTC200 calorimeter (Malvern Panalytical), with stirring at

1,000 rpm at 25°C. TEX264 peptides were prepared as injection samples from

the syringe. The concentration of TEX264 peptides was 0.5 mM. In total, the cell

was filled with 200 µL LC3B and GABARAP with a 10-times lower concentration

than the syringe samples. The titration involved 18 injections of 2 µL syringe

sample at intervals of 120 s into a cell after one injection of 0.4 µL syringe

sample. The datasets obtained from the titration of the same syringe sample to

the cell filled with buffer were used as reference data for subtraction of the heat

of the dilution. MicroCal Origin 7.0 software was used for data analysis. Thermal

measurement of the first injection of the syringe samples was removed from the

analysis. Thermal titration data were fitted to a single-site binding model, which

determines the thermodynamic parameters enthalpy (ΔH), dissociation constant

(K_d), and stoichiometry of binding (N). When the fitting was not convergent due

to weak interactions, N was fixed to 1.0 or that of the control experiment. The

error of each parameter shows the fitting error.

Doxycycline treatment

HeLa cells stably expressing the ER-phagy probe were cultured with 0.5 µg/mL

doxycycline (D3447; Sigma-Aldrich) for 24 h. Thereafter, cells were washed with

PBS twice and incubated in DMEM (D6546; Sigma-Aldrich) supplemented with

10% fetal bovine serum (S1820-500; Biowest) and 2 mM L-glutamine

(25030-081; Gibco), or amino acid-free DMEM (048-33575; Wako Pure

Chemical Industries) without serum followed by confocal microscopy or

biochemical analysis.

Quantification and statistical analysis

Two groups of data were evaluated by an unpaired two-tailed Student's *t*-test,

and multiple comparison tests were performed by one-way analysis of variance

(ANOVA) followed by Tukey's multiple comparison test. The data were assumed

to have a normal distribution, but this was not formally tested.

Data availability

Coordinates and structure factors for the GABARAP-TEX264 LIR[p-S271,272]

complex and GABARAP fused to TEX264 LIR[S272D] have been deposited in the

PDB under accession codes 7VEC and 7VED, respectively.

Acknowledgments

We thank Toshio Kitamura for providing pMXs-IP, Shoji Yamaoka for pMRXIP,

Teruhito Yasui for pCG-VSV-G and pCG-gag-pol, Robert A. Weinberg for

pCMV-VSV-G, Didier Trono for psPAX2, Toshiya Endo for the anti-RFP antibody,

Emi Izumi and Kaori Suzuki for technical assistance with cellular studies and

Yuki Ishii for help with protein preparation and crystallization. This study was

supported by Exploratory Research for Advanced Technology (ERATO; grant

number JPMJER1702 to N.M; JPMJER2001 to H.R.U.) and Core Research for

Evolutional Science and Technology (CREST; grant number JPMJCR20E3 to

N.N.N.) from the Japan Science and Technology Agency, by JSPS KAKENHI

(grant number 19H05707 to N.N.N.; 17K18339 and 20K06637 to A.Y.;

JP25221004 to H.R.U.), by HFSP Research Grant Program RGP0019/2018 to

H.R.U., and by the Takeda Science Foundation (to N.N.N. and A.Y.).

Author contributions

H.C. and N.M. designed the project. H.C. performed most cell biology

experiments. K.L.O. and H.R.U. performed mass spectrometry. A.Y. and N.N.N.

performed ITC analysis and crystal structural analysis. H.C., N.N.N., and N.M.

wrote the manuscript. All authors analyzed and discussed the results and

commented on the manuscript.

Conflict of Interest

References

An H, Ordureau A, Paulo JA, Shoemaker CJ, Denic V, Harper JW (2019)
TEX264 Is an Endoplasmic Reticulum-Resident ATG8-Interacting Protein
Critical for ER Remodeling during Nutrient Stress. *Mol Cell* 74:
891-908.e810

Anding AL, Baehrecke EH (2017) Cleaning House: Selective Autophagy of
Organelles. *Dev Cell* 41: 10-22

Chino H, Hatta T, Natsume T, Mizushima N (2019) Intrinsically disordered protein
TEX264 mediates ER-phagy. *Mol Cell* 74: 909-921.e906

Chino H, Mizushima N (2020) ER-Phagy: Quality Control and Turnover of
Endoplasmic Reticulum. *Trends Cell Biol* 30: 384-398

Cooper JA, Sefton BM, Hunter T (1983) Detection and quantification of
phosphotyrosine in proteins. *Methods Enzymol* 99: 387-402

Dikic I, Elazar Z (2018) Mechanism and medical implications of mammalian
autophagy. *Nat Rev Mol Cell Biol* 19: 349-364

Emsley P, Lohkamp B, Scott WG, Cowtan K (2010) Features and development
of Coot. *Acta crystallogr D Biol Crystallogr* 66: 486-501

Fielden J, Wiseman K, Torrecilla I, Li S, Hume S, Chiang SC, Ruggiano A,
Narayan Singh A, Freire R, Hassanieh S *et al* (2020) TEX264 coordinates
p97- and SPRTN-mediated resolution of topoisomerase 1-DNA adducts.
Nat Commun 11: 1274

Gatica D, Lahiri V, Klionsky DJ (2018) Cargo recognition and degradation by
selective autophagy. *Nat Cell Biol* 20: 233-242

Gomes LC, Dikic I (2014) Autophagy in antimicrobial immunity. *Mol Cell* 54:
224-233

Hübner CA, Dikic I (2020) ER-phagy and human diseases. *Cell Death Differ* 27:
833-842

Johansen T, Lamark T (2020) Selective Autophagy: ATG8 Family Proteins, LIR
Motifs and Cargo Receptors. *J Mol Biol* 432: 80-103

Kabsch W (2010) Integration, scaling, space-group assignment and
post-refinement. *Acta crystallogr D Biol Crystallogr* 66: 133-144

Kaizuka T, Morishita H, Hama Y, Tsukamoto S, Matsui T, Toyota Y, Kodama A,
Ishihara T, Mizushima T, Mizushima N (2016) An Autophagic Flux Probe
that Releases an Internal Control. *Mol Cell* 64: 835-849

Keegan RM, Winn MD (2008) MrBUMP: an automated pipeline for molecular
replacement. *Acta crystallogr D Biol Crystallogr* 64: 119-124

Kim J, Kim YC, Fang C, Russell RC, Kim JH, Fan W, Liu R, Zhong Q, Guan KL
(2013) Differential regulation of distinct Vps34 complexes by AMPK in
nutrient stress and autophagy. *Cell* 152: 290-303
Kirkin V, Rogov VV (2019) A Diversity of Selective Autophagy Receptors
Determines the Specificity of the Autophagy Pathway. *Mol Cell* 76:
268-285
Kitamura T, Koshino Y, Shibata F, Oki T, Nakajima H, Nosaka T, Kumagai H
(2003) Retrovirus-mediated gene transfer and expression cloning: powerful
tools in functional genomics. *Exp Hematol* 31: 1007-1014
Li J, Zhu R, Chen K, Zheng H, Zhao H, Yuan C, Zhang H, Wang C, Zhang M
(2018) Potent and specific Atg8-targeting autophagy inhibitory peptides
from giant ankyrins. *Nat Chem Biol* 14: 778-787
Liebschner D, Afonine PV, Baker ML, Bunkóczi G, Chen VB, Croll TI, Hintze B,
Hung LW, Jain S, McCoy AJ *et al* (2019) Macromolecular structure
determination using X-rays, neutrons and electrons: recent developments
in Phenix. *Acta crystallogr D Struct Biol* 75: 861-877
Meggio F, Marin O, Pinna LA (1994) Substrate specificity of protein kinase CK2.
Cell Mol Biol Res 40: 401-409
Mizushima N (2020) The ATG conjugation systems in autophagy. *Curr Opin
Cell Biol* 63: 1-10
Mizushima N, Levine B (2020) Autophagy in Human Diseases. *N Engl J Med*
383: 1564-1576
Mochida K, Yamasaki A, Matoba K, Kirisako H, Noda NN, Nakatogawa H (2020)
Super-assembly of ER-phagy receptor Atg40 induces local ER remodeling
at contacts with forming autophagosomal membranes. *Nat Commun* 11:
3306
Morita K, Hama Y, Izume T, Tamura N, Ueno T, Yamashita Y, Sakamaki Y,
Mimura K, Morishita H, Shihoya W *et al* (2018) Genome-wide CRISPR
screen identifies TMEM41B as a gene required for autophagosome
formation. *J Cell Biol* 217: 3817-3828
Murshudov GN, Skubák P, Lebedev AA, Pannu NS, Steiner RA, Nicholls RA,
Winn MD, Long F, Vagin AA (2011) REFMAC5 for the refinement of
macromolecular crystal structures. *Acta crystallogr D Biol Crystallogr*
67: 355-367
Nakatogawa H (2020) Mechanisms governing autophagosome biogenesis. *Nat
Rev Mol Cell Biol* 21: 439-458

Noda NN, Ohsumi Y, Inagaki F (2010) Atg8-family interacting motif crucial for
selective autophagy. *FEBS letters* 584: 1379-1385

Palikaras K, Lionaki E, Tavernarakis N (2018) Mechanisms of mitophagy in
cellular homeostasis, physiology and pathology. *Nat Cell Biol* 20:
1013-1022

Pinna LA (2002) Protein kinase CK2: a challenge to canons. *J Cell Sci* 115:
3873-3878

Rappsilber J, Mann M, Ishihama Y (2007) Protocol for micro-purification,
enrichment, pre-fractionation and storage of peptides for proteomics using
StageTips. *Nat Protoc* 2: 1896-1906

Rekha N, Srinivasan N (2003) Structural basis of regulation and substrate
specificity of protein kinase CK2 deduced from the modeling of
protein-protein interactions. *BMC Struc Biol* 3: 4

Rogov VV, Suzuki H, Fiskin E, Wild P, Kniss A, Rozenknop A, Kato R, Kawasaki
M, McEwan DG, Löhr F *et al* (2013) Structural basis for
phosphorylation-triggered autophagic clearance of Salmonella. *Biocheml*
J 454: 459-466

Rogov VV, Suzuki H, Marinković M, Lang V, Kato R, Kawasaki M, Buljubašić M,
Šprung M, Rogova N, Wakatsuki S *et al* (2017) Phosphorylation of the
mitochondrial autophagy receptor Nix enhances its interaction with LC3
proteins. *Sci Rep* 7: 1131

Rozenknop A, Rogov VV, Rogova NY, Löhr F, Güntert P, Dikic I, Dötsch V (2011)
Characterization of the interaction of GABARAPL-1 with the LIR motif of
NBR1. *J Mol Biol* 410: 477-487

Sakurai S, Tomita T, Shimizu T, Ohto U (2017) The crystal structure of mouse
LC3B in complex with the FYCO1 LIR reveals the importance of the
flanking region of the LIR motif. *Acta Crystallogr F Struct Biology*
Commun 73: 130-137

Søreng K, Neufeld TP, Simonsen A (2018) Membrane Trafficking in Autophagy.
Int Rev Cell Mol Biol 336: 1-92

Sou YS, Waguri S, Iwata J, Ueno T, Fujimura T, Hara T, Sawada N, Yamada A,
Mizushima N, Uchiyama Y *et al* (2008) The Atg8 conjugation system is
indispensable for proper development of autophagic isolation membranes
in mice. *Mol Biol Cell* 19: 4762-4775

Strickfaden SC, Winters MJ, Ben-Ari G, Lamson RE, Tyers M, Pryciak PM
(2007) A mechanism for cell-cycle regulation of MAP kinase signaling in a

838   yeast differentiation pathway. ***Cell*** 128: 519-531

839 Suzuki H, Tabata K, Morita E, Kawasaki M, Kato R, Dobson RC, Yoshimori T,
840   Wakatsuki S (2014) Structural basis of the autophagy-related LC3/Atg13
841   LIR complex: recognition and interaction mechanism. ***Structure*** 22: 47-58

842 Thorsness PE, Koshland DE, Jr (1987) Inactivation of isocitrate dehydrogenase
843   by phosphorylation is mediated by the negative charge of the phosphate. ***J***
844   ***Biol Chem*** 262: 10422-10425

845 Tsuboyama K, Koyama-Honda I, Sakamaki Y, Koike M, Morishita H, Mizushima
846   N (2016) The ATG conjugation systems are important for degradation of
847   the inner autophagosomal membrane. ***Science*** 354: 1036-1041

848 Wild P, Farhan H, McEwan DG, Wagner S, Rogov VV, Brady NR, Richter B,
849   Korac J, Waidmann O, Choudhary C *et al* (2011) Phosphorylation of the
850   autophagy receptor optineurin restricts Salmonella growth. ***Science*** 333:
851   228-233

852 Winn MD, Ballard CC, Cowtan KD, Dodson EJ, Emsley P, Evans PR, Keegan
853   RM, Krissinel EB, Leslie AG, McCoy A *et al* (2011) Overview of the CCP4
854   suite and current developments. ***Acta crystallogr D Biol crystallogr*** 67:
855   235-242

856 Wirth M, Mouilleron S, Zhang W, Sjøttem E, Princely Abudu Y, Jain A, Lauritz
857   Olsvik H, Bruun JA, Razi M, Jefferies HBJ *et al* (2021) Phosphorylation of
858   the LIR Domain of SCOC Modulates ATG8 Binding Affinity and Specificity.
859   ***J Mol Biol*** 433: 166987

860 Zhou J, Tien AC, Alberta JA, Ficarro SB, Griveau A, Sun Y, Deshpande JS, Card
861   JD, Morgan-Smith M, Michowski W *et al* (2017) A Sequentially Priming
862   Phosphorylation Cascade Activates the Gliomagenic Transcription Factor
863   Olig2. ***Cell Rep*** 18: 3167-3177

864 Zhu Y, Massen S, Terenzio M, Lang V, Chen-Lindner S, Eils R, Novak I, Dikic I,
865   Hamacher-Brady A, Brady NR (2013) Modulation of serines 17 and 24 in
866   the LC3-interacting region of Bnip3 determines pro-survival mitophagy
867   versus apoptosis. ***J Biol Chem*** 288: 1099-1113

Figure legends

Figure 1. Phosphorylation of Ser271 and Ser272 upstream of the TEX264 LIR is required for ATG8 binding and autophagosomal localization

(A) Alignment of the LIR motif of TEX264 homologs in vertebrates. Asterisks show the phosphorylation sites identified by mass spectrometry.

(B) HEK293T cells transiently expressing FLAG and FLAG-tagged TEX264 were subjected to immunoprecipitation (IP). Inputs (20% of the lysates) and immunoprecipitates (from 80% of the lysates) were analyzed by immunoblotting using the indicated antibodies.

(C) HEK293T cells transiently expressing WT or mutated TEX264-FLAG were subjected to immunoprecipitation with an anti-FLAG antibody and detected with the indicated antibodies.

(D and E) MEFs stably expressing TEX264-GFP or its mutant were cultured in starvation media and immunostained with an anti-LC3 antibody. Bars: 10 μm and 1 μm (insets) (D). Quantification of the number of TEX264 puncta per cell. Solid bars indicate the medians, boxes the interquartile range (25th to 75th percentile), and whiskers the 0th to 100th percentile. Differences were statistically analyzed by one-way ANOVA and Tukey's multiple comparison test. Data were collected from 48 cells for each cell type (E).

(F and G) WT and TEX264-KO HeLa cells (with or without the indicated TEX264 mutants with a C-terminal FLAG tag) stably expressing the ER-phagy reporter (ss-RFP-GFP-KDEL) were cultured in the presence of doxycycline for 24 h to induce the reporter. After doxycycline was removed, cells were cultured in starvation medium lacking amino acids and serum for 9 h (F). The band intensities of RFP and RFP-GFP were quantified and the ratio of RFP:RFP-GFP (normalized to WT) is shown. Data represent the mean ± standard error of the mean (SEM) of three independent experiments. Differences were statistically analyzed by one-way ANOVA and Tukey's multiple comparison test (G).

Figure 2. Phosphorylated TEX264 localizes to autophagosomes

(A and B) MEFs stably expressing TEX264-GFP were cultured in starvation

media with or without bafilomycin A_1 (Baf A_1) and immunostained with

anti-pTEX264 and anti-LC3 antibodies. Bars: 10 µm and 1 µm (insets) (A).

Quantification of the number of pTEX264 puncta per cell. Solid bars indicate the

medians, boxes the interquartile range (25th to 75th percentile), and whiskers

the 0th to 100th percentile. Differences were statistically analyzed by one-way

ANOVA and Tukey's multiple comparison test. Data were collected from 35 cells

for each cell type (B).

(C) HeLa cells were treated with 250 nM Torin 1 and Baf A_1 for 16 h, and cell

homogenates were subjected to an OptiPrep membrane flotation analysis.

(D, E and F) WT, ATG3-KO, and FIP200-KO HeLa cells were cultured in

starvation media lacking amino acids and serum with or without Baf A_1 for 1 or 5

h. Cell lysates were analyzed by immunoblotting using the indicated antibodies

(D). Relative changes of the ratio of band intensities of phosphorylated TEX264

to HSP90 and total TEX264 during starvation are shown (E and F). Data

represent the mean ± SEM of three independent experiments.

Figure 3. CK2 is important for starvation-induced ER-phagy through

TEX264 phosphorylation

(A) HeLa cells were cultured with 20 nM CK1 inhibitor (D4476) or CK2 inhibitor

(CX4945) for 3 h, followed by starvation (with the inhibitors) for 3 h. Cell lysates

were analyzed by immunoblotting using the indicated antibodies.

(B) Recombinant TEX264-FLAG was incubated with or without CK2 and the CK2

inhibitor (CX4945) in the presence of 200 µM ATP for 1 h.

(C) HEK293T cells transiently expressing TEX264-FLAG and one of the three

Halo-CK2A isoforms or Halo-CK2B were subjected to immunoprecipitation with

an anti-FLAG antibody and detected with anti-Halo and anti-FLAG antibodies.

(D and E) HEK293T cells transiently expressing WT or TEX264-FLAG mutants

were subjected to immunoprecipitation (IP). Inputs (20% of the lysates) and

immunoprecipitates (from 80% of the lysates) were analyzed by immunoblotting

using the indicated antibodies (D). The band intensities in the IP fractions were

quantified, and the phospho-serine:FLAG ratio was calculated. Data represent the mean ± SEM of five independent experiments. Differences were statistically analyzed by one-way ANOVA and Tukey's multiple comparison test (E).

(F and G) MEFs stably expressing TEX264-GFP were cultured with the CK2 inhibitor for 3 h followed by starvation (with the CK2 inhibitor) and immunostained with an anti-LC3 antibody. Scale bars represent 10 μm and 1 μm (insets) (F). Quantification of the number of TEX264 puncta per cell. Solid bars indicate the medians, boxes the interquartile range (25th to 75th percentile), and whiskers the 0th to 100th percentile. Differences were statistically analyzed by an unpaired two-tailed Student's t-test. Data were collected from 85 cells for each cell type (G).

(H and I) WT HeLa cells stably expressing the ER-phagy reporter were cultured in the presence of doxycycline for 24 h to induce the reporter. After doxycycline was removed, cells were cultured with 20 nM CK2 inhibitor for 3 h followed by starvation (with the CK2 inhibitor) for 6 h (H). The band intensities of RFP and RFP-GFP were quantified and the ratio of RFP:RFP-GFP (normalized to WT) is shown. Data represent the mean ± SEM of six independent experiments. Differences were statistically analyzed by an unpaired two-tailed Student's t-test (I).

(J) Model of the CK2-mediated phosphorylation of TEX264 in ER-phagy.

Figure 4. Phosphorylation of TEX264 LIR increases its affinity to ATG8s

(A) Binding affinity of TEX264 LIR peptides (TEX264$^{269-278}$) to LC3 and GABARAP was measured by ITC. The error bars show the fitting error.

(B) HEK293T cells transiently expressing WT or mutated TEX264-FLAG were subjected to immunoprecipitation with an anti-FLAG antibody and detected with the indicated antibodies.

(C and D) MEFs stably expressing TEX264-GFP or its mutants were cultured in starvation media and immunostained with an anti-LC3 antibody. Bars: 10 μm and 5 μm (insets) (C). The intensity of the GFP signal under starvation conditions was quantified at TEX264-GFP puncta and the ER. The puncta:ER

signal ratio was calculated. Solid bars indicate the medians, boxes the

interquartile range (25th to 75th percentile), and whiskers the 0th to 100th

percentile. Differences were statistically analyzed by one-way ANOVA and

Tukey's multiple comparison test. Data were collected from 70 puncta for each

cell type (D).

(E and F) WT and TEX264-KO (expressing WT or TEX264-FLAG mutants) HeLa

cells stably expressing the ER-phagy reporter were cultured in the presence of

doxycycline for 24 h to induce the reporter. After doxycycline was removed, the

cells were cultured in starvation medium lacking amino acids and serum for 9 h

(E). The band intensities of RFP and RFP-GFP were quantified and the ratio of

RFP:RFP-GFP (normalized to WT) is shown. Data represent the mean ± SEM of

four independent experiments. Differences were statistically analyzed by

one-way ANOVA and Tukey's multiple comparison test (F).

Figure 5. Phosphorylation of the LIR in TEX264 produces hydrogen bonds to interact with GABARAP

(A) Crystal structure of GABARAP complexed with TEX264$^{269–278}$ peptide

containing p-Ser271 and p-Ser272. hp1 and hp2 indicate hydrophobic pockets 1

and 2, respectively. TEX264 peptide is colored green.

(B) Crystal structure of GABARAP fused with TEX264$^{271–281}$ peptide with the

S272D mutation. TEX264 peptide is colored green.

(C and D) Close-up view of the interaction between GABARAP and the

upstream region of LIR$^{p\text{-}S271,p\text{-}S272}$ (C) or LIR^{S272D} (D). Broken lines indicate

possible hydrogen bonds between GABARAP and LIR (colored back) and within

GABARAP (colored gray).

(E) Summary of the distances (Å) of possible hydrogen bonds mediated by Tyr5,

His9, and Lys48 of GABARAP with TEX264 LIR.

(F) Structural comparison between the GABARAP-TEX264 LIR$^{p\text{-}S271,p\text{-}S272}$

complex and GABARAPL1-SCOC LIR complex (PDB 7AA7). Possible hydrogen

bonds are shown as in C and D. F_0 and p-S_{-2} indicate Phe located at position Θ_0

and phosphorylated Ser located at position X_{-2}, respectively.

Expanded View Figure legends

Figure EV1. Phosphorylated TEX264 can be detected by an anti-phosphoTEX264 antibody

(A) A representative MS^2 spectrum of an LIR-containing peptide of TEX264 G280K. B-ions (red), Y-ions (blue) and their neutral losses of water and ammonia are highlighted.

(B and C) MEFs stably expressing TEX264-GFP or its S266A mutant were cultured in starvation media and immunostained with an anti-LC3 antibody. Bars: 10 μm and 2 μm (insets) (B). Quantification of the number of TEX264 puncta per cell. Solid bars indicate the medians, boxes the interquartile range (25th to 75th percentile), and whiskers the 0th to 100th percentile. Differences were statistically analyzed by student t-test. Data were collected from 35 cells for each cell type (C).

(D) Immunoblotting of the lysates of WT and TEX264-KO HeLa cells stably expressing WT TEX264 or the TEX264^{S271A,S272A} mutant using the indicated antibodies.

(E) HeLa cells stably expressing TEX264-FLAG were cultured in starvation media lacking amino acids and serum with or without bafilomycin A_1 for 2 h. Cell lysates were subjected to immunoprecipitation and treated with or without lambda and alkaline phosphatases for 1 h. Inputs (20% of the lysates) and immunoprecipitates (from 80% of the lysates) were analyzed by immunoblotting using the indicated antibodies.

Figure EV2. CK2 inhibitor does not affect starvation-induced autophagy

HEK293T cells stably expressing the GFP-LC3-RFP autophagic flux reporter were cultured with 10 μM or 20 μM CK2 inhibitor for 3 h followed by starvation (with the same concentration of inhibitor) for 6 h and analyzed by flow cytometry. The GFP:RFP ratio was quantified. Data represent the mean ± SEM of four independent experiments. Differences were statistically analyzed by one-way ANOVA and Tukey's multiple comparison test.

1026

1027 **Figure EV3. Phosphorylation of TEX264 LIR increases its affinity to ATG8s**

1028 Binding affinity of TEX264 LIR peptides (TEX264$^{269–278}$ (A) or TEX264$^{266–278}$ (B))

1029 containing the indicated replacements to LC3B or GABARAP was measured by

1030 ITC.

1031

Figure EV4. Multiple copies of GABARAP-TEX264 LIR complexes observed

in the crystal.

(A) Ribbon representation of 12 copies of GABARAP observed in the

asymmetric unit of the crystal (left) and their superimposition (right). The

side-chains of p-Ser271, p-Ser272, Phe273, and Leu276 are shown with a stick

model.

(B) Ribbon representation of two copies of GABARAP observed in the

asymmetric unit and two symmetrically related GABARAPs in the crystal (left)

and their superimposition (right). The side-chains of Asp272, Phe273, and

Leu276 are shown with a stick model.

(C and D) 2Fo-Fc map (C) and simulated annealing Fo-Fc omit map (D) of the

TEX264 LIR peptides bound to GABARAP together with the structural model,

where carbon, nitrogen, oxygen, and sulfur atoms are colored yellow, blue, red,

and orange, respectively. Counter levels are indicated at each image.

(E and F) HEK293T cells transiently expressing WT or FLAG-GABARAP

mutants were subjected to immunoprecipitation (IP). Inputs (20% of the lysates)

and immunoprecipitates (from 80% of the lysates) were analyzed by

immunoblotting using the indicated antibodies (F). The band intensities in the IP

fractions were quantified. Data represent the mean ± SEM of three independent

experiments. Differences were statistically analyzed by one-way ANOVA and

Tukey's multiple comparison test (G).

1053

Figure EV5. Structural comparison of GABARAP/GABARAPL1-LIR

complexes deposited in the PDB.

Each structural model is shown in the same orientation as that in Fig 5C.

Possible hydrogen bonds are shown as in Fig 5C. Green letters below the PDB

code indicate the amino acid sequence in the upstream region of each LIR.

Numbers in subscript indicate positions in the LIR sequence.

1060

1061

1062

1063

Figure 1. Phosphorylation of Ser271 and Ser272 upstream of the TEX264 LIR are required for ATG8 binding and autophagosomal localization

(A) Alignment of the LIR motif of TEX264 homologs in vertebrates. Asterisks show phosphorylation sites identified by mass spectrometry.

(B) HEK293T cells transiently expressing FLAG and FLAG-tagged TEX264 were subjected to immunoprecipitation (IP). Inputs (20% of the lysates) and immunoprecipitates (from 80% of the lysates) were analyzed by immunoblotting using the indicated antibodies.

(C) HEK293T cells transiently expressing WT or mutated TEX264-FLAG were subjected to immunoprecipitation with anti-FLAG antibody and detected with the indicated antibodies.

(D and E) Mouse embryonic fibroblasts (MEFs) stably expressing TEX264-GFP or its mutant were cultured in starvation media and immunostained with anti-LC3 antibody. Bars: 10 μm and 1 μm (insets) (D). Quantification of the number of TEX264 puncta per cell. Solid bars indicate the medians, boxes the interquartile range (25th to 75th percentile), and whiskers the 0th to 100th percentile. Differences were statistically analyzed by one-way ANOVA and Tukey's multiple comparison test. Data were collected from 48 cells for each cell type (E).

(F and G) WT and TEX264-KO HeLa cells (with or without the indicated TEX264 mutants tagged with FLAG at the C terminus) stably expressing the ER-phagy reporter (ss-RFP-GFP-KDEL) were cultured in the presence of doxycycline for 24 h to induce the reporter. After doxycycline was removed, cells were cultured in starvation medium lacking amino acids and serum for 9 h (F). The band intensities of RFP and RFP-GFP were quantified and the ratio of RFP:RFP-GFP (normalized to WT) is shown. Data represent the mean ± SEM of three independent experiments. Differences were statistically analyzed by one-way ANOVA and Tukey's multiple comparison test (G).

Figure 2. Phosphorylated TEX264 localizes to autophagosome

(A and B) MEFs stably expressing TEX264-GFP were cultured in starvation media with or without bafilomycin A_1 (Baf A_1) and immunostained with anti-pTEX264 and anti-LC3 antibodies. Bars: 10 μm and 1 μm (insets) (A). Quantification of the number of pTEX264 puncta per cell. Solid bars indicate the medians, boxes the interquartile range (25th to 75th percentile), and whiskers the 0th to 100th percentile. Differences were statistically analyzed by one-way ANOVA and Tukey's multiple comparison test. Data were collected from 35 cells for each cell type (B).
(C) HeLa cells were treated with 250 nM Torin 1 and Baf.A_1 for 16 h, and cell homogenates were subjected to OptiPrep floatation analysis.
(D, E and F) WT, ATG3-KO, and FIP200-KO HeLa cells were cultured in starvation media lacking amino acids and serum with or without Baf A_1 for 1 or 5 h. Cell lysates were analyzed by immunoblotting using the indicated antibodies (D). Relative changes of the ratio of band intensities of phosphorylated TEX264 to HSP90 and total TEX264 during starvation are shown (E and F). Data represent the mean ± SEM of three independent experiments.

Figure 3. Casein kinase 2 is important for starvation-induced ER-phagy through phosphorylation of TEX264

(A) HeLa cells were cultured with 20 nM CK1 inhibitor (D4476) or CK2 inhibitor (CX4945) for 3 h, followed by starvation (with the inhibitors) for 3 h. Cell lysates were analyzed by immunoblotting using the indicated antibodies.

(B) Recombinant TEX264-FLAG was incubated with or without CK2 ± CK2 inhibitor (CX4945) in addition with 200μM ATP for 1 h. Reaction solutions were analyzed by immunoblotting using the indicated antibodies.

(C) HEK293T cells transiently expressing TEX264-FLAG and Halo-CK2A or Halo-CK2B were subjected to immunoprecipitation with anti-FLAG antibody and detected with anti-Halo and anti-FLAG antibodies.

(D and E) HEK293T cells transiently expressing WT or TEX264-FLAG mutants were subjected to immunoprecipitation (IP). Inputs (20% of the lysates) and immunoprecipitates (from 80% of the lysates) were analyzed by immunoblotting using the indicated antibodies (D). The band intensities in IP fractions were quantified, and the phospho-serine: FLAG ratio was calculated. Data represent the mean ± SEM of five independent experiments (E).

 (F and G) MEFs stably expressing TEX264-GFP were cultured with CK2 inhibitor for 3 h followed by starvation (with the CK2 inhibitor) and immunostained with anti-LC3 antibody. Scale bars represent 10 μm and 1 μm (insets) (F). Quantification of the number of TEX264 puncta per cell. Solid bars indicate the medians, boxes the interquartile range (25th to 75th percentile), and whiskers the 0th to 100th percentile. Differences were statistically analyzed by unpaired two-tailed Student's t-test. Data were collected from 85 cells for each cell type (G).

(H and I) WT HeLa cells stably expressing the ER-phagy reporter were cultured in the presence of doxycycline for 24 h to induce the reporter. After doxycycline was removed, cells were cultured with 20 nM CK2 inhibitor for 3 h followed by starvation (with the CK2 inhibitor) for 6 h (H). The band intensities of RFP and RFP-GFP were quantified and the ratio of RFP:RFP-GFP (normalized to WT) is shown. Data represent the mean ± SEM of six independent experiments. Differences were statistically analyzed by an unpaired two-tailed Student's t-test (I).

(J) Model of CK2-mediated phosphorylation of TEX264 in ER-phagy.

Figure 4. Phosphorylation of TEX264 LIR increases the affinity to ATG8s

(A) Binding affinity of TEX264 LIR peptides (TEX264$^{269-278}$) to LC3 and GABARAP was measured by isothermal titration calorimetry. The error bars show the fitting error.

(B) HEK293T cells transiently expressing WT or mutated TEX264-FLAG were subjected to immunoprecipitation with anti-FLAG antibody and detected with the indicated antibodies.

(C and D) MEFs stably expressing TEX264-GFP or its mutants were cultured in starvation media and immunostained with anti-LC3 antibody. Bars: 10 μm and 5 μm (insets) (C). The intensity of GFP signal under starvation conditions was quantified at TEX264-GFP puncta and the ER. The puncta:ER signal ratio was calculated. Solid bars indicate the medians, boxes the interquartile range (25th to 75th percentile), and whiskers the 0th to 100th percentile. Differences were statistically analyzed by one-way ANOVA and Tukey's multiple comparison test. Data were collected from 70 puncta for each cell type (D).

(E and F) WT and TEX264-KO (expressing WT or TEX264-FLAG mutants) HeLa cells stably expressing the ER-phagy reporter were cultured in the presence of doxycycline for 24 h to induce the reporter. After doxycycline was removed, cells were cultured in starvation medium lacking amino acids and serum for 9 h (E). The band intensities of RFP and RFP-GFP were quantified and the ratio of RFP:RFP-GFP (normalized to WT) is shown. Data represent the mean ± SEM of four independent experiments. Differences were statistically analyzed by one-way ANOVA and Tukey's multiple comparison test (F).

Figure 5. Phosphorylation of the LIR in TEX264 produces hydrogen bonds to interact with GABARAP

(A) Crystal structure of GABARAP complexed with TEX264[269-278] peptide containing p-Ser271 and p-Ser272. hp1 and hp2 indicate hydrophobic pockets 1 and 2, respectively. TEX264 peptide is colored green.

(B) Crystal structure of GABARAP fused with TEX264[271-281] peptide with S272D mutation. TEX264 peptide is colored green.

(C and D) Close-up view of the interaction between GABARAP and the upstream region of LIR [p-S271,p-S272] (C) or LIR[S272D] (D). Broken lines indicate possible hydrogen bonds between GABARAP and LIR (colored back) and within GABARAP (colored gray).

(E) Summary of the distances (Å) of possible hydrogen bonds mediated by Tyr5, His9, and Lys48 of GABARAP with TEX264 LIR.

(F) Structural comparison between GABARAP-TEX264 LIR [p-S271,p-S272] complex and GABARAPL1-SCOC LIR complex (PDB 7AA7). Possible hydrogen bonds are shown as in C and D. F_0 and p-S_{-2} indicate Phe located at position Θ_0 and phosphorylated Ser located at position X_{-2}, respectively.

A

TEX264^{G280K} 259- S E H S Y S E pS G A pS G pS pS F E E L D L E K -280

y_8 y_7 y_6 y_5 y_4 y_3 y_2 y_1

b_2 b_3 b_5 b_6

Precursor ion = 898.96515 m/z (z = +3); [MH+] = 2694.88089 (Theoretical [MH+] = 2694.88106)

Expanded View Figure 1. Phosphorylated TEX264 can be detected by anti-phosphoTEX264 antibody

(A) A representative MS^2 spectrum of an LIR-containing peptide of TEX264 G280K. B-ions (red), Y-ions (blue) and their neutral losses of water and ammonia are highlighted.

(B and C) MEFs stably expressing TEX264-GFP or its mutant were cultured in starvation media and immunostained with an anti-LC3 antibody. Bars: 10 μm and 2 μm (insets) (B). Quantification of the number of TEX264 puncta per cell. Solid bars indicate the medians, boxes the interquartile range (25th to 75th percentile), and whiskers the 0th to 100th percentile. Differences were statistically analyzed by student t-test. Data were collected from 35 cells for each cell type (C).

(D) Immunoblotting of the lysates of WT and TEX264-KO HeLa cells stably expressing WT TEX264 or the TEX264^{S271A,S272A} mutant using the indicated antibodies.

(E) HeLa cells stably expressing TEX264-FLAG were cultured in starvation media lacking amino acids and serum with or without bafilomycin A1 for 2 h. Cell lysates were subjected to immunoprecipitation and treated with or without lambda and alkaline phosphatases for 1 h. Inputs (20% of the lysates) and immunoprecipitates (from 80% of the lysates) were analyzed by immunoblotting using the indicated antibodies.

Expanded View Figure 2. The CK2 inhibitor does not affect starvation-induced autophagy

HEK293T cells stably expressing the GFP-LC3-RFP autophagic flux reporter were cultured with 10 μM or 20 μM CK2 inhibitor for 3 h followed by starvation (with the same inhibitors) for 6 h and analyzed by flow cytometry. The GFP/RFP ratio was quantified. Data represent the mean ± SEM of four independent experiments. Differences were statistically analyzed by one-way ANOVA and Tukey's multiple comparison test.

Expanded View Figure 3. Phosphorylation of TEX264 LIR increases the affinity to ATG8s

(A and B) Binding affinity of TEX264 LIR peptides (TEX264$^{269-278}$(A) or TEX264$^{266-278}$ (B)) containing the indicated replacements to LC3B or GABARAP was measured by isothermal titration calorimetry.

Expanded View Figure 4. Multiple copies of GABARAP-TEX264 LIR complexes observed in the crystal.

(A) Ribbon representation of 12 copies of GABARAP observed in the asymmetric unit of the crystal (left) and their superimposition (right). The side-chains of p-Ser271, p-Ser272, Phe273, and Leu276 are shown with a stick model.

(B) Ribbon representation of two copies of GABARAP observed in the asymmetric unit and two symmetrically related GABARAPs in the crystal (left) and their superimposition (right). The side-chains of Asp272, Phe273, and Leu276 are shown with a stick model.

(C and D) 2Fo-Fc map (C) and simulated annealing Fo-Fc omit map (D) of the TEX264 LIR peptides bound to GABARAP together with the structural model, where carbon, nitrogen, oxygen, and sulfur atoms are colored yellow, blue, red, and orange, respectively. Counter levels are indicated at each image.

(E and F) HEK293T cells transiently expressing WT or FLAG-GABARAP mutants were subjected to immunoprecipitation (IP). Inputs (20% of the lysates) and immunoprecipitates (from 80% of the lysates) were analyzed by immunoblotting using the indicated antibodies (E). The band intensities in the IP fractions were quantified, and the phospho-serine:FLAG ratio was calculated. Data represent the mean ± SEM of five independent experiments. Differences were statistically analyzed by one-way ANOVA and Tukey's multiple comparison test (F).

Expanded View Figure 5. Structural comparison of GABARAP/GABARAL1-LIR complexes deposited in the Protein Data Bank.

Each structural model is shown in the same orientation as that in Figure 5C. Possible hydrogen bonds are shown as in Figure 5C. Green letters below the PDB code indicate the amino acid sequence in the upstream region of each LIR. Numbers in subscript indicate positions in the LIR sequence.

Table EV1; Summary of ITC results

	LC3B		GABARAP	
Sequence	Kd (µM)	N (site)	Kd (µM)	N (site)
SGSSFEELDLY	21±6	0.641	47±7	1.0 (fixed)
SGAAFEELDLY	46±6	1.0 (fixed)	72±29	1.03
SGDDFEELDLY	7.6±1.2	0.641	2.1±0.6	0.728
SGEEFEELDLY	5.2±0.8	0.639	2.3±0.5	0.696
DGDDFEELDLY	6.8±1.3	0.690	2.2±0.4	0.721
EGEEFEELDLY	5.0±0.8	0.681	2.2±0.3	0.684
SGS[pS]FEELDLY	0.57±0.08	0.710	2.1±0.3	0.704
SG[pS][pS]FEELDLY	0.37±0.05	0.742	0.52±0.06	0.766
[pS]G[pS][pS]FEELDLY	0.25±0.03	0.730	0.25±0.04	0.780

Table EV2; Data collection and refinement statistics

	GABARAP-TEX264 LIR[p-S271,272]	GABARAP-TEX264 LIR[S272D] fusion
Data collection		
Space group	*F*222	*P*2₁2₁2₁
Cell dimensions		
a, *b*, *c* (Å)	200.48 201.38 211.82	38.36 82.63 102.43
Resolution (Å)	30.48 - 3.0 (3.11 - 3.00)*	41.31 - 2.02 (2.093 - 2.02)*
R_{merge}	0.368 (1.77)	0.0877 (0.826)
$I / \sigma I$	6.5 (1.2)	14.1 (1.7)
Completeness (%)	99.9 (100.0)	99.29 (93.92)
Redundancy	8.7 (8.0)	6.4 (6.2)
CC1/2	0.964 (0.608)	0.998 (0.696)
Refinement		
Resolution (Å)	30.48 - 3.0 (3.11 - 3.00)	41.31 - 2.02 (2.093 - 2.02)
No. reflections	41789 (4234)	21898 (1991)
R_{work} / R_{free}**	0.257 / 0.301	0.188 / 0.225
No. atoms		
GABARAP	11559	2134
TEX264 peptides	806	
Sulfate		15
Water		150
B-factors (Å²)		
GABARAP	33.00	41.04
TEX264 peptides	43.65	
Sulfate		72.96
Water		48.74
R.m.s. deviations		
Bond lengths (Å)	0.003	0.007
Bond angles (°)	0.68	0.94

*Values in parentheses are for highest-resolution shell.

**R_{work} and R_{free} correspond to R-factors calculated using 95% (used for refinement) and 5% (not used for refinement) reflections, respectively.