Discovery of Flaviviridae-derived endogenous viral elements in shrew genomes provide novel insights into *Pestivirus* ancient history

Li YQ[1], Bletsa M[1], Zisi Z[1], Boonen I[1], Gryseels S[1,2], Kafetzopoulou L[1], Webster J[3], Pybus OG[3], Van de Perre F[2], Li HT[4], Li YY[4], Li YC[4], Abramov A[5], Lymberakis P[6], Lemey P[1], Lequime S[1,7]*

Department of Microbiology, Immunology and Transplantation, KU Leuven, Rega Institute, KU Leuven, Leuven, Belgium

Evolutionary Ecology Group, University of Antwerp, Antwerp, Belgium

Department of Pathobiology and Population Science, Royal Veterinary College, University of London, London, UK

Marine College, Shandong University (Weihai), Weihai, China

Laboratory of Theriology, Zoological Institute of the Russian Academy of Sciences, Saint Petersburg, Russia

Natural History Museum of Crete, Greece

Cluster of Microbial Ecology, Groningen Institute for Evolutionary Life Sciences, University of Groningen, Groningen, The Netherlands.

* corresponding author: Sebastian Lequime (s.j.j.lequime@rug.nl)

Keywords: Endogenous viral element, pestivirus, *Flaviviridae*, *Crocidura*, host range, paleovirology

Abstract

As viral genomic imprints in host genomes, endogenous viral elements (EVEs) shed light on

the deep evolutionary history of viruses, ancestral host ranges, and ancient viral-host

interactions. In addition, they may provide crucial information for calibrating viral

evolutionary timescales. In this study, we conducted a comprehensive *in silico* screening of a

large dataset of available mammalian genomes for EVEs deriving from members of the viral

family *Flaviviridae*, an important group of viruses including well-known human pathogens.

We identified two novel pestivirus-like EVEs in the reference genome of the Indochinese

shrew (*Crocidura indochinensis*). Homologs of these novel EVEs were subsequently detected

in vivo by molecular detection and sequencing in 27 shrew species, including 26 species

representing a wide distribution within the Crocidurinae subfamily and one in the Soricinae

subfamily. Based on this wide distribution, we estimate that the integration event occurred

before the last common ancestor of the subfamily, about 10.8 million years ago, attesting to an

ancient origin of pestiviruses and *Flaviviridae* in general. Moreover, we provide the first

description of *Flaviviridae*-derived EVEs in mammals even though the family encompasses

numerous mammal-infecting members, including major human pathogens such as Zika,

dengue, or hepatitis C viruses. This also suggests that shrews were past and perhaps also current

natural reservoirs of pestiviruses. Taken together, our results expand the current known

Pestivirus host range and provide novel insight into the ancient evolutionary history of

pestiviruses and the *Flaviviridae* family in general.

1 – Introduction

Endogenous viral elements (EVEs) are integrations of partial or full-length viral genomic material into the host genome (Katzourakis and Gifford, 2010a). In addition to retroviruses, which incorporate their genomic sequences into their host genome as an essential part of their replication cycle, many eukaryotic viruses have been found endogenized in various hosts (Feschotte and Gilbert, 2012). These non-retroviruses can derive from dsDNA (Aswad and Katzourakis, 2014; Li and Li, 2015; Liu et al., 2020), ssDNA (Belyi et al., 2010a; Katzourakis and Gifford, 2010a; Kobayashi et al., 2019), dsRNA (Horie et al., 2010; Katzourakis and Gifford, 2010b; Liu et al., 2012), and even ssRNA viruses (Crochu et al., 2004; Flynn and Moreau, 2019; Horie et al., 2010; Lequime and Lambrechts, 2017). Non-retroviral RNA virus-derived EVEs arise from a conjunction of relatively rare events: i) production of DNA genomic material, using retrotransposon encoded reverse transcriptase (Horie, 2019; Horie et al., 2010), ii) integration in the host chromosome of germ-line cells, and iii) overcoming genetic drift and/or natural selection at the population until fixation (Aiewsakun and Katzourakis, 2015a; Holmes, 2011). EVEs thus reflect long-term and intimate interactions of viruses with their hosts, and their identification can reveal insights into past and present host distributions of viral genera and families. The detection of endogenous bornavirus-like elements in invertebrate genomes (Horie et al., 2013), for example, suggested a broader host range for bornaviruses than previously thought. The study of filovirus-like EVEs in some small mammals offer predictive value for further identifying filovirus reservoirs (Taylor et al., 2010). Similarly, the discovery of flavivirus-derived EVEs in *Anopheles* mosquito genomes supported the idea that *Anopheles* mosquitoes could also be the natural hosts of flaviviruses (Lequime and Lambrechts, 2017), as confirmed by other studies (Colmant et al., 2017; Öncü et al., 2018).

Aside from qualitative insights into host ranges of viruses, EVEs can also shed light on deep evolutionary histories of viruses. EVEs are significant traces of past virus-host interactions; unlike animals or plants, viruses do not leave physical fossil records, limiting our ability to study their deep evolutionary histories. EVEs could thus be considered as "genomic fossil records" that can help to unravel long-term evolutionary dynamics between hosts and viral families. The presence of EVE homologs in different host species hints at an integration event before speciation, here the time to the most recent common ancestor. It thus provides a minimum age estimate for the integration event, and therefore a minimum age for the existence of a specific viral taxonomic group (Aiewsakun and Katzourakis, 2015b). For example, EVEs derived from adeno-associated viruses appear to be orthologous in African and Asian elephants,

indicating an integration event more than 6 million years ago (Kobayashi et al., 2019).

Similarly, the discovery of abundant bornavirus-like EVEs across vertebrates reveals that

ancient bornaviral infections occurred over a timeframe of about 100 million years before

present (Kawasaki et al., 2021). These studies can also provide genetic fossil calibration points

for further gauging ancient viral timescales using phylogenetics (Feschotte and Gilbert, 2012).

Members of the *Flaviviridae* family are linear, positive-sense, single-stranded RNA viruses

currently classified in four recognized genera (*Flavivirus*, *Hepacivirus*, *Pestivirus*, and

Pegivirus). They encompass many significant pathogens, such as dengue, Zika, hepatitis C, or

bovine viral diarrhea virus. In addition to human and livestock infections, *Flaviviridae* viruses

have been detected in a broad range of hosts, e.g. non-human primates (Mares-Guia et al.,

2020), rodents (Bletsa et al., 2021), bats (Wu et al., 2018), birds (Strand et al., 2018), arthropods

(Shi et al., 2016), and fish (Hartlage, 2016; Soto et al., 2020). Divergence date estimates on a

genus-wide scale are relatively rare, with the oldest, for the *Flavivirus* genus, being estimated

at 200,000 years ago (Pettersson and Fiz-Palacios, 2014). Although *Flaviviridae* viruses have

been detected in various hosts, especially in small mammals (Bletsa et al., 2021; de Lamballerie

et al., 2002; Wu et al., 2020, 2018), current published studies have only identified *Flaviviridae*-

derived EVEs in arthropods, including mosquitos (Crochu et al., 2004; Lequime and

Lambrechts, 2017; Roiz et al., 2009), ticks (Maruyama et al., 2014), and crustaceans (Parry

and Asgari, 2019), and these are all related to the *Flavivirus* genus. A recent study also

identified *Flaviviridae*-derived EVEs in a wide variety of hosts, including various invertebrates

and fish (Bamford et al., 2021). Currently, however, convincing evidence for *Flaviviridae*-

derived EVEs in vertebrates remains lacking. Interestingly, potential integration has been

suggested in medaka fish (Belyi et al., 2010b) as well as in rabbit and hare genomes (Silva et

al., 2015, 2012). While this raises the hypothesis that *Flaviviridae* viruses have integrated in

vertebrate hosts, the origin of these specific genomic sequences remains inconclusive due to

their short size and low sequence similarity to the *Flaviviridae* (*Flavivirus* and *Hepacivirus*

respectively).

In this study, we explored *in silico* the presence of *Flaviviridae*-derived EVEs in a

comprehensive set of mammalian genomes, and we discovered two novel pesti-like EVEs in

the genome of the Indochinese shrew *Crocidura indochinensis*. We subsequently identified

and characterized homologs of these EVEs *in vivo* in 26 species of the Crocidurinae subfamily

and one member of the Soricinae subfamily, establishing the integration event at least 10.8

million years ago. Our results provide the first evidence for an ancient origin of pestiviruses
and also contribute to a better understanding of the evolutionary history of the *Flaviviridae*
family in general.
115
2 – Material and Methods
2.1 – *In silico* survey
2.1.1 – Data collection
To screen for *Flaviviridae*-like EVEs, 689 mammalian genomes (57 bats, 9 insectivores, 177
rodents, 101 nonhuman primates, 207 even-toed ungulates, 15 odd-toed ungulates, 108
carnivores, and 15 marsupials), were retrieved from the National Center for Biotechnology
Information (NCBI) Whole Genome Shotgun (WGS) database (last accessed in November
2020). A detailed list of all the surveyed mammalian genomes is provided in Supplementary
Table S1. A representative group of 306 *Flaviviridae* or *Flaviviridae*-like polyprotein
sequences was compiled from the NCBI non-redundant protein database (accessed in February
2019). We provide a list of the nucleotide/protein accession numbers in Supplementary Table
S2.
128
2.1.2 – Genome screening
Flaviviridae polyprotein sequences were used as queries in tBLASTn (BLAST+ v2.6.0)
(Camacho et al., 2009) searches with mammalian genomes as targets. To avoid potential
artifacts, only hits with E-value < 10^{-4} and length >= 250 nt were extracted from mammalian
genomes based on the reported position by BLAST in the host contig. These putative EVEs
were then used as query in a reciprocal tBLASTx (BLAST+ v2.6.0) (Camacho et al., 2009)
against a local NCBI nucleotide (nt) database (accessed in October 2018) and BLASTx
(BLAST+ v2.6.0) (Camacho et al., 2009) against a non-redundant protein (nr) database
(accessed in October 2018). EVEs were confirmed if the best hits contained *Flaviviridae* family
members with an E-value < 10^{-4} and length >= 250 nt. The presence of conserved viral genetic
features within the hits was assessed using the NCBI Conserved Domain Database (Marchler-
Bauer et al., 2015).
141
2.1.3 – EVE characterization
Upon identification of the EVEs, they were translated and aligned with corresponding
polyprotein sequences from several representative *Flaviviridae* species using MAFFT v7.453
(Katoh et al., 2002). All alignments were trimmed in BMGE v1.12 (Criscuolo and Gribaldo,

2010) in order to select for phylogenetic informative regions. The best substitution models

were PMB+G4 for the EVE1 alignment, LG+F+G4 for the EVE2 alignment, and LG+F+I+G4

for the concatenated EVEs alignment according to the BIC criterion and were used to construct

phylogenetic ML trees with IQ-TREE v1.6.12 (Nguyen et al., 2015).

2.1.4 – Flanking region analysis

To characterize the EVEs loci and identify potential transposable elements or other genetic

features, flanking regions of the identified EVEs were extracted from the host contigs and used

as BLAST queries to screen against the NCBI nucleotide (nt) and non-redundant protein (nr)

databases (both accessed in October 2018).

2.1.5 – Metagenomic screening

According to the WGS screening results above, some *Flaviviridae*-related hits were detected

in a shrew (*Crocidura indochinensis*) genome. However, apart from the *Crocidura*

indochinensis and *Sorex araneus* complete genomes, only a limited number of shrew genomes

are currently available in the NCBI WGS database. Therefore, 73 DNA experimental genomic

data sets and 6 RNA-Seq transcriptome data sets (Supplementary Table S3) from the Soricidae

family were retrieved from NCBI Sequence Read Archive (SRA) database using SRA Toolkit

v2.10.8 (Leinonen et al., 2011). Reads were mapped to the identified EVEs nucleotide

references using Bowtie2 v2.3.5.1 (Langmead and Salzberg, 2012). Alignment files were

processed with SAMtools v1.10 (Li et al., 2009) and coverage was determined using bedtools

v2.27.1 (Quinlan and Hall, 2010) and visualized in RStudio v1.1.463.

2.2 – *In vivo* validation

2.2.1 – Sample collection

Based on the screening results, to further verify the presence of *Flaviviridae*-related EVEs *in*

vivo, a total of 65 tissue and DNA samples from species belonging to the *Crocidura* genus and

6 other related genera of the Soricidae family, namely *Paracrocidura*, *Scutisorex*, *Suncus*,

Sylvisorex (subfamily Crocidurinae), *Neomys,* and *Sorex* (subfamily Soricinae), were screened

for the presence of the identified EVEs. These samples were previously collected in China,

Vietnam, Africa, and the Eastern Mediterranean (Supplementary Table S4) as part of other

studies (Bannikova et al., 2011; de Perre et al., 2019; Jenkins et al., 2013; van de Perre et al.,

2018)

2.2.2 – Target EVEs and cytochrome b (Cytb) amplification

DNA was extracted from tissue samples using the DNeasy Blood & Tissue Kit (Qiagen)

following the manufacturer's instructions.

To screen for the presence of EVEs *in vivo*, we designed 18 PCR primers (Supplementary

Table S5) spanning the 2 EVEs region and a section of the intermediate flanking region from

the host genome. Amplicons were generated with DreamTaq DNA Polymerase (ThermoFisher

Scientific) using the following cycling conditions: (i) 3 min of denaturation at 95°C, (ii) 35

cycles of 95°C for 30s, 56°C for 30s, 72°C extension for 1min/kb, and (iii) 10 min final

extension at 72°C.

To confirm the host species, and to complement available specimen information, the

mitochondrial Cytb gene was amplified using general primers (Supplementary Table S5) of the

Crocidurinae subfamily. PCR reactions were conducted using the DreamTaq DNA Polymerase

(ThermoFisher Scientific) with the following thermal cycling conditions: (i) 3 min of

denaturation at 95°C, (ii) 35 cycles of 95°C for 30s, 56°C for 30s, 72°C extension for 1min/kb,

and (iii) 10 min final extension at 72°C.

All PCR products were purified using the ExoSAP-IT PCR Product Cleanup (ThermoFisher

Scientific) or Zymoclean Gel DNA Recovery Kits (ZYMO Research) to remove primer dimers

and unspecific products, following the manufacturer's instructions.

2.2.3 – Sanger sequencing

The generated PCR products were sequenced by Macrogen Europe. The amplicons were

mapped to the whole EVEs region (2,235nt) from the WGS *Crocidura indochinensis* contig,

and concatenated based on consensus sequence to get the complete EVEs in Geneious Prime®

v2020.2.4. Cytb amplicons (~1,140nt) were forward and reverse sequenced and a consensus

sequence was generated using Geneious Prime® v2020.2.4.

2.2.4 – MinION sequencing

For 12 samples with relatively low-quality Sanger sequencing chromatograms (additional

information provided in Supplementary Table S6), MinION sequencing was performed to

obtain the complete EVEs region (~2,235nt) together with the Cytb gene (~1,140nt). The

Oxford Nanopore Technologies (ONT) 1D Native barcoding genomic DNA protocol was used

without the DNA fragmentation step and the barcoded amplicons were loaded onto the

MinION device. We used the MinKNOW software v19.13.5 on the MinIT companion for data

acquisition and basecalling. Qcat v1.1.0 (ONT, https://github.com/nanoporetech/qcat) was

used to demultiplex reads under the epi2me algorithm and to trim bad quality reads and

adapters with min score of 90. The EVE regions extracted from the WGS *Crocidura*

indochinensis contig were used as references to map the reads with Minimap2 v.2.22 (Li, 2018)

using -ax map-ont parameters. Alignments were converted and indexed using SAMtools v1.10

(Li et al., 2009) and consensus sequences were generated using a custom Python script

(Kafetzopoulou, 2019).

2.3 – Phylogenetic analysis and visualization

All generated EVEs sequences were translated and aligned with homologous polyproteins from

available *Pestivirus* species using MAFFT v7.453 (Katoh et al., 2002). Sequences of dengue-

2 and Zika virus (*Flavivirus* genus) were used as an outgroup. The alignment was trimmed

using BMGE v1.12 (Criscuolo and Gribaldo, 2010) and the filtered regions were used to

construct maximum-likelihood (ML) phylogenetic trees using IQ-TREE v1.6.12 (Nguyen et

al., 2015) under the best-fitting models (according to the Bayesian information criterion):

WAG+G4 (for EVE1), LG+F+I+G4 (for EVE2), and LG+G4 (for complete concatenated

EVEs). Phylogenies were visualized and annotated using FigTree v1.4.4 (A. Rambaut;

http://tree.bio.ed.ac.uk/software/figtree/). Percent identity matrices were generated using

Clustal Omega (Sievers et al., 2011) via EMBL-EBI web services (Madeira et al., 2019).

Cytb sequences of EVEs-positive specimens (n=48) were aligned using MAFFT v7.453 (Katoh

et al., 2002) together with a dataset of *n*=393 Soricidae nucleotide sequences downloaded from

NCBI. The generated alignment was trimmed in BMGE v1.12 (Criscuolo and Gribaldo, 2010)

and an ML phylogeny was reconstructed using IQ-TREE v1.6.12 (Nguyen et al., 2015) with

the best-fitting model (TIM2+F+I+G4).

To highlight the evolutionary relationships of our newly discovered EVEs and their hosts,

EVEs and Cytb phylogenies were annotated in ggtree v1.14.6 (Yu et al., 2017) and treeio v1.6.2

(Wang et al., 2020) R packages, followed by the estimation of a co-phylogenetic plot

(tanglegram) using the ape v5.0 (Paradis and Schliep, 2019) and dendextend v1.14.0 (Galili,

2015) R packages.

2.4 – Characterization of selective pressure

We only characterized the selective pressure acting on the EVE 1 locus, as the EVE 2 locus

exhibits widespread stop codons and translation frame shifts. We aligned the open reading

frame of the complete EVE1 region (318 nt) using MEGA v11.0.9 (Tamura et al., 2021). An

ML tree was build based on this alignment in IQ-TREE v1.6.12 (Nguyen et al., 2015) under

the best-fitting model (HKY+F). We then conducted two site-specific selection analyses to

characterize the selective pressure on each site using estimates of the ratio of non-

synonymous/synonymous substitution rate ($\omega = d_N/d_S$): 1) fixed effects likelihood (FEL)

analyses (Kosakovsky Pond and Frost, 2005) using MG94xREV model available in HyPhy

software v2.5.3 (Kosakovsky Pond et al., 2005), and 2) the Bayesian renaissance counting

method (Lemey et al., 2012) implemented in BEAST v1.10.5 (Suchard et al., 2018). The value

of ω quantifies the selective pressure, with $\omega > 1$ suggesting positive selection, $\omega = 1$ neutral

evolution and $\omega < 1$ negative or purifying selection.

3 – Results

3.1 – False positive Flaviviridae-like hits from mammalian genome screening

Our initial screening of 689 available mammalian genomes using *Flaviviridae* and

Flaviviridae-like polyproteins yielded 66 positive hits from 49 species, including rodents, non-

human primates, marsupials, insectivores, carnivores, and bats. Detailed information about our

in silico screening results can be found in Supplementary Table S7. All hits were similar to

three pestiviruses, namely border disease virus, bovine viral diarrhea virus 1, and bovine viral

diarrhea virus 2. With the exception of one shrew species (see section below), the position of

all hits in the corresponding viral genomic sequence was in the ubiquitin-homolog domain

between the nonstructural proteins NS2 and NS3, while some of the hits slightly expanded the

alignment to the NS3 region (shown in Supplementary Fig. 1). The ubiquitin domain in bovine

viral diarrhea virus is however predicted to originate from cellular derived insertions

in cytopathogenic pestivirus (Agapov et al., 2004; Becher and Tautz, 2011). The similarity

between this viral genomic region and ubiquitin poses a considerable risk for false positives

when searching for pestivirus-derived EVEs, and these hits were therefore not further

considered.

3.2 – *In silico* identification of *Crocidura indochinensis* pesti-like EVEs

Besides the ubiquitin-related false positive results, our *in silico* screening identified a series of

five *Flaviviridae*-related EVEs fragments in a single contig of the *Crocidura indochinensis*

reference genome PVKC01 (Table 1). The first EVE (EVE1) is 318 nt long, with its closest

BLAST hit being the Linda virus (Pestivirus) envelope glycoprotein E2 region (tBLASTx, 25.5%

identity, e-value 3.31E[-35]), without any stop codon (Fig. 1). The remaining four EVEs

fragments are 1,053 nt, 84 nt, 114 nt and 87 nt long respectively, with their closest BLAST hits

being a classical swine fever virus and a rodent pestivirus (with minimum amino-acid identity

24.9%, maximum 62.1%). These four fragments are separated by very short gaps, with lengths

of 16, 1 and 4 nucleotides, respectively. The two central fragments of EVE2, fragments 3 (84

nt) and fragment 4 (114 nt), are in a different translation frame than the two others, but in the

same orientation. Their arrangement in the host contig reflects their relative position in the

pestiviral genome, which partially spans the non-structural NS2 and NS3 genes (Fig 1). For

these reasons, in our further analyses, we considered these four fragments as the result of a

single pestiviral integration event, and thus a unique EVE (EVE2). Phylogenetic

reconstructions of the identified *Crocidura indochinensis* EVE1, EVE2 and entire

concatenated sequence with exogenous *Flaviviridae* viruses supports the pestivirus-origin of

the EVEs (Supplementary Fig. 2).

In addition, a fragment prior to EVE1 in the contig shows a strong similarity (tBLASTx, 45.2%

identity, e-value $5.51E^{-19}$, supplementary Table 7) with the pestivirus ribonuclease T2 gene.

However, considering that this enzyme exists in a wide range of organisms (Luhtala and Parker,

2010), the virus-derived origin of this sequence in *Crocidura indochinensis* is not guaranteed.

The position of all pestivirus-like hits, including the ribonuclease T2 gene, in the host contig

corresponds to their relative organization in the pestivirus genome. No additional features were

detected after a tBLASTx search of the whole contig encompassing the two identified EVEs

(Supplementary Table S7). In addition, we did not detect the EVE sequences in reads of

publicly available experimental genomic and transcriptomic data from Soricidae species.

contig accession no.	contig length (nt)	EVE	GenBank accession no.	EVEs fragments	EVE length (nt)	position in host contig		translation frame	Closest BLAST hit	Conserved-domain search
						start	end			
PVKC010097735.1	6104	EVE1	BK014483	No. 1	318	2329	2646	1	Linda virus	Pestivirus envelope glycoprotein E2
		EVE2		No. 2	1053	3205	4257	1	Classical swine fever virus	Peptidase_C74: Pestivirus NS2 peptidase
				No. 3	84	4274	4357	2	Rodent pestivirus	Peptidase_S31: Pestivirus NS3 polyprotein peptidase S31
				No. 4	114	4359	4472	2	Rodent pestivirus	Peptidase_S31: Pestivirus NS3 polyprotein peptidase S31
				No. 5	87	4477	4563	1	Rodent pestivirus	Peptidase_S31: Pestivirus NS3 polyprotein peptidase S31

Table 1. Newly detected EVEs in *Crocidura indochinensis* genome

Fig. 1. The positions of the newly detected *Crocidura indochinensis* EVEs are shown relative to an archetypal *Pestivirus* genome

(classical swine fever virus, NC_002657). N^pro, N-terminal protease; C, nucleocapsid core protein; E^rns, envelope glycoprotein E^rns;

E1, envelope glycoprotein E1; E2, envelope glycoprotein E2; p7, nonstructural protein p7; NS2, nonstructural protein NS2; NS3,

nonstructural protein NS3; NS4A, nonstructural protein NS4A; NS4B, nonstructural protein NS4B; NS5A, nonstructural protein

NS5A; NS5B, nonstructural protein NS5B.

3.3 – Identification and distribution of pesti-like EVEs in other Soricidae species

To expand our screening and evaluate the distribution of these newly identified EVEs in

additional species that are phylogenetically close to *Crocidura indochinensis*, we undertook a

PCR-based screening of 65 samples from 29 species of the Soricidae family (Supplementary

Table S4). These samples belonged to 7 different genera (*Crocidura*, *Paracrocidura*,

Scutisorex, *Suncus*, *Sylvisorex*, *Neomys,* and *Sorex*), encompassing two subfamilies,

Crocidurinae and Soricinae. Cytb genomic sequences were also generated to confirm the

species identification (Supplementary Table S6 & Supplementary Fig. 3).

In total, 58 samples derived from 27 species contained the newly identified pesti-like EVEs,

and 48 samples yielded complete or nearly complete EVEs sequences, representing 22 species

in the Crocidurinae subfamily and one species (*Neomys anomalus*) in the Soricinae subfamily

(Fig. 2). All novel EVEs sequences were highly similar to the *Crocidura indochinensis* EVEs

sequences, with a mean identity of 90.38% on amino acid level, and 95.12% on nucleotide

level, respectively (Supplementary Table S8). Phylogenetic reconstruction indicated that all

Crocidurinae pesti-like EVEs clustered together as a sister-lineage of currently recognized

Pestivirus species (Fig. 3, Supplementary Fig. 4).

Though collected in different locations, nearly all species tested in the Crocidurinae subfamily

harbored the pesti-like EVEs sequences. For some species however, such as *Crocidura* cf.

zimmermanni, *C. leucodon*, *C. suaveolens* and *Suncus etruscus*, we could not always detect or

sequence the EVEs in all samples. The failed detection could be explained by the genomic

template being of poor quality due to storage conditions associated with the museum specimens.

Interestingly, the pesti-like EVEs were also detected in one *Neomys anomalus* sample, while

the remaining Soricinae specimens yielded negative results. The widespread nature of these

homolog EVEs in the Crocidurinae species suggests a single endogenization event before their

common ancestor about 10.8 million years ago (Dubey et al., 2007). Since we did not manage

to sequence the pesti-like EVEs found in *Neomys anomalus*, their phylogenetic relationship to

the Crocidurinae pesti-like EVEs is unclear and may not necessarily derive from the same

endogenization event. To compare the evolutionary history between the EVEs and the Cytb

gene, we constructed a tanglegram based on their respective phylogenies, which illustrates

specific phylogenetic discordances that may also be expected between mitochondrial and

nuclear genes (Fig 4).

3.4 – Selective pressure on pesti-like EVE1 locus

For testing the indication of function of the EVEs region, we further processed selective

pressure analysis. We only characterized the selective pressure acting on the EVE1 locus, as

the EVE2 locus exhibits widespread stop codons and translation frame shifts. We measured

the selective pressure acting on EVE1 by estimating the ratio (ω) of non-synonymous

substitution rate (d_N) and synonymous substitutions rate (d_S) in protein-coding sequences using

two methods: 1) fixed effects likelihood (FEL) (Kosakovsky Pond and Frost, 2005), and 2)

Bayesian renaissance counting (Lemey et al., 2012). It is expected that non-functional region

should conform to neutral evolution while functional region would appear to be under purifying

selection. Based on the 48 pesti-like EVE1 potentially coding sequences the FEL method

indicates an overall neutral evolution with $\omega = 0.94$. Likewise, the Bayesian renaissance

counting model yields a ratio at 1.098 [95% credible interval: 0.658, 1.624], reflecting neutral

evolution.

Fig. 2. Maximum likelihood phylogeny of the Soricidae (shrews) family based on the Cytb gene and the available samples' species

distribution in this study, highlighted in pink: (A) The phylogeny of Crocidurinae subfamily and sample distribution; (B) Subfamilies

relationships within the Soricidae family; (C) The phylogeny of Soricinae subfamily and sample distribution.

Fig. 3. Phylogenetic relationships of pesti-like EVEs with representative *Pestivirus* species and with Dengue and Zika virus

(*Flavivirus*) as outgroup. Clades are colored based on viral species. Node labels indicate Shimodaira-Hasegawa (SH)-like branch

support (%, only values > 80% are shown). Scale bars indicate the number of amino acid substitutions per site.

365
366 **Fig. 4 Tanglegram of the cytochrome b phylogeny (A) and the corresponding EVEs phylogeny (B). The cytochrome b tree was inferred**
367 **for gene sequences from 22 shrew species, and the EVEs tree (right) was inferred using the newly generated EVEs sequences. The**
368 **clade nodes with SH-like branch support < 50% were collapsed as polytomies. The black circles indicate the SH-like branch support >**
369 **80%. Lines connect corresponding tips in the two phylogenies. Scale bars indicate the number of nucleotide substitutions per site.**
370

4 – Discussion

Non-retroviral endogenous viral elements (EVEs) are rare traces of the ancient evolutionary history of viruses. These genomic fossils offer valuable insights into host range, ancestral genetic diversity and can provide invaluable information for dating viral evolutionary history (Aiewsakun and Katzourakis, 2015a; Feschotte and Gilbert, 2012). In our study we screened a comprehensive set of mammalian genomes to discover such *Flaviviridae*-derived EVEs. We uncovered two *Flaviviridae*-derived EVEs sequences in the genome of the Indochinese shrew and confirmed their presence in a broad range of shrew species belonging to the Crocidurinae subfamily.

The EVEs we identified are related to extant viruses within the *Pestivirus* genus. Viruses belonging to this genus were initially detected in a variety of artiodactylous hosts, such as ruminants and swine, in which they cause subclinical or clinical infections including hemorrhagic syndrome, abortion, acute fatal mucosal disease. Recent metagenomic studies extended the host range towards rodents (Wu et al., 2020, 2018), bats (Wu et al., 2018), fish

(Shi et al., 2018), and ticks (Sameroff et al., 2019), but to some extent, the restricted sampling
beyond agriculturally important animals limits our understanding of the real host range. Shrews,
for example, have been recently identified as host of hepaciviruses, another genus in the
Flaviviridae family (Guo et al., 2019; Wu et al., 2020), but to date not of pestiviruses. The
broad detection of pestivirus-derived EVEs reported in our study strongly supports that
ancestors of the Crocidurinae shrew subfamily have been hosts of pestiviruses and suggests
that their descendants might still be. Indeed, considering the extremely low probability of a
non-retroviral endogenization event to occur in the germline, EVEs are strong indicators of
frequent interactions between the original exogenous viruses and their hosts (Aiewsakun and
Katzourakis, 2015a; Feschotte and Gilbert, 2012). Although direct detection and
characterization of pestiviruses from shrews are still required to formally demonstrate that they
are natural hosts of pestiviruses, our study provides indirect support for a wider and more
diverse host range of pestiviruses.

Given the low probability of endogenization events of non-retroviral RNA viruses and the
contiguous nature of the two EVEs on the host and viral genome, our results suggest a single
endogenization event followed by genetic drift. One or several insertion events separated the
original EVE in two fragments, EVE1 and EVE2. EVE1 shows a short but intact open reading
frame to be evolving under neutral evolution while EVE2 exhibits multiple stop codons and
frame-shifts due to additional insertions. Many studies have identified the important roles that
EVEs can play in host antiviral immunity, both in vertebrates and invertebrates (Blair et al.,
2020; Ophinni et al., 2019; Skirmuntt et al., 2020). Flavivirus-like EVEs in *Aedes* mosquitoes,
for example, can produce P-element-induced wimpy testis (PIWI)-interacting RNAs (piRNAs)
which limit the cognate virus replication (Suzuki et al., 2020). It is highly unlikely that the
pesti-like sequences we discovered have a function in shrews because of the absence of
negative selection and the disruption of the original viral coding region.

The evolutionary history of the EVEs sequences after integration remains unclear. Interestingly,
they do not show any appreciable patterns of phylogenetic consistency with the host Cytb gene
sequences. The EVE sequences might not be a reliable genetic marker to discriminate species
(Tobe et al., 2010), as highlighted by the low amount of genetic diversity compared to Cytb
sequences (Fig 4). Additionally, the discrepancy might be explained by differences in the
genetic inheritance of the Cytb gene and EVEs: the Cytb gene is a mitochondrial gene whereas
EVEs are integrated in the nuclear genome. This can lead to different observed evolutionary

patterns, especially in the case of weak reproductive isolation within species or species
complex, allowing hybridization, as has been suggested for some *Crocidura* species (Dubey et
al., 2008, 2006; Vogel et al., 2004).

Dating the ancient evolutionary history of ssRNA viruses such as pestiviruses and *Flaviviridae*
in general is challenging. The most commonly used method for inferring viral divergence time
is based on the estimation of evolutionary rates derived from sequence data and their collection
dates. However, the applicability of this method is often limited by heterogeneous substitution
rates though time (Aiewsakun and Katzourakis, 2016) and among viral lineages (Duffy et al.,
2008; Sanjuán, 2012). Not accounting for the former leads to recent estimates for the origins
of ssRNA viruses that are often in conflict with other phylogenetic evidence (Holmes, 2003).
Using suitable molecular clock models, the powerful combination of both tip and node
calibrations may help to recover more accurate evolutionary timescales (O'Reilly et al., 2015).
Node calibration is however challenging for viruses as no fossil evidence can be found. It thus
often relies on known phylogeographic events and other indirect calibrations point, such as
ecological events or assumptions of co-divergence as alternative (Bamford et al., 2021;
Moureau et al., 2015; Pettersson and Fiz-Palacios, 2014). The discovery of ssRNA virus-
related EVEs thus enable a direct estimation for a robust long-term timeline of virus evolution
history by co-phyletic analysis of EVE's orthologs in different host (Gilbert and Feschotte,
2010).

The pesti-like EVE sequences characterized in our study are widespread in Crocidurinae
species, are monophyletic and exhibit high sequence similarity. Considering the low
probability of endogenization events of non-retroviral RNA viruses, this suggests that the pesti-
like EVE got integrated before the most-recent common ancestor of the subfamily, which is
estimated to be over 10.8 million years ago (Dubey et al., 2007). There are only a handful of
molecular dating estimates for pestiviruses and they mostly focus on viral species or clades
that are associated with economic losses. Diversification of bovine viral diarrhea virus 1
(Pestivirus A) subtypes was estimated to have started about 363 years ago (Weber et al., 2021),
and the divergence of HoBi-like pestivirus (Pestivirus H) was dated back to the 16[th] century
(Silveira et al., 2020). Our results provide the first robust ancient time node for pestiviruses
based on the estimated EVEs integration date and suggest that pestiviruses were already
circulating in mammals more than 10.8 million years ago.

In conclusion, we discovered and characterized the first *Flaviviridae*-related EVEs records
from mammalian reference genomes, which derived from pestiviruses. The wide EVEs
distribution in shrew Crocidurinae subfamily indicates they are a historical host group of
pestiviruses and further suggests a robust ancient origin time of the *Pestivirus* genus. Our
results show the key role of EVEs not only in expanding our knowledge about ancient viral-
host interactions, but also their importance in reconstructing the viral evolutionary history,
which contributes to our understanding of viral evolutionary dynamics from ancient times to
the present.
461

462   **5 – Acknowledgement**
463   The samples from Vietnam were collected during biodiversity surveys carried by the Joint
464   Vietnam-Russian Tropical Research and Technological Centre.
465

6 – Funding
The research leading to these results has received funding from the European Research Council
under the European Union's Horizon 2020 research and innovation programme (grant
agreement no. 725422-ReservoirDOCS). PL acknowledges support by the Research
Foundation -- Flanders (`Fonds voor Wetenschappelijk Onderzoek -- Vlaanderen', G066215N,
G0D5117N and G0B9317N). YQ LI acknowledges the support of China Scholarship Council.
AVA acknowledges the support by Ministry of Science and Higher Education of the Russian
Federation (grant no. 075-15-2021-1069). YC LI acknowledges the support of National Natural
Science Foundation of China (grant ID. 31672254). F.V.d.P. acknowledges the support of the
Ph.D. fellowship from the Research Foundation– Flanders. M.R.W.

References

Agapov, E. v., Murray, C.L., Frolov, I., Qu, L., Myers, T.M., Rice, C.M., 2004. Uncleaved NS2-3 Is Required for Production of Infectious Bovine Viral Diarrhea Virus. Journal of Virology 78, 2414–2425. https://doi.org/10.1128/jvi.78.5.2414-2425.2004

Aiewsakun, P., Katzourakis, A., 2016. Time-Dependent Rate Phenomenon in Viruses. Journal of Virology 90, 7184–7195. https://doi.org/10.1128/jvi.00593-16

Aiewsakun, P., Katzourakis, A., 2015a. Endogenous viruses: Connecting recent and ancient viral evolution. Virology 479–480, 26–37. https://doi.org/10.1016/j.virol.2015.02.011

Aiewsakun, P., Katzourakis, A., 2015b. Endogenous viruses: Connecting recent and ancient viral evolution. Virology. https://doi.org/10.1016/j.virol.2015.02.011

Aswad, A., Katzourakis, A., 2014. The First Endogenous Herpesvirus, Identified in the Tarsier Genome, and Novel Sequences from Primate Rhadinoviruses and Lymphocryptoviruses. PLoS Genetics 10. https://doi.org/10.1371/journal.pgen.1004332

Bannikova, A.A., Abramov, A. v., Borisenko, A. v., Lebedev, V.S., Rozhnov, V. v., 2011. Mitochondrial diversity of the white-toothed shrews (Mammalia, Eulipotyphla, Crocidura) in Vietnam. Zootaxa 1–20. https://doi.org/10.11646/zootaxa.2812.1.1

Becher, P., Tautz, N., 2011. RNA recombination in pestiviruses: Cellular RNA sequences in viral genomes highlight the role of host factors for viral persistence and lethal disease. RNA Biology 8, 216–224. https://doi.org/10.4161/rna.8.2.14514

Belyi, V.A., Levine, A.J., Skalka, A.M., 2010a. Sequences from Ancestral Single-Stranded DNA Viruses in Vertebrate Genomes: the Parvoviridae and Circoviridae Are More than 40 to 50 Million Years Old. Journal of Virology 84, 12458–12462. https://doi.org/10.1128/jvi.01789-10

Belyi, V.A., Levine, A.J., Skalka, A.M., 2010b. Unexpected inheritance: Multiple integrations of ancient bornavirus and ebolavirus/marburgvirus sequences in vertebrate Genomes. PLoS Pathogens 6, 1–13. https://doi.org/10.1371/journal.ppat.1001030

Blair, C.D., Olson, K.E., Bonizzoni, M., 2020. The widespread occurrence and potential biological roles of endogenous viral elements in insect genomes. Current Issues in Molecular Biology 34, 13–29. https://doi.org/10.21775/CIMB.034.013

Bletsa, M., Vrancken, B., Gryseels, S., Boonen, I., Fikatas, A., Li, Y., Laudisoit, A., Lequime, S., Bryja, J., Makundi, R., Meheretu, Y., Akaibe, B.D., Mbalitini, S.G., van de Perre, F., van Houtte, N., Těšíková, J., Wollants, E., van Ranst, M., Pybus, O.G., Drexler, J.F., Verheyen, E., Leirs, H., Gouy De Bellocq, J., Lemey, P., 2021. Molecular detection and genomic characterization of diverse hepaciviruses in African rodents. Virus Evolution 7, 1–14. https://doi.org/10.1093/ve/veab036

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., Madden, T.L., 2009. BLAST+: Architecture and applications. BMC Bioinformatics 10, 1–9. https://doi.org/10.1186/1471-2105-10-421

Colmant, A.M.G., Hobson-Peters, J., Bielefeldt-Ohmann, H., Hurk, A.F. van den, Hall-Mendelin, S., Chow, W.K., Johansen, C.A., Fros, J., Simmonds, P., Watterson, D., Cazier, C., Etebari, K., Asgari, S., Schulz, B.L., Beebe, N., Vet, L.J., Piyasena, T.B.H., Nguyen, H.-D., Barnard, R.T., Halla, R.A., 2017. A New Clade of Insect-Specific Flaviviruses from Australian Anopheles Mosquitoes Displays Species-Specific Host Restriction 2, 1–19.

Criscuolo, A., Gribaldo, S., 2010. BMGE (Block Mapping and Gathering with Entropy): A new software for selection of phylogenetic informative regions from multiple sequence alignments. BMC Evolutionary Biology 10. https://doi.org/10.1186/1471-2148-10-210

Crochu, S., Cook, S., Attoui, H., Charrel, R.N., de Chesse, R., Belhouchet, M., Lemasson, J.J., de Micco, P., de Lamballerie, X., 2004. Sequences of flavivirus-related RNA

viruses persist in DNA form integrated in the genome of Aedes spp. mosquitoes. Journal
of General Virology 85, 1971–1980. https://doi.org/10.1099/vir.0.79850-0

de Lamballerie, X., Crochu, S., Billoir, F., Neyts, J., de Micco, P., Holmes, E.C., Gould,
E.A., 2002. Genome sequence analysis of Tamana bat virus and its relationship with the
genus Flavivirus. Journal of General Virology 83, 2443–2454.
https://doi.org/10.1099/0022-1317-83-10-2443

de Perre, F. van, Leirs, H., Cigar, J., Mbalitini, S.G., Itoka, J.C.M., Verheyen, E., 2019.
Shrews (Soricidae) of the lowland forests around kisangani (DR Congo). Biodiversity
Data Journal 7. https://doi.org/10.3897/BDJ.7.E46948

Dubey, S., Diker, E., Kurtonur, C., Vogel, P., 2008. Secondary contact zones and
hybridizations: The case of the lesser white-toothed shrew (Crocidura suaveolens group,
Soricidae). Biological Journal of the Linnean Society 95, 557–565.
https://doi.org/10.1111/j.1095-8312.2008.01070.x

Dubey, S., Salamin, N., Ohdachi, S.D., Barrière, P., Vogel, P., 2007. Molecular
phylogenetics of shrews (Mammalia: Soricidae) reveal timing of transcontinental
colonizations. Molecular Phylogenetics and Evolution 44, 126–137.
https://doi.org/10.1016/j.ympev.2006.12.002

Dubey, S., Zaitsev, M., Cosson, J.F., Abdukadier, A., Vogel, P., 2006. Pliocene and
Pleistocene diversification and multiple refugia in a Eurasian shrew (Crocidura
suaveolens group). Molecular Phylogenetics and Evolution 38, 635–647.
https://doi.org/10.1016/j.ympev.2005.11.005

Duffy, S., Shackelton, L.A., Holmes, E.C., 2008. Rates of evolutionary change in viruses:
Patterns and determinants. Nature Reviews Genetics 9, 267–276.
https://doi.org/10.1038/nrg2323

Feschotte, C., Gilbert, C., 2012. Endogenous viruses: Insights into viral evolution and impact
on host biology. Nature Reviews Genetics 13, 283–296. https://doi.org/10.1038/nrg3199

Flynn, P.J., Moreau, C.S., 2019. Assessing the diversity of endogenous viruses throughout
ant genomes. Frontiers in Microbiology 10. https://doi.org/10.3389/fmicb.2019.01139

G. G.Bamford, C., de Souza, W., Parry, R., J. Gifford, R., 2021. Comparative analysis of
genome-encoded viral sequences reveals the evolutionary history of Flaviviridae.
bioRxiv 2021.09.19.460981 (preprint).

Galili, T., 2015. dendextend: An R package for visualizing, adjusting and comparing trees of
hierarchical clustering. Bioinformatics 31, 3718–3720.
https://doi.org/10.1093/bioinformatics/btv428

Gilbert, C., Feschotte, C., 2010. Genomic fossils calibrate the long-term evolution of
hepadnaviruses. PLoS Biology 8. https://doi.org/10.1371/journal.pbio.1000495

Guo, H., Cai, C., Wang, B., Zhuo, F., Jiang, R., Wang, N., Li, B., Zhang, W., Zhu, Y., Fan,
Y., Chen, Wushen, Chen, Weihong, Yang, X., Shi, Z., 2019. Novel hepacivirus in Asian
house shrew, China. Science China Life Sciences 62, 701–704.
https://doi.org/10.1007/s11427-018-9435-7

Hartlage, A.S., 2016. The Strange, Expanding World of Animal Hepaciviruses. Annu Rev
Virol 176, 53–75. https://doi.org/10.1146/annurev-virology-100114-055104.The

Holmes, E.C., 2011. The evolution of endogenous viral elements. Cell Host and Microbe 10,
368–377. https://doi.org/10.1016/j.chom.2011.09.002

Holmes, E.C., 2003. Molecular Clocks and the Puzzle of RNA Virus Origins. Journal of
Virology 77, 3893–3897. https://doi.org/10.1128/jvi.77.7.3893-3897.2003

Horie, M., 2019. Interactions among eukaryotes, retrotransposons and riboviruses:
Endogenous riboviral elements in eukaryotic genomes. Genes and Genetic Systems 94,
253–267. https://doi.org/10.1266/ggs.18-00049

Horie, M., Honda, T., Suzuki, Y., Kobayashi, Y., Daito, T., Oshida, T., Ikuta, K., Jern, P., Gojobori, T., Coffin, J.M., Tomonaga, K., 2010. Endogenous non-retroviral RNA virus elements in mammalian genomes. Nature 463, 84–87. https://doi.org/10.1038/nature08695

Horie, M., Kobayashi, Y., Suzuki, Y., Tomonaga, K., 2013. Comprehensive analysis of endogenous bornavirus-like elements in eukaryote genomes. Philosophical Transactions of the Royal Society B: Biological Sciences 368. https://doi.org/10.1098/rstb.2012.0499

Jenkins, P.D., Abramov, A. v., Bannikova, A.A., Rozhnov, V. v., 2013. Bones and genes: Resolution problems in three Vietnamese species of Crocidura (Mammalia, Soricomorpha, Soricidae) and the description of an additional new species. ZooKeys 313, 61–79. https://doi.org/10.3897/zookeys.313.4823

Kafetzopoulou, L.E., 2019. Metagenomic sequencing at the epicenter of the Nigeria 2018 Lassa fever outbreak. Science 176, 139–148. https://doi.org/10.1016/j.physbeh.2017.03.040

Katoh, K., Misawa, K., Kuma, K.I., Miyata, T., 2002. MAFFT: A novel method for rapid multiple sequence alignment based on fast Fourier transform. Nucleic Acids Research 30, 3059–3066. https://doi.org/10.1093/nar/gkf436

Katzourakis, A., Gifford, R.J., 2010a. Endogenous viral elements in animal genomes. PLoS Genetics 6. https://doi.org/10.1371/journal.pgen.1001191

Katzourakis, A., Gifford, R.J., 2010b. Endogenous viral elements in animal genomes. PLoS Genetics 6. https://doi.org/10.1371/journal.pgen.1001191

Kawasaki, J., Kojima, S., Mukai, Y., Tomonaga, K., Horie, M., 2021. 100-My history of bornavirus infections hidden in vertebrate genomes. Proceedings of the National Academy of Sciences 118. https://doi.org/10.1073/PNAS.2026235118

Kobayashi, Y., Shimazu, T., Murata, K., Itou, T., Suzuki, Y., 2019. An endogenous adeno-associated virus element in elephants. Virus Research 262, 10–14. https://doi.org/10.1016/j.virusres.2018.04.015

Kosakovsky Pond, S.L., Frost, S.D.W., 2005. Not so different after all: A comparison of methods for detecting amino acid sites under selection. Molecular Biology and Evolution 22, 1208–1222. https://doi.org/10.1093/molbev/msi105

Kosakovsky Pond, S.L., Frost, S.D.W., Muse, S. v., 2005. HyPhy: Hypothesis testing using phylogenies. Bioinformatics 21, 676–679. https://doi.org/10.1093/bioinformatics/bti079

Langmead, B., Salzberg, S.L., 2012. Fast gapped-read alignment with Bowtie 2. Nature Methods 9, 357–359. https://doi.org/10.1038/nmeth.1923

Leinonen, R., Sugawara, H., Shumway, M., 2011. The sequence read archive. Nucleic Acids Research 39, 2010–2012. https://doi.org/10.1093/nar/gkq1019

Lemey, P., Minin, V.N., Bielejec, F., Pond, S.L.K., Suchard, M.A., 2012. A counting renaissance: Combining stochastic mapping and empirical Bayes to quickly detect amino acid sites under positive selection. Bioinformatics 28, 3248–3256. https://doi.org/10.1093/bioinformatics/bts580

Lequime, S., Lambrechts, L., 2017. Discovery of flavivirus-derived endogenous viral elements in Anopheles mosquito genomes supports the existence of Anopheles - associated insect-specific flaviviruses . Virus Evolution 3, vew035. https://doi.org/10.1093/ve/vew035

Li, H., 2018. Minimap2: Pairwise alignment for nucleotide sequences. Bioinformatics 34, 3094–3100. https://doi.org/10.1093/bioinformatics/bty191

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R., 2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics 25, 2078–2079. https://doi.org/10.1093/bioinformatics/btp352

Li, N., Li, Q., 2015. Identification and characterization of endogenous viral elements for the three key schistosomes of humans. Pakistan journal of pharmaceutical sciences 28, 375–382.

Liu, H., Fu, Y., Xie, J., Cheng, J., Ghabrial, S.A., Li, G., Yi, X., Jiang, D., 2012. Discovery of novel dsRNA viral sequences by in silico cloning and implications for viral diversity, host range and evolution. PLoS ONE 7, 1–10. https://doi.org/10.1371/journal.pone.0042147

Liu, S., Coates, B.S., Bonning, B.C., 2020. Endogenous viral elements integrated into the genome of the soybean aphid, Aphis glycines. Insect Biochemistry and Molecular Biology 123, 103405. https://doi.org/10.1016/j.ibmb.2020.103405

Luhtala, N., Parker, R., 2010. T2 Family ribonucleases: ancient enzymes with diverse roles. Trends in Biochemical Sciences 35, 253–259. https://doi.org/10.1016/j.tibs.2010.02.002

Madeira, F., Park, Y.M., Lee, J., Buso, N., Gur, T., Madhusoodanan, N., Basutkar, P., Tivey, A.R.N., Potter, S.C., Finn, R.D., Lopez, R., 2019. The EMBL-EBI search and sequence analysis tools APIs in 2019. Nucleic Acids Research 47, W636–W641. https://doi.org/10.1093/nar/gkz268

Marchler-Bauer, A., Derbyshire, M.K., Gonzales, N.R., Lu, S., Chitsaz, F., Geer, L.Y., Geer, R.C., He, J., Gwadz, M., Hurwitz, D.I., Lanczycki, C.J., Lu, F., Marchler, G.H., Song, J.S., Thanki, N., Wang, Z., Yamashita, R.A., Zhang, D., Zheng, C., Bryant, S.H., 2015. CDD: NCBI's conserved domain database. Nucleic Acids Research 43, D222–D226. https://doi.org/10.1093/nar/gku1221

Mares-Guia, M.A.M.D.M., Horta, M.A., Romano, A., Rodrigues, C.D.S., Mendonça, M.C.L., dos Santos, C.C., Torres, M.C., Araujo, E.S.M., Fabri, A., de Souza, E.R., Ribeiro, R.O.R., Lucena, F.P., Junior, L.C.A., da Cunha, R. v., Nogueira, R.M.R., Sequeira, P.C., de Filippis, A.M.B., 2020. Yellow fever epizootics in non-human primates, Southeast and Northeast Brazil (2017 and 2018). Parasites and Vectors 13, 1–8. https://doi.org/10.1186/s13071-020-3966-x

Maruyama, S.R., Castro-Jorge, L.A., Ribeiro, J.M.C., Gardinassi, L.G., Garcia, G.R., Brandão, L.G., Rodrigues, A.R., Okada, M.I., Abrão, E.P., Ferreira, B.R., da Fonseca, B.A.L., de Miranda-Santos, I.K.F., 2014. Characterisation of divergent flavivirus NS3 and NS5 protein sequences detected in Rhipicephalus microplus ticks from Brazil. Memorias do Instituto Oswaldo Cruz 109, 38–50. https://doi.org/10.1590/0074-0276130166

Moureau, G., Cook, S., Lemey, P., Nougairede, A., Forrester, N.L., Khasnatinov, M., Charrel, R.N., Firth, A.E., Gould, E.A., de Lamballerie, X., 2015. New insights into flavivirus evolution, taxonomy and biogeographic history, extended by analysis of canonical and alternative coding sequences e0117849. PLoS ONE 10. https://doi.org/10.1371/journal.pone.0117849

Nguyen, L.T., Schmidt, H.A., von Haeseler, A., Minh, B.Q., 2015. IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Molecular Biology and Evolution 32, 268–274. https://doi.org/10.1093/molbev/msu300

Öncü, C., Brinkmann, A., Günay, F., Kar, S., Öter, K., Sarıkaya, Y., Nitsche, A., Linton, Y.M., Alten, B., Ergünay, K., 2018. West Nile virus, Anopheles flavivirus, a novel flavivirus as well as Merida-like rhabdovirus Turkey in field-collected mosquitoes from Thrace and Anatolia. Infection, Genetics and Evolution 57, 36–45. https://doi.org/10.1016/j.meegid.2017.11.003

Ophinni, Y., Palatini, U., Hayashi, Y., Parrish, N.F., 2019. piRNA-Guided CRISPR-like Immunity in Eukaryotes. Trends in Immunology 40, 998–1010. https://doi.org/10.1016/j.it.2019.09.003

O'Reilly, J.E., dos Reis, M., Donoghue, P.C.J., 2015. Dating Tips for Divergence-Time
Estimation. Trends in Genetics 31, 637–650. https://doi.org/10.1016/j.tig.2015.08.001
Paradis, E., Schliep, K., 2019. Ape 5.0: An environment for modern phylogenetics and
evolutionary analyses in R. Bioinformatics 35, 526–528.
https://doi.org/10.1093/bioinformatics/bty633
Parry, R., Asgari, S., 2019. Discovery of Novel Crustacean and Cephalopod Flaviviruses:
Insights into the Evolution and Circulation of Flaviviruses between Marine Invertebrate
and Vertebrate Hosts. Journal of Virology 93, 1–20. https://doi.org/10.1128/jvi.00432-
19
Pettersson, J.H.O., Fiz-Palacios, O., 2014. Dating the origin of the genus Flavivirus in the
light of Beringian biogeography. Journal of General Virology 95, 1969–1982.
https://doi.org/10.1099/vir.0.065227-0
Quinlan, A.R., Hall, I.M., 2010. BEDTools: A flexible suite of utilities for comparing
genomic features. Bioinformatics 26, 841–842.
https://doi.org/10.1093/bioinformatics/btq033
Roiz, D., Vázquez, A., Seco, M.P.S., Tenorio, A., Rizzoli, A., 2009. Detection of novel insect
flavivirus sequences integrated in Aedes albopictus (Diptera: Culicidae) in Northern
Italy. Virology Journal 6, 1–6. https://doi.org/10.1186/1743-422X-6-93
Sameroff, S., Tokarz, R., Charles, R.A., Jain, K., Oleynik, A., Che, X., Georges, K.,
Carrington, C. v., Lipkin, W.I., Oura, C., 2019. Viral Diversity of Tick Species
Parasitizing Cattle and Dogs in Trinidad and Tobago. Scientific Reports 9, 1–10.
https://doi.org/10.1038/s41598-019-46914-1
Sanjuán, R., 2012. From molecular genetics to phylodynamics: Evolutionary relevance of
mutation rates across viruses. PLoS Pathogens 8, 1–5.
https://doi.org/10.1371/journal.ppat.1002685
Shi, M., Lin, X.-D., Chen, X., Tian, J.-H., Chen, L.-J., Li, K., Wang, W., Eden, J.-S., Shen,
J.-J., Liu, L., Holmes, E.C., Zhang, Y.-Z., 2018. The evolutionary history of vertebrate
RNA viruses. Nature 561, E6–E6. https://doi.org/10.1038/s41586-018-0310-0
Shi, M., Lin, X.-D., Vasilakis, N., Tian, J.-H., Li, C.-X., Chen, L.-J., Eastwood, G., Diao, X.-
N., Chen, M.-H., Chen, X., Qin, X.-C., Widen, S.G., Wood, T.G., Tesh, R.B., Xu, J.,
Holmes, E.C., Zhang, Y.-Z., 2016. Divergent Viruses Discovered in Arthropods and
Vertebrates Revise the Evolutionary History of the Flaviviridae and Related Viruses.
Journal of Virology 90, 659–669. https://doi.org/10.1128/jvi.02036-15
Sievers, F., Wilm, A., Dineen, D., Gibson, T.J., Karplus, K., Li, W., Lopez, R., McWilliam,
H., Remmert, M., Söding, J., Thompson, J.D., Higgins, D.G., 2011. Fast, scalable
generation of high-quality protein multiple sequence alignments using Clustal Omega.
Molecular Systems Biology 7. https://doi.org/10.1038/msb.2011.75
Silva, E., Marques, S., Osório, H., Carvalheira, J., Thompson, G., 2012. Endogenous
Hepatitis C Virus Homolog Fragments in European Rabbit and Hare Genomes Replicate
in Cell Culture. PLoS ONE 7. https://doi.org/10.1371/journal.pone.0049820
Silva, E., Osório, H., Thompson, G., 2015. Hepatitis C-like viruses are produced in cells from
rabbit and hare DNA. Scientific Reports 5, 1–10. https://doi.org/10.1038/srep14535
Silveira, S., Cibulski, S.P., Junqueira, D.M., Mósena, A.C.S., Weber, M.N., Mayer, F.Q.,
Canal, C.W., 2020. Phylogenetic and evolutionary analysis of HoBi-like pestivirus:
Insights into origin and dispersal. Transboundary and Emerging Diseases 67, 1909–
1917. https://doi.org/10.1111/tbed.13520
Skirmuntt, E.C., Escalera-Zamudio, M., Teeling, E.C., Smith, A., Katzourakis, A., 2020. The
Potential Role of Endogenous Viral Elements in the Evolution of Bats as Reservoirs for
Zoonotic Viruses. Annual Review of Virology 7, 103–119.
https://doi.org/10.1146/annurev-virology-092818-015613

Soto, E., Camus, A., Yun, S., Kurobe, T., Leary, J.H., Rosser, T.G., Dill-Okubo, J.A., Nyaoke, A.C., Adkison, M., Renger, A., Ng, T.F.F., 2020. First Isolation of a Novel Aquatic Flavivirus from Chinook Salmon (Oncorhynchus tshawytscha) and Its In Vivo Replication in a Piscine Animal Model . Journal of Virology 94. https://doi.org/10.1128/jvi.00337-20

Strand, T.M., Lundkvist, Å., Olsen, B., Gustafsson, L., 2018. Breeding consequences of flavivirus infection in the collared flycatcher. BMC Evolutionary Biology 18, 1–9. https://doi.org/10.1186/s12862-018-1121-5

Suchard, M.A., Lemey, P., Baele, G., Ayres, D.L., Drummond, A.J., Rambaut, A., 2018. Bayesian phylogenetic and phylodynamic data integration using BEAST 1.10. Virus Evolution 4, 1–5. https://doi.org/10.1093/ve/vey016

Suzuki, Y., Baidaliuk, A., Miesen, P., Frangeul, L., Crist, A.B., Merkling, S.H., Fontaine, A., Lequime, S., Moltini-Conclois, I., Blanc, H., van Rij, R.P., Lambrechts, L., Saleh, M.C., 2020. Non-retroviral Endogenous Viral Element Limits Cognate Virus Replication in Aedes aegypti Ovaries. Current Biology 30, 3495-3506.e6. https://doi.org/10.1016/j.cub.2020.06.057

Tamura, K., Stecher, G., Kumar, S., 2021. MEGA11: Molecular Evolutionary Genetics Analysis Version 11. Molecular Biology and Evolution 38, 3022–3027. https://doi.org/10.1093/molbev/msab120

Taylor, D.J., Leach, R.W., Bruenn, J., 2010. Filoviruses are ancient and integrated into mammalian genomes. BMC Evolutionary Biology 10. https://doi.org/10.1186/1471-2148-10-193

Tobe, S.S., Kitchener, A.C., Linacre, A.M.T., 2010. Reconstructing mammalian phylogenies: A detailed comparison of the cytochrome b and cytochrome oxidase subunit i mitochondrial genes. PLoS ONE 5. https://doi.org/10.1371/journal.pone.0014156

van de Perre, F., Willig, M.R., Presley, S.J., Andemwana, F.B., Beeckman, H., Boeckx, P., Cooleman, S., de Haan, M., de Kesel, A., Dessein, S., Grootaert, P., Huygens, D., Janssens, S.B., Kearsley, E., Kabeya, P.M., Leponce, M., van den Broeck, D., Verbeeck, H., Würsten, B., Leirs, H., Verheyen, E., 2018. Reconciling biodiversity and carbon stock conservation in an Afrotropical forest landscape. Science Advances 4. https://doi.org/10.1126/sciadv.aar6603

Vogel, P., Maddalena, T., Sarà, M., 2004. Crocidura cossyrensis Contoli, 1989 (Mammalia, Soricidae): Karyotype, biochemical genetics and hybridization experiments. Revue Suisse de Zoologie 111, 925–934. https://doi.org/10.5962/bhl.part.80277

Wang, L.G., Lam, T.T.Y., Xu, S., Dai, Z., Zhou, L., Feng, T., Guo, P., Dunn, C.W., Jones, B.R., Bradley, T., Zhu, H., Guan, Y., Jiang, Y., Yu, G., 2020. Treeio: An R Package for Phylogenetic Tree Input and Output with Richly Annotated and Associated Data. Molecular Biology and Evolution 37, 599–603. https://doi.org/10.1093/molbev/msz240

Weber, M.N., Wolf, J.M., da Silva, M.S., Mosena, A.C.S., Budaszewski, R.F., Lunge, V.R., Canal, C.W., 2021. Insights into the origin and diversification of bovine viral diarrhea virus 1 subtypes. Archives of Virology 166, 607–611. https://doi.org/10.1007/s00705-020-04913-y

Wu, Z., Han, Y., Liu, B., Li, H., Zhu, G., Latinne, A., Dong, J., Sun, L., Du, J., Zhou, S., Chen, M., Kritiyakan, A., Jittapalapong, S., Chaisiri, K., Buchy, P., Duong, V., Yang, J., Jiang, J., Xu, X., Zhou, H., Yang, F., Morand, S., Daszak, P., Jin, Q., 2020. Decoding the RNA viromes of rodent lungs provides new visions into the origin and evolution pattern of rodent-borne diseases in Mainland Southeast Asia. https://doi.org/10.21203/rs.3.rs-17323/v1

Wu, Z., Liu, B., Du, J., Zhang, J., Lu, L., Zhu, G., Han, Y., Su, H., Yang, L., Zhang, S., Liu, Q., Jin, Q., 2018. Discovery of diverse rodent and bat pestiviruses with distinct genomic

and phylogenetic characteristics in several Chinese provinces. Frontiers in Microbiology
9, 1–8. https://doi.org/10.3389/fmicb.2018.02562
Yu, G., Smith, D.K., Zhu, H., Guan, Y., Lam, T.T.Y., 2017. Ggtree: an R Package for
Visualization and Annotation of Phylogenetic Trees With Their Covariates and Other
Associated Data. Methods in Ecology and Evolution 8, 28–36.
https://doi.org/10.1111/2041-210X.12628

779