Omicron-Specific and Bivalent Omicron-Containing Vaccine Candidates Elicit Potent Virus Neutralisation in the Animal Model

Asghar Abdoli[a,b], Hamidreza Jamshidi[c], Mohammad Taqavian[d], Mehdi Lari Baghal[e], Hasan Jalili[f,*]

[a]Department of Hepatitis and AIDS, Pasteur Institute of Iran, Tehran, Iran

[b]Amirabad Virology Laboratory, Vaccine Unit, Tehran, Iran

[c]Department of Pharmacology, Faculty of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

[d]Research and Development Unit, Biosunpharmed Co, Karaj, Iran

[e]Research and Development Unit, Shifa Pharmed Co, Karaj, Iran

[f]Department of Life Science Engineering, Faculty of New Sciences and Technologies, University of Tehran, Tehran, Iran

[*]Corresponding author: hjalili@ut.ac.ir

Abstract

Background: Omicron variant (B.1.1.529) is able to escape from naturally acquired and

vaccine-induced immunity, which mandates updating the current COVID-19 vaccines. Here, we

investigated and compared the neutralising antibody induction of the ancestral variant-based

BIV1-CovIran vaccine, the Omicron variant-based BIV1-CovIran Plus vaccine, and the novel

bivalent vaccine candidate, BBIV1-CovIran, against the Omicron and ancestral Wuhan variants

on the rat model.

Methods: Viruses were isolated from a clinical specimen and virus characterisation performed.

After inactivating the viral particles, the viruses were purified and formulated. Bivalent vaccines

were a composition of 2.5 µg (5 µg total) or 5 µg (10 µg total) doses of each ansectral-based and

Omicron-based monovalent vaccine. Subsequently, the potency of the monovalent and bivalent

vaccines was investigated using the virus neutralisation test (VNT).

Results: The group that received three doses of the Omicron-specific vaccine demonstrated

neutralisation activity against the Omicron variant with a geometric mean titer of 337.8.

However, three doses of the Wuhan variant-specific vaccine could neutralise the Omicron

variant at a maximum of 1/32 serum dilution. The neutralisation activity of the Omicron-specific

vaccine, when administered as the booster dose after two doses of the Wuhan variant-specific

vaccine, was 100% against the Omicron variant and the Wuhan variant at 1/64 and 1/128 serum

dilution, respectively. Three doses of 5 µg bivalent vaccine could effectively neutralise both

variants at the minimum of 1/128 serum dilution. The 10 µg bivalent vaccine at three doses

showed even higher neutralisation titers: geometric mean titer of 338.0 against Omicron and

445.7 against Wuhan).

Conclusion: It is shown that the candidate bivalent vaccine could elicit a potent immune

response against both Wuhan-Hu-1 and Omicron BA.1 variants. Therefore, we plan to evaluate

the updated vaccine in the clinical trial setting.

Keywords: Bivalent vaccine; Virus Neutralisation Test; Neutralising antibody; Omicron variant;

SARS-CoV-2

Introduction

The Omicron lineage of SARS-CoV-2, first described in November 2021, spread rapidly to become globally dominant and has split into many sublineages (1,2). With its high transmissibility and multiple novel spike (S) protein mutations, Omicron could escape from naturally acquired immunity while also challenging the effectiveness of current COVID-19 vaccines (2,3). According to the WHO weekly epidemiological update on COVID-19 released on 19 January 2023, globally, the Omicron variant of concern, accounted for 99.9% of reported SARS-Cov-2 viral sequences in the past 30 days (4).

The health impacts of future SARS-CoV-2 transmission depend on evolving mitigation strategies by health systems and societies. World Health Organization has reportedly declared that vaccination strategies based on repeated booster doses are unlikely to be sustainable. In this sense, the current COVID-19 vaccines originally developed against ancestral strains may need to be updated (5). The earliest attempts of boosting with Omicron-based vaccines had controversial results, some indicating similar efficacy to ancestral variant-based vaccine boosters in neutralising the Omicron variant (6), while others showed a 20-fold higher neutralising titers against the Omicron variant compared to wild-type boosters (7). Later findings indicated that bivalent vaccines containing both Omicron and ancestral strain-matched vaccines could elicit a superior neutralising antibody response against omicron and cross-reactivity with other variants, without evident safety concerns (8,9).

Since the beginning of the pandemic, Iran has endeavoured to develop and administer safe domestic vaccines, and considering the successful experiences of the country in the mass production of inactivated vaccines, applying this vaccine platform seemed the most feasible (10). BIV1-CovIran is an inactivated virus vaccine developed against the Wuhan variant of SARS-

CoV-2. In phase I and II clinical trials, all adverse events were mild or moderate and transient. The seroconversion rate of neutralising-antibody was 82.8% and the vaccine neutralised SARS-CoV-2 at 1/64 serum titers among 82.5% of 280 participants who received two doses of 5 µg vaccine in a 28-day interval (10). In the phase III clinical trial, which is currently under peer review, a two-dose vaccine regimen was well tolerated, with no safety concerns, and conferred 70.5% and 83.1% efficacy against hospitalization and ICU admission, respectively. A historical cohort study also showed the vaccine has reduced hospitalization by 86.4% and deaths by 98.3% (11).

Exploiting our previous experience in constructing the Wuhan variant-based vaccine and commensurate with the global efforts in updating previous vaccines against the omicron variant, the inactivated virus vaccine platform was re-employed to develop an Omicron variant-based vaccine titled BIV1-CovIran Plus within a month of virus isolation (12). In the clinical trial study, A 5µg booster dose of the Omicron-based vaccine induced superior Omicron-specific total IgG antibody and more potent neutralisation titer compared to ancestral strain-based vaccines (neutralisation rate of 66% vs. 25% in 1/16 serum dilution).

Considering the enhanced immunogenicity and cross-reactivity shown in bivalent SARS-CoV-2 vaccines, a bivalent vaccine candidate was also developed containing an equal amount of inactivated virus particles of the ancestral SARS-CoV-2 (Wuhan-Hu-1) and the Omicron variant (BA.1) in two doses of 5 µg (2.5 µg each) and 10 µg (5 µg each). Here, we investigated and compared the neutralising antibody induction of the ancestral variant-based BIV1-CovIran vaccine, the Omicron variant-based BIV1-CovIran Plus vaccine, and the novel bivalent vaccine candidate, BBIV1-CovIran , against the Omicron and ancestral Wuhan variants on the rat model.

Materials and methods

Overview

This study was conducted following our previous study on the preclinical evaluation of the

BIV1-CovIran vaccine in the mice model (13). Here, we conducted an animal study in the rat

model to evaluate and compare the Omicron-specific vaccine (BIV1-CovIran Plus) and the

bivalent vaccine (BBIV1-CovIran) with the ancestral variant-based vaccine (BIV1-CovIran)

against Wuhan-Hu-1 and BA.1 SARS-CoV-2 variants. Since the construction method of

Omicron-specific vaccine introduced here has not changed from the previously generated BIV1-

CovIran vaccine (13). Briefly, the Omicron variant was isolated and an inactivated vaccine was

generated against it. The 5-μg and 10-μg doses of bivalent vaccine consisted of 2.5 μg or 5 μg

from each of the ancestral Wuhan-Hu-1 or Omicron inactivated vaccines was also generated.

Subsequently, the safety and immunogenicity of the generated vaccines against the Wuhan and

Omicron variant were evaluated and compared with the ancestral-based vaccine in animal

models.

Virus isolation and vaccines preparation

In the biosafety level□3 facility, the Omicron variant virus was isolated from a

nasopharyngeal swab and subsequently infected mono-layer Vero cells (ATCC#

CCL81). After incubation, virus growth was detected and confirmed by cytopathic

effects (CPE), gene detection and electron microscopy. After passage, the viral RNAs

were extracted and converted to cDNAs to be sequenced and characterized.

Multiplication kinetics analysis of viral particles in Vero cells were demonstrated by

TCID50, 72 hours post-infection.

The virus particles were inactivated by BPL (1:1400 dilution) at 2–8°C for 20–24

hours. Moreover, to eliminate the possibility of remaining live virus particles, three

blind passages were done and no CPE was detected (14). Finally, the inactivated

viruses were purified by ultracentrifugation and chromatography and then, formulated

with Alhydrogel ® adjuvant 2% (Brenntag Biosector).

In this study, two types of monovalent and bivalent vaccines were prepared. The

monovalent vaccines were composed of a 5 µg dose of the Wuhan-specific or the

Omicron-specific vaccines. On the other hand, bivalent vaccines were a composition of

2.5 µg or 5 µg doses of each vaccine.

Animal Studies

Nine groups of five female, eight weeks old Wistar rat models provided by the

Laboratory Animal Science Department, Pasteur Institute of Iran, Karaj, Iran were

prepared. A total of 45 rats were divided into nine groups (each containing 5 rats).

Some groups received two doses of vaccines on days 0 and 28, while others received

three doses on days 0, 28, and 56. A detailed description of the study groups stratified

by the vaccine they had been received is demonstrated in Table 1. All animal studies

were conducted under the approval of animal ethics committee guidelines of the

Ministry of Health and Medical Education (Tehran, Iran; ethical code:

IR.ACECR.IBCRC.REC.1399.016).

Table 1. The vaccine type, dose numbers, and the days of injection among study groups

Group	Vaccine	Number of doses	Days of injection
WW	BIV1-CovIran (monovalent ancestral-based)	2	0-28

OO	BIV1-CovIran Plus (monovalent Omicron-based)	2	0-28
WWW	BIV1-CovIran	3	0-28-56
OOO	BIV1-CovIran Plus	3	0-28-56
WWO	2xBIV1-CovIran + 1xBIV1-CovIran Plus	3	0-28-56
BB (2.5)	BBIV1-CovIran (bivalent vaccine: 2.5μg +2.5μg)	2	0-28
BB (5)	BBIV1-CovIran (bivalent vaccine: 5μg +5μg)	2	0-28
BBB (2.5)	BBIV1-CovIran (2.5μg +2.5μg)	3	0-28-56
BBB (5)	BBIV1-CovIran (5μg +5μg)	3	0-28-56

Virus neutralisation test

The conventional virus neutralising test (cVNT) was conducted on sera acquired from

rat groups (15–17). The heat-inactivated sera were mixed with Omicron and Wuhan

variants of SARS-CoV-2 in the Dulbecco's Modified Eagle Medium (DMEM). The

mixtures were then inoculated onto Vero cells in 96☐well plates and the supernatants

(containing neutralising antibodies) were removed at one hour post infection. The

infected Vero cells were kept in DMEM and incubated for 48 h at 37 °C in a 5% CO_2

incubator. CPEs were recorded under microscopes 72 h post☐infection and the

neutralising antibody titers were measured as values of the highest dilution that

inhibited CPE formation in each well.

Results

Viral particles successfully infected the Vero cells to prepare the vaccine seed

The Variant of Concern (VOC) Omicron GRA (B.1.1.529+BA) SARS-CoV-2 strain

was successfully isolated in Vero cells from the clinical specimen. The isolated strain

had more than 99.9% identity to the original Omicron variant. Multiplication kinetics

analysis of viral particles in Vero cells demonstrated the peak of multiplication with

10^6 TCID$_{50}$, 72 hours post-infection (Figure 1-A). After four passages, the vaccine seed

was prepared, and the virus growth was recorded by the presence of CPE (Figures 1-B

and C).

The monovalent Omicron-specific vaccine demonstrated neutralisation activity
against the Omicron at various dose intervals

Among five rats that received two doses of the Omicron-specific on days 0 and 28,

four of them (80%) demonstrated neutralisation activity against the Omicron variant at

the 1/128 dilution (Figure 2A). On the other hand, two doses of Wuhan-specific could

neutralise the Omicron variant at maximum 1/8 serum dilution (i.e., geometric mean:

97 vs. 7 against Omicron) (Figure 3). All of the samples' sera (100%) in OO group

neutralised the Wuhan variant at the 1/16 dilution (Figure 2B). In addition, among

samples that received three doses of Omicron-specific on days 0, 28, and 56, the

maximum dilution that successfully neutralised the Omicron variant was 1/512. Indeed,

3 out of five rats (60%) at 1/256 and 2 out of five (40%) at 1/512 sera dilution

demonstrated neutralisation activity (Figure 2C). Three doses of Omicron-specific were

also able to neutralise the Wuhan variant at 1/32 (80%) and 1/64 (20%) dilutions

(Figure 2D). We also examined the neutralisation activity of the Omicron-specific

vaccine when administered as the booster dose on day 56 after two doses of original

the Wuhan-specific vaccine (i.e., BIV1-CovIran) on days 0 and 28. In this group, all

five samples (100%) neutralised the Omicron variant and the Wuhan variant at 1/64

and 1/128 serum dilution, respectively (Figures 2C and D). The neutralisation potency

in this group was also higher than three doses of Wuhan-specific vaccine against the

Omicron variant (i.e., geometric mean: 64 vs. 27.9) (Figure 3).

The bivalent vaccine demonstrated neutralisation activity against both the

Omicron and the Wuhan variants

Two different doses (i.e., 2.5 and 5 mg) of the Omicron-specific and the Wuhan-

specific vaccines were mixed and injected into rats two or three times at 28-day

intervals. The first group that had received two doses of the bivalent vaccine consisting

of 2.5 mg from each vaccine on days 0 and 28 demonstrated similar neutralising

activity against both variants (Figure 2). Accordingly, the sera of three rats (60%)

could neutralise both the Omicron and Wuhan variants at 1/128 dilution, and the other

two (40%) had neutralising activity at 1/64 serum dilution (i.e., geometric mean: 97)

(Figure 3). When it came to the group that had received two doses of bivalent made

from 5 mg of each vaccine, the minimum neutralising serum dilution was 1/128. 80%

of the samples' sera in this group neutralised the Omicron variant at 1/128 dilution, and

20% neutralised it at 1/256 (i.e., geometric mean: 147). On the other hand, 60% of

them neutralised the Wuhan variant at 1/128 and 40% at 1/256 serum dilutions (i.e.,

geometric mean: 168.9). Finally, when we tried three doses of the bivalent vaccines on

days 0, 28, and 56, the neutralising potency of sera was approximately one-fold

increased. Four out of five rats (80%) that had received three doses of the 2.5mg

mixture demonstrated neutralising activity against both the Omicron and the Wuhan

variants at 1/256 serum dilution (Figure 4), and the other one at 1/512 dilution. In the

last group, three doses of 5mg mixture were tested and the maximum neutralising

serum dilution was 1/256. In this group, 60% and 80% of the samples' sera could

neutralise the Omicron and Wuhan variants, respectively at 1/512 dilution (i.e., the

geometric mean of 445.7 against Wuhan and 388 against Omicron) (Figure 3).

Discussion

Omicron is associated with a marked ability of reinfection or breakthrough infection even in

populations with wide vaccine coverage, benefiting from its heavily altered antigenicity (1,18).

Since the identification of the variant, evidence prompted the need for developing Omicron-

specific vaccines (7). Recent Omicron sublineages, BA.4/5, are currently causing most cases of

COVID-19 in many countries, and have been shown to resist even more than BA.1 and BA.2

against neutralisation by triple-dosed vaccinee serums (19). It is evident that protection against

symptomatic BA.4/5 reinfection in those with a previous pre-omicron SARS-CoV-2 infection

was 35.5%, while in those with a history of infection with a post-Omicron subvariant (including

BA.1 or BA.2), the protection was 76.2% (20). This highlights the role of prior exposure to

Omicron subvariants in having high protection against the BA.4/5 (21). The BQ and XBB

subvariants, which are now rapidly expanding, are shown to further compromise the efficacy of

current COVID-19 vaccines. The three shots of the ancestral-based vaccine showed far lower

neutralization titers against BQ.1, BQ.1.1, XBB, and XBB.1 compared to the ancestral strain,

with reductions of >37-fold to >71-fold. The variants also showed alarmingly high resistance to

neutralization by serum of persons recently boosted with the new bivalent (covering BA.5)

mRNA vaccines (22). Nonetheless, CDC data showed bivalent booster doses provided 73%

additional protection against COVID-19 hospitalization compared to the ancestral-based

monovalent vaccine through September-November 2022 when BQ.1 was circulating in the US

(23).

To date (as of January 21, 2023), more than 13.2 billion doses of vaccine have been delivered to

people, and 5.52 billion people, i.e. 69.3% of the global population, received at least one dose of

COVID-19 vaccine (Our World in Data; Coronavirus (COVID-19) Vaccinations). However,

almost all those vaccines were designed based on the antigens from the ancestral virus. In this

study, we showed that even three doses of the ancestral-based vaccine could neutralise Omicron

with a geometric mean titer of 27.9; whereas, three doses of the Omicron-based vaccine can

result in 12-fold higher titer. This was in line with our previous pilot animal study in the mice

model to evaluate the neutralization potency of the Omicron-specific vaccine, BIV1-CovIran,

against Omicron variant, which its outline are presented in supplementary material

(Supplementary material 1). We also showed delivering the third dose of the Omicron-based

booster instead of repeating an ancestral-based booster dose can result in a 2- folds higher

neutralisation titer (geometric mean of 27.9 vs. 64). Considering that Omicron and its

sublineages may become the persistent variants that humans might face even when the pandemic

ends (24), this study overwhelmingly recommends including an omicron component in COVID-

19 booster vaccines. Updating current vaccines would shape immune antigenic memory to have

the lowest antigenic distance to the current circulating SARS-CoV-2 subvariants, and makes the

immune system responses compatible with the circulating epitopes. The updated vaccines could

also be applicable in immunologically naïve populations, such as children who reach the age of

eligibility for getting COVID-19 vaccines (6).

On the other hand, viral evolution continues and escape variants emerge, vaccine neutralization

potency against multiple variants is highly desirable (8). In this study, the bivalent vaccine,

without sacrificing potency against the Wuhan variant, achieved significantly higher neutralising

titers against Omicron than the ancestral-based vaccine. Two doses of 5 µg bivalent vaccine (2.5

µg each) neutralised both Omicron and Wuhan variants by the geometric mean titer of 97.

Interestingly, the titer against Omicron equals the titer elicited by two doses of the Omicron-

based vaccine and is 13-fold higher than the titer elicited by two doses of the Wuhan-based

vaccine. Delivering the third dose of bivalent vaccine increased the titer by more than 2-folds for

both Omicron and Wuhan variants. Moreover, delivering the 10 µg bivalent vaccine (5 µg each)

resulted in even higher neutralisation titers against Omicron and Wuhan variants in both two-

dose and three-dose regimens. The broad neutralising antibody response shown in this study is in

accordance with findings from bivalent vaccine boosters of Moderna in preclinical studies (25).

The bivalent Omicron-containing vaccine candidate of Moderna not only elicited higher spike-

binding antibody responses against omicron BA.1 than mRNA-1273 (based on ancestral strain)

but also showed a superior response against BA.4/5 subvariants and ancestral Wuhan-1 variant.

Mechanisms underlying the increased antibody responses with bivalent vaccines have yet to be

elucidated but may include generating new memory immune responses (26).

This study is subjected to some limitations. First, the study evaluated neutralization titers against

the Wuhan-Hu-1 and BA.1 sublineage; but the neutralization potency of the vaccines against

BA.2, BQ.1, and XBB.1 subvariants remained unexplored. However, the prospering preclinical

and real-world results of the Wuhan variant-based (BIV1-CovIran) against pre-Omicron variants,

combined with the high anti-Omicron neutralization titers elicited by the Omicron-based vaccine

(BIV1-CovIran Plus), would be a promise that the bivalent candidate may enhance immunity

against all currently circulating variants (11,13). As the second limitation, non-neutralizing

antibodies and cross-reactive T-cell responses, both of which could influence protective

immunity, were not taken into account in our analysis (27). Future experiments using other

circulating and emerging strains and evaluating other components of immune system may be

informative to determine the breadth of the vaccine protection.

Immunity induced by existing COVID-19 vaccines is subjected to waning over time (28,29),

which in the case of the Omicron is particularly concerning since it is coupled with the variant's

ability to escape the humoral immune response elicited by ancestral variant-based vaccines

(30,31). Altogether, the study iterate that the population's waning immunity against SARS-CoV-

2 should be recovered using Omicron-specific vaccines. This is especially important for Iran's

case, in which the new peak of COVID-19 caused by BA.2, BQ.1, and XBB was officially

declared by the ministry of health on December 30, 2022 (32). Our findings indicate that

Omicron-specific and bivalent vaccines may be a new tool in response to emerging variants,

conserving neutralisation potency against Omicron sublineages. Accordingly, we are planning to

evaluate the Omicron-based and bivalent vaccines according to the latest FDA guidance on the

emergency use authorization for COVID-19 vaccines (33).

Ethical approval

All animal models and maintenance procedures were under the approval of animal ethics

committee guidelines of the Ministry of Health and Medical Education (Tehran, Iran; ethical

code: IR.ACECR.IBCRC.REC.1399.016).

Contributors

Conceptualization: A.A., Hamidreza J., M.T., M.LB., Hassan J.; Data curation: A.A., M.T.;

Formal Analysis: A.A., M.T.,; Funding acquisition: A.A., M.T.; Investigation: Hamidreza J.,

Hassan J.; Methodology: A.A., M.T., Hassan J.; Project administration: Hassan J.; Resources:

Hassan J., Hamidreza J.; Supervision: Hassan J., Hamidreza J.; Validation: Hassan J., Hamidreza

J.; Writing – original draft: A.A.; Writing – review & editing: A.A., Hamidreza J., M.T., M.LB.,

Hassan J.

Funding

Shifa-Pharmed Industrial Group

Competing interests

Mohammad Taqavian and Mehdi Lari Baghal are employees of Shifa Pharmed, with no stock

options or incentives. Hamidreza Jamshidi and Hasan Jalili are the chairman and managing

directors of the vaccine project in Shifa Pharmed, respectively. Asghar Abdoli is the founder and

scientific director of Amirabad Virology Lab.

References

1. Collie S, Champion J, Moultrie H, Bekker L-G, Gray G. Effectiveness of BNT162b2 Vaccine against Omicron Variant in South Africa. N Engl J Med [Internet]. 2021 Dec 29 [cited 2022 Jan 31];386(5):494–6. Available from: https://pubmed.ncbi.nlm.nih.gov/34965358/

2. Tseng HF, Ackerson BK, Luo Y, Sy LS, Talarico CA, Tian Y, et al. Effectiveness of mRNA-1273 against SARS-CoV-2 Omicron and Delta variants. Nat Med 2022 285 [Internet]. 2022 Feb 21 [cited 2022 Nov 17];28(5):1063–71. Available from: https://www.nature.com/articles/s41591-022-01753-y

3. Bok K, Sitar S, Graham BS, Mascola JR. Accelerated COVID-19 vaccine development: milestones, lessons, and prospects. Immunity. 2021 Aug 10;54(8):1636–51.

4. World Health Organization. Weekly epidemiological update on COVID-19 - 19 January 2023. Available from: https://www.who.int/publications/m/item/weekly-epidemiological-update-on-covid-19---19-january-2023

5. WHO. Interim Statement on COVID-19 vaccines in the context of the circulation of the Omicron SARS-CoV-2 Variant from the WHO Technical Advisory Group on COVID-19 Vaccine Composition (TAG-CO-VAC) [Internet]. 2022 [cited 2022 Feb 2]. Available from: https://www.who.int/news/item/11-01-2022-interim-statement-on-covid-19-vaccines-in-the-context-of-the-circulation-of-the-omicron-sars-cov-2-variant-from-the-who-technical-advisory-group-on-covid-19-vaccine-composition

6. Gagne M, Moliva JI, Foulds KE, Andrew SF, Flynn BJ, Werner AP, et al. mRNA-1273 or mRNA-Omicron boost in vaccinated macaques elicits similar B cell expansion, neutralizing responses, and protection from Omicron. Cell [Internet]. 2022 Apr 28 [cited

2022 Nov 5];185(9):1556-1571.e18. Available from:

https://pubmed.ncbi.nlm.nih.gov/35447072/

7.   Fang Z, Peng L, Filler R, Suzuki K, McNamara A, Lin Q, et al. Omicron-specific mRNA

vaccination alone and as a heterologous booster against SARS-CoV-2. Nat Commun 2022

131 [Internet]. 2022 Jun 6 [cited 2022 Dec 4];13(1):1–12. Available from:

https://www.nature.com/articles/s41467-022-30878-4

8.   Chalkias S, Harper C, Vrbicky K, Walsh SR, Essink B, Brosz A, et al. A Bivalent

Omicron-Containing Booster Vaccine against Covid-19. N Engl J Med [Internet]. 2022

Oct 6 [cited 2022 Nov 5]; Available from:

https://www.nejm.org/doi/10.1056/NEJMoa2208343

9.   Lee I-J, Sun C-P, Wu P-Y, Lan Y-H, Wang I-H, Liu W-C, et al. A booster dose of

Delta□×□Omicron hybrid mRNA vaccine produced broadly neutralizing antibody against

Omicron and other SARS-CoV-2 variants. J Biomed Sci [Internet]. 2022;29(1):49.

Available from: https://doi.org/10.1186/s12929-022-00830-1

10.   Mohraz M, Salehi M, Tabarsi P, Abbasi-Kangevari M, Ghamari SH, Ghasemi E, et al.

Original research: Safety and immunogenicity of an inactivated virus particle vaccine for

SARS-CoV-2, BIV1-CovIran: findings from double-blind, randomised, placebo-

controlled, phase I and II clinical trials among healthy adults. BMJ Open [Internet]. 2022

Apr 8 [cited 2022 Nov 5];12(4). Available from: /pmc/articles/PMC8995575/

11.   Mirahmadizadeh A, Heiran A, Lankarani KB, Serati M, Habibi M, Eilami O, et al.

Effectiveness of Coronavirus Disease 2019 Vaccines in Preventing Infection, Hospital

Admission, and Death: A Historical Cohort Study Using Iranian Registration Data During

Vaccination Program. Open forum Infect Dis [Internet]. 2022 Jun 1 [cited 2022 Nov

5];9(6). Available from: https://pubmed.ncbi.nlm.nih.gov/35615300/

12.   Mohraz M, Abbasi-Kangevari M, Ghamari S-H, Ghasemi E, Malekpour M-R, Rezaei N,

et al. Immunogenicity and Safety of a Variant-Specific COVID-19 Vaccine Booster,

BIV1-Coviran Plus: Findings from a Non-Inferiority, Parallel-Design Non-Randomised

Clinical Trial [Internet]. 2022 [cited 2022 Nov 17]. Available from:

https://papers.ssrn.com/abstract=4189405

13.   Abdoli A, Aalizadeh R, Aminianfar H, Kianmehr Z, Teimoori A, Azimi E, et al. Safety

and potency of BIV1-CovIran inactivated vaccine candidate for SARS-CoV-2: A

preclinical study. Rev Med Virol [Internet]. 2022 May [cited 2022 Feb 9];32(3):e2305.

Available from: https://pubmed.ncbi.nlm.nih.gov/34699647/

14.   World Health Organization. Recommendations for the production and control of

poliomyelitis vaccine (oral). WHO Tech Rep Ser [Internet]. 2002 [cited 2022 Feb

18];(No. 904). Available from:

https://www.who.int/publications/m/item/recommendations-for-the-production-and-

control-of-poliomyelitis-vaccine-(oral)-(addendum-2000)-annex-1

15.   van Tol S, Mögling R, Li W, Godeke GJ, Swart A, Bergmans B, et al. Accurate serology

for SARS-CoV-2 and common human coronaviruses using a multiplex approach. Emerg

Microbes Infect. 2020 Jan;9(1):1965–73.

16.   Meyer B, Reimerink J, Torriani G, Brouwer F, Godeke GJ, Yerly S, et al. Validation and

clinical evaluation of a SARS-CoV-2 surrogate virus neutralisation test (sVNT).

https://doi.org/101080/2222175120201835448. 2020 Jan;9(1):2394–403.

17.   Rijkers G, Murk JL, Wintermans B, van Looy B, van den Berge M, Veenemans J, et al.

Differences in Antibody Kinetics and Functionality Between Severe and Mild Severe

Acute Respiratory Syndrome Coronavirus 2 Infections. J Infect Dis. 2020

Oct;222(8):1265–9.

18.   Pulliam JRC, van Schalkwyk C, Govender N, von Gottberg A, Cohen C, Groome MJ, et

al. Increased risk of SARS-CoV-2 reinfection associated with emergence of Omicron in

South Africa. Science (80-) [Internet]. 2022 May 6 [cited 2022 Dec 5];376(6593).

Available from: https://www.science.org/doi/10.1126/science.abn4947

19.   Tuekprakhon A, Nutalai R, Dijokaite-Guraliuc A, Zhou D, Ginn HM, Selvaraj M, et al.

Antibody escape of SARS-CoV-2 Omicron BA.4 and BA.5 from vaccine and BA.1

serum. Cell. 2022 Jul 7;185(14):2422-2433.e13.

20.   Altarawneh HN, Chemaitelly H, Ayoub HH, Hasan MR, Coyle P, Yassine HM, et al.

Protective Effect of Previous SARS-CoV-2 Infection against Omicron BA.4 and BA.5

Subvariants. N Engl J Med [Internet]. 2022 Oct 5;387(17):1620–2. Available from:

https://doi.org/10.1056/NEJMc2209306

21.   Hansen CH, Friis NU, Bager P, Stegger M, Fonager J, Fomsgaard A, et al. Risk of

reinfection, vaccine protection, and severity of infection with the BA.5 omicron

subvariant: a nation-wide population-based study in Denmark. Lancet Infect Dis

[Internet]. 2022 Oct [cited 2022 Dec 5];0(0). Available from:

http://www.thelancet.com/article/S1473309922005953/fulltext

22.   Wang Q, Iketani S, Li Z, Liu L, Guo Y, Huang Y, et al. Alarming antibody evasion

properties of rising SARS-CoV-2 BQ and XBB subvariants. Cell. 2022 Dec;

23.   Surie D, DeCuir J, Zhu Y, Gaglani M, Ginde AA, Douin DJ, et al. Early Estimates of

Bivalent mRNA Vaccine Effectiveness in Preventing COVID-19–Associated

Hospitalization Among Immunocompetent Adults Aged ≥65 Years — IVY Network, 18

States, September 8–November 30, 2022. MMWR Morb Mortal Wkly Rep [Internet].

2022 Dec 30 [cited 2023 Jan 20];71(5152):1625–30. Available from:

https://www.cdc.gov/mmwr/volumes/71/wr/mm715152e2.htm

24.   Fauci: Omicron-specific vaccines "prudent" but may be unnecessary | TheHill [Internet].

[cited 2022 Jan 28]. Available from: https://thehill.com/policy/healthcare/591241-fauci-

omicron-specific-vaccines-prudent-but-may-be-unnecessary

25.   Scheaffer SM, Lee D, Whitener B, Ying B, Wu K, Liang C-Y, et al. Bivalent SARS-CoV-

2 mRNA vaccines increase breadth of neutralization and protect against the BA.5

Omicron variant in mice. Nat Med [Internet]. 2022; Available from:

https://doi.org/10.1038/s41591-022-02092-8

26.   Gaebler C, Wang Z, Lorenzi JCC, Muecksch F, Finkin S, Tokuyama M, et al. Evolution

of antibody immunity to SARS-CoV-2. Nature [Internet]. 2021;591(7851):639–44.

Available from: https://doi.org/10.1038/s41586-021-03207-w

27.   Errico JM, Adams LJ, Fremont DH. Antibody-mediated immunity to SARS-CoV-2 spike.

Adv Immunol. 2022;154:1–69.

28.   Levin EG, Lustig Y, Cohen C, Fluss R, Indenbaum V, Amit S, et al. Waning Immune

Humoral Response to BNT162b2 Covid-19 Vaccine over 6 Months. N Engl J Med

[Internet]. 2021 Dec 9 [cited 2022 Dec 6];385(24):e84. Available from:

https://pubmed.ncbi.nlm.nih.gov/34614326/

29.   Doria-Rose N, Suthar MS, Makowski M, O'Connell S, McDermott AB, Flach B, et al.

Antibody Persistence through 6 Months after the Second Dose of mRNA-1273 Vaccine

for Covid-19. N Engl J Med [Internet]. 2021 Jun 10 [cited 2022 Dec 6];384(23):2259–61.

Available from: https://pubmed.ncbi.nlm.nih.gov/33822494/

30.   Cao Y, Wang J, Jian F, Xiao T, Song W, Yisimayi A, et al. Omicron escapes the majority

of existing SARS-CoV-2 neutralizing antibodies. Nature [Internet]. 2022 Feb 24 [cited

2022 Dec 6];602(7898):657–63. Available from:

https://pubmed.ncbi.nlm.nih.gov/35016194/

31.   Planas D, Saunders N, Maes P, Guivel-Benhassine F, Planchais C, Buchrieser J, et al.

Considerable escape of SARS-CoV-2 Omicron to antibody neutralization. Nature

[Internet]. 2022 Feb 24 [cited 2022 Dec 6];602(7898):671–5. Available from:

https://pubmed.ncbi.nlm.nih.gov/35016199/

32.   Financial Tribune. Medical Community on High Alert to Tackle New Covid Wave

[Internet]. 2023. Available from: https://financialtribune.com/node/116667

33.   Food and Drug Administration. Emergency Use Authorization for Vaccines to Prevent

COVID-19. 2022 [cited 2022 Feb 9];(March):1–31. Available from:

https://www.fda.gov/media/142749/download

Figure legends

Figure 1 (A) The virus growth curve analysed by TCID50, (B) positive CPE resulting from virus infection, (C) CPE in the control sample.

Figure 2 The neutralisation titer of samples against the Omicron and Wuhan variants (A) The neutralisation titer against the Wuhan variant in groups that received two doses of vaccines (B) The neutralisation titer against the Omicron variant in groups that received two doses of vaccines (C) The neutralisation titer against the Wuhan variant in groups that received three doses of vaccines (D) The neutralisation titer against the Omicron variant in groups that received three doses of vaccines

WW: Two doses of BIV1-CovIran; OO: Two doses of BIV1-CovIran Plus; WWW: Three doses of BIV1-CovIran; OOO: Three doses of BIV1-CovIran Plus; WWO: BIV1-CovIran Plus booster after two doses of BIV1-CovIran; BB (2.5): Two doses of 5µg BBIV1- CovIran; BB (5): Two doses of 10µg BBIV1- CovIran; BBB (2.5): Three doses of 5µg BBIV1- CovIran; BBB (5): Three doses of 10µg BBIV1- CovIran

Figure 3 The geometric means of the neutralisation activity of study groups against the Omicron and Wuhan variants.

WW: Two doses of BIV1-CovIran; OO: Two doses of BIV1-CovIran Plus; WWW: Three doses of BIV1-CovIran; OOO: Three doses of BIV1-CovIran Plus; WWO: BIV1-CovIran Plus booster after two doses of BIV1-CovIran; BB (2.5): Two doses of 5µg BBIV1- CovIran; BB (5): Two doses of 10µg BBIV1- CovIran; BBB (2.5): Three doses of 5µg BBIV1- CovIran; BBB (5): Three doses of 10µg BBIV1- CovIran