Kinase Suppressor of Ras 2 promotes self-renewal and clonogenicity of small-cell lung carcinoma

Dianna H. Huisman[1], Deepan Chatterjee[1], Robert A. Svoboda[2], Heidi M. Vieira[1], Abbie S. Ireland[3], Sydney Skupa[1], James W. Askew[1], Danielle E. Frodyma[1#], Luc Girard[4], Kurt W. Fisher[2], Michael S. Kareta[5], John D. Minna[4,6], Trudy G. Oliver[7], Robert E. Lewis[1]*

[1]Eppley Institute, Fred & Pamela Buffett Cancer Center, University of Nebraska Medical Center, Omaha, NE 68198, USA.

[2]Department of Pathology, Microbiology, and Immunology, University of Nebraska Medical Center, Omaha, NE 68198, USA.

[3]Deparment of Molecular Cancer Biology, Duke University, Durham, NC, USA.

[4]Hamon Center for Therapeutic Oncology Research, Department of Pharmacology, Simmons Cancer Center, University of Texas Southwestern Medical Center, Dallas, Texas.

[5]Cancer Biology and Immunotherapies Group, Sanford Research, Sioux Falls, South Dakota.

[6]Department of Internal Medicine, University of Texas Southwestern Medical Center, Dallas, Texas.

[7]Deparment of Pharmacology and Cancer Biology, Duke University, Durham, NC, USA.

[#]Current address: Porterhouse Medical, Holmdel, NJ

*Correspondence: rlewis@unmc.edu

Abstract

Small-cell lung carcinoma (SCLC) tumors are heterogeneous, with a subpopulation of cells primed for tumor initiation. Here, we show that Kinase Suppressor of Ras 2 (KSR2) promotes the self-renewal and clonogenicity of SCLC cells. KSR2 is a molecular scaffold that promotes Raf/MEK/ERK signaling. KSR2 is preferentially expressed in the ASCL1 subtype of SCLC (SCLC-A) tumors and is expressed in pulmonary neuroendocrine cells, one of the identified cells of origin for SCLC-A tumors. The expression of KSR2 in SCLC and pulmonary neuroendocrine cells (PNECs) was previously unrecognized and serves as a novel model for understanding the role of KSR2-dependent signaling in normal and malignant tissues. Disruption of KSR2 in SCLC-A cell lines inhibits the colony forming ability of tumor propagating cells (TPCs) *in vitro* and their tumor initiating capacity *in vivo.* The effect of KSR2 depletion on self-renewal and clonogenicity is dependent on the interaction of KSR2 with ERK. These data indicate that the expression of KSR2 is an essential driver of SCLC-A tumor propagating cell function, and therefore may play a role in SCLC tumor initiation. These findings shed light on a novel effector promoting initiation of ASCL1-subtype SCLC tumors, and a potential subtype-specific therapeutic target.

Impact Statement

Manipulation of the molecular scaffold KSR2 in ASCL1-subtype small-cell lung cancer cells reveals its contribution to self-renewal, clonogenicity, and tumor initiation via ERK signaling.

Introduction

Small-cell lung carcinoma (SCLC) affects current and former heavy smokers, accounting for 13% of all lung cancers[1]. There have been few improvements in SCLC detection, treatment, and survival in almost 40 years, leading to its classification as a recalcitrant cancer in 2012[1]. The five-year relative survival rates for SCLC patients with localized, regional, and distant disease are 30%, 18%, and 3%, respectively (American Cancer Society, 2024). Currently, SCLC patients are treated with first line therapy (cisplatin or carboplatin combined with etoposide chemotherapy, plus anti-PDL1 antibody, atezolizumab), second (topotecan), and third line (PD1 antagonist, nivolumab) therapies, and for some patients thoracic radiation therapy[2]. Although SCLC tumors are responsive to therapy initially, residual disease quickly develops resistance leading to the low five-year survival[3]. Substantial efforts have been made to characterize SCLC tumors and identify targets that may be selectively toxic to tumor cells while preserving normal lung tissue[4-8]. Rigorous and innovative basic science using state-of-the-art genetically-engineered mouse (GEM) models, and an extensive set of cell lines have led to key discoveries regarding the cells-of-origin and the common recurring mutations that underlie SCLC[7, 9-18]. These discoveries have yielded comprehensive genomic profiles and a durable classification of SCLC subtypes based on the differential expression of four key transcription factors, ASCL1 (SCLC-A), NEUROD1 (SCLC-N), POU2F3 (SCLC-P) and YAP1 (SCLC-Y)[7, 13, 17, 18].

Lineage tracing and single cell RNA sequencing (scRNA-seq) in genetically modified mice showed that a rare pulmonary neuroendocrine cell (PNEC) subpopulation, NE^{stem} cells, actively responds to lung injury of the epithelia by expanding, migrating, and undergoing Notch-dependent transit amplification to regenerate the damaged epithelium[19]. This effort additionally identified NE^{stem} as a cell-of-origin for SCLC following Trp53, Rb1 and Notch mutations causing constitutive activation of stem cell renewal and altered deprogramming[19]. SCLC tumors in some GEM models have a small population of tumor propagating cells (TPCs) essential to the initiation, long-term propagation, and metastatic capacity of the tumor, while the bulk non-TPC population is highly proliferative but incapable of establishing tumors *in vivo*[20]. TPCs have been implicated in the initiation and growth of SCLC as well as therapy resistance[21-27]. Tumor propagating cells are also implicated in epithelial-to-mesenchymal transition (EMT) and metastasis[27]. Their slower cycling and self-renewing ability enhances DNA repair, rendering these cells resistant to DNA damage-dependent chemo and radiation therapy[23, 24, 27, 28]. Thus, SCLC TPCs offer a unique population within which to search for new targets, which in combination with current standard-of-care therapies may yield a durable and effective strategy for therapy.

Kinase Suppressor of Ras proteins KSR1 and KSR2 are molecular scaffolds for Raf/MEK/ERK signaling[29, 30]. While KSR1 is widely expressed, KSR2 expression is restricted to the brain, pituitary, adrenal glands, and neuroendocrine tissues[31, 32]. KSR2 is abundantly expressed in the brain, and its disruption reduces body temperature, promotes cold intolerance, impairs glucose homeostasis, elevates fasting insulin and free fatty acid levels in the blood, and causes obesity[30]. ChIP-seq analysis also revealed *KSR2* as a transcriptional target in human ASCL1-subtype SCLC (SCLC-A) cell lines[33, 34]. KSR2 was identified as one of 24 druggable and overexpressed target genes of ASCL1 identified by ChIP-seq, providing a rationale for studying the role of KSR2 in SCLC-A[35]. Our work shows that KSR2 is expressed in PNECs and SCLC-A tumors and cell lines. Expression of KSR2 promotes the self-renewal and clonogenicity of SCLC-A TPCs and their tumor initiating capacity *in vitro* and *in vivo*. While KSR2 disruption is rescued by an intact KSR2 transgene, the inability of an ERK-binding deficient KSR2 construct to restore clonogenicity and tumor initiation implicates its scaffolding function in TPC formation. This result defines a novel mechanism of tumor initiation in SCLC-A and a potential therapeutic vulnerability.

Results

Pulmonary neuroendocrine cells and ASCL1-subtype SCLC tumors express KSR2. ASCL1-subtype (SCLC-A) tumors can arise from PNECs[19, 33, 35, 36]. PNECs are heterogeneous, including a small subpopulation termed NE^{stem} cells that respond to injury of the lung epithelia by expanding, migrating to the location of the damage, and undergoing Notch-dependent transit amplification to regenerate the damaged epithelium[19]. SCLC tumors may arise from PNECs following *Trp53*, *Rb1*, and *Notch* mutations causing constitutive activation of stem cell renewal and altered deprogramming[19]. Although *KSR2* mRNA is not detectable in normal epithelial lung tissue (Fig. 1A), it is present in PNECs (Fig. 1B). We examined expression of *Ksr2* in an established *Ascl1*-high *Rb1/p53/Rbl2* (RPR2) model as well as a MYC-driven SCLC model (*Rb1/p53/Myc*, RPM) associated with reduced *Ascl1* and the acquisition of other SCLC subtypes including *Neurod1* and *Yap1* (Fig.

1C)[37, 38]. *Ksr2* was highly enriched in RPR2 tumors and significantly depleted in RPM tumors when *Ascl1* was genetically depleted (RPMA), demonstrating a dependency of *Ksr2* expression on ASCL1. Consistent with the mouse tumor data, analysis of human SCLC tumors showed that ASCL1-high and MYC-low tumors[13] preferentially express *KSR2* mRNA (Fig. 1D). Analysis of *KSR2* mRNA by subtype reveals that *KSR2* is highly expressed in SCLC-A, but has varying expression in NEUROD1, POU2F3, and YAP1 subtypes (Fig. 1E). While *KSR2* mRNA is preferentially expressed in SCLC-A compared to other subtypes[7], *KSR1* mRNA is detected across SCLC subtypes (Supplementary Table 1). SCLC-A and -N cells express KSR1, but only SCLC-A cells express KSR2 by western blot analysis (Fig. 1F, 1G).

Depletion of Kinase Suppressor of Ras 2 reduces SCLC clonogenicity and self-renewal. SCLC-A tumor propagating cells (TPCs) are essential for tumor initiation and metastasis and are enriched by selection for high expression of CD24 and EpCAM, but low expression of CD44[20]. Transplantation assays show that TPCs are a minor but highly tumorigenic subpopulation of SCLC cells characterized by their colony forming activity *in vitro*, which measures their capacity for clonogenicity and self-renewal[20, 39]. Mouse SCLC cell lines KP1 and KP3 were derived from spontaneous mouse models of SCLC with knockout of Rb1, Trp53 (KP1) or Rb1, Trp53 and p130 (KP3), which replicate the ASCL1 subtype of SCLC tumors[7, 11]. KP1 and KP3 cells expressing one of three doxycycline (dox)-inducible shRNAs targeting *Ksr2* (sh5, sh6, sh7) were treated with or without doxycycline (Fig. 2A). Human ASCL1-subtype SCLC cell line NCI-H209 (H209) expressing dox-inducible shRNA targeting *KSR2* (sh5) was also treated with or without doxycycline (Fig. 2A). KP1 and H209 cells infected with control (CTRL) or KSR2 knockdown (sh5) shRNA encoding viruses were stained for CD24^{high}CD44^{low}EpCAM^{high} (Fig. 2B, C). Dox-induced targeting of *KSR2* reduced the proportion of CD24^{high}CD44^{low}EpCAM^{high} TPCs detected by FACS in H209 cells from 42.4% to 0.4%, and in KP1 cells from 39.1 to 5.4% (Fig. 2D). Following downregulation of KSR2 expression, KP3 cells were stained for TPC markers CD24^{high}CD44^{low}EpCAM^{high}, isolated by FACS and plated as single cells in 96-well plates to be analyzed for colony formation by CellTiter-Glo®. Robust colony formation, a measure of self-renewing capability of an individual cell[40-42], was observed in KP3 control TPCs. Viability was reduced by 58% and 45% with *Ksr2* RNAi by sh6 and sh7, respectively, which reflects their effectiveness at targeting *Ksr2* (Fig. 2A). In KP1 cells, colony formation was assessed with or without dox-induced RNAi by hairpin sh5 after TPC isolation, which inhibited colony formation by 86% (Fig. 2F). These data demonstrate that KSR2 is a significant contributor to the clonogenic and self-renewing properties of SCLC TPCs.

KSR2 disruption inhibits the tumor initiating capacity of murine SCLC cells *in vivo*. Extreme limiting dilution analysis (ELDA) is a statistical method optimized for estimating the stem cell frequency from limiting dilution analysis[43]. Stem cell frequency within the bulk tumor cell population is determined from the frequency of tumor-positive and tumor-negative injections arising from varying doses of xenografted cells. Dox-inducible shRNA targeted KP1 cells were injected with successive dilutions into NOD-*Prkd*^{cem26Cd52}*Il2rg*^{em26Cd22}/NjuCrl (NCG) mice, and the mice were provided drinking water with sucrose, or sucrose plus doxycycline (2 mg/kg). ELDA was performed by scoring each tumor that arose as "1". The absence of tumor formation was scored as "0". KSR2 disruption reduced frequency of TPCs 10-fold, from 1/255 control KP1 cells to 1/2530 KP1 cells with KSR2 KD (Fig. 3, Table 1). These data indicate that KSR2 is a critical effector of tumor initiation of SCLC tumor propagating cells *in vivo*.

Expression of a KSR2 transgene restores colony formation and increases TPC frequency. To confirm on-target effect of our inducible shRNA system for KSR2 knockdown, a construct containing wildtype *Ksr2* mutated to be resistant to binding hairpin sh5 (KSR2r) was expressed in KP1 sh5 cells (Fig. 4A). Rescue of KSR2 expression restored colony formation to wildtype levels in KP1 sh5 cells confirming the on-target effect of hairpin sh5 (Fig. 4B). *In vitro* ELDA was used to estimate the relative frequency of TPCs. KSR2 disruption reduced TPC frequency from 1/19 in control KP1 cells to 1/166 in KP1 cells with KSR2 KD (Fig. 4C, D). The expression of wildtype *Ksr2* construct KSR2r restored the TPC frequency to 1/18 (Fig. 4C, D). These data demonstrate the on-target action of the inducible *Ksr2* shRNA system to regulate TPC abundance.

KSR2 disruption inhibits the tumor initiating capacity of human SCLC cell lines *in vitro*. *In vitro* ELDA was used to test the effect of doxycycline (dox)-induced *KSR2* disruption on human SCLC cell line H209. The TPC frequency of control H209 cells was reduced 4- to 25-fold from 1/3-1/5 in control cells to 1/12-1/76 in H209 sh5 cells (Fig. 5A, Supplementary Fig. 1A, B). KSR2 was also targeted using a CRISPR/Cas9 in conjunction with a *KSR2* sgRNA (CR3), or non-targeting control (NTC) (Fig. 5B). The TPC frequency was

reduced 3- to 12-fold from 1/10-1/39 in NTC to 1/112-1/140 in KSR2 KO cells (Fig. 5C, Supplementary Fig. 1C, D). These data indicate that KSR2 is a critical effector of tumor initiating capacity in human SCLC tumor propagating cells.

KSR2-ERK interaction is necessary for SCLC clonogenicity. KSR2 knockdown reduces activation of ERK (Fig. 6A). With serum starvation, control and KSR2 knockdown samples are reduced to a basal level of ERK activation (Fig. 6A). ERK can be stimulated with addition of serum in control cells, while KSR2 knockdown cells activate ERK poorly after serum addition (Fig. 6A). Treatment with calcium ionophore ionomycin also induces ERK activation in control cells, but KSR2 knockdown inhibits this response (Fig. 6A). Cells expressing KSR2r rescue ERK activation in each condition (Fig. 6B). To assess the importance of the interaction between KSR2 and ERK, a DEF domain mutant (570 FIFP/AAAP 573)[29] KSR2 construct deficient in its ability to bind ERK (FIF570) was further mutated to be resistant to binding hairpin sh5 (FIF570r) and expressed in KP1 sh5 cells. Cells expressing FIF570r are unable to rescue ERK activation when endogenous KSR2 is depleted (Fig. 6C). Disruption of KSR2/ERK interaction prevents ERK activation in response to serum stimulus or treatment with ionomycin. Immunoprecipitation of the FLAG epitope tag on this construct and the full length KSR2 demonstrated reduced 6hosphor-ERK associated with the ERK-binding mutant construct (FIF570r) (Fig. 6D). Colony formation was significantly reduced in the FIF570r cells after dox-induced targeting of endogenous KSR2, suggesting that KSR2 interaction with ERK is necessary for the clonogenic capacity of SCLC TPCs (Fig. 6E). Disruption of KSR2 significantly reduced the number of colonies formed by TPCs (Fig. 6F). KSR2r rescued colony formation (Fig. 6F), while FIF570r was unable to restore colony formation to KP1 cells (Fig. 6F). These data indicate that KSR2-dependent ERK signaling is an important contributor to colony formation by TPCs.

Discussion

Kinase Suppressor of Ras proteins KSR1 and KSR2 have unique and overlapping functions. Both function as scaffolds for Raf/MEK/ERK signaling[29, 30] promoting phosphorylation of ERK. KSR2 is expressed in the brain, pituitary gland, and adrenal gland, and some neuroendocrine cells and tissues[31, 32, 44]. Here we show that disruption of KSR2 in SCLC-A reduces TPC clonogenicity and self-renewal *in vitro*, and tumor initiation *in vivo*. Further work demonstrates that the clonogenic capacity of TPCs depends upon the interaction of KSR2 with ERK. Activating Ras mutations are rare in SCLC tumors, therefore Raf/MEK/ERK signaling is likely activated by extracellular stimuli, suggesting that a molecular scaffold, such as KSR2, is required to amplify this signaling pathway[45]. The role of ERK signaling in SCLC is incompletely understood, but is implicated in cell proliferation, differentiation, survival, and drug resistance[46]. Our data reveal KSR2-dependent ERK signaling is involved in self-renewal and clonogenicity of the SCLC-A TPC population. This observation contrasts with previous studies that tested the efficacy of targeting Raf/MEK/ERK signaling in SCLC and concluded that ERK activation can be both pro- and anti-proliferative toward SCLC[45]. Treatment with an ERK inhibitor was unable to induce apoptosis in human SCLC-A cell lines H209 and H69[47]. In contrast, our data are consistent with observations that SCLC-A cell line proliferation was significantly reduced *in vitro* and in *in vivo* tumor xenografts by ARHGEF19 disruption and downstream reduced Raf/MEK/ERK signaling[48]. Activation of Raf/MEK/ERK by endoplasmic reticulum (ER) stress has also been reported to promote SCLC cell survival[49]. Raf/MEK/ERK signaling may play an essential role in promoting metastasis of SCLC tumors[45]. CXCL12 induces ERK activation in SCLC cells, which correlates with increased invasion through extracellular matrix[50]. Increased Raf/MEK/ERK signaling induced by expression of an activated Raf construct has been reported to cause growth arrest and reduce expression of neuroendocrine markers in SCLC[51, 52]. These contrasting observations may suggest that the intensity, frequency, localization, or duration of ERK activation and its effects may be context dependent. Further, our findings suggest that Raf/MEK/ERK signaling may be selectively important for the TPC population, while dispensable for the bulk tumor population.

The effect of KSR2 depletion on ERK activation is surprising as SCLC-A cells express both KSR2 and KSR1. We do not detect KSR1 compensation for KSR2 loss, suggesting that KSR2 is required for TPC function in SCLC-A cells. Our data suggest that either KSR2 is contributing to TPC function by promoting signaling that KSR1 cannot influence, or that KSR2 and KSR1 coordinate function to conduct efficient signaling. KSR1 and KSR2 are capable of heterodimerization with Raf, resulting in a conformational change of KSR proteins that allows phosphorylation of MEK[53-55]. The dimerization of KSR proteins with Raf orients the Raf protein so that its catalytic site is not in close proximity to the phosphorylation site on MEK to complete the phosphorylation,

which necessitates MEK phosphorylation by Raf through a *trans* interaction[53]. KSR2-BRAF heterodimerization results in an increase of MEK phosphorylation via the KSR2-mediated relay of signal from BRAF to release the activation segment of MEK for phosphorylation[53]. KSR2 can also homodimerize via the same interface that interacts with BRAF, however this creates a different quaternary structure when interacting with MEK and it's unknown how this may affect the availability of MEK for phosphorylation[53]. BRAF homodimers, or KSR1-BRAF heterodimers, should generate a quaternary structure sufficient for availability of MEK for phosphorylation by an additional BRAF protein[53]. Via the same interaction interface, it is possible that KSR1 and KSR2 heterodimerize. Our incomplete understanding of how KSR2 homodimers or KSR1-KSR2 heterodimers may impact MEK phosphorylation makes it difficult to predict the effect of KSR2 loss on MEK phosphorylation and downstream ERK activation. However, KSR2 depletion has a deleterious effect on ERK activation in SCLC-A cells. It is notable that we fail to detect KSR2 expression in the NEUROD1-subtype of SCLC (SCLC-N), while they retain expression of KSR1 (Fig. 1E, F). This observation suggests that SCLC-N cells are capable of compensating for loss of KSR2 with KSR1 expression, or that transition to the NEUROD1-subtype SCLC results in upregulation of other compensatory signaling pathways that negate the dependency on KSR2 for ERK signaling.

ASCL1 and NEUROD1 drive distinct transcriptional profiles in SCLC[14, 56, 57]. SCLC-A cells can be converted to SCLC-N cells by elevation of MYC[57]. MYC amplification has been associated with poor patient outcome, therapy resistance, and tumor progression[58-60]. KSR2 expression seems to be lost during transition from SCLC-A to SCLC-N. The initiation of switching from SCLC-A subtype to SCLC-N subtype is incompletely understood, although recent studies implicate that epigenetic mechanisms are involved[61]. The expression of KSR2 in SCLC-A is thought to be driven by its defining transcription factor ASCL1, as *KSR2* is a transcriptional target of ASCL1[33, 34]. ASCL1 and KSR2 are also expressed in SCLC-A cell-of-origin, PNECs. PNECs respond to injury of the lung epithelia by expanding, migrating, and undergoing Notch-dependent transit amplification to regenerate the damaged epithelium[19]. Although KSR2 plays a role in SCLC TPC function, its role in normal PNECs is undetermined. KSR2 knockout mice have metabolic defects[30, 31] but have no reports of abnormal lung pathology. Studies determining the effect of KSR2 manipulation on lung repair are required to determine if KSR2 plays a role in the function of PNECs in response to damage.

If KSR2 is dispensable for normal lung repair, it represents a potential therapeutic target for SCLC-A tumors. The discovery that cell surface proteins CD24, CD44, and EpCAM can be used to enrich and isolate SCLC-A TPCs significantly advanced investigation of potential therapeutic vulnerabilities within this population[19]. Cisplatin treatment does not enrich TPCs indicating that TPCs are not a reservoir for drug tolerant persister cells[19], however TPCs are capable of repopulating tumors after treatment suggesting that they are not preferentially sensitive to the chemotherapeutic. Isolation of TPCs allows for interrogation of potential therapies that may be selectively detrimental to the TPC population. The distinct transcriptional profiles of the SCLC subtypes are therapeutically relevant, as there are a number of MYC-associated vulnerabilities in ASCL1 low tumors which differ from vulnerabilities identified in ASCL1 high tumors[62]. Subtype specific vulnerabilities have been reported for all four SCLC subtypes[63] suggesting that further investigation into subtype specific therapeutic options, and their clinical relevance, is necessary. The preferential expression of KSR2 in SCLC-A, and the role it plays in TPC function provide rationale for further interrogating KSR2 as a SCLC-A specific therapeutic target.

JQ1, an inhibitor of bromodomain and extra-terminal (BET) proteins, reduced the frequency of TPCs *in vivo* and suppressed the ability of single TPCs to form colonies *in vitro*[19]. Our data demonstrate that KSR2 disruption reduces frequency of TPCs and reduces colony formation *in vitro* and tumor initiation *in vivo*. The effect of KSR2 on TPC clonogenicity is dependent on interaction with ERK, revealing a role for ERK activation in TPC function. These results suggest that targeting of KSR2 or its effector ERK may be TPC-specific therapeutic strategies and may create a more durable therapeutic response when used in conjunction with standard of care chemotherapies. The data provides a rationale for interrogating the effects of targeting KSR2 in genetically engineered mouse models of SCLC-A[12, 14, 64]. KSR2 disruption may work well in combination with cisplatin to target TPCs, reducing tumor burden and recurrence.

Materials and Methods

Cell culture

Murine small-cell lung carcinoma cell lines KP1 and KP3 were a gift from J. Sage (Stanford University). Human small-cell lung carcinoma cell lines H209 and H1963 were a gift of John Minna (UT Southwestern). The cells were cultured in RPMI 1640 medium containing 10% fetal bovine serum (FBS) and grown at 37°C with ambient O_2 and 5 % CO_2. Serum starved cells were grown in above conditions, with serum removed for 16 hours prior to cell lysis. Serum shocked cells were grown normally followed by boosting FBS concentration to 50% for 15 minutes prior to cell lysis. Ionomycin (ThermoFisher Scientific I24222) treatment was given at a 1 uM dose for 15 minutes prior to cell lysis. All cells were routinely tested for mycoplasma. No further authentication of cell lines was performed by the authors.

Generation of KSR2 shRNA knockdown and CRISPR/Cas9 knockout cell lines
Individual SMARTvector human inducible lentiviral shRNAs suitable for targeting murine and human KSR2 expressed in piSMART hEF1a/TurboGFP vector were stably transfected into KP1, KP3, and H209 SCLC cell lines with PEI. Cells were selected for expression of the shRNAs using 0.25 μg/mL of puromycin. 48 hours after doxycycline induction, cells were selected again by flow cytometry sorting for GFP+ cells. Knockdown of KSR2 was confirmed by western blot. KSR2 cDNA (MSCV KSR2 IRES YFP) and ERK binding deficient KSR2 cDNA (MSCV KSR2 FIF570 IRES YFP) was made resistant to binding of hairpin sh5 by introducing three point mutations in the binding region. Point mutations were introduced using the QuikChange Lightning Site Directed Mutagenesis Kit (Agilent #210518) according to the manufacturer's protocol. Sequencing of the construct confirmed the correct point mutations were made with no additional mutations. MSCV KSR2 IRES YFP resistant to binding hairpin sh5 (KSR2r) and MSCV KSR2 FIF570 IRES YFP resistant to binding hairpin sh5 (FIF570r) were transfected into HEK-293T cells using trans-lentiviral packing system (ThermoFisher Scientific). The virus was collected 48 hours post transfection and used to infect KP1 sh5 cells with 8 μg/mL Polybrene for 72 hours. KP1 sh5 cells expressing the KSR2r or FIF570r construct were selected for using flow cytometry sorting YFP+ cells. Presence of the KSR2r or FIF570r expression after doxycycline induced downregulation on endogenous KSR2 was confirmed via western blotting. To generate knockout cells, sgRNA sequences targeting KSR2 or non-targeting control (NTC) were inserted into lentiCRISPR v2 (Addgene #52961). The constructs were PEI transfected into HEK293T cells along with psPAX2 lentiviral packaging construct (Addgene #12259) and pMD2.G envelope construct (Addgene #12259). Lentivirus-containing media was harvested at 72-h and used to infect the H209 cells with 8 μg/mL Polybrene. Cells were selected for expression of the sgRNAs using 2 ug/mL of puromycin, and western blot was used to confirm KSR2 knockout.

Cell lysis and western blot analysis
Whole cell lysate was extracted in radioimmunoprecipitation assay (RIPA) buffer containing 50 mM Tris-HCl, 1% NP-40, 0.5 % Na deoxycholate, 0.1 % Na dodecyl sulfate, 150 mM NaCl, 2 mM EDTA, 2 mM EGTA, and protease and phosphatase inhibitors aprotinin (0.5 U/mL), leupeptin (20 mM), and NA_3VO_4 (0.5 mM). The estimation of protein concentration was done using BCA protein assay (Promega #PI-23222, PI-23224). Samples were diluted using 1 X sample buffer (4 X stock, LI-COR #928–40004) with 100 mM dithiothreitol (DTT) (10 X stock, 1 mM, Sigma #D9779-5G). The protein was separated using 8–12% SDS-PAGE and transferred to nitrocellulose membrane. The membrane was blocked with Odyssey TBS blocking buffer (LICOR-Biosciences #927–50003) for 45 min at room temperature, then incubated with primary antibodies (*Key Resources Table*) at least overnight at 4°C. IRDye 800CW and 680RD secondary antibodies (LI-COR Biosciences # 926–32211, # 926–68072) were diluted 1:10,000 in 0.1% TBS-Tween and imaged on the Odyssey Classic Scanner (LI-COR Biosciences).

Analysis of KSR2 expression in normal tissues and mouse tumors
GTEx portal was used to display the relative expression of human *KSR2* mRNA (TPM) in brain-cortex and lung tissue. In a murine model, neuroendocrine specific reporter, *Chga*-GFP was used to identify PNECs. GFP+ neuroendocrine cells from 3 mouse lungs were isolated by flow cytometry. qPCR was performed to measure mRNA expression of *Ksr2*, *Cgrp*, *Chga*, *Syp*, and *Spc* in GFP+ neuroendocrine cells and GFP- lung epithelial cells. RPR2, RPM and RPMA mouse tumor data is derived from Ireland *et al.*, Cancer Cell, 2020[37] and Olsen *et al.*, Genes Dev, 2021[38] and deposited in NCI GEO: GSE149180 and GSE155692.

SCLC sequencing analysis
RNA sequencing data from human primary tumor samples[13] was analyzed for *KSR2* expression based on high and low *ASCL1* expression or high and low *MYC* expression. RNA sequencing data of human SCLC cell lines (DepMap Portal, Broad Institute MIT) was segregated by SCLC subtype and analyzed for *KSR2* mRNA levels.

RNA samples from SCLC (n = 69) and NSCLC-NE (n = 14) cell lines were analyzed by paired-end RNA sequencing as previously described (McMillan *et al.*, 2018, Cell 173, 864–878; dbGaP study accession: phs001823.v1.p1). Briefly, reads were aligned to the human reference genome GRCh38 using STAR-2.7 (https://github.com/alexdobin/STAR [github.com]) and FPKM values were generated with cufflinks-2.2.1 (http://cole-trapnell-lab.github.io/cufflinks/ [cole-trapnell-lab.github.io]). All data were then pooled, normalized to Transcripts Per Millions (TPM), and log-transformed. RNAseq from the SCLC primary tumors (George *et al.*, 2015, Nature 524, 47-53) were included in the analysis.

Fluorescence Activated Cell Sorting

SCLC cell lines were incubated 20 minutes on ice in PBS with DAPI (3uM), PE-CD24 (1:400) PE-Cy7-CD44 (1:300) and APC-EpCAM (1:100). Cells were resuspended in PBS and flow cytometry of SCLC cell lines was performed using a 100 µm nozzle on a BD FACSAria II using FACSDiva software. Debris were excluded by gating on forward scatter area versus side scatter area. Doublets were excluded by gating on forward scatter area versus side scatter height. Viable cells were identified by exclusion of DAPI stained cells. CD24^{high}CD44^{low} cells were included by sequential gating followed by EpCAM^{high} TPCs. Compensation was performed using single stain and fluorescence- one (FMO) controls. Positive gates were set based on the negative unstained sample. Data were analyzed using FlowJo software.

Colony Formation Assay

For colony formation assays, SCLC cells were dissociated by gentle pipetting. Live TPCs were sorted using a 100 µm nozzle on a BD FACSAria II. TPCs were sorted individually into 96 well plates filled with regular media (200 µl/well) containing DMSO or doxycycline (DOX) (1 µg/mL). 50 µL fresh media with or without DOX was added to the wells every 10 days. Three weeks later, colony numbers were assessed using CellTiter-Glo 2.0 reagent (Promega #G9242) and luminescence was measured (POLARstar Optima plate reader) according to the manufacturer's protocol. CellTiter-Glo® readings greater than 300 Relative Luminescence Units (RLUs) in colony forming assays were considered colonies.

Extreme Limiting Dilution Analysis

In vivo

The viable cell number was assessed by replicate cell counts on a hemocytometer using Trypan Blue exclusion. Viable cell number was used to derive a titration of cell numbers for implantation. Cells were diluted in 50 µl media (RPMI +10% FBS) and 50 µl Cultrex PathClear BME (Trevigen # 3632-005-02). Six eight-week-old NCG mice were injected subcutaneously into the shoulders and flanks. 3 replicates for each dilution were used. Mice were provided drinking water with 5% sucrose, or sucrose plus doxycycline (2 mg/kg). Injection sites were palpated biweekly to monitor for tumor growth and all mice were sacrificed when any one tumor reached 1 cm. Tumors that formed were scored as 1 and the absence of tumor formation was scored as 0 for the extreme limiting dilution analysis (ELDA). Tumors that formed were analyzed for expression of KSR2 by western blot and tumors in the doxycycline treated group which did not maintain knockdown were scored as 0. ELDA software was used to estimate TPC frequency for control and doxycycline treated groups.

In vitro

Cells were seeded in 96-well plates (Fisher #12556008) at decreasing cell concentrations (1000 cells/well – 1 cell/well) at half log intervals (1000, 300, 100, 30, 10, 3, 1), 12 wells per condition. Cells were cultured for 14 days, and wells with spheroids >100 µM were scored as spheroid positive. TPC frequency and significance between groups was calculated by ELDA software.

Bibliography

1. Gazdar, A.F., Bunn, P.A. & Minna, J.D. Small-cell lung cancer: what we know, what we need to know and the path forward. *Nat Rev Cancer* **17**, 765 (2017).
2. Petty, W.J. & Paz-Ares, L. Emerging Strategies for the Treatment of Small Cell Lung Cancer: A Review. *JAMA Oncol* **9**, 419-429 (2023).
3. Gay, C.M. *et al.* Patterns of transcription factor programs and immune pathway activation define four major subtypes of SCLC with distinct therapeutic vulnerabilities. *Cancer Cell* **39**, 346-360 e347 (2021).
4. Byers, L.A. *et al.* Proteomic profiling identifies dysregulated pathways in small cell lung cancer and novel therapeutic targets including PARP1. *Cancer Discov* **2**, 798-811 (2012).
5. Gazdar, A.F., Bunn, P.A. & Minna, J.D. Small-cell lung cancer: what we know, what we need to know and the path forward. *Nat Rev Cancer* **17**, 725-737 (2017).
6. Gazdar, A.F., Carney, D.N., Nau, M.M. & Minna, J.D. Characterization of variant subclasses of cell lines derived from small cell lung cancer having distinctive biochemical, morphological, and growth properties. *Cancer Res* **45**, 2924-2930 (1985).
7. George, J. *et al.* Comprehensive genomic profiles of small cell lung cancer. *Nature* **524**, 47-53 (2015).
8. McFadden, David G. *et al.* Genetic and Clonal Dissection of Murine Small Cell Lung Carcinoma Progression by Genome Sequencing. *Cell* **156**, 1298-1311 (2014).
9. Sutherland, Kate D. *et al.* Cell of Origin of Small Cell Lung Cancer: Inactivation of Trp53 and Rb1 in Distinct Cell Types of Adult Mouse Lung. *Cancer Cell* **19**, 754-764 (2011).
10. Yang, D. *et al.* Intertumoral Heterogeneity in SCLC Is Influenced by the Cell Type of Origin. *Cancer Discov* **8**, 1316-1331 (2018).
11. Park, K.S. *et al.* Characterization of the cell of origin for small cell lung cancer. *Cell Cycle* **10**, 2806-2815 (2011).
12. Schaffer, B.E. *et al.* Loss of p130 accelerates tumor development in a mouse model for human small-cell lung carcinoma. *Cancer Res* **70**, 3877-3883 (2010).
13. Rudin, C.M. *et al.* Molecular subtypes of small cell lung cancer: a synthesis of human and mouse model data. *Nat Rev Cancer* **19**, 289-297 (2019).
14. Borromeo, M.D. *et al.* ASCL1 and NEUROD1 Reveal Heterogeneity in Pulmonary Neuroendocrine Tumors and Regulate Distinct Genetic Programs. *Cell Rep* **16**, 1259-1272 (2016).
15. Gazdar, A.F. & Minna, J.D. Cell lines as an investigational tool for the study of biology of small cell lung cancer. *Eur J Cancer Clin Oncol* **22**, 909-911 (1986).
16. Carney, D.N. *et al.* Establishment and identification of small cell lung cancer cell lines having classic and variant features. *Cancer Res* **45**, 2913-2923 (1985).
17. Rudin, C.M. *et al.* Comprehensive genomic analysis identifies SOX2 as a frequently amplified gene in small-cell lung cancer. *Nat Genet* **44**, 1111-1116 (2012).
18. George, J. *et al.* Evolutionary trajectories of small cell lung cancer under therapy. *Nature* **627**, 880-889 (2024).
19. Ouadah, Y. *et al.* Rare Pulmonary Neuroendocrine Cells Are Stem Cells Regulated by Rb, p53, and Notch. *Cell* **179**, 403-416 e423 (2019).
20. Jahchan, N.S. *et al.* Identification and Targeting of Long-Term Tumor-Propagating Cells in Small Cell Lung Cancer. *Cell Rep* **16**, 644-656 (2016).
21. Stewart, C.A. & Byers, L.A. Altering the Course of Small Cell Lung Cancer: Targeting Cancer Stem Cells via LSD1 Inhibition. *Cancer Cell* **28**, 4-6 (2015).
22. Sancho, P., Barneda, D. & Heeschen, C. Hallmarks of cancer stem cell metabolism. *Br J Cancer* **114**, 1305-1312 (2016).
23. Ailles, L.E. & Weissman, I.L. Cancer stem cells in solid tumors. *Curr Opin Biotechnol* **18**, 460-466 (2007).
24. Borah, A., Raveendran, S., Rochani, A., Maekawa, T. & Kumar, D.S. Targeting self-renewal pathways in cancer stem cells: clinical implications for cancer therapy. *Oncogenesis* **4**, e177 (2015).

25. Codony-Servat, J., Verlicchi, A. & Rosell, R. Cancer stem cells in small cell lung cancer. *Transl Lung Cancer Res* **5**, 16-25 (2016).

26. Hu, J. *et al.* Cancer stem cell self-renewal as a therapeutic target in human oral cancer. *Oncogene* (2019).

27. Yang, F., Xu, J., Tang, L. & Guan, X. Breast cancer stem cell: the roles and therapeutic implications. *Cell Mol Life Sci* **74**, 951-966 (2017).

28. Geradts, J. *et al.* Human lung cancer cell lines exhibit resistance to retinoic acid treatment. *Cell Growth Differ* **4**, 799-809 (1993).

29. Fernandez, M.R., Henry, M.D. & Lewis, R.E. Kinase suppressor of Ras 2 (KSR2) regulates tumor cell transformation via AMPK. *Mol Cell Biol* **32**, 3718-3731 (2012).

30. Costanzo-Garvey, D.L. *et al.* KSR2 is an essential regulator of AMP kinase, energy expenditure, and insulin sensitivity. *Cell Metab* **10**, 366-378 (2009).

31. Guo, L. *et al.* Kinase Suppressor of Ras 2 (KSR2) expression in the brain regulates energy balance and glucose homeostasis. *Mol Metab* **6**, 194-205 (2017).

32. Dougherty, M.K. *et al.* KSR2 is a calcineurin substrate that promotes ERK cascade activation in response to calcium signals. *Mol Cell* **34**, 652-662 (2009).

33. Borges, M. *et al.* An achaete-scute homologue essential for neuroendocrine differentiation in the lung. *Nature* **386**, 852-855 (1997).

34. Kim, E.J., Ables, J.L., Dickel, L.K., Eisch, A.J. & Johnson, J.E. Ascl1 (Mash1) defines cells with long-term neurogenic potential in subgranular and subventricular zones in adult mouse brain. *PLoS One* **6**, e18472 (2011).

35. Augustyn, A. *et al.* ASCL1 is a lineage oncogene providing therapeutic targets for high-grade neuroendocrine lung cancers. *Proc Natl Acad Sci U S A* **111**, 14788-14793 (2014).

36. Sutherland, K.D. *et al.* Cell of origin of small cell lung cancer: inactivation of Trp53 and Rb1 in distinct cell types of adult mouse lung. *Cancer Cell* **19**, 754-764 (2011).

37. Ireland, A.S. *et al.* MYC Drives Temporal Evolution of Small Cell Lung Cancer Subtypes by Reprogramming Neuroendocrine Fate. *Cancer Cell* **38**, 60-78 e12 (2020).

38. Olsen, R.R. *et al.* ASCL1 represses a SOX9(+) neural crest stem-like state in small cell lung cancer. *Genes Dev* **35**, 847-869 (2021).

39. Johnson, S., Chen, H. & Lo, P.K. In vitro Tumorsphere Formation Assays. *Bio Protoc* **3** (2013).

40. Gou, S. *et al.* Establishment of clonal colony-forming assay for propagation of pancreatic cancer cells with stem cell properties. *Pancreas* **34**, 429-435 (2007).

41. Rajendran, V. & Jain, M.V. In Vitro Tumorigenic Assay: Colony Forming Assay for Cancer Stem Cells. *Methods Mol Biol* **1692**, 89-95 (2018).

42. Chang, K.J., Yin, J.Z., Huang, H., Li, B. & Yang, M.H. Arsenic trioxide inhibits the growth of cancer stem cells derived from small cell lung cancer by downregulating stem cell-maintenance factors and inducing apoptosis via the Hedgehog signaling blockade. *Transl Lung Cancer Res* **9**, 1379-1396 (2020).

43. Hu, Y. & Smyth, G.K. ELDA: extreme limiting dilution analysis for comparing depleted and enriched populations in stem cell and other assays. *J Immunol Methods* **347**, 70-78 (2009).

44. Guo, L. *et al.* Cell non-autonomous regulation of hepatic IGF-1 and neonatal growth by Kinase Suppressor of Ras 2 (KSR2). *Sci Rep* **6**, 32093 (2016).

45. Cristea, S. & Sage, J. Is the Canonical RAF/MEK/ERK Signaling Pathway a Therapeutic Target in SCLC? *J Thorac Oncol* **11**, 1233-1241 (2016).

46. Barbosa, R., Acevedo, L.A. & Marmorstein, R. The MEK/ERK Network as a Therapeutic Target in Human Cancer. *Mol Cancer Res* **19**, 361-374 (2021).

47. Tsurutani, J., West, K.A., Sayyah, J., Gills, J.J. & Dennis, P.A. Inhibition of the phosphatidylinositol 3-kinase/Akt/mammalian target of rapamycin pathway but not the MEK/ERK pathway attenuates laminin-mediated small cell lung cancer cellular survival and resistance to imatinib mesylate or chemotherapy. *Cancer Res* **65**, 8423-8432 (2005).

48.   Wang, Z. *et al.* ARHGEF19 regulates MAPK/ERK signaling and promotes the progression of small cell lung cancer. *Biochem Biophys Res Commun* **533**, 792-799 (2020).

49.   Liu, Y. *et al.* Resistance to BH3 mimetic S1 in SCLC cells that up-regulate and phosphorylate Bcl-2 through ERK1/2. *Br J Pharmacol* **169**, 1612-1623 (2013).

50.   Burger, M. *et al.* Functional expression of CXCR4 (CD184) on small-cell lung cancer cells mediates migration, integrin activation, and adhesion to stromal cells. *Oncogene* **22**, 8093-8101 (2003).

51.   Ravi, R.K. *et al.* Raf-1 causes growth suppression and alteration of neuroendocrine markers in DMS53 human small-cell lung cancer cells. *Am J Respir Cell Mol Biol* **20**, 543-549 (1999).

52.   Ravi, R.K. *et al.* Activated Raf-1 causes growth arrest in human small cell lung cancer cells. *J Clin Invest* **101**, 153-159 (1998).

53.   Brennan, D.F. *et al.* A Raf-induced allosteric transition of KSR stimulates phosphorylation of MEK. *Nature* **472**, 366-369 (2011).

54.   Dhawan, N.S., Scopton, A.P. & Dar, A.C. Small molecule stabilization of the KSR inactive state antagonizes oncogenic Ras signalling. *Nature* **537**, 112-116 (2016).

55.   Rajakulendran, T., Sahmi, M., Lefrancois, M., Sicheri, F. & Therrien, M. A dimerization-dependent mechanism drives RAF catalytic activation. *Nature* **461**, 542-545 (2009).

56.   Bragelmann, J. *et al.* Family matters: How MYC family oncogenes impact small cell lung cancer. *Cell Cycle* **16**, 1489-1498 (2017).

57.   Mollaoglu, G. *et al.* MYC Drives Progression of Small Cell Lung Cancer to a Variant Neuroendocrine Subtype with Vulnerability to Aurora Kinase Inhibition. *Cancer Cell* **31**, 270-285 (2017).

58.   Sos, M.L. *et al.* A framework for identification of actionable cancer genome dependencies in small cell lung cancer. *Proc Natl Acad Sci U S A* **109**, 17034-17039 (2012).

59.   Johnson, B.E. *et al.* myc family oncogene amplification in tumor cell lines established from small cell lung cancer patients and its relationship to clinical status and course. *J Clin Invest* **79**, 1629-1634 (1987).

60.   Brennan, J. *et al.* myc family DNA amplification in 107 tumors and tumor cell lines from patients with small cell lung cancer treated with different combination chemotherapy regimens. *Cancer Res* **51**, 1708-1712 (1991).

61.   Duplaquet, L. *et al.* KDM6A epigenetically regulates subtype plasticity in small cell lung cancer. *Nat Cell Biol* **25**, 1346-1358 (2023).

62.   Sutherland, K.D., Ireland, A.S. & Oliver, T.G. Killing SCLC: insights into how to target a shapeshifting tumor. *Genes Dev* **36**, 241-258 (2022).

63.   Poirier, J.T. *et al.* New Approaches to SCLC Therapy: From the Laboratory to the Clinic. *J Thorac Oncol* **15**, 520-540 (2020).

64.   Meuwissen, R. *et al.* Induction of small cell lung cancer by somatic inactivation of both Trp53 and Rb1 in a conditional mouse model. *Cancer Cell* **4**, 181-189 (2003).

65.   Lim, J.S. *et al.* Intratumoural heterogeneity generated by Notch signalling promotes small-cell lung cancer. *Nature* **545**, 360-364 (2017).

Figure Legends

Fig. 1. A) *KSR2* mRNA in brain tissue and lung epithelial tissue (GTEx Portal). B) GFP[+] cells isolated from the lungs of mice expressing GFP from the promoter for the neuroendocrine (NE)-specific *Cgrp.* mRNA expression of *Ksr2* (*left*), neuroendocrine markers (*Chga*, *Syp*) (*middle*), and lung epithelial marker surfactant protein C (*Spc*) (*right*). C) *Ksr2* expression from bulk RNA-sequencing of indicated mouse tumors. D) *KSR2* expression in human ASCL-1 subtype SCLC tumors (*left*) and tumors with elevated *MYC* expression (*right*) from RNA-seq analysis of human SCLC tumors. **, p<0.01 E) *KSR2* RNA-seq (DepMap) in SCLC segregated by subtype. *p=0.03; **p<0.004; ****p<0.0001. F) KSR2 and KSR1 expression, with ACTIN loading control, in western blots of ASCL1-subtype SCLC cell lines and non-transformed human bronchial epithelial cells (HBEC). G) KSR2 and KSR1 expression, with ACTIN loading control, in western blots of ASCL1 (A) or NEUROD1 (N) SCLC lines and non-transformed human bronchial epithelial cells (HBEC).

Fig. 2. A) Dox-inducible RNAi of *Ksr2* in KP1 and KP3 cells from an SCLC mouse model. Dox-inducible RNAi of *KSR2* in human SCLC cell line H209. NTC, non-targeting control. FACS of H209 (B) and KP1 (C) cells for CD24^{high} CD44^{low} (left) and EPCAM^{high} (right) with (sh5) and without (CTRL) dox-inducible RNAi of *KSR2*. D) Percent TPCs detectable by FACS in control or *KSR2*-targeted (sh5) H209 and KP1 cells. Clonogenicity of Dox-treated KP3 (E) and KP1 (F) SCLC TPCs with and without the indicated shRNAs targeting *Ksr2*. Colony viability was measured by CellTiter-Glo. ****, p<0.0001, n=3.

Fig. 3. Quantification (left) and raw data (right) of in vivo extreme limiting dilution analysis of KP1 cells without and with Dox-induced RNAi of *Ksr2*. **, p=0.00698.

Fig. 4. A) A *Ksr2* cDNA resistant to the sh5 shRNA (KSR2r) or the empty expression vector were introduced into KP1 cells expressing the Dox-inducible sh5 shRNA. B) Clonogenicity of KP1 SCLC TPC colonies within wildtype (WT), dox-induced RNAi of *Ksr2* (sh5) and dox-induced RNAi of *Ksr2* + expression of shRNA-resistant *Ksr2* (KSR2r). Colony viability was measured by CellTiter-Glo. ****, p<0.0001. C-D) *In vitro* ELDA of KP1 sh5 or KSR2r KP1 cells with dox-induced RNAi of KSR2 was used to determine the estimated frequency of TPCs. ****, p<0.0001.

Fig. 5. A) *In vitro* ELDA was used to determine the proportion of TPCs in H209 cells with dox-induced RNAi of *KSR2* (sh5) cells. B) Western blot of H209 cells following CRISPR/Cas9 expression with non-targeting (NTC1) control gRNA or sgRNA3 (KSR2KO) targeting *KSR2*. C) *In vitro* ELDA of H209 non-targeting control (NTC) cells or *KSR2*-targeted (KSR2KO) cells. ****, p<0.0001 (one representative image is shown of an experiment completed in triplicate).

Fig. 6. Western blot showing phosphorylation of ERK in KP1 sh5 (A), KSR2r (B), and FIF570r (C) cells with and without dox-induced KSR2 depletion with no treatment (NT), serum starvation, serum shock, or ionomycin treatment. D) Immunoprecipitation of FLAG epitope-tagged wildtype KSR2 (KSR2r) and ERK-binding mutant KSR2 (FIF570r). E) Clonogenicity of KP1 SCLC TPC colonies with dox-induced RNAi of *Ksr2* (sh5) and dox-induced RNAi of *Ksr2* + expression of hairpin resistant *Ksr2* (KSR2r) or hairpin resistant ERK-binding mutant *Ksr2* (FIF570r). F) Quantification of colonies greater than 300 RLU per 96/well plate, sample size is as indicated, ****, p<0.0001.

Fig. 1

Fig. 2

KP1

Table 1. ELDA KP1 cells ± KSR2			
Cells	Mice	Tumor	Group
10	3	0	control
50	3	1	control
100	3	2	control
200	3	3	control
500	3	1	control
10	3	0	dox
50	3	0	dox
100	3	1	dox
200	3	0	dox
500	3	0	dox

Group	TPC Estimate
Dox	2530
control	255
Chisq	P value
7.28	0.00698

Fig. 3

A

Group	TPC Estimate
Control	1/5.49
KSR2 KD	1/49.86

Chisq	P value
56.4	5.84e-14

B

C

Group	TPC Estimate
NTC	1/10.7
KSR2KO	1/131.8

Chisq	P value
40.2	2.29e-10

Fig. 5

Fig. 6

Key Resources

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Cell line (*Mus musculus*)	Small-cell lung carcinoma	Obtained from Julien Sage (Stanford University)	KP1	
Cell line (*Mus musculus*)	Small-cell lung carcinoma	Obtained from Julien Sage (Stanford University)	KP3	
Cell line (*Homo sapiens*)	Small-cell lung carcinoma	Obtained from John Minna (UT Southwestern)	NCI-H209	
Cell line (*Homo sapiens*)	Small-cell lung carcinoma	Obtained from John Minna (UT Southwestern)	NCI-H1963	
Cell line (*Homo sapiens*)	Small-cell lung carcinoma	Obtained from John Minna (UT Southwestern)	NCI-H711	
Cell line (*Homo sapiens*)	Small-cell lung carcinoma	Obtained from John Minna (UT Southwestern)	NCI-H196	
Cell line (*Homo sapiens*)	Small-cell lung carcinoma	Obtained from John Minna (UT Southwestern)	NCI-H1836	
Cell line (*Homo sapiens*)	Small-cell lung carcinoma	Obtained from John Minna (UT Southwestern)	NCI-H82	
Cell line (*Homo sapiens*)	Small-cell lung carcinoma	Obtained from John Minna (UT Southwestern)	NCI-H146	
Cell line (*Homo sapiens*)	Small-cell lung carcinoma	Obtained from John Minna (UT Southwestern)	NCI-H524	
Cell line (*Homo sapiens*)	Small-cell lung carcinoma	Obtained from John Minna (UT Southwestern)	HCC33	
Cell line (*Homo sapiens*)	Small-cell lung carcinoma	Obtained from John Minna (UT Southwestern)	HBEC	
Transfected construct	piSMART hEF1a/TurboGFP-KSR2 shRNA 5	Dharmacon	Cat# V3SH11252-225640245	GGAGCAAATCCCACGAGTT
Transfected construct	piSMART hEF1a/TurboGFP-KSR2 shRNA 6	Dharmacon	Cat# V3SH11252-227968197	GCACATCAGTCAGACGCTC
Transfected construct	piSMART hEF1a/TurboGFP-KSR2 shRNA 7	Dharmacon	Cat# V3SH11252-228320208	CCATCAAGCACAGGTTTTC
Transfected construct	lentiCRISPR v2-KSR2 sgRNA3	Addgene	Plasmid #52961	KSR2_3: 5' CCGAATCGTCGATGTGCGCA 3'

Transfected construct	lentiCRISPR v2- NTC	Addgene	Plasmid #52961	NTC: 5' CCATATCGGGGCGAGACA TG 3'
Recombinant DNA reagent	KSR2r	(adapted from Fernandez et al. 2012)[29]	MSCV KSR2 IRES YFP	QuikChange Lightning Mutagenesis: Forward 5' CGCTGCACAGGAGTAAGT CCCAGTAATTCCAGCTCGG G 3' Reverse 5'CCCGAGCGTGAATTCAT GGGACTTACTCCTGTGCAG CG 3'
Recombinant DNA reagent	FIF570r	(adapted from Fernandez et al. 2012)[29]	MSCV KSR2 FIF570 IRES YFP	QuikChange Lightning Mutagenesis: Forward 5' CGCTGCACAGGAGTAAGT CCCAGTAATTCCAGCTCGG G 3' Reverse 5'CCCGAGCGTGAATTCAT GGGACTTACTCCTGTGCAG CG 3'
Sequence based reagent	Ksr2 (PCR primer)	(Guo et al. 2017)[31]		Forward primer 5'-TGGATGTCCGAAAGGAAGT C-3' Reverse primer 5'-CTTCTCCACGGTCTCACAC A-3'
Sequence based reagent	Crgp (PCR primer)	(Lim et al. 2017)[65]		
Sequence based reagent	Chga (PCR primer)	(Lim et al. 2017)[65]		
Sequence based reagent	Syp (PCR primer)	(Lim et al. 2017)[65]		
Sequence based reagent	Spc (PCR primer)	(Lim et al. 2017)[65]		
Antibody	Anti-B ACTIN, mouse monoclonal	Santa Cruz	Cat# 47778	WB 1:1000
Antibody	Anti-KSR2 (MO8), mouse monoclonal	Abnova	Cat# H00283455-M08	WB 1:1000
Antibody	Phospho-p44/42 MAPK (Erk1/2) (Thr202/Tyr204) (E10), mouse monoclonal	Cell Signaling	Cat# 9106	WB 1:1000
Antibody	p44/42 MAPK (Erk1/2), rabbit polyclonal	Cell Signaling	Cat# 9102	WB 1:1000
Antibody	Anti-Flag (M2), mouse monoclonal	Sigma	Cat# F3165	WB 1:1000

Antibody	Anti-HDAC2 (Y461), rabbit monoclonal	Abcam	Cat# ab32117	WB: 1:1200
Antibody	APC anti-mouse CD326 (EpCAM), mouse monclonal	Biolegend	Cat# 118213	FACS: 1:100
Antibody	PE anti-mouse CD24, mouse monoclonal	Biolegend	Cat# 101807	FACS: 1:400
Antibody	PE Cyanine7 anti-mouse/human CD44, mouse monoclonal	Biolegend	Cat# 103029	FACS: 1:300
Nuclear Stain	DAPI (4',6-Diamidino-2-Phenylindole, Dilactate)	Biolegend	Cat# 422801	FACS: 3 uM

Acknowledgements
The authors thank the UNMC Flow Cytometry Research Facility for help isolating SCLC TPCs.

We declare no conflicts of interest.

Funding Information
This work was supported by the Cancer and Smoking Disease Research Program (NE DHHS – LB 506), R21CA256638, NIH GM121316, and Lung Cancer Research Program (LCRP) LC210123 to R.E.L., NIH U24CA213274 to T.G.O., and P50 CA070907, U24 CA213274 award to J.D.M.

The funders had no role in the study design, data collection and interpretation, or the decision to submit the work for publication.