Title: Strength of stabilizing selection on the amino-acid sequence is associated with the amount of non-additive variance in gene expression

Authors: Margarita Takou[1,2], Daniel J. Balick[3], Kim A. Steige[1,4], Hannes Dittberner[1], Ulrike Göbel[1,5], Holger Schielzeth[6], Juliette de Meaux[1]

Addresses:

1 Institute of Botany, University of Cologne, Cologne, Germany

2 Department of Biology, Pennsylvania State University, PA, United States of America

3 Department of Biomedical Informatics, Harvard Medical School, Boston, Massachusetts, United States of America

4 State Plant Breeding Institute, University of Hohenheim, Stuttgart, Germany

5 Bioinformatics Core Facility, CECAD Research Center, Cologne, Germany

6. Institute of Ecology and Evolution, Friedrich Schiller University Jena, Jena, Germany

Abstract:

The distinction between the additive and non-additive components of genetic trait variance has long been known to be crucial for predicting the potential of a population to respond to selection. Yet we know surprisingly little about the genomic and evolutionary factors that influence non-additive genetic variance in natural populations. Here, we use a quantitative genetic breeding design to delineate the additive and non-additive components of expression variance of 17,657 gene transcripts in the outcrossing plant *Arabidopsis lyrata*. We find that most of the genetic variance in gene expression is from non-additive variance. Transcripts showing the highest levels of non-additive variance tend to be longer, clustered in larger groups of co-expressed genes, and enriched in functions related to cellular differentiation or to the epigenetic reprogramming of gene expression. Furthermore, we find that genes with the most non-additive variation not only display a markedly lower rate of synonymous variation, they are also exposed to stronger stabilizing selective forces than genes with high additive variance. To the best of our knowledge, this study is the first to show that stabilizing selection on amino-acid sequences is associated with a decrease in a population's potential for adaptation of gene expression levels.

Main

Genetic variation is a prerequisite for populations to respond to selection[1]. However, in each
generation, alleles within the same or different loci are reshuffled, so that the phenotypic effects
resulting from inter-allelic interactions cannot be transmitted from parents to offspring.
Consequently, only a fraction of the genetic variation in a trait exposed to selection can sustain a
response to selection. To account for this fraction, Fisher divided genetic variance into additive
variance, to refer to components that are directly transmitted to the next generation, and non-
additive variance, to describe the remaining variance[2]. The main non-additive component is
termed dominance variance (V_D) in quantitative genetics studies, and although strictly defined
for intra-locus interactions, the use of the term typically also includes inter-locus allelic
interactions that give rise to epistatic genetic variance. The amount of non-additive variance
inferred in quantitative genetics studies is a statistical component of variation. It cannot be
simply derived from the recessive/dominant relationships between alleles that can be dissected
genetically[3]. Because most classic genome-wide association studies (GWAS) assume co-
dominant allelic relationships, the amount of trait heritability in a population that has the
potential to respond to selection remains unknown[4–9] (but see[3,10]). This amount also cannot be
derived from dissecting interactions between alleles at different loci because a fraction of
functional epistasis will contribute to additive variance[3,11,12]. Therefore, quantitative genetics
approaches are required to understand what controls how much genetic variance present in a
natural population is non-additive and thus not available for selection.

Several factors are expected to influence the amount of non-additive genetic variance. First, the
amount depends on the genetic architecture of the phenotype. As the number of loci controlling a
phenotype increases, so will the likelihood that new allelic variants interact with other variants of
the same or other loci to define a phenotypic value that differs from the sum of effects of the
individual variants[13]. Second, the amount of non-additive variance is assumed to depend on the
population's recent history. Indeed, non-additive variance is expected to be enriched in traits that
have been depleted of their additive genetic variance by recent natural selection[14,15]. Indeed,
additive variance will decrease after the advantageous alleles have fixed in the population[7]. By
contrast, when genetic drift is the primary force affecting allele frequencies, the proportion of

additive genetic variance is expected to increase[7,16,17]. The amount of non-additive variance thus

depends on the molecular architecture of a trait as well as on the selective and demographic

history of the population.

Despite the crucial importance of non-additive variance for predicting the adaptive potential of

single traits, few studies have investigated the evolutionary factors underlying the amount of

non-additive variance present in natural populations, and even less is known about the molecular

and genomic features that promote the emergence of non-additive genetic variation[16].

Transcriptome studies generate powerful opportunities to study the evolutionary properties of a

large collection of quantitative molecular traits[18]. In addition, such approaches can deliver crucial

information on the extent to which genetic variation in gene expression can contribute to

adaptation[19–21]. Most transcriptome studies assume that genetic variance is additive[18,21].

However, when quantified over appropriate pedigrees, transcriptome variation provides ideal

data for exploring the genomic and evolutionary properties shaping the amount of non-additive

variance in natural populations. Indeed, transcript level variance in full- and half-sib families of

threespine sticklebacks showed that non-additive variance is pervasive in the transcriptome[22].

Conducting such studies in plants can be particularly powerful because environmental and

maternal variance can be easily distinguished, preventing them from having a confounding

effect. Here, we quantify the additive and non-additive components of genetic variance in gene

expression during the exponential growth phase of the outcrossing plant species *Arabidopsis*

lyrata. We show that additive and non-additive variance in gene expression varies with the level

of negative selection detected at the protein level. Whereas additive variance tends to be

associated with genes experiencing relaxed selection, we find that non-additive variance is

pervasive, dependent on transcript length, and promoted by the complexity of gene regulation.

Results

We took advantage of a set of 10 full- and half-sib families of the outcrossing plant species

Arabidopsis lyrata ssp. *petraea*, which were obtained by crossing individuals from two locally

adapted populations [23–25] (Supplementary Table S1). Given the amount of shared variation

between the populations and their similar frequency distribution, this set of inter-population

hybrids is expected to show levels of non-additive variance similar to those within populations

(Supplementary Figure S1)[26]. Full- and half-sib relationships allowed us to partition the total

phenotypic variance (V_P) of gene expression into additive variance (V_A), non-additive (V_D),

maternal (V_M), and residual (V_R) components of variance in the expression for each transcript in

the genome (Figure 1a, Supplementary Table 1). We refer to V_D here as the non-additive

variance because the statistical variance due intra- and inter-locus, i.e. dominance and epistatic

variance, cannot be disentangled. In 67.7% of the 17,657 genes analyzed, more than 50% of total

expression variance (V_P) was explained by the significant contribution of the two genetic

components of variance, V_D (mean = 0.375) and V_A (mean = 0.215). The maternal variance (V_M)

was close to zero for most transcripts (mean = 0.088). Finally, the population of origin of the

mother plant did not affect gene expression variance (Bonferroni corrected threshold of p =

2.486e-06).

Delineating overall genetic variance showed that V_D contributed the most (25.4% of 17,657

genes had more than 50% of gene expression variance V_P due to V_D), in contrast to the additive

genetic component V_A (6.3% of these genes had more than 50% of their variance in expression,

Figure 1b-c). Hence, gene expression variance V_P is due largely to genetics, but only a limited

fraction of that genetic variance is inherited additively. Although amounts of non-additive

genetic variance are rarely quantified in natural populations, the amount we detected is not

unexpected. Similar fractions of non-additive variance have been reported in phenotypic traits[16],

and non-additive variance was as high if not higher than additive variance for most transcripts

expressed in threespine stickleback[22]. In a control dataset generated by permuting family labels,

V_R explained more than half of the V_P for more than 73% of transcripts (mean = 0.717), and V_D

was not significant (mean = 0.135), confirming that the levels of genetic variance we report are

not random.

Most genetic variance cannot be explained by large effect regulatory variants

We used our data to investigate the genomic and evolutionary features that are associated with

the relative importance of additive and non-additive variance in gene expression. We first

explored the regulatory architecture of gene expression, using a hierarchical clustering approach.

We used a k-clustering method and tested a number of clusters k ranging from 23 to 300 clusters

of co-expressed genes (Supplementary Figure S2). Setting the number of clusters, k, to 200,

cluster size ranged between 5 and 2,829 genes (median=39.5 genes per cluster). Thus, and

perhaps unsurprisingly given what is known about the regulatory networks controlling gene
expression[18,27], our dataset revealed that many genes form co-expressed clusters of variable size.

To evaluate the number and effect sizes of regulatory variants explaining the expression and co-
expression of the genes, we used allelic segregation across 131 individuals of the transcriptome
sample; this allowed us to implement a GWAS approach, controlling for the family structure of
the mapping population (see Methods). In total, we identified 156 Single Nucleotide
Polymorphisms (SNPs) associated with the expression level of only 174 of the 17,656 expressed
gene transcripts and these 156 SNPs never co-localized with the transcript (Supplementary
Figure S2). Thus, for 99% of expressed transcripts, no main causal genetic variant could be
detected with the approach. Importantly, most of the 156 SNPs (76.4%) were associated with the
expression of a single gene and, thus, do not explain the formation of any of the clusters. This
result suggests that most of the genetic variation in expression has a polygenic basis. A major
genetic incompatibility, which can lead to phenotypic aberration and broad regulatory alterations
in inter-population crosses, is unlikely to explain the data[28–31].

Differences in the strength of negative selection are associated with the mode of regulatory
inheritance

Both drift and natural selection are predicted to shape the amount of non-additive variance in a
population [11,12]. To determine whether the non-additive variance in gene expression is linked to
the evolutionary forces acting on the encoded protein, we examined rates of amino-acid
evolution in the transcripts that were expressed in our dataset. We partitioned expressed
transcripts in three non-overlapping sets based on their relative excess of V_D, V_A, or V_R (Figure
2a). As a control, we included a fourth set of 3,500 genes sampled at random (Fig. 2a). We then
estimated the rate of amino acid divergence that evolved between *A. lyrata* and *A. thaliana*
across these four groups of genes. The four sets of transcripts differed significantly in their rate
of amino-acid divergence. Genes with high V_R had the lowest rate of amino-acid divergence,
whereas genes with highest V_A had the fastest rate (Figure 2b-c). Interestingly, genes with the
largest V_D displayed a rate of amino-acid divergence that was intermediate between those shown
by genes with large V_A and genes with large V_R.

The rate of amino-acid divergence is expected to increase if positive selection is acting on the
encoded protein, but it can also arise if the evolutionary constraints on the amino-acid sequence

are weak. In order to differentiate these alternatives, the segregation of molecular variation

within species can be used to infer the fitness effect of new mutations and quantify the strength

of the evolutionary constraints imposed on the amino-acid sequence. We took advantage of a

previous study, which determined the demographic history of the same two parental populations

of *A. lyrata*[25,32]. We inferred the distribution of fitness effects (DFE) of new non-synonymous

variants in these gene sets, following a method described in[25] that incorporates the specific

demographic parameters of each population (Supplementary Figure S3). Both the inferred

demographic model and the DFE fitted the site-frequency spectra (SFS) of synonymous and non-

synonymous variants well (Supplementary Figure S3a), indicating that the model provides a

good representation of the evolutionary and selective history of these populations.

The results showed that the inferred strength of evolutionary constraint (Figure 2d;

Supplementary Table S2) co-varies with differences in the mode of genetic variance in transcript

expression level. Compared to the other sets of genes, the set of genes with low genetic variance

in expression (i.e., high residual variance, V_R) showed a significantly lower proportion of nearly

neutral non-synonymous mutations ($0 < N_{anc}s < 1$, $p = 0.01$), as well as a significantly higher

proportion of mutations under strong purifying selection ($N_{anc}s > 10$, $p = 0.01$). This observation

indicates that genes evolving under strong constraint at the amino-acid level also tended to show

low levels of genetic variation at the transcript expression level. Stabilizing selection has indeed

been shown to be associated with decreased rates of expression variation or divergence [18,19,33–39].

Our analysis further revealed that selective constraints on genes with large V_A (threshold 0.5) are

comparatively lower than on other genes (Figure 2c, Supplementary Figure S3b-c). The DFE of

the set of genes with large V_A in transcript expression indicated that, compared to other gene

sets, these genes hosted a significantly larger fraction of new non-synonymous mutations that are

nearly neutral ($0 < N_{anc}s < 1$, $p= 0.03$). At the same time, a significantly smaller fraction of new

mutations is under strong purifying selection in genes with large V_A ($N_{anc}s > 10$, $p = 0.01$).

Furthermore, genes with high V_A were less likely to map to genomic regions harboring footprints

of selective sweeps. Although regions with signatures of selective sweeps extended over 27% of

the genome, only 17% of the high V_A genes mapped to these regions (pairwise t-test, $p= 2.115e-$

12), further confirming that these genes are unlikely to contain variants that have been under

directional selection. Therefore, both the high rate of amino-acid divergence observed in these

genes and their inferred DFE indicate that genes with additive variance in gene expression tend
to be exposed to lower levels of purifying selection than are the other genes.

The evolutionary constraints detected on the amino-acid sequence of genes with high levels of
V_D (threshold 0.5) did not differ significantly from those detected in the control group of random
genes (Fig. 2c, Supplementary Figure S3d). However, the rate of nucleotide divergence at
synonymous sites, Ks, was markedly lower in genes with high V_D, than in all other gene groups
(Supplementary Figure S3). A lower K_S is expected if the mutation rate is lower or the
background selection is stronger[40].

Gene architecture associates with the level of non-additive variance in transcripts

In order to better understand what modulates the relative fractions of non-additive and additive
variance, we compiled a total of 200 descriptors of genetic variation at each gene, including the
number of known transcription-factor binding sites in the upstream non-coding region, the length
of the genes, and various descriptors of nucleotide diversity (see Methods). We trained a random
forest model to identify the genomic architecture or population genetic parameters that best
explain variation in the prevalence of V_A or V_D in gene expression. We first performed the
analysis using the 200 descriptors to predict variation in V_A, but the model's ability to classify
high V_A genes was poor ($R^2 = 0.009$ and mean square error of 0.021).

For variation in V_D, however, the model was accurate at classifying genes with high V_D ($R^2 =$
0.18 and mean square error of 0.026), highlighting that the most important factors for classifying
genes according to their level of V_D are descriptors of gene structure (Figure 3a; Supplementary
Table S3). Specifically, the number of exons per gene (importance: 53.021; $p = 0.0164$), total
gene length (importance: 26.92; $p = 0.0164$), and transcript length (importance: 14.27; $p =$
0.0164) were the three factors that contributed most to predict levels of non-additive variance.
The importance of exon number persisted even when gene length was corrected for. Transcript
length was indeed significantly correlated with V_D (Spearman coefficient $\rho = 0.258$, $p = 1.22e-$
266, Supplementary Table S4; Figure 3b). We used a linear model to confirm that the effect of
transcript length on V_D ($p < 2.2e-16$) remained significant even after correcting for median gene
expression ($p = 0.0176$) and V_G ($p < 2.2e-16$). The model did not suggest a significant interaction
between transcript length and median gene expression ($p = 0.224$). The number of exons per
gene correlated significantly with the level of non-additive variance ($\rho = 0.358$, $p < 2.2e-16$;

Figure 3c). The random forest analysis further revealed that the density of SNPs within a gene
was a predictor of high V_D. Strikingly, all significant descriptors tend to increase the mutational
target size of the transcript. We thus conclude that non-additive variance is more likely to be
observed as the mutational target size increases.

Interestingly, the analysis further revealed that the total number of transcription-factor binding
sites explained V_D well (p = 0.0163; importance 4.34; Figure 3d). When we corrected for gene
length, the total number of transcription-factor binding sites becomes the most important
predictor in the random forest analysis (Importance 28.72; p =0.0163), which explains the level
of V_D in the dataset. In particular, the transcription factor binding sites associated with high V_D
were targets of a DOF zinc finger protein expressed in young rosettes (DOF 5.3; p = 0.0163),
that was recently shown to determine stem cell division necessary for plant thickening[41]
(AT5G60200 TMO6; p = 0.01639). It is likely that this regulator was active in the tissue samples
we collected, because we sampled during the exponential phase of rosette growth.

We further quantified the distribution of genes with high V_D across Gene Ontology (GO)
categories of expressed genes. We found that genes with high V_D were enriched for GO
categories reflecting cell differentiation. The top enriched GO terms for the highest V_D estimates
were "chromatin silencing" (p = 4.5e-16) and "cytokinesis by cell plate formation" (p = 2e-11),
"DNA methylation" (p = 4.3e-11; Supplementary Table S5). Genes with high V_A were also not
randomly distributed among molecular functions, but, in contrast to V_D genes, their functional
enrichment was less striking than for high V_D genes and not related to cell differentiation.
Among the top GO categories most enriched among genes with high V_A values were several
functions linked to central metabolism such as "glucose catabolic process" (p = 8.3-05) and
"glycolytic process" (p = 1.6e-05), but we also detected enrichments among genes involved in
the "response to cadmium ion" (p = 0.00015) and "proteasome core complex assembly" (p =
0.00016; Supplementary Table S6).

Non-additive variance increases with the size of the co-regulatory network

Our results indicate that two processes may increase the opportunities for the inter-allelic
interactions that favor the emergence of non-additive variance: the mutational target size of a
transcript (its length and number of exon-intron boundaries) and the complexity of the regulatory
network. The observations that transcription-factor binding sites as well as the DOF zinc finger

protein TMO6 contribute significantly to the amount of non-additive variance in gene expression

confirm this observation. More generally, we found that the median V_D in a cluster of co-

regulated genes increased with the number of genes contained in a cluster ($\rho = 0.49$, $p = 1.253\text{e-}$

13); additionally, the correlation between the gene number per cluster and median V_A was

negative ($\rho = -0.277$, $p = 7.101\text{e-}05$). The median V_D per cluster was positively correlated with

the median transcript length ($\rho = 0.201$, $p = 0.0042$) and median gene exon number ($\rho = 0.439$, p

$= 7.752\text{e-}11$). Accordingly, our data suggest that inter-allelic and inter-genic interactions both

increase the amount of non-additive variance in gene expression.

Interestingly, there was a core set of 2,829 genes that always clustered together in the k-

clustering analysis. Genes in this cluster had median V_D and V_A values of 0.501 and 0.141,

respectively. GO enrichment analysis highlights 294 GO terms (significance threshold $p =$

0.0077; Supplementary Table S7). The top processes are related to developmental functions,

such as "gravitropism" ($p = 6.4\text{e-}28$) and "trichome morphogenesis" ($p = 8.9\text{e-}15$).

Genes with high V_A in gene expression show higher levels of nucleotide variation

The random forest analysis indicated that none of the population genetic parameters of the

original populations, such as F_{ST} ($\rho = 0.0128$, $p = 0.164$), had a significant impact on the level of

non-additive variance (Figure 3a; Supplemental Table S3). In contrast, we observed that the

amount of V_A was inversely correlated with genetic divergence between the two parental

populations. The correlation was detected both at the individual gene level and at the level of

gene clusters (Supplemental Table S4). The median V_A of each cluster was negatively correlated

with the median value of relative population divergence F_{ST} ($\rho = -0.201$, $p = 0.0041$) and

positively correlated with the absolute measure of population divergence d_{xy} ($\rho = 0.320$, $p =$

2.59e-06) among parental populations. The median V_A was also correlated with the median π

values in both populations (SP: $\rho = 0.34$, $p = 8.643\text{e-}07$, PL: $\rho = 0.33$, $p = 1.627\text{e-}07$), and high

V_A genes were among the genes that displayed the highest divergence at both synonymous and

non synonymous sites (Supplementary Figure 3b-c). Taken together, our observations indicate

that V_A evolves preferentially in transcripts displaying a higher mutation rate[42].

We used a generalized linear mixed model to test whether values of V_A, V_D, or V_G were

predicted by F_{ST} and/or local signals of positive selection while controlling for gene identity. We

also tested in the same model whether effects of selection signals differed among the three gene

sets. The variance distributions are all significantly different from each other ($p < 2.2e\text{-}16$), and
the presence of a signature of a selective sweep in the vicinity of the expressed gene has a
significant impact on the components of genetic variance of this gene ($p = 1.319e\text{-}4$). We further
focused on the 156 SNPs displaying a significant association with the expression level of a gene.
Because these SNPs did not co-locate with the expressed transcript, we can assume that they act
as trans-regulatory variants. Overall, we find that when such a trans-regulatory SNP mapped to a
genomic region carrying a signature of a selective sweep, the transcript it is associated with
displayed significantly higher V_A and lower V_D than when this trans-regulator was located
outside of a sweep area. Thus, recent selective sweeps seem to associate preferentially with
higher additive variance in gene expression and to lower levels of non-additive variance. This
observation accounts only for the 1% of transcripts for which a significant association could be
detected, it thus does not contradict the observation reported above that genes with high V_A tend
to show relaxed selective constraint. Clearly additive variance can sometimes be targeted by
positive selection.

Discussion

Delineating the additive and non-additive components of genetic variance is crucial for
understanding the adaptive potential of a trait, because only additive genetic variance can fuel a
response to selection. Plants are particularly well suited for the dissection of genetic variance
because both genetic crosses and environmental variance can be completely controlled; such
control is necessary for a reliable estimation of the non-additive components of variance[13,43]. Our
analysis of non-additive variance in 17,657 gene transcripts provides a unique opportunity to
determine the factors that promote the emergence of variance in natural populations. Our data
show that a substantial fraction of regulatory variation is not available to respond to selection
despite being genetically determined. This work further documents three striking features of non-
additive variance in gene expression: i) there is a strong relationship between mode of genetic
inheritance of expression levels and the rate of amino acid evolution, ii) there are identifiable
genomic features that favor non-additive variation in gene expression, and iii) non-additive
variance is particularly elevated among genes related to the epigenetic regulation of gene
expression.

First, we report a strong association between two independent sources of genetic variation: the
composition of genetic variance in expression and the strength of selective constraint on amino-
acid evolution. Gene regulation, which is central to the formation of phenotypes and variation in
gene expression, is evidently subject to natural selection[21,44–47], and non-additive variance is
expected to be higher in traits exposed to natural selection. A meta-analysis of non-additive
variance in 338 traits in various organisms suggested that traits linked to fitness often have larger
levels of non-additive variance in natural populations[16]. In maize, non-additive variance in 18
traits selected during domestication was also reported to be greater in the domesticated
population than in its wild ancestor[43]. The scale of our study provides a compelling
documentation of the relationship between the strength of selection and the amount of non-
additive variance. Genes whose genetic variance at the transcript level is predominantly non-
additive were significantly more constrained than genes showing the largest fraction of additive
variance. The most strongly constrained genes, in turn, were found among the genes that were
almost completely depleted of any genetic variance in transcript level, additive and non-additive
variance alike. Although it has long been known that levels of genetic variation are low in genes
with constrained expression[18,19,33–37,48], this study is, as far as we know, the first report that non-
additive transcript variance increases when the strength of stabilizing selection increases but
ultimately decreases in genes exposed to the strongest stabilizing selective forces. These
differences are revealed by a skew in the frequency spectrum of non-synonymous mutations as
well as in the number of amino-acid substitutions that ultimately fix in a population. The method
used to estimate the DFE assumes that allelic effects are purely additive. If instead variants had
recessive effects on fitness, the method would underestimate the magnitude of their effect and
thus should lead to a pattern opposite to the one we observe, where variants in high V_A genes
experienced weaker selection than variants in high V_D genes. Our results are therefore robust to
hypothesized fitness effects of alleles. Although non-additive genetic variance emerges as a
result of intra- and inter-locus interactions, no such direct relationship exists between allelic
effects and the amount of non-additive variance[3]. This is because non-additive variance is a
property of the population resulting from the frequency at which genetic variants segregate as
well as the molecular interactions between them[13,49]. It thus would not make much sense to
model the DFE assuming distinct values of the allelic dominance h for the different groups of
genes. We note however, that our data reveals an important additional feature: the rate of

synonymous divergence and polymorphism is significantly lower in genes with high non-additive variance compared to all other genes, suggesting that background selection or the mutation rate may also play a role on the composition of genetic variance for expression.

Second, our data indicate that non-additive variance is promoted by transcript length, suggesting that the mutational target size of the expressed locus favors the emergence of such variance. Our study of transcript-level variance further indicates that non-additive variance increases with the size of the network of co-regulated genes, which confirms the observation that proteins that are more central tend to be more constrained at the amino-acid level[15,50–52]. Nevertheless, this work also shows that what promotes non-additive phenotypic variance is distinct from the molecular interaction between alleles. Indeed, additive molecular effects on fitness (e.g. co-dominant alleles) were inferred, in the same species, to be more frequent in highly connected genes, a result opposite to our findings[53].

Finally, we report that non-additive variance is particularly predominant among genes influencing the epigenetic remodeling of chromatin in differentiating cells. Interestingly, we also associated non-additive variance in expression with the presence of sites bound by a regulator of cell differentiation associated with stem thickening during growth[54]. Although epigenetic variation has been often hypothesized to facilitate adaptation[55], our results indicate that given the genetic architecture of epigenetic regulation, it is comparatively less likely to produce the additive variance needed to fuel adaptation in *A. lyrata*. Differentiating between additive and non-additive sources of variance appears to be crucial to understand the role of gene expression variation for past and future adaptation.

Acknowledgments

We would like to thank Kirsten Bell for help with the laboratory work, the Cologne Genome Center for the transcriptome sequences and Emily Wheeler, Harwich, MA, for editorial help. Furthermore, we would like to thank F. He, S. Sunyaev, O. Savolainen, T. Hämälä, and M. Stetter for feedback, and F. Hospital for inspiring discussions ahead of this project. This research was funded by ERC Grant No. 648617 Adaptoscope.

Data availability

The whole-genome and transcriptome sequences are available via ENA. The whole-genome
sequences of the parental lines are under the number PRJEB33206, and the transcriptome
sequences are under the number PRJEB49153. The gene counts per individual and the sql
database compiling descriptors of genomic variation will be deposited in dryad upon acceptance
of the manuscript. All the original scripts are in git repository
https://github.com/mtakou/Quantitative-analysis-of-gene-expression.

References

1. Savolainen, O., Lascoux, M. & Merilä, J. Ecological genomics of local adaptation. *Nat Rev Genet* **14**, 807–820 (2013).

2. Fisher, R. A. The correlation between relatives on the supposition of mendelian inheritance. *Trans. R. Soc. Edinb* **53**, 399–433 (1918).

3. Huang, W. & Mackay, T. F. C. The Genetic Architecture of Quantitative Traits Cannot Be Inferred from Variance Component Analysis. *PLOS Genetics* **12**, e1006421 (2016).

4. Manolio, T. A. *et al.* Finding the missing heritability of complex diseases. *Nature* **461**, 747–753 (2009).

5. Boyle, E. A., Li, Y. I. & Pritchard, J. K. An Expanded View of Complex Traits: From Polygenic to Omnigenic. *Cell* **169**, 1177–1186 (2017).

6. Zhu, Z. *et al.* Dominance genetic variation contributes little to the missing heritability for human complex traits. *Am J Hum Genet* **96**, 377–385 (2015).

7. Hill, W. G., Goddard, M. E. & Visscher, P. M. Data and Theory Point to Mainly Additive Genetic Variance for Complex Traits. *PLoS Genet* **4**, e1000008 (2008).

8. Crow, J. F. On epistasis: why it is unimportant in polygenic directional selection. *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**, 1241–1244 (2010).

9. Visscher, P. M., Hill, W. G. & Wray, N. R. Heritability in the genomics era — concepts and misconceptions. *Nature Reviews Genetics* **9**, 255–266 (2008).

10. Mäki-Tanila, A. & Hill, W. G. Influence of Gene Interaction on Complex Trait Variation with

Multilocus Models. *Genetics* **198**, 355–367 (2014).

11. Willis, J. H. & Orr, H. A. Increased Heritable Variation Following Population Bottlenecks: The Role of

Dominance. *Evolution* **47**, 949–957 (1993).

12. Wang, J., Caballero, A., Keightley, P. D. & Hill, W. G. Bottleneck effect on genetic variance. A

theoretical investigation of the role of dominance. *Genetics* **150**, 435–447 (1998).

13. Lynch, M. & Walsh, B. *Genetics and Analysis of Quantitative Traits*. (Sinauer Associates, Sunderland,

Massachusetts, 1998).

14. Mousseau, T. A. & Roff, D. A. Natural selection and the heritability of fitness components. *Heredity*

59, 181–197 (1987).

15. Orr, H. A. Fitness and its role in evolutionary genetics. *Nat Rev Genet* **10**, 531–539 (2009).

16. Crnokrak, P. & Roff, D. A. Dominance variance: associations with selection and fitness. *Heredity* **75**,

530–540 (1995).

17. Waldmann, P. Additive and non-additive genetic architecture of two different-sized populations of

Scabiosa canescens. *Heredity* **86**, 648–657 (2001).

18. Ayroles, J. F. *et al.* Systems genetics of complex traits in Drosophila melanogaster. *Nat Genet* **41**,

299–307 (2009).

19. Romero, I. G., Ruvinsky, I. & Gilad, Y. Comparative studies of gene expression and the evolution of

gene regulation. *Nat Rev Genet* **13**, 505–516 (2012).

20. Oleksiak, M. F., Churchill, G. A. & Crawford, D. L. Variation in gene expression within and among

natural populations. *Nature Genetics* **32**, 261–266 (2002).

21. Groen, S. C. *et al.* The strength and pattern of natural selection on gene expression in rice. *Nature*

1–5 (2020) doi:10.1038/s41586-020-1997-2.

22. Leder, E. H. *et al.* The Evolution and Adaptive Potential of Transcriptional Variation in Sticklebacks—

Signatures of Selection and Widespread Heritability. *Mol Biol Evol* **32**, 674–689 (2015).

23. Mattila, T. M., Tyrmi, J., Pyhäjärvi, T. & Savolainen, O. Genome-Wide Analysis of Colonization

History and Concomitant Selection in Arabidopsis lyrata. *Mol Biol Evol* **34**, 2665–2677 (2017).

24. Hämälä, T., Mattila, T. M. & Savolainen, O. Local adaptation and ecological differentiation under

selection, migration, and drift in Arabidopsis lyrata. *Evolution* **72**, 1373–1386 (2018).

25. Takou, M. *et al.* Maintenance of Adaptive Dynamics and No Detectable Load in a Range-Edge

Outcrossing Plant Population. *Molecular Biology and Evolution* **38**, 1820–1836 (2021).

26. Reif, J. C., Gumpert, F.-M., Fischer, S. & Melchinger, A. E. Impact of Interpopulation Divergence on

Additive and Dominance Variance in Hybrid Populations. *Genetics* **176**, 1931–1934 (2007).

27. Bullard, J. H., Mostovoy, Y., Dudoit, S. & Brem, R. B. Polygenic and directional regulatory evolution

across pathways in Saccharomyces. *PNAS* **107**, 5058–5063 (2010).

28. Lafon-Placette, C. & Köhler, C. Epigenetic mechanisms of postzygotic reproductive isolation in

plants. *Current Opinion in Plant Biology* **23**, 39–44 (2015).

29. Todesco, M. *et al.* Hybridization and extinction. *Evolutionary Applications* **9**, 892–908 (2016).

30. Alcázar, R. *et al.* Natural variation at Strubbelig Receptor Kinase 3 drives immune-triggered

incompatibilities between Arabidopsis thaliana accessions. *Nat Genet* **42**, 1135–1139 (2010).

31. Bomblies, K. *et al.* Local-Scale Patterns of Genetic Variability, Outcrossing, and Spatial Structure in

Natural Stands of Arabidopsis thaliana. *PLOS Genetics* **6**, e1000890 (2010).

32. He, F. *et al.* Cis-regulatory evolution spotlights species differences in the adaptive potential of gene

expression plasticity. *Nat Commun* **12**, 3376 (2021).

33. Fay, J. C. & Wittkopp, P. J. Evaluating the role of natural selection in the evolution of gene

regulation. *Heredity* **100**, 191–199 (2008).

34. Steige, K. A., Reimegård, J., Koenig, D., Scofield, D. G. & Slotte, T. Cis-Regulatory Changes Associated

with a Recent Mating System Shift and Floral Adaptation in Capsella. *Mol Biol Evol* **32**, 2501–2514

(2015).

35. Hämälä, T. *et al.* Gene Expression Modularity Reveals Footprints of Polygenic Adaptation in

Theobroma cacao. *Mol Biol Evol* **37**, 110–123 (2020).

36. Whitehead, A. & Crawford, D. L. Neutral and adaptive variation in gene expression. *PNAS* **103**,

5425–5430 (2006).

37. Hodgins-Davis, A., Rice, D. P. & Townsend, J. P. Gene Expression Evolves under a House-of-Cards

Model of Stabilizing Selection. *Mol Biol Evol* **32**, 2130–2140 (2015).

38. Metzger, B. P. H., Wittkopp, P. J. & Coolon, J. D. Evolutionary Dynamics of Regulatory Changes

Underlying Gene Expression Divergence among Saccharomyces Species. *Genome Biol Evol* **9**, 843–

854 (2017).

39. Khaitovich, P., Enard, W., Lachmann, M. & Pääbo, S. Evolution of primate gene expression. *Nat Rev*

Genet **7**, 693–702 (2006).

40. Schrider, D. R. Background Selection Does Not Mimic the Patterns of Genetic Diversity Produced by

Selective Sweeps. *Genetics* **216**, 499–519 (2020).

41. Miyashima, S. *et al.* Mobile PEAR transcription factors integrate positional cues to prime cambial

growth. *Nature* **565**, 490–494 (2019).

42. Cruickshank, T. E. & Hahn, M. W. Reanalysis suggests that genomic islands of speciation are due to

reduced diversity, not reduced gene flow. *Mol Ecol* **23**, 3133–3157 (2014).

43. Yang, C. J. *et al.* The genetic architecture of teosinte catalyzed and constrained maize domestication.

Proc Natl Acad Sci USA **116**, 5643–5652 (2019).

44. Kremling, K. A. G. *et al.* Dysregulation of expression correlates with rare-allele burden and fitness

loss in maize. *Nature* **555**, 520–523 (2018).

45. Studer, A., Zhao, Q., Ross-Ibarra, J. & Doebley, J. Identification of a functional transposon insertion

in the maize domestication gene tb1. *Nat Genet* **43**, 1160–1163 (2011).

46. Clark, R. M., Wagler, T. N., Quijada, P. & Doebley, J. A distant upstream enhancer at the maize

domestication gene tb1 has pleiotropic effects on plant and inflorescent architecture. *Nat Genet* **38**,

594–597 (2006).

47. Jacob, F. & Monod, J. Genetic regulatory mechanisms in the synthesis of proteins. *Journal of*

Molecular Biology **3**, 318–356 (1961).

48. Bedford, T. & Hartl, D. L. Optimization of gene expression by natural selection. *PNAS* **106**, 1133–

1138 (2009).

49. Falconer, D. S. & Mackay, T. F. C. *Introduction to Quantitative Genetics*. (Prentice Hall, 1996).

50. Hahn, M. W. & Kern, A. D. Comparative Genomics of Centrality and Essentiality in Three Eukaryotic

Protein-Interaction Networks. *Mol Biol Evol* **22**, 803–806 (2005).

51. Josephs, E. B., Wright, S. I., Stinchcombe, J. R. & Schoen, D. J. The Relationship between Selection,

Network Connectivity, and Regulatory Variation within a Population of Capsella grandiflora. *Genome*

Biol Evol **9**, 1099–1109 (2017).

52. Masalia, R. R., Bewick, A. J. & Burke, J. M. Connectivity in gene coexpression networks negatively

correlates with rates of molecular evolution in flowering plants. *PLOS ONE* **12**, e0182289 (2017).

53. Huber, C. D., Durvasula, A., Hancock, A. M. & Lohmueller, K. E. Gene expression drives the evolution

of dominance. *Nature Communications* **9**, 2750 (2018).

54. Lister, R. *et al.* A Transcriptomic and Proteomic Characterization of the Arabidopsis Mitochondrial

Protein Import Apparatus and Its Response to Mitochondrial Dysfunction. *Plant Physiol* **134**, 777–

789 (2004).

55. Schmid, M. W. *et al.* Contribution of epigenetic variation to adaptation in Arabidopsis. *Nat Commun*

9, 4446 (2018).

Figure 1: Phenotypic variance of gene expression is mostly comprised of non-additive genetic variance. **a.** We used 2 populations of *A. lyrata*, performed a set of inter-population crosses, and generated a set of 131 individuals. The two populations separated from each other approximately 61 kyears ago, with the SP populations being more isolated and more severely bottlenecked than the PL population (redrawn from [25]). In total, 5 individuals sampled in Plech (PL) were reciprocally crossed with 6 individuals from Spiterstulen (SP). The crosses were done in such a way that plants were reciprocally crossed with 2 individuals from the other population in order to produce full- and half-sib families. **b.** The total phenotypic variance of each one of the 17,656 genes included in the analysis were partitioned into non-additive, additive, maternal, and residual components of variance. The genes were ordered by increasing values of the fraction of total phenotypic variance in transcript level that was attributed to genetic variance (V_G). The dashed line marks the 0.5 point where half of the variance is explained. **c.** The distribution of total genetic variance values for each transcript (V_G), as a fraction of the total observed variance in expression level. The dashed blue line shows the median value (0.604) of the distribution; the dashed red line indicates the mean value (0.585).

Figure 2: Constraints on the amino-acid level depend on the predominant component of genetic variance. **a.** We grouped the genes based on the partitioning of total phenotypic variance in transcript level into non-additive, additive, or residual variance. The threshold for each case is 50% of the total phenotypic variance. **b.** We computed the mean and variance of average pi/ps values for each group based on population genomics datasets for PL. **c.** The Ka/Ks ratio for the different gene groups. The confidence intervals and p values for b through d were estimated based on 100 bootstrap iterations. **d.** Inferred distribution of fitness effects of new mutations in the four gene groups. Gray: control group of 3500 random genes; beige: high V_A genes; orange: high V_D genes; purple: high V_R genes; green: intermediate levels of V_G. In all plots, the letters indicate whether the values differ significantly from each other.

Figure 3: Gene architecture controls the level of non-additive variance. **a.** The significance and importance of tested factors in the random forest analysis that explains the level of non-additive variance each gene has. The twenty factors that explain the most variance are shown. The predictive power of the model is $R^2 = 0.18$. **b.** The non-additive variance correlates significantly with the transcript length of each gene (p=1.22e-266, ρ=0.258). **c.** The fraction of non-additive variance correlates significantly with the number of exons that a gene has ($p < 2.2$e-16, ρ=0.385). **d.** The non-additive variance correlates significantly with the total number of transcription-factor binding sites detected in the upstream vicinity of the gene ($p < 2.2$e-16, ρ=0.162).

514