Enhanced Sodium Channel Inactivation by Temperature and FHF2 Deficiency Blocks Heat Nociception

Christopher Marra[1,2], Timothy V. Hartke[3], Matthias Ringkamp[3], Mitchell Goldfarb[1,2,*]

[1] Department of Biological Sciences, Hunter College of City University, NY, NY 10065 USA

[2] Program in Biology, Graduate Center of City University, NY, NY 10016 USA

[3] Department of Neurosurgery, Neurosurgery Pain Research Institute, Johns Hopkins University, Baltimore, MD 21287 USA

* Corresponding author. Address: Department of Biological Sciences, Hunter College, 695 Park Avenue, New York, NY 10065 USA. Tel.: +1 212-772-5295 Email: goldfarb@genectr.hunter.cuny.edu

Abstract

Transient voltage-gated sodium currents are essential for the initiation and conduction of action potentials in neurons and cardiomyocytes. The amplitude and duration of sodium currents are tuned by intracellular fibroblast growth factor homologous factors (FHFs/iFGFs) that associate with the cytoplasmic tails of voltage-gated sodium channels (Na$_v$s), and genetic ablation of *Fhf* genes disturbs neurological and cardiac functions. Amongst reported phenotypes, *Fhf2^{null}* mice undergo lethal hyperthermia-induced cardiac conduction block attributable to the combined effects of FHF2 deficiency and elevated temperature on the cardiac sodium channel (Na$_v$1.5) inactivation rate. *Fhf2^{null}* mice also display a lack of heat nociception, while retaining other somatosensory capabilities. Here we employ electrophysiological and computational methods to show that the heat nociception deficit can be explained by the combined effects of elevated temperature and FHF2 deficiency on the inactivation gating of sodium channels Na$_v$1.7 and Na$_v$1.8 expressed in dorsal root ganglion C-fibers. Hence, neurological and cardiac heat-associated deficits in *Fhf2^{null}* mice derive from shared impacts of FHF deficiency and temperature towards Na$_v$ inactivation gating kinetics in distinct tissues.

Keywords

Heat nociception, DRG, sodium channel, inactivation, FHF2, modeling

1. Introduction

Noxious cutaneous chemical, mechanical, and thermal stimuli open transient receptor potential (TRP) cationic channels at nociceptor afferent terminals [30], with sodium influx triggering the activation of recessed voltage-gated sodium channels ($Na_v1.7$ and $Na_v1.8$) that mediate initiation and orthodromic conduction of action potentials [15]. Loss-of-function mutations in genes encoding $Na_v1.7$ and $Na_v1.8$ are hypoalgesic, reducing or ablating peripheral nociceptor activation by noxious stimuli [1,3,6,12,20], while gain-of-function missense mutations in these channel genes are hyperalgesic [7,10,36].

The differential voltage-dependent gating properties of $Na_v1.7$ and $Na_v1.8$ collaborate to promote action potential initiation. $Na_v1.7$ opens more rapidly and at more negative membrane potential than does $Na_v1.8$, but $Na_v1.7$ also inactivates at more negative voltages than $Na_v1.8$ [2,8]. During a slow rise in voltage driven by stimulus-induced TRP channel currents at afferent terminals, most $Na_v1.7$ channels inactivate from the closed state, but residual $Na_v1.7$ availability provides a "primer" current depolarizing membrane sufficiently to open $Na_v1.8$ channels that further drive action potential initiation [5]. The orthodromic action potential rapidly depolarizes downstream membrane, allowing for a greater contribution of $Na_v1.7$ to spike conduction. Reduced availability of $Na_v1.7$ near afferent terminals may explain why distal afferents harbor greater $Na_v1.8$ density [17].

Sodium channel gating is regulated by fibroblast growth factor homologous factors (FHFs), also termed intracellular fibroblast growth factors (iFGFs), that bind to the carboxyl-terminal domain of Na_vs [13,18]. Most FHF isoforms induce a depolarizing shift in voltage dependence of Na_v inactivation without altering the voltage for activation [9,19,33], thereby increasing sodium current generated by Na_vs. Mutations in FHF genes result in various disorders of excitable tissues [11,14,21,22,24,27,29,31,34,35]. *Fhf2^null* mice are sensitive to ventricular cardiac conduction block when exposed to various stressors, including sodium or calcium channel blockers, uncoupling agents, or hyperthermia [21,22]. FHF2 deficiency

and elevated core body temperature collaborate to reduce availability and speed inactivation of cardiac $Na_v1.5$ leading to failure of action potential conduction through myocardium [21].

Genetic ablation of mouse FHF2 in peripheral neurons also results in loss of cutaneous heat nociception [35]. In this paper, we report that FHF2 modulates $Na_v1.7$ and $Na_v1.8$ inactivation gating in a manner analogous to its effects on cardiac $Na_v1.5$. Empirical and computational studies demonstrate that FHF2 modulation of Na_v inactivation provides a transient sodium current reserve in heated afferent terminals to enable action potential initiation and conduction. Overall, these data show that the limiting of sodium currents by FHF2 deficiency and elevated temperature constitutes a common gating mechanism underlying heat nociception and cardiac conduction deficits in *Fhf2^null* mice.

2. Materials and Methods

2.1 Mouse genetics and genotyping

All work conducted with mice complied with protocol approved by the Hunter College Institutional Animal Care and Use Committee (IACUC). The murine embryonic stem cell clone EPD0339-4-F09 (International Mouse Phenotype Consortium) carrying a knockout first with conditional potential cassette bears loxP sites flanking (floxing) *Fhf2* exon 3 and Frt sites flanking the embedded G418-resistance and beta-galactosidase markers; these cells were used to derive mice carrying a *Fhf2^{targeted}* allele by blastocyst injection and outbreeding [21]. The *Fhf2^{targeted}* allele was converted to an *Fhf2^{null}* allele by injection of Cre recombinase-encoding plasmid into fertilized *Fhf2$^{targeted/+}$* eggs followed by reimplantation [21]. Alternatively, the *Fhf2^{targeted}* allele was converted to the conditional *Fhf2^{flox}* allele by mating with mice that globally express a nuclear-targeted FLP recombinase ROSA26Sor$^{tm2(FLP*)Sor}$ (Jackson Labs Stock #012930) [25] to excise the embedded G418-resistance and beta-galactosidase markers. The *Fhf2^{null}* and *Fhf2^{flox}* alleles were backcrossed for four generations onto the 129S2 strain background. By crossing heterozygous *Fhf2$^{flox/+}$* females with male transgenic Advillin-Cre driver mice (Jackson Labs Stock #032536), which express Cre recombinase exclusively in the peripheral sensory neurons [37,38], *Fhf2$^{flox/Y}$*:Adv-Cre test mice and *Fhf2$^{+/Y}$*:Adv-Cre negative control littermates were generated. *Fhf2^{flox}* mice are publicly accessible through Jackson Labs (Stock #036280).

PCR genotyping was performed using PhireII Hot-Start DNA polymerase (Fisher Sci.). The *Fhf2^{flox}* and *Fhf2^{+}* alleles were distinguishable upon PCR with primers IVS2-For (5'-GCCAGGAGTCTGCTCAACTCT) and IVS3-Rev (5'-GACTTTGGTGGGAGCATCCTGA), yielding products of 533 base pairs versus 388 base pairs, respectively.

2.2 Immunoblot detection of FHF2

Detergent-solubilized brain and DRG tissue lysates were electrophoresed through 4%–20% precast polyacrylamide gradient gel (Invitrogen), transferred to activated PVDF (Immobilion-P, Millipore), probed with rabbit anti-FHF2 polyclonal antibodies

(Schoorlemmer 2002), and detected with peroxidase-conjugated secondary antibodies and enhanced chemiluminescence.

2.3 Footpad Immunofluorescence

Footpads tissue punch samples harvested from adult wild-type ($Fhf2^{+/Y}$) and $Fhf2^{null}$ ($Fhf2^{-/Y}$) mice were submersion fixed in 4% paraformaldehyde for 1 hour at 4°C, treated with cryoprotectant 30% sucrose in phosphate-buffer saline for 24 hours at 4°C, and embedded and frozen in OCT blocks for cryosections. Cryosections 30μm thick were permeabilized with 0.5% Triton X100, preabsorbed with 10% fetal calf serum, and incubated overnight at 4°C with chicken anti-PGP9 antibodies (EnCor Biotechnology) at 0.5 μg mL^{-1} each. After three wash cycles, the tissue was subsequently incubated with 1:200 dilution of flurophore-conjugated anti-chicken IgY at room temperature for 2 hours. Images were captured with a Leica TCS2 confocal microscope and images spanning 2–5 mm through the z-axis were merged.

2.4 Sensory assays

All mice were conditioned to test apparati, environments and modes of restraint for at least two consecutive days prior to data collection. *Tail-flick:* Heat nociception was evaluated using a tail-flick apparatus [Ugo Basile Tail Flick 7360]. Mice were held with tail resting unrestrained on disk heat source, which upon test initiation emitted infrared radiant heat 2 cm from the tip of the tail. The instrument measures the latency to tail withdrawal from the heat source. *Paw pinch:* Temperature independent nociception was assayed using a force-calibrated forceps pressure transducer [Ahlborn - Almemo 2450] to determine exactly how much pressure applied to the hind paw of a mouse elicited a pain induced withdrawal response. Mice were placed in a tubular restraint with their hind limbs exposed. Force was applied to each hind paw and the threshold response was recorded. *Von Frey filament assay:* Basal somatosensory response was measured using a set of sensory evaluator filaments [NC Medical - NC12775-99]. Mice were placed in an elevated mesh-floored cage and allowed to explore and adapt for 20-30 minutes. Stimulation was manually applied to the hind footpads of the mouse starting with the thinnest softest filament and proceeding through filaments of greater thickness, performing five trials for each filament. Paw

withdrawal was scored as a positive response, while bending of the filament without paw withdrawal constituted a negative response.

2.5 Voltage clamp analysis of $Na_v1.7$ sodium currents in transfected HEK293 cells

Human embryonic kidney HEK293 cells (QBiogene) were transiently co-transfected with a 2:1 mixture of $Na_v1.7$-expressing plasmid [7] and a pIRES2-ZsGreen bicistronic plasmid (Clontech) expressing ZsGreen and mouse FHF2B proteins [9]. The same pIRES2-ZsGreen plasmid without FHF2 coding sequence served as control. Cells were trypsinized 3 hours post-transfection, seeded onto gelatinized coverslips, and were used for recording after 48 hr. For sodium current recordings, coverslips were transferred to recording chamber containing carbogen-buffered bath solution (115 mM NaCl, 26 mM $NaHCO_3$, 3 mM KCl, 10 mM glucose, 4 mM $MgCl_2$, 2 mM $CaCl_2$, 0.2 mM $CdCl_2$, 3 mM myoinositol, 2 mM Na pyruvate, 7 mM NaOH-buffered HEPES pH 7.2) at 25 °C and green fluorescent cells were whole-cell patched with pipettes filled with 104 mM CsF, 50 mM tetraethylamine chloride, 10 mM HEPES pH 7.2, 5 mM glucose, 2 mM $MgCl_2$, 10 mM EGTA, 2 mM ATP, 0.2 mM GTP and having 1–2 MΩ resistance. All voltage commands and current recordings were made using a MultiClamp 700 amplifier, digital/analog converter Digidata 1440, and pClamp10 software (Molecular Devices – Axon Instruments). For all recording protocols, voltage-gated current was isolated during data acquisition by P/N subtraction of leak and capacitive currents (N = -6). To ensure adequate voltage clamping during sodium channel activation was adequate cells were subjected to a 19-sweep series of voltage steps from a hold of -120 mV to between -80 mV and -10 mV in 5 mV increments. As criterion for adequate clamp, transient current peaks for all voltage commands were nested within the larger current trace of a following or preceding voltage step command.

Steady-state channel inactivation protocol: To determine the voltage dependence of steady state channel inactivation, a 19-sweep protocol used a -120 mV holding command, a 60 ms variable test voltage step (-120 + 5(n - 1) mV), followed by a 25 mV reporting pulse.

Voltage ramp protocol: As a measure of closed state channel inactivation rate, a 10-sweep protocol used -120 mV hold command followed by depolarization to -30 mV either instantaneously (voltage step) or as a ramp ranging in time from 2ms (= 45 mV ms⁻¹) to 18ms (= 5mV ms⁻¹). After recording at 25 °C, temperature was ramped to 35 °C and later to

40 °C. At elevated temperatures, the cell was first tested as above to ensure maintenance of tight clamp, after which the voltage ramp protocol was conducted. For many cells, voltage ramp protocols could be successfully run at all three temperatures.

2.6 Voltage clamp analysis of TTX-resistant sodium currents in acutely dissociated DRG neurons

DRGs were plucked from spinal columns of 8-12 week-old mice and trimmed free of associated nerves. Neurons were dissociated by treatment with 1% trypsin and 150 µg/ml collagenase in Hank's balanced salt solution at 37°C for 30 minutes followed by washing into Dulbecco's modified Eagle's medium (DMEM) + 10% fetal bovine serum (FBS) + 250 µg/ml DNaseI and trituration with fire-polished Pasteur pipettes. Dissociated cells were washed three times with DMEM + 10% FBS and seeded on to 12 mm diameter coverslips precoated with poly-D-lysine and laminin at a density of 3×10^4 cells/coverslip and cultured for 18-40 hours prior to recording. Recordings were conducted in extracellular solution consisting of 24 mM NaCl, 3mM KCl, 3 mM CsCl, 26 mM NaHCO$_3$, 50 mM Choline Cl, 11 mM Glucose, 25 mM TEA-Cl, 2 mM Na Pyruvate, 3 mM Myoinositol, 10 mM HEPES, 5 mM 4-aminopyridine, 2 mM MgCl$_2$, 1 mM BaCl$_2$, 1 mM CaCl$_2$ and 0.2 mM CdCl$_2$ and 0.5 µM tetrodotoxin (TTX) at pH 7.2 using patch pipettes of 1-1.5 MΩ resistance when filled with 120 mM K gluconate, 4 mM NaCl, 5 mM KOH-buffered HEPES, 5 mM KOH-buffered EDTA, 15 mM glucose, 1 mM MgSO$_4$, 3 mM Mg-ATP, 0.1 mM Na$_2$-GTP at pH 7.2. The voltage dependence of TTX-resistant sodium channel steady state inactivation was assayed in a 19-sweep protocol used a -70 mV holding command, a 60 ms variable test voltage step (-70 + 5(n - 1) mV), followed by a 25 mV reporting pulse.

2.7 Temperature-induced excitation of acutely dissociated DRG nociceptive neurons

Voltage-sensitive dye recordings. DRG neurons grown on coverslips as described above were first treated with a blocking solution consisting of sterile HBSS with 10 mM HEPES + 1% horse serum for 5 minutes at room temperature followed by staining with HBSS/HEPES + 70 mM Di-8-ANEPPS (Biotium) for 15 minutes room temperature and recovery in HBSS/HEPES + 10% horse serum for 5 minutes. Coverslips of stained neurons were transferred to recording chamber containing carbogen-bubbled extracellular solution: 115 mM NaCl, 26 mM NaHCO$_3$, 3 mM KCl, 1.2 mM KH$_2$PO$_4$, 3 mM glucose, 2 mM

myoinositol, 2 mM Na pyruvate, 7 mM HEPES, 1.2 mM $MgSO_4$, 2 mM $CaCl_2$, 0.2 mM $CdCl_2$ at pH 7.2. Using a 40X lens objective, SOLA-SE LED illumination (Lumencor) and a filter cube consisting of a 520±45 nm excitation interference filter, a dichroic mirror with 570 nm central wavelength, and 610nm barrier emission filter, fluorescence images were captured at 2 Hz with a NeuroCCD-SM camera (Redshirt Imaging) while heating the bath from 32°C to 44°C at a rate of approximately 4 degrees C per minute. Action potentials were detected as transient fluorescence changes of 2-3% resting fluorescence. Excitable cells were then imaged by high-resolution bright field illumination in order to assess the neuronal cell diameter.

Current-clamp recordings. Recordings were performed using the same extracellular solution as above and patch pipettes of 2-3 MΩ resistance when filled with 120 mM K gluconate, 4 mM NaCl, 5 mM KOH-buffered HEPES, 5 mM KOH-buffered EDTA, 15 mM glucose, 1 mM $MgSO_4$, 3 mM Mg-ATP, 0.1 mM Na_2-GTP at pH 7.2. Cells approximately 12 microns in diameter were selected for patching based upon prior optical imaging that had determined the size of heat responsive cells. Excitability was first assessed at 32°C by applying 200 millisecond current pulses ranging from 0-400 pA in 25 pA intervals. Excitable cells were then held with 0 pA current injection while heating the chamber to 45°C at approximately 4 degrees C per minute. Voltages were recorded at 10 kHz acquisition rate.

2.8 Temperature-induced modulation of C-fiber compound action potential conduction

Saphenous nerve segments approximately 10 mm in length were used for recording C-fiber compound action potentials (CCAPs) as previously described [17]. Each end of a nerve was drawn into a glass pipette and electrically isolated from the central nerve with petroleum jelly to establish high resistance electrical seals. The central nerve was bathed in continuously replenished carbogen-bubble synthetic interstitial fluid. One nerve end was stimulated with a 100 microsecond current pulse every nine seconds, while a recording electrode placed at the other end of the nerve monitored evoked potentials. Recordings were initiated at 27°C with the stimulus strength adjusted sufficient to yield a maximal evoked response. Thereafter, a heating element in the buffer flow path was used to heat

the bath up to 45°C at a rate of approximately 0.2 degrees Celsius per second, and then cooled back to 27°C at the same rate. Only CCAPs from the declining temperature ramp were used for analysis.

2.9 Computational modeling of wild-type and Fhf2^{null} nociceptor excitation

A computational model of a nociceptive neuron with realistic afferent terminal branching and TTX-sensitive (Na$_v$1.7) and –resistant (Na$_v$1.8) transient voltage-gated sodium conductances constructed on the NEURON simulation platform [4,16] was adapted for our construction of temperature-responsive wild-type and *Fhf2^{null}* nociceptor models. For all gated ion conductance models within the neuron, we applied a rate scaling factor Q10 of N^log$_{10}$[localtemp(°C) – 37], where N ranged from 2.3 to 3.0 and where localtemp is the temperature in a particular segment of the neuron, thereby allowing for localized temperature modulation. The *Fhf2^{null}* nociceptor model was identical to the wild-type model, except for altering the parameters within the TTX-sensitive and –resistant sodium conductances to induce left shifts in V$_{1/2}$ inactivation and accelerated inactivation to an extent consistent with experimental findings. Another modification in the models was to increase Na$_v$1.8 density in the branched terminal afferents, consistent with empirical data [17]. These wild-type and *Fhf2^{null}* nociceptor models will be deposited at the publicly accessible repository ModelDB (https://senselab.med.yale.edu/modeldb/).

To assess axonal conduction of action potentials, a segment of the main axon was stimulated with a 100 microsecond 400 pA current pulse while setting the entire neuron to either 37°C or 43°C. TRP channel activation was simulated as inward current ramps delivered into all termini of the afferent tree. For simulation of mechanosensation, the entire neuron was maintained at 37°C, while for simulation of heat nociception, all segments of the afferent tree within 250 microns of termini were raised to 43°C before injection of current ramps. In order to simulate heat stimulation of an acutely dissociated model neuron, the axon was disconnected from the soma, the density of Na$_v$1.8 was increased to that found at branched termini, and a current ramp was injected into the soma set to 45°C.

3. Results

FHF2 Deficiency in Sensory Neurons Causes Loss of Heat Nociception with Lesser Effects on Other Somatosensory Modalities

Fhf2^{null} mice (*Fhf2$^{-/Y}$* male and *Fhf2$^{-/-}$* female) maintained on the 129S2 background strain are viable and fertile, but are sensitive to cardiac conduction block when core body temperature rises above 41°C [21]. When *Fhf2$^{-/Y}$* and *Fhf2$^{-/-}$* mice were tested for heat nociception by tail flick assay, they were uniformly nonresponsive over 12 second exposure of the tail to the infrared heating beam, whereas wild-type *Fhf2$^{+/Y}$* and *Fhf2$^{+/+}$* mice sensed heat and flicked their tails in 3.2 +/- 0.3 seconds (Fig. 1A). In order to determine whether heat nociception requires FHF2 expression in nociceptive neurons, *Fhf2$^{flox/+}$* mice heterozygous for a loxP-flanked (*flox*) *Fhf2* conditional allele (see Methods) were mated with Advillin-Cre transgenic mice that express Cre recombinase exclusively in peripheral neurons [37,38]. Male *Fhf2$^{flox/Y}$*:*AdvCre* progeny showed complete absence of FHF2 protein in DRG neurons while retaining normal FHF2 levels in the brain (Fig. 1B). When tested in the tail flick assay, *Fhf2$^{flox/Y}$*:*AdvCre* mice were uniformly nonresponsive, while control animals retaining peripheral FHF2 expression (*Fhf2$^{flox/Y}$* and *Fhf2$^{+/Y}$*:*AdvCre*) responded to heat stimulus in 3.0 +/- 0.2 seconds (Fig. 1C).

Fhf2^{null} and wild-type mice were tested for mechanical nociception by the paw pinch assay, which entails slowly increasing the compression force on the footpad until a response is elicited. *Fhf2$^{+/Y}$* mice responded to compression at a force of 199 +/- 25 gm (n = 7), while *Fhf2$^{-/Y}$* mice responded at 252 +/- 30 gm (n = 9; P < 0.21)(Fig. 1D). Therefore, *Fhf2^{null}* mice retain relatively near-normal mechanosensitivity. *Fhf2^{null}* and wild-type mice were also tested for paw sensitivity to touch using graded diameter Von Frey nylon filaments. *Fhf2$^{-/Y}$* mice were moderately but significantly less sensitive to touch compared to *Fhf2$^{+/Y}$* littermates (Fig. 1E). In order to qualitatively assess sensory fiber distal innervation, cryosections of *Fhf2$^{+/Y}$* and *Fhf2$^{-/Y}$* footpads were stained with antibodies for

the neural marker PGP9.5. Figure 1F shows similar projection of afferents through the pad and close to the surface in wild-type and mutant animals.

In summary, *Fhf2^null* mice have multiple somatosensory deficits, with heat nociception completely absent. FHF2 is functionally required in heat sensitive, nociceptive neurons, with mutant afferents showing apparently normal projections. Our findings are consistent with those reported previously for nociception when peripheral neurons were rendered *Fhf2^null* on a different background strain [35].

FHF2 Modulation of Inactivation Gating For Sodium Channels Expressed in Nociceptive Neurons

$Na_v1.7$ is the most abundantly expressed sodium channel in nociceptive neurons [5], and the $Na_v1.7$ cytoplasmic tail has been shown to bind FHF2 [35]. Indeed, the crystal structure of FHF2 in complex with the tail of cardiac $Na_v1.5$ revealed that the channel residues that interface with FHF2 are conserved within all nine mammalian sodium channel alpha subunits [32]. In order to assess potential effects of FHF2 on $Na_v1.7$ inactivation gating, HEK293 cells (from QBiogene) were cotransfected with $Na_v1.7$ expression plasmid [7] together with either an FHF2B/GFP bicistronic expression plasmid or GFP-only plasmid [9]. Fluorescent cells were selected for patching and voltage clamp recording of sodium currents (see Methods). The short B isoform of FHF2 protein was chosen for our analyses based upon immunoblotting data showing that FHF2B predominates in peripheral neurons, whereas higher molecular weight FHF2 species are more abundant in the brain (Fig. 1B) and heart [21].

The voltage dependence of steady-state $Na_v1.7$ inactivation was $V_{1/2}$ = -86.8 +/- 1.0 mV in the absence of FHF2B (n=10) and was right-shifted to $V_{1/2}$ = -72.0 +/- 1.9 mV when FHF2B was present (n=10, *P<0.00001*) (Fig. 2A). By contrast, FHF2B did not significantly alter the voltage dependence of $Na_v1.7$ activation (Fig. 2A). We then examined $Na_v1.7$ inactivation rate from the open state at 25°C, 35°C and 40°C in the presence vs. absence of FHF2B upon depolarization to -20 mV. At 25°C, $Na_v1.7$ inactivated more slowly in the

presence of FHF2B (τ = 0.55 +/- 0.04 msec) vs. its absence (τ = 0.40 +/- 0.02 msec, P<0.006, Fig. 2B,C). $Na_v1.7$ inactivated progressively faster at 35°C and 40°C, but at each temperature, inactivation was significantly slowed by the presence of FHF2B (Fig. 2B,C). Hence, the shortest duration of $Na_v1.7$ current we measured was at 40°C in the absence of FHF2B (τ = 0.13 +/- 0.05 msec, Fig. 2B,C) due to the independent effects of elevated temperature and FHF vacancy.

The rate of closed-state $Na_v1.7$ inactivation was assayed at different temperatures in the presence vs. absence of FHF2B by a voltage ramp protocol, as previously described [21]. The voltage clamp was raised from -110 mV to -20 mV either instantaneously (step) or linearly over times ranging from 2 to 18 msec (ramps), and the $Na_v1.7$ peak sodium conductance for each voltage ramp was expressed as percentage of the peak conductance that had been achieved by direct voltage step. Under each transfection and temperature condition, $Na_v1.7$ peak conductance diminished as voltage ramp speed was reduced (Fig. 2D-F), reflecting increased closed-state inactivation. At any voltage ramp speed, inactivation was more pronounced by both elevated temperature and the absence of FHF2B *(P<0.003)*(Fig. 2D-F). More than 50% of channels underwent closed-state inactivation during a rapid 45 mV/msec ramp at 40°C in the absence of FHF2B (Fig. 2F).

In small-diameter nociceptive neurons, $Na_v1.7$ is the principal TTX-sensitive sodium channel, while $Na_v1.8$ is responsible for the transient TTX-resistent sodium current. Transient TTX-resistant sodium currents in acutely dissociated wild-type and *Fhf2^null* DRG neurons (inferred to be $Na_v1.8$ currents) were analyzed by voltage clamp. Steady-state inactivation occurred with $V_{1/2}$ = -31 mV (n = 7) in wild-type neurons vs. $V_{1/2}$ = -24.5 mV (n = 7, *P<0.025*) in *Fhf2^null* neurons (Fig. 2G), demonstrating that FHF2 modulates inactivation gating of the two principal sodium channel isoforms in nociceptive neurons.

FHF2 Is Not Required For Heat-Induced Excitation of Acutely Dissociated DRG Neurons

FHF2 is required in peripheral neurons for heat nociception and FHF2 deficiency facilitates inactivation of the Na$_v$s expressed in DRGs. Therefore, we first sought to analyze heat-induced excitation of acutely dissociated $Fhf2^{WT}$ and $Fhf2^{null}$ DRG neurons. To determine which cell sizes in heterogeneous DRG population to focus on for patch clamp analysis, we first surveyed fields of adult acutely dissociated $Fhf2^{WT}$ neurons for heat-induced fluorescence spikes following extracellular staining with voltage-sensitive dye Di8-ANEPPS.   In one visual field containing several stained cells (Fig.3A), one of four cells commenced firing action potentials upon heating to 43^{o}C (Fig. 3B). This cell (Fig. 3C) and other optically identified heat excitable neurons had a diameter of approximately 12 microns.

For electrical recordings, $Fhf2^{WT}$ DRG neurons were dissociated, cultured overnight, and ~12 micron diameter cells selected for patching. Patched cells were first analyzed in voltage clamp for inward sodium and outward potassium currents in response to command depolarization, followed by excitability in response to inward current injection. Current-excitable cells were then subjected to temperature ramping from 32^{o}C to 45^{o}C followed by cooling while recording voltage. Figure 3D shows an example of a heat-excitable $Fhf2^{WT}$ neuron which commenced firing at 42^{o}C. Figure 3E shows heat-induced excitation of an $Fhf2^{null}$ neuron. The fraction of $Fhf2^{WT}$ neurons that were excitable by heating (39 out of 96 cells) was statistically indistinghishable from the fraction of heat-excitable $Fhf2^{null}$ neurons (13 out of 31 cells). By a range of criteria, including current stimulus-evoked spike amplitude (Fig 3F), temperature threshold for spike induction (Fig 3G), maximum temperature-evoked spike frequency (Fig 3H), and temperature-evoked spike amplitude (Fig 3I), $Fhf2^{null}$ neurons were as excitable as their wild-type counterparts. The only parameter significantly affected by $Fhf2$ deletion was a small reduction in the amplitude of peak inward current upon voltage-clamp depolarization from -70 mV to 0 mV (Fig 3J). This likely reflects a hyperpolarizing shift in $V_{1/2}$ steady state inactivation of Na$_v$1.7, rendering some of the channels inactivated at -70 mV prior to depolarization.

We suspect that acutely dissociated $Fhf2^{null}$ DRG neurons retain heat-induced excitability due to the transient reorganization and expression of nociceptor- and voltage-gated channels after the severing of axonal processes (see Discussion).

Rapid and Reversible Heat-Induced Conduction Failure in $Fhf2^{null}$ C-Fibers

If altered voltage dependence and kinetics of sodium channel inactivation in $Fhf2^{null}$ nociceptors account for loss of heat nociception, it should be possible to detect a rapidly reversible impairment in $Fhf2^{null}$ sensory action potential generation or conduction upon heating. Towards this end, we next tested the ability of unmyelinated sensory C-type fibers to conduct action potentials at different temperatures. Electrical stimulation of $Fhf2^{WT}$ and $Fhf2^{null}$ saphenous nerves *ex vivo* [17] generated C-fiber compound action potentials (CCAPs) (Fig. 4A,B). Each nerve was then stimulated every nine seconds while being warmed to 45°C and cooled back to 27°C, and CCAPs from the cooling phase of the temperature ramp were analyzed. Elevated temperature increased CCAP conduction velocity and decreased amplitude (Fig. 4C-F), reflecting faster rates of voltage-gated channel activation and inactivation. In $Fhf2^{WT}$ nerves, CCAP amplitudes declined linearly from 27°C to 44°C (Fig.4C,E,F). CCAP amplitudes in $Fhf2^{null}$ nerves showed a similar decrement up to 40°C, but declined precipitously above 40°C (Fig. 4D-F), reflecting conduction block or severe attenuation in at least half of all mutant fibers. Figure panels 4D,E illustrate that $Fhf2^{null}$ CCAP impairment was reversed within 27 seconds (3 stimuli) as nerves were cooled from 44°C to 40°C.

The rapid reversibility of conduction deficit upon nerve cooling is strong evidence that sodium channels in $Fhf2^{null}$ axons at high temperature are not denatured or internalized, but fail to generate sufficient inward current for impulse propagation due to accelerated inactivation. These findings parallel prior demonstration that hyperthermia-induced cardiac conduction block in $Fhf2^{null}$ mice is ameliorated upon cooling [21].

Heat Excitation Deficit in a Computational Model of $Fhf2^{null}$ Nociceptor

In order to assess the requirement of FHF2 in mature heat sensitive nociceptors, we employed computational modeling and simulations. For these studies, the wild-type (*Fhf2^WT^*) nociceptor model was taken from the model cell described by Barkai et al [4]. Their model has a highly branched terminal axon with uniformly distributed leak and voltage-gated channels except for the distal afferent terminals, which lack voltage-gated sodium channels but can generate inward currents mimicking the opening of TRP receptor channels [4]. The compartmental segregation of excitatory currents from voltage-gated channels was based upon recordings showing that action potentials initiate at recessed sites on nociceptive afferents [15]. We simulated FHF2 loss-of-function by modifying the gating of the embedded TTX-sensitive ($Na_v1.7$) and TTX-resistant ($Na_v1.8$) conductances to reflect shifts in inactivation rates and voltage dependence commensurate with our recording data (Figure 2) without altering conductance densities. All channel gating transitions in both the *Fhf2^WT^* and *Fhf2^null^* nociceptor models, including $Na_v1.7$ and $Na_v1.8$, were accelerated with increasing temperature by employing Q10 rate scaling factors.

We first tested whether the *Fhf2^null^* model axon suffers conduction block at elevated temperature, analogous to ex vivo conduction block (Figure 4). Wild-type and *Fhf2^null^* model axons were point-stimulated by current pulse and action potential conduction was monitored over a distance of 5 millimeters. The wild-type model axon conducted an action potential at either 37°C (Fig. 5A) or 43°C (Fig. 5B). In contrast, the *Fhf2^null^* model axon conducted at 37°C (Fig. 5C), but suffered conduction block at 43°C (Fig. 5D).

Simulations were next run by current injection into afferent terminals of nociceptor models at 37°C, which recapitulates the effects of noxious chemical or mechanical stimulation (Fig. 5E). The *Fhf2^WT^* model fired a train of action potentials in response to such stimulation (Fig. 5F). The *Fhf2^null^* nociceptor model also generated action potentials, albeit fewer in number, during current injection (Fig. 5G), consistent with preservation of mechanosensation (Fig. 1C) and chemosensation [35] in *Fhf2^null^* mice. Heat nociception was simulated by injecting current into afferent terminals while the temperature of distal axon segments were elevated to 43°C while leaving the rest of the model neuron at 37°C (Fig.

5H). Under these conditions, the *Fhf2^{WT}* nociceptor model still generated a train of action potentials (Fig. 5I), while the *Fhf2^{null}* model failed to be excited (Fig. 5J).

In an attempt to explain why acutely dissociated *Fhf2^{null}* nociceptor neurons display relatively normal heat-induced excitation, we simulated acute dissociation of nociceptive neurons by severing the entire axons in the nociceptor models and performing simulated somatic current injection at 45°C (Fig. 5K). Under these conditions, both wild-type and *Fhf2^{null}* models generated a train of action potentials during current injection (Fig. 5L,M). These findings emphasize the importance of FHF2 to preserve sodium channel availability at elevated temperature during passive propagation of depolarization from nociceptor axon terminal to trigger zone and during conduction down the axon.

4. Discussion

We have shown here that FHF2 is a powerful inactivation gating modulator for the principle TTX-sensitive and TTX-resistant sodium channel isoforms expressed in nociceptive neurons. FHF2 induces depolarizing shifts in $V_{1/2}$ inactivation for $Na_v1.7$ and $Na_v1.8$, and slows the rate of $Na_v1.7$ inactivation from closed and open states. Elevating temperature acts as an independent accelerant of inactivation, speeding closed and open state $Na_v1.7$ inactivation regardless of FHF2 association. Hence, the transient sodium flux upon membrane depolarization is most compromised at elevated temperature in the absence of FHF2. These gating principles offer a plausible explanation for the specificity of heat nociception deficit in *Fhf2^null* mice, despite the broad expression of FHF2 in all classes of sensory neurons. These principles also provide a rationale for preservation of mechanosensation in *Fhf2^null* mice, which is mediated, at least partially, by the same nociceptors that respond to noxious temperatures [30].

Loss of heat nociception was observed following either global or peripheral neuron-restricted genetic ablation of the *Fhf2* gene, demonstrating the requirement of FHF2 function in heat sensitive nociceptors. We were initially surprised to find that heat-induced excitability of acutely dissociated *Fhf2^null* DRG neurons was unimpaired. This paradox is likely explained by the fact that acutely dissociated neurons lack axonal processes and are electrotonically compact; with heat-inducible TRP channels and voltage-gated channels comingled on the somatic plasma membrane, inward TRP channel currents directly depolarize the somatic membrane bearing voltage-gated sodium channels. By contrast, depolarization of heated nociceptor afferent terminals produces retrograde axial current that must passively depolarize a peri-terminal trigger zone [15] subject to cable property delay, reducing sodium channel availability in the absence of FHF. Furthermore, reduced sodium current duration in the *Fhf2^null* afferent may provide insufficient charge to ensure spike conduction, consistent with our observed failure of unmyelinated *Fhf2^null* sensory axons to sustain C-fiber compound action potentials upon heating. Our computational modeling further supports this hypothesis. While current influx at afferent terminals of the

Fhf2^null model neuron failed to evoke action potentials at 43°C, simulated injection of current into the heated axon-severed soma was able to trigger excitation. It is also possible that axotomy accompanying preparation of acutely dissociated DRG neurons triggers upregulation of channel expression to provide sufficient sodium current in mutant cells at elevated temperature.

While peripheral sensory axons express FHF2, myelinated and unmyelinated axons in the central nervous system show little or no FHF expression [9]. This natural deficiency may explain why spike conduction in brain axons is blocked under hyperthermal conditions [23].

We employed plasmid-transfected HEK293 cells to assay the effects of FHF2B expression on the inactivation of $Na_v1.7$. The voltage dependence of $Na_v1.7$ steady-state inactivation was markedly right-shifted by 15 mV from $V_{1/2}$ of -87 mV in the absence of FHF2B to -72 mV in its presence. As discussed above, strong modulation of Na_v inactivation by FHF2 is central to the heat-associated nociception and cardiac deficits in *Fhf2^null* mice. Our findings are in sharp contrast to Yang et al. (2019), who reported only a minor shift in $Na_v1.7$ $V_{1/2}$ inactivation from -76 mV in HEK cells without FHF cotransfection to -73 mV in presence of FHF2B. These disparate findings reflect the expression of endogenous FHF2 protein in some sublineages of HEK293 cells that result in a significant depolarizing shift in $Na_v1.7$ $V_{1/2}$ inactivation in the absence of FHF transfection. Alternatively, the expression vector employed by Yang and colleagues may have produced a lesser concentration of FHF2 insufficient to saturate binding to $Na_v1.7$. It is possible that the failure by Yang and colleagues to detect a strong inactivation modulatory effect of FHF2 led them to emphasize other mechanistic bases for the *Fhf2^null* heat nociception deficit [35].

In the central nervous system, FHF2 is expressed in both excitatory and inhibitory neurons [24]. *Fhf2^+/-* female mice show enhanced susceptibility to epileptic seizures, with most juvenile mice undergoing seizures when core body temperature was elevated above 42°C [24]. As FHF2 expression is subject to X-linked chromosome inactivation, 50% of all

Fhf2+/- neurons are functionally *Fhf2null*, as confirmed by immunofluorescence (C. Marra, unpublished data). We suggest that core hyperthermia combined with FHF2 deficiency triggers seizures by impairing firing of some inhibitory neurons by the same Na_V inactivation gating principles that dictate heat and FHF2 modulation of cardiac and sensory sodium channels.

5. Conclusions

FHF2 expression in sensory neurons is absolutely essential for heat nociception. FHF2 modulation of $Na_v1.7$ and $Na_v1.8$ inactivation gating allows for sufficient inward sodium currents in response to temperature elevation and opening of TRP channels to enable axonal initiation and propagation of antidromic action potentials. This conclusion is supported by voltage clamp analysis of sodium currents, by C-fiber action potential conduction, and by initiation and conduction of action potentials in a nociceptive neuron computational model as a function of temperature and FHF2 expression status. The heat nociception deficit is mechanistically equivalent to cardiac conduction deficit in *Fhf2^{null}* mice exposed to core hyperthermia.

Conflicts of Interest

The authors have no conflicts of interest regarding the research described in this paper.

Acknowledgments

We wish to thank Sergio Solinas (University of Sassari, Sardinia) for expert assistance in implementing modifications to the nociceptor neuron computational model of Barkai et al [4]. We thank Glenn I. Fishman (NYU Langone Medical Center) for his encouragement and support. This work was supported by R01HL142498 to G.I.F., a CUNY research grant to M.G., and the Blaustein Pain Research Foundation and the Neurosurgery Pain Research Institute at Johns Hopkins (to M.R.).

References

[1] Ahmad S, Dahllund L, Eriksson AB, Hellgren D, Karlsson U, et al. A stop codon mutation in SCN9A causes lack of pain sensation. Hum. Mol. Genet. 207; 16:2114-21.

[2] Akopian AN, Sivilotti L, Wood JN. A tetrodotoxin-resistant voltage-gated sodium channel expressed in sensory neurons. Nature 1996; 379:257-262.

[3] Akopian AN, Souslova V, England S, Okuse K, Ogata N, Ure J, Smiath A, Kerr BJ, McMahon SB, Boyce S, Hill R, Stanfa LC, Dickenson AH, Wood JN. The tetrodotoxin-resistant sodium channel SNS has a specialized function in pain pathways. Nature Neurosci. 1999; 2:541-8.

[4] Barkai O, Butterman R, Katz B, Lev S, Binshtok AS. The input-output relation of primary nociceptive neurons is determined by the morphology of the peripheral nociceptive terminals. J. Neurosci. 2020; 40:9346-63.

[5] Blair NT, Bean BP. Roles of tetrodotoxin (TTX)-sensitive Na+ current, TTX-resistant Na+ current, and Ca+2 current in the action potentials of nociceptive sensory neurons. J. Neurosci. 2002; 22:10277-90.

[6] Cox JJ, Reimann F, Nicholas AK, Thornton G, Roberts E, et al. An SCN9A channelopathy causes congenital inability to experience pain. Nature 206; 444:894-8.

[7] Cummins TR, Dib-Hajj SD, Waxman SG. Electrophysiological properties of mutant Na$_v$1.7 sodium channels in a painful inherited neuropathy. J. Neurosci. 2004; 24:8232-6.

[8] Cummins TR, Howe JR, Waxman SG. Slow closed-state inactivation: a novel mechanism underlying ramp currents in cells expressing the hNE/PN1 sodium channel. J. Neurosci. 1998; 18:9607-19.

[9] Dover K, Solinas S, D'Angelo ED, Goldfarb M. Long-term inactivation particle for voltage-gated sodium channels. J. Physiol. 2010; 588:3695-711.

[10] Faber CG, Lauria G, Merkies IS, Cheng X, Han C, Ahn HS, Persson AK, Hoeijmakers JG, Gerrits MM, Pierro T, Lombardi R, Kapetis D, Dib-Hajj SD, Waxman SG. Gain-of-function Nav1.8 mutations in painful neuropathy. Proc Natl Acad Sci U S A. 2012;109:19444–9.

[11] Fry AE, Marra C, Derrick AV, Pickrell WO, Higgins AT et al. Missense variants in the N-terminal domain of the A isoform of FHF2/FGF13 cause an X-linked developmental and epileptic encephalopathy. Am. J. Hum. Genet. 2021; 108:176-85.

[12] Gingras J, Smith S, Matson DJ, Johnson D, Nye K, Couture L, Feric E, Yin R, Moyer BD, Peterson ML, Rottman JB, Beiler RJ, Malmberg AB, McDonough SI. PLoS One 2014; 9:e105895.

[13] Goetz R, Dover K, Laezza F, Shtraizent N, Huang X, Tchetchik D, Eliseenkova AV, Xu C-F, Neubert TA, Ornitz DM, Goldfarb M, Mohammadi M. Crystal structure of a fibroblast growth factor homologous factor (FHF) defines a conserved surface on FHFs for binding and modulation of voltage-gated sodium channels. J. Biol. Chem. 2009; 284:17883-96.

[14] Goldfarb M, Schoorlemmer J, Williams A, Diwakar S, Wang Q, Huang X, Giza J, Tchetchik D, Kelley K, Vega A, Matthews G, Rossi P, Ornitz DM, D'Angelo E. Fibroblast growth factor homologous factors control neuronal excitability through modulation of voltage-gated sodium channels. Neuron 2007; 55:449-63.

[15] Goldstein RH, Barkai O, Inigo-Portuges A, Katz B, Lev S, Binshtok AM. Location and plasticity of the spike initiation zone in nociceptive terminals in vivo. Neuron 2019; 102:801-12.

[16] Hines ML, Carnevale NT. NEURON: a tool for neuroscientists. Neuroscientist 2001; 7:123-35.

[17] Klein AH, Vyshnevska A, Hartke TV, De Col R, Mankowski JL, Turnquist B, Bosmans F, Reeh PW, Schmelz M, Carr RW, Ringkamp M. Sodium channel Nav1.8 underlies TTX-resistant axonal potential conduction in somatosensory C-fibers of distal cutaneous nerves. J. Neurosci. 2017; 37:5204-14.

[18] Liu C-J, Dib-Hajj SD, Waxman SG. Fibroblast growth factor homologous factor 1B binds to the C terminus of the tetrodotoxin-resistant sodium channel rNav1.9a (NaN). J. Biol. Chem. 2001; 276:18925-33.

[19] Lou J-Y, Laezza F, Gerber BR, Xiao M, Yamada KA, Hartmann H, Craig AM, Nerbonne JM, Ornitz DM. Fibroblast growth factor 14 is an intracellular modulator of voltage-gated sodium channels. J. Physiol. 2005; 569:179-93.

[20] Nassar MA, Stirling LC, Forlani G, Baker MD, Matthews EA, Dickenson AH, Wood JN. Nociceptor-specific gene deletion reveals a major role for Nav1.7 (PN1) in acute and inflammatory pain. Proc. Natl. Acad. Sci. USA 2004; 101:12706-11.

[21] Park DS, Shekhar A, Marra C, Lin X, Vasquez C, Solinas S, Kelley K, Morley G, Goldfarb M, Fishman GI. Fhf2 gene deletion causes temperature-sensitive cardiac conduction failure. Nature Commun. 2016; 7:12966.

[22] Park DS, Shekhar A, Santucci J, Redel-Traub G, Solinas S, Mintz S, et al. Ionic mechanisms of impulse propagation failure in the FHF2-deficient heart. Circ. Res. 2020; 127; 1536-48.

[23] Pekala D, Szkudlarek H, Raastad M. Typical gray matter axons in mammalian brain fail to conduct action potentials faithfully at fever-like temperatures. Physiol. Rep. 2016; 4:e12981.

[24] Puranam RS, He XP, Yao L, Le T, Jang W, Rehder CW, Lewis DV, McNamara JO. Disruption of Fgf13 causes synaptic excitatory-inhibitory imbalance and Genetic Epilepsy and Seizures Plus. J. Neurosci. 2015; 35:8866-81.

[25] Raymond CS, Soriano P. High-efficiency FLP and φ31 site-specific recombination in mammalian cells. PLoS One 2007; 2:e162.

[26] Schoorlemmer J, Goldfarb M. FGF homologous factors and islet brain-2 scaffold protein regulate activation of a stress-activated protein kinase. J. Biol. Chem. 2002; 277:49111-9.

[27] Siekierska A, Isrie M, Liu Y, Scheldeman C, Vanthillo N, Lagae L, de Witte PAM, Van Esch H, Goldfarb M, Buyse G. Gain-of-function FHF1 mutation causes early-onset epileptic encephalopathy with cerebellar atrophy. Neurology 2016; 86:2162-70.

[28] Schoorlemmer J, Goldfarb M. FGF homologous factors and the islet brain-2 scaffold protein regulate activation of a stress-activated protein kinase. J. Biol. Chem. 2002; 277:49111-9.

[29] Sinden DS, Holman CD, Bare CJ, Sun X, Gade AR, Cohen DE, Pitt GS. Knockout of the X-linked *Fgf13* in the hypothalamic paraventricular nucleus impairs sympathetic output to brown fat and causes obesity. FASEB J. 2019; 33:11579-94.

[30] Tominaga T, Caterina MJ, Malmberg AB, Rosen TA, Gilbert H, Skinner K, Raumann BE, Basbaum AI, Julius D. The cloned capsaicin receptor integrates multiple pain-producing stimuli. Neuron 1998; 21:531-43.

[31] Trivisano M, Ferretti A, Bebin E, Huh L, Lesca G, Siekierska A, et al. Defining the phenotype of FHF1 developmental and epileptic encephalopathy. Epilepsia 2020; 61:e71-81.

[32] Wang C, Chung BC, Yan H, Lee S-Y, Pitt GS. Crystal structure of the ternary complex of a NaV C-terminal domain, a fibroblast growth factor homologous factor, and calmodulin. Structure 2012; 20:1167-76.

[33] Wang C, Hennessey JA, Kirkton RD, Wang W, Graham V, Puranam RS, Rosenberg PB, Bursac N, Pitt GS. Fibroblast growth factor homologous factor 13 regulates Na+ channels and conduction velocity in murine hearts. Circ. Res. 2011; 109:775-82.

[34] Wang Q, Bardgett ME, Wong M, Wozniak DF, Lou J, McNeil BD, Chen C, Nardi A, Reid DC, Yamada K, Ornitz DM. Ataxia and paroxysmal dyskinesia in mice lacking axonally transported FGF14. Neuron 2002; 35:25-38

[35] Yang L, Dong F, Yang Q, Yang P-F, Wu R, Wu Q-F, et al. FGF13 selectively regulates heat nociception by interacting with Nav1.7. Neuron 2017; 93; 806-21.

[36] Yang Y, Wang Y, Li S, Xu Z, Li H, Ma L, Fan J, Bu D, Liu B, Fan Z, Wu G, Jin J, Ding B, Zhu X, Shen Y. Mutations in SCN9A, encoding a sodium channel alpha subunit, in patients with primary erythermalgia. J. Med. Genet. 2004; 41:171-4.

[37] Zhou X, Wang L, Hasegawa H, Amin P, Han BX, Kaneko S, He Y, Wang F. Deletion of Pik3C3/Vps34 in sensory neurons causes rapid neurodegeneration by disrupting the endosomal but not the autophagic pathway. Proc. Natl. Acad. Sci. USA 2010; 107:9424-9.

[38] Zurborg S, Piszczek A, Martinez C, Hublitz P, Al Banchaabouchi M, Moreira P, Perlas E, Heppenstall PA. Generation and characterization of an *Advillin-Cre* driver mouse line. Molec. Pain 2011; 7:66.

Figure Legends

Figure 1. Somatosensation in mice harboring global or peripheral neuron-specific *Fhf2* gene ablation. (A) Tail flick assay of heat nociception in wild-type and *Fhf2^null^* mice. Cumulative percentage plot for tail flick as function of tail heating time for wild-type female (*Fhf2^+/+^*), wild-type male (*Fhf2^+/Y^*), *Fhf2^null^* female (*Fhf2^-/-^*) and *Fhf2^null^* male (*Fhf2^-/Y^*) mice. Male and female *Fhf2^null^* mice were not responsive to heating. *n*, number of mice. (B) FHF-2 protein expression in wild-type and mutant mice. Post-nuclear protein extracts from whole brain, cerebral cortex, and DRGs were subjected to SDS polyacrylamide gel electrophoresis and immunoblotting with antibodies directed against the C-terminus of FHF2 shared by all FHF2 isoforms. Lower molecular weight FHF2 (FHF2B) and higher molecular weight FHF2 isoforms were present in wild-type, but not *Fhf2^null^*, cerebral cortex (left). Male mice carrying a floxed *Fhf2* allele (*Fhf2^fl/Y^*) and the advillin-Cre transgene (*Adv-Cre*) showed complete loss of FHF2B expression in DRG, while retaining FHF2 expression in brain. (C) Tail flick assay in mice with DRG-specific ablation of FHF2 expression. Cumulative percentage tail flick plot shows that mice with DRG-specific ablation of FHF-2 expression (*Fhf2^fl/Y^:Adv-Cre*) were insensitive to heating, while control mice (*Fhf2^+/Y^:Adv-Cre* and *Fhf2^fl/Y^*) all rapidly responded to tail heating. *n*, number of mice. (D) Paw pinch assay of mechanical nociception in wild-type and *Fhf2^null^* mice. For each animal, the footpad was squeezed with a force-calibrated forceps until a withdrawal response was elicited. n.s., not significant impairment of mechanical nociception. (E) Von Frey filament assay of touch sensation in wild-type and *Fhf2^null^* mice. Five male wild-type (*Fhf2^+/Y^*) and mutant (*Fhf2^-/Y^*) mice were placed without restraint on a mesh platform, a hind paw footpad was touched with nylon filaments of graded thickness/force, and sensation was scored by paw withdrawal. Five trials were conducted for each filament per mouse (25 trials per genotype cohort). Graph plots number of trials (out of 25) that a response was elicited. *, $P < 0.004$; **, $P < 0.0003$. (F) Footpad sensory fiber innervation in wild-type and *Fhf2^null^* mice. Sensory fibers in footpad cryosections were visualized by immunofluorescence using antibodies to PGP-9. Dashed line indicates outer surface.

Figure 2. FHF-2 modulation of sensory neuronal sodium channel inactivation. (A) Voltage dependence of $Na_v1.7$ steady-state inactivation and activation with or without FHF2B at 25^oC. The presence of FHF2B induced a 15 mV depolarizing shift in $V_{1/2}$ inactivation without significantly impacting activation. n, number cells analyzed. (B) $Na_v1.7$ current traces with or without FHF2B at 25^oC and 40^oC. . (C) $Na_v1.7$ current decay time constants at -20 mV +/- FHF2B at 25^oC, 35^oC , and 40^oC. FHF2B slows channel inactivation at all temperatures.. (D-F) $Na_v1.7$ peak conductance upon ramped vs. step depolarization +/- FHF2 at (D) 25^oC, (E) 35^oC, and (F) 40^oC. Slower voltage ramp rates reduce sodium channel opening via closed-state inactivation, with greater reduction in the absence of FHF2B. (G) Steady state inactivation of TTX-resistant sodium channels in wild-type and *Fhf2^null* DRG neurons. $V_{1/2}$ inactivation of TTX-resistant currents undergoes a hyperpolarizing shift in the absence of FHF2.

Figure 3. Excitation by heat of acutely dissociated DRG neurons. (A-C) Optical detection of heat sensitive nociceptor excitation. Acutely dissociated wild-type DRG neurons were cultured overnight and stained with di-8-ANEPPS. A field of cells was imaged with high speed CCD camera while heating to 43^oC. (A) A single image shows fluorescence of four neurons (color circled) along with debris including myelin. (B) Fluorescence intensity monitored at 2 kHz shows action potentials generated in cell 2. (C) Bright field and fluorescent high magnification imaging of cell 2. Scale bar, 2 microns. (D) Voltage recording of a patched 12 micron diameter wild-type neuron during heating and cooling. (E) Voltage recording of a patched 12 micron diameter *Fhf2^null* neuron. (F-J) Excitation properties of wild-type and *Fhf2^null* heat sensitive cells. (F) Stimulus-evoked spike amplitude, (G) temperature threshold for spike initiation, (H) maximum temperature-evoked spike frequency, (I) spike amplitude at 44.5^oC, and (J) inward current density upon voltage clamp step from -70 mV to 0 mV. Number of recorded cells and standard errors are indicated for each recording parameter. *n.s.*, not significant.

Figure 4. Temperature modulation of C-fiber compound action action potential conduction. (A) Stimulus response of wild-type saphenous nerve. The nerve maintained at 30-31°C was stimulated at one end at time = 0 msec. The recording electrode detected an instantaneous stimulus artifact, fast arriving voltage deflections (1-3 msec) reflecting activation and conduction of myelinated axons, and slow C-fiber compound action potential (CCAP) from 10-20 msec post-stimulus. (B) Stimulus response of *Fhf2null* saphenous nerve. A similar CCAP was detected at 30-31°C. (C) Superimposed CCAP traces of a wild-type nerve stimulated every 9 sec during cooling from 44.3°C down to ~30°C. CCAPs displayed a near-linear increase in amplitude and decrease in velocity across the temperature range. (D) Superimposed CCAP traces of an *Fhf2null* nerve stimulated every 9 sec during cooling from 44.3°C down to ~30°C. The amplitude of the CCAP negative deflection fell precipitously when nerve temperature exceeded 41°C. (E) Plot of CCAP amplitudes versus temperature. For each of four wild-type and four *Fhf2null* nerves, the positive-to-negative peak-to-peak CCAP amplitudes at each temperature were plotted as a percentage of the amplitude recorded at 27°C. Wild-type nerves all showed near-linear amplitude decrement from minimal to maximal temperature, while *Fhf2null* nerves displayed enhanced decrements above 41°C. (F) Histogram representation of CCAP amplitude decrements with increasing temperature. For each of seven wild-type and six *Fhf2null* nerve recordings, CCAP amplitude ratios for 32°C vs. 28°C, 36°C vs. 32°C, 40°C vs. 36°C, and 44°C vs. 40°C. The decrement in CCAP amplitude is significantly greater for *Fhf2null* nerves versus wild-type nerves for 44°C vs. 40°C.

Figure 5. Computational modeling of wild-type and *Fhf2null* nociceptors. Wild-type and *Fhf2null* nociceptor models differ only in the inactivation gating parameters of the TTX-sensitive and TTX-resistant voltage-gated sodium conductances. All gated conductances have temperature-dependent state transition rates. (A-D) Simulation of axonal conduction. The long axon of each model was stimulated at one point with a 100 μsec current pulse,

and membrane voltage was monitored at 0, 1, 2, 3, and 4 mm from the site of stimulation. (A) Wild-type model at 37°C. (B) Wild-type model at 43°C. (C) *Fhf2^{null}* model at 37°C. (D) *Fhf2^{null}* model at 43°C; note conduction failure beyond 1 mm from point of stimulation.   (E-G) Simulation of mechanical nociception. (E) Branching morphology of the afferent terminal. The terminal architecture, as published elsewhere [4], was incorporated into wild-type and *Fhf2^{null}* models. Temperature throughout nociceptor was set to 37°C. Arrows at all terminal branch tips indicate sites of up-ramping current injection delivered from 0.5 to 2.2 seconds of simulation followed by down-ramping from 2.2 to 3 seconds. This protocol simulated opening of TRP channels at branch tips due to mechanical stimulation. (F) Central axonal voltage recording of wild-type model neuron during simulation in (E). The current ramp (light blue trace) evoked a train of action potentials that initiated peripherally and propagated antidromically. (G) Voltage recording of *Fhf2^{null}* model neuron during simulation in (E). A train of action potentials was generated, though fewer in number than in the wild-type model. (H-J) Simulation of heat excitability. (H) Terminal branching architecture. Branch segments within 200 microns of tips were set to 43°C (red), while remainder of neuron was maintained at 37°C. Current ramp injection into branch tips simulated heat stimulation. (I) Central axonal voltage recording of wild-type neuron during simulation in (H). Stimulation evoked a train of antidromic action potentials. (J) Voltage recording of *Fhf2^{null}* neuron during simulation in (H). The mutant neuron was not excited by stimulation, simulating the heat nociception deficit observed in vivo. (K-M) Simulation of heat-induced excitation in "acutely dissociated" model soma. (K) Model soma were disconnected from axons, somatic temperature was set to 45°C (red), and up-ramped somatic inward current simulated heat-induced excitation of the isolated soma. (L) Wild-type and (M) *Fhf2^{null}* isolated soma fired trains of action potentials in response to current injections (light blue traces).

Figure 1

Figure 2

Figure 3

Figure 5