Pervasive species-specific repulsion among adult tropical trees

Authors: Michael Kalyuzhny[1*], Jeffrey K. Lake[2], S. Joseph Wright[3], Annette M. Ostling[1].

Affiliations:

[1] Department of Integrative Biology, The University of Texas at Austin; 2415 Speedway #C0930, Austin, TX 78712.

[2] Department of Biology, Adrian College; Adrian, MI.

[3] Smithsonian Tropical Research Institute; Apartado 0843□03092, Balboa, Republic of Panama.

*Corresponding author. Email: michael.kalyuzhny@utexas.edu

Abstract:

For species to coexist, performance must decline as the density of conspecific individuals increases. Although evidence for such Conspecific Negative Density Dependence (CNDD) exists in forests, the spatial repulsion it should produce has not been demonstrated in adults. Here we show that in comparison to a null model of stochastic birth, death and limited dispersal, the adults of dozens of tropical forest tree species show almost ubiquitous and strong spatial repulsion, some to surprising distances of ~100 meters. We use simulations to show such strong repulsion can only occur if CNDD considerably exceeds Heterospecific Negative Density Dependence, an even stronger condition required for coexistence, and that large-scale repulsion can indeed result from small-scale CNDD. These results highlight the power of limited dispersal spatial null models.

One-Sentence Summary: Spatial distributions of tropical trees reflect a strong negative effect of conspecifics, exceeding that of heterospecifics

Main Text:

How can hundreds of species coexist on small spatial scales? Some tropical forests contain over 250 species of trees in a single hectare, or over 1,100 species of trees and shrub in 25 hectares (*1*), begging for mechanistic explanation. According to coexistence theory, a necessary condition for species coexistence is that the performance of species decreases as they become more common, preventing any one species from taking over, a phenomenon known as Conspecific Negative Density Dependence (CNDD) (*2, 3*). A stronger necessary condition is that CNDD is stronger than Heterospecific Negative Density Dependence (HNDD) (*2, 3*), implying that species mostly 'suffer' from their own kind due to niche differences, that stabilize population size. Coexistence is assured if, in addition to the aforementioned condition, species are sufficiently similar in their competitive ability (*3, 4*).

Considerable evidence has been presented for CNDD acting on early life stages in tropical and temperate forests, and variation in CNDD has been hypothesized to underlie the latitudinal gradient from high tree species diversity at the equator to lower diversity at higher latitudes (*3, 5-7*). For example, the abundance of seedlings, seeds and adults have been shown by numerous works to reduce the number and survival of seedlings and saplings (*8-11*) and several works have also presented evidence that CNDD is stronger than HNDD (*8, 11*). Nevertheless, the validity of many of these findings has been questioned on methodological grounds (*2, 10*). Crucially, current evidence suggests that CNDD has very limited effect on adults (*12, 13*). Moreover, adult tropical trees can live for hundreds of years (*14*) and integrate density dependent effects throughout their life. CNDD at one life stage can be offset by positive effects at another (*2, 15*), or have a weak effect on lifetime performance, begging the question – what is the overall integrated effect of conspecifics on populations? Importantly, this integrated effect is what matters for coexistence (*2*).

A potential approach to evaluating the integrated effect of CNDD across life stages is through the study of the spatial distribution of adult trees. Because CNDD has been shown to operate at short scales around adults (*3, 8, 9*), it is theorized to generate spatial 'repulsion' between adults, potentially leading to spatial overdispersion (*16, 17*). Alas, classical studies of the spatial distributions of adult trees found them to be almost ubiquitously *aggregated* in tropical forests (*18-20*). Hence, individuals are typically closer to each other than expected, especially at small scales where CNDD is expected to be the strongest (*18-20*). Furthermore, many spatial patterns in tropical forests resemble the expectations of neutral models (*21-23*), or reflect habitat heterogeneity (*24-26*), with little signs of CNDD. These findings suggest that sensitivity to CNDD at early life stages is severely diluted, overwhelmed or obscured before reaching adulthood by processes that create aggregation such as habitat specificity, facilitation and dispersal limitation (*2, 3*). If CNDD at early life stages does not propagate to affect the adults (as evidence for this is lacking (*27*)), how can it maintain species diversity?

Fig. 1. Spatial point patterns of adult *Beilschmiedia pendula* trees; The (**A**) observed pattern appears aggregated compared to (**B**) realizations of the Complete Spatial Randomness (CSR) null model, but overdispersed compared to (**C**) the standard variant of the Dispersal Limitation (DL) null model. Both nulls use the same abundance as the observed. The mean dispersal distance of *B. pendula* is estimated at 10.7 meters (*28*). Fig. S1 presents realizations of the other variants of DL, while Fig. S2 zooms in on the clump of trees at the top-left of (**C**).

Classical studies on spatial patterns of adult trees compared their distributions to the expectation of the Complete Spatial Randomness (CSR) null model (*16, 18, 19*). This model distributes individuals with equal probability across the (often arbitrary) survey area, inevitably assigning some to sites that they would not be able to reach due to dispersal limitation (Fig. 1a,b). We believe that a more ecologically sound null model is Dispersal Limitation (DL, spatially explicit neutral theory) (*21, 29*), which assumes that abundance and distribution are determined by stochastic births and deaths and random, yet limited, dispersal. Using DL as a null model should incorporate the aggregation created by dispersal limitation into the null

expectations (Fig 1c, Fig. S1-2), potentially revealing signals of biotic interactions and habitat specificity.

Here we compare the spatial patterns of tropical tree species in a Panamanian forest to their expectation under DL, using previously estimated species-specific dispersal distance distributions (*28, 30*). Our goal is to examine the prevalence, magnitude and scale of overdispersion and aggregation with respect to DL. Since spatial repulsion is the signature of CNDD (*16, 31*), which is caused by local interactions and has been shown to operate on small spatial scales (~20 m., *3, 8, 9*), we hypothesize that some species, for which CNDD propagates to the adult stage, would show overdispersion at small scales. We then use a spatially explicit simulation model to examine whether the spatial patterns we find provide insight into the relative strength of CNDD versus HNDD. Finally, we examine the implications of our findings for tropical tree diversity.

We perform the analysis with 41 species that have reliable estimates of dispersal distance distributions, obtained using known locations of seed-bearing trees and seed arrivals (*28*) in the Barro Colorado Island 50 ha forest plot (BCI) (*32*). These estimates have been corroborated using genotyped seeds and adults (*33*), highlighting their reliability. DL simulations, parameterized with the dispersal distance distributions, are run for every species separately, and the observed spatial distribution of adults in 2015 is compared to the expectation under DL. For that comparison we develop analogs of the two most used point-pattern statistics: the Clark-Evans nearest neighbor statistic and the relative neighborhood density (a.k.a. pair correlation function) (*16, 18, 19*) statistic. We compare the observed mean (or median, see methods) distance of trees to their nearest conspecific neighbor, NND_{obs}, to its value under the null (NND_{null}) using the Excess Distance (*ED*) statistic:

(eq. 1) $ED = \log(NND_{obs}) - \overline{\log(NND_{null})}$,

where the averaging (denoted by an overbar in eqs. 1 and 2) is over samples of the DL model. We also compute the Excess neighborhood Abundance (*EA(r)*), namely the observed number of conspecific neighbors of an average tree at distances r to $r + \Delta r$, $N_{obs}(r)$, relative to its value under the null $N_{null}(r)$:

(eq. 2) $EA(r) = \log(N_{obs}(r)) - \overline{\log(N_{null}(r))}$.

To examine the robustness of our results ('Standard'), we repeat this analysis for the standard DL null hypothesis and five additional variants of the DL null model. The five variants 1) use earlier estimates of dispersal, for a broader set of 81 species ('H2001' (*30*)); 2) add 10% of global dispersal, representing widespread long-distance dispersal ('LDD'); these variants consider different types of dispersal distance distributions and increase the dispersion under the null due to longer dispersal distances, making the analysis more conservative; 3) incorporate a lag of one generation from dispersal to recruitment, likewise increasing dispersion under the null since parent and offspring often do not co-occur; 4) start with the distribution of conspecific trees

in 1985 and examine only the recruits to the adult class appearing by 2015 ('Recruits'); 5) Fix the observed locations of all adult trees ('Fixed'), while allowing their species identity to vary dynamically, controlling the spatial distribution of trees of all species. Finally, we also show that clustered dispersal that may result from animals depositing clusters of seeds (*34*) would typically produce aggregation compared to our null models with independent seeds (see Supplementary Text).

	Excess Abundance (EA) at 20 meters			Excess nearest neighbor Distance (ED)		
Null	Values	Overdispersed	Aggregated	Values	Overdispersed	Aggregated
Standard	-1.5 ± 1.2 [-3.9, 1.3]	38 (34)	3 (2)	0.9 ± 0.61 [-0.65, 2.1]	39 (36)	2 (1)
H2001	-1.1 ± 1.3 [-4.7, 1.3]	69 (47)	12 (3)	0.69 ± 0.67 [-0.45, 2.8]	75 (60)	6 (3)
Lag	-0.99 ± 1.1 [-3.3, 1.5]	33 (28)	8 (2)	0.57 ± 0.52 [-0.72, 1.7]	36 (31)	5 (2)
LDD	-0.97 ± 1.1 [-3.4, 1.4]	35 (27)	6 (3)	0.77 ± 0.71 [-1.1, 2.1]	37 (32)	4 (2)
Recruits	-0.65 ± 1.1 [-3.1, 1.9]	34 (22)	7 (1)	0.69 ± 0.66 [-1.2, 2.6]	38 (33)	3 (1)
Fixed	-1.3 ± 1.1 [-3.7, 1.3]	37 (33)	4 (2)	0.86 ± 0.68 [-1.1, 2.1]	38 (36)	3 (2)

Table 1. Distributions and significance of the spatial statistics of Barro Colorado tree species in 2015 using the six variants of the DL null model as references. We compare the abundance of conspecific neighbors within 20 meters (left, eq. 2) and the nearest neighbor distance (right, eq. 1) to their expectations. For abundances (left), negative values represent repulsion and vice versa. For distances (right), positive values represent repulsion. Exponentiating the statistics would give the factor by which the observed density or distance exceeds expectations. The mean ± SD and the range (in square brackets) of the statistics over all species are presented, as well as the number of overdispersed and aggregated species, along with the number of statistically significant (with $\alpha = 0.05$) species in parentheses. For the LDD and Recruits nulls, median nearest neighbor distance is used, while the mean is calculated in the other cases.

Contrary to previous findings based on CSR (*18, 19*), BCI tree species are almost ubiquitously and strongly overdispersed on short scales (~20 meters) when compared to all six variants of the Dispersal Limitation null model (Table 1). On average, trees have 50% - 80% fewer conspecific neighbors than expected within 20 meters, and their nearest conspecific neighbor is 1.8 - 2.5 times more distant than expected (these numbers are the values of the mean statistics in Table 1, exponentiated). Even the most common shrubs, treelets and trees (e.g. *Trichilia tuberculata*, Fig. 2), are overdispersed, as are relatively rare species (Data S1) and these patterns are qualitatively similar under the different DL null model variants (Table 1). Fig. 2

shows typical, monotonically increasing *EA*(*r*)s for four species, along with their distribution maps in comparison with the expectations of the DL null, while Fig. S3-4 show *EA*(r) for all non-rare species we analyzed. Not surprisingly, some light-demanding and habitat specialist species deviate from this typical pattern, exhibiting increased aggregation reflective of the limited spatial extent of the conditions they require (Fig. S5).

Fig. 2. Spatial distributions of four typical species and their Excess neighbor abundance at distance *r*, *EA*(*r*) (eq. 2). Each row presents a different species, while the columns present for each species the observed distribution (left) versus a single realization of the Standard DL null (center). *EA*(*r*) is presented in the rightmost, shaded column, with the orange region indicating the 95% simulation envelope of the DL null. *EA*(*r*) of 0 indicates agreement with DL while values < 0 indicate overdispersion. All distribution maps are 1000 by 500 meters. Compared to the DL null, trees have considerably fewer neighbors than expected at distances up to 100 m.

What underlies this widespread, strong and robust (under different assumptions of the null models) overdispersion? Theoretical studies show that while aggregation can be caused by dispersal limitation (*17, 35*), habitat specificity (*31*) and facilitation (*36*), overdispersion is the signature of negative density dependence (*16, 17*). Hence, we infer that the overdispersion we

detect reflects the spatial signature of density dependence acting on juveniles, for which much evidence on BCI exists (*8, 10, 11, 37*), propagating to affect the adults.

Fig. 3. Excess neighborhood Abundance at 20 meters, *EA*(20) (eq. 2) in simulations. In each simulation, the magnitude of CNDD (Q_C, Y axis) and HNDD (Q_H, X axis) were controlled, and the color of each square represents the mean *EA*(20) of simulated populations. In the simulations, seeds are dispersed an average distance of $D = 7$ meters (**a**) or $D = 20$ meters (**b**) and the survival of each is determined by both the density of conspecific adults and the density of heterospecific adults in a way that gives large weight to adults within ~ 7 meters of where the seed lands. For perspective, a seed landing right near a single conspecific would have a recruitment probability of $1/(1+Q_C)$, and more nearby conspecifics and heterospecifics would reduce this probability further. See Methods and eq. S3 for more details. Results are not shown for cases where HNDD > CNDD (below the diagonal) since priority effects lead to one species taking over these communities. When CNDD = HNDD (on the diagonal) overdispersion is weak, and the strongest overdispersion is observed when CNDD >> HNDD.

     Importantly, we show here using a simulation model that the observed overdispersion is only consistent with a scenario when CNDD >> HNDD. Figures 3 and S6 show the degree of overdispersion generated in spatially explicit stochastic simulations of community dynamics where CNDD and HNDD both reduce the survival of dispersing juveniles and are set to be strong out to a distance of about 7m. (Fig. S10). First, when the magnitude of CNDD = HNDD (along the diagonal in Fig. 3), overdispersion in the simulations quickly asymptotes with the magnitude of density dependence, at values indicating much lower overdispersion than observed on BCI: while *EA*(20) is ~ -1.5 in BCI (Table 1), the simulated *EA*(20) asymptotes at much lower values (~-0.18 or ~-0.55, Fig. 3, S7-8). Intuitively, when HNDD = CNDD, the competitive effect of neighbors is equal regardless of their species, and so density dependence cannot generate a strong repulsion from conspecifics. Second, only the scenario CNDD >> HNDD (upper left

corners in Fig. 3) is compatible with the magnitude of overdispersion observed on BCI. Finally, our results using the 'Fixed' null variant (Table 1), in which the locations of all adults are preserved, are similar to the analysis using the standard null variant. The 'Fixed' variant preserves the spacing between all trees of all species, resulting from competition for light, soil resources and other non-species-specific factors, hence incorporating HNDD and some CNDD into the null expectation. The strong overdispersion compared with this variant further confirms the weakness of HNDD compared to CNDD.

While we expected overdispersion to occur primarily at the short distances at which CNDD operates (*8, 11*), we find several species are significantly and substantially overdispersed at distances up to ~100 meters, with *all* species having on average ~ 45% fewer neighbors 75 – 125 meters away from a typical tree (under the Standard null and all variants except the 'LDD' or 'Recruits' null, Figure 2, Table S1). Our simulations show that such long-distance overdispersion can result from short-distance CNDD (acting to seven meters) (Fig. S9). In hindsight, this result appeals to intuition. While each tree "repels" only its neighbors, they in turn repel their neighbors and so forth, creating large-scale patterns from short-distance interactions.

Previous analyses aimed at detecting CNDD from spatial patterns have reached contradictory conclusions. Many did not detect evidence of CNDD (*21-26*). Several revealed spatial repulsion of juveniles from conspecific adults (e.g., *7, 9*). Others fit complex process-based models to all individuals (including subadults) finding evidence for CNDD (*35, 38*). However, none of these studies provide evidence that the effects of CNDD propagate to affect the adult stage. Focusing on larger trees and community-level spatial pattern, May et al. (*21*) found that *all* trees ≥ 10 CM DBH (*not only conspecifics*) are overdispersed compared to CSR at distances < 10 m, indicating general density dependence among all trees of all species. Several studies found that conspecific aggregation decreases with tree size (*18, 20, 39*), but those that considered more than a handful of species found significant reductions in only 15% - 22% of species studied (*18, 20*). In contrast, we found that almost all species are overdispersed at the adult stage. This difference might occur as the effects of CNDD accumulate through ontogeny with the strongest effects at early life stages (*12*) before the sapling stage considered by references 18, 20. This highlights the utility of comparison to the DL null model, which uses seed rain as the reference. Finally, no previous study of spatial pattern evaluated the relative strength of CNDD and HNDD, while we show that species-specific spatial patterns of adults are only consistent with a scenario when CNDD >> HNDD across life stages.

These findings have important implications for the maintenance of species diversity. First, our finding that CNDD must be substantially greater than HNDD for many species implies large niche differences among species, which may stabilize diversity because species that increase in abundance suffer reduced fitness. It has recently been debated whether differences in species-level fitness can overwhelm the stabilizing effects of CNDD (*40, 41*), so the potential of CNDD to maintain species diversity needs further investigation. However, our work provides a key missing ingredient in this line of research on diversity maintenance in forests — namely evidence that the overall forces of density dependence *across life stages* set up substantial stabilizing niche differences, a key requirement for stable coexistence. Moreover, given that the

forest is overall saturated with individuals (*29*), the finding that individuals of most species have fewer *conspecific* neighbors up to long distances implies an increase in *heterospecific* neighbors and an increase in compositional turnover at these distances. This is consistent with the reported increased spatial turnover (beta diversity) at scales of 20-200 meters in BCI compared with a neutral model (*22*). Beta diversity has been extensively studied and is believed to be driven mostly by spatial heterogeneity, dispersal limitation and interspecific competition (*22, 42, 43*). Our results suggest that the role of CNDD in shaping turnover should also be carefully considered.

More broadly, patterns of species distributions at local scales have been investigated and compared to CSR for over one hundred years (*44*), often with the aim of detecting the signals of habitat specificity and biotic interactions (*18, 19, 26, 45*). However, theoretical models predict that dispersal limitation alone could override and obscure the signal of biotic interactions and dominate spatial patterns, creating ubiquitous aggregated distributions (*17*). Indeed, spatial patterns have been found to be ubiquitously aggregated with respect to the CSR null (*18, 19, 45*). In contrast, a richness of spatial patterns is revealed with respect to the DL null: most species are overdispersed, owing to CNDD, while some habitat and gap specialists are more aggregated (Fig. S5). This demonstrates the power of the DL null to filter out much of the effect of dispersal limitation and to reveal the overall effect of biotic interactions and habitat specificity on the spatial distributions of adult individuals. Our work thus promotes a complementary definition of the classical terms "aggregation" and "overdispersion", rooted in ecological processes and theory. We believe that a more widespread use of this approach will have much capacity to shed light on the processes shaping the assembly of ecological communities.

References and Notes

1. S. J. Wright, Plant diversity in tropical forests: a review of mechanisms of species coexistence. *Oecologia* **130**, 1-14 (2002).
2. M. J. E. Broekman *et al.*, Signs of stabilisation and stable coexistence. *Ecology Letters* **22**, 1957-1975 (2019).
3. L. Hülsmann, R. A. Chisholm, F. Hartig, Is Variation in Conspecific Negative Density Dependence Driving Tree Diversity Patterns at Large Scales? *Trends in Ecology & Evolution*, (2020).
4. P. Chesson, Mechanisms of maintenance of species diversity. *Annual Review of Ecology and Systematics* **31**, 343-366 (2000).
5. J. A. LaManna *et al.*, Plant diversity increases with the strength of negative density dependence at the global scale. *Science* **356**, 1389-1392 (2017).
6. D. H. Janzen, Herbivores and the number of tree species in tropical forests. *American Naturalist* **104**, 501-528 (1970).
7. J. A. LaManna, S. A. Mangan, J. A. Myers, Conspecific negative density dependence and why its study should not be abandoned. *Ecosphere* **12**, (2021).
8. L. S. Comita, H. C. Muller-Landau, S. Aguilar, S. P. Hubbell, Asymmetric density dependence shapes species abundances in a tropical tree community. *Science* **329**, 330-332 (2010).
9. P. Marchand *et al.*, Seed-to-seedling transitions exhibit distance-dependent mortality but no strong spacing effects in a Neotropical forest. *Ecology* **101**, (2020).
10. M. Detto, M. D. Visser, S. J. Wright, S. W. Pacala, Bias in the detection of negative density dependence in plant communities. *Ecology Letters* **22**, 1923-1939 (2019).
11. E. Lebrija-Trejos, P. B. Reich, A. Hernandez, S. J. Wright, Species with greater seed mass are more tolerant of conspecific neighbours: a key driver of early survival and future abundances in a tropical forest. *Ecology Letters* **19**, 1071-1080 (2016).
12. Y. Zhu, L. S. Comita, S. P. Hubbell, K. P. Ma, Conspecific and phylogenetic density-dependent survival differs across life stages in a tropical forest. *Journal of Ecology* **103**, 957-966 (2015).
13. P. T. Green, K. E. Harms, J. H. Connell, Nonrandom, diversifying processes are disproportionately strong in the smallest size classes of a tropical forest. *Proceedings of the National Academy of Sciences of the United States of America* **111**, 18649-18654 (2014).
14. J. Q. Chambers, N. Higuchi, J. P. Schimel, Ancient trees in Amazonia. *Nature* **391**, 135-136 (1998).
15. M. J. O'Brien *et al.*, Demographic consequences of heterogeneity in conspecific density dependence among mast-fruiting tropical trees. *Proceedings of the Royal Society B-Biological Sciences* **289**, (2022).
16. T. Wiegand, K. A. Moloney, *Handbook of spatial point-pattern analysis in ecology*. (CRC press, 2013).
17. J. Chave, H. C. Muller-Landau, S. A. Levin, Comparing classical community models: Theoretical consequences for patterns of diversity. *American Naturalist* **159**, 1-23 (2002).
18. R. Condit *et al.*, Spatial patterns in the distribution of tropical tree species. *Science* **288**, 1414-1418 (2000).

19. S. P. Hubbell, Tree dispersion, abundance, and diversity in a tropical dry forest. *Science* **203**, 1299-1309 (1979).

20. R. Bagchi *et al.*, Spatial patterns reveal negative density dependence and habitat associations in tropical trees. *Ecology* **92**, 1723-1729 (2011).

21. F. May, A. Huth, T. Wiegand, Moving beyond abundance distributions: neutral theory and spatial patterns in a tropical forest. *Proceedings of the Royal Society B-Biological Sciences* **282**, (2015).

22. R. Condit *et al.*, Beta-diversity in tropical forest trees. *Science* **295**, 666-669 (2002).

23. E. Seri, Y. E. Maruvka, N. M. Shnerb, Neutral Dynamics and Cluster Statistics in a Tropical Forest. *American Naturalist* **180**, E161-E173 (2012).

24. T. Wiegand, S. Gunatilleke, N. Gunatilleke, Species associations in a heterogeneous Sri lankan dipterocarp forest. *American Naturalist* **170**, E77-E95 (2007).

25. G. C. Shen *et al.*, Species-area relationships explained by the joint effects of dispersal limitation and habitat heterogeneity. *Ecology* **90**, 3033-3041 (2009).

26. J. B. Plotkin *et al.*, Species-area curves, spatial aggregation, and habitat specialization in tropical forests. *Journal of Theoretical Biology* **207**, 81-99 (2000).

27. G. G. Mittelbach, B. J. McGill, *Community ecology*. (Oxford University Press, 2019).

28. H. C. Muller-Landau, S. J. Wright, O. Calderon, R. Condit, S. P. Hubbell, Interspecific variation in primary seed dispersal in a tropical forest. *Journal of Ecology* **96**, 653-667 (2008).

29. S. P. Hubbell, *The unified neutral theory of biodiversity and biogeography*. Monographs in Population Biology (Princeton University Press, 2001), vol. 32.

30. H. C. Muller-Landau, Seed dispersal in a tropical forest: empirical patterns, their origins, and their consequences for community dynamics. (2001).

31. C. Brown, J. B. Illian, D. Burslem, Success of spatial statistics in determining underlying process in simulated plant communities. *Journal of Ecology* **104**, 160-172 (2016).

32. R. Condit *et al.*, Complete data from the Barro Colorado 50-ha plot: 423617 trees, 35 years. *URL https://doi. org/10.15146/5xcp-0d46*, (2019).

33. F. A. Jones, H. C. Muller-Landau, Measuring long-distance seed dispersal in complex natural environments: an evaluation and integration of classical and genetic methods. *Journal of Ecology* **96**, 642-652 (2008).

34. R. S. Manasse, H. F. Howe, Competition For Dispersal Agents Among Tropical Trees - Influences Of Neighbors. *Oecologia* **59**, 185-190 (1983).

35. M. Detto, H. C. Muller-Landau, Fitting Ecological Process Models to Spatial Patterns Using Scalewise Variances and Moment Equations. *American Naturalist* **181**, E68-E82 (2013).

36. J. Molofsky, J. D. Bever, J. Antonovics, Coexistence under positive frequency dependence. *Proceedings of the Royal Society B-Biological Sciences* **268**, 273-277 (2001).

37. D. L. Forrister, M. J. Endara, G. C. Younkin, P. D. Coley, T. A. Kursar, Herbivores as drivers of negative density dependence in tropical forest saplings. *Science* **363**, 1213-+ (2019).

38. M. Detto, H. C. Muller☐Landau, Rates of formation and dissipation of clumping reveal lagged responses in tropical tree populations. *Ecology* **97**, 1170-1181 (2016).

39. R. W. Sterner, C. A. Ribic, G. E. Schatz, Testing for Life Historical Changes in Spatial Patterns of 4 Tropical Tree Species. *Journal of Ecology* **74**, 621-633 (1986).

40. R. A. Chisholm, T. Fung, Janzen-Connell Effects Are a Weak Impediment to Competitive Exclusion. *American Naturalist* **196**, 649-661 (2020).

41.	D. J. B. Smith, The functional form of specialised predation affects whether Janzen-Connell effects can prevent competitive exclusion. *Ecology Letters* **25**, 1458-1470 (2022).

42.	J. M. Chase, J. A. Myers, Disentangling the importance of ecological niches from stochastic processes across scales. *Philosophical Transactions of the Royal Society B-Biological Sciences* **366**, 2351-2363 (2011).

43.	H. Segre *et al.*, Competitive exclusion, beta diversity, and deterministic vs. stochastic drivers of community assembly. *Ecology Letters* **17**, 1400-1408 (2014).

44.	H. A. Gleason, Some applications of the quadrat method. *Bulletin of the Torrey Botanical Club* **47**, 21-33 (1920).

45.	J. M. Levine, D. J. Murrell, The community-level consequences of seed dispersal patterns. *Annual Review of Ecology Evolution and Systematics* **34**, 549-574 (2003).

46.	R. Nathan, E. Klein, J. J. Robledo-Arnuncio, E. Revilla, Dispersal kernels. *Dispersal ecology and evolution*, 187-210 (2012).

47.	S. P. Hubbell *et al.*, Light-gap disturbances, recruitment limitation, and tree diversity in a neotropical forest. *Science* **283**, 554-557 (1999).

48.	R. Condit, *Tropical Forest Census Plots: Methods and Results from Barro Colorado Island, Panama and a Comparison with Other Plots*. (Springer, 1998).

49.	J. S. Clark, M. Silman, R. Kern, E. Macklin, J. HilleRisLambers, Seed dispersal near and far: Patterns across temperate and tropical forests. *Ecology* **80**, 1475-1494 (1999).

50.	S. J. Wright *et al.*, Understanding strategies for seed dispersal by wind under contrasting atmospheric conditions. *Proceedings of the National Academy of Sciences of the United States of America* **105**, 19084-19089 (2008).

51.	H. F. Howe, Scatter-dispersal And Clump-dispersal And Seedling Demography - Hypothesis And Implications. *Oecologia* **79**, 417-426 (1989).

52.	N. Ruger *et al.*, Beyond the fast-slow continuum: demographic dimensions structuring a tropical tree community. *Ecology Letters* **21**, 1075-1084 (2018).

53.	S. J. Wright *et al.*, Functional traits and the growth-mortality trade-off in tropical trees. *Ecology* **91**, 3664-3674 (2010).

54.	Y. C. Lin, L. W. Chang, K. C. Yang, H. H. Wang, I. F. Sun, Point patterns of tree distribution determined by habitat heterogeneity and dispersal limitation. *Oecologia* **165**, 175-184 (2011).

55.	P. J. Clark, F. C. Evans, Distance to nearest neighbor as a measure of spatial relationships in populations. *Ecology* **35**, 445-453 (1954).

56.	S. P. Hubbell, J. A. Ahumada, R. Condit, R. B. Foster, Local neighborhood effects on long-term survival of individual trees in a neotropical forest. *Ecological Research* **16**, 859-875 (2001).

57.	M. Uriarte, C. D. Canham, J. Thompson, J. K. Zimmerman, N. Brokaw, Seedling recruitment in a hurricane-driven tropical forest: light limitation, density-dependence and the spatial distribution of parent trees. *Journal of Ecology* **93**, 291-304 (2005).

58.	T. Wiegand, I. Martinez, A. Huth, Recruitment in Tropical Tree Species: Revealing Complex Spatial Patterns. *American Naturalist* **174**, E106-E140 (2009).

59.	M. Uriarte, C. D. Canham, J. Thompson, J. K. Zimmerman, A neighborhood analysis of tree growth and survival in a hurricane-driven tropical forest. *Ecological Monographs* **74**, 591-614 (2004).

60.	S. Grote, R. Condit, S. Hubbell, C. Wirth, N. Ruger, Response of Demographic Rates of Tropical Trees to Light Availability: Can Position-Based Competition Indices Replace Information from Canopy Census Data? *Plos One* **8**, (2013).

61. R. A. Chisholm, J. W. Lichstein, Linking dispersal, immigration and scale in the neutral theory of biodiversity. *Ecology Letters* **12**, 1385-1393 (2009).

62. I. Volkov, J. R. Banavar, S. P. Hubbell, A. Maritan, Neutral theory and relative species abundance in ecology. *Nature* **424**, 1035-1037 (2003).

63. E. G. Leigh, *Tropical Forest Ecology : A View from Barro Colorado Island.* (Oxford University Press, USA, 1999).

64. M. K. Kalyuzhny, R. Kadmon, N. M. Shnerb, A neutral theory with environmental stochasticity explains static and dynamic properties of ecological communities. *Ecology Letters* **18**, 572-580 (2015).

65. K. E. Harms, R. Condit, S. P. Hubbell, R. B. Foster, Habitat associations of trees and shrubs in a 50-ha neotropical forest plot. *Journal of Ecology* **89**, 947-959 (2001).

Acknowledgments:

We thank R. Condit and R. Kadmon for comments. BCI forest dynamics research project was made possible by National Science Foundation grants to Stephen P. Hubbell: DEB-0640386, DEB-0425651, DEB-0346488, DEB-0129874, DEB-00753102, DEB-9909347, DEB-9615226, DEB-9615226, DEB-9405933, DEB-9221033, DEB-9100058, DEB-8906869, DEB-8605042, DEB-8206992, DEB-7922197, support from the Forest Global Earth Observatory, the Smithsonian Tropical Research Institute, the John D. and Catherine T. MacArthur Foundation, the Mellon Foundation, the Small World Institute Fund, and numerous private individuals, and through the hard work of over 100 people from 10 countries over the past three decades. The plot project is part the Forest Global Earth Observatory (ForestGEO), a global network of large-scale demographic tree plots.

Funding:

Michigan Life Sciences Fellowship (MK)

Zuckerman STEM Leadership Program (MK)

Sabbatical support from the University of Michigan and Adrian College (JKL)

Mcubed, The University of Michigan (MK, AMO)

Associate Professor Support Fund (MK, AMO)

Author contributions:

Conceptualization: MK

Planning: MK, JKL, AMO

Data generation: SJW

Data analysis: MK

Writing – original draft: MK

Interpreting results: MK, JKL, SJW, AMO

Writing – review & editing: MK, JKL, SJW, AMO

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: All data used in this analysis is publicly available and referenced to. All the code used to generate the results is included in the supplementary.

Supplementary Materials

Materials and Methods

Supplementary Text

Figs. S1 to S11

Tables S1

References (*46–65*)

Data S1