Working title

Spike-dependent dynamic partitioning of the *Locus coeruleus* network through noradrenergic volume release in a simulation of nucleus core

Shristi Baral[1,2]*, Hassan Hosseini[1]* , Kaushik More[1], Thomaz M.C. Fabrin[1], Jochen Braun[2+] & Matthias Prigge[1,3++]

1 Research Group Neuromodulatory Networks, Leibniz Institute for Neurobiology, 39118 Magdeburg, Germany
2 Cognition Biology, Otto-von Guericke University, 39118 Magdeburg, Germany
3 Center for Behavioral Brain Sciences, 39118 Magdeburg, Germany
* contributed equally

[+] e-mail:jochen.braun@ovgu.de
[++] e-mail:prigge@lin-magdeburg.de

Abstract

The *Locus coeruleus* (LC) modulates various neuronal circuits throughout the brain. Its unique architectural organization encompasses a net of axonal innervation that spans the entire brain; while its somatic core is highly compact. Recent research revealed an unexpected cellular input specificity within the nucleus that can give rise to various network states that either broadcast norepinephrine signals throughout the brain or pointedly modulate specific brain areas. Such adaptive input-output functions likely surpass our existing network models that build upon a given synaptic wiring configuration between neurons. As the distances between noradrenergic neurons in the core of the LC are unusually small, neighboring neurons could theoretically impact each other via volume transmission of NE. We therefore set out to investigate if such interaction could be mediated through noradrenergic alpha2-receptors in a spiking neuron model of the LC. We validate our model of LC neurons through comparison with experimental patch-clamp data and identify key variables that impact alpha2-mediated inhibition of neighboring LC neurons. Our simulation confirms a reliable autoinhibition of LC neurons after episodes of high neuronal activity that continue even after neuronal activity subsided. Also, dendro-somatic synapses inhibit spontaneous spiking in the somatic compartment of connected neurons in our model. We determined the exact position of hundreds of LC neurons in the mouse brain stem via a tissue clearing approach and, based on this, further determined that 25 percent of noradrenergic neurons have a neighboring LC neuron within less than a 25-micrometer radius. By modeling NE diffusion, we estimate that more than 15 percent of alpha2-adrenergic receptors fraction can bind NE within such diffusion radius. Our spiking neuron model of LC neurons predicts that repeated or long-lasting episodes of high neuronal activity induce partitioning of the gross LC network, and reduces the spike rate in neighboring neurons at distances smaller than 25 micrometers.
As these volume-mediate neighboring effects are challenging to test with the current methodology, our findings can guide future experimental approaches to test this phenomenon and its physiological consequences.

Introduction

The *Locus coeruleus* (LC) is a brain stem nucleus that is the primary source of norepinephrine (NE) in the central nervous system. It comprises a unique neuronal architecture with an extensive and diffuse axonal innervation throughout the entire brain and spinal cord, while the somatic core consisting of several hundred neurons is highly compact (Szabadi 2013). Research on the noradrenergic system has revealed astonishing insights on how a single neuromodulator such as NE can pointedly impact diverse brain functions ranging from arousal to pain sensation to memory encoding and sensory processing (Carter et al. 2010; Hirschberg et al. 2017; Takeuchi et al. 2016; Martins and Froemke 2015). In addition, much research is focused on understanding how expression profiles of adrenergic receptors in their respective axonal terminal fields are correlated to mental disorders such as attention deficit, clinical depression, or drug abuse; reviewed in (Poe et al. 2020). Distinct catecholaminergic receptor subtypes can be activated through a dynamic modulation of an extracellular NE gradient in the vicinity of a synaptic release site (Takeuchi et al. 2016; Zhang et al. 2013). In this respect, the type of NE release site in axonal terminals has been controversially discussed in the past; to one extrem a pure volume transmission was assume as main mode of action, and on the other

extrem a purely wiring-like transmission (Descarries, Watkins, and Lapierre 1977; Latsari et
al. 2002; Olschowka et al. 1981; Pickel et al. 1981; Nirenberg et al. 1996).
On the opposite, less research has focused on the synaptic wiring modes within the LC core
itself. Primarily, such research interest was hampered by the canonical view that LC activity is
mainly based on autonomous pacemaking of a homogenous noradrenergic cell population.
Yet, recent technological advancements in single-cell genetics, viral tracing, and optical
manipulation of neurons begin to challenge this view. Already a surprising heterogeneity in the
LC cell population regarding their projection-specificity and physiological function has been
described (Chandler, Gao, and Waterhouse 2014; Uematsu et al. 2017; Hirschberg et al.
2017). Nevertheless, we are still missing a framework on how individual LC neurons are
computing their brain-wide inputs to the nucleus and how this information is relayed within the
nucleus and thereby to the entire brain (Poe et al. 2020). As the somas in the nucleus are in
unusually close proximity to each other, volume-mediated interactions are perceivable.
Previous work on catecholaminergic neurons in the substantia nigra and ventral tegmental
area argue for such volume interaction between neurons as a key mechanism to fully describe
neuronal activities in these areas (Rice and Cragg 2008; Rice and Patel 2015; Cragg and Rice
2004). Early work in LC focused primarily on electronic coupling via gap junction as the main
mode of connection (Ishimatsu and Williams 1996). Gap junctions are most prominent in the
dendritic compartment and decrease in abundance with aging (Christie and Jelinek 1993;
Christie, Williams, and North 1989). They give rise to subthreshold oscillation that can
synchronize the entire nucleus during postnatal periods (Ishimatsu and Williams 1996). Yet,
recent high-density silicon probes recordings confirm that at older age neurons from different
subsets of the LC exhibit asynchronous activity (Totah, Logothetis, and Eschenko 2020). Here,
correlated activity patterns are observed within these subsets of LC neurons at different
timescales: from sub-milliseconds to milliseconds to periodic oscillation in the range of
seconds. Such a wide range of temporal interactions between LC neurons is beyond a pure
gap junction coupled neuronal network.
Indeed, several neuromodulators such as galanin, serotonin, neuropeptides and NE itself are
known to also impact LC network activity on the timescale of seconds and even minutes
(Tillage et al. 2020; Bai et al. 2018; Segal 1979; Aghajanian, Cedarbaum, and Wang 1977).
Electrophysiological studies clearly demonstrated a strong NE-mediated control of the LC
network via high-affinity alpha2-receptor (Singewald 1998; Singewald et al. 1994; Lakhlani et
al. 1997; Wagner-Altendorf, Fischer, and Roeper 2019). Infusion of antagonists leads to
hyperpolarization that is mediated through a NE-sensitive potassium conductance (Swanson
1976; Lee, Rosin, and Bockstaele 1998; Torrecilla et al. 2013). This decrease in excitability is
mediated via $G_{i/o}$ pathways that trigger the G-protein coupled inward rectifying potassium
channel GIRK2 (also called Kirk3) thereby controlling afferent activation of LC neurons as well
as the network state of the entire nucleus (Travagli, Dunwiddie, and Williams 1995; Travagli,
Wessendorf, and Williams 1996; Williams, North, and Tokimasa 1988; Aghajanian,
Cedarbaum, and Wang 1977; Lüscher and Slesinger 2010). So far, these questions have been
solely approached through a drug-based experimental design, where a drug effect is either
studied with conventional patch-clamp recordings on a single cell level or at the other extrem,
on the level of the entire LC network. As our view of the LC network shifts to a microcircuitry
level of functional specific subnetworks within the LC, spiking patterns of individual neurons in
their respective subnetworks become more critical for our understanding. Episodes of high
spiking rate (burst-like activity) in individual neurons can evoke elevated intracellular calcium
levels in soma that surpass the baseline level seen during tonic activity (Huang et al. 2012).
These high calcium levels can trigger the release of NE-containing vesicles from dendritic and

somatic compartments (Huang et al. 2007; 2012). It is challenging to evaluate how synaptic
and extracellular NE levels influence the LC network and neighboring neurons with the current
technologies. We, therefore, set out to employ a Hodgkin-Huxley-based model to study NE-
mediated local effects with the nucleus.
146

Methods

148
Acute brain slice preparation
All procedures were approved by the Landesverwaltungsamt / Halle, and conformed to
German regulation. All studies were performed in compliance to animal protocol 42502-2-1545
LIN. Acute brain slices for electrophysiology where obtained from Ai9 x DbH-Cre mice (Jax#
007905 and 033951) on a C57Bl6/J background. For tissue clearing mice from an Ai9 x PV-
Cre were used (Jax# 007905 and 17320).
Mice were injected intraperitoneally with pentobarbital (Narcoren®, Boehringer Ingelheim
Vetmedica GmbH, D-55216 Ingelheim, Germany,130 mg/kg, i.p.) and perfused with
carbonated (95% O_2, 5% CO_2) ice-cold slicing solution (in mM: 2.5 KCl, 11 glucose, 234
sucrose, 26 $NaHCO_3$, 1.25 NaH_2PO_4, 10 $MgSO_4$, 2 $CaCl_2$; 340 mOsm). After decapitation,
300-µm coronal brain stem slices were prepared in carbonated ice-cold slicing solution using
a vibratome (Leica VT 1200S) and allowed to recover for 20 min at 33 °C in carbonated high-
osmolarity artificial cerebrospinal fluid (high-Osm ACSF; in mM: 3.2 KCl, 11.8 glucose, 132
NaCl, 27.9 $NaHCO_3$, 1.34 NaH_2PO_4, 1.07 $MgCl_2$, 2.14 $CaCl_2$; 320 mOsm) followed by 40
min incubation at 33 °C in carbonated ACSF (in mM: 3 KCl, 11 glucose, 123 NaCl, 26
$NaHCO_3$, 1.25 NaH_2PO_4, 1 $MgCl_2$, 2 $CaCl_2$; 300 mOsm). Subsequently, slices were kept at
RT in carbonated ACSF until use. The recording chamber was perfused with carbonated ACSF
at a rate of 2 ml min^{-1} and maintained at 32 °C.

Brain slice electrophysiology

Whole-cell patch-clamp recordings were performed under visual control using oblique
illumination on a Slicecope Pro2000 (Scientifica) equipped with a 12-bit monochrome CMOS
camera (Hamamatsu Model OrcaFlash). Borosilicate glass pipettes (Sutter Instrument BF100-
58-10) with resistances ranging from 3–7 MΩ were pulled using a laser micropipette puller
(Sutter Instrument Model P-2000). Pipettes were filled using standard intracellular solution (in
mM: 135 potassium gluconate, 4 KCl, 2 NaCl, 10 HEPES, 4 EGTA, 4 Mg-ATP, 0.3 Na-GTP;
280 mOsm kg^{-1}; pH adjusted to 7.3 with KOH) Whole-cell voltage-clamp recordings were
performed using a MultiClamp 700B amplifier, filtered at 8 kHz and digitized at 20 kHz using a
Digidata 1550A digitizer (Molecular Devices).

177
HEK cells electrophysiology

Human embryonic kidney (HEK) cells stably expressing G-protein rectifying potassium
channels (GIRK 1/2 subunits), kindly provided by Dr. A. Tinker UCL London, GB, and Dr. S.
Herlitze were maintained at 37 °C in Dulbecco's modified Eagle's medium, 4.5 g/l D-glucose,
supplemented with 10% fetal bovine serum (Gibco) and penicillin/streptomycin incubated
under 5% CO_2. Cells were cultured in a 24-well plate, on 12 mm glass coverslips.

Whole-cell patch-clamp recordings were performed under visual control using a Slicecope
Pro2000 (Scientifica) equipped with a 12-bit monochrome CMOS camera (Hamamatsu Model
OrcaFlash). Borosilicate glass pipettes (Sutter Instrument BF100-58-10, 2-5 MΩ) were pulled

using a laser micropipette puller (Sutter Instrument Model P-2000). Pipettes were filled using
the internal solution as follows: 100 mM potassium aspartate, 40 mM KCl, 5 mM MgATP,
10mM HEPES-KOH, 5 mM NaCl, 2 mM EGTA, 2 mM MgCl2, 0.01 mM GTP, pH 7.3 (KOH).
The bath was filled using the following external solution: 20 mM NaCl, 120mM KCl, 2 mM
CaCl2,1mM MgCl2, 10 mM HEPES-KOH, pH 7.3 (KOH). Whole-cell voltage-clamp recordings
were performed using a MultiClamp 700B amplifier, filtered at 8 kHz and digitized at 20 kHz
using a Digidata 1550A digitizer (Molecular Devices). The following protocol was used during
the IV curve recordings: fifteen sweeps starting from -80 mV (Δ +10 mV). The cells were
clamped on a holding potential at -60 mV.

Immunohistochemistry
Mice were transcardially first perfused with phosphate-buffered saline (PBS 1x) and then with
4% paraformaldehyde. The brains were extracted and stored overnight in 4% PFA followed by
soaking in 30% sucrose to avoid forming water-crystals during slice sectioning at 4°C. Brains
were sliced on a microtome (LEICA SM2010R) to 40 μm thick coronal sections. Afterward,
brain slices were first permeabilized and blocked by incubating in PBST (Phosphate-buffered
saline with 0.3% Triton-X) containing 10% horse serum for 2 hours at room temperature to
prevent non-specific binding of antibodies. Sections were incubated in a solution containing
the primary antibodies, 2% horse serum, and 0.3% Triton-X for 24hrs at 4°C. The primary
antibodies used were guinea pig anti-TH (Synaptic Systems, Catalog No. 213104) with 1:500
dilution and rabbit anti-DBH (ImmunoStar, Catalog No. 22806) with 1:1000 dilution. Following
three washes in 1x PBS for 10 min the sections were then incubated in a solution containing
Cy3 anti-guinea pig IgG, (Jackson ImmunoResearch, Catalog No. 706-165-148), Cy5
AffiniPure Donkey Anti-Rabbit IgG, (Jackson ImmunoResearch, Catalog No. 711-175-152)
both in 1:200, 5% horse serum and 0.05% Triton-X for 2 hours at room temperature. The
sections were then washed thrice with 1xPBS for 10 min and then incubated in a solution
containing DAPI at a dilution of 1:30000. All the above-mentioned solutions were made in 1x
PBS. The sections were mounted on glass slides and coverslips containing DPX mounting
media. Images were acquired on a Leica SP8 confocal microscope using 63x oil-immersion
objectives (NA 1.4). Images were analyzed using ImageJ (https://imagej.nih.gov/ij/). For
images in figure 1, we overlay the reference atlas form *The Mouse Brain in Stereotaxic*
Coordinates 3rd Edition Franklin & Paxinos at anterior-posterior axis -5.40mm.

Tissue-clearing
Sectioning of 4% PFA fixated mouse brains were performed in a coronal configuration using
a brain mold to obtain 2-mm thick slices that contain the major part of the LC nucleus (AP 4.4
to 6.4mm) (brain matrix, 3D printed). Brain slices were then washed in PBS for 1h and
incubated at different series of mixtures prepared from tetrahydrofuran (THF, Sigma-Aldrich,
Catalog No.186562-1L):dH2O + 2% Tween-20 to dehydrate brain slices. The pH of the THF
solution was adjusted to 9.0 by adding triethylamine. The slices were washed in 50% THF
solution for 4h at 4°C followed by 70%, 90%, and 100% THF solutions under the same
conditions. To ensure complete dehydration, we washed the section with 100% THF twice.
Slices were then bleached in chilled 5% H_2O_2 in THF overnight at 4°C. The next day, the slices
were rehydrated in the reverse order by washing them in 100%, 90%, 70%, and 50% THF
solutions as above for 4h at 4°C. Following this, slices were washed twice in a PBS / 0.2%
TritonX for 1h each wash. Slices were then permeabilized in PBS / 0.2% TritonX / 20% DMSO
/ 0.3 M glycine at 4°C for 2 days and then blocked in PBS/0.2% TritonX / 10% DMSO / 6%
horse serum at 4°C for another 2 days. Samples were kept in primary antibody (guinea pig
anti-TH Synaptic Systems, Catalog No. 213104) dilution in PBS / 0.2% Tween-20 / 5% DMSO

/ 3% horse serum with 10 μg/ml heparin for two days at 4°C. Primary antibodies were diluted
at 1:400. Following this treatment, samples were washed in PBS / 0.2% Tween-20 with 10
μg/ml heparin for 1h at 4°C for 4x and then kept in secondary antibody (Cy3 anti-guinea pig
IgG, Jackson Immuno Research, Catalog No. 706-165-148) in 1:150 dilution in PBS / 0.2%
Tween-20 / 3% horse serum with 10 μg/ml heparin at 4°C for 2 days and washed in PBS /
0.2% Tween-20 with 10 μg/ml heparin for 1h at 4°C for 4x. For imaging, samples were
embedded in 0.8% phytagel and dehydrated in the THF/dH20 series exactly as described
before. After the dehydration, the samples were transferred to Ethyl Cinnamate (ECi) at room
temperature for clearing. ECi was replaced with fresh ECi the following day, and samples were
stored at room temperature.

Light-sheet microscopy
The light-sheet microscopy was performed using LaVision Biotech Ultramicroscope II (Miltenyi
Biotec, Bergisch-Gladbach, Germany) which was equipped with a Zyla 4.2 PLUS sCMOS
camera (Oxford Instruments, Abingdon-on-Thames, UK). The embedded brain slice was
placed in the sample chamber containing Ethyl Cinnamate (ECi). The brain volume was
sampled through a 4x objective (LaVision BioTec LVMI-Flour 4x/0.3). The objective was
configured to match the refractive index of ECi (1.558), for optimal imaging. For excitation, we
used light from a EXW-12 extreme laser (NKT Photonics, Birkerød, Denmark) which was
guided through the appropriate excitation filters (AHF Analysentechnik AG, Tübingen,
Germany) for Cy3 fluorescence. We started acquiring the image using ImSpector software
(Miltenyi Biotec, Bergisch-Gladbach, Germany) after setting the parameters such as sheet
width, z-step sizes, number of tiles and zoom. For post-processing and segmentation we used
Imaris software (version 9.72 Bitplane, Zurich, Switzerland).

Modeling
Based on cortical membrane kinetics, Alvarez et al. 2002
proposed a two-compartment mathematical model of LC cells, with soma and dendrite
connected via a gap junction on the dendrite. We employed this model as an initial starting
point and extended the model to incorporate quantal release of NE, which eventually leads to
the opening of GIRK channels and inhibition of the neighboring cell and/ or itself via alpha2-
receptor located in the somatic area. The release event is coupled with the amount of cytosolic
calcium [Ca^{2+}] in the cell.

Dynamic equation of state variables
268 Voltage change in the soma is driven by the balance of currents, including electrode current
269 I_{elec}, HH-sodium current I_{Na}, HH-potassium current I_K, persistent sodium current I_P, Ca^{2+}-
270 driven potassium current I_{AHP}, voltage-driven calcium current I_{Ca}, synaptic current from GIRK
271 conductance I_{GIRK}, current from dendrite I_{dend}, leak current I_L, and independent random
272 current η(t).

$$c_m \frac{dV_S^i}{dt} = I_{elec} - I_{Na} - I_K - I_P - I_{AHP} - I_{Ca} - I_{GIRK} - I_{dend} - I_L + \eta(t)$$

$$\frac{\mu F}{area} \frac{mV}{ms} = \frac{\mu A}{area} = \frac{mS}{area} mV, \qquad area\ [cm^2]$$

(eq. 1)

280 Voltage change in the dendrites are driven by current from soma I_{soma}, gap junctions from other
281 dendrites I_{GJ}, and dendritic leak current I_{LD}

$$c_m \frac{dV_D^i}{dt} = -I_{soma} - I_{GJ} - I_{LD} \qquad \text{(eq. 2)}$$

$$I_{AHP} = g_{AHP}\, r\, (V - V_K); \qquad \text{(eq. 3)}$$

290
291 The Ca^{2+} concentration changes over time as a function of voltage.
292

$$r = \frac{[Ca]}{[Ca] + 1\mu M} \qquad \text{(eq. 4)}$$

298
299 Specifically, membrane depolarization during a spike allows Ca^{2+} to enter the neuron. As
300 described above, the opening and closing of the g_{AHP} potassium conductance shadow the
301 cytosolic Ca^{2+} concentration.
302
303 The steady-state of the calcium is

$$[Ca]_\infty = f_{in}(V)\,\tau_{Ca} = \frac{f_{Ca}\,k_{Ca}\,\tau_{Ca}\,(V_{Ca} - V)}{1 + exp(-(V + 25)/2.5))}, \qquad k_{ca} = 2,\; f_{Ca} = 0.002\,\frac{\mu M}{mV\,ms} \qquad \text{(eq. 5)}$$

308 And the update rule is:

$$[Ca]_\infty = \frac{M_{Ca}}{132.6\,mV} \frac{V_{Ca} - V}{1 + exp(-(V + 25)/2.5))}$$

309 $\qquad \text{(eq. 6.1)}$

$$[Ca]_{i+1} = [Ca]_\infty + ([Ca]_i - [Ca]_\infty) \exp\left(-\frac{dt}{\tau_{Ca}}\right)$$

310 $\qquad \text{(eq. 6.2)}$
311

312 As the voltage-dependent factor in the expression of $[Ca]_\infty$ peaks at 132.6mV for $V_{Ca} =$
313 120mV , it is convenient to define a new parameter, M_{ca} = 42.4 µM and τ_{Ca} = 80ms
314 (supplemental figure 1)

Dendro-somatic and soma-somatic NE release and alpha2-receptor activation, and GIRK conductance

Action potentials in the presynaptic neuron results in NE release of dense core vesicles from presynaptic sites, which are found at the soma and at nearby dendrites. The release fraction R_{rel} depends on the level of presynaptic calcium at the time of the presynaptic action potential.

$$R_{rel}\left([Ca^j_{AP}]\right) = \frac{2 * [Ca^j_{AP}]^k}{[Ca]^k_{high} + [Ca^j_{AP}]^k},$$

(eq. 7)

$[Ca^j_{AP}]$ is presynaptic cytosolic calcium at the time of the action potential. $[Ca]_{high}$=1.3 µM is the amount of cytosolic calcium accumulated when the cell fires at 20 Hz. The exponent, k=1, modulates the release.

Once released, NE binds to alpha-2 receptors in the postsynaptic membrane, which through G-protein signaling opens an inwardly-rectifying potassium channel (GIRK: G-protein-coupled inward-rectifying potassium channel). The effective time course of the opening and closing of this synaptic conductance is comparatively slow (Williams, Bobker, and Harris 1991). We approximate this time-course with a double-exponential function with τ_{fast}= 300 ms and τ_{slow}= 350 ms as well as measure the voltage-dependence of GIRK current in a HEK cells line that is stably expressing GIRK1/2 subunit channel (supplemental figure 2) (Williams, Bobker, and Harris 1991; Eickelbeck et al. 2020).

$$\begin{aligned}
P_{open}(t) &= B\left[f^j_{slow}(t) - f^j_{fast}(t)\right] \\
f^j_{fast}(t) &= \exp\left(-\frac{t-t^j_{AP}}{\tau_{fast}}\right) \\
f^j_{slow}(t) &= \exp\left(-\frac{t-t^j_{AP}}{\tau_{slow}}\right)
\end{aligned}$$

(eqs. 8)

$$t_{peak} = \frac{\tau_{fast}\tau_{slow}}{\tau_{slow} - \tau_{fast}} \log\frac{\tau_{slow}}{\tau_{fast}}, \qquad B = \frac{1}{e^{-t_{peak}/\tau_{slow}} - e^{-t_{peak}/\tau_{fast}}}$$

$$B = \left[\left(\frac{\tau_{fast}}{\tau_{slow}}\right)^{\left(\frac{\tau_{fast}}{\tau_{slow}-\tau_{fast}}\right)} - \left(\frac{\tau_{fast}}{\tau_{slow}}\right)^{\left(\frac{\tau_{slow}}{\tau_{slow}-\tau_{fast}}\right)}\right]^{-1}$$

The absolute value G_{rect} of this rectifying conductance is not constant, but changes with the postsynaptic membrane potential. Accordingly, we need to consider this factor in the equation.

$$G_{rect}(V^i) = \frac{0.2}{1 + \exp\left(0.05(V^i - V_{GIRK})\right)}$$

(eq. 9)

372 G_{rect} is given in μS per mm^2.

373 Combining all three factors for multiple successive presynaptic action potentials AP_1, AP_2,...,

374 the GIRK conductance is given then by:

$$g^{ij}_{GIRK}\left(V^i, t, [Ca^j_{AP1}], [Ca^j_{AP2}], \ldots\right) = G_{rect}(V^i)\left[P_{open}(t - t_{AP1})\, R_{rel}\left([Ca^j_{AP1}]\right) + \right.$$
$$+ \; P_{open}(t - t_{AP2})\, R_{rel}\left([Ca^j_{AP2}]\right) + $$
$$\left. + \; \ldots\right]$$

(eq. 10)

381 Release from or near soma and binding near soma. This hyperpolarizes the membrane

382 potential of the neighboring soma. S_{ij} is the strength of the synaptic connection from neuron j

383 to neuron i.

$$I^i_{GIRK} = \left(V^i_S - V_{GIRK}\right) \sum_j S_{ij}\, g^{ij}_{GIRK}\left(V^i_S, t, [Ca^j_{AP1}], [Ca^j_{AP2}], \ldots\right)$$
$$= \left(V^i_S - V_{GIRK}\right) G_{rect}(V^i) \times$$
$$\times \sum_j S_{ij}\left[P_{open}(t - t_{AP1})\, R_{rel}\left([Ca^j_{AP1}]\right) + P_{open}(t - t_{AP2})\, R_{rel}\left([Ca^j_{AP2}]\right) + \ldots\right]$$

(eq. 11)

392 **NE diffusion**

394 To estimate the effect of NE released into the extracellular space, we computed the time-

395 course of NE concentration at some distance from a release site as by:

397 $$C_{NE}(r, t) = \frac{Exp\left(-\frac{r^2}{4\,D\,t} - k\,t\right)}{\alpha(4\,D\,\pi\,t)^{3/2}}$$

(eq. 12)

399 where r is distance, t is time, D=3.4e-6 cm^2/s is the diffusion constant, and k = 20 Hz is the

400 reuptake constant (Rice and Nicholson 1995). The fraction of NE-binding alpha2-receptors

401 was computed from

403 $$\tau_{a2R}\frac{dx}{dt} = -x + x_{a2R}, \quad \tau_{a2R} = \frac{1}{200\,nM\,k_{on} + k_{off}}, \quad x_{a2R} = \frac{200\,nM\,k_{on}}{200\,nM\,k_{on} + k_{off}},$$

(eq. 13)

where k_{on} = 1.0 and k_{off} = 1.25 nM are the kinetic constants for NE binding and unbinding.

The resulting time course and voltage dependence of GIRK conductance was fitted with a

double-exponential function with τ_{fast}= 1400 ms and τ_{slow}= 1450 ms and an attenuation factor

of 0.07 (supplemental figure 2).

To take into account the kinetics of transmitter binding and release, the resulting time-course

was convolved with a low-pass filter ('alpha function') $g(t) = \frac{t}{\tau}exp(-t/\tau)$ with characteristic

time $\tau = 50ms$.

Data analysis

All data were analyzed in Python. For extraction of action potential features (simulated and experimental), we used the Allen Institute ephys package. The generalized leaky integrate and fire models (GLIF) were applied. Additional analyses were performed using custom Python scripts, all of which are available on the laboratory GitHub account.

Results

We first set out to confirm that LC neurons express both key enzymes to catalyze NE in somatic and dendritic compartments throughout their lifetime. Immunostaining against tyrosine hydroxylase (TH), as well as dopamine-beta-hydroxylase (DbH) in young mice (P14) and old mice (older than P500), revealed expression of both enzymes in all cells at both ages (figure 1 a, b). Both stainings also highlight the compact nature of the LC nucleus itself. Furthermore, a dense dendritic arbor directed towards the medial-rostral axis becomes visible (Ishimatsu and Williams 1996). Image overlay of the fluorescence channel from TH and DbH confirms an overall co-expression of both enzymes throughout the somatic compartment within the LC nucleus (see figure 1 a and b overlay). Surprisingly, we observed a differential spatial expression levels of TH and DbH in the dendritic arbor. Here, both poles of the LC (dorsal and ventral) appear to have protruding dendrites that mainly display a TH immunosignal. The central part of the nucleus seems to extend dendrites that preferentially express DbH (figure 1 c, 3 animals). The dendritic expression profile for the DbH enzymes was also similar for young and old mice along the anterior-posterior axis (supplemental figure 3)

As it is difficult to experimentally approach somatic or dendritic NE release from a single LC neuron and investigate how it affects spike activity in neighboring LC neurons, we decided to develop a computational model that allows us to study three modes of NE-mediated interactions. The first mode was via dendro-somatic synapse (Nirenberg et al. 1996); secondly, we considered a somatic autoinhibition (Andrade and Aghajanian 1984; Wagner-Altendorf, Fischer, and Roeper 2019), and thirdly we investigated if a NE volume diffusion from a somatic release point can lead to an inhibition of a neighboring neuron.

Figure 1. *Tyrosine hydroxylase and Dopamine-beta-Hydroxylase expression largely overlap in somatic and dendritic compartments in the Locus coeruleus.*

(**a**) Shows immunostaining against key enzymes in the synthesis pathway of NE. White dotted lines show an overlay mask of the corresponding section from The Mouse Brain in Stereotaxic Coordinates 3rd Edition Franklin & Paxinos (**a**) Confocal image of

immunofluorescence against tyrosine hydroxylase (TH / *green*) from a brain stem slice of a young P14 mouse (anterior-posterior axis: -5.40 mm). The fluorescence signal is present in the soma as well as in the dendritic arbor (here more dominantly in the medial part); We observed a similar spatial distribution for the fluorescence signal from immunostainings against the dopamine-beta-hydroxylase (*magenta*). **(b)** depict the same immunostainings as in (a) on brain slices obtained from an old P500 mice. A similar spatial expression profile for both enzymes is visible when compared to slices from a young P14 mouse (a). **(c)** Shows a quantification of average pixel (px) intensities from the *yellow-colored ROIs* displayed in TH/DbH overlay image in a and b. ROIs are placed above the somatic or dendritic compartments, respectively. **(c)** Shows average intensity of pixel values from ROIs along the dorsal-ventral axis for soma or dendrites. Scale bar 100µm

To realistically model the distribution of the distances between LC neurons, we employed a tissue clearing approach to map the positions of noradrenergic neurons in the LC. First, we prepared 2-mm thick brain stem slices from PV-Cre x Ai9 animals. We then used a modified iDISCO protocol to clear the entire brain stem slices and image the slice volume via fluorescence light sheet microscopy (see methods for details). After acquiring images of the entire brain stem, we visualize anti-TH+ and PV-dTomato neurons with high contrast (figure 2 a). We segmented the LC and resolved positions of 220 to 439 anti-TH+ neurons (4 nuclei, 2 animals) (figure 2 b). Based on the x-y-z coordinates of individual anti-TH+ neurons, we compute the nearest-neighbor (NN) distances for each cell in all four analyzed nuclei. The distribution of the NN-distances peaks at distances of beta-hydroxylase around 25µm (median 41.34µm) (see figure 2 c).

To investigate the possible effects of a noradrenergic interaction between LC neurons on a single cell level, we choose as a starting point a previously published mathematical model that describes the electronic coupling between two LC neurons via gap junctions (Alvarez et al. 2002). This model recapitulates frequency-dependent synchronization between LC neurons as an electrotonic coupling strength between dendrites. Here, strong gap-junction coupling between dendritic compartments, as described experimentally in young mice (Christie and Jelinek 1993; Christie, Williams, and North 1989; Ishimatsu and Williams 1996), lead to synchronization of spontaneous spiking for all frequencies; while weak dendritic coupling promotes spike synchrony only at higher spiking frequencies. Comparison of our initial model under a weak electrotonic coupling regime reiterates our current-clamp recordings from noradrenergic neurons from acute brain slices prepared from mice older than P70. The spontaneous spike rate at resting potential was 4.2±1.77Hz (n = 6 neurons, 2 animals) while our model estimates 4.3 ± 0.03 Hz (figure 3 a). Overall features of action potentials between our model and experimental data largely coincide (figure 3 b). To emulate different excitability levels of LC neurons, we recorded the spike rate during a three hundred millisecond long current pulse at various amplitudes (figure 3 c, *upper trace*). Also, in our model, current steps evoked an increased spike rate (figure 3 c, *lower trace*). As expected, spike rates increased with amplitudes of injected current. Quantification of spike rate at various current step amplitudes revealed an overall linear dependence in the frequency-current relation (FI-curve) for experimental and simulated data (compare figure 3 d1 and d2). Interestingly, we observed a higher spike rate for the two initial spikes and an overall steeper slope in the FI-curve for these early spikes compared to the following spikes (supplemental figure 3 a). We also found such an increased spike rate for the initial spikes in our model; this effect gets more pronounced at higher current amplitudes for experimental and simulated data (compare supplemental figures 4 a and b).

To simulate a noradrenergic modulation between LC neurons, we extended our model with a calcium-dependent release of NE. We considered a NE release site as a point source of NE secretion that either mimics a dendro-somatic synapse or a somatic release site that can either retroact on the secreting neuron (self-inhibition) or effect neighboring neurons via diffusion (neighboring inhibition). High spiking rates lead to membrane fusion of NE-containing large-dense vesicles at the soma (Huang et al. 2007; 2012). The voltage-dependent rise in calcium is modeled by equations 3, 4, 5 and 6 (see Methods). Based on amperometric recordings in acute brain slices, we assumed an initial extracellular NE concentration after vesicle fusion of 200nM (Huang et al. 2012). We combined the time courses of several molecular events in equation 10 i.e. binding of NE to alpha2-receptors, initiation of $G_{i/o}$ signaling cascade, and the subsequent release of $G_{\beta\gamma}$ dimer that increases GIRK channel conductance. The dependence of GIRK conductance and intracellular calcium levels is shown in figure 4a. For a given calcium level (dotted line in figure 4a) the temporal development of GIRK open probability (P_{open}) is modeled based on electrophysiological characterization of GIRK current in LC neurons (supplemental figure 2) (Williams, Bobker, and Harris 1991) .

Figure 3. *LC neuron model recapitulates essential spontaneous spiking, action potential characteristics and current-evoked frequency dependency.*

(a) Spontaneous neuronal spiking in an LC neuron recorded in an acute brain slice from P70 mice via whole-cell patch-clamp (*upper black trace*). Action potentials show typical long after-hyperpolarization (AHP) that give rise to a 4.2 Hz spontaneous spiking rate. Trace below shows simulated LC neuron that exhibit similar spontaneous spiking rate and slow AHP. **(b)** Overlay of experimental and simulated action potentials exhibit a similar temporal profile. **(c)** The current-evoked increase in membrane potential evokes a barrage of action potentials in a recorded and simulated neuron (300ms current injections). **(d1)** Illustrates the spike rate of recorded LC neurons under different current step amplitudes (error margin as SEM; n = 12 cells, 2 animals); **(d2)** Simulated LC neurons exhibit an increase in spike rate for different amounts of injected current. Note the different current units in d1 and d2 due to the different cell sizes of experimental and simulated neurons. Scale bars 10mV for experimental neurons and 20mV for simulated neurons.

Based on these assumptions, we first tested our model for a synaptic-like dendro-somatic interaction between two LC neurons. To mimic episodes of high activity, we injected current amplitudes for 1-sec that lead to spike rates close to 20Hz in neuron 1 (N1) (figure 4 c1). As expected, an increased spike rate led to an uprise of intracellular calcium levels (figure 4 c1 *magenta trace* and supplemental figure 5 a,b). As described in equation 5, this increase translates into a delayed, slowly developing onset of GIRK conductance equation 10, (figure 4 b, c1 *blue trace*). The generated GIRK conductance decreases with Tau = 350ms after the current injection step ends. In our model, the increased GIRK conductance during and after episodes of high spike activity led to complete inhibition of spontaneous spiking in the dendro-somatically connected neuron 2 (N2). Only after GIRK conductance returns to basal levels, N2 resumes spontaneous spiking.

In a different scenario, we investigated how the somatic release of NE impacts the neuronal activity of the releasing neuron (N1) itself. Similar to our previous constraints, we induced phasic-like spiking through a current-step injection in our modeled neuron. The high spike rate induces an increase of intracellular calcium followed by a delayed onset GIRK conductance (figure 4 c2). Also, here, we assumed quantal release concentration as in the previous scenario, i.e., an initial extracellular NE concentration of 200nM. Despite the slowly rising of GIRK conductance in N1, the calcium concentration in N1 only shows a subtle decline in the average intracellular calcium levels during these current-induced bouts of high neuronal activity (*compare 4 c magenta traces upper versus lower trace*). The spike rate in N1 remains stable despite rising potassium leak through GIRK channels. Yet, we observed that after the current step subsided, the slow-decaying GIRK conductance impede spontaneous firing in N1; 500ms after the current pulse ended, spontaneous spiking recurred. Such an observation agrees with the typical quiescent period of LC neurons after episodes of high activity (Williams et al. 1984).

Figure 4. *Simulation of spike rate from two LC neurons with a built-in NE-dependent GIRK activity leads to a reduced spiking rate in LC neurons that are connected via dendro-somatic synapses or a somatic release that retroact onto the secreting LC neuron.*

(a) R_{rel} is the release fraction that is dependent on the level of presynaptic calcium at the time of action potential. The gray vertical line is the amount of calcium accumulated in a cell when firing at 20Hz. The relation is defined by equation 7. **(b)** P_{open} shows the effective time course of the opening and closing of the synaptic conductance of GIRK channel.

(equation 8 $P_{open}(t)$). **(c1)** Depicts a simulation scenario where neuron 1 (N1) connects to neuron 2 (N2) via dendro-somatic synaptic interaction. *The upper diagram* shows the spike rates of N1 and N2 throughout the simulated 5-sec. Here, the *gray line* represents the time course of current injections (1s, I= 270nA) in the respective neuron. *The lower diagram* shows the development of intracellular calcium levels in N1 (*magenta*) during a phasic-like activation; the *blue line* represents the time course of GIRK conductance in N2. **(c2)** Cartoon illustrates the autoinhibition scenario in which alpha2-adrenergic autoreceptors in N1 activate through local NE-release from the soma. One-second bouts of current injections (I=270nA) lead to an increased spike rate in N1, while N2 displays a spontaneous firing pattern (*upper diagram*). *The lower diagram* shows time courses of intracellular calcium concentration and GIRK conductance in N1.

We then began to examine if somatic NE release and the spatial confinements of the LC architecture can give rise to volume-mediated interaction between LC neurons. Here, we hypothesize that the LC neurons during episodes of high activity can modulate spontaneous spiking of neighboring neurons via bystander-inhibition. The spatiotemporal NE concentration profile originating from a point source can be estimated through equation 12 (Cragg and Rice 2004; Rice and Cragg 2008).

The attenuation factor (C_0 / $C(r,t)$) relative to the initial NE concentration shows a spatiotemporal profile with an attenuation of 0.001 for distances of around 20 to 25μm after 200 to 800ms (figure 5 a). The fraction of NE bound alpha2-receptors (K_D = 1.25nM) is shown as a temporal profile for various distances in figure 5 b. The onset and fractions of activated alpha2-receptor are slowed down and decreased through the time course of diffusion, respectively. In our neural model of LC neurons, we aim to estimate the neighbor inhibition of freely diffusing NE onto a neuron (N3) at a distance of 25μm. Evoking a phasic-like activity in neuron N1 through repeated current-step injections evoked an uprise of intracellular calcium level that surpasses the threshold for NE vesicle release. The onset of a slow-rising GIRK conductance in N3 is delayed by 1400 ms compared to previous scenarios (dendro-somatic and autoinhibition) (compare figure 4 c1, c2 versus figure 5 c). The decelerated rise of GIRK conductance is due to the small number of active alpha2-receptors. This translates into a slow and small increase of GIRK conductance in N3. As GIRK activation was delayed, a second current step two seconds after the first one ended, continued to build up GIRK conductance on the already existing conductance from the previous high activity bout (figure 5 c). Shorter intervals between the current step further facilitates this effect. This cumulated increase in GIRK conductance in N3 leads to an enlarged interspike interval between spontaneous spikes in N3 (figure 5 c and supplemental figure 6 a).

Figure 5. *Spike modulation in the neighboring neuron of somatically release NE cumulatively affects neighboring neurons within a 25μm radius.* **(a)** Spatiotemporal diffusion profiles of NE from a point source. The diffusion constant for NE was 3.4×10^{-6} cm^2 s^{-1} (Rice and Cragg 2008; Cragg and Rice 2004). **(b)** Time course of fractions of NE bound alpha2-receptor at 1, 5 and 20μm distance from release site. Kinetics of transmitter binding and release was modeled by an alpha function (think black line, τ=50 ms). **(c)** Simulation of spike train of current injected neurons (N1) and nearby neurons (N3 / 25μm distance). The lower plot shows the development of intracellular calcium levels in N1, while the blue line represents GIRK conductance in N3.

Discussion

Noradrenergic modulation exerts robust control over neuronal circuits throughout the entire brain. The densely packed core of the LC integrates various inputs from cortical, subcortical, and brain stem areas. These inputs are computed within the nucleus and evoke various network states in the LC. Early work highlights the importance of electronic coupling between noradrenergic neurons that can synchronize sub-threshold membrane fluctuation and spiking rates in the entire LC network (Christie and Jelinek 1993; Christie, Williams, and North 1989; Ishimatsu and Williams 1996; Alvarez et al. 2002). While later research points to multiple asynchronous neuronal ensembles that dynamically switch spiking patterns within the LC (Uematsu et al. 2017; Hirschberg et al. 2017; Totah et al. 2018). As the specific neuronal architecture and synaptic wiring onto and between LC neurons are unknown, the space of potential network states and their respective output function remains elusive. Recently, the identification of local inhibitory neurons that control the phasic activity of specific neuronal subsets in the LC adds another layer of computational capacity to this brain stem nucleus (Breton-Provencher and Sur 2019; Jin et al. 2016; Kuo et al. 2020). Yet, to achieve a deeper understanding of the functional role of noradrenergic subnetworks, we need experimental approaches that target these ensembles.

Here, we developed an experimentally validated neuronal model of noradrenergic cells that allowed us to investigate NE-mediated modulatory effects between LC neurons. First, our immunostaining against DbH and TH enzymes confirms the expression of both key enzymes for synthesize of NE in the vast majority of LC neurons in young and old mice (figure 1). As recent studies demonstrated that noradrenergic axons in the dorsal hippocampus predominantly release dopamine (Kempadoo et al. 2016; Takeuchi et al. 2016), our results demonstrate that in somatic and dendritic compartments the NE can be produced in all LC neurons. Still, such partition between dopamine and NE release site could appear on the level of axonal varicosities; either through targeting of DbH enzymes to distinct projection areas or a differential expression of monoamine transporters that bias uptake of a specific catecholamine from the extracellular space.

Our reconstruction of LC geometry reproduces noradrenergic cell numbers that are five times lower than earlier histological studies (Sturrock and Rao 1985;O'Neil et al. 2007). As previous studies use stereological cell counts of 6 to 50µm brain slices, overestimation through double counting of neurons is possible. Yet, it is likely that these large differences are also due to insufficient penetration of TH antibodies in our cleared tissue preparation that limit cell count compared to classical histology. Nevertheless, a possible underestimation of the cell count in our LC preparation would only affect the overall fraction of LC neurons that are in a 25µm-radius, but the resolved absolute number of LC neuron pairs would stay constant or be larger. In our diffusion model, we have not incorporated noradrenaline transporters activity (NET) that can limit the diffusion radius of NE. Despite studies showing the presence of transporters in the LC core (Klimek et al. 1997; Isaias et al. 2011), previous diffusion modeling studies for the Substantia nigra argue that the impact of monoamine transporter on the diffusion radius of dopamine is small (Cragg and Rice 2004; Rice and Cragg 2008). Direct electrophysiological recordings in the LC neurons confirm that increasing electrical evoked-release of NE scales with the rate of alpha2-receptor evoked inhibitory potentials in noradrenergic neurons; arguing that monoamine transporters are not drawing excessive NE out of the extracellular space (Courtney and Ford 2014). In contrast, in similar dose measurements in the ventral tegmental area, D1-receptor evoked inhibitory responses that did not show dependence on stimulation

strength. Only after treatment with a DAT antagonist responses are scaling with stimulation
strength (Courtney and Ford 2014). These experimental findings argue that NET activity within
the LC is not limiting the diffusion of NE.
The observed reduction in spontaneous spike rate in our model through the diffusion of
extracellular NE has so far not been described to our knowledge; it has been discussed as a
plausible mechanism for dopaminergic neurons in the substantia nigra as well as dopaminergic
axons (Cragg and Rice 2004; Rice and Cragg 2008; Walters et al. 2014). Key factors for the
observed bystander-inhibition are distances and geometrical orientation of the release site
between LC neurons. Our model deduces that extracellular NE can accumulate through
episodes of long-lasting phasic activity or repeated periods of bursting discharge. This
accumulation is dependent on the speed of G_i signaling cascade, GIRK channel kinetic and
dispersion of extracellular NE through the densely packed LC core. Such bystander-inhibition
adds another layer of interaction between LC neurons. The physiological role of such type of
inhibition is strongly interwoven with a topographical organization of the LC.
The idea of topographic organization of the LC has been put forward decades ago; Most
studies at that time used local injection of retrograde dyes (Waterhouse et al. 1993; Loughlin,
Foote, and Bloom 1986; Loughlin, Foote, and Fallon 1982). Despite controversial findings, it
is generally accepted that the LC has a dorsal-ventral organization in which neurons in the
dorsal part of the LC are preferential projecting to frontal areas while the ventral part projects
to brain stem areas. For an example, LC neurons that project to hippocampal areas or olfactory
bulb are more likely found in the dorsal part, while ventral LC neurons are more probable to
project to medulla or spinal cord (Mason and Fibiger 1979; Loughlin, Foote, and Bloom 1986;
Schwarz et al. 2015). Based on new viral strategies and high-throughput axonal mapping
technologies, a better understanding of the brain-wide input-output organization of the LC
became available. These findings helped to develop new concepts that now allow also to
understand the controversial findings from earlier tracing studies. For example, one of these
concepts argues that axonal projections from a single LC neuron forms an extensive axonal
field to a specific brain area, while also

possessing collaterals to other brain areas at lower density. A variation of such a concept entails the distinction between broadcasting LC neurons that form equally dense innervation to many areas, versus exclusive LC neurons that form specific innervation to a single brain area. Both concepts would blur retrograde tracing experiments, and underestimate a topographical organization of the LC. Interestingly, functional studies highlight heterogeneity of projection-defined LC neurons in terms of intrinsic excitability, after-hyperpolarization and their proteomic make-up (Wagner-Altendorf, Fischer, and Roeper 2019; Chandler, Gao, and Waterhouse 2014). Here, LC neurons that preferentially project to the prefrontal cortex or hippocampus show a different sensitivity to alpha2-receptor agonists. Therefore, a topographic organization and differential expression of adrenergic receptors could fine-tune the impact of neigboring inhibition. It is interesting to speculate how the here postulated volume-mediated interaction of spatially clustered LC neurons with different projection targets could amplify differences in extracellular NE in their respective axonal fields. The increase in contrast of NE levels in two target regions could increase sensory saliency in one region.

Figure 6: *Illustration of functional consequences of a volume-mediated inhibition of neighboring neurons with different projection targets.*
Strong enduring input to neuron 3, N3, leads to an increase in neuronal spiking (*arrow up*) and subsequently increase release of NE within the core of the LC. Based on distances and receptor portfolio of neuron 1 (N1) from the release sites of N3, the spontaneous spiking activity in N1 will be decrease through activation of GIRK (arrow down); while for example activity in neuron 2 (N2) is not altered (arrow horizontal). Consequently, axonal terminals from N3 to projection-target area A increase spike-dependent release of NE, while axons in area B from N1 decrease NE release due to decrease spiking in N1. It is conceivable that the volume-mediated local NE inhibition in the core of the LC, leads to contrasting functional effects in the neuronal circuits in region A and B i.e. facilitating sensory salicieny in region A versus region B.

References

Aghajanian, G.K., J.M. Cedarbaum, and R.Y. Wang. 1977. "Evidence for Norepinephrine-Mediated Collateral Inhibition of Locus Coeruleus Neurons." *Brain Research* 136 (3): 570–77. https://doi.org/10.1016/0006-8993(77)90083-X.

Alvarez, Veronica A., Carson C. Chow, Elisabeth J. Van Bockstaele, and John T. Williams. 2002. "Frequency-Dependent Synchrony in Locus Ceruleus: Role of Electrotonic Coupling." *Proceedings of the National Academy of Sciences* 99 (6): 4032–36. https://doi.org/10.1073/pnas.062716299.

Andrade, Rodrigo, and George K. Aghajanian. 1984. "Intrinsic Regulation of Locus Coeruleus Neurons: Electrophysiological Evidence Indicating a Predominant Role for Autoinhibition." *Brain Research* 310 (2): 401–6. https://doi.org/10.1016/0006-8993(84)90170-7.

Bai, Yun-Fei, Hai-Tao Ma, Li-Na Liu, Hui Li, Xiao-Xiao Li, Yu-Tao Yang, Bing Xue, Di Wang, and Zhi-Qing David Xu. 2018. "Activation of Galanin Receptor 1 Inhibits Locus Coeruleus Neurons via GIRK Channels." *Biochemical and Biophysical Research Communications* 503 (1): 79–85. https://doi.org/10.1016/j.bbrc.2018.05.181.

Breton-Provencher, Vincent, and Mriganka Sur. 2019. "Active Control of Arousal by a Locus Coeruleus GABAergic Circuit." *Nature Neuroscience* 22 (2): 218–28. https://doi.org/10.1038/s41593-018-0305-z.

Carter, Matthew E, Ofer Yizhar, Sachiko Chikahisa, Hieu Nguyen, Antoine Adamantidis, Seiji Nishino, Karl Deisseroth, and Luis de Lecea. 2010. "Tuning Arousal with Optogenetic Modulation of Locus Coeruleus Neurons." *Nature Neuroscience* 13 (12): 1526–33. https://doi.org/10.1038/nn.2682.

Chandler, D. J., W.-J. Gao, and B. D. Waterhouse. 2014. "Heterogeneous Organization of the Locus Coeruleus Projections to Prefrontal and Motor Cortices." *Proceedings of the National Academy of Sciences* 111 (18): 6816–21. https://doi.org/10.1073/pnas.1320827111.

Christie, M.J., and H.F. Jelinek. 1993. "Dye-Coupling among Neurons of the Rat Locus Coeruleus during Postnatal Development." *Neuroscience* 56 (1): 129–37. https://doi.org/10.1016/0306-4522(93)90568-Z.

Christie, M.J., J.T. Williams, and Ra North. 1989. "Electrical Coupling Synchronizes Subthreshold Activity in Locus Coeruleus Neurons in Vitro from Neonatal Rats." *The Journal of Neuroscience* 9 (10): 3584–89. https://doi.org/10.1523/JNEUROSCI.09-10-03584.1989.

Courtney, N. A., and C. P. Ford. 2014. "The Timing of Dopamine- and Noradrenaline-Mediated Transmission Reflects Underlying Differences in the Extent of Spillover and Pooling." *Journal of Neuroscience* 34 (22): 7645–56. https://doi.org/10.1523/JNEUROSCI.0166-14.2014.

Cragg, Stephanie J., and Margaret E. Rice. 2004. "DAncing Past the DAT at a DA Synapse." *Trends in Neurosciences* 27 (5): 270–77. https://doi.org/10.1016/j.tins.2004.03.011.

Descarries, Laurent, Kenneth C. Watkins, and Yves Lapierre. 1977. "Noradrenergic Axon Terminals in the Cerebral Cortex of Rat. III. Topometric Ultrastructural Analysis." *Brain Research* 133 (2): 197–222. https://doi.org/10.1016/0006-8993(77)90759-4.

Eickelbeck, Dennis, Till Rudack, Stefan Alexander Tennigkeit, Tatjana Surdin, Raziye Karapinar, Jan-Claudius Schwitalla, Brix Mücher, et al. 2020. "Lamprey Parapinopsin ('UVLamP'): A Bistable UV-Sensitive Optogenetic Switch for Ultrafast Control of GPCR Pathways." *ChemBioChem* 21 (5): 612–17. https://doi.org/10.1002/cbic.201900485.

Hirschberg, Stefan, Yong Li, Andrew Randall, Eric J Kremer, and Anthony E Pickering. 2017. "Functional Dichotomy in Spinal- vs Prefrontal-Projecting Locus Coeruleus Modules Splits Descending Noradrenergic Analgesia from Ascending Aversion and Anxiety in Rats." *ELife* 6 (October). https://doi.org/10.7554/eLife.29808.

Huang, H.P., S.-R. Wang, W. Yao, C. Zhang, Y. Zhou, X.-W. Chen, B. Zhang, et al. 2007.

845   "Long Latency of Evoked Quantal Transmitter Release from Somata of Locus
846    Coeruleus Neurons in Rat Pontine Slices." *Proceedings of the National Academy of*
847    *Sciences* 104 (4): 1401–6. https://doi.org/10.1073/pnas.0608897104.
848 Huang, H.P., Fei-Peng Zhu, Xiao-Wei Chen, Zhi-Qing David Xu, Claire Xi Zhang, and Zhuan
849    Zhou. 2012. "Physiology of Quantal Norepinephrine Release from Somatodendritic
850    Sites of Neurons in Locus Coeruleus." *Frontiers in Molecular Neuroscience* 5.
851    https://doi.org/10.3389/fnmol.2012.00029.
852 Isaias, Ioannis U, Giorgio Marotta, Gianni Pezzoli, Osama Sabri, Johannes Schwarz, Paolo
853    Crenna, Joseph Classen, and Paolo Cavallari. 2011. "Enhanced Catecholamine
854    Transporter Binding in the Locus Coeruleus of Patients with Early Parkinson
855    Disease." *BMC Neurology* 11 (1): 88. https://doi.org/10.1186/1471-2377-11-88.
856 Ishimatsu, Masaru, and J.T. Williams. 1996. "Synchronous Activity in Locus Coeruleus
857    Results from Dendritic Interactions in Pericoerulear Regions." *The Journal of*
858    *Neuroscience* 16 (16): 5196–5204. https://doi.org/10.1523/JNEUROSCI.16-16-
859    05196.1996.
860 Jin, Xin, Shanshan Li, Brian Bondy, Weiwei Zhong, Max F. Oginsky, Yang Wu, Christopher
861    M. Johnson, Shuang Zhang, Ningren Cui, and Chun Jiang. 2016. "Identification of a
862    Group of GABAergic Neurons in the Dorsomedial Area of the Locus Coeruleus."
863    Edited by Benjamin Arenkiel. *PLOS ONE* 11 (1): e0146470.
864    https://doi.org/10.1371/journal.pone.0146470.
865 Kempadoo, Kimberly A., Eugene V. Mosharov, Se Joon Choi, David Sulzer, and Eric R.
866    Kandel. 2016. "Dopamine Release from the Locus Coeruleus to the Dorsal
867    Hippocampus Promotes Spatial Learning and Memory." *Proceedings of the National*
868    *Academy of Sciences* 113 (51): 14835–40. https://doi.org/10.1073/pnas.1616515114.
869 Klimek, Violetta, Craig Stockmeier, James Overholser, Herbert Y. Meltzer, Sheila Kalka,
870    Ginny Dilley, and Gregory A. Ordway. 1997. "Reduced Levels of Norepinephrine
871    Transporters in the Locus Coeruleus in Major Depression." *The Journal of*
872    *Neuroscience* 17 (21): 8451–58. https://doi.org/10.1523/JNEUROSCI.17-21-
873    08451.1997.
874 Kuo, Chao-Cheng, Jung-Chien Hsieh, Hsing-Chun Tsai, Yu-Shan Kuo, Hau-Jie Yau, Chih-
875    Cheng Chen, Ruei-Feng Chen, Hsiu-Wen Yang, and Ming-Yuan Min. 2020.
876    "Inhibitory Interneurons Regulate Phasic Activity of Noradrenergic Neurons in the
877    Mouse Locus Coeruleus and Functional Implications." *The Journal of Physiology* 598
878    (18): 4003–29. https://doi.org/10.1113/JP279557.
879 Lakhlani, P. P., L. B. MacMillan, T. Z. Guo, B. A. McCool, D. M. Lovinger, M. Maze, and L. E.
880    Limbird. 1997. "Substitution of a Mutant 2a-Adrenergic Receptor via 'Hit and Run'
881    Gene Targeting Reveals the Role of This Subtype in Sedative, Analgesic, and
882    Anesthetic-Sparing Responses in Vivo." *Proceedings of the National Academy of*
883    *Sciences* 94 (18): 9950–55. https://doi.org/10.1073/pnas.94.18.9950.
884 Latsari, Maria, Ioanna Dori, John Antonopoulos, Maria Chiotelli, and Athanasios Dinopoulos.
885    2002. "Noradrenergic Innervation of the Developing and Mature Visual and Motor
886    Cortex of the Rat Brain: A Light and Electron Microscopic Immunocytochemical
887    Analysis." *The Journal of Comparative Neurology* 445 (2): 145–58.
888    https://doi.org/10.1002/cne.10156.
889 Lee, Amy, Diane L Rosin, and Elisabeth J Van Bockstaele. 1998. "A2A-Adrenergic
890    Receptors in the Rat Nucleus Locus Coeruleus: Subcellular Localization in
891    Catecholaminergic Dendrites, Astrocytes, and Presynaptic Axon Terminals," 13.
892 Loughlin, S.E., S.L. Foote, and F.E. Bloom. 1986. "Efferent Projections of Nucleus Locus
893    Coeruleus: Topographic Organization of Cells of Origin Demonstrated by Three-
894    Dimensional Reconstruction." *Neuroscience* 18 (2): 291–306.
895    https://doi.org/10.1016/0306-4522(86)90155-7.
896 Loughlin, S.E., Stephen L. Foote, and James H. Fallon. 1982. "Locus Coeruleus Projections
897    to Cortex: Topography, Morphology and Collateralization." *Brain Research Bulletin* 9
898    (1–6): 287–94. https://doi.org/10.1016/0361-9230(82)90142-3.
899 Lüscher, Christian, and Paul A. Slesinger. 2010. "Emerging Roles for G Protein-Gated

900   Inwardly Rectifying Potassium (GIRK) Channels in Health and Disease." *Nature*
901   *Reviews Neuroscience* 11 (5): 301–15. https://doi.org/10.1038/nrn2834.
902 Martins, Ana Raquel O, and Robert C Froemke. 2015. "Coordinated Forms of Noradrenergic
903   Plasticity in the Locus Coeruleus and Primary Auditory Cortex." *Nature Neuroscience*
904   18 (10): 1483–92. https://doi.org/10.1038/nn.4090.
905 Mason, Stephen T., and Hans C. Fibiger. 1979. "Regional Topography within Noradrenergic
906   Locus Coeruleus as Revealed by Retrograde Transport of Horseradish Peroxidase."
907   *The Journal of Comparative Neurology* 187 (4): 703–24.
908   https://doi.org/10.1002/cne.901870405.
909 Nirenberg, Melissa J., June Chan, Yongjian Liu, Robert H. Edwards, and Virginia M. Pickel.
910   1996. "Ultrastructural Localization of the Vesicular Monoamine Transporter-2 in
911   Midbrain Dopaminergic Neurons: Potential Sites for Somatodendritic Storage and
912   Release of Dopamine." *The Journal of Neuroscience* 16 (13): 4135–45.
913   https://doi.org/10.1523/JNEUROSCI.16-13-04135.1996.
914 Olschowka, J A, M E Molliver, R Grzanna, F L Rice, and J T Coyle. 1981. "Ultrastructural
915   Demonstration of Noradrenergic Synapses in the Rat Central Nervous System by
916   Dopamine-Beta-Hydroxylase Immunocytochemistry." *Journal of Histochemistry &*
917   *Cytochemistry* 29 (2): 271–80. https://doi.org/10.1177/29.2.7019303.
918 O'Neil, Jahn N., Peter R. Mouton, Yousef Tizabi, Mary Ann Ottinger, De-Liang Lei, Donald K.
919   Ingram, and Kebreten F. Manaye. 2007. "Catecholaminergic Neuronal Loss in Locus
920   Coeruleus of Aged Female Dtg APP/PS1 Mice." *Journal of Chemical Neuroanatomy*
921   34 (3–4): 102–7. https://doi.org/10.1016/j.jchemneu.2007.05.008.
922 Pickel, Virginia M., Susan C. Beckley, Tong H. Joh, and Donald J. Reis. 1981.
923   "Ultrastructural Immunocytochemical Localization of Tyrosine Hydroxylase in the
924   Neostriatum." *Brain Research* 225 (2): 373–85. https://doi.org/10.1016/0006-
925   8993(81)90843-X.
926 Poe, Gina R., Stephen Foote, Oxana Eschenko, Joshua P. Johansen, Sebastien Bouret,
927   Gary Aston-Jones, Carolyn W. Harley, et al. 2020. "Locus Coeruleus: A New Look at
928   the Blue Spot." *Nature Reviews Neuroscience* 21 (11): 644–59.
929   https://doi.org/10.1038/s41583-020-0360-9.
930 Rice, Margaret E., and Stephanie J. Cragg. 2008. "Dopamine Spillover after Quantal
931   Release: Rethinking Dopamine Transmission in the Nigrostriatal Pathway." *Brain*
932   *Research Reviews* 58 (2): 303–13. https://doi.org/10.1016/j.brainresrev.2008.02.004.
933 Rice, Margaret E., and Charles Nicholson. 1995. "Diffusion and Ion Shifts in the Brain
934   Extracellular Microenvironment and Their Relevance for Voltammetric
935   Measurements." In *Voltammetric Methods in Brain Systems*, edited by Alan A.
936   Boulton, Glen B. Baker, and Ralph N. Adams, 27–79. Neuromethods. Totowa, NJ:
937   Humana Press. https://doi.org/10.1385/0-89603-312-0:27.
938 Rice, Margaret E., and Jyoti C. Patel. 2015. "Somatodendritic Dopamine Release: Recent
939   Mechanistic Insights." *Philosophical Transactions of the Royal Society B: Biological*
940   *Sciences* 370 (1672): 20140185. https://doi.org/10.1098/rstb.2014.0185.
941 Schwarz, Lindsay A., Kazunari Miyamichi, Xiaojing J. Gao, Kevin T. Beier, Brandon
942   Weissbourd, Katherine E. DeLoach, Jing Ren, et al. 2015. "Viral-Genetic Tracing of
943   the Input–Output Organization of a Central Noradrenaline Circuit." *Nature* 524 (7563):
944   88–92. https://doi.org/10.1038/nature14600.
945 Segal, M. 1979. "Serotonergic Innervation of the Locus Coeruleus from the Dorsal Raphe
946   and Its Action on Responses to Noxious Stimuli." *The Journal of Physiology* 286 (1):
947   401–15. https://doi.org/10.1113/jphysiol.1979.sp012628.
948 Singewald, N. 1998. "Release of Neurotransmitters in the Locus Coeruleus." *Progress in*
949   *Neurobiology* 56 (2): 237–67. https://doi.org/10.1016/S0301-0082(98)00039-2.
950 Singewald, N., C. Schneider, A. Pfitscher, and A. Philippu. 1994. "In Vivo Release of
951   Catecholamines in the Locus Coeruleus." *Naunyn-Schmiedeberg's Archives of*
952   *Pharmacology* 350 (4). https://doi.org/10.1007/BF00178948.
953 Sturrock, R. R., and K. A. Rao. 1985. "A QUANTITATIVE HISTOLOGICAL STUDY OF
954   NEURONAL LOSS FROM THE LOCUS COERULEUS OF AGEING MICE."

Neuropathology and Applied Neurobiology 11 (1): 55–60.
https://doi.org/10.1111/j.1365-2990.1985.tb00004.x.

Swanson, L W. 1976. "THE LOCUS COERULEUS: A CYTOARCH1TECTONIC, GOLGI
AND IMMUNOHISTOCHEMICAL STUDY IN THE ALBINO RAT," no. 110: 39–56.

Szabadi, Elemer. 2013. "Functional Neuroanatomy of the Central Noradrenergic System."
Journal of Psychopharmacology 27 (8): 659–93.
https://doi.org/10.1177/0269881113490326.

Takeuchi, Tomonori, Adrian J. Duszkiewicz, Alex Sonneborn, Patrick A. Spooner, Miwako
Yamasaki, Masahiko Watanabe, Caroline C. Smith, et al. 2016. "Locus Coeruleus
and Dopaminergic Consolidation of Everyday Memory." *Nature* 537 (7620): 357–62.
https://doi.org/10.1038/nature19325.

Tillage, Rachel P., Genevieve E. Wilson, L. Cameron Liles, Philip V. Holmes, and David
Weinshenker. 2020. "Chronic Environmental or Genetic Elevation of Galanin in
Noradrenergic Neurons Confers Stress Resilience in Mice." *The Journal of
Neuroscience* 40 (39): 7464–74. https://doi.org/10.1523/JNEUROSCI.0973-20.2020.

Torrecilla, María, Irrintzi Fernández-Aedo, Aurora Arrue, Mercedes Zumarraga, and Luisa
Ugedo. 2013. "Role of GIRK Channels on the Noradrenergic Transmission in Vivo:
An Electrophysiological and Neurochemical Study on GIRK2 Mutant Mice."
International Journal of Neuropsychopharmacology 16 (5): 1093–1104.
https://doi.org/10.1017/S1461145712000971.

Totah, Nelson K., Nikos K. Logothetis, and Oxana Eschenko. 2020. "Synchronous Spiking
Associated with High Gamma Oscillations in Prefrontal Cortex Exerts Top-down
Control over a 5Hz-Rhythmic Modulation of Spiking in Locus Coeruleus." Preprint.
Neuroscience. https://doi.org/10.1101/2020.04.26.061085.

Totah, Nelson K., Ricardo M. Neves, Stefano Panzeri, Nikos K. Logothetis, and Oxana
Eschenko. 2018. "The Locus Coeruleus Is a Complex and Differentiated
Neuromodulatory System." *Neuron* 99 (5): 1055-1068.e6.
https://doi.org/10.1016/j.neuron.2018.07.037.

Travagli, R. A, Thomas V. Dunwiddie, and John T. Williams. 1995. "Opioid Inhibition in Locus
Coeruleus." *Journal of Neurophysiology* 74 (2): 519–28.
https://doi.org/10.1152/jn.1995.74.2.519.

Travagli, R. A., M. Wessendorf, and J. T. Williams. 1996. "Dendritic Arbor of Locus
Coeruleus Neurons Contributes to Opioid Inhibition." *Journal of Neurophysiology* 75
(5): 2029–35. https://doi.org/10.1152/jn.1996.75.5.2029.

Uematsu, Akira, Bao Zhen Tan, Edgar A Ycu, Jessica Sulkes Cuevas, Jenny Koivumaa,
Felix Junyent, Eric J Kremer, Ilana B Witten, Karl Deisseroth, and Joshua P
Johansen. 2017. "Modular Organization of the Brainstem Noradrenaline System
Coordinates Opposing Learning States." *Nature Neuroscience* 20 (11): 1602–11.
https://doi.org/10.1038/nn.4642.

Wagner-Altendorf, Tobias A., Beatrice Fischer, and Jochen Roeper. 2019. "Axonal
Projection-specific Differences in Somatodendritic A2 Autoreceptor Function in Locus
Coeruleus Neurons." *European Journal of Neuroscience* 50 (11): 3772–85.
https://doi.org/10.1111/ejn.14553.

Walters, Seth H., I. Mitch Taylor, Zhan Shu, and Adrian C. Michael. 2014. "A Novel
Restricted Diffusion Model of Evoked Dopamine." *ACS Chemical Neuroscience* 5 (9):
776–83. https://doi.org/10.1021/cn5000666.

Waterhouse, Barry D., Barbara Border, Linda Wahl, and Gregory A. Mihailoff. 1993.
"Topographic Organization of Rat Locus Coeruleus and Dorsal Raphe Nuclei:
Distribution of Cells Projecting to Visual System Structures." *The Journal of
Comparative Neurology* 336 (3): 345–61. https://doi.org/10.1002/cne.903360304.

Williams, J.T., D.H. Bobker, and G.C. Harris. 1991. "Synaptic Potentials in Locus Coeruleus
Neurons in Brain Slices." In *Progress in Brain Research*, 88:167–72. Elsevier.
https://doi.org/10.1016/S0079-6123(08)63806-6.

Williams, J.T., R.A. North, S.A. Shefner, S. Nishi, and T.M. Egan. 1984. "Membrane
Properties of Rat Locus Coeruleus Neurones." *Neuroscience* 13 (1): 137–56.

https://doi.org/10.1016/0306-4522(84)90265-3.
Williams, J.T., Ra North, and T Tokimasa. 1988. "Inward Rectification of Resting and Opiate-
Activated Potassium Currents in Rat Locus Coeruleus Neurons." *The Journal of*
Neuroscience 8 (11): 4299–4306. https://doi.org/10.1523/JNEUROSCI.08-11-
04299.1988.
Zhang, Zizhen, Steven Cordeiro Matos, Sonia Jego, Antoine Adamantidis, and Philippe
Séguéla. 2013. "Norepinephrine Drives Persistent Activity in Prefrontal Cortex via
Synergistic A1 and A2 Adrenoceptors." Edited by Kevin Currie. *PLoS ONE* 8 (6):
e66122. https://doi.org/10.1371/journal.pone.0066122.

Data availability

All code used for all analyses and plots are publicly available on GitHub at
https://github.com/teamprigge

Acknowledgments

We are thankful to the Center for Brain and Behavioral Science for financing K.M. M.P is supported by the Best Minds programs from the Leibniz Association. We are in depth to Menachem Segal for discussion and Oliver Kobler from the Combinatorial NeuroImaging (CNI) core for support with light sheet and confocal microscopy.

Author contributions

SB, HH, KM, TF, JB and MP conceived experiment(s) and developed the analyses. MP wrote first draft of manuscript with support SB, KM and JB. MP and JB supervised the work, and acquired funding.

Competing interests the authors declare no competing interests.

1043   **Supplement**

1044
1045   Supp. Figure 1:

1046
The calcium current is responsible for the spontaneous spiking of the cell. It is unclear in

Alvarez et al if this is a transient conductance (with activation and inactivation gates) or a

persistent conductance (with two kinds of activation gates). If we take the formulas as given,

The 'fast' gating variable mCa opens with membrane depolarization. Specifically, it

approaches unity for membrane potentials above −40 mV (see Supp Figure 1). The 'slow'

gating variable y_∞ closes with membrane depolarization. Specifically, it approaches zero for

membrane potentials above −60 mV . In the same regime, the characteristic time to

approaches 20 ms. Taken together, this current is active in a window between −80mV and

−40mV , being strongest around −60mV . As a result, this current can cause a slow Ca-

spike, on which multiple fast Na-spikes can occur. Activation variables for y_∞ (left) and m_ca

(middle) and I_ca (right) for calcium current, as a function of membrane voltage, for the formulas as

given by Alvarez et al.

1059
1060

1073   Supp. Figure 2:

1074

1090

1091   Inward-rectification presented in a current-voltage plot derived from HEK293 that heterologus
1092   express GIRK2 channels. GIRK time course are used in equation 13.

1123 Supp. Figure 3:

1124
1125
1126 Confocal images of immunostained brainstem slices against tyrosine hydroxylase (TH / green)
1127 and dopamin-beta-hydroxylase (DbH / magenta) at three different position from bregma: - 5.40
1128 , - 5.52 and -5.68mm. **(a)** Shows LCs slices from an old animal (P500), while **(b)** overviews LC
1129 slices from a young animal (P14). Somatic as well as dendritic compartments exhibit different
1130 expression levels for TH and DbH within the LC topography.

1155 Supp. Figure 4:

1156
1157 Spike rate dependence on 300-ms current-injection steps separated for early spikes (spike 1 and 2 /
1158 dotted line) versus later spikes (> spike 2 / full line). **(a)** FI Curve of an experimental neuron. It is
1159 firing at almost 6.1Hz at spontaneous state. **(b)** FI curve of a simulated neuron. It is firing at almost
1160 4.5Hz at spontaneous state.

1161
1162 Supp. Figure 5:

1163
1164

1165
1166 Action potential triggers calcium accumulation in the cell which facilitates the release of
1167 neurotransmitters along with many other calcium-triggered events. When the simulated neuron
1168 is firing at 4Hz, a small amount of calcium is accumulated and equibrilates at constant level as
1169 shown in fig 3 **(a)**. When the same neuron is firing at 20Hz, a higher level of calcium is
1170 accumulated after several action potentials **(b)**.

1186 Supp. Figure 6:
1187

1188
1189
Effect of extracellular NE that diffuses freely in the extracellular space. The released NE
binds to a receptor site in a nearby neuron and triggers opening of GIRK. Unlike synaptic
inhibition, inhibition by diffusion is delayed in nature as it takes time to reach the binding site
when diffused in extracellular space. Here ISI (binned 2s) are calculated for neuron N2 in
figure 5. Inhibition slowly builds up (1s-7s) after N1 releases NE during several intervals
which leads to a significantly longer ISI in N3. N3 slowly gets back to the steady state after
N1 stops releasing NE(9s).