1
2 **Main Manuscript for**
3 The V-type H$^+$-ATPase is targeted in anti-diuretic hormone control of the
4 Malpighian 'renal' tubules
5

Farwa Sajadi[1], María Fernanda Vergara-Martínez[1,2] and Jean-Paul V. Paluzzi[1]*
1. Department of Biology, York University, 4700 Keele Street, Toronto, Ontario, M3J 1P3, Canada.
2. Departamento de Biología Celular y Fisiología, Instituto de Investigaciones Biomédicas, Universidad
Nacional Autónoma de México, Mexico City, Mexico.

*Corresponding author: Jean-Paul V. Paluzzi
Email: paluzzi@yorku.ca

12
13 PNAS strongly encourages authors to supply an ORCID identifier for each author. Do not include ORCIDs
14 in the manuscript file; individual authors must link their ORCID account to their PNAS account at
15 www.pnascentral.org. For proper authentication, authors must provide their ORCID at submission and are
16 not permitted to add ORCIDs on proofs.
17
18 **Author Contributions:** J.-P.P. and F.S. designed research; F.S. and M.F.V. performed research; F.S. and
19 J.P.P. analyzed data; and J.-P.P. and F.S. wrote the manuscript.
20 **Competing Interest Statement:** The authors declare no competing interest.
21 **Classification:** Biological Sciences; Physiology
22 **Keywords:** excretory system, *Aedes aegypti*, mosquito, anti-diuresis, proton pump, disease vector
23
24 **This PDF file includes:**
25   Main Text
26   Figures 1 to 6; S1 to S7

Abstract

Like other insects, secretion by mosquito Malpighian tubules (MTs) is driven by the V-type H^+-ATPase (VA) localized in the apical membrane of principal cells. The anti-diuretic neurohormone CAPA inhibits secretion by MTs stimulated by select diuretic hormones; however, the cellular effectors of this inhibitory signaling cascade remain unclear. Herein, we demonstrate that the VA inhibitor bafilomycin selectively inhibits serotonin (5HT)- and calcitonin-related diuretic hormone (DH_{31})-stimulated secretion. VA activity increases in DH_{31}-treated MTs, whereas CAPA abolishes this increase through a NOS/cGMP/PKG signaling pathway. A critical feature of VA activation involves the reversible association of the cytosolic (V_1) and membrane (V_o) complexes. Indeed, higher V_1 protein abundance was found in membrane fractions of DH_{31}-treated MTs whereas CAPA significantly decreased V_1 abundance in membrane fractions while increasing it in cytosolic fractions. Immunolocalization of V_1 was observed strictly in the apical membrane of MTs treated with DH_{31} alone whereas immunoreactivity was dispersed following CAPA treatment. VA complexes colocalized apically in female MTs shortly after a blood-meal consistent with the peak and post-peak phases of diuresis. Comparatively, V_1 immunoreactivity in MTs was more dispersed and did not colocalize with the V_o complex in the apical membrane at 3 hours post blood-meal, representing a timepoint after the late phase of diuresis has concluded. Therefore, CAPA inhibition of MTs involves reducing VA activity and promotes complex dissociation hindering secretion. Collectively, these findings reveal a key target in hormone-mediated inhibition of MTs countering diuresis that provides a deeper understanding of this critical physiological process necessary for hydromineral balance.

Significance Statement

The V-type H^+ ATPase (VA), also known as the proton pump, provides the driving force for transepithelial ion and fluid secretion in insect Malpighian tubules (MTs). While studies have shown that diuretic stimulation activates various signaling pathways promoting increased VA activity, our understanding of anti-diuretic signaling and its potential regulation of the VA remains rudimentary. Here we show that a CAPA neuropeptide acts through the NOS/cGMP/PKG pathway to inhibit VA activity, supporting the notion that the anti-diuretic regulation is achieved by promoting dissociation of the VA complexes. These results demonstrate a critical role of VA inhibition and trafficking necessary for anti-diuretic signaling and advances our understanding of the complex neuroendocrine control of the MTs in this important human disease-vector mosquito.

Main Text

Introduction

Insect post-prandial diuresis is under rigorous control by neuroendocrine factors (1) acting on the Malpighian 'renal' tubules (MTs) to regulate primary urine production. In the yellow fever mosquito, *Aedes aegypti,* several diuretics have been identified that regulate urine production including serotonin (5HT), calcitonin-related diuretic hormone (DH_{31}), corticotropin-releasing factor-related diuretic hormone (DH_{44}) and leucokinin-related (LK) diuretic hormone (2–5). An anti-diuretic peptidergic neurohormone, CAPA, selectively inhibits DH_{31}- and 5HT-stimulated secretion of MTs (4, 6, 7). Insect CAPA

neuropeptides are produced in the central nervous system and are evolutionarily related to

the vertebrate neuromedin U peptides (8). In *Drosophila melanogaster*, CAPA peptides

have been shown to act through a conserved nitridergic signaling pathway to stimulate

diuresis by MTs (9, 10); however, a few other studies have alluded to an anti-diuretic role

(11, 12). In contrast, in both larval and adult *A. aegypti*, CAPA peptides inhibit fluid

secretion through a signaling cascade involving the NOS/cGMP/PKG pathway (6, 7).

Despite this, the anti-diuretic signaling mechanism and downstream cellular targets, such

as the ion channels and transporters, remain elusive.

In insect MTs, including *A. aegypti*, the bafilomycin-sensitive V-type H^+ ATPase

(VA), also known as the proton pump, functions as an electrogenic pump allowing the

transport of protons from the cytoplasm to the tubule lumen, thus generating a cell-

negative membrane voltage (13, 14). This membrane voltage can then drive secondary

transport processes such as the cation/H^+ exchanger or anion/H^+ cotransporter (15, 16).

Originally found in vacuolar membranes of animals and plants, the VA has since been

found to be essential in cell function in both invertebrates and vertebrates (17). In insects,

the VA is densely located in the apical brush border membrane of tubule principal cells

(13, 18), which is rich in mitochondria and endoplasmic reticula, fueling the ATP-

consuming proton pump (1, 19). Previous studies have shown VA localization within the

apical membrane of principal cells along the entire length of the MTs (19), but absent in

stellate cells that express relatively higher levels of the P-type Na^+/K^+ ATPase (NKA)

(19). Due to stronger VA immunoreactivity observed in MTs (19), and greater ATPase

activity by electrophysiological assays (13, 14), the VA is categorized as serving mainly,

but not exclusively (20), stimulated transport mechanisms, whereas the NKA serves basic

cell housekeeping functions when MTs are undergoing low unstimulated rates of

secretion (13, 20, 21).

Stimulation of distinct diuretic hormone receptors can activate various signaling

pathways, including elevation of cyclic AMP (cAMP) levels which is known to increase

VA activity and assembly in insects (22). *In A. aegypti,* DH_{31}, identified as the mosquito

natriuretic peptide (23), selectively activates transepithelial secretion of Na^+ in the MTs,

using cAMP as a second messenger (1), and upregulating the VA function to stimulate

fluid secretion (24). Similarly, 5HT-stimulated diuresis is also thought to be mediated (at

least in part) through the cAMP second messenger pathway (25), activating protein

kinase A (PKA) to increase the transepithelial voltage of the basolateral membrane in

tubule principal cells (1, 26). In contrast, DH_{44} has been shown to initiate diuresis via the

paracellular and transcellular pathways, with higher nanomolar concentrations increasing

cAMP and Ca^{2+}, influencing both paracellular and transcellular transport, and lower

nanomolar concentrations acting through the paracellular pathway only, via intracellular

Ca^{2+} (27). Thus, due to its predominant role in fluid secretion, the VA could be a likely

target for both diuretic and anti-diuretic factors.

Eukaryotic V-ATPases are a multi-subunit protein composed of up to 14 different

polypeptides, which form two major structural complexes. The peripheral V_1 complex

(400-600 kDa), is invariably present in the cytoplasm and interacts with ATP, ADP, and

inorganic phosphate (28). The cytosolic V_1 complex consists of eight different subunits

(A-H): a globular headpiece with three alternating subunits A and B forming a hexamer

with nucleotide binding sites located at their interface, a central rotor stalk with single

copies of subunits D and F, and lastly, a peripheral stalk made up of subunits C, E, G, and

H (28). The B subunit, shown to have high sequence similarity amongst several species

from fungi to mammals (29, 30), is a 56 kDa polypeptide, and is one of the two sites

(along with subunit A) (31) in the V_1 complex that binds ATP. In contrast, the

membrane-integrated V_o complex (150-350 kDa) mediates the transport of H^+ across the

membrane (28) and is composed of at least six different subunits, which collectively

function in the proton translocation pathway (13, 28). Although the proton channel of the

VA can be blocked pharmacologically by the macrolide antibiotic, bafilomycin (32),

there are two known intrinsic mechanisms for VA regulation: firstly, through oxidation of

the cystine residue on the A subunit of the V_1 complex, thus preventing ATP hydrolysis;

secondly, through reversible disassembly of the V_1 complex from the holoenzyme (13,

33). While the role and regulation of the VA by diuretic hormones in insect MTs has

been studied (34–36), research examining anti-diuretic signaling mechanisms involving

the VA remain in their infancy.

This study aimed to identify the cellular targets necessary for CAPA-mediated

inhibition of fluid secretion by MTs stimulated by select diuretic factors in adult female

A. aegypti. Our results provide evidence that CAPA neuropeptides inhibit fluid secretion

by VA complex dissociation, thus hindering VA function and activity that is essential for

driving rapid post-prandial diuresis.

Results

Bafilomycin inhibits DH_{31}- and 5HT-stimulated fluid secretion rate. To determine the

appropriate concentration of bafilomycin to test on adult *A. aegypti* MTs, several doses

were applied against DH_{31}-stimulated tubules (Fig. S1). Higher doses of bafilomycin (10^-

^{4} M and 10^{-5} M) resulted in significant inhibition of fluid secretion rate, with maximal

inhibition leading to a five-fold decrease, observed with treatment of 10^{-5} M bafilomycin.

Next, to determine whether inhibiting the VA would decrease fluid secretion rate

stimulated by other diuretic hormones, including 5HT and DH_{44}, the effect of 10^{-5} M

bafilomycin on adult tubules stimulated with these diuretics was tested (Fig. 1). Fluid

secretion rates were measured over 30 min under control (stimulated) conditions, and

then at 10-min intervals in the presence of bafilomycin. Treatment of MTs with 10^{-5} M

bafilomycin against DH_{31} led to a decrease of fluid secretion over the treatment interval.

Specifically, 30 min after treatment with bafilomycin, fluid secretion rate was

significantly reduced by over two-fold to 0.438 ± 0.041 nL min^{-1}, compared to DH_{31} alone,

0.941 ± 0.077 nL min^{-1} (Fig. 1A). Similar results were seen with 5HT-stimulated MTs;

however, a decrease in fluid secretion was observed 30 min post-bafilomycin, with a

significant inhibition 40 min after treatment (0.522 ± 0.072 nL min^{-1}, 5HT alone vs.

0.182 ± 0.045 nL min^{-1}, 5HT + bafilomycin) (Fig. 1B). Distinct from DH_{31} and 5HT-

stimulated tubules, DH_{44}-stimulated secretion was insensitive to bafilomycin treatment

(Fig. 1C). To confirm whether *Aedae*CAPA-1 anti-diuresis is mediated by VA inhibition,

adult female MTs were treated with either DH_{31} alone or in combination with

*Aedae*CAPA-1, bafilomycin, or both (Fig. 1D). MTs treated with either *Aedae*CAPA-1 or

bafilomycin resulted in a significant inhibition of DH_{31}-stimulated secretion, and similar

inhibition was observed when both *Aedae*CAPA-1 and bafilomycin were applied together

with no evidence of any additive inhibitory effects (Fig. 1D).

***Aedae*CAPA-1 and bafilomycin alkalinizes secreted fluid in DH$_{31}$- and 5HT-**

stimulated MTs. The VA pumps protons from the cell into the tubule lumen thus

generating an electromotive potential (20, 31) and providing energy to drive the secretion

of cations via Na$^+$/H$^+$ and/or K$^+$/H$^+$ antiporters (13). An indirect way to measure whether

*Aedae*CAPA-1 and bafilomycin inhibits VA activity involves measuring the pH of the

secreted fluid from diuretic-stimulated MTs treated with *Aedae*CAPA-1 or bafilomycin

(Fig. 1E-G). In DH$_{31}$-stimulated MTs treated with *Aedae*CAPA-1, there was an

immediate significantly higher pH in the secreted fluid (7.479±0.030) at 40 min relative

to control, increasing up to 7.73±0.038 at 60 min (Fig. 1E). Similarly, pH levels in DH$_{31}$-

stimulated MTs treated with bafilomycin were significantly higher (7.66±0.064) relative

to control at 40 min, increasing up to 7.855±0.074 at 60 min. Comparable to DH$_{31}$,

addition of *Aedae*CAPA-1 or bafilomycin, significantly increased the pH of secreted fluid

from 5HT-stimulated MTs to 7.75±0.061 and 7.82±0.083 respectively, at the 60 min

mark (Fig. 1F). In contrast, unlike the effects observed with DH$_{31}$- and 5HT-stimulated

MTs, *Aedae*CAPA-1 or bafilomycin did not alter the pH of the secreted fluid in DH$_{44}$-

stimulated MTs (Fig. 1G). The pH increased from 7.4 to 7.9 during the 30-min DH$_{44}$

incubation; however, pH did not change following the addition of *Aedae*CAPA-1 or

bafilomycin. Separately, we conducted measurements in unstimulated tubules to verify

pH in these small droplets did not drift over a time frame consistent with our above

experiments. Unstimulated MTs were allowed to secrete, and the droplets were isolated

and their pH was measured over the course of 60 min (Fig. S2). Over this incubation

period, no change was observed in the pH of secreted droplets from unstimulated MTs,

upholding the notion that the alkalinization of secreted fluid observed following

*Aedae*CAPA-1 (or bafilomycin) treatment of DH_{31}- and 5HT-stimulated MTs is a result

of VA inhibition. Additionally, unstimulated MTs treated with either *Aedae*CAPA-1 or

bafilomycin resulted in no significant changes in either secretion rate (Fig. S3A) or pH

(Fig. S3B).

***Aedae*CAPA-1 increases cGMP and decreases cAMP levels in DH_{31}-treated MTs.** To

further clarify the CAPA signaling pathway involving the second messengers, cGMP and

cAMP, we sought to determine changes in levels of these cyclic nucleotides in MTs

incubated in DH_{31} alone or combined with *Aedae*CAPA-1. Treatment of MTs with DH_{31}

alone had basal levels of cGMP, 10.91 ± 0.109 $pmol^{-1}$/tubule, comparable to saline treated

MTs. Treatment of MTs with *Aedae*CAPA-1 resulted in a significant increase in cGMP

levels compared to DH_{31}- incubated tubules, increasing to 11.39 ± 0.101 $pmol^{-1}$/tubule

(Fig. 1I). Similar results were observed with MTs treated with both DH_{31} + *Aedae*CAPA-

1, with significantly increased cGMP levels of 11.39 ± 0.123 $pmol^{-1}$/tubule (Fig. 1I)

compared to MTs treated with DH_{31} alone. In contrast, treatment of MTs with DH_{31} alone

led to significantly higher levels of cAMP, 9.153 ± 0.039 $pmol^{-1}$/tubule, while baseline

levels of this second messenger were observed in saline, DH_{31} + *Aedae*CAPA-1, and

*Aedae*CAPA-1 treated tubules (Fig. 1J). To further confirm the stimulatory role of cAMP

and inhibitory role of cGMP, tubules were treated with either cyclic nucleotide alone and

secretion rates were measured (Fig. 1H). Unstimulated fluid secretion rates were

measured over the first 30 min, and then at 10-min intervals with either cAMP or cGMP.

Treatment of MTs with 10^{-4} M cAMP led to a significant increase over the treatment

interval, with fluid secretion rates increasing to 0.615 ± 0.096 nL min^{-1} at 60 min,

compared to 10^{-8} M cGMP (0.052 ± 0.091 nL min^{-1}) and unstimulated (0.065 ± 0.091 nL

min^{-1}) (Fig. 1H). Finally, to establish whether these cyclic nucleotide second messengers

elicit antagonistic control of the MTs in adult *A. aegypti*, tubules were treated initially

with cAMP over the first 30 min and then cGMP was added in the presence of cAMP for

a subsequent 30 min (Fig. 1K). Similarly, we also tested the opposite treatment regimen

where MTs were treated initially with cGMP and subsequently with cAMP added along

with cGMP. Treatment of cAMP-stimulated MTs with 10^{-8} M cGMP led to a significant

decrease (~4-fold) over the treatment interval, with secretion rates decreasing to

0.231 ± 0.113 nL min^{-1} at 60 min, compared to 10^{-4} M cAMP alone (0.931 ± 0.134 nL min^{-}

^{1}). In contrast, cGMP-incubated tubules treated with 10^{-4} M cAMP led to a significant

increase (~10-fold) in secretion rate (0.537 ± 0.072 nL min^{-1}) compared to MTs treated

with 10^{-8} M cGMP alone (0.045 ± 0.018 nL min^{-1}).

Parallel studies examining cAMP levels were measured in DH$_{44}$-incubated MTs.

Treatment of MTs with DH$_{44}$ alone had high levels of cAMP, $9.115\pm0.0.061$ pmol^{-}

^{1}/tubule, compared to saline treated MTs, 8.709 ± 0.081 pmol^{-1}/tubule (Fig. S4A). Levels

of cAMP remained unchanged in tubules treated with both DH$_{44}$ + *Aedae*CAPA-1,

8.954 ± 0.108 pmol^{-1}/tubule. To further resolve the cAMP signaling pathway downstream

of DH$_{31}$ and DH$_{44}$ stimulated diuresis, a PKA inhibitor (KT5720) was tested against

diuretic-stimulated MTs. KT5720 abolished the stimulatory effect of DH$_{31}$, whereas

secretion by DH$_{44}$-treated MTs remained unchanged (Fig. S4B).

_Aedae_CAPA-1 decreases VA activity in DH$_{31}$-stimulated MTs through the

NOS/cGMP/PKG pathway. In *Aedes* MTs, 50-60% of the total ATPase activity can be

attributed to a bafilomycin- and nitrate-sensitive component that reflects the activity of

the VA pump (13). The remaining ATPase activity may be due to nucleotide cyclases,

protein kinases, myosin, DNA helicases, and other ATP-consuming processes such as the

NKA (13). As such, to determine whether CAPA inhibits VA and/or NKA function,

female diuretic-stimulated MTs were challenged with *Aedae*CAPA-1 to measure the

resultant NKA and VA activity. As expected, given its role as the natriuretic diuretic

hormone, adult female MTs treated with DH_{31} resulted in a significant (> two-fold)

increase of VA activity, 0.0329 ± 0.0007 μmoles ADP/μg protein/hour, compared to saline

controls, 0.0151 ± 0.0021 μmoles ADP/μg protein/hour (Fig. 2A). Importantly, MTs

incubated with both DH_{31} and *Aedae*CAPA-1 had significantly lower VA activity,

resulting in activity levels indistinguishable from saline controls. In contrast, neither 5HT

nor DH_{44} influenced VA activity (p>0.05) when compared with saline controls, while co-

treatment with *Aedae*CAPA-1 also resulted in indistinguishable VA activity. Similar VA

activity levels were observed between 5HT and DH_{44} (0.0255 ± 0.0078 and 0.0208 ± 0.0042

μmoles ADP/μg protein/hour) and with co-application of *Aedae*CAPA-1 (0.0150 ± 0.0036

and 0.0154 ± 0.0070 μmoles ADP/μg protein/hour). Unlike changes observed in VA

activity following treatment with DH_{31}, diuretic-stimulation or *Aedae*CAPA-1 treatment

did not perturb NKA activity, with levels similar to that of unstimulated MTs (Fig. 2B).

To confirm the actions of CAPA are mediated through the NOS/cGMP/PKG

pathway, pharmacological blockers, including inhibitors of NOS (L-NAME) and PKG

(KT5823), were tested against DH_{31}-stimulated MTs treated with either *Aedae*CAPA-1 or

cGMP (Fig. 2C). Application of L-NAME or KT5823 abolished the inhibitory effect of

*Aedae*CAPA-1, resulting in high levels of VA activity, 0.02671 ± 0.0025 and

0.03653±0.0051 µmoles ADP/µg protein/hour respectively, compared to MTs treated

with DH_{31} + *Aedae*CAPA-1. As expected, treatment of DH_{31}-stimulated MTs with cGMP

resulted in a significant decrease in VA activity, 0.006±0.0026 µmoles ADP/µg

protein/hour, similar to *Aedae*CAPA-1-treated MTs, while co-treatment with KT5823,

abolished the inhibitory effect of cGMP, resulting in an increase in VA activity.

_Aedae_CAPA-1 leads to VA holoenzyme dissociation in DH_{31}-treated MTs. The

reversible dissociation of the V_1 complex from the V_o membrane-integrated complex is a

well-known mechanism for regulating VA transport activity (22, 31, 34, 35, 37). To

determine whether *Aedae*CAPA-1 influences VA complex dissociation, membrane and

cytosolic protein fractions were isolated from DH_{31} and DH_{31} + *Aedae*CAPA-1 incubated

MTs, and a polyclonal V_1 antibody (13) was used to measure protein abundance. First,

membrane and cytosolic protein isolation was verified with specific cytosolic (beta-actin)

and membrane (AQP1) markers (Fig. S5). Western blot analysis revealed three protein

bands, with calculated molecular masses of 74 kDa, 56 kDa, and 32 kDa (13, 38) (Fig.

S6). The V_1 complex is composed of eight subunits (A-H), which includes the A

(~74kDa) and B subunit (~56Da) that are arranged in a ring forming the globular

headpiece for ATP binding and hydrolysis (31). Additionally, studies have suggested that

subunit D (~32kDa) alongside subunit F constitute the central rotational stalk of the V_1

complex (17). There was no difference in abundance observed for the A subunit (74 kDa

band) in either membrane or cytosolic fractions between saline and DH_{31} treatments

whereas DH_{31} + *Aedae*CAPA-1 incubated MTs had increased A subunit (74 kDa band)

protein abundance in cytosolic fractions compared to saline treatment (Fig. 3A) and

decreased abundance in the membrane fraction compared to MTs treated solely with

DH_{31} (Fig. 3B). Similarly, the V_1 complex B subunit abundance (56 kDa band) was

similar in all treatments within the cytosolic protein fraction (Fig. 3C) whereas DH_{31} +

*Aedae*CAPA-1 incubated MTs had significantly lower abundance in membrane fractions

compared to MTs treated with DH_{31} alone (Fig. 3D). Finally, there was no difference in

abundance of the V_1 complex subunit D (32 kDa band) between saline and DH_{31} treated

MTs in neither cytosolic or membrane fractions. However, as observed for the A and B

subunit bands (74 and 56 kDa band, respectively) DH_{31} + *Aedae*CAPA-1 incubated MTs

showed a significant increase in the D subunit in cytosolic fractions compared to saline

treated MTs (Fig. 3E) and a decrease in its abundance in membrane fraction compared to

MTs treated with DH_{31} alone (Fig. 3F). In summary, all three of the V1 complex

immunoreactive bands corresponding to subunits A, B and D (74, 56 and 32 kDa,

respectively) showed significantly higher abundance in cytosolic protein fractions and

lower abundance in membrane fractions in DH_{31} + *Aedae*CAPA-1 incubated MTs.

To visualize this potential endocrine-mediated reorganization of the VA

holoenzyme in this simple epithelium, we immunolocalized the membrane-integrated V_o

and cytosolic V_1 complex in the female *A. aegypti* MTs. Transverse sections of saline-

incubated (control) MTs demonstrated moderate enrichment of V_o (Fig. 4A, red), and V_1

(Fig 4B, green) complexes in the apical membrane of principal cells (Fig. 4C, S7A-D).

Comparatively, DH_{31}-incubated MTs revealed intense localization of the V_o (Fig. 4D,

red), and V_1 (Fig 4E, green) complexes within principal cells, where V_1 staining was

strictly co-localized with V_o staining on the apical membrane (Fig. 4F, Fig. S7E-H).

Interestingly, although V_o immunolocalization was restricted to the apical membrane, V_1

immunoreactivity was observed in both the apical membrane and cytosolic region in

DH_{31} + *Aedae*CAPA-1 co-treated MTs (Fig. 4G-I, Fig. S7I-T). Immunostaining was

absent in control preparations probed with only secondary antibodies (not shown)

confirming the specific detection of the VA complexes with each primary antibody.

To investigate this endocrine-mediated phenomenon *in vivo*, we immunolocalized

the membrane-integrated V_o and cytosolic V_1 complex in blood-fed females at different

time points. Whole body sections of non-blood-fed similarly aged females (control)

demonstrated moderate enrichment of both V_o (Fig. 5A, red), and V_1 (Fig. 4B, green) in

the MTs, with minimal co-localization (Fig. 5C), resembling saline-incubated MTs.

Interestingly, blood-fed female MTs revealed strong co-localization of the V_o and V_1

complexes at 10 min (Fig. 5D-F) and 30 min (Fig. 5G-I) post blood-meal, whereas V_1

immunoreactivity was more dispersed in both the apical membrane and cytosolic area in

MTs 3 hrs post blood-meal (Fig. 5J-L), comparable to non-blood-fed females.

Discussion

The MTs of the *Aedes* mosquito are the main organs responsible for the secretion

of water and solutes, thereby contributing towards hydromineral homeostasis of the

animal (39). Active ion transport in *A. aegypti* MTs is accomplished mainly by the V-

ATPases (VA) densely localized in the apical brush-border membrane of principal cells,

that energize the apical and basolateral membrane as well as the paracellular pathway,

allowing for transepithelial secretion of NaCl, KCl, and other solutes (40). In animal

cells, V-ATPase molecules in the plasma membrane, especially on the apical membrane

of epithelial cells, contribute to extracellular acidification or alkalization, intracellular pH

homeostasis, or energize the plasma membrane for secondary active transport (41, 42). In

insect MTs, the VA plays a major role in fluid secretion, thus serving as a primary target

for both diuretic and, as this study demonstrates, anti-diuretic hormonal regulation of the

insect 'renal' tubules. Although the structure and function of the VA has been elucidated

in some detail (13, 20, 22, 31, 43–45), the regulation of the proton pump remains unclear.

Of the various regulatory mechanisms for VA activity, the most studied is the reversible

dissociation of the cytosolic V_1 complex from the membrane-integrated V_o complex, first

established in the midgut of the tobacco hornworm, *Manduca sexta* and yeast,

Saccharomyces cerevisiae (44, 46). In this study, the activity and regulation of the VA

was investigated under both diuretic and anti-diuretic hormone control of the adult female

A. aegypti MTs. Notably, the current results advance our knowledge of the anti-diuretic

control of the *A. aegypti* MTs, revealing a cellular mechanism for CAPA inhibition of the

MTs by targeting the VA to block fluid secretion stimulated by select diuretic factors.

This includes inhibition of the DH_{31}-related mosquito natriuretic peptide, which is critical

for the post-haematophagy diuresis that eliminates excess water and sodium originating

from the bloodmeal-derived plasma.

In insects, water excretion is tightly regulated to maintain homeostasis of ions and

water (1, 47, 48). Female *A. aegypti* engorge a salt- and water-rich bloodmeal to obtain

the necessary nutrients and proteins for their eggs (1), with about 40% of the ingested

water eliminated in the first hour post feeding (49). The high rates of water excretion

along with the high rates of primary urine production post bloodmeal suggest a highly

coordinated and defined hormonal regulation of the signaling processes and downstream

cellular targets for ion and water transport (50). In *Aedes* MTs, fluid secretion increases at

least three-fold after stimulation with mosquito natriuretic peptide (identified as DH_{31}),

via cAMP as a second messenger (1), activating PKA, which subsequently activates V-

ATPase-driven cation transport processes (22, 35, 38). Herein we show that DH_{31}-

stimulated secretion is inhibited by bafilomycin, thought to block the proton channel of

the VA (32). Moreover, the addition of either bafilomycin or *Aedae*CAPA-1 caused

alkalization of the secreted fluid, indicating inhibition of the VA, which may lead to

constrained entry of cations across the apical membrane through a proposed alkali

cation/proton antiporter (15, 16). Thus, since bafilomycin inhibits DH_{31}-stimulated

secretion, this supports the VA as a target in the inhibition of fluid secretion.

Consequently, the driving force for ion movement and osmotically-obliged water is

reduced, but select Na^+ channels and cotransporters remain unaffected in the presence of

*Aedae*CAPA-1, as observed by the unchanged natriuretic effect of DH_{31} despite reduced

secretion rates in response to *Aedae*CAPA-1 (4). Similar results were seen in 5HT-

stimulated secretion, albeit a partial inhibition. An earlier study demonstrated that Ca^{2+}-

mediated diuresis does not require the assembly and activation of the VA (37, 38). The

cAMP effect on the VA is implemented by protein kinase A (PKA), with inhibitors of

PKA abolishing hormone-induced assembly and activation of the VA (34). Although the

endogenous 5HT receptor expressed within the *A. aegypti* MTs necessary for diuretic

activity remains elusive, in the kissing bug, *Rhodnius prolixus*, both cAMP and Ca^{2+} have

been shown to initiate diuresis in response to 5HT (51), which could explain the partial

inhibitory response of *Aedae*CAPA-1 inhibition on 5HT-stimulated tubules as Ca^{2+}-

mediated diuresis is independent of the VA (38). Notably, the anticipated 5HT type 2

receptor subtype expressed in the principal cells of the MTs is predicted to couple

through a Gq/11 signaling mechanism (52) and likely excludes the type 7 Gs-coupled

receptor localized to tracheolar cells associated with the MTs (53, 54) as well as the type

1 Gi-coupled receptor localized to principal cells in larval stage mosquitoes (55).

Interestingly, DH_{44}-mediated stimulation was observed to be independent of the

VA, as bafilomycin had no effect on the secretion rate or pH of the secreted fluid

following application of this CRF-related diuretic peptide. Previous studies have noted

that low nanomolar concentrations of a DH_{44}-related peptide were linked to the

stimulation of the paracellular pathway only (27), mediating this action through

intracellular Ca^{2+} as a second messenger (56). In contrast, high nanomolar concentrations

of a DH_{44}-related peptide were shown to influence both paracellular and transcellular

transport, increasing intracellular Ca^{2+} and cAMP (56). Although haemolymph

concentrations of diuretic peptides have yet to be determined in mosquitoes, DH_{31} is

immediately released into circulation post bloodmeal, stimulating rapid secretion of Na^+

and excess water (23, 49). In contrast, DH_{44}-stimulated diuresis in *A. aegypti* involves

non-selective transport Na^+ and K^+ cations (4), this supports a delayed release of this

diuretic hormone post-feeding to maintain production (albeit reduced) of primary urine

whilst conserving Na^+ ions.

In unstimulated adult female *Aedes* MTs isolated *in vitro*, the VA exhibits

variable rates of enzyme activity, consistent with highly variable rates of secretion, as

found also in various other insect species (26, 57, 58). The VA is the main energizer in

MTs as 60% of total ATPase activity can be linked to the VA (13), whereas the NKA,

with around 28% of ATPase activity, also plays a role in membrane energization,

denoting a more important role in the function of MTs than was previously assumed (19,

38–40). Here we show a significant two-fold increase in VA activity in MTs treated with

DH_{31} compared to the unstimulated MTs, with no change in VA activity in 5HT and

DH_{44}-treated MTs. Notably, *Aedae*CAPA-1 treatment blocked the DH_{31}-driven increase

in VA activity, which corroborates the reduced fluid secretion rate and alkalization of the

secreted fluid. Additionally, we sought to establish the importance of the

NOS/cGMP/PKG pathway in the inhibitory actions of *Aedae*CAPA-1 on VA association.

In stimulated MTs treated with *Aedae*CAPA-1 along with NOS inhibitor, L-NAME, or

PKG inhibitor, KT5823, the inhibitory activity of *Aedae*CAPA-1 and its second

messenger cGMP was abolished, resulting in elevated VA activity as a result of DH_{31}

treatment. The present study examined the effects of *Aedae*CAPA-1 on DH_{31}-stimulated

VA activation for the first time in insects. Stimulation of DH_{31} causes an increase in

cAMP production, which activates Na^+ channels and the $Na^+/K^+/2Cl^-$ cotransporter in the

basolateral membrane (59) and up-regulates VA activity (as shown herein and

previously) critical for increased fluid secretion (24). The DH_{31} receptor

(*Aaeg*GPRCAL1) is expressed in a distal-proximal gradient in the MTs, with greater

expression in principal cells where the VA in the apical membrane is highly expressed

(19, 60). The co-localization of the DH_{31} receptor, VA, and cation exchangers (61–63) in

the distal segment of the MTs, along with the CAPA receptor (6), collectively supports

the major roles DH_{31} and CAPA play in post-prandial diuresis and anti-diuresis,

respectively. In contrast to the marked changes in VA activity in response to diuretic and

anti-diuretic hormones, NKA activity remained unchanged in response to treatments

conducted herein. Further studies should examine the potential role of the NKA in

diuretic and anti-diuretic processes.

431   The reversible dissociation of the V_1 and V_o complexes is currently thought as a

432 universal regulatory mechanism of V-ATPases, appearing to be widely conserved from

433 yeast to animal cells (13, 22, 35, 64). Although previously shown with cAMP (24), it

434 remained unclear whether other second messengers (eg. Ca^{2+}, cGMP, and nitric oxide)

435 affect the assembly/disassembly of the V_1V_o complexes in insect MTs. In this study, VA

436 protein abundance in membrane and cytosolic fractions of MTs was confirmed by

437 western blot analyses. The 56 kDa band represents the B subunit (29), while the 74 kDa

438 and 32 kDa bands are suggested to be the A and D subunits, respectively, of the V_1

439 complex (36). The higher abundance of these V_1 complex protein subunits in the

440 cytosolic fraction and lower abundance in membrane fraction in *Aedae*CAPA-1-treated

441 MTs provides novel evidence of hormonally-regulated V_1 dissociation from the

442 holoenzyme in *A. aegypti* MTs. This was further confirmed with V_1 staining found both

443 in the apical membrane and cytosol of the MTs treated with DH_{31} and *Aedae*CAPA-1 in

444 contrast to the strict co-localization of the V_1 and V_o complex in the apical membrane of

445 MTs treated with DH_{31} alone. In unstimulated *A. aegypti* MTs, 40-73% of the V_1 subunits

446 were found to be membrane associated, with reassembly of the V_1V_o complex observed

447 upon stimulation with cAMP analogues (38). Although studies have revealed that

448 hormonal regulation can activate the assembly of the holoenzyme, the signaling

449 mechanisms achieving this control are unclear. In this study, the data provides evidence

450 of VA assembly in DH_{31}-treated MTs, with V_1 complex protein subunit enrichment found

451 in the membrane fractions, confirming the crucial role of the VA in DH_{31}-stimulated

452 secretion. Studies in *A. aegypti* have demonstrated the involvement of PKA in the

453 activation and assembly of the VA upon natriuretic hormone (i.e. DH_{31}) stimulation and

indicate the phosphorylation of the VA subunits by PKA in the MTs (38). These studies

indicate a regulatory role of PKA in VA assembly and its activation that may be

independent or in addition to phosphorylation (38). In line with these earlier observations,

the current results indicate PKA is critical for DH_{31}-stimulated fluid secretion by MTs

whereas DH_{44}-stimulated diuresis was found to be PKA-independent.

Together, these results indicate that *Aedae*CAPA-1 binding to its cognate receptor

in principal cells of the MTs (6), targets the NOS/cGMP/PKG pathway (4, 6) to inhibit

DH_{31}-mediated elevation of cAMP (23, 59), which blocks PKA-activated VA association

and prevents protons from being pumped across the apical membrane, resulting in a more

alkaline lumen. Our study provides novel evidence that the anti-diuretic activity of CAPA

is mediated through the dissociation of the VA holoenzyme involving the removal of the

V_1 complex from the apical membrane, hindering luminal flux of protons that in turn

starves cation/H^+ exchange, which ultimately reduces fluid secretion (Fig. 8). In *R.*

prolixus MTs, the physiological roles of cGMP and cAMP were examined (65)

suggesting cGMP inhibits fluid secretion by activating a phosphodiesterase (PDE) that

degrades cAMP elevated following 5HT and diuretic hormone stimulation of MTs.

Indeed, the current results demonstrated the addition of cAMP reversed the inhibitory

effects of cGMP, while the addition of cGMP reduced the stimulatory response of cAMP,

supporting that these two cyclic nucleotides facilitate two opposing physiological roles in

the MTs of adult *A. aegypti*. The data herein reveals cGMP levels increase in MTs treated

with CAPA alone or in combination with DH_{31} while cAMP levels decrease in MTs

treated with CAPA in combination with DH_{31} compared to tubules stimulated with DH_{31}

alone, which upholds the roles of cAMP and cGMP in diuretic and anti-diuretic signaling

pathways, respecitively. Interestingly, mid-nanomolar concentrations of DH_{44} also led to

increased levels of cAMP, with levels unchanging in response to *Aedae*CAPA-1, raising

doubt regarding the involvement of a PDE. Treatment of a PKA inhibitor, KT5720,

abolished DH_{31}-stimulated secretion but had no effect on DH_{44}-mediated stimulation. It is

well established that the effects of cAMP are mediated by activation of cAMP-dependent

protein kinase (PKA), a major cAMP target, followed by phosphorylation of target

proteins (66). More recently, in *D. melanogaster* MTs, two distinct cAMP pathways have

been elucidated to sustain fluid secretion; a PKA-dependent pathway, shown to regulate

basal fluid secretion in principal cells; and a PKA-independent pathway, specifically a

stimulatory principal EPAC (exchange proteins directly activated by cAMP) pathway,

stimulating fluid secretion above basal levels (67). Future studies should examine the

potential DH_{44}-stimulated PKA-independent pathway leading to secretion in *A. aegypti*

MTs.

In summary, our study highlights a novel target in the anti-diuretic signaling

pathway of adult female *A. aegypti* MTs, emphasizing the intricate and precise regulatory

mechanism of anti-diuresis. Although a plethora studies have investigated the process of

hydromineral balance in terrestrial insects from a diuretic perspective (1, 10, 23, 68–70),

these current findings advance our understanding of anti-diuretic hormone control while

providing further evidence of a previously elusive endocrine regulatory mechanism of the

VA in mosquitoes (Fig. 8). Given that many terrestrial insects are recognized as

agricultural pests or disease vectors, further investigating the complex regulation of their

ionic and osmotic balance may aid in lessening their burden on human health and

prosperity through development of improved management strategies that, at least in part,

impede their neuroendocrine control of hydromineral homeostasis.

Materials and Methods

Animal rearing

Eggs of *Aedes aegypti* (Liverpool strain) were collected from an established

laboratory colony described previously (4, 71). All mosquitoes were raised under a 12:12

light:dark cycle. Non-blood fed female insects (three-six days post-eclosion) were used

for bioassays, dissected under physiological saline (*Aedes* saline) adapted from (59) that

contained (in $mmol^{-1}$): 150 NaCl, 25 HEPES, 3.4 KCl, 7.5 NaOH, 1.8 $NaHCO_3$, 1

$MgSO_4$, 1.7 $CaCl_2$, and 5 glucose, and titrated to pH 7.1.

MT fluid secretion assay

In order to determine fluid secretion rates, modified Ramsay assays were

performed as described previously (4, 72). Female adults (3-6 day old) were dissected

under physiological *Aedes* saline prepared as described above, and MTs were removed

and placed in a Sylgard-lined Petri dish containing 20 μL bathing droplets (1:1 mixture of

Schneider's Insect Medium (Sigma-Aldrich): *Aedes* saline, immersed in hydrated mineral

oil to prevent evaporation. The proximal end of each tubule was wrapped around a

Minutien pin to allow for fluid secretion measurements. To investigate the effects of

second messengers, cAMP and cGMP, on fluid secretion rate, 10^{-4} M 8 bromo-cAMP

(cAMP) (23, 65) and 10^{-8} M 8 bromo-cGMP (cGMP) (4) (Sigma-Aldrich, Oakville, ON,

Canada) were used against unstimulated MTs. To test the effects of the pharmacological

blocker KT5720 (7) (protein kinase A (PKA) inhibitor), a dosage of 5 μmol l^{-1}

(manufacturer's recommended dose) was used against 25 nmol l^{-1} DH$_{31}$- and 10 nmol l^{-1}

DH$_{44}$-stimulated MTs.

Time course inhibition of bafilomycin

Dosage of bafilomycin A$_1$ was based on a dose-response analysis of bafilomycin

against DH$_{31}$-stimulated tubules (Fig. S1). In the interest of determining whether

bafilomycin inhibits the effects of the diuretic factors, dosages of 25 nmol l^{-1} *Drome*DH$_{31}$

(~84% identical to *Aedae*DH$_{31}$) (4, 23, 73), 100 nmol l^{-1} 5HT (4, 69, 74), and 10 nmol l^{-1}

*Rhopr*DH (CRF-related diuretic peptide, DH$_{44}$) (~48% overall identity; ~65% identity

and ~92% similarity within the highly-conserved N-terminal region to *Aedae*DH$_{44}$) (4,

56, 75, 76), were applied to the isolated MTs. Neurohormone receptors, including those

for 5HT, and the peptides DH$_{31}$, DH$_{44}$, and CAPA, are localized to the basolateral

membrane of principal cells (6, 61, 82), while the LK receptor is localized exclusively to

stellate cells (5). As a result, the effects of bafilomycin were tested on diuretics known to

act on the principal cells of the MTs. After incubating with the individual diuretics for 30

min (using the modified Ramsay assay), diuretic peptide was added alone (controls) or in

combination with bafilomycin (final concentration 10^{-5} M). The fluid secretion rate was

recorded every 10 min for a total of 80 min. In order to determine whether inhibition of

the VA was involved in the anti-diuretic activity of CAPA peptides on adult MTs, the

effects of 1 fmol l^{-1} *Aedae*CAPA-1 (4, 6) were investigated in combination with DH$_{31}$

and bafilomycin.

Measurement of pH of secreted fluid

The pH of secreted fluid was measured by using ion-selective microelectrodes

(ISME) pulled from glass capillaries (TW-150-4, World Precision Instruments, Sarasota,

Fl, USA) using a Sutter P-97 Flaming Brown pipette puller (Sutter Instruments, San

Raffael, CA, USA). Microelectrodes were silanized with *N,N*-

dimethyltrimethylsilylamine (Fluka, Buchs, Switzerland) pipetted onto the interior of a

glass dish inverted over the group of microelectrodes. A 1:2 ratio of number of

microelectrodes to amount of silanization solution (in µl) was used. The microelectrodes

were left to silanize for 75 min at 350°C and left to cool before use. The microelectrodes

were back-filled with a solution containing 100 mmol l^{-1} NaCl and 100 mmol l^{-1} sodium

citrate that was titrated to pH 6.0 and front-filled using Hydrogen Ionophore I – cocktail

B (Fluka, Buchs, Switzerland). The electrode tips were then coated with ~3.5% (w/v)

polyvinyl chloride (PVC) dissolved in tetrahydrofuran, to avoid displacement of the

ionophore cocktail when submerged in the paraffin oil (77). The H^+-selective

microelectrodes were calibrated in *Aedes* saline titrated to either pH 7.0 or pH 8.0.

Reference electrodes were prepared from glass capillaries (1B100F-4, World Precision

Instruments) using a pipette puller described above and were backfilled with 500 mmol l^{-1} KCl. Secreted droplet pH measurements were done immediately after collection to

prevent alkalization of the droplet due to carbon dioxide diffusion into the paraffin oil.

Microelectrodes and reference electrodes were connected to an electrometer through

silver chloride wires where voltage signals were recorded through a data acquisition

system (Picolog for Windows, version 5.25.3). In order to measure pH of the secreted

fluid, tubules were set up using the Ramsay assay, and pH measurements were recorded

every 10 min for a total of 60 min.

NKA and VA activities

The Na^+/K^+-ATPase (NKA) and VA activity in the MTs was determined using a

modified 96-well microplate method (78, 79), which relies on the enzymatic coupling of

ouabain- or bafilomycin-sensitive hydrolysis of ATP to the oxidation of reduced

nicotinamide adenine dinucleotide (NADH). The microplate spectrophotometer is

therefore able to directly measure the disappearance of NADH. Adult female MTs (three

to six day old) were dissected and incubated for 30 min in *Aedes* saline, diuretic (DH_{31},

5HT, or DH_{44}) alone or combined with *Aedae*CAPA-1. Following 30-min incubation,

MTs were collected into 1.5 mL microcentrifuge tubes (40-50 sets of MTs per tube =

200-250 MTs per treatment), flash frozen in liquid nitrogen and stored at -80°C. To

investigate the effects of the pharmacological blockers, a nitric oxide synthase (NOS)

inhibitor, N_ω-Nitro-L-arginine methyl ester hydrochloride (L-NAME), and protein kinase

G (PKG) inhibitor, KT5823, were used against DH_{31}-stimulated MTs treated with

*Aedae*CAPA-1 or 10^{-8} M cGMP. Dosages of 2 μmol l^{-1} L-NAME (manufacturer's

recommended dose) and 5 μmol l^{-1} KT582336 were applied to the MTs (4, 6, 7).

Tissues were thawed on ice and 150 μL of homogenizing buffer (four parts of SEI

with composition (in mmol l^{-1}): 150 sucrose, 10 EDTA, and 50 imidazole; pH 7.3 and

one part of SEID with composition: 0.5% of sodium deoxycholic acid in SEI) and MTs

were then sonicated on ice for 10 sec (two pulses of 5 sec) and subsequently centrifuged

at 10,000 x *g* for 10 min at 4°C. The supernatant was transferred into a fresh

microcentrifuge tube and stored on ice.

Prior to the assay, three solutions were prepared (Solution A, B, and C) and all

stored on ice. Solution A contained 4 units mL^{-1} lactate dehydrogenase (LDH), 5 units

mL^{-1} pyruvate kinase (PK), 50 mmol l^{-1} imidazole, 2.8 mmol l^{-1} phosphoenolpyruvate

(PEP), 0.22 mmol l^{-1} ATP, and 50 mmol l^{-1} NADH, pH 7.5. The solution was

subsequently mixed with a salt solution in a 4:1 ratio (salt solution composition (in mmol

l^{-1}): 189 NaCl, 10.5 MgCl$_2$, 42 KCl, and 50 imidazole, pH 7.5. Working solution B

consisted of solution A with 5 mmol l^{-1} ouabain and solution C consisted of solution A

with 10 μmol l^{-1} bafilomycin. The concentrations of ouabain and bafilomycin were based

on previous studies (78). To ensure the batch of assay mixture (Solution A) was effective,

an adenosine diphosphate (ADP) standard curve was run. ADP standards were prepared

as follows: 0 nmol 10μL^{-1} (200μL of 50mmol l^{-1} imidazole buffer (IB), pH 7.5); 5 nmol

10μL^{-1} (25μL of 4mmol l^{-1} ADP stock and 175μL of IB); 10 nmol 10uL^{-1} (50μL of

4mmol l^{-1} ADP stock/ 150 μL of IB); 20 nmol 10 μL^{-1} (100μL of 4mmol l^{-1} ADP

stock/100μL of IB); 40 nmol 10 μL^{-1} (40 mmol l^{-1} ADP stock). The standards were added

to a 96-well polystyrene microplate in duplicates of 10 μL per well, followed by the

addition of 200 μL of solution A. The plate was placed in a Thermo Multiscan Spectrum

microplate spectrophotometer set at 25°C and a linear rate of NADH disappearance was

measured at 340 nm. The absorbance spectra were recorded and analyzed using the

Multiscan Spectrum data acquisition system with SkanIt version 2.2 software. The ADP

standards (0 to 40 nmoles well^{-1}) should yield an optical density (OD) between 0.9 and

0.2 OD units, while the slope of the curve should result in -0.012 to -0.014 OD nmol

ADP^{-1}. Homogenized MT samples were thawed and added to the microplate (kept on ice)

in six replicates of 10 μL per well. Next, two wells per sample were filled with 200 μL of

working solution A, two wells with 200 μL of working solution B and two wells with 200

μL of working solution C (Fig. 4B). The microplate was quickly placed in the microplate

spectrophotometer and the decrease in NADH absorbance was measured for 30 min at

340 nm. NKA and VA activity was calculated using the following equation:

618   NKA or VA activity = $(((\Delta ATPase/S)/[P]) \times 60$ (min),

where ΔATPase is the difference in ATP hydrolysis in the absence and presence of

ouabain or bafilomycin, S is the slope of the ADP standard curve, [P] is the protein

concentration of the sample. Protein was quantified using a Bradford assay (Sigma-

Aldrich Canada, Ltd.,) according to manufacturer's guidelines with bovine serum

albumin (Bioshop Canada Inc., Burlington, ON, Canada) as a standard. Final activity was

expressed as micromoles of ADP per milligram of protein per hour.

Protein processing and western blot analyses

MTs were isolated under physiological saline from 40-50 female *A. aegypti* for

each biological replicate (defined as n = 1) and incubated for 60 min in the following

three treatments: *Aedes* saline, 25 nmol l^{-1} DH$_{31}$, or 25 nmol l^{-1} DH$_{31}$ + 1 fmol l^{-1}

*Aedae*CAPA-1. Following the incubation, tissues were stored at -80ºC until processing.

To separate the membrane and cytosolic proteins, a membrane protein extraction kit was

used (ThermoFisher Scientific) following recommended guidelines for soft tissue with

minor modifications including 200 μL of permeabilization and solubilization buffer and a

1:200 protease inhibitor cocktail (Sigma Aldrich) in both buffers. Final protein

concentrations were calculated by Bradford assay (Sigma-Aldrich Canada, Ltd.,)

according to manufacturer's guidelines with bovine serum albumin (BioRad

Laboratories) as a standard and quantified using an A_o Absorbance Microplate Reader

(Azure Biosystems) at 595 nm.

Protein samples were denatured by heating for 5 min at 100ºC with 6X loading

buffer (225 mmol l^{-1} Tris-HCl pH 6.8, 3.5% (w/v) SDS, 35% glycerol, 12.5% (v/v) ß-

mercaptoethanol and 0.01% (w/v) Bromophenol blue). Into each lane, 5 μg of protein

was loaded onto a 4% stacking and 12% resolving sodium dodecyl sulphate

polyacrylamide gel electrophoresis (SDS-PAGE) gel. Protein samples were migrated

initially at 80 V for 30 min and subsequently at 110 V for 90 min before being transferred

onto a polyvinylidene difluoride (PVDF) membrane using a wet transfer method at 100 V

for 60 min in a cold transfer buffer. Following transfer, PVDF membranes were blocked

with 5% skim milk powder in Tris-buffered saline (TBS-T; 9.9 mmol l^{-1} Tris, 0.15 mmol

l^{-1} NaCl, 0.1 mmol l^{-1} Tween-20, 0.1 mmol l^{-1} NP-40 pH 7.4) for 60 min at RT, and

incubated on a rocking platform overnight at 4ºC with a guinea pig polyclonal anti-VA

(Ab 353-2 against the V_1 complex of the VA (13), a kind gift from Profs. Wieczorek and

Tiburcy, University of Osnabruck, Germany, used at a 1:2000 dilution in 5% skim milk

in TBS-T. The next day, PVDF membranes were washed for 60 min in TBST-T,

changing the wash buffer every 15 min. Immunoblots were then incubated with a goat

anti-guinea pig HRP conjugated secondary antibody (1:2500 in 5% skim milk in TBS-T)

(Life Technologies, Burlington, ON) for 60 min at RT and subsequently washed three

times for 15 min with TBS-T. Lastly, blots were incubated with the Clarity Western ECL

substrate and images were acquired using a ChemiDoc MP Imaging System (Bio-Rad).

Molecular weight measurements were performed using Image Lab 5.0 software (Bio-

Rad). PVDF membranes were then probed with Coomassie brilliant blue, since total

protein normalization is now considered the benchmark method for quantitative analysis

of western blot data (80, 81) and has been used in studies involving *A. aegypti* protein

normalization (82). ImageJ software (NIH, USA) was used to quantify protein

abundance. (Fig. 3 and S5 show a saturated blot of the 56kDa band to ensure

visualization of the 74kDa and 32kDa band, however protein quantification was

measured using a pre-saturated blot).

To confirm successful separation of membranes from cytosol using the membrane

protein extraction kit, saline-incubated MTs were incubated in either anti-beta-tubulin

(cytosolic marker, 1:5000) or -AaAQP1 affinity purified rabbit polyclonal antibody

(generous gift from Dr. Andrew Donini, York University, Canada) (83) (membrane

marker, 1:1000). Blots were then incubated with a goat anti-mouse (for beta-tubulin) or

goat anti-rabbit (for AQP1) HRP-conjugated secondary antibody (1:5000 in 5% skim

milk in TBS-T) (Life Technologies, Burlington, ON) for 60 min at RT (beta-tubulin).

Immunolocalization of VA complexes in MTs

Immunohistochemistry of the MTs localizing the VA complexes was conducted

following a previously published protocol (84). Adult female MTs (three to six day old)

were dissected out in *Aedes* saline and incubated following similar conditions described

in the western blot section above. After the incubation, the MTs were immersed in

Bouin's solution and fixed for two hours in small glass vials. To test *in vivo* changes of

the VA complexes, five to six day old females were allowed to blood-feed for 20 min

(71), after which female mosquitoes were isolated at 10 min, 30 min, and 3 hr post blood-

meal. Similarly aged, non blood-fed (sucrose-fed) females were isolated as controls.

Following the bloodmeal, whole body females were immersed in Bouin's solution and

fixed for three hours in small glass vials. Tissues/whole body females were then rinsed

three times and stored in 70% ethanol at 4ºC until further processing. Fixed samples were

dehydrated through a series of ethanol washes: 70% ethanol for 30 min, 95% ethanol for

30 min, and 100% ethanol three times for 30 min. The samples were cleared with xylene

(ethanol:xylene for 30 min then 100% xylene three times for 30 min), and infiltrated in

Paraplast Plus Tissue Embedding Medium (Oxford Worldwide, LLC, Memphis, USA) at

xylene:paraffin wax for 60 min at 60ºC, then rinsed in pure paraffin wax twice for 1 hour

for 60ºC. Following the last rinse, the samples were embedded in the paraffin wax and

left to solidify at 4ºC until further processing. Sections (5 µm) were cut using a Leica RM

2125RT manual rotary microtome (Leica Microsystems Inc., Richmond Hill, Canada)

and slides were incubated overnight on a slide warmer at 45ºC.

The following day, sections were deparaffinized with xylene (two rinses for 5 min

each), and rehydrated via a descending series of ethanol washes (100% ethanol twice for

2 min, 95% ethanol for 2 min, 70% ethanol for 2 min, 50% ethanol for 2 min) and finally

in distilled water for 20 min. Next, sections were subjected to a heat-induced epitope

retrieval (HIER) by immersing slides in a sodium citrate buffer (10 nmol l^{-1}, pH 6.0) and

heating both slides and solution in a microwave oven for 4 min. The solution and slides

were allowed to cool for 20 min, reheated for 2 min, and left to stand at room temperature

(RT) for 15 min. Slides were then washed three times in phosphate-buffered saline (PBS)

pH 7.4, 0.4% Kodak Photo-Flo 200 in PBS (PBS/PF, 10 min), 0.05% Triton X-100 in

PBS (PBS/TX, 10 min), and 10% antibody dilution buffer (ADB; 10% goat serum, 3%

BSA and 0. 05% Triton X-100 in PBS) in PBS (PBS/ADB, 10 min). Slides were

incubated overnight at RT with a guinea pig polyclonal anti-V_1 (Ab 353-2 against the V_1

complex, identical antibody used in western blot analyses, at 1:5000 dilution in ADB) in

combination with a 1:100 mouse polyclonal anti-ATP6V0A1 antibody for V_o (Abnova,

Taipei, Taiwan).

Following overnight primary antibody incubation, slides were washed briefly in

distilled water, with sequential washes with PBS/PF, PBS/TX, and PBS/ADB for 10 min

each as described above. A goat anti-guinea pig antibody (for V_1 detection) conjugated to

AlexaFluor 488 (1:500 in ADB, Jackson Immunoresearch) and sheep anti-mouse

antibody (for V_0 detection) conjugated to AlexaFluor 594 (1:500 in ADB, Jackson

Immunoresearch) were applied to visualize the VA complexes. The slides were left to

incubate in the secondary antibody for 60 min at RT. For negative controls, slides were

processed as described above with primary antibodies omitted. Slides were then washed

in distilled water/0.4% PF three times for 1 min each and finally in distilled water for 1

min. Slides were air dried for 30 min and mounted using ProLong^{TM} Gold antifade

reagent with DAPI (Life Technologies, Burlington, ON, Canada). Fluorescence images

were captured using an Olympus IX81 inverted fluorescent microscope (Olympus

Canada, Richmond Hill, ON, Canada).

cGMP and cAMP Measurements

A competitive cGMP ELISA kit (Cell Signaling Technology, #4360) and cAMP

ELISA kit (Cell Signaling Technology, #4339) were used to measure the effect of DH_{31},

DH$_{44}$ and *Aedae*CAPA-1 on cGMP and cAMP levels in the MTs. Adult MTs were

isolated under physiological saline from 50 female *A. aegypti* for each biological

replicate (defined as n = 1). To prevent cGMP degradation, tubules were incubated first

with a phosphodiesterase inhibitor, 0.1 mmol l^{-1} zaprinast, for 10 min (85) or 0.5 mmol l^{-1}

3-isobutyl-1-methylxanthine (IBMX) for 10 min (23) before any of the experimental

treatments, specifically *Aedes* saline, 25 nmol^{-1} DH$_{31}$, 10 nmol l^{-1} DH$_{44}$, 1 fmol^{-1}

*Aedae*CAPA-1, 25 nmol^{-1} DH$_{31}$ + 1 fmol^{-1} *Aedae*CAPA-1, or 10 nmol^{-1} DH$_{44}$ + 1 fmol^{-1}

*Aedae*CAPA-1 for a further 20 min. After incubation was complete, tissues were stored at

-80ºC until processing. To measure cGMP and cAMP concentrations, frozen tubule

samples were thawed on ice and 125 µL of 1X cell lysis buffer (CLB, #9803) was added

to each tube (1 mmol^{-1} phenylmethylsulfonyl fluoride (PMSF) was added to 1X CLB

fresh each time). Tissue samples were kept on ice for 10 min, sonicated for 10 s (similar

conditions as described for NKA/VA activity assay), centrifuged for 3 min at 10,000 rpm,

and the supernatant was isolated and kept on ice. Using a commercial 96-well microtitre

plate precoated with either a cGMP or cAMP rabbit monoclonal antibody, the cGMP-

HRP (or cAMP-HRP) conjugate was added in triplicate wells with 50 µL per well. This

was followed by the addition of 50 µL of either tubule samples or cGMP (or cAMP)

standards ranging from 100 nmol^{-1} to 0.25 nmol^{-1}. The plate was covered and incubated

at RT for 3 h on a horizontal orbital plate shaker. Following incubation, plate contents

were discarded, and wells were washed four times with 200 µL/well of 1X wash buffer.

Next, 100 µL of 3,3',5,5'-tetramethylbenzidine (TMB) substrate was added to each well,

and the plate was covered and kept for 10 min at RT. The enzymatic reaction was

quenched by adding 100 µL of 2 mmol^{-1} HCl and absorbance read at 450 nm using a

Synergy 2 Microplate Reader (Biotek).

Statistical analyses

Data was compiled using Microsoft Excel and transferred to Graphpad Prism

software v.7 to create figures and conduct all statistical analyses. Data was analyzed

accordingly using a one-way or two-way ANOVA and a Bonferroni post-test, or

Student's t-test, with differences between treatments considered significant if $p < 0.05$.

Acknowledgments

The authors sincerely thank Prof. Helmut Wieczorek and Prof. Felix Tiburcy (University

of Osnabrück, Germany) for providing the V_1 antibody and Prof. Andrew Donini (York

University, Canada) for providing the *A. aegypti* AQP1 antibody that were used in this

study. The authors are also grateful to Prof. Ian Orchard (University of Toronto

Mississauga, Canada) and Prof. Michael J. O'Donnell (McMaster University, Canada) for

providing synthetic peptides, *Rhopr*DH and *Drome*DH$_{31}$ respectively, used in this study.

Funding

This research was funded by: Natural Sciences and Engineering Research Council of

Canada (NSERC) Discovery Grant (JPP)**,** Ontario Ministry of Research Innovation Early

Researcher Award (JPP), and NSERC CGS-D (FS) and the Carswell Scholarships in the

Faculty of Science, York University (FS) along with a MITACS Globalink Research

Internship (MFV).

Data and materials availability

All data are available in the main text or the supplementary materials. Additional data

related to this paper may be requested from the authors.

Figures and Tables

Fig. 1. Effect of bafilomycin on fluid secretion rates and pH along with cyclic nucleotide

second messengers on adult *A. aegypti* MTs. Tubules were treated with either (A, E)

DH_{31} (B, F) 5HT or (C, G) DH_{44} and secreted droplets were measured at 10-min intervals

for 30 min. Immediately following measurement of the 30-min point fluid droplet (solid

arrow), MTs were treated with (A, E) DH_{31} (B, F) 5HT (C, G) or DH_{44} alone (blue

circles) or in combination with *Aedae*CAPA-1 (purple triangles) or bafilomycin (yellow

squares). (A-C). Significant differences between bafilomycin-treated and the

corresponding time point controls are denoted by an asterisk, as determined by a two-way

ANOVA and Bonferroni multiple comparison post-hoc test ($p<0.05$). Data represent the

mean ± standard error (n=12-23). (D) MTs were treated with DH_{31} alone or in

combination with *Aedae*CAPA-1, bafilomycin, or both *Aedae*CAPA-1 and bafilomycin

for 60 min. Bars labeled with different letters are significantly different from each other

(mean± SEM; one-way ANOVA with Bonferroni multiple comparison, $p<0.05$, n=7-17).

(E-G) Significant differences between *Aedae*CAPA-1- and bafilomycin-treated and the

corresponding time point controls are denoted by a double asterisk (mean± SEM; two-

way ANOVA with Bonferroni multiple comparison, $p<0.05$, n=14-34). (H, K) Fluid

secretion rates were measured at 10 min intervals initially over a 30 min time interval

(unstimulated, cAMP or cGMP) (solid arrow) and then over a second 30 min interval

after the addition of 10^{-4} M cAMP or 10^{-8} M cGMP. (H) Significant differences between

cAMP-treated MTs and the corresponding time point controls (or cGMP- treated MTs)

are denoted by a double asterisk (mean± SEM; two-way ANOVA with Bonferroni

multiple comparison, $p<0.05$, n=5-6). To monitor activity of *Aedae*CAPA-1 on cGMP (I)

and cAMP (J) production in stimulated *A. aegypti* MTs, tubules were incubated in

zaprinast (cGMP) or IBMX (cAMP) for 10 min before incubation with the following

treatments; *Aedes* saline, DH_{31}, *Aedae*CAPA-1 alone, or in combination with DH_{31} for 20

min before collection. Bars labeled with different letters are significantly different from

each other (mean± SEM; one-way ANOVA with Bonferroni multiple comparison,

$p<0.05$, n=50 sets of MTs for all treatments (n=3 per treatment)). (K) Significant

differences between cAMP-treated MTs (yellow, solid) and corresponding time point

after addition at 30 min (downward arrow) of cGMP (yellow, dash) are denoted by an

asterisk (similar with cGMP alone (purple, solid) and cAMP added at 30 min (purple,

dash)), mean± SEM; one-way ANOVA with Bonferroni multiple comparison, $p<0.05$,

n=5-9.

Fig. 2. Effect of *Aedae*CAPA-1 and NOS/PKG inhibitors on VA and NKA activity in

diuretic stimulated *A. aegypti* MTs. MTs were incubated in *Aedes* saline, diuretics (DH_{31},

5HT, and DH_{44}) alone or in combination with *Aedae*CAPA-1 for 30 min before collection

to measure (A) VA and (B) NKA activity. (C) MTs were treated with pharmacological

blockers, NOS inhibitor ($_L$-NAME) and PKG inhibitor (KT5923) in combination with

either *Aedae*CAPA-1 or cGMP. Bars labeled with different letters are significantly

different from each other (mean± SEM; one-way ANOVA with Bonferroni multiple

comparison, p<0.05). N.S. denotes no statistical significance. For each treatment, 50 sets

of MTs were collected with n=3 biological replicates per treatment.

Fig. 3. Membrane and cytosolic protein abundance of the V_1 complex in MTs of *A.*

aegypti. The MTs (n=40-50) were incubated in *Aedes* saline, DH_{31}, or DH_{31} +

*Aedae*CAPA-1 for one hour before collection. Protein abundance was measured in the

(A) 74 kDa band, A subunit (B) 56 kDa band, B subunit and (c) 32 kDa band, D subunit

of the V_1 complex. Individual band densities were normalized to total protein using

Coomassie staining, and graphed relative to saline-treated controls. Bars labeled with

different letters are significantly different from each other (mean± SEM; one-way

ANOVA with Bonferroni multiple comparison, p<0.05, n=3-4 replicates).

Fig. 4. Immunolocalization of the V_o and V_1 complexes in transverse sections of

stimulated *A. aegypti* MTs. Representative paraffin-embedded sections of *A. aegypti* MTs

incubated in either (A-C) *Aedes* saline alone, (D-F) DH_{31} and (G-I) DH_{31} + *Aedae*CAPA-

1 for 30 min. Panels (A,D,G) show V_o staining (red), (B,E,H) show V_1 staining (green),

and panels (C,F,I) show merged images with staining highly colocalized in DH_{31}

treatment but less evident in saline and *Aedae*CAPA-1 added treatments. Solid white

arrows denote apical VA staining, and empty arrows indicate cytosolic VA staining.

Where visible in sections, DAPI nuclear staining is shown in blue. Scale bar 100 μm, n =

4 biological replicates (SC = stellate cell).

Fig. 5. Immunolocalization of the V_o and V_1 complexes in blood-fed *A. aegypti* females.

Representative paraffin-embedded sections of whole body non-blood-fed females (A-C),

blood-fed females isolated (D-F) 10 min (G-I) 30 min and (J-L) 3 hr post blood-meal.

Panels (A,D,G,J) show V_o staining (red), (B,E,H,K) show V_1 staining (green), and panels

(C,F,I,L) show merged immunoreactive staining. DAPI nuclear staining is shown in blue.

Scale bar 100 μm, n = 4 biological replicates.

Fig. 6. Schematic diagram summarizing the signaling pathway of diuretic and anti-

diuretic control of adult *A. aegypti* MTs. The principal cells in *A. aegypti* MTs are

responsible for the transport of Na^+ and K^+ cations via secondary active transport,

energized by the V-type H^+-ATPase (VA), localized in the brush border of the apical

membrane. The movement of protons creates a gradient, driving the exchange of Na^+ and

K^+ through cation/H^+ antiporters (NHA). Neurohormone receptors for DH_{31}, 5HT, DH_{44},

and CAPA are localized to the basolateral membrane of the principal cells, while the

kinin receptor is localized exclusively in the stellate cells. The current results together

with previous data indicates that DH_{31} stimulates diuresis through activation and

assembly of the VA in the apical membrane, with no effect on the Na^+/K^+-ATPase

(NKA). The anti-diuretic effect of *Aedae*CAPA-1, facilitated by the NOS/cGMP/PKG

pathway, causes V_1 dissociation from the membrane, hindering activity, and thus

reducing fluid secretion. The biogenic amine, 5HT, has also been shown to stimulate

activation of the VA, however, to a lesser extent. DH_{44}-related peptide receptor activation

increases Ca^{2+} (as well as cAMP at higher doses) but its action was found to be

independent of PKA and VA. Lastly, it was shown earlier that cGMP inhibits kinin-

stimulated diuresis, suggesting an additional anti-diuretic factor may exist that acts

specifically on stellate cells.

Fig. S1. Dose response of bafilomycin on fluid secretion rates of DH_{31}-

stimulated MTs *in vitro* isolated from adult female *A. aegypti*. Doses of 10^{-4} M to 10^{-10} M

bafilomycin were applied to MTs together with DH_{31} for 60 min. Bars labeled with

different letters are significantly different from each other (mean± SEM; one-way

ANOVA with Bonferroni multiple comparison, $p<0.05$, n=7-13).

Fig. S2. Measurement of pH in fluid secreted by unstimulated MTs in vitro from adult

female *A. aegypti*. Secreted droplets were isolated and measured at 10-min intervals for

60 min using an ion-selective microelectrode. No significant differences were observed,

(mean± SEM; two-way ANOVA with Bonferroni multiple comparison, $p<0.05$, n=10);

ns denotes no statistical significance.

Fig. S3. Effect of bafilomycin and *Aedae*CAPA-1 on the secretion rate and pH of the

secreted fluid by unstimulated MTs *in vitro* from adult female *A. aegypti*. (A) Fluid

secretion rates were measured at 10 min intervals over a 30 min control period and then

after the addition (solid arrow) of 10^{-5} M bafilomycin (yellow), 1 fmol l^{-1} *Aedae*CAPA-1

(purple), or unstimulated alone (blue). (B) The pH of the secreted droplets was measured

at 10-min intervals for 60 min using an ion-selective microelectrode. No significant

differences were observed in the measurements (mean± SEM; two-way ANOVA with

Bonferroni multiple comparison, $p<0.05$, n=5-8); ns denotes no statistical significance

Fig S4. Intracellular levels of cAMP in DH_{44}-treated *A. aegypti* MTs. (A) To monitor

activity of *Aedae*CAPA-1 on cAMP levels in DH_{44}-stimulated *A. aegypti* MTs, tubules

were incubated in IBMX for 10 min prior to the following treatments; *Aedes* saline, DH_{44}

alone, or in combination with *Aedae*CAPA-1 for 20 min before collection. Bars labeled

with different letters are significantly different from each other (mean± SEM; one-way

ANOVA with Bonferroni multiple comparison, $p<0.05$. For each treatment, 50 sets of

MTs were isolated and combined (n=3 biological replicates per treatment). (B) Female

MTs were treated with either DH_{31} or DH_{44} alone or in combination with the PKA

inhibitor, KT5720 for 60 min to measure fluid secretion rate. Significant differences are

denoted by an asterisk (mean± SEM; one-way ANOVA with Bonferroni multiple

comparison, $p<0.05$, n=8-10).

Fig S5. Representative western blot of membrane and cytosolic protein fraction isolation

validated using established membrane (aquaporin-1 in *A. aegypti* MTs) and cytosolic

(beta-tubulin) protein markers. (A) Aquaporin-1 (AQP1) detection shown in membrane

fractions of *Aedes* saline-incubated MTs at 26 kDa, and (B) beta-tubulin in cytosolic

fractions of *Aedes* saline-incubated MTs at 55 kDa (n=40-50 MTs per replicate, n = 3

biological replicates).

Fig S6. Representative western blot image of membrane and cytosolic protein abundance

of the V_1 complex in *A. aegypti* MTs. The MTs (n=40-50) were incubated in *Aedes* saline

alone, DH_{31}, or DH_{31} + *Aedae*CAPA-1 for one hour before collection. Western blot

analysis revealed three protein bands, with calculated molecular masses of 74 kDa, 56

kDa, and 32 kDa (S = *Aedes* saline, D = DH_{31}, C = DH_{31} + *Aedae*CAPA-1).

Fig S7. Immunolocalization of the V_o and V_1 complexes in cross sections of stimulated

A. aegypti MTs. Representative paraffin-embedded sections of *A. aegypti* MTs

incubated in either (A-D) *Aedes* saline, (E-H) DH_{31} and (I-T) DH_{31} + *Aedae*CAPA-1.

Panels (A,E,I-K) show membrane-integrated V_o immunoreactive staining (red), (B,F,L-

N) demonstrate V_1 immunoreactive staining (green), (C,G,O-Q) show merged images of

V_o and V_1 staining, while (D,H,R-T) show brightfield images. Solid white arrowheads

denote apical VA staining, and empty arrowheads mark cytosolic VA staining. DAPI

nuclear staining is shown in blue. Scale bar 100 μm, n=3-4 biological replicates (PC =

principal cell).

References

1.   K. W. Beyenbach, Transport mechanisms of diuresis in Malpighian tubules of

insects. *J. Exp. Biol.* **206**, 3845–3856 (2003).

2.   T. M. Clark, T. J. Bradley, Malpighian tubules of larval *Aedes aegypti* are

hormonally stimulated by 5-hydroxytryptamine in response to increased salinity.

Arch. Insect Biochem. Physiol. **34**, 123–141 (1997).

3.   G. M. Coast, C. S. Garside, Neuropeptide control of fluid balance in insects, *Ann.*

N.Y. Acad. Sci. **1040**: 1–8 (2005).

4.   F. Sajadi, C. Curcuruto, A. Al Dhaheri, J.-P. V Paluzzi, Anti-diuretic action of a

CAPA neuropeptide against a subset of diuretic hormones in the disease vector,

Aedes aegypti. J. Exp. Biol. (2018).

5.   H. L. Lu, C. Kersch, P. V. Pietrantonio, The kinin receptor is expressed in the

Malpighian tubule stellate cells in the mosquito *Aedes aegypti* (L.): A new model

needed to explain ion transport? *Insect Biochem. Mol. Biol.* **41**, 135–140 (2011).

6.   F. Sajadi, *et al.*, CAPA neuropeptides and their receptor form an anti-diuretic

hormone signaling system in the human disease vector, *Aedes aegypti. Sci. Rep.* **10**

(2020).

7.   A. Ionescu, A. Donini, *Aedes*CAPA-PVK-1 displays diuretic and dose dependent

antidiuretic potential in the larval mosquito *Aedes aegypti* (Liverpool). *J. Insect*

Physiol. **58**, 1299–1306 (2012).

8.   R. Jurenka, The PRXamide neuropeptide signalling system. Conserved in animals,

Adv. Insect Phys. (Academic Press, 2015), pp. 123–170.

9.   S. A. Davies, *et al.*, Signaling by *Drosophila* capa neuropeptides. *Gen. Comp.*

Endocrinol. **188**, 60–66 (2012).

10. S. Terhzaz, *et al.*, Mechanism and function of *drosophila* capa GPCR: A

desiccation stress-responsive receptor with functional homology to human

neuromedinu receptor. *PLoS One* **7** (2012).

11. H. A. MacMillan, *et al.*, Anti-diuretic activity of a CAPA neuropeptide can

compromise *Drosophila* chill tolerance. *J. Exp. Biol.* **221**, jeb185884 (2018).

12. A. R. Rodan, M. Baum, C. L. Huang, The *drosophila* NKCC Ncc69 is required for

normal renal tubule function. *Am. J. Physiol. - Cell Physiol.* **303**, C883–C894

(2012).

13.   X. H. Weng, M. Huss, H. Wieczorek, K. W. Beyenbach, The V-type H^+-ATPase

in Malpighian tubules of *Aedes aegypti*: localization and activity. *J. Exp. Biol.* **206**,

2211–2219 (2003).

14.   H. Wieczorek, K. W. Beyenbach, M. Huss, O. Vitavska, Vacuolar-type proton

pumps in insect epithelia. *J. Exp. Biol.* **212**, 1611–1619 (2009).

15.   H. Wieczorek, The insect V-ATPase, a plasma membrane proton pump energizing

secondary active transport: molecular analysis of electrogenic potassium transport

in the tobacco hornworm midgut. *J. Exp. Biol.* **172**, 335–343 (1992).

16.   H. Wieczorek, M. Putzenlechnere, U. Klein, A Vacuolar-type proton pump

energizes K^+/H^+ antiport in an animal plasma membrane. *J. Biol. Chem.* **266**,

15340–15347 (1991).

17.   W. R. Harvey, S. H. P. Maddrell, W. H. Telfer, H. Wieczorek, H^+ V-ATPases

energize animal plasma membranes for secretion and absorption of ions and fluids.

Am. Zool. **38**, 426–441 (1998).

18.   Y. Li, *et al.*, RNA-Seq comparison of larval and adult malpighian tubules of the

yellow fever mosquito *Aedes aegypti* reveals life stage-specific changes in renal

function. *Front. Physiol.* **8**, 1–13 (2017).

19.   M. L. Patrick, K. Aimanova, H. R. Sanders, S. S. Gill, P-type Na^+/K^+-ATPase and

V-type H^+-ATPase expression patterns in the osmoregulatory organs of larval and

adult mosquito *Aedes aegypti*. *J. Exp. Biol.* **209**, 4638–4651 (2006).

20.   K. W. Beyenbach, T. L. Pannabecker, W. Nagel, Central role of the apical

membrane H^+-ATPase in electrogenesis and epithelial transport in Malpighian

tubules. *J. Exp. Biol.* **203**, 1459–1468 (2000).

21.   R. M. Hine, *et al.*, The excretion of NaCl and KCl loads in mosquitoes. *Am. J.*

Physiol. - Regul. Integr. Comp. Physiol. **307**, R837–R849 (2014).

22.   P. Dames, *et al.*, cAMP regulates plasma membrane vacuolar-type H^+-ATPase

assembly and activity in blowfly salivary glands. *Proc. Natl. Acad. Sci. U. S. A.*

103, 3926–3931 (2006).

23.   G. M. Coast, C. S. Garside, S. G. Webster, K. M. Schegg, D. A. Schooley,

Mosquito natriuretic peptide identified as a calcitonin-like diuretic hormone in

Anopheles gambiae (Giles). *J. Exp. Biol.* **208**, 3281–3291 (2005).

24.   K. Karas, P. Brauer, D. Petzel, Actin redistribution in mosquito Malpighian

tubules after a blood meal and cyclic AMP stimulation. *J. Insect Physiol.* **51**,

1041–1054 (2005).

25.   C. Cady, H. H. Hagedorn, Effects of putative diuretic factors on intracellular

second messenger levels in the Malpighian tubules of *Aedes aegypti*. *J. Insect*

Physiol. **45**, 327–337 (1999).

26.   D. H. Petzel, P. T. Pirotte, E. Van Kerkhove, Intracellular and luminal pH

measurements of malpighian tubules of the mosquito *Aedes aegypti*: the effects of

cAMP. *J. Insect Physiol.* **45**, 973–982 (1999).

27.   T. M. Clark, T. K. Hayes, K. W. Beyenbach, Dose-dependent effects of CRF-like

diuretic peptide on transcellular and paracellular transport pathways. *Am. J.*

Physiol. - Ren. Physiol. **274** (1998).

28.   K. W. Beyenbach, H. Wieczorek, The V-type H$^+$ ATPase: molecular structure and

function, physiological roles and regulation. *J. Exp. Biol.* **209**, 577–589 (2006).

29.   F. J. S. Novak, *et al.*, Primary structure of V-ATPase subunit B from *Manduca*

sexta midgut. BBA - Gene Struct. Expr. **1132**, 67–71 (1992).

30.   Y. X. Pan, H. H. Gu, J. Xu, G. E. Dean, *Saccharomyces cerevisiae* expression of

exogenous vacuolar ATPase subunits B. *BBA - Biomembr.* **1151**, 175–185 (1993).

31.   H. Wieczorek, D. Brown, S. Grinstein, J. Ehrenfeld, W. R. Harvey, Animal plasma

membrane energization by proton-motive V-ATPases. *BioEssays* **21**, 637–648

(1999).

32.   J. Zhang, Y. Feng, M. Forgac, Proton conduction and bafilomycin binding by the

V0 domain of the coated vesicle V-ATPase. *J. Biol. Chem.* **269**, 23518–23523

(1994).

33.   W. J. A. Dschida, B. J. Bowman, The vacuolar ATPase: sulfite stabilization and

the mechanism of nitrate inactivation. *J. Biol. Chem.* **270**, 1557–1563 (1995).

34.   J. Rein, M. Voss, W. Blenau, B. Walz, O. Baumann, Hormone-induced assembly

and activation of V-ATPase in blowfly salivary glands is mediated by protein

kinase A. *Am. J. Physiol. - Cell Physiol.* **294**, C56–C65 (2008).

35.   B. Zimmermann, P. Dames, B. Walz, O. Baumann, Distribution and serotonin-

1016  induced activation of vacuolar-type H$^+$-ATPase in the salivary glands of the

1017  blowfly *Calliphora vicina*. *J. Exp. Biol.* **206**, 1867–1876 (2003).

1018 36. H. Wieczorek, G. Grüber, W. R. Harvey, M. Huss, H. Merzendorfer, The plasma

1019  membrane H$^+$-V-ATPase from tobacco hornworm midgut. *J. Bioenerg. Biomembr.*

1020  **31**, 67–74 (1999).

1021 37. M. C. Quinlan, N. J. Tublitz, M. J. O'Donnell, Anti-diuresis in the blood-feeding

1022  insect *Rhodnius prolixus* Stal: The peptide CAP(2b) and cyclic GMP inhibit

1023  Malpighian tubule fluid secretion. *J. Exp. Biol.* **200**, 2363–2367 (1997).

1024 38. F. Tiburcy, K. W. Beyenbach, H. Wieczorek, Protein kinase A-dependent and -

1025  independent activation of the V-ATPase in Malpighian tubules of *Aedes aegypti*. *J.*

1026  *Exp. Biol.* **216**, 881–891 (2013).

1027 39. K. W. Beyenbach, P. M. Piermarini, Transcellular and paracellular pathways of

1028  transepithelial fluid secretion in Malpighian (renal) tubules of the yellow fever

1029  mosquito *Aedes aegypti*. *Acta Physiol. (Oxf)*. **202**, 387–407 (2011).

1030 40. K. W. Beyenbach, Energizing epithelial transport with the vacuolar H$^+$-ATPase.

1031  *News Physiol. Sci.* **16**, 145–151 (2001).

1032 41. T. Nishi, M. Forgac, The vacuolar (H$^+$)-ATPases - nature's most versatile proton

1033  pumps. *Nat. Rev. Mol. Cell Biol.* **3**, 94–103 (2002).

1034 42. N. Nelson, W. R. Harvey, Vacuolar and plasma membrane proton-

1035  adenosinetriphosphatases. *Physiol. Rev.* **79**, 361–385 (1999).

1036 43. K. W. Beyenbach, H. Skaer, J. A. T. Dow, The developmental, molecular, and

transport biology of malpighian tubules. *Annu. Rev. Entomol.* **55**, 351–374 (2010).

44.   P. M. Kane, Disassembly and reassembly of the yeast vacuolar H$^+$-ATPase in vivo.

J. Biol. Chem. **270**, 17025–17032 (1995).

45.   M. Voss, O. Vitavska, B. Walz, H. Wieczorek, O. Baumann, Stimulus-induced

phosphorylation of vacuolar H$^+$-ATPase by protein kinase A. *J. Biol. Chem.* **282**,

33735–33742 (2007).

46.   J. P. Sumner, *et al.*, Regulation of plasma membrane V-ATPase activity by

dissociation of peripheral subunits. *J. Biol. Chem.* **270**, 5649–5653 (1995).

47.   G. Coast, The endocrine control of salt balance in insects. *Gen. Comp. Endocrinol.*

152, 332–338 (2007).

48.   M. J. O'Donnell, J. H. Spring, Modes of control of insect Malpighian tubules:

synergism, antagonism, cooperation and autonomous regulation. *J. Insect Physiol.*

46, 107–117 (2000).

49.   J. C. Williams, H. H. Hagedorn, K. W. Beyenbach, Dynamic changes in flow rate

and composition of urine during the post-bloodmeal diuresis in *Aedes aegypti* (L.).

J. Comp. Physiol. **153**, 257–265 (1983).

50.   C. L. Jagge, P. V. Pietrantonio, Diuretic hormone 44 receptor in Malpighian

tubules of the mosquito *Aedes aegypti*: evidence for transcriptional regulation

paralleling urination. *Insect Mol. Biol.* **17**, 413–426 (2008).

51.   P. Gioino, B. G. Murray, J. P. Ianowski, Serotonin triggers cAMP and PKA-

mediated intracellular calcium waves in Malpighian tubules of *Rhodnius prolixus*.

Am. J. Physiol. - Regul. Integr. Comp. Physiol. **307**, R828–R836 (2014).

52.   F. Sajadi, J. P. V. Paluzzi, Hormonal regulation and functional role of the 'renal'

tubules in the disease vector, *Aedes aegypti. Vitamins and Hormones*, (Academic

Press, 2021), pp. 189–225.

53.   P. V. Pietrantonio, C. Jagge, C. McDowell, Cloning and expression analysis of a

5HT7-like serotonin receptor cDNA from mosquito *Aedes aegypti f*emale

excretory and respiratory systems. *Insect Mol. Biol.* **10**, 357–369 (2001).

54.   D. W. Lee, P. V. Pietrantonio, In vitro expression and pharmacology of the 5-HT7-

like receptor present in the mosquito *Aedes aegypti* tracheolar cells and hindgut-

associated nerves. *Insect Mol. Biol.* **12**, 561–569 (2003).

55.   A. Petrova, D. F. Moffett, Comprehensive immunolocalization studies of a

putative serotonin receptor from the alimentary canal of *Aedes aegypti* larvae

suggest its diverse roles in digestion and homeostasis. *PLoS One* **11**, e0146587

(2016).

56.   K. . Clark, T M, Hayes, T K., Holman, G.M., Beyenbach, The concentration-

dependence of CRF-like diuretic peptide: mechanisms of action. *J. Exp. Biol.* **201**,

1753–1762 (1998).

57.   D. H. Petzel, H. H. Hagedorn, K. W. Beyenbach, Preliminary isolation of

mosquito natriuretic factor. *Am. J. Physiol. - Regul. Integr. Comp. Physiol.* **18**

(1985).

58.   J. A. Dow, *et al.*, The malpighian tubules of *Drosophila melanogaster*: a novel

phenotype for studies of fluid secretion and its control. *J. Exp. Biol.* **197**, 421–428

(1994).

59.   D. H. Petzel, M. M. Berg, K. W. Beyenbach, Hormone-controlled cAMP-mediated

fluid secretion in yellow-fever mosquito. *Am. J. Physiol. - Regul. Integr. Comp.*

Physiol. **253** (1987).

60.   H. Kwon, H. L. Lu, M. T. Longnecker, P. V. Pietrantonio, Role in diuresis of a

calcitonin receptor (GPRCAL1) expressed in a distal-proximal gradient in renal

organs of the mosquito *Aedes aegypti* (L.). *PLoS One* **7**, e50374 (2012).

61.   W. Kang'ethe, K. G. Aimanova, A. K. Pullikuth, S. S. Gill, NHE8 mediates

amiloride-sensitive Na^+/H^+ exchange across mosquito Malpighian tubules and

catalyzes Na^+ and K^+ transport in reconstituted proteoliposomes. *Am. J. Physiol. -*

Ren. Physiol. **292**, 1501–1512 (2007).

62.   A. K. Pullikuth, K. Aimanova, W. Kang'ethe, H. R. Sanders, S. S. Gill, Molecular

characterization of sodium/proton exchanger 3 (NHE3) from the yellow fever

vector, *Aedes aegypti*. *J. Exp. Biol.* **209**, 3529–3544 (2006).

63.   P. M. Piermarini, D. Weihrauch, H. Meyer, M. Huss, K. W. Beyenbach, NHE8 is

an intracellular cation/H^+ exchanger in renal tubules of the yellow fever mosquito

Aedes aegypti. *Am. J. Physiol. - Ren. Physiol.* **296**, 730–750 (2009).

64.   H. Merzendorfer, M. Huss, R. Schmid, W. R. Harvey, H. Wieczorek, A novel

insect V-ATPase subunit M9.7 is glycosylated extensively. *J. Biol. Chem.* **274**,

17372–17378 (1999).

65.   M. C. Quinlan, M. J. O'Donnell, Anti-diuresis in the blood-feeding insect

1101   *Rhodnius prolixus* Stal: antagonistic actions of cAMP and cGMP and the role of

1102   organic acid transport. *J. Insect Physiol.* **44**, 561–568 (1998).

1103   66.   S. Seino, T. Shibasaki, PKA-dependent and PKA-independent pathways for

1104   cAMP-regulated exocytosis. *Physiol. Rev.* **85**, 1303–1342 (2005).

1105   67.   M. Efetova, *et al.*, Separate roles of PKA and EPAC in renal function unraveled by

1106   the optogenetic control of cAMP levels in vivo. *J. Cell Sci.* **126**, 778–788 (2013).

1107   68.   K. W. Beyenbach, A. Oviedo, D. J. Aneshansley, Malpighian tubules of *Aedes*

1108   *aegypti*: five tubules, one function. *J. Insect Physiol.* **39**, 639–648 (1993).

1109   69.   J. A. Veenstra, Effects of 5-hydroxytryptamine on the Malpighian tubules of *Aedes*

1110   *aegypti*. *J. Insect Physiol.* **34**, 299–304 (1988).

1111   70.   J. P. V. Paluzzi, W. Naikkhwah, M. J. O'Donnell, Natriuresis and diuretic

1112   hormone synergism in *R. prolixus* upper Malpighian tubules is inhibited by the

1113   anti-diuretic hormone, *Rhopr*CAPA-α2. *J. Insect Physiol.* **58**, 534–542 (2012).

1114   71.   D. A. Rocco, D. H. Kim, J. P. V. Paluzzi, Immunohistochemical mapping and

1115   transcript expression of the GPA2/GPB5 receptor in tissues of the adult mosquito,

1116   *Aedes aegypti*. *Cell Tissue Res.* **369**, 313–330 (2017).

1117   72.   A. Lajevardi, F. Sajadi, A. Donini, J. P. V. Paluzzi, Studying the activity of

1118   neuropeptides and other regulators of the excretory system in the adult mosquito.

1119   *J. Vis. Exp.* **2021** (2021).

1120   73.   M. Vanderveken, M. J. O'Donnell, Effects of diuretic hormone 31, *droso*kinin, and

1121   allatostatin A on transepithelial K^+ transport and contraction frequency in the

1122   midgut and hindgut of larval *Drosophila melanogaster*. *Arch. Insect Biochem.*

1123   *Physiol.* **85**, 76–93 (2014).

1124   74.   T. M. Clark, T. J. Bradley, Additive effects of 5-HT and diuretic peptide on *Aedes*

1125   malpighian tubule fluid secretion. *Comp. Biochem. Physiol. - A Mol. Integr.*

1126   *Physiol.* **119**, 599–605 (1998).

1127   75.   V. Te Brugge, J. P. Paluzzi, D. A. Schooley, I. Orchard, Identification of the

1128   elusive peptidergic diuretic hormone in the blood-feeding bug *Rhodnius prolixus*: a

1129   CRF-related peptide. *J. Exp. Biol.* **214**, 371–381 (2011).

1130   76.   H. R. Lee, M. Zandawala, A. B. Lange, I. Orchard, Isolation and characterization

1131   of the corticotropin-releasing factor-related diuretic hormone receptor in *Rhodnius*

1132   *prolixus*. *Cell. Signal.* **28**, 1152–1162 (2016).

1133   77.   M. R. Rheault, M. J. O'Donnell, Organic cation transport by Malpighian tubules of

1134   *Drosophila melanogaster*: application of two novel electrophysiological methods.

1135   *J. Exp. Biol.* **207**, 2173–2184 (2004).

1136   78.   S. Jonusaite, S. P. Kelly, A. Donini, The physiological response of larval

1137   *Chironomus riparius* (Meigen) to abrupt brackish water exposure. *J. Comp.*

1138   *Physiol. B Biochem. Syst. Environ. Physiol.* **181**, 343–352 (2011).

1139   79.   S. D. McCormick, Methods for nonlethal gill biopsy and measurement of Na^+, K^+-

1140   ATPase activity. *Can. J. Fish. Aquat. Sci.* **50**, 656–658 (1993).

80.   A. J. Fosang, R. J. Colbran, Transparency is the key to quality. *J. Biol. Chem.* **290**,

29692–29694 (2015).

81.   T. A. J. Butler, J. W. Paul, E. C. Chan, R. Smith, J. M. Tolosa, Misleading

westerns: common quantification mistakes in western blot densitometry and

proposed corrective measures. *Biomed Res. Int.* **2019** (2019).

82.   A. C. Durant, A. Donini, Ammonium transporter expression in sperm of the

disease vector *Aedes aegypti* mosquito influences male fertility. *Proc. Natl. Acad.*

Sci. U. S. A. **117**, 29712–29719 (2020).

83.   L. Misyura, E. G. Guardian, A. C. Durant, A. Donini, A comparison of aquaporin

expression in mosquito larvae (*Aedes aegypti*) that develop in hypo-osmotic

freshwater and iso-osmotic brackish water. *PLoS One* **15**, e0234892 (2020).

84.   H. Chasiotis, S. P. Kelly, Occludin immunolocalization and protein expression in

goldfish. *J. Exp. Biol.* **211**, 1524–1534 (2008).

85.   L. Kean, *et al.*, Two nitridergic peptides are encoded by the gene capability in

Drosophila melanogaster. *Am. J. Physiol. - Regul. Integr. Comp. Physiol.* **282**

(2002).

1157

| Non-fed | 10min PBF | 30min PBF | 3hr PBF |