1

Main Manuscript for

The V-type H[+]-ATPase is targeted in anti-diuretic hormone control of the Malpighian 'renal' tubules in *Aedes aegypti*

Farwa Sajadi[1] and Jean-Paul V. Paluzzi[1]*

1. Department of Biology, York University, 4700 Keele Street, Toronto, Ontario, M3J 1P3, Canada.

*Corresponding author: Jean-Paul V. Paluzzi

Email: paluzzi@yorku.ca

Author Contributions: J.-P.P. and F.S. designed research; F.S. performed research; F.S. and J.P.P. analyzed data; and J.-P.P. and F.S. wrote the manuscript.

Competing Interest Statement: The authors declare no competing interest.

Classification: Biological Sciences; Physiology

Keywords: excretory system, *Aedes aegypti*, mosquito, anti-diuresis, proton pump, disease vector

This PDF file includes:

Main Text
Figures 1 to 8; S1 to S3

Abstract

Like other insects, secretion by mosquito MTs is driven by the V-type H^+-ATPase (VA), which is densely packed in the apical membrane of principal cells. The anti-diuretic neurohormone CAPA inhibits tubule secretion stimulated by select diuretic hormones; however, the cellular effectors of this inhibitory signaling cascade remain unclear. Herein, we show that the VA inhibitor bafilomycin selectively inhibits serotonin (5HT)- and calcitonin-related diuretic hormone (DH_{31})-stimulated secretion. VA activity increases in DH_{31}-treated MTs, whereas CAPA abolishes this increase. A critical feature of VA activation involves the reversible association of the cytosolic (V_1) and membrane (V_o) complexes and results revealed higher V_1 protein abundance in membrane fractions of DH_{31}-treated MTs whereas CAPA significantly decreased V_1 protein abundance in membrane fractions while increasing it in cytosolic fractions. Immunolocalization of V_1 was observed strictly in the apical membrane of MTs treated with DH_{31} alone whereas immunoreactivity was dispersed following CAPA treatment. Elevated levels of cGMP in CAPA-treated MTs suggests activation of a phosphodiesterase (PDE), resulting in decreased levels of DH_{31}-stimulated cAMP. Therefore, CAPA inhibition of MTs involves reducing VA activity and promotes complex dissociation hindering secretion. Collectively, these findings reveal a key target necessary for hormone-mediated inhibition of MTs required in countering diuresis providing a deeper understanding of this critical physiological process.

Significance Statement

The V-type H^+-ATPase (VA) proton pump, provides the driving force for transepithelial ion and fluid secretion in insect Malpighian tubules (MTs). While studies have shown that diuretic stimulation activates various signaling pathways that culminate with increased VA activity, our understanding of anti-diuretic signaling and its potential regulation involving the VA remains rudimentary. Here we show that a CAPA neuropeptide acts through the NOS/cGMP/PKG pathway that inhibits VA activity, supporting the notion that anti-diuretic regulation is mediated through reversible dissociation of the VA complexes. Our results demonstrate a critical role of VA inhibition and trafficking necessary for anti-diuretic signaling, advancing our understanding of the complex neuroendocrine regulatory mechanisms of the MTs in an important human disease vector mosquito.

Main Text

Introduction

Insect post-prandial diuresis is under rigorous control by neuroendocrine factors (1), acting on the Malpighian 'renal' tubules (MTs) to regulate primary urine production. In the yellow fever mosquito, *Aedes aegypti,* several diuretics have been identified that regulate urine production including serotonin (5HT), calcitonin-related diuretic hormone (DH_{31}), corticotropin-releasing factor-related diuretic hormone (DH_{44}) and leucokinin-related (LK) diuretic hormone (2–5). Recently, an anti-diuretic peptidergic neurohormone, CAPA, was found to selectively inhibit DH_{31}- and 5HT-stimulated secretion of MTs (4, 6, 7). Insect CAPA neuropeptides are produced in the central

64 nervous system and are evolutionarily related to the vertebrate neuromedin U peptides

65 (8). In *Drosophila melanogaster* and other dipterans, CAPA peptides act through a

66 conserved nitridergic signaling pathway to stimulate diuresis by MTs (9, 10); however, in

67 both larval and adult *A. aegypti*, CAPA peptides inhibit fluid secretion through a

68 signaling cascade involving the NOS/cGMP/PKG pathway (6, 7). Despite this, the anti-

69 diuretic signaling mechanism and downstream cellular targets, such as the ion channels

70 and transporters, remain elusive.

71  In insect MTs, including *A. aegypti*, the bafilomycin-sensitive V-type H^+ ATPase

72 (VA), also known as the proton pump, functions as an electrogenic pump allowing the

73 transport of protons from the cytoplasm to the tubule lumen, thus generating a cell-

74 negative membrane voltage (11, 12). This membrane voltage can then drive secondary

75 transport processes such as the cation/H^+ exchanger or anion/H^+ cotransporter (13, 14).

76 Originally found in vacuolar membranes of animals and plants, the VA has since been

77 found to be essential in cell function in both invertebrates and vertebrates (15). In insects,

78 the VA is densely located in the apical brush border membrane of tubule principal cells

79 (11, 16), which is rich in mitochondria and endoplasmic reticulum, fueling the ATP-

80 consuming proton pump (1, 17). Previous studies have shown VA localization within the

81 apical membrane of principal cells along the entire length of the MTs (17), but absent in

82 stellate cells that express relatively higher levels of the P-type Na^+/K^+ ATPase (NKA)

83 (17). Due to stronger VA immunoreactivity observed in MTs (17), and greater ATPase

84 activity by electrophysiological assays (11, 18), the VA is categorized as serving mainly -

85 stimulated transport mechanisms, whereas the NKA serves basic cell housekeeping

86 functions when MTs are undergoing low unstimulated rates of secretion (11, 19).

Stimulation of distinct diuretic hormone receptors can activate various signaling

pathways, including elevation of cyclic AMP (cAMP) levels which is known to increase

VA activity and assembly in insects (20). *In A. aegypti,* DH_{31}, identified as the mosquito

natriuretic peptide (21), selectively activates transepithelial secretion of Na^+ in the MTs,

using cAMP as a second messenger (1), and upregulating the VA function to stimulate

fluid secretion (22). Similarly, 5HT-stimulated diuresis is also be mediated (at least in

part) through the cAMP second messenger pathway (23), activating protein kinase A

(PKA) to increase the transepithelial voltage of the basolateral membrane in tubule

principal cells (1, 24). Thus, due to its predominant role in fluid secretion, the VA could

be a likely target for both diuretic and anti-diuretic factors. The multi-subunit protein is

composed of two major functional domains, a membrane-integrated V_o complex with

properties of a proton channel (16, 25), and a cytoplasmic V_1 complex, which is the site

for ATP binding and hydrolysis (11). Although the proton channel of the VA can be

blocked pharmacologically by the macrolide antibiotic, bafilomycin (26), there are two

known instrinsic mechanisms for VA regulation: firstly, through oxidation of the cystine

residue on the A subunit of the V_1 complex, thus preventing ATP hydrolysis; secondly,

through reversible disassembly of the V_1 complex from the holoenzyme (11, 27). While

the role and regulation of the VA by diuretic hormones in insect MTs has been studied

(28–30), research examining anti-diuretic signaling mechanisms involving the VA remain

in their infancy.

The goal of this study was to identify the cellular targets necessary for CAPA-

mediated inhibition of tubule fluid secretion stimulated by select diuretic factors in adult

female *A. aegypti*. Our results provide evidence that CAPA neuropeptides inhibit fluid

secretion by VA complex dissociation, thus hindering VA function and activity that is

essential for driving rapid post-prandial diuresis.

Results

Bafilomycin inhibits DH$_{31}$- and 5HT-stimulated fluid secretion rate. To determine the

appropriate concentration of bafilomycin to test on adult *A. aegypti* MTs, dosages of 10^{-4}

M to 10^{-10} M were applied against DH$_{31}$-stimulated tubules (Fig. S1). Higher doses of

bafilomycin (10^{-4} M and 10^{-5} M) resulted in significant inhibition of fluid secretion rate,

with maximal inhibition leading to a five-fold decrease, observed with treatment of 10^{-5}

M bafilomycin. Next, to determine whether inhibiting the VA would decrease fluid

secretion rate stimulated by other diuretic hormones, including 5HT and DH$_{44}$, the effect

of 10^{-5} M bafilomycin on adult tubules stimulated with these diuretics was tested (Fig. 1).

Fluid secretion rates were measured over 30 min under control (stimulated) conditions,

and then at 10-min intervals in the presence of bafilomycin. Treatment of MTs with 10^{-5}

M bafilomycin against DH$_{31}$ led to a decrease of fluid secretion over the treatment

interval. Specifically, 30 min after treatment with bafilomycin, fluid secretion rate was

significantly reduced by over two-fold to 0.438±0.041 nL min^{-1}, compared to the DH$_{31}$

control, 0.941±0.077 nL min^{-1} (Fig. 1A). Similar results were seen with 5HT-stimulated

MTs; however, a decrease in fluid secretion was observed 30 min post-bafilomycin, with

a significant inhibition 40 min after treatment (0.522±0.072 nL min^{-1}, 5HT vs.

0.182±0.045 nL min^{-1}, 5HT + bafilomycin) (Fig. 1B). Distinct from DH$_{31}$ and 5HT-

stimulated tubules, DH$_{44}$-stimulated secretion was insensitive to bafilomycin treatment

(Fig. 1C). To confirm whether *Aedae*CAPA-1 anti-diuresis is mediated by VA inhibition,

adult female MTs were treated with either DH_{31} alone or in combination with

*Aedae*CAPA-1, bafilomycin, or both (Fig. 1D). MTs treated with either *Aedae*CAPA-1 or

bafilomycin resulted in a significant inhibition of DH_{31}-stimulated secretion,

(0.092 ± 0.027 nL min^{-1} and 0.155 ± 0.036 nL min^{-1} respectively) and similar inhibition of

fluid secretion was observed when both *Aedae*CAPA-1 and bafilomycin were applied

together (0.131 ± 0.043 nL min^{-1}) with no evidence of any additive inhibitory effects (Fig.

1D).

_Aedae_CAPA-1 and bafilomycin alkalinizes secreted fluid in DH_{31}- and 5HT-

stimulated MTs. The VA pumps protons from the cell into the tubule lumen thus

generating an electromotive potential (31, 32) and providing energy to drive the secretion

of cations via Na^{+}/H^{+} and/or K^{+}/H^{+} antiporters (11). An indirect way to measure whether

*Aedae*CAPA-1 and bafilomycin inhibits VA activity involves measuring the pH of the

secreted fluid from diuretic-stimulated MTs treated with *Aedae*CAPA-1 or bafilomycin

(Fig. 2). In DH_{31}-stimulated MTs treated with *Aedae*CAPA-1, there was an immediate

significantly higher pH (7.479 ± 0.030) at 40 min relative to control, increasing up to

7.73 ± 0.038 at 60 min (Fig. 2A). Similar, yet slightly higher pH levels were seen with

DH_{31}-stimulated MTs treated with bafilomycin, with significantly higher pH

(7.66 ± 0.064) relative to control at 40 min, increasing up to 7.855 ± 0.074 at 60 min.

Comparable to DH_{31}, after addition of *Aedae*CAPA-1 or bafilomycin, the pH of 5HT-

stimulated MTs was seen to significantly increase up to 7.75 ± 0.061 and 7.82 ± 0.083

respectively, at the 60 min mark (Fig. 2B). In contrast, unlike the effects observed with

DH_{31}- and 5HT-stimulated MTs, *Aedae*CAPA-1 or bafilomycin did not alter the pH of

the secreted fluid in DH_{44}-stimulated MTs (Fig. 2C). The pH increased from 7.4 to 7.9

during the 30-min DH_{44} incubation; however, pH did not change following the addition

of $Aedae$CAPA-1 or bafilomycin. Separately, we conducted measurements in

unstimulated tubules to verify pH in these small droplets did not drift over a time frame

consistent with our above experiments. Unstimulated MTs were allowed to secrete, and

the droplets were isolated and their pH was measured over the course of 60 min (Fig. S2).

Over this incubation period, no change was observed in the pH of secreted droplets from

unstimulated MTs, upholding the notion that the alkalinization of secreted fluid observed

following $Aedae$CAPA-1 (or bafilomycin) treatment of DH_{31}- and 5HT-stimulated MTs

is a result of VA inhibition.

$Aedae$CAPA-1 decreases VA activity in DH_{31}-stimulated MTs. In $Aedes$ MTs,

between 50% and 60% of the total ATPase activity can be attributed to a bafilomycin-

and nitrate-sensitive component that reflects the activity of the VA pump (11). The

remaining ATPase activity may be due to nucleotide cyclases, protein kinases, myosin,

DNA helicases, and other ATP-consuming processes such as the NKA (11). As such, to

determine whether CAPA inhibits VA and/or NKA function, female diuretic-stimulated

MTs were challenged with $Aedae$CAPA-1 to measure the resultant NKA and VA

activity. As expected given its role as the natriuretic diuretic hormone, adult female MTs

treated with DH_{31} resulted in a significant (> two-fold) increase of VA activity,

0.0329 ± 0.0007 µmoles ADP/µg protein/hour, compared to saline controls,

0.0151 ± 0.0021 µmoles ADP/µg protein/hour (Fig. 3A). Importantly, MTs incubated with

both DH_{31} and $Aedae$CAPA-1 had a significantly lower VA activity, 0.00617 ± 0.00347

μmoles ADP/μg protein/hour, resulting in activity levels indistinguishable from saline

controls. In contrast, neither 5HT nor DH_{44} influenced VA activity ($p>0.05$) when

compared with saline controls, while co-treatment with *Aedae*CAPA-1 also resulted in

indistinguishable VA activity. Similar VA activity levels were observed between 5HT

and DH_{44} (0.0255 ± 0.0078 and 0.0208 ± 0.0042 μmoles ADP/μg protein/hour) and with

co-application of *Aedae*CAPA-1 (0.0150 ± 0.0036 and 0.0154 ± 0.0070 μmoles ADP/μg

protein/hour). Unlike changes observed in VA activity following treatment with DH_{31},

diuretic-stimulation or *Aedae*CAPA-1 treatment did not perturb NKA activity, with levels

similar to that of unstimulated MTs (Fig. 3B).

***Aedae*CAPA-1 leads to increased cGMP levels in DH_{31}-treated MTs.** To further

understand the CAPA signaling pathway involving the second messenger, cGMP, we

sought to determine changes of cGMP levels in MTs incubated in DH_{31} alone or

combined with *Aedae*CAPA-1 (Fig. 3C). Treatment of MTs with DH_{31} alone had low

levels of cGMP, 10.91 ± 0.1096 $pmol^{-1}$/tubule, comparable to saline treated MTs,

10.94 ± 0.1108 $pmol^{-1}$/tubule. Treatment of MTs with *Aedae*CAPA-1 resulted in a

significant increase in cGMP levels compared to saline ($p=0.0543$) and DH_{31} ($p<0.05$)

incubated tubules, increasing to 11.39 ± 0.1012 $pmol^{-1}$/tubule (Fig. 3C). Similar results

were observed with MTs treated with both DH_{31} + *Aedae*CAPA-1, with significantly

increased cGMP levels of 11.39 ± 0.1232 $pmol^{-1}$/tubule (Fig. 3C).

_Aedae_CAPA-1 leads to VA holoenzyme dissociation in DH$_{31}$-treated MTs. The

reversible dissociation of the V$_1$ complex from the V$_o$ membrane-integrated complex is a

well-known mechanism for regulating VA transport activity (20, 28, 29, 32, 33). To

determine whether _Aedae_CAPA-1 influences VA complex dissociation, membrane and

cytosolic protein fractions were isolated from DH$_{31}$ and DH$_{31}$ + _Aedae_CAPA-1 incubated

MTs, and a polyclonal V$_1$ antibody (11) was used to measure protein abundance. Western

blot analysis revealed three protein bands, with calculated molecular masses of 74 kDa,

56 kDa, and 32 kDa (11, 34). The V$_1$ complex is composed of eight subunits (A-H),

which includes the A (~74kDa) and B subunit (~56Da) that form a ring structure and

allow for ATP binding (32). Additionally, studies have suggested that subunit D

(~32kDa) alongside subunit F constitute the central rotational stalk of the V$_1$ complex

(35). There was no difference in abundance observed for the 74 kDa band between

membrane and cytosolic fractions in saline and DH$_{31}$ treatments whereas DH$_{31}$ +

_Aedae_CAPA-1 incubated MTs had reduced 74 kDa band protein abundance in membrane

fractions relative to the cytosolic fraction (Fig. 4A-C). Abundance of the 56 kDa band

was higher in the membrane protein isolations of DH$_{31}$-incubated MTs, whereas DH$_{31}$ +

_Aedae_CAPA-1 incubated MTs had significantly lower abundance in membrane compared

to cytosolic fractions. (Fig. 4D-F). Finally, there was no difference in abundance of the

32 kDa band between cytosolic and membrane fractions in MTs treated with saline or

DH$_{31}$ alone; however, as observed for the 74 and 56 kDa band, DH$_{31}$ + _Aedae_CAPA-1

incubated MTs showed a significant decrease in the membrane protein abundance of the

32 kDa band compared to the cytosolic fractions (Fig. 4G-I). In summary, all three of the

VA immunoreactive bands (74 kDa, 56 kDa, and 32 kDa) showed significantly lower

abundance in membrane fractions relative to cytosolic protein fractions in DH_{31} +

*Aedae*CAPA-1 incubated MTs.

We next immunolocalized the membrane-integrated V_o and cytosolic V_1 complex

in the female *A. aegypti* MTs. Transverse sections of saline-incubated (control) MTs

demonstrated moderate enrichment of V_o (Fig. 5A, red), and V_1 (Fig 5B, green)

complexes in the apical membrane of principal cells (Fig. 6A-B, Fig. S3A-B).

Comparatively, DH_{31}-incubated MTs revealed intense co-localization of the V_o (Fig. 5D,

red), and V_1 (Fig 5E, green) complexes strictly to the apical membrane of principal cells

(Fig. 5F, solid arrow), identified by their larger nuclei (blue DAPI staining). To further

demonstrate the apical co-localization of the V_o/V_1 subunits, transverse and cross-

sectional images of MTs were co-labelled with both V_o and V_1 antibodies. In these tubule

preparations treated with DH_{31} alone, V_1 staining was strictly co-localized with V_o

staining (Fig. 5F, Fig. 6C, yellow, solid arrow) on the apical membrane of principal cells

(Fig. 6D, Fig. S3C-D), resulting in a significant increase in V_1 apical membrane

immunolocalization (Fig. 7A) with no change in the staining intensity in the cytosolic

portion of the MTs (Fig. 7B) compared to saline treated tubule preparations.

Interestingly, although V_o immunolocalization was restricted to the apical membrane, V_1

immunoreactivity was observed in both the apical membrane and cytosolic region in

DH_{31} + *Aedae*CAPA-1 co-treated MTs (Fig. 5G-I, Fig. 6E-J, Fig. S3E-J), with a

significantly higher immunoreactive staining intensity found in the cytosolic region of the

MTs compared to control (saline) or DH_{31}-treated MTs (Fig. 7A). Notably, the significant

increase in V_1 apical membrane immunolocalization following DH_{31} treatment was

abolished when MTs were co-incubated with *Aedae*CAPA-1 (Fig. 7B). Immunostaining

was absent in control preparations probed with only secondary antibodies (not shown)

confirming the specific detection of the VA complexes with each primary antibody.

Discussion

Eukaryotic V-ATPases are a multi-subunit protein composed of up to 14 different

polypeptides, which form two major structural complexes. The peripheral V_1 complex

(400-600 kDa), is invariably present in the cytoplasm and interacts with ATP, ADP, and

inorganic phosphate (36). The cytosolic V_1 complex consists of eight different subunits

(A-H): (1) a globular headpiece with three alternating subunits A and B forming a ring,

(2) a central rotor stalk with single copies of subunits D and F, and lastly, (3) a peripheral

stalk made up of subunits C, E, G, and H (36). The B subunit, shown to have high

sequence similarity amongst several species from fungi to mammals (37, 38), is a 56 kDa

polypeptide, and is one of the two sites (along with subunit A) (32) in the V_1 complex

that binds ATP. In contrast, the membrane-integrated V_o complex (150-350 kDa)

mediates the transport of H^+ across the membrane (36). The V_o complex is composed of

five different subunits (a-d), which collectively function in the proton translocation

pathway (11, 36).

The MTs of the *Aedes* mosquito are the main organs responsible for the secretion

of water and solutes, thereby contributing towards hydromineral homeostasis of the

animal (39). Active ion transport in *A. aegypti* MTs is accomplished mainly by the V-

ATPases (VA) densely localized in the apical brush-border membrane of principal cells,

that energize the apical and basolateral membrane as well as the paracellular pathway,

allowing for transepithelial secretion of NaCl, KCl, and other solutes (40). The VA is

widely distributed in both eukaryotes and prokaryotes and, due to their involvement in

basic cellular mechanisms, serve as critical components of virtually every eukaryotic cell

(36). In animal cells, V-ATPase molecules in the plasma membrane, especially on the

apical membrane of epithelial cells, contribute to extracellular acidification or

alkalization, intracellular pH homeostasis, or energize the plasma membrane for

secondary active transport (41, 42). In insect MTs, the VA plays a major role in fluid

secretion, thus serving as a primary target for both diuretic and, as our data demonstrates,

anti-diuretic hormonal regulation of the insect 'kidney'. Although the structure and

function of the VA has been elucidated in some detail (11, 20, 31, 32, 43–46), the

regulation of the proton pump remains unclear. Of the various regulatory mechanisms for

VA activity, the most studied is the reversible dissociation of the cytosolic V_1 complex

from the membrane-integrated V_o complex, first established in the midgut of the tobacco

hornworm, *Manduca sexta* and yeast, *Saccharomyces cervisiae* (45, 47). In this study, we

investigated the activity and regulation of the VA under both diuretic and anti-diuretic

hormone control of the adult female *A. aegypti* MTs. In particular, the current results

advance our knowledge of the anti-diuretic control of the *A. aegypti* MTs, revealing a

cellular mechanism for CAPA inhibition of the MTs by targeting the VA to block fluid

secretion stimulated by select diuretic factors. This includes inhibition of the DH_{31}-

related mosquito natriuretic peptide, which is critical for the post-haematophagy diuresis

that eliminates excess water and sodium originating from the bloodmeal-derived plasma.

In insects, water excretion is tightly regulated to maintain homeostasis of ions and

water (1, 48, 49). Female *A. aegypti* engorge a salt- and water-rich bloodmeal to obtain

the necessary nutrients and proteins for their eggs (1), with about 40% of the ingested

water eliminated in the first hour post feeding (50). The high rates of water excretion

along with the high rates of primary urine production post bloodmeal suggest a highly

coordinated and defined hormonal regulation of the signaling processes and downstream

cellular targets for ion and water transport (51). In *Aedes* MTs, fluid secretion increases

three-fold after stimulation with mosquito natriuretic peptide (identified as DH_{31}), via

cAMP as a second messenger (1), which activates V-ATPase-driven cation transport

processes (20, 29, 34). Herein we show that DH_{31}-stimulated secretion is inhibited by

bafilomycin, thought to block the proton channel of the VA (26). Moreover, the addition

of either bafilomycin or *Aedae*CAPA-1 caused alkalization of the secreted fluid,

indicating inhibition of the VA, which may lead to constrained entry of cations across the

apical membrane through proposed alkali cation/proton antiporter (13, 14). Thus, since

bafilomycin inhibits DH_{31}-stimulated secretion while the natriuretic activity of DH_{31}

remains unaffected in response to *Aedae*CAPA-1 (4), this supports the VA as a target in

the inhibition of fluid secretion. Consequently, the driving force of ion movement and

osmotically-obliged water is reduced, but select Na^+ channels and cotransporters remain

unaffected in the presence of *Aedae*CAPA-1. Similar results were seen in 5HT-stimulated

secretion, albeit slightly delayed. A study by Tiburcy et al. (34) showed that Ca^{2+}-

mediated diuresis does not require the assembly and activation of the VA (33). The

cAMP effect is implemented by protein kinase A (PKA), with inhibitors of PKA

abolishing the response of VA to stimulation (28). Although the endogenous 5HT

receptor expressed within the *A. aegypti* MTs necessary for diuretic activity remains

elusive, in the kissing bug, *Rhodnius prolixus*, both cAMP and Ca^{2+} have been shown to

initiate diuresis in response to 5HT (52), which could explain the delayed response of

*Aedae*CAPA-1 inhibition on 5HT-stimulated tubules as Ca^{2+}-mediated diuresis is

independent of the VA (34).

Interestingly, DH_{44}-mediated stimulation was observed to be independent of the

VA, as bafilomycin had no effect on the secretion rate or pH of the secreted fluid

following application with this CRF-related diuretic peptide. Previous studies have noted

that low nanomolar concentrations of a DH_{44}-related peptide were linked to the

stimulation of the paracellular pathway only (53), mediating this action through

intracellular Ca^{2+} as a second messenger (54). In contrast, high nanomolar concentrations

of a DH_{44}-related peptide were shown to influence both paracellular and transcellular

transport, increasing intracellular Ca^{2+} and cAMP (54). In female *A. aegypti*, post-

prandial diuresis is observed in three phases: peak phase (within 10 min of feeding), post-

peak phase (10 – 50 min), and late phase (50 – 120 min), with significant water loss

during the peak phase (50). Although haemolymph concentrations of diuretic peptides

have yet to be determined in mosquitoes, DH_{31} is immediately released into circulation

post bloodmeal, stimulating rapid secretion of Na^+ and excess water (21, 50). Since

DH_{44}-stimulated diuresis in *A. aegypti* involves non-selective transport Na^+ and K^+

cations (4), this supports a delayed release of this diuretic hormone post-feeding to

maintain production (albeit reduced) of primary urine whilst conserving Na^+ ions. This

coincides with the high DH_{44} receptor transcript abundance one hour post-feeding,

suggesting a role for DH_{44} during post-peak and late phases of diuresis (51), in which

anti-diuretic CAPA neurohormones may not be involved. Additionally, highest DH_{44}

receptor transcript abundance was observed 24 hours post-bloodmeal (51), which could

reflect the transport of nitrogenous wastes (55, 56), suggesting a crucial, yet relatively

elusive role for DH_{44} in mosquito tubule fluid secretion.

In unstimulated adult female *Aedes* MTs isolated *in vitro*, the VA exhibits

variable rates of enzyme activity, consistent with highly variable rates of secretion, as

found also in various other insect species (24, 57, 58). The VA is the main energizer in

MTs as 60% of total ATPase activity can be appointed to the VA (11), whereas the NKA,

with around 28% of ATPase activity, also plays a role in membrane energization,

denoting a more important role in the function of MTs than was previously assumed (17,

34, 39, 40). Indeed, the NKA inhibitor ouabain was shown to inhibit low rates of

secretion and alter cation levels in unstimulated adult female MTs in *A. aegypti* (19). In

R. prolixus, NKA activity in MTs is inhibited by 5HT via cAMP (59), whereas our results

showed no change in NKA activity with each tested diuretic (DH_{31}, 5HT, or DH_{44}). Thus,

the NKA may be involved in cell housekeeping functions and determining the Na^+/K^+

ratio of the secreted fluid by MTs (59). Here we show a significant two-fold increase in

VA activity in MTs treated with DH_{31} compared to the unstimulated MTs, with no

change in VA activity in 5HT and DH_{44}-treated MTs. Notably, *Aedae*CAPA-1

diminished the DH_{31}-stimulated VA activity, which corroborates the reduced fluid

secretion rate and alkalization of the secreted fluid. The present study examined the

effects of *Aedae*CAPA-1 on DH_{31}-stimulated VA activation for the first time in insects.

Stimulation of DH_{31} causes an increase in cAMP production, which activates Na^+

channels and the $Na^+/K^+/2Cl^-$ cotransporter in the basolateral membrane (60) and up-

regulates the VA activity (as shown here) for increased fluid secretion (22). The DH_{31}

receptor (*Aaeg*GPRCAL1) is expressed in a distal-proximal gradient in the MTs, with

greater expression in principal cells in which the VA in the apical membrane is highly

expressed (17, 61). The co-localization of the DH_{31} receptor, VA, and cation exchangers

(62–64) in the distal segment of the MTs, along with the CAPA receptor (6), collectively

supports the major roles DH_{31} and CAPA play in post-prandial diuresis and anti-diuresis,

respectively. In contrast to the marked changes in VA activity in response to diuretic and

anti-diuretic hormones, NKA activity remained unchanged in response to treatments

conducted herein. Similarly, previous studies have noted no measurable effects of 1

mmol ouabain on transepithelial resistance and voltage of isolated MTs (65); however,

approximately 50% of fluid secretion was inhibited by ouabain when treated for an

extended period (66). Thus, although NKA inhibition would have no instantaneous effect

on transepithelial transport, it may compromise transport after normal cell housekeeping

functions are impaired (11). In unstimulated adult female MTs, ouabain inhibited the low

rates of fluid secretion by ~33% by reducing secretion of K^+ and Cl^- into the lumen (19).

Further studies should examine the potential role of the NKA in diuretic and anti-diuretic

processes.

The reversible dissociation of the V_1 and V_o complexes is currently thought as a

universal regulatory mechanism of V-ATPases, appearing to be widely conserved from

yeast to animal cells (11, 20, 29, 67). Although previously shown with cAMP (22), it

remains unknown on whether the second messengers, Ca^{2+}, cGMP, and nitric oxide (NO)

affect the assembly/disassembly of the V_1V_o complexes in insect MTs. In this study, VA

protein abundance was confirmed by western blot analyses. The 56 kDa band represents

the B subunit (37), while the 74 kDa is suggested to be the A subunit and the 32 kDa

band is suggested to be the D subunit of the V_1 complex (30). The higher abundance of

the V_1 complex in the cytosolic fraction of the *Aedae*CAPA-1-treated MTs compared to

membrane fractions provide novel evidence of hormonally regulated V_1 dissociation from

the holoenzyme in *Aedes* MTs. This was further confirmed with V_1 staining found both in

the apical membrane and cytosol of the MTs treated with DH_{31} and *Aedae*CAPA-1 in

contrast to the strict co-localization of the V_1 and V_o complex in the apical membrane of

MTs treated alone with DH_{31}. In unstimulated *A. aegypti* MTs, 40-73% of the V_1 subunits

were found to be membrane associated, with reassembly of the V_1V_o complex observed

upon stimulation with cAMP analogues (34). Currently, studies are still unraveling the

molecular details of reversible V_1/V_o dissociation as a regulatory mechanism of VA

activity (68, 69). Although studies have revealed that hormonal regulation can activate

the assembly of the holoenzyme, the signaling mechanisms achieving this control are

unclear. Upon dissociation of the V_1 complex, subunit C is released from the complex

(45, 67, 70, 71), suggesting a central role in holoenzyme assembly/disassembly. Subunit

C binds to subunits E and G of the V_1 complex, and to subunit a of the V_o complex,

bridging the two complexes together (72) and serving as a good candidate for modulating

VA stability. In yeast, the cAMP/PKA pathway is proposed to be involved in the

reversible dissociation of the VA; however, it is still unknown on whether an active PKA

enhances the reassembly or prevents dissociation of the VA (12). In the *Calliphora*

blowfly, 5HT stimulates transepithelial secretion of a KCl-rich primary saliva (73). It was

shown that the reversible activation and assembly of the VA was dependent on the 5HT-

stimulated cAMP and activation of PKA (20, 28). The V_1 subunit C is one likely target of

PKA in the blowfly salivary glands, as it was shown to be phosphorylated in a cAMP-

dependent manner (46). In this study, the data provides direct evidence of VA assembly

in DH_{31}-incubated MTs, with high V_1 enrichment found in the membrane fractions,

confirming the crucial role of the VA in DH_{31}-stimulated secretion. Studies in *A. aegypti*

have demonstrated the involvement of PKA in the activation and assembly of the VA

upon natriuretic hormone (i.e. DH_{31}) stimulation and indicate the phosphorylation of the

VA subunits by PKA in the MTs (34). These studies indicate a regulatory role of PKA in

VA assembly and activation that may be independent or in addition to phosphorylation

(34).

Together, these results indicate that *Aedae*CAPA-1 binding to its cognate receptor

in principal cells of the MTs (6), targets the NOS/cGMP/PKG pathway (4, 6) to inhibit

DH_{31}-mediated elevation of cAMP (21, 74), which blocks VA activation and prevents

protons from being pumped across the apical membrane, creating a more alkaline lumen.

Our study provides novel evidence that the anti-diuretic activity of CAPA is mediated

through the dissociation of the VA holoenzyme involving the removal of the V_1 complex

from the apical membrane, hindering luminal flux of protons that in turn starves

cation/H^+ exchange, which ultimately reduces fluid secretion (Fig. 8). However, it has yet

to be determined how CAPA-activated PKG leads to the reversible dissociation of the

VA. In *R. prolixus* MTs, the physiological roles of cGMP and cAMP were examined (75)

suggesting cGMP inhibits fluid secretion by activating a phosphodiesterase (PDE) that

degrades cAMP that is elevated following 5HT and diuretic hormone (DH) stimulation of

MTs. Indeed, the addition of cAMP at high concentrations reversed the negative effects

of cGMP (75), supporting that these two cyclic nucleotides facilitate two opposing

physiological roles in the MTs. The data herein reveals cGMP levels increase in MTs

treated with CAPA alone or in combination with DH_{31} compared to tubules stimulated

with DH_{31} alone, which upholds the role of cGMP in the CAPA anti-diuretic signaling

pathway. Taken together, this data suggests that CAPA hormone activation of its receptor

in principal cells of the MTs elevates cGMP levels that activate a cAMP-selective PDE to

reduce levels of the diuretic hormone second messenger, cAMP. This in turn reduces the

activity of the PKA-activated VA, causing V_1V_o disassembly and ceasing transport of

protons into the tubule lumen.

In summary, our study highlights a novel target in the anti-diuretic signaling

pathway of adult female *A. aegypti* MTs, emphasizing the intricate and precise regulatory

mechanism of anti-diuresis. Although extensive studies have investigated the process of

hydromineral balance in terrestrial insects from a diuretic perspective (1, 10, 21, 76–78),

these current findings advance our understanding of anti-diuretic hormone control while

providing further evidence of a previously elusive endocrine regulatory mechanism of the

VA in mosquitoes (Fig. 8). Given that many terrestrial insects are recognized as

agricultural pests or disease vectors, further investigating their complex regulation of

ionic and osmotic balance may aid in lessening the burden of insects that are a hazard to

human health and prosperity through development of novel pest management strategies

that, at least in part, impede their neuroendocrine control of hydromineral homeostasis.

Materials and Methods

Animal rearing

Eggs of *Aedes aegypti* (Liverpool strain) were collected from an established

laboratory colony described previously (4, 79). All mosquitoes were raised under a 12:12

light:dark cycle. Non-blood fed female insects (three-six days post-eclosion) were used

for physiological assays, dissected under physiological saline (*Aedes* saline) adapted from

(60) that contained (in mmol^{-1}): 150 NaCl, 25 HEPES, 3.4 KCl, 7.5 NaOH, 1.8 NaHCO$_3$,

1 MgSO$_4$, 1.7 CaCl$_2$, and 5 glucose, and titrated to pH 7.1.

MT fluid secretion assay

In order to determine fluid secretion rates, modified Ramsay assays were

performed as described previously (4, 80). Female adults (3-6 day old) were dissected

under physiological *Aedes* saline prepared as previously described (60), and MTs were

removed and placed in a Sylgard-lined Petri dish containing 20 µL bathing droplets (1:1

mixture of Schneider's Insect Medium (Sigma-Aldrich): *Aedes* saline, immersed in

hydrated mineral oil to prevent evaporation. The proximal end of the MT was wrapped

around a Minutein pin to allow for fluid secretion measurements.

Time course inhibition of bafilomycin

Dosage of bafilomycin A$_1$ was based on a dose-response analysis of bafilomycin

against DH$_{31}$-stimulated tubules (Fig. S1). In the interest of determining whether

bafilomycin inhibits the effects of the diuretic factors, dosages of 25 nmol l^{-1} *Drome*DH$_{31}$

(4, 21), 100 nmol l^{-1} 5HT (4, 77, 81), and 10 nmol l^{-1} *Rhopr*DH (CRF-related diuretic

peptide, DH$_{44}$) (4, 54) were applied to the isolated MTs. Neurohormone receptors,

including those for 5HT, and the peptides DH$_{31}$, DH$_{44}$, and CAPA, are localized to the

basolateral membrane of principal cells (6, 61, 82), while the LK receptor is localized

exclusively to stellate cells (83). As a result, the effects of bafilomycin were tested on

diuretics known to act on the principal cells of the MTs. After incubating with the

individual diuretics for 30 min (using the modified Ramsay assay), diuretic peptide was

added alone (controls) or in combination with bafilomycin (final concentration 10^{-5} M).

The fluid secretion rate was recorded every 10 min for a total of 80 min. In order to

determine whether inhibition of the VA was involved in the anti-diuretic activity of

CAPA peptides on adult MTs, the effects of 1 fmol l^{-1} *Aedae*CAPA-1 (4, 6) were

investigated in combination with DH_{31} and bafilomycin.

Measurement of pH of secreted fluid

The pH of secreted fluids was measured by using ion-selective microelectrodes

(ISME) pulled from glass capillaries (TW-150-4, World Precision Instruments, Sarasota,

Fl, USA) using a Sutter P-97 Flaming Brown pipette puller (Sutter Instruments, San

Raffael, CA, USA). Microelectrodes were silanized with *N,N*-

dimethyltrimethylsilylamine (Fluka, Buchs, Switzerland) pipetted onto the interior of a

glass dish inverted over the group of microelectrodes. A 1:2 ratio of number of

microelectrodes to amount of silanization solution (in µl) was used. The microelectrodes

were left to silanize for 75 min at 350°C and left to cool before use. The microelectrodes

were back-filled with a solution containing 100 mmol l^{-1} NaCl and 100 mmol l^{-1} sodium

citrate that was titrated to pH 6.0 and front-filled using Hydrogen Ionophore I – cocktail

B (Fluka, Buchs, Switzerland). The electrode tips were then coated with ~3.5% (w/v)

polyvinyl chloride (PVC) dissolved in tetrahydrofuran, to avoid displacement of the

ionophore cocktail when submerged in the paraffin oil (84). The H^+-selective

microelectrodes were calibrated in *Aedes* saline titrated to either pH 7.0 or pH 8.0.

Reference electrodes were prepared from glass capillaries (1B100F-4, World Precision

Instruments) using a pipette puller described above and were backfilled with 500 mmol l^-

^{1} KCl. Secreted droplet pH measurements were done immediately after collection to

prevent alkalization of the droplet due to carbon dioxide diffusion into the paraffin oil.

Microelectrodes and reference electrodes were connected to an electrometer through

silver chloride wires where voltage signals were recorded through a data acquisition

system (Picolog for Windows, version 5.25.3). In order to measure pH of the secreted

fluid, tubules were set up using the Ramsay assay, and pH measurements were recorded

every 10 min for a total of 60 min.

NKA and VA activities

The Na^+/K^+-ATPase (NKA) and VA activity in the MTs was determined using a

modified 96-well microplate method (85, 86), which relies on the enzymatic coupling of

ouabain- or bafilomycin-sensitive hydrolysis of ATP to the oxidation of reduced

nicotinamide adenine dinucleotide (NADH). The microplate spectrophotometer is

therefore able to directly measure the disappearance of NADH. Adult female MTs (three

to six day old) were dissected and incubated for 30 min in *Aedes* saline, diuretic (DH_{31},

5HT, or DH_{44}) alone or combined with *Aedae*CAPA-1. Following 30-min incubation,

MTs were collected into 1.5 mL microcentrifuge tubes (40-50 sets of MTs per tube =

200-250 MTs per treatment), flash frozen in liquid nitrogen and stored at -80°C. Tissues

were thawed on ice and 150 μL of homogenizing buffer (four parts of SEI with

composition (in mmol l^{-1}): 150 sucrose, 10 EDTA, and 50 imidazole; pH 7.3 and one part

of SEID with composition: 0.5% of sodium deoxycholic acid in SEI) and MTs were then

sonicated on ice for 10 sec (two pulses of 5 sec) and subsequently centrifuged at 10,000 x

g for 10 min at 4°C. The supernatant was transferred into a fresh microcentrifuge tube

and stored on ice.

Prior to the assay, three solutions were prepared (Solution A, B, and C) and all

stored on ice. Solution A contained 4 units mL^{-1} lactate dehydrogenase (LDH), 5 units

mL^{-1} pyruvate kinase (PK), 50 mmol l^{-1} imidazole, 2.8 mmol l^{-1} phosphoenolpyruvate

(PEP), 0.22 mmol l^{-1} ATP, and 50 mmol l^{-1} NADH, pH 7.5. The solution was

subsequently mixed with a salt solution in a 4:1 ratio (salt solution composition (in mmol

l^{-1}): 189 NaCl, 10.5 MgCl$_2$, 42 KCl, and 50 imidazole, pH 7.5. Working solution B

consisted of solution A with 5 mmol l^{-1} ouabain and solution C consisted of solution A

with 10 μmol l^{-1} bafilomycin. The concentrations of ouabain and bafilomycin were based

on previous studies (85). To ensure the batch of assay mixture (Solution A) was effective,

an adenosine diphosphate (ADP) standard curve was run. ADP standards were prepared

as follows: 0 nmol 10μL^{-1} (200μL of 50mmol l^{-1} imidazole buffer (IB), pH 7.5); 5 nmol

10μL^{-1} (25μL of 4mmol l^{-1} ADP stock and 175μL of IB); 10 nmol 10uL^{-1} (50μL of

4mmol l^{-1} ADP stock/ 150 μL of IB); 20 nmol 10 μL^{-1} (100μL of 4mmol l^{-1} ADP

stock/100μL of IB); 40 nmol 10 μL^{-1} (40 mmol l^{-1} ADP stock). The standards were added

to a 96-well polystyrene microplate in duplicates of 10 μL per well, followed by the

addition of 200 μL of solution A. The plate was placed in a thermo Multiscan Spectrum

microplate spectrophotometer set at 25°C and a linear rate of NADH disappearance was

measured at 340 nm. The absorbance spectra were recorded and analyzed using the

Multiscan Spectrum data acquisition system with SkanIt version 2.2 software. The ADP

standards (0 to 40 nmoles well^{-1}) should yield an optical density (OD) between 0.9 and

0.2 OD units, while the slope of the curve should result in -0.012 to -0.014 OD nmol

ADP^{-1}. Homogenized MT samples were thawed and added to the microplate (kept on ice)

in six replicates of 10 µL per well. Next, two wells per sample were filled with 200 µL of

working solution A, two wells with 200 µL of working solution B and two wells with 200

µL of working solution C. The microplate was quickly placed in the microplate

spectrophotometer and the decrease in NADH absorbance was measured for 30 min at

340 nm. NKA and VA activity was calculated using the following equation:

550   NKA or VA activity = $(((\Delta\text{ATPase}/S)/[P]) \times 60$ (min),

where ΔATPase is the difference in ATP hydrolysis in the absence and presence of

ouabain or bafilomycin, S is the slope of the ADP standard curve, [P] is the protein

concentration of the sample. Protein was quantified using a Bradford assay (Sigma-

Aldrich Canada, Ltd.,) according to manufacturer's guidelines with bovine serum

albumin (Bioshop Canada Inc., Burlington, ON, Canada) as a standard. Final activity was

expressed as micromoles of ADP per milligram of protein per hour.

Protein processing and western blot analyses

MTs were isolated under physiological saline from 40-50 female *A. aegypti* for

each biological replicate (defined as n = 1) and incubated for 60 min in the following

treatments: *Aedes* saline, 25 nmol l^{-1} DH$_{31}$, or 25 nmol l^{-1} DH$_{31}$ and 1 fmol l^{-1}

*Aedae*CAPA-1. Following the incubation, tissues were stored at -80ºC until processing.

To separate the membrane and cytosolic proteins, a membrane protein extraction kit was

used (ThermoFisher Scientific) following recommended guidelines for soft tissue with

minor modifications including 200 μL of permeabilization and solubilization buffer and a

1:200 protease inhibitor cocktail (Sigma Aldrich) in both buffers. Final protein

concentrations were calculated using a Bradford assay (Sigma-Aldrich Canada, Ltd.,)

according to manufacturer's guidelines with bovine serum albumin (BioRad

Laboratories) as a standard and quantified using an A_o Absorbance Microplate Reader

(Azure Biosystems) at 595 nm.

Protein samples were denatured by heating for 5 min at 100ºC with 6X loading

buffer (225 mmol l^{-1} Tris-HCl pH 6.8, 3.5% (w/v) SDS, 35% glycerol, 12.5% (v/v) ß-

mercaptoethanol and 0.01% (w/v) Bromophenol blue). Into each lane, 5 μg of protein

was loaded onto a 4% stacking and 12% resolving sodium dodecyl sulphate

polyacrylamide gel electrophoresis (SDS-PAGE) gel. Protein samples were migrated

initially at 80 V for 30 min and subsequently at 110 V for 90 min before being transferred

onto a polyvinylidene difluoride (PVDF) membrane using a wet transfer method at 100 V

for 60 min in a cold transfer buffer (0.225 g Tris. 1.05 g glycine in 20% methanol).

Following transfer, PVDF membranes were blocked with 5% skim milk powder in Tris-

buffered saline (TBS-T; 9.9 mmol l^{-1} Tris, 0.15 mmol l^{-1} NaCl, 0.1 mmol l^{-1} Tween-20,

0.1 mmol l^{-1} NP-40 pH 7.4) for 60 min at RT, and incubated on a rocking platform

overnight at 4ºC with a guinea pig polyclonal anti-VA (Ab 353-2 against the V_1 complex

of the VA (11), a kind gift from Prof. Wieczorek, University of Osnabruck, Germany,

used at a 1:2000 dilution in 5% skim milk in TBS-T. The next day, PVDF membranes

were washed for 60 min in TBST-T, changing the wash buffer every 15 min. Blots were

then incubated with a goat anti-guinea pig HRP conjugated secondary antibody (1:2500

in 5% skim milk in TBS-T) (Life Technologies, Burlington, ON) for 60 min at RT and

subsequently washed three times for 15 min with TBS-T. Lastly, blots were incubated

with the Clarity Western ECL substrate and images were acquired using a ChemiDoc MP

Imaging System (Bio-Rad). Molecular weight measurements were performed using

Image Lab 5.0 software (Bio-Rad). PVDF membranes were then probed with Coomassie

brilliant blue, since total protein normalization is now considered the benchmark method

for quantitative analysis of western blot data (87, 88) and has been used in studies

involving *A. aegypti* protein normalization (89). ImageJ software (NIH, USA) was used

to quantify protein abundance.

Immunolocalization of VA complexes in MTs

Immunohistochemistry of the MTs localizing the VA complexes was conducted

following a previously published protocol (90). Adult female MTs (three to six day old)

were dissected out in *Aedes* saline and incubated in similar conditions described in the

western blot section above. Following the incubation, the MTs were immersed in Bouin's

solution and fixed for two hours in small glass vials. Tissues were then rinsed three times

and stored in 70% ethanol at 4ºC until further processing. Fixed tissues were dehydrated

through a series of ethanol washes: 70% ethanol for 30 min, 95% ethanol for 30 min, and

100% ethanol three times for 30 min. The tissues were cleared with xylene

(ethanol:xylene for 30 min then 100% xylene three times for 30 min), and infiltrated in

Paraplast Plus Tissue Embedding Medium (Oxford Worldwide, LLC, Memphis, USA) at

xylene:paraffin wax for 60 min at 60ºC, then rinsed in pure paraffin wax twice for 1 hour

for 60ºC. Following the last rinse, the tissues were embedded in the paraffin wax and left

to solidify at 4ºC until further processing. Sections (5 μm) were cut using a Leica RM

2125RT manual rotary microtome (Leica Microsystems Inc., Richmond Hill, Canada)

and slides were incubated overnight on a slide warmer at 45ºC.

The following day, sections were deparaffinized with xylene (two rinses for 5 min

each), and rehydrated via a descending series of ethanol washes (100% ethanol twice for

2 min, 95% ethanol for 2 min, 70% ethanol for 2 min, 50% ethanol for 2 min) and finally

in distilled water for 20 min. Next, sections were subjected to a heat-induced epitope

retrieval (HIER) by immersing slides in a sodium citrate buffer (10 nmol l^{-1}, pH 6.0) and

heating both slides and solution in a microwave oven for 4 min. The solution and slides

were allowed to cool for 20 min, reheated for 2 min, and left to stand at room temperature

(RT) for 15 min. Slides were then washed three times in phosphate-buffered saline (PBS)

pH 7.4, 0.4% Kodak Photo-Flo 200 in PBS (PBS/PF, 10 min), 0.05% Triton X-100 in

PBS (PBS/TX, 10 min), and 10% antibody dilution buffer (ADB; 10% goat serum, 3%

BSA and 0. 05% Triton X-100 in PBS) in PBS (PBS/ADB, 10 min). Slides were

incubated overnight at RT with a guinea pig polyclonal anti-V_1 (Ab 353-2 against the V_1

complex, identical antibody used in western blot analyses, at 1:5000 dilution in ADB) in

combination with a 1:100 mouse polyclonal anti-ATP6V0A1 antibody for V_o (Abnova,

Taipei, Taiwan).

Following overnight primary antibody incubation, slides were washed briefly in

distilled water, with sequential washes with PBS/PF, PBS/TX, and PBS/ADB for 10 min

each as described above. A goat anti-guinea pig antibody conjugated to AlexaFluor 488

(1:500 in ADB, Jackson Immunoresearch) and sheep anti-mouse antibody conjugated to

AlexaFluor 594 or (1:500 in ADB, Jackson Immunoresearch) were applied to visualize

the VA complexes. The slides were left to incubate in the secondary antibody for 60 min

at RT. For negative controls, slides were processed as described above with primary

antibodies omitted. Slides were then washed in distilled water/0.4% PF three times for 1

min each and finally in distilled water for 1 min. Slides were air dried for 30 min and

mounted using ProLong^{TM} Gold antifade reagent with DAPI (Life Technologies,

Burlington, ON, Canada). Fluorescence images were captured using an Olympus IX81

inverted fluorescent microscope (Olympus Canada, Richmond Hill, ON, Canada). Using

ImageJ, V_1 cell fluorescence was measured as described previously (91, 92) where

sections of interest were selected and their area, integrated density, and mean grey values

measured. A total of 14 – 16 preps were selected, consisting of both transverse and cross-

sections, and three sections of interest were selected from both membrane and cytosolic

staining (the size of the selected area was kept consistent to reduce bias). Background

values were also recorded by selecting regions of the preps with no fluorescence near the

sections of interest. The corrected cell fluorescence (CCF) was calculated using the

equation: CCF = integrated density – (area of selected cell x mean fluorescence of

background readings (modified from original equation since only portions of the cell

were selected).

cGMP Measurements

A competitive cGMP ELISA kit (Cell Signaling Technology, #4360) was used to

measure the effect of DH$_{31}$ and *Aedae*CAPA-1 on cGMP levels in the MTs. Adult MTs

were isolated under physiological saline from 50 female *A. aegypti* for each biological

replicate (defined as n = 1). To prevent cGMP degradation, tubules were incubated first

with a phosphodiesterase inhibitor, 10^{-4} mol^{-1} zaprinast, for 10 min (93) before any of the

experimental treatments, specifically *Aedes* saline, 25 $nmol^{-1}$ DH_{31}, 1 $fmol^{-1}$

*Aedae*CAPA-1, or 25 $nmol^{-1}$ DH_{31} + 1 $fmol^{-1}$ *Aedae*CAPA-1 for a further 20 min. After

incubation was complete, tissues were stored at -80ºC until processing. To measure

cGMP concentrations, frozen tubule samples were thawed on ice and 125 µL of 1X cell

lysis buffer (CLB, #9803) was added to each tube (1 $mmol^{-1}$ phenylmethylsulfonyl

fluoride (PMSF) was added to 1X CLB fresh each time). Tissue samples were kept on ice

for 10 min, sonicated for 10 s (similar conditions as described for NKA/VA activity

assay), centrifuged for 3 min at 10,000 x rpm, and the supernatant was isolated and kept

on ice. Using a commercial 96-well microtitre plate precoated with cGMP rabbit

monoclonal antibody, the cGMP-HRP conjugate was added in triplicate wells with 50 µL

per well. This was followed by the addition of 50 µL of either tubule samples or cGMP

standards ranging from 100 $nmol^{-1}$ to 0.25 $nmol^{-1}$. The plate was covered and incubated

at RT for 3 h on a horizontal orbital plate shaker. Following incubation, plate contents

were discarded, and wells were washed four times with 200 µL/well of 1X wash buffer.

Next, 100 µL of 3,3',5,5'-tetramethylbenzidine (TMB) substrate was added to each well,

and the plate was covered and kept for 10 min at RT. The enzymatic reaction was

quenched by adding 100 µL of 2 $mmol^{-1}$ HCl and absorbance read at 450 nm using a

Synergy 2 Microplate Reader (Biotek).

Statistical analyses

Data was compiled using Microsoft Excel and transferred to Graphpad Prism

software v.7 to create figures and conduct all statistical analyses. Data was analyzed

accordingly using a one-way or two-way ANOVA and a Bonferroni post-test, or

Student's T-test, with differences between treatments considered significant if $p<0.05$.

Acknowledgments

The authors sincerely thank Prof. Helmut Wieczorek and Prof. Felix Tiburcy (University

of Osnabrück, Germany) for providing the V_1 antibody used in this study. The authors are

also grateful to Professors Ian Orchard (University of Toronto Mississauga, Canada) and

Michael J. O'Donnell (McMaster University, Canada) for providing aliquots of *Rhopr*DH

and *Drome*DH$_{31}$ used in this study.

Funding

This research was funded by: Natural Sciences and Engineering Research Council of

Canada (NSERC) Discovery Grant (JPP)**,** Ontario Ministry of Research Innovation Early

Researcher Award (JPP), and NSERC CGS-D (FS) and the Carswell Scholarships in the

Faculty of Science (FS).

Data and materials availability

All data are available in the main text or the supplementary materials. Additional data

related to this paper may be requested from the authors.

Figures and Tables

Fig. 1. Effect of 10^{-5} M bafilomycin on fluid secretion rates by *A. aegypti* MTs stimulated

with different diuretic hormones. MTs were treated with either (A) 25 nmol l^{-1} DH_{31} (B)

100 nmol l^{-1} 5HT or (C) 10 nmol l^{-1} DH_{44}, and secreted droplets were measured at 10-min

intervals for 30 min. Immediately following measurement of the 30-min point fluid

droplet (solid arrow), MTs were treated with (A) DH_{31} (B) 5HT (C) or DH_{44} alone (blue

circles) or in combination with 10^{-5} M bafilomycin (yellow squares). Droplets were

measured at similar intervals for an additional 50 min. Significant differences between

bafilomycin-treated and the corresponding time point controls are denoted by an asterisk,

as determined by a two-way ANOVA and Bonferroni multiple comparison post-hoc test

($p<0.05$). Data represent the mean ± standard error (n=12-23). (D) MTs were treated with

25 nmol l^{-1} DH_{31} alone or in combination with 1 fmol l^{-1} *Aedae*CAPA-1, bafilomycin, or

both *Aedae*CAPA-1 and bafilomycin for 60 min. Data labelled with different letters are

significantly different from each other (mean± SEM; one-way ANOVA with Bonferroni

multiple comparison, $p<0.05$, n=7-17).

Fig. 2. Effect of bafilomycin and *Aedae*CAPA-1 on pH of the secreted fluid by *A.*

aegypti MTs stimulated with distinct diuretic hormones. MTs were treated with either (A)

DH_{31} (B) 5HT (C) or DH_{44}, and secreted droplets were measured at 10-min intervals for

30 min using an ion-selective microelectrode. Immediately following measurement of the

30-min point fluid droplet (solid arrow), MTs were treated with (A) DH_{31} (B) 5HT (C) or

DH_{44} alone (blue circles) or in combination with *Aedae*CAPA-1 (purple triangles) or

bafilomycin (yellow squares). Droplets were measured at similar intervals for an

additional 30 min. Significant differences between *Aedae*CAPA-1- and bafilomycin-

treated and the corresponding time point controls are denoted by a double asterisk,

(mean± SEM; two-way ANOVA with Bonferroni multiple comparison, $p<0.05$, n=14-

34).

Fig. 3. Effect of *Aedae*CAPA-1 on VA and NKA activity and cGMP levels in stimulated

A. aegypti MTs. MTs were incubated in *Aedes* saline, diuretics (DH$_{31}$, 5HT, and DH$_{44}$)

alone or in combination with *Aedae*CAPA-1 for 30 min before collection to measure (A)

VA and (B) NKA activity. Data labelled with different letters are significantly different

from each other (mean± SEM; one-way ANOVA with Bonferroni multiple comparison,

$p < 0.05$). N.S. denotes no statistical significance. (n=50 sets of MTs for all treatments

(n=3 per treatment)). To monitor activity of *Aedae*CAPA-1 on cGMP production in

stimulated *A. aegypti* MTs (C), tubuules were incubated in zaprinast for 10 min before

incubation with following treatments; *Aedes* saline, DH$_{31}$, *Aedae*CAPA-1 alone, or in

combination with DH$_{31}$ for 20 min before collection. An asterisk is denoted for

significance within treatments (mean± SEM; one-way ANOVA with Bonferroni multiple

comparison, $p < 0.05$, n=50 sets of MTs for all treatments (n=3 per treatment)).

Fig. 4. Membrane and cytosolic protein abundance of the V_1 complex in *A. aegypti*

MTs. The MTs (n=40-50) were incubated in *Aedes* saline, DH_{31}, or DH_{31} and

*Aedae*CAPA-1 for one hour before collection. Protein abundance was measured in the

(A-C) 74 kDa (D-F) 56 kDa and (G-I) 32 kDa bands of the V_1 complex. Individual band

densities were normalized to total protein using Coomassie staining, and graphed relative

to saline-treated controls. An asterisk denotes significance within cytosolic and

membrane fractions (mean± SEM; Student's T-test, $p < 0.05$, n=3-4 replicates) whereas ns

denotes no significance.

Fig. 5. Immunolocalization of the V_o and V_1 complexes in transverse sections of

stimulated *A. aegypti* MTs. Representative paraffin-embedded sections of *A. aegypti* MTs

incubated in either (A-C) *Aedes* saline, (D-F) DH_{31} and (G-I) DH_{31} and *Aedae*CAPA-1.

Panels (A,D,G) show V_o staining (red), (B,E,H) show V_1 staining (green), and panels

(C,F,I) show merged antibody staining highly prominent in DH_{31} treatment but less

evident in saline and *Aedae*CAPA-1 treatments. Solid white arrows denote apical VA

staining, and empty arrows indicate cytosolic VA staining. DAPI nuclear staining is

shown in blue. Scale bar 100 μm (PC = principal cell, SC = stellate cell).

Fig. 6. Immuno co-localization of the V_o and V_1 complexes in cross sections of

stimulated *A. aegypti* MTs. Representative paraffin-embedded sections of *A. aegypti* MTs

incubated in either (A, B) *Aedes* saline, (C, D) DH_{31} and (E-J) DH_{31} and *Aedae*CAPA-1.

Images A, C and E-G show merged V_o and V_1 antibody staining while B, D and H-J

show brightfield images. Solid white arrows denote apical VA staining, and empty

arrows indicate cytosolic VA staining. DAPI nuclear staining is shown in blue. Scale bar

100 μm. (PC = principal cell).

Fig. 7. Corrected cell fluorescence (CTCF) of V_1 staining in stimulated *A.*

aegypti MTs. (A) Membrane and (B) cytosolic V_1 staining was measured using

immunohistochemistry in MTs incubated in either *Aedes* saline, DH_{31}, or DH_{31} and

*Aedae*CAPA-1. Data labelled with different letters are significantly different from each

other (mean± SEM; one-way ANOVA with Bonferroni multiple comparison, $p<0.05$,

n=14-16).

Fig. 8. Schematic diagram summarizing the signaling pathway of diuretic and anti-

diuretic control of adult *A. aegypti* MTs. The principal cells in *A. aegypti* MTs are

responsible for the transport of Na^+ and K^+ cations via secondary active transport,

energized by the V-type H^+-ATPase (VA), localized in the brush border of the apical

membrane. The movement of protons creates a gradient, driving the exchange of Na^+ and

K^+ through cation/H^+ antiporters (NHA). Neurohormone receptors for DH_{31}, 5HT, DH_{44},

and CAPA are localized to the basolateral membrane of the principal cells, while the

kinin receptor is localized exclusively in the stellate cells. Our data together with

previous work indicates that DH_{31} stimulates diuresis through activation and assembly of

the VA in the apical membrane, with no effect on the Na^+/K^+-ATPase (NKA). The anti-

diuretic effect of *Aedae*CAPA-1, facilitated by the NOS/cGMP/PKG pathway, causes V_1

dissociation from the membrane, hindering activity, and thus reducing fluid secretion.

The biogenic amine, 5HT, has also been shown to stimulate activation of the VA,

however, to a lesser extent. DH_{44}-related peptide receptor activation increases Ca^{2+} (as

well as cAMP at higher doses) but it's action was found to be independent of the VA.

Lastly, it was shown earlier that cGMP inhibits kinin-stimulated diuresis, suggesting an

additional anti-diuretic factor may exist that acts specifically on stellate cells.

Fig. S1. Dose response of bafilomycin on *in vitro* fluid secretion rates of DH_{31}-

stimulated MTs of adult female *A. aegypti*. Doses of 10^{-4} M to 10^{-10} M bafilomycin were

applied to MTs together with DH_{31} for 60 min. Data labelled with different letters are

significantly different from each other (mean± SEM; one-way ANOVA with Bonferroni

multiple comparison, p<0.05, n=5-13).

Fig. S2. Measurement of pH in fluid secreted by unstimulated *A. aegypti* adult female

MTs. Secreted droplets were isolated and measured at 10-min intervals for 60 min using

an ion-selective microelectrode. No significant differences were observed in the

measurements, (mean± SEM; two-way ANOVA with Bonferroni multiple comparison,

p<0.05, n=10); ns denotes no statistical significance.

Fig. S3. Immunolocalization of the V_o and V_1 complexes in stimulated *A. aegypti* MTs

(cross sections). Representative paraffin-embedded sections of *A. aegypti* MTs

incubated in either (A, B) *Aedes* saline, (C, D) DH_{31} and (E-J) DH_{31} and *Aedae*CAPA-1.

Panels (A,C, E-G) show membrane-integrated V_o staining (red) and (B,D,H-J)

demonstrate V_1 staining (green). Solid white arrows denote apical VA staining, and

empty arrows indicate cytosolic VA staining. Scale bar 100 μm (PC = principal cell).

References

1. K. W. Beyenbach, Transport mechanisms of diuresis in Malpighian tubules of insects. *J. Exp. Biol.* **206**, 3845–3856 (2003).

2. T. M. Clark, T. J. Bradley, Malpighian tubules of larval *Aedes aegypti* are hormonally stimulated by 5-hydroxytryptamine in response to increased salinity. *Arch. Insect Biochem. Physiol.* **34**, 123–141 (1997).

3. G. M. Coast, C. S. Garside, Neuropeptide control of fluid balance in insects, *Ann. N.Y. Acad. Sci.* **1040**: 1–8 (2005).

4. F. Sajadi, C. Curcuruto, A. Al Dhaheri, J.-P. V Paluzzi, Anti-diuretic action of a CAPA neuropeptide against a subset of diuretic hormones in the disease vector, *Aedes aegypti*. *J. Exp. Biol.* (2018).

5. H. L. Lu, C. Kersch, P. V. Pietrantonio, The kinin receptor is expressed in the Malpighian tubule stellate cells in the mosquito *Aedes aegypti* (L.): A new model needed to explain ion transport? *Insect Biochem. Mol. Biol.* **41**, 135–140 (2011).

6. F. Sajadi, *et al.*, CAPA neuropeptides and their receptor form an anti-diuretic hormone signaling system in the human disease vector, *Aedes aegypti*. *Sci. Rep.* **10** (2020).

7. A. Ionescu, A. Donini, *Aedes*CAPA-PVK-1 displays diuretic and dose dependent antidiuretic potential in the larval mosquito *Aedes aegypti* (Liverpool). *J. Insect Physiol.* **58**, 1299–1306 (2012).

8.   R. Jurenka, The PRXamide Neuropeptide Signalling System. Conserved in

Animals, *Adv. Insect Phys.* (Academic Press, 2015), pp. 123–170.

9.   S. A. Davies, *et al.*, Signaling by *Drosophila* capa neuropeptides. *Gen. Comp.*

Endocrinol. **188**, 60–66 (2012).

10.   S. Terhzaz, *et al.*, Mechanism and function of *drosophila* capa GPCR: A

desiccation stress-responsive receptor with functional homology to human

neuromedinu receptor. *PLoS One* **7** (2012).

11.   X. H. Weng, M. Huss, H. Wieczorek, K. W. Beyenbach, The V-type H^+-ATPase

in Malpighian tubules of *Aedes aegypti*: Localization and activity. *J. Exp. Biol.*

206, 2211–2219 (2003).

12.   H. Wieczorek, K. W. Beyenbach, M. Huss, O. Vitavska, Vacuolar-type proton

pumps in insect epithelia. *J. Exp. Biol.* **212**, 1611–1619 (2009).

13.   H. Wieczorek, The insect V-ATPase, a plasma membrane proton pump energizing

secondary active transport: molecular analysis of electrogenic potassium transport

in the tobacco hornworm midgut. *J. Exp. Biol.* **172**, 335–343 (1992).

14.   H. Wieczorek, M. Putzenlechnere, U. Klein, A Vacuolar-type proton pump

energizes K^+/H^+ antiport in an animal plasma membrane. *J. Biol. Chem.* **266**,

15340–15347 (1991).

15.   W. R. Harvey, S. H. P. Maddrell, W. H. Telfer, H. Wieczorek, H^+ V-ATPases

energize animal plasma membranes for secretion and absorption of ions and fluids.

Am. Zool. **38**, 426–441 (1998).

16.   Y. Li, *et al.*, RNA-Seq comparison of larval and adult malpighian tubules of the

yellow fever mosquito *Aedes aegypti* reveals life stage-specific changes in renal

function. *Front. Physiol.* **8**, 1–13 (2017).

17.   M. L. Patrick, K. Aimanova, H. R. Sanders, S. S. Gill, P-type Na^+/K^+-ATPase and

V-type H^+-ATPase expression patterns in the osmoregulatory organs of larval and

adult mosquito *Aedes aegypti*. *J. Exp. Biol.* **209**, 4638–4651 (2006).

18.   H. Wieczorek, K. W. Beyenbach, M. Huss, O. Vitavska, Vacuolar-type proton

pumps in insect epithelia. *J. Exp. Biol.* **212**, 1611–1619 (2009).

19.   R. M. Hine, *et al.*, The excretion of NaCl and KCl loads in mosquitoes. *Am. J.*

Physiol. - Regul. Integr. Comp. Physiol. **307**, R837–R849 (2014).

20.   P. Dames, *et al.*, cAMP regulates plasma membrane vacuolar-type H^+-ATPase

assembly and activity in blowfly salivary glands. *Proc. Natl. Acad. Sci. U. S. A.*

103, 3926–3931 (2006).

21.   G. M. Coast, C. S. Garside, S. G. Webster, K. M. Schegg, D. A. Schooley,

Mosquito natriuretic peptide identified as a calcitonin-like diuretic hormone in

Anopheles gambiae (Giles). *J. Exp. Biol.* **208**, 3281–3291 (2005).

22.   K. Karas, P. Brauer, D. Petzel, Actin redistribution in mosquito Malpighian

tubules after a blood meal and cyclic AMP stimulation. *J. Insect Physiol.* **51**,

1041–1054 (2005).

23. C. Cady, H. H. Hagedorn, Effects of putative diuretic factors on intracellular

second messenger levels in the Malpighian tubules of *Aedes aegypti*. *J. Insect*

Physiol. **45**, 327–337 (1999).

24. D. H. Petzel, P. T. Pirotte, E. Van Kerkhove, Intracellular and luminal pH

measurements of malpighian tubules of the mosquito *Aedes aegypti*: The effects of

cAMP. *J. Insect Physiol.* **45**, 973–982 (1999).

25. V. Müller, G. Grüber, ATP synthases: Structure, function and evolution of unique

energy converters. *Cell. Mol. Life Sci.* **60**, 474–494 (2003).

26. J. Zhang, Y. Feng, M. Forgac, Proton conduction and bafilomycin binding by the

V0 domain of the coated vesicle V-ATPase. *J. Biol. Chem.* **269**, 23518–23523

(1994).

27. W. J. A. Dschida, B. J. Bowman, The vacuolar ATPase: Sulfite stabilization and

the mechanism of nitrate inactivation. *J. Biol. Chem.* **270**, 1557–1563 (1995).

28. J. Rein, M. Voss, W. Blenau, B. Walz, O. Baumann, Hormone-induced assembly

and activation of V-ATPase in blowfly salivary glands is mediated by protein

kinase A. *Am. J. Physiol. - Cell Physiol.* **294**, C56–C65 (2008).

29. B. Zimmermann, P. Dames, B. Walz, O. Baumann, Distribution and serotonin-

induced activation of vacuolar-type H+-ATPase in the salivary glands of the

blowfly *Calliphora vicina. J. Exp. Biol.* **206**, 1867–1876 (2003).

30. H. Wieczorek, G. Grüber, W. R. Harvey, M. Huss, H. Merzendorfer, The plasma

membrane H^+-V-ATPase from tobacco hornworm midgut. *J. Bioenerg. Biomembr.*

31, 67–74 (1999).

31.   K. W. Beyenbach, T. L. Pannabecker, W. Nagel, Central role of the apical

membrane H^+-ATPase in electrogenesis and epithelial transport in Malpighian

tubules. *J. Exp. Biol.* **203**,1459–1468 (2000).

32.   H. Wieczorek, D. Brown, S. Grinstein, J. Ehrenfeld, W. R. Harvey, Animal plasma

membrane energization by proton-motive V-ATPases. *BioEssays* **21**, 637–648

(1999).

33.   O. Baumann, B. Walz, The blowfly salivary gland – A model system for analyzing

the regulation of plasma membrane V-ATPase. *J. Insect Physiol.* **58**, 450–458

(2012).

34.   F. Tiburcy, K. W. Beyenbach, H. Wieczorek, Protein kinase A-dependent and -

independent activation of the V-ATPase in Malpighian tubules of *Aedes aegypti. J.*

Exp. Biol. **216**, 881–891 (2013).

35.   W. R. Harvey, S. H. P. Maddrell, W. H. Telfer, H. Wieczorek, H^+ V-ATPases

energize animal plasma membranes for secretion and absorption of ions and fluids.

Am. Zool. **38**, 426–441 (1998).

36.   K. W. Beyenbach, H. Wieczorek, The V-type H^+ ATPase: Molecular structure and

function, physiological roles and regulation. *J. Exp. Biol.* **209**, 577–589 (2006).

37.   F. J. S. Novak, *et al.*, Primary structure of V-ATPase subunit B from *Manduca*

994   *sexta* midgut. *BBA - Gene Struct. Expr.* **1132**, 67–71 (1992).

995 38. Y. X. Pan, H. H. Gu, J. Xu, G. E. Dean, *Saccharomyces cerevisiae* expression of

996   exogenous vacuolar ATPase subunits B. *BBA - Biomembr.* **1151**, 175–185 (1993).

997 39. K. W. Beyenbach, P. M. Piermarini, Transcellular and paracellular pathways of

998   transepithelial fluid secretion in Malpighian (renal) tubules of the yellow fever

999   mosquito *Aedes aegypti*. *Acta Physiol. (Oxf)*. **202**, 387–407 (2011).

1000 40. K. W. Beyenbach, Energizing epithelial transport with the vacuolar H^+-ATPase.

1001   *News Physiol. Sci.* **16**, 145–151 (2001).

1002 41. T. Nishi, M. Forgac, The vacuolar (H^+)-ATPases - Nature's most versatile proton

1003   pumps. *Nat. Rev. Mol. Cell Biol.* **3**, 94–103 (2002).

1004 42. N. Nelson, W. R. Harvey, Vacuolar and plasma membrane proton-

1005   adenosinetriphosphatases. *Physiol. Rev.* **79**, 361–385 (1999).

1006 43. K. W. Beyenbach, H. Skaer, J. A. T. Dow, The developmental, molecular, and

1007   transport biology of Malpighian tubules. *Annu. Rev. Entomol.* **55**, 351–374 (2010).

1008 44. K. W. Beyenbach, H. Skaer, J. A. T. Dow, The developmental, molecular, and

1009   transport biology of Malpighian tubules. *Annu. Rev. Entomol.* **55**, 351–374 (2010).

1010 45. P. M. Kane, Disassembly and reassembly of the yeast vacuolar H^+-ATPase in vivo.

1011   *J. Biol. Chem.* **270**, 17025–17032 (1995).

1012 46. M. Voss, O. Vitavska, B. Walz, H. Wieczorek, O. Baumann, Stimulus-induced

phosphorylation of vacuolar H$^+$-ATPase by protein kinase A. *J. Biol. Chem.* **282**, 33735–33742 (2007).

47. J. P. Sumner, *et al.*, Regulation of plasma membrane V-ATPase activity by dissociation of peripheral subunits. *J. Biol. Chem.* **270**, 5649–5653 (1995).

48. G. Coast, The endocrine control of salt balance in insects. *Gen. Comp. Endocrinol.* **152**, 332–338 (2007).

49. M. J. O'Donnell, J. H. Spring, Modes of control of insect Malpighian tubules: Synergism, antagonism, cooperation and autonomous regulation. *J. Insect Physiol.* **46**, 107–117 (2000).

50. J. C. Williams, H. H. Hagedorn, K. W. Beyenbach, Dynamic changes in flow rate and composition of urine during the post-bloodmeal diuresis in *Aedes aegypti* (L.). *J. Comp. Physiol.* **153**, 257–265 (1983).

51. C. L. Jagge, P. V. Pietrantonio, Diuretic hormone 44 receptor in Malpighian tubules of the mosquito *Aedes aegypti*: evidence for transcriptional regulation paralleling urination. *Insect Mol. Biol.* **17**, 413–426 (2008).

52. P. Gioino, B. G. Murray, J. P. Ianowski, Serotonin triggers cAMP and PKA-mediated intracellular calcium waves in Malpighian tubules of *Rhodnius prolixus*. *Am. J. Physiol. - Regul. Integr. Comp. Physiol.* **307**, R828–R836 (2014).

53. T. M. Clark, T. K. Hayes, K. W. Beyenbach, Dose-dependent effects of CRF-like diuretic peptide on transcellular and paracellular transport pathways. *Am. J.*

Physiol. - Ren. Physiol. **274** (1998).

54. K. . Clark, T M, Hayes, T K., Holman, G.M., Beyenbach, The concentration-

dependence of CRF-like diuretic peptide: mechanisms of action. *J. Exp. Biol.* **201**,

1753–1762 (1998).

55. S. J. Cole, J. D. Gillett, The influence of the brain hormone on retention of blood

in the mid-gut of the mosquito *Aedes aegypti* (L.). III. The involvement of the

ovaries and ecdysone. *Proc. R. Soc. London - Biol. Sci.* **205**, 411–422 (1979).

56. E. Van Handel, M. J. Klowden, Defecation by the mosquito, *Aedes aegypti*, is

controlled by the terminal abdominal ganglion. *J. Insect Physiol.* **42**, 139–142

(1996).

57. D. H. Petzel, H. H. Hagedorn, K. W. Beyenbach, Preliminary isolation of

mosquito natriuretic factor. *Am. J. Physiol. - Regul. Integr. Comp. Physiol.* **18**

(1985).

58. J. A. Dow, *et al.*, The malpighian tubules of *Drosophila melanogaster*: a novel

phenotype for studies of fluid secretion and its control. *J. Exp. Biol.* **197**, 421–428

(1994).

59. M. A. B. Grieco, A. G. Lopes, 5-Hydroxytryptamine regulates the

(Na^+/K^+)ATPase activity in Malpighian tubules of *Rhodnius prolixus*: evidence for

involvement of G-Protein and cAMP-dependent protein kinase. *Arch. Insect*

Biochem. Physiol. **36**, 203–214 (1997).

60.   D. H. Petzel, M. M. Berg, K. W. Beyenbach, Hormone-controlled cAMP-mediated

fluid secretion in yellow-fever mosquito. *Am. J. Physiol. - Regul. Integr. Comp.*

Physiol. **253** (1987).

61.   H. Kwon, H. L. Lu, M. T. Longnecker, P. V. Pietrantonio, Role in diuresis of a

calcitonin receptor (GPRCAL1) expressed in a distal-proximal gradient in renal

organs of the mosquito *Aedes aegypti* (L.). *PLoS One* **7**, e50374 (2012).

62.   W. Kang'ethe, K. G. Aimanova, A. K. Pullikuth, S. S. Gill, NHE8 mediates

amiloride-sensitive Na^+/H^+ exchange across mosquito Malpighian tubules and

catalyzes Na^+ and K^+ transport in reconstituted proteoliposomes. *Am. J. Physiol. -*

Ren. Physiol. **292**, 1501–1512 (2007).

63.   A. K. Pullikuth, K. Aimanova, W. Kang'ethe, H. R. Sanders, S. S. Gill, Molecular

characterization of sodium/proton exchanger 3 (NHE3) from the yellow fever

vector, *Aedes aegypti*. *J. Exp. Biol.* **209**, 3529–3544 (2006).

64.   P. M. Piermarini, D. Weihrauch, H. Meyer, M. Huss, K. W. Beyenbach, NHE8 is

an intracellular cation/H^+ exchanger in renal tubules of the yellow fever mosquito

Aedes aegypti. *Am. J. Physiol. - Ren. Physiol.* **296**, 730–750 (2009).

65.   J. C. Williams, K. W. Beyenbach, Differential effects of secretagogues on the

electrophysiology of the Malpighian tubules of the yellow fever mosquito. *J.*

Comp. Physiol. B **154**, 301–309 (1984).

66.   J. L. Hegarty, *et al.*, Dibutyryl cAMP activates bumetanide-sensitive electrolyte

transport in Malpighian tubules. *Am. J. Physiol. - Cell Physiol.* **261** (1991).

67. H. Merzendorfer, M. Huss, R. Schmid, W. R. Harvey, H. Wieczorek, A novel

insect V-ATPase subunit M9.7 is glycosylated extensively. *J. Biol. Chem.* **274**,

17372–17378 (1999).

68. Q. Song, B. Meng, H. Xu, Z. Mao, The emerging roles of vacuolar-type ATPase-

dependent Lysosomal acidification in neurodegenerative diseases. *Transl.*

Neurodegener. **9**, 1–14 (2020).

69. R. A. Oot, S. Couoh-Cardel, S. Sharma, N. J. Stam, S. Wilkens, Breaking up and

making up: The secret life of the vacuolar H$^+$-ATPase. *Protein Sci.* **26**, 896–909

(2017).

70. R. Gräf, W. R. Harvey, H. Wieczorek, Purification and properties of a cytosolic

V1-ATPase. *J. Biol. Chem.* **271**, 20908–20913 (1996).

71. O. Vitavska, H. Wieczorek, H. Merzendorfer, A novel role for subunit C in

mediating binding of the H$^+$-V-ATPase to the actin cytoskeleton. *J. Biol. Chem.*

278, 18499–18505 (2003).

72. T. Inoue, M. Forgac, Cysteine-mediated cross-linking indicates that subunit C of

the V-ATPase is in close proximity to subunits E and G of the V1 domain and

subunit a of the V0 domain. *J. Biol. Chem.* **280**, 27896–27903 (2005).

73. M. J. Berridge, B. D. Lindley, W. T. Prince, Studies on the mechanism of fluid

secretion by isolated salivary glands of *Calliphora*. *J. Exp. Biol.* **64**, 311–322

(1976).

74. D. H. Petzel, M. M. Berg, K. W. Beyenbach, Hormone-controlled cAMP-mediated

fluid secretion in yellow-fever mosquito. *Am. J. Physiol. - Regul. Integr. Comp.*

Physiol. **253**, R701–R711 (1987).

75. M. C. Quinlan, M. J. O'Donnell, Anti-diuresis in the blood-feeding insect

Rhodnius prolixus Stal: Antagonistic actions of cAMP and cGMP and the role of

organic acid transport. *J. Insect Physiol.* **44**, 561–568 (1998).

76. K. W. Beyenbach, A. Oviedo, D. J. Aneshansley, Malpighian tubules of *Aedes*

aegypti: Five tubules, one function. *J. Insect Physiol.* **39**, 639–648 (1993).

77. J. A. Veenstra, Effects of 5-hydroxytryptamine on the Malpighian tubules of *Aedes*

aegypti. *J. Insect Physiol.* **34**, 299–304 (1988).

78. J. P. V. Paluzzi, W. Naikkhwah, M. J. O'Donnell, Natriuresis and diuretic

hormone synergism in *R. prolixus* upper Malpighian tubules is inhibited by the

anti-diuretic hormone, *Rhopr*CAPA-α2. *J. Insect Physiol.* **58**, 534–542 (2012).

79. D. A. Rocco, D. H. Kim, J. P. V. Paluzzi, Immunohistochemical mapping and

transcript expression of the GPA2/GPB5 receptor in tissues of the adult mosquito,

Aedes aegypti. Cell Tissue Res. **369**, 313–330 (2017).

80. A. Lajevardi, F. Sajadi, A. Donini, J. P. V. Paluzzi, Studying the activity of

neuropeptides and other regulators of the excretory system in the adult mosquito.

J. Vis. Exp. **2021** (2021).

81. T. M. Clark, T. J. Bradley, Additive effects of 5-HT and diuretic peptide on *Aedes*

malpighian tubule fluid secretion. *Comp. Biochem. Physiol. - A Mol. Integr.*

Physiol. **119**, 599–605 (1998).

82.   P. V. Pietrantonio, C. Jagge, C. McDowell, Cloning and expression analysis of a

5HT7-like serotonin receptor cDNA from mosquito *Aedes aegypti* female

excretory and respiratory systems. *Insect Mol. Biol.* **10**, 357–369 (2001).

83.   H. L. Lu, C. Kersch, P. V. Pietrantonio, The kinin receptor is expressed in the

Malpighian tubule stellate cells in the mosquito *Aedes aegypti* (L.): A new model

needed to explain ion transport? *Insect Biochem. Mol. Biol.* **41**, 135–140 (2011).

84.   M. R. Rheault, M. J. O'Donnell, Organic cation transport by Malpighian tubules of

Drosophila melanogaster: Application of two novel electrophysiological methods.

J. Exp. Biol. **207**, 2173–2184 (2004).

85.   S. Jonusaite, S. P. Kelly, A. Donini, The physiological response of larval

Chironomus riparius (Meigen) to abrupt brackish water exposure. *J. Comp.*

Physiol. B Biochem. Syst. Environ. Physiol. **181**, 343–352 (2011).

86.   S. D. McCormick, Methods for Nonlethal Gill Biopsy and Measurement of Na^+,

K^+-ATPase Activity. *Can. J. Fish. Aquat. Sci.* **50**, 656–658 (1993).

87.   A. J. Fosang, R. J. Colbran, Transparency is the key to quality. *J. Biol. Chem.* **290**,

29692–29694 (2015).

88.   T. A. J. Butler, J. W. Paul, E. C. Chan, R. Smith, J. M. Tolosa, Misleading

westerns: Common quantification mistakes in western blot densitometry and

proposed corrective measures. *Biomed Res. Int.* **2019** (2019).

89. A. C. Durant, A. Donini, Ammonium transporter expression in sperm of the

disease vector *Aedes aegypti* mosquito influences male fertility. *Proc. Natl. Acad.*

Sci. U. S. A. **117**, 29712–29719 (2020).

90. H. Chasiotis, S. P. Kelly, Occludin immunolocalization and protein expression in

goldfish. *J. Exp. Biol.* **211**, 1524–1534 (2008).

91. J. Schindelin, *et al.*, Fiji: An open-source platform for biological-image analysis.

Nat. Methods **9**, 676–682 (2012).

92. O. I. Kubrak, O. V. Lushchak, M. Zandawala, D. R. Nässel, Systemic corazonin

signalling modulates stress responses and metabolism in *Drosophila*. *Open Biol.* **6**

(2016).

93. L. Kean, *et al.*, Two nitridergic peptides are encoded by the gene capability in

Drosophila melanogaster. Am. J. Physiol. - Regul. Integr. Comp. Physiol. **282**

(2002).

1148

1149