1 **Manuscript title:**
2 The V-type H$^+$-ATPase is targeted in anti-diuretic hormone control of the
3 Malpighian 'renal' tubules
4

Farwa Sajadi[1], María Fernanda Vergara-Martínez[1,2] and Jean-Paul V. Paluzzi[1]*
1. Department of Biology, York University, 4700 Keele Street, Toronto, Ontario, M3J 1P3, Canada.
2. Departamento de Biología Celular y Fisiología, Instituto de Investigaciones Biomédicas, Universidad
Nacional Autónoma de México, Mexico City, Mexico.

*Corresponding author: Jean-Paul V. Paluzzi
Email: paluzzi@yorku.ca

17 **Author Contributions:** J.-P.P. and F.S. designed research; F.S. and M.F.V. performed research; F.S. and
18 J.P.P. analyzed data; and J.-P.P. and F.S. wrote and revised the manuscript.
19 **Competing Interest Statement:** The authors declare no competing interest.
20 **Classification:** Biological Sciences; Physiology
21 **Keywords:** excretory system, *Aedes aegypti*, mosquito, anti-diuresis, proton pump, disease vector

22
23 **This PDF file includes:**
24   Main Text
25   Figures 1 to 6
26
27

Abstract

Like other insects, secretion by mosquito Malpighian tubules (MTs) is driven by the V-type H^+-ATPase (VA) localized in the apical membrane of principal cells. In *Aedes aegypti*, the anti-diuretic neurohormone CAPA inhibits secretion by MTs stimulated by select diuretic hormones; however, the cellular effectors of this inhibitory signaling cascade remain unclear. Herein, we demonstrate that the VA inhibitor bafilomycin selectively inhibits serotonin (5HT)- and calcitonin-related diuretic hormone (DH_{31})-stimulated secretion. VA activity increases in DH_{31}-treated MTs, whereas CAPA abolishes this increase through a NOS/cGMP/PKG signaling pathway. A critical feature of VA activation involves the reversible association of the cytosolic (V_1) and membrane (V_o) complexes. Indeed, higher V_1 protein abundance was found in membrane fractions of DH_{31}-treated MTs whereas CAPA significantly decreased V_1 abundance in membrane fractions while increasing it in cytosolic fractions. V_1 immunolocalization was observed strictly in the apical membrane of DH_{31} treated MTs whereas immunoreactivity was dispersed following CAPA treatment. VA complexes colocalized apically in female MTs shortly after a blood-meal consistent with the peak and post-peak phases of diuresis. Comparatively, V_1 immunoreactivity in MTs was more dispersed and did not colocalize with the V_o complex in the apical membrane at 3 hours post blood-meal, representing a timepoint after the late phase of diuresis has concluded. Therefore, CAPA inhibition of MTs involves reducing VA activity and promotes complex dissociation hindering secretion. Collectively, these findings reveal a key target in hormone-mediated inhibition of MTs countering diuresis that provides a deeper understanding of this critical physiological process necessary for hydromineral balance.

Significance Statement

The V-type H^+ ATPase (VA), or proton pump, provides the driving force for transepithelial ion and fluid secretion in insect Malpighian tubules (MTs). While studies have shown diuretic stimulation activates various signaling pathways, including cAMP and downstream effectors promoting increased VA activity, our understanding of anti-diuretic signaling and its potential regulation of the VA remains rudimentary. Herein, we show that CAPA neuropeptide acts through the NOS/cGMP/PKG pathway to inhibit DH_{31}-stimulated VA activity, supporting the notion that the anti-diuretic regulation is achieved through dissociation of the VA complexes. These results demonstrate a critical role of VA inhibition and trafficking necessary for anti-diuretic signaling and advances our understanding of the complex neuroendocrine control of the MTs in this important human disease-vector mosquito.

Main Text

Introduction

Insect post-prandial diuresis is under rigorous control by neuroendocrine factors (1) acting on the Malpighian 'renal' tubules (MTs) to regulate primary urine production. In the yellow fever mosquito, *Aedes aegypti,* several diuretics have been identified that regulate urine production including serotonin (5HT), calcitonin-related diuretic hormone (DH_{31}), corticotropin-releasing factor-related diuretic hormone (DH_{44}) and leucokinin-related (LK) diuretic hormone (2–5). In female *Aedes* mosquitoes, an anti-diuretic peptidergic neurohormone, CAPA, selectively inhibits DH_{31}- and 5HT-stimulated secretion of MTs (4, 6, 7). Insect CAPA neuropeptides are produced in the central nervous system and are evolutionarily related to neuromedin U in vertebrates (8). In the

fruit fly, *Drosophila melanogaster*, CAPA peptides have been shown to act through a

conserved nitridergic signaling pathway to stimulate diuresis by MTs (9, 10); however, a

few other studies have alluded to an anti-diuretic role (11, 12). In contrast, in both larval

and adult *A. aegypti*, CAPA peptides inhibit fluid secretion through a signaling cascade

involving the NOS/cGMP/PKG pathway (6, 7). Despite this, the anti-diuretic signaling

mechanism and downstream cellular targets, such as the ion channels and transporters,

remain elusive.

In insect MTs, including *A. aegypti*, the bafilomycin-sensitive V-type H^+ ATPase

(VA), also known as the proton pump, functions as an electrogenic pump allowing the

transport of protons from the cytoplasm to the tubule lumen, thus generating a cell-

negative membrane voltage (13, 14). This membrane voltage can then drive secondary

transport processes such as the cation/H^+ exchanger or anion/H^+ cotransporter (15, 16).

Originally found in vacuolar membranes of animals and plants, the VA has since been

found to be essential in cell function in both invertebrates and vertebrates (17). In insects,

the VA is densely located in the apical brush border membrane of tubule principal cells

(13, 18), which is rich in mitochondria and endoplasmic reticula, fueling the ATP-

consuming proton pump (1, 19). Previous studies have shown VA localization within the

apical membrane of principal cells along the entire length of the MTs (19), but absent in

stellate cells that express relatively higher levels of the P-type Na^+/K^+ ATPase (NKA)

(19). Due to stronger VA immunoreactivity observed in MTs (19), and greater ATPase

activity by electrophysiological assays (13, 14), the VA is categorized as serving mainly,

but not exclusively (20), stimulated transport mechanisms, whereas the NKA serves basic

cell housekeeping functions when MTs are undergoing low unstimulated rates of

secretion (13, 20, 21).

Stimulation of distinct diuretic hormone receptors can activate various signaling

pathways, including elevation of cyclic AMP (cAMP) levels which is known to increase

VA activity and assembly in insects (22). *In A. aegypti,* DH_{31}, identified as the mosquito

natriuretic peptide (23), selectively activates transepithelial secretion of Na^+ in the MTs,

using cAMP as a second messenger (1), and upregulating the VA function to stimulate

fluid secretion (24). Similarly, 5HT-stimulated diuresis is also thought to be mediated (at

least in part) through the cAMP second messenger pathway (25), activating protein

kinase A (PKA) to increase the transepithelial voltage of the basolateral membrane in

tubule principal cells (1, 26). In contrast, DH_{44} has been shown to initiate diuresis via the

paracellular and transcellular pathways, with higher nanomolar concentrations increasing

cAMP and Ca^{2+}, influencing both paracellular and transcellular transport, and lower

nanomolar concentrations acting through the paracellular pathway only, via intracellular

Ca^{2+} (27). Thus, due to its predominant role in fluid secretion, the VA could be a likely

target for both diuretic and anti-diuretic factors.

Eukaryotic V-ATPases are a multi-subunit protein composed of up to 14 different

polypeptides, which form two major structural complexes. The peripheral V_1 complex

(400-600 kDa), is invariably present in the cytoplasm and interacts with ATP, ADP, and

inorganic phosphate (28). The cytosolic V_1 complex consists of eight different subunits

(A-H): a globular headpiece with three alternating subunits A and B forming a hexamer

with nucleotide binding sites located at their interface, a central rotor stalk with single

copies of subunits D and F, and lastly, a peripheral stalk made up of subunits C, E, G, and

H (28). The B subunit, shown to have high sequence similarity amongst several species

from fungi to mammals (29, 30), is a 56 kDa polypeptide, and is one of the two sites

(along with subunit A) (31) in the V_1 complex that binds ATP. In contrast, the

membrane-integrated V_o complex (150-350 kDa) mediates the transport of H^+ across the

membrane (28) and is composed of at least six different subunits, which collectively

function in the proton translocation pathway (13, 28). Although the proton channel of the

VA can be blocked pharmacologically by the macrolide antibiotic, bafilomycin (32),

there are two known intrinsic mechanisms for VA regulation: firstly, through oxidation of

the cystine residue on the A subunit of the V_1 complex, thus preventing ATP hydrolysis;

secondly, through reversible disassembly of the V_1 complex from the holoenzyme (13,

33). While the role and regulation of the VA by diuretic hormones in insect MTs has

been studied (34–36), research examining anti-diuretic signaling mechanisms involving

the VA remains in its infancy.

Considering that the *A. aegypti* mosquito is recognized as a disease vector for

several viruses, further investigating the regulatory mechanisms of diuresis and anti-

diuresis in the MTs can allow for useful insights into novel methods to control disease

transmission. This study aimed to identify the cellular targets necessary for CAPA-

mediated inhibition of fluid secretion by MTs stimulated by select diuretic factors in

adult female *A. aegypti*. Our results provide evidence that CAPA neuropeptides inhibit

fluid secretion in the mosquito by VA complex dissociation, thus hindering VA function

and activity that is essential for driving rapid post-prandial diuresis.

Results

Bafilomycin inhibits DH_{31}- and 5HT-stimulated fluid secretion rate.

To determine the appropriate concentration of bafilomycin to test on adult *A. aegypti*

MTs, several doses were applied against DH_{31}-stimulated tubules (**S1A Fig**). Higher

doses of bafilomycin (10^{-4} M and 10^{-5} M) resulted in significant inhibition of fluid

secretion rate, with maximal inhibition leading to a five-fold decrease, observed with

treatment of 10^{-5} M bafilomycin. Next, to determine whether inhibiting the VA would

decrease fluid secretion rate stimulated by other diuretic hormones, including 5HT and

DH_{44}, the effect of 10^{-5} M bafilomycin on adult tubules stimulated with these diuretics

was tested (**Fig 1**). Fluid secretion rates were measured over 30 min under control

(stimulated) conditions, and then at 10-min intervals in the presence of bafilomycin.

Treatment of MTs with 10^{-5} M bafilomycin against DH_{31} led to a decrease of fluid

secretion over the treatment interval. Specifically, 30 min after treatment with

bafilomycin, fluid secretion rate was significantly reduced by over two-fold to

0.438 ± 0.041 nL min^{-1}, compared to DH_{31} alone, 0.941 ± 0.077 nL min^{-1} (**Fig 1A**). Similar

results were seen with 5HT-stimulated MTs; however, a decrease in fluid secretion was

observed 30 min post-bafilomycin, with a significant inhibition 40 min after treatment

(0.522 ± 0.072 nL min^{-1}, 5HT alone vs. 0.182 ± 0.045 nL min^{-1}, 5HT + bafilomycin) (**Fig**

1B). Distinct from DH_{31} and 5HT-stimulated tubules, DH_{44}-stimulated secretion was

insensitive to bafilomycin treatment (**Fig 1C**). To confirm whether *Aedae*CAPA-1 anti-

diuresis is mediated by VA inhibition, adult female MTs were treated with either DH_{31}

alone or in combination with *Aedae*CAPA-1, bafilomycin, or both (**Fig 1D**). MTs treated

with either *Aedae*CAPA-1 or bafilomycin resulted in a significant inhibition of DH_{31}-

stimulated secretion, and similar inhibition was observed when both *Aedae*CAPA-1 and

bafilomycin were applied together with no evidence of any additive inhibitory effects

(**Fig 1D**).

***Aedae*CAPA-1 and bafilomycin alkalinizes secreted fluid in DH$_{31}$- and 5HT-**

stimulated MTs.

The VA pumps protons from the cell into the tubule lumen thus generating an

electromotive potential (20, 31) and providing energy to drive the secretion of cations via

Na^+/H^+ and/or K^+/H^+ antiporters (13). An indirect way to measure whether *Aedae*CAPA-

1 and bafilomycin inhibits VA activity involves measuring the pH of the secreted fluid

from diuretic-stimulated MTs treated with *Aedae*CAPA-1 or bafilomycin (**Fig 1E–1G**).

In DH$_{31}$-stimulated MTs treated with *Aedae*CAPA-1, there was an immediate

significantly higher pH in the secreted fluid (7.479±0.030) at 40 min relative to control,

increasing up to 7.73±0.038 at 60 min (**Fig 1E**). Similarly, pH levels in DH$_{31}$-stimulated

MTs treated with bafilomycin were significantly higher (7.66±0.064) relative to control

at 40 min, increasing up to 7.855±0.074 at 60 min. Comparable to DH$_{31}$, addition of

*Aedae*CAPA-1 or bafilomycin, significantly increased the pH of secreted fluid from 5HT-

stimulated MTs to 7.75±0.061 and 7.82±0.083 respectively, at the 60 min mark (**Fig 1F**).

In contrast, unlike the effects observed with DH$_{31}$- and 5HT-stimulated MTs,

*Aedae*CAPA-1 or bafilomycin did not alter the pH of the secreted fluid in DH$_{44}$-

stimulated MTs (**Fig 1G**). The pH increased from 7.4 to 7.9 during the 30-min DH$_{44}$

incubation; however, pH did not change following the addition of *Aedae*CAPA-1 or

bafilomycin. Separately, we conducted measurements in unstimulated tubules to verify

pH in these small droplets did not drift over a time frame consistent with our above

experiments. Unstimulated MTs were allowed to secrete, and the droplets were isolated

and their pH was measured over the course of 60 min. Over this incubation period, no

change was observed in the pH of secreted droplets from unstimulated MTs (**Fig 1H**),

upholding the notion that the alkalinization of secreted fluid observed following

*Aedae*CAPA-1 (or bafilomycin) treatment of DH_{31}- and 5HT-stimulated MTs is a result

of VA inhibition. Additionally, unstimulated MTs treated with either *Aedae*CAPA-1 or

bafilomycin resulted in no significant changes in either secretion rate (**S1B Fig**) or pH

(**S1C Fig**).

***Aedae*CAPA-1 increases cGMP and decreases cAMP levels in DH_{31}-treated MTs.**

To further clarify the CAPA signaling pathway involving the second messengers, cGMP

and cAMP, we sought to determine changes in levels of these cyclic nucleotides in MTs

incubated in DH_{31} alone or combined with *Aedae*CAPA-1. Treatment of MTs with DH_{31}

alone had basal levels of cGMP, 10.91 ± 0.109 $pmol^{-1}$/tubule, comparable to saline treated

MTs (**Fig 1I**). Treatment of MTs with *Aedae*CAPA-1 resulted in a significant increase in

cGMP levels compared to DH_{31}- incubated tubules, increasing to 11.39 ± 0.101 $pmol^{-}$

^{1}/tubule. Similar results were observed with MTs treated with both DH_{31} + *Aedae*CAPA-

1, with significantly increased cGMP levels of 11.39 ± 0.123 $pmol^{-1}$/tubule (**Fig 1I**)

compared to MTs treated with DH_{31} alone. In contrast, treatment of MTs with DH_{31} alone

led to significantly higher levels of cAMP, 9.153 ± 0.039 $pmol^{-1}$/tubule, while baseline

levels of this second messenger were observed in saline, DH_{31} + *Aedae*CAPA-1, and

*Aedae*CAPA-1 treated tubules (**Fig 1J**). To further confirm the stimulatory role of cAMP

and inhibitory role of cGMP, tubules were treated with either cyclic nucleotide alone and

secretion rates were measured (**Fig 1K**). Unstimulated fluid secretion rates were

measured over the first 30 min, and then at 10-min intervals with either cAMP or cGMP.

Treatment of MTs with 10^{-4} M cAMP led to a significant increase over the treatment

interval, with fluid secretion rates increasing to 0.615 ± 0.096 nL min^{-1} at 60 min,

compared to 10^{-8} M cGMP (0.052 ± 0.091 nL min^{-1}) and unstimulated (0.065 ± 0.091 nL

min^{-1}) (**Fig 1K**). Finally, to establish whether these cyclic nucleotide second messengers

elicit antagonistic control of the MTs in adult *A. aegypti*, tubules were treated initially

with cAMP over the first 30 min and then cGMP was added in the presence of cAMP for

a subsequent 30 min (**Fig 1L**). Similarly, we also tested the opposite treatment regime

where MTs were treated initially with cGMP and subsequently with cAMP added along

with cGMP. Treatment of cAMP-stimulated MTs with 10^{-8} M cGMP led to a significant

decrease (~4-fold) over the treatment interval, with secretion rates decreasing to

0.231 ± 0.113 nL min^{-1} at 60 min, compared to 10^{-4} M cAMP alone (0.931 ± 0.134 nL min^{-}

^{1}). In contrast, cGMP-incubated tubules treated with 10^{-4} M cAMP led to a significant

increase (~10-fold) in secretion rate (0.537 ± 0.072 nL min^{-1}) compared to MTs treated

with 10^{-8} M cGMP alone (0.045 ± 0.018 nL min^{-1}) (**Fig 1L**).

Parallel studies examining cAMP levels were measured in DH$_{44}$-incubated MTs.

Treatment of MTs with DH$_{44}$ alone had high levels of cAMP, $9.115 \pm 0.0.061$ pmol^{-}

^{1}/tubule, compared to saline treated MTs, 8.709 ± 0.081 pmol^{-1}/tubule (**S2A Fig**). Levels

of cAMP remained unchanged in tubules treated with both DH$_{44}$ + *Aedae*CAPA-1,

8.954 ± 0.108 pmol^{-1}/tubule. To further resolve the cAMP signaling pathway downstream

of DH$_{31}$ and DH$_{44}$ stimulated diuresis, a PKA inhibitor (KT5720) was tested against

diuretic-stimulated MTs. KT5720 abolished the stimulatory effect of DH_{31}, whereas

secretion rates by DH_{44}-treated MTs remained unchanged (**S2B Fig**).

In *D. melanogaster*, the effects of CAPA peptides is dependent on both

intracellular and extracellular sources of Ca^{2+} (9, 37). Thus, we sought to test the

involvement of Ca^{2+} in the CAPA intracellular second messenger pathway in the *Aedes*

mosquito. The tubules were tested in a modified Ca^{2+}-free saline containing L-glutamine,

resulting in secretion rates similar to MTs incubated in Ca^{2+}-free saline with Schneider's

medium and *Aedes* saline used previously (**S3A Fig**). Treatment of MTs in the presence

of the Ca^{2+}-chelating agent EGTA had no effect on the inhibition of DH_{31}-stimulated

secretion by *Aedae*CAPA-1 (**S3B Fig**). Similarly, inhibition of intracellular Ca^{2+} levels

with TMB-8 or membrane permeable BAPTA-AM alone or in combination with EGTA

had no effect on the anti-diuretic activity of *Aedae*CAPA-1 (**S3B Fig**).

***Aedae*CAPA-1 decreases VA activity in DH_{31}-stimulated MTs through the**

NOS/cGMP/PKG pathway.

In *Aedes* MTs, 50-60% of the total ATPase activity can be attributed to a bafilomycin-

and nitrate-sensitive component that reflects the activity of the VA pump (13). The

remaining ATPase activity may be due to nucleotide cyclases, protein kinases, myosin,

DNA helicases, and other ATP-consuming processes such as the NKA (13). As such, to

determine whether CAPA inhibits VA and/or NKA function in the female mosquito,

diuretic-stimulated MTs were challenged with *Aedae*CAPA-1 to measure the resultant

NKA and VA activity. As expected, given its established role as the natriuretic diuretic

hormone, adult female MTs treated with DH_{31} resulted in a significant (over two-fold)

increase of VA activity, 0.0329±0.0007 µmoles ADP/µg protein/hour, compared to saline

controls, 0.0151±0.0021 µmoles ADP/µg protein/hour (**Fig 2A**). Importantly, MTs

incubated with both DH$_{31}$ and *Aedae*CAPA-1 had significantly lower VA activity,

resulting in activity levels indistinguishable from saline controls. In contrast, neither 5HT

nor DH$_{44}$ influenced VA activity (p>0.05) when compared with saline controls, while co-

treatment with *Aedae*CAPA-1 also resulted in indistinguishable VA activity. Similar VA

activity levels were observed between 5HT and DH$_{44}$ (0.0255±0.0078 and 0.0208±0.0042

µmoles ADP/µg protein/hour) and with co-application of *Aedae*CAPA-1 (0.0150±0.0036

and 0.0154±0.0070 µmoles ADP/µg protein/hour). Unlike changes observed in VA

activity following treatment with DH$_{31}$, diuretic-stimulation or *Aedae*CAPA-1 treatment

did not perturb NKA activity, with levels similar to unstimulated MTs (**Fig 2B**).

To confirm the actions of CAPA are mediated through the NOS/cGMP/PKG

pathway, pharmacological blockers, including inhibitors of NOS ($_L$-NAME) and PKG

(KT5823), were tested against DH$_{31}$-stimulated MTs treated with either *Aedae*CAPA-1 or

cGMP (**Fig 2C**). Application of $_L$-NAME or KT5823 abolished the inhibitory effect of

*Aedae*CAPA-1, resulting in high levels of VA activity, 0.02671±0.0025 and

0.03653±0.0051 µmoles ADP/µg protein/hour respectively, compared to MTs treated

with DH$_{31}$ + *Aedae*CAPA-1. As expected, treatment of DH$_{31}$-stimulated MTs with cGMP

resulted in a significant decrease in VA activity, 0.006±0.0026 µmoles ADP/µg

protein/hour, similar to *Aedae*CAPA-1-treated MTs, while co-treatment with KT5823,

abolished the inhibitory effect of cGMP, resulting in an increase in VA activity (**Fig 2C**).

AedaeCAPA-1 leads to VA holoenzyme dissociation in DH$_{31}$-treated MTs.

The reversible dissociation of the V_1 complex from the V_o membrane-integrated complex

is a well-known mechanism for regulating VA transport activity (22, 31, 34, 35, 38). To

determine whether *Aedae*CAPA-1 influences VA complex dissociation, membrane and

cytosolic protein fractions were isolated from DH_{31} and DH_{31} + *Aedae*CAPA-1 incubated

MTs, and a polyclonal V_1 antibody (13) was used to measure protein abundance. First,

membrane and cytosolic protein isolation was verified with specific membrane (AQP1,

S4A Fig) and cytosolic (beta-tubulin, **S4B Fig**) markers. Western blot analysis revealed

three protein bands, with calculated molecular masses of 74 kDa, 56 kDa, and 32 kDa

(13, 39) (**S4C Fig**). The V_1 complex is composed of eight subunits (A-H), which includes

the A (~74kDa) and B subunit (~56Da) that are arranged in a ring forming the globular

headpiece for ATP binding and hydrolysis (31). Additionally, studies have suggested that

subunit D (~32kDa) alongside subunit F constitute the central rotational stalk of the V_1

complex (17). There was no difference in abundance observed for the A subunit (74 kDa

band) in either membrane or cytosolic fractions between saline and DH_{31} treatments

whereas DH_{31} + *Aedae*CAPA-1 incubated MTs had increased A subunit (74 kDa band)

protein abundance in cytosolic fractions compared to saline treatment (**Fig 3A**) and

decreased abundance in the membrane fraction compared to MTs treated with DH_{31} alone

(**Fig 3B**). Similarly, the V_1 complex B subunit abundance (56 kDa band) was similar in

all treatments within the cytosolic protein fraction (**Fig 3C**) whereas DH_{31} +

*Aedae*CAPA-1 incubated MTs had significantly lower abundance in membrane fractions

compared to MTs treated with DH_{31} alone (**Fig 3D**). Finally, there was no difference in

abundance of the V_1 complex subunit D (32 kDa band) between saline and DH_{31} treated

MTs in neither cytosolic nor membrane fractions. However, as observed for the A and B

subunit bands (74 and 56 kDa band, respectively) DH_{31} + *Aedae*CAPA-1 incubated MTs

showed a significant increase in subunit D abundance in cytosolic fractions compared to

saline treated MTs (**Fig 3E**) and a decrease in its abundance in membrane fractions

compared to MTs treated with DH_{31} alone (**Fig 3F**). In summary, all three of the V1

complex immunoreactive bands corresponding to subunits A, B and D (74, 56 and 32

kDa, respectively) showed significantly lower abundance in membrane fractions in DH_{31}

+ *Aedae*CAPA-1 incubated MTs.

To visualize this potential endocrine-mediated reorganization of the VA

holoenzyme in this simple epithelium, we immunolocalized the membrane-integrated V_o

and cytosolic V_1 complex in the female *A. aegypti* MTs. Transverse sections of saline-

incubated (control) MTs demonstrated moderate enrichment of V_o (**Fig 4A**, **S5A Fig**),

and V_1 (**Fig 4B**, **S5B Fig**) complexes in the apical membrane of principal cells (**Fig 4C**,

S5C–S5D Fig). Comparatively, DH_{31}-incubated MTs revealed intense localization of the

V_o (**Fig 4D**, **S5E Fig**), and V_1 (**Fig 4E**, **S5F Fig**) complexes within principal cells, where

V_1 staining was strictly co-localized with V_o staining on the apical membrane (**Fig 4F**,

S5G–S5H Fig). Interestingly, although V_o immunolocalization was restricted to the

apical membrane (**Fig 4G**, **S5I-K Fig**), V_1 immunoreactivity was dispersed in both the

apical membrane and cytosolic region (**Fig 4H**, **S5L-N Fig**) in DH_{31} + *Aedae*CAPA-1

treated MTs with little evidence of apical co-localization (**Fig 4I**, **S5O-T Fig**) as

observed in MTs treated with DH31 alone. Immunostaining was absent in control

preparations probed with only secondary antibodies (**S6A-B Fig**) confirming the specific

detection of the VA complexes with each primary antibody. Additionally, to further

resolve the involvement of the cGMP/PKG pathway in VA holoenzyme organization,

DH$_{31}$- and cAMP- stimulated tubules were incubated with cGMP alone or with PKG

blocker, KT5823 (**S7 Fig**). As expected, transverse sections of both DH$_{31}$- and cAMP-

incubated MTs demonstrated strict colocalization of V$_o$ (**S7A, S7J Fig**) and V$_1$ (**S7B,**

S7K Fig) in the apical membrane of principal cells (**S7C, S7L Fig**), while V$_1$ and V$_o$

immunoreactivity was found in both the apical membrane and cytosol regions in cAMP +

cGMP and DH$_{31}$ + cGMP co-treated MTs (**S7D–F, S7M–O Fig**). However,

immunoreactivity of the V$_1$ complex was found in the apical membrane alongside the V$_o$

complex in preparations treated with the PKG blocker, KT5823 (**S7G–I, S7P–R Fig**).

To investigate this endocrine-mediated phenomenon *in vivo*, we immunolocalized

the membrane-integrated V$_o$ and cytosolic V$_1$ complex in blood-fed females at different

time points. Whole body sections of non-blood-fed similarly aged females (control)

demonstrated moderate enrichment of both V$_o$ (**Fig 5A**), and V$_1$ (**Fig 5B**) in the MTs,

with minimal co-localization (**Fig 5C**), resembling saline-incubated MTs. Interestingly,

blood-fed female MTs revealed strong co-localization of the V$_o$ and V$_1$ complexes at 10

min (**Fig 5D–5F**) and 30 min (**Fig 5G–5I**) post blood-meal, whereas V$_1$

immunoreactivity was more dispersed in both the apical membrane and cytosolic area in

MTs 3 hrs post blood-meal (**Fig 5J–5L**), comparable to non-blood-fed females.

Discussion

The MTs of the *Aedes* mosquito are the main organs responsible for the secretion

of water and solutes, thereby contributing towards hydromineral homeostasis of the

animal (40). Active ion transport in *A. aegypti* MTs is accomplished mainly by the V-

ATPases (VA) densely localized in the apical brush-border membrane of principal cells,

that energize the apical and basolateral membrane as well as the paracellular pathway,

allowing for transepithelial secretion of NaCl, KCl, and other solutes (41). In animal

cells, VA localized to the plasma membrane, especially on the apical membrane of

epithelial cells, contribute to extracellular acidification or alkalization, intracellular pH

homeostasis, or energize the plasma membrane for secondary active transport (42, 43). In

insect MTs, the VA plays a major role in fluid secretion, thus serving as a primary target

for both diuretic and, as this study demonstrates, anti-diuretic hormonal regulation of the

mosquito 'renal' tubules. Although the structure and function of the VA has been

elucidated in some detail (13, 20, 22, 31, 44–46), the regulation of the proton pump

remains unclear. Of the various regulatory mechanisms for VA activity, the most studied

is the reversible dissociation of the cytosolic V_1 complex from the membrane-integrated

V_o complex, first established in the midgut of the tobacco hornworm, *Manduca sexta* and

yeast, *Saccharomyces cerevisiae* (45, 47). In this study, the activity and regulation of the

VA was investigated under both diuretic and anti-diuretic hormone control of the adult

female *A. aegypti* MTs. Notably, the current results advance our knowledge of the anti-

diuretic control of the *A. aegypti* MTs, revealing a cellular mechanism for CAPA

inhibition of the MTs by targeting the VA to block fluid secretion stimulated by select

diuretic factors. This includes inhibition of the DH_{31}-related mosquito natriuretic peptide,

which is critical for the post-haematophagy diuresis that eliminates excess water and

sodium originating from the bloodmeal-derived plasma.

In insects, water excretion is tightly regulated to maintain homeostasis of ions and

water (1, 48, 49). Female *A. aegypti* engorge a salt- and water-rich bloodmeal to obtain

the necessary nutrients and proteins for their eggs (1), with about 40% of the ingested

water eliminated in the first hour post feeding (50). The high rates of water excretion

along with the high rates of primary urine production post bloodmeal suggest a highly

coordinated and defined hormonal regulation of the signaling processes and downstream

cellular targets for ion and water transport (51). In *Aedes* MTs, fluid secretion increases at

least three-fold after stimulation with mosquito natriuretic peptide (identified as DH_{31}),

using cAMP as a second messenger (1), activating PKA, which subsequently activates V-

ATPase-driven cation transport processes (22, 35, 39). Herein we show that DH_{31}-

stimulated secretion is inhibited by bafilomycin, which blocks the proton channel of the

VA (32). Moreover, the addition of either bafilomycin or *Aedae*CAPA-1 caused

alkalization of the secreted fluid, indicating inhibition of the VA, which may lead to

constrained entry of cations across the apical membrane through a proposed alkali

cation/proton antiporter (15, 16). Thus, since bafilomycin inhibits DH_{31}-stimulated

secretion, this supports the VA as a target in the inhibition of fluid secretion.

Consequently, the driving force for ion movement and osmotically-obliged water is

reduced, but select Na^+ channels and cotransporters remain unaffected in the presence of

*Aedae*CAPA-1, as observed by the unchanged natriuretic effect of DH_{31} despite reduced

secretion rates in response to *Aedae*CAPA-1 (4). Similar results were seen in 5HT-

stimulated secretion, albeit a partial inhibition. An earlier study demonstrated that Ca^{2+}-

mediated diuresis does not require the assembly and activation of the VA (38, 39). The

cAMP effect on the VA is implemented by protein kinase A (PKA), with inhibitors of

PKA abolishing hormone-induced assembly and activation of the VA (34). Although the

endogenous 5HT receptor expressed within the *A. aegypti* MTs necessary for diuretic

activity remains elusive, in the kissing bug, *Rhodnius prolixus*, both cAMP and Ca^{2+} have

been shown to initiate diuresis in response to 5HT (52), which might explain the partial

inhibitory response of *Aedae*CAPA-1 inhibition on 5HT-stimulated tubules as Ca^{2+}-

mediated diuresis is independent of the VA (39). Notably, the anticipated 5HT type 2

receptor subtype expressed in the principal cells of the MTs is predicted to couple

through a Gq/11 signaling mechanism (53) and likely excludes the type 7 Gs-coupled

receptor localized to tracheolar cells associated with the MTs (54, 55) as well as the type

1 Gi-coupled receptor localized to principal cells in larval stage mosquitoes (56).

Interestingly, DH_{44}-mediated stimulation was observed to be independent of the

VA, as bafilomycin had no effect on the secretion rate or pH of the secreted fluid

following application of this CRF-related diuretic peptide. Previous studies have noted

that low nanomolar concentrations of a DH_{44}-related peptide were linked to the

stimulation of the paracellular pathway only (27), mediating this action through

intracellular Ca^{2+} as a second messenger (57). In contrast, high nanomolar concentrations

of a DH_{44}-related peptide were shown to influence both paracellular and transcellular

transport, increasing intracellular Ca^{2+} and cAMP (57). Although haemolymph

concentrations of diuretic peptides have yet to be determined in mosquitoes, DH_{31} is

immediately released into circulation post bloodmeal, stimulating rapid secretion of Na^+

and excess water (23, 50). In contrast, DH_{44}-stimulated diuresis in *A. aegypti* involves

non-selective transport of Na^+ and K^+ cations (4); this supports a delayed release of this

diuretic hormone post-feeding to maintain production (albeit reduced) of primary urine

whilst conserving Na^+ ions.

In unstimulated adult female *Aedes* MTs isolated *in vitro*, the VA exhibits

variable rates of enzyme activity, consistent with highly variable rates of secretion, as

found also in various other insect species (26, 58, 59). The VA is the main energizer in

MTs as 60% of total ATPase activity can be linked to the VA (13), whereas the NKA,

with around 28% of ATPase activity, also plays a role in membrane energization,

denoting a more important role in the function of MTs than was previously assumed (19,

39–41). Here we show a significant two-fold increase in VA activity in MTs treated with

DH_{31} compared to the unstimulated MTs, with no change in VA activity in 5HT and

DH_{44}-treated MTs. Notably, *Aedae*CAPA-1 treatment blocked the DH_{31}-driven increase

in VA activity, which corroborates the reduced fluid secretion rate and alkalization of the

secreted fluid. Additionally, we sought to establish the importance of the

NOS/cGMP/PKG pathway in the inhibitory actions of *Aedae*CAPA-1 on VA association.

In stimulated MTs treated with *Aedae*CAPA-1 along with NOS inhibitor, $_L$-NAME, or

PKG inhibitor, KT5823, the inhibitory activity of *Aedae*CAPA-1 and its second

messenger cGMP was abolished, resulting in elevated VA activity as a result of DH_{31}

treatment. The present study examined the effects of *Aedae*CAPA-1 on DH_{31}-stimulated

VA activation for the first time in insects. Stimulation of DH_{31} causes an increase in

cAMP production, which activates Na^+ channels and the $Na^+/K^+/2Cl^-$ cotransporter in the

basolateral membrane (60) and up-regulates VA activity (as shown herein and

previously) critical for increased fluid secretion (24). The DH_{31} receptor

(*Aaeg*GPRCAL1) is expressed in a distal-proximal gradient in the MTs, with greater

expression in principal cells where the VA in the apical membrane is highly expressed

(19, 61). The co-localization of the DH_{31} receptor, VA, and cation exchangers (62–64) in

the distal segment of the MTs, along with the CAPA receptor (6), collectively supports

the major roles DH_{31} and CAPA play in post-prandial diuresis and anti-diuresis,

respectively. In contrast to the marked changes in VA activity in response to diuretic and

anti-diuretic hormones, NKA activity remained unchanged in response to treatments

conducted herein. Further studies should examine the potential role of the NKA in

diuretic and anti-diuretic processes.

The reversible dissociation of the V_1 and V_o complexes is currently thought of as

a universal regulatory mechanism of V-ATPases, appearing to be widely conserved from

yeast to animal cells (13, 22, 35, 65). Although previously shown with cAMP (24), it

remained unclear whether other second messengers (eg. Ca^{2+}, cGMP, and nitric oxide)

affect the assembly/disassembly of the V_1V_o complexes in insect MTs. In this study, VA

protein abundance in membrane and cytosolic fractions of MTs was confirmed by

immunoblot analyses. The 56 kDa band represents the B subunit (29), while the 74 kDa

and 32 kDa bands are suggested to be the A and D subunits, respectively, of the V_1

complex (36). The higher abundance of these V_1 complex protein subunits in the

cytosolic fraction and lower abundance in membrane fraction in *Aedae*CAPA-1-treated

MTs provides novel evidence of hormonally-regulated V_1 dissociation from the

holoenzyme in *A. aegypti* MTs. This was further confirmed with V_1 staining found both

in the apical membrane and cytosol of the MTs treated with DH_{31} and *Aedae*CAPA-1 in

contrast to the strict co-localization of the V_1 and V_o complex in the apical membrane of

MTs treated with DH_{31} alone. This strict co-localization of the VA complex in both

DH_{31}- and cAMP-treated MTs was further abolished in the presence of cGMP, however,

rescued with the addition of the PKG inhibitor, KT5823. In unstimulated *A. aegypti* MTs,

40-73% of the V_1 subunits were found to be membrane associated, with reassembly of

the V_1V_o complex observed upon stimulation with cAMP analogues (39). Although

studies have revealed that hormonal regulation can activate the assembly of the

holoenzyme, the signaling mechanisms achieving this control are unclear. In this study,

the data provides strong evidence of VA assembly in DH_{31}-treated MTs, with V_1 complex

protein subunit enrichment found in the membrane fractions, confirming the crucial role

of the VA in DH_{31}-stimulated secretion. Studies in *A. aegypti* have demonstrated the

involvement of PKA in the activation and assembly of the VA upon natriuretic hormone

(i.e. DH_{31}) stimulation and indicate the phosphorylation of the VA subunits by PKA in

the MTs (39). These studies indicate a regulatory role of PKA in VA assembly and its

activation that may be independent or in addition to phosphorylation (39). In line with

these earlier observations, the current results indicate PKA is critical for DH_{31}-stimulated

fluid secretion by MTs.

Together, these results indicate that *Aedae*CAPA-1 binds to its cognate receptor in

principal cells of the adult *A. aegypti* mosquito MTs (6), targets the NOS/cGMP/PKG

pathway (4, 6) to inhibit DH_{31}-mediated elevation of cAMP (23, 60), which blocks PKA-

activated VA association and prevents protons from being pumped across the apical

membrane, resulting in a more alkaline lumen. Additionally, in female *Aedes* MTs, the

effect of CAPA peptides are independent of Ca^{2+}, which is in contrast to *Drosophila*

MTs, where CAPA peptide diuretic activity involves modulation of intracellular

concentrations of Ca^{2+}, which can derive from both extracellular and intracellular sources

of this important signaling molecule (see 9). Our study provides novel evidence that the

anti-diuretic activity of CAPA is mediated through the dissociation of the VA

holoenzyme involving the removal of the V_1 complex from the apical membrane,

hindering luminal flux of protons that in turn starves cation/H^+ exchange, which

ultimately reduces fluid secretion (**Fig. 6**). CAPA peptides are known to elicit both

diuretic and anti-diuretic actions in different insects, whereby a stimulatory role has been

established in *D. melanogaster* (9, 10) and inhibitory in the *R. prolixus* (65) and *D.*

melanogaster (11, 12) indicating a species-specific role of this neuropeptide family,

which may be due to their different diets and lifestyles. Given the unique blood-feeding

stress the female mosquito is subjected to, a carefully controlled mechanism of both

diuretic and anti-diuretic hormones is warranted to rid the haemolymph of excess salts

and water while also preventing the female from excessive secretion.

In *R. prolixus* MTs, the physiological roles of cGMP and cAMP were examined

(66) suggesting cGMP inhibits fluid secretion by activating a phosphodiesterase (PDE)

that degrades cAMP elevated following 5HT and diuretic hormone stimulation of MTs.

Indeed, the current results demonstrated the addition of cAMP reversed the inhibitory

effects of cGMP, while the addition of cGMP reduced the stimulatory response of cAMP,

supporting that these two cyclic nucleotides facilitate two opposing regulatory roles in the

MTs of adult *A. aegypti*. The data herein reveals cGMP levels increase in MTs treated

with CAPA alone or in combination with DH_{31} while cAMP levels decrease in MTs

treated with CAPA in combination with DH_{31} compared to tubules stimulated with DH_{31}

alone, which upholds the roles of cAMP and cGMP in diuretic and anti-diuretic signaling

pathways, respectively. Interestingly, mid-nanomolar concentrations of DH_{44} also led to

increased levels of cAMP, with levels unchanging in response to *Aedae*CAPA-1, raising

doubt regarding the involvement of a PDE. While treatment of a PKA inhibitor, KT5720,

abolished DH_{31}-stimulated secretion, no effect was observed in DH_{44}-mediated

stimulation. Thus, despite DH_{44}-stimulated diuresis was found to involve increased

cAMP, it appears to be PKA-independent. It is well established that the effects of cAMP

are mediated by activation of cAMP-dependent protein kinase (PKA), a major cAMP

target, followed by phosphorylation of target proteins (67). More recently, in *D.*

melanogaster MTs, two distinct cAMP pathways have been elucidated to sustain fluid

secretion; a PKA-dependent pathway, shown to regulate basal fluid secretion in principal

cells; and a PKA-independent pathway, specifically a stimulatory principal EPAC

(exchange proteins directly activated by cAMP) pathway, stimulating fluid secretion

above basal levels (68). Future studies should examine the DH_{44} diuretic pathway leading

to secretion in *A. aegypti* MTs that appears to be PKA-independent and functions in the

absence of VA activation.

In summary, our study highlights a novel target in the anti-diuretic signaling

pathway of adult female *A. aegypti* MTs, emphasizing the intricate and precise regulatory

mechanism of anti-diuresis. Although a plethora studies have investigated the process of

hydromineral balance in terrestrial insects from a diuretic perspective (1, 10, 23, 65, 69–

71), these current findings advance our understanding of anti-diuretic hormone control

while providing further evidence of a previously elusive endocrine regulatory mechanism

of the VA in mosquitoes (**Fig 6**). Given that many insects are recognized as agricultural

pests or disease vectors, further investigating the complex regulation of their ionic and

osmotic balance may aid in lessening their burden on human health and prosperity

through development of improved management strategies that, at least in part, impede

their neuroendocrine control of hydromineral homeostasis.

Materials and Methods

Animal rearing

Eggs of *Aedes aegypti* (Liverpool strain) were collected from an established

laboratory colony described previously (4, 72). All mosquitoes were raised under a 12:12

light:dark cycle. Non-blood fed female insects (three-six days post-eclosion) were used

for bioassays, dissected under physiological saline (*Aedes* saline) adapted from (60) that

contained (in $mmol^{-1}$): 150 NaCl, 25 HEPES, 3.4 KCl, 7.5 NaOH, 1.8 $NaHCO_3$, 1

$MgSO_4$, 1.7 $CaCl_2$, and 5 glucose, and titrated to pH 7.1.

MT fluid secretion assay

In order to determine fluid secretion rates, modified Ramsay assays were

performed as described previously (4, 73). Female adults (3-6 day old) were dissected

under physiological *Aedes* saline prepared as described above, and MTs were removed

and placed in a Sylgard-lined Petri dish containing 20 μL bathing droplets (1:1 mixture of

Schneider's Insect Medium (Sigma-Aldrich): *Aedes* saline, immersed in hydrated mineral

oil to prevent evaporation. The proximal end of each tubule was wrapped around a

Minutien pin to allow for fluid secretion measurements. To investigate the effects of

second messengers, cAMP and cGMP, on fluid secretion rate, 10^{-4} M 8 bromo-cAMP

(cAMP) (23, 66) and 10^{-8} M 8 bromo-cGMP (cGMP) (4) (Sigma-Aldrich, Oakville, ON,

Canada) were used against unstimulated MTs. To test the effects of the pharmacological

blocker KT5720 (7) (protein kinase A (PKA) inhibitor), a dosage of 5 μmol l^{-1}

(manufacturer's recommended dose) was used against 25 nmol l^{-1} DH_{31}- and 10 nmol l^{-1}

DH_{44}-stimulated MTs. To determine the role of second messenger, Ca^{2+}, in the CAPA

signaling pathway, MTs were treated in a modified Ca^{2+}-free saline (with and without 9.1

$mmol^{-1}$ L-glutamine) containing (in $mmol^{-1}$): 150 NaCl, 25 HEPES, 3.4 KCl, 7.5 NaOH,

1.8 $NaHCO_3$, 1 $MgSO_4$, and 5 glucose, and titrated to pH 7.1. The role of Ca^{2+} was tested

using 1 mmol l^{-1} Ca^{2+} chelator EGTA, 50 μmol l^{-1} (in DMSO) membrane-permeable

chelator BAPTA-AM, and 1 nmol l^{-1} intracellular Ca^{2+} chelator 8-(*N,N*-

diethylamino)octyl 3,4,5-trimethoxybenzoate hydrochloride (TMB-8-HCl) (74).

Time course inhibition of bafilomycin

Dosage of bafilomycin A_1 was based on a dose-response analysis of bafilomycin

against DH_{31}-stimulated tubules (S1 Fig). In the interest of determining whether

bafilomycin inhibits the effects of the diuretic factors, dosages of 25 nmol l^{-1} *Drome*DH_{31}

(~84% identical to *Aedae*DH_{31}) (4, 23, 75), 100 nmol l^{-1} 5HT (4, 70, 76), and 10 nmol l^{-1}

*Rhopr*DH (CRF-related diuretic peptide, DH_{44}) (~48% overall identity; ~65% identity

and ~92% similarity within the highly-conserved N-terminal region to *Aedae*DH_{44}) (4,

57, 77, 78), were applied to the isolated MTs. Neurohormone receptors, including those

for 5HT, and the peptides DH_{31}, DH_{44}, and CAPA, are localized to the basolateral

membrane of principal cells (6, 55, 60, 79), while the LK receptor is localized exclusively

to stellate cells (5). As a result, the effects of bafilomycin were tested on diuretics known

to act on the principal cells of the MTs. After incubating with the individual diuretics for

30 min (using the modified Ramsay assay), diuretic peptide was added alone (controls) or

in combination with bafilomycin (final concentration 10^{-5} M). The fluid secretion rate

was recorded every 10 min for a total of 80 min. In order to determine whether inhibition

of the VA was involved in the anti-diuretic activity of CAPA peptides on adult MTs, the

effects of 1 fmol l^{-1} *Aedae*CAPA-1 (4, 6) were investigated in combination with DH_{31}

and bafilomycin.

NKA and VA activities

The Na^+/K^+-ATPase (NKA) and VA activity in the MTs was determined using a

modified 96-well microplate method (80, 81), which relies on the enzymatic coupling of

ouabain- or bafilomycin-sensitive hydrolysis of ATP to the oxidation of reduced

nicotinamide adenine dinucleotide (NADH). The microplate spectrophotometer is

therefore able to directly measure the disappearance of NADH. Adult female MTs (three

to six day old) were dissected and incubated for 30 min in *Aedes* saline, diuretic (DH_{31},

5HT, or DH_{44}) alone or combined with *Aedae*CAPA-1. Following 30-min incubation,

MTs were collected into 1.5 mL microcentrifuge tubes (40-50 sets of MTs per tube =

200-250 MTs per treatment), flash frozen in liquid nitrogen and stored at -80°C. To

investigate the effects of the pharmacological blockers, a nitric oxide synthase (NOS)

inhibitor, N_ω-Nitro-L-arginine methyl ester hydrochloride (L-NAME), and protein kinase

G (PKG) inhibitor, KT5823, were used against DH_{31}-stimulated MTs treated with

*Aedae*CAPA-1 or 10^{-8} M cGMP. Dosages of 2 µmol l^{-1} L-NAME (manufacturer's

recommended dose) and 5 µmol l^{-1} KT5823 were applied to the MTs (4, 6, 7) (see *SI*

Materials and Methods for assay preparation).

Protein processing and western blot analyses

MTs were isolated under physiological saline from 40-50 female *A. aegypti* for
each biological replicate (defined as n = 1) and incubated for 60 min in the following
three treatments: *Aedes* saline, 25 nmol l^{-1} DH_{31}, or 25 nmol l^{-1} DH_{31} + 1 fmol l^{-1}
*Aedae*CAPA-1. Following the incubation, tissues were stored at -80ºC until processing.
To separate the membrane and cytosolic proteins, a membrane protein extraction kit was
used (ThermoFisher Scientific) following recommended guidelines for soft tissue with
minor modifications including 200 μL of permeabilization and solubilization buffer and a
1:200 protease inhibitor cocktail (Sigma Aldrich) in both buffers. Final protein
concentrations were calculated by Bradford assay (Sigma-Aldrich Canada, Ltd.,)
according to manufacturer's guidelines with bovine serum albumin (BioRad
Laboratories) as a standard and quantified using an A_o Absorbance Microplate Reader
(Azure Biosystems) at 595 nm (see *SI Materials and Methods for western blot analyses*).

Immunolocalization of VA complexes in MTs

Immunohistochemistry of the MTs localizing the VA complexes was conducted
following a previously published protocol (82). Adult female MTs (three to six day old)
were dissected out in *Aedes* saline and incubated following similar conditions described
in the western blot section above. After the incubation, the MTs were immersed in
Bouin's solution and fixed for two hours in small glass vials. To test *in vivo* changes of
the VA complexes, five to six day old females were allowed to blood-feed for 20 min
(72), after which female mosquitoes were isolated at 10 min, 30 min, and 3 hr post blood-

meal. Similarly aged, non blood-fed (sucrose-fed) females were isolated as controls.

Following the bloodmeal, whole body females were immersed in Bouin's solution and

fixed for three hours in small glass vials. To test role of second messengers in VA

localization, similarly aged female MTs were incubated in solutions containing 10^{-4} M

cAMP, 10^{-8} M cGMP, and 5 µmol l^{-1} KT5823 before being immersed in Bouin's solution

and fixed for two hours. Tissues/whole body females were then rinsed three times and

stored in 70% ethanol at 4ºC until further processing. Fixed samples were dehydrated

through a series of ethanol washes: 70% ethanol for 30 min, 95% ethanol for 30 min, and

100% ethanol three times for 30 min. The samples were cleared with xylene

(ethanol:xylene for 30 min then 100% xylene three times for 30 min), and infiltrated in

Paraplast Plus Tissue Embedding Medium (Oxford Worldwide, LLC, Memphis, USA) at

xylene:paraffin wax for 60 min at 60ºC, then rinsed in pure paraffin wax twice for 1 hour

for 60ºC. Following the last rinse, the samples were embedded in the paraffin wax and

left to solidify at 4ºC until further processing. Sections (5 µm) were cut using a Leica RM

2125RT manual rotary microtome (Leica Microsystems Inc., Richmond Hill, Canada)

and slides were incubated overnight on a slide warmer at 45ºC for preparation of

immunohistochemistry (see *SI Materials and Methods*).

cGMP and cAMP Measurements

A competitive cGMP ELISA kit (Cell Signaling Technology, #4360) and cAMP

ELISA kit (Cell Signaling Technology, #4339) were used to measure these cyclic

nucleotide second messengers from adult *A. aegypti* MTs following different treatments

(see *SI Materials and Methods for further details*).

Statistical analyses

Data was compiled using Microsoft Excel and transferred to Graphpad Prism

software v.7 to create figures and conduct all statistical analyses. Data was analyzed

accordingly using a one-way or two-way ANOVA and a Bonferroni post-test, or

Student's t-test, with differences between treatments considered significant if $p<0.05$.

Acknowledgments

The authors sincerely thank Prof. Helmut Wieczorek and Prof. Felix Tiburcy (University

of Osnabrück, Germany) for providing the V_1 antibody and Prof. Andrew Donini (York

University, Canada) for providing the *A. aegypti* AQP1 antibody that were used in this

study. The authors are also grateful to Prof. Ian Orchard (University of Toronto

Mississauga, Canada) and Prof. Michael J. O'Donnell (McMaster University, Canada) for

providing synthetic peptides, *Rhopr*DH and *Drome*DH$_{31}$ respectively, used in this study.

The authors would like to thank the reviewers for their insightful comments and feedback

which helped improve an earlier version of this manuscript.

Funding

This research was funded by: Natural Sciences and Engineering Research Council of

Canada (NSERC) Discovery Grant (JPP)**,** Ontario Ministry of Research Innovation Early

Researcher Award (JPP), and NSERC CGS-D (FS) and the Carswell Scholarships in the

Data and materials availability

All data are available in the main text of the article or the supplementary information.

References

1.   K. W. Beyenbach, Transport mechanisms of diuresis in Malpighian tubules of

insects. *J. Exp. Biol.* **206**, 3845–3856 (2003).

2.   T. M. Clark, T. J. Bradley, Malpighian tubules of larval *Aedes aegypti* are

hormonally stimulated by 5-hydroxytryptamine in response to increased salinity.

Arch. Insect Biochem. Physiol. **34**, 123–141 (1997).

3.   G. M. Coast, C. S. Garside, Neuropeptide control of fluid balance in insects, *Ann.*

N.Y. Acad. Sci. **1040**: 1–8 (2005).

4.   F. Sajadi, C. Curcuruto, A. Al Dhaheri, J.-P. V Paluzzi, Anti-diuretic action of a

CAPA neuropeptide against a subset of diuretic hormones in the disease vector,

Aedes aegypti. J. Exp. Biol. (2018).

5.   H. L. Lu, C. Kersch, P. V. Pietrantonio, The kinin receptor is expressed in the

Malpighian tubule stellate cells in the mosquito *Aedes aegypti* (L.): A new model

needed to explain ion transport? *Insect Biochem. Mol. Biol.* **41**, 135–140 (2011).

6.   F. Sajadi, *et al.*, CAPA neuropeptides and their receptor form an anti-diuretic

hormone signaling system in the human disease vector, *Aedes aegypti. Sci. Rep.* **10**

(2020).

7.   A. Ionescu, A. Donini, *Aedes*CAPA-PVK-1 displays diuretic and dose dependent

antidiuretic potential in the larval mosquito *Aedes aegypti* (Liverpool). *J. Insect*

Physiol. **58**, 1299–1306 (2012).

8.   R. Jurenka, The PRXamide neuropeptide signalling system. Conserved in animals,

Adv. Insect Phys. (Academic Press, 2015), pp. 123–170.

9.   S. A. Davies, *et al.*, Signaling by *Drosophila* capa neuropeptides. *Gen. Comp.*

Endocrinol. **188**, 60–66 (2012).

10.   S. Terhzaz, *et al.*, Mechanism and function of *drosophila* capa GPCR: A

desiccation stress-responsive receptor with functional homology to human

neuromedinu receptor. *PLoS One* **7** (2012).

11.   H. A. MacMillan, *et al.*, Anti-diuretic activity of a CAPA neuropeptide can

compromise *Drosophila* chill tolerance. *J. Exp. Biol.* **221**, jeb185884 (2018).

12.   A. R. Rodan, M. Baum, C. L. Huang, The *drosophila* NKCC Ncc69 is required for

normal renal tubule function. *Am. J. Physiol. - Cell Physiol.* **303**, C883–C894

(2012).

13.   X. H. Weng, M. Huss, H. Wieczorek, K. W. Beyenbach, The V-type H^+-ATPase in

Malpighian tubules of *Aedes aegypti*: localization and activity. *J. Exp. Biol.* **206**,

2211–2219 (2003).

14.   H. Wieczorek, K. W. Beyenbach, M. Huss, O. Vitavska, Vacuolar-type proton

pumps in insect epithelia. *J. Exp. Biol.* **212**, 1611–1619 (2009).

15.   H. Wieczorek, The insect V-ATPase, a plasma membrane proton pump energizing

secondary active transport: molecular analysis of electrogenic potassium transport

in the tobacco hornworm midgut. *J. Exp. Biol.* **172**, 335–343 (1992).

16.   H. Wieczorek, M. Putzenlechnere, U. Klein, A Vacuolar-type proton pump

energizes K^+/H^+ antiport in an animal plasma membrane. *J. Biol. Chem.* **266**,

15340–15347 (1991).

17. W. R. Harvey, S. H. P. Maddrell, W. H. Telfer, H. Wieczorek, H[+] V-ATPases

energize animal plasma membranes for secretion and absorption of ions and fluids.

Am. Zool. **38**, 426–441 (1998).

18. Y. Li, *et al.*, RNA-Seq comparison of larval and adult malpighian tubules of the

yellow fever mosquito *Aedes aegypti* reveals life stage-specific changes in renal

function. *Front. Physiol.* **8**, 1–13 (2017).

19. M. L. Patrick, K. Aimanova, H. R. Sanders, S. S. Gill, P-type Na[+]/K[+]-ATPase and

V-type H[+]-ATPase expression patterns in the osmoregulatory organs of larval and

adult mosquito *Aedes aegypti*. *J. Exp. Biol.* **209**, 4638–4651 (2006).

20. K. W. Beyenbach, T. L. Pannabecker, W. Nagel, Central role of the apical

membrane H[+]-ATPase in electrogenesis and epithelial transport in Malpighian

tubules. *J. Exp. Biol.* **203**, 1459–1468 (2000).

21. R. M. Hine, *et al.*, The excretion of NaCl and KCl loads in mosquitoes. *Am. J.*

Physiol. - Regul. Integr. Comp. Physiol. **307**, R837–R849 (2014).

22. P. Dames, *et al.*, cAMP regulates plasma membrane vacuolar-type H[+]-ATPase

assembly and activity in blowfly salivary glands. *Proc. Natl. Acad. Sci. U. S. A.*

103, 3926–3931 (2006).

23. G. M. Coast, C. S. Garside, S. G. Webster, K. M. Schegg, D. A. Schooley,

Mosquito natriuretic peptide identified as a calcitonin-like diuretic hormone in

Anopheles gambiae (Giles). *J. Exp. Biol.* **208**, 3281–3291 (2005).

24. K. Karas, P. Brauer, D. Petzel, Actin redistribution in mosquito Malpighian

tubules after a blood meal and cyclic AMP stimulation. *J. Insect Physiol.* **51**,

1041–1054 (2005).

25. C. Cady, H. H. Hagedorn, Effects of putative diuretic factors on intracellular

second messenger levels in the Malpighian tubules of *Aedes aegypti*. *J. Insect*

Physiol. **45**, 327–337 (1999).

26. D. H. Petzel, P. T. Pirotte, E. Van Kerkhove, Intracellular and luminal pH

measurements of Malpighian tubules of the mosquito *Aedes aegypti*: the effects of

cAMP. *J. Insect Physiol.* **45**, 973–982 (1999).

27. T. M. Clark, T. K. Hayes, K. W. Beyenbach, Dose-dependent effects of CRF-like

diuretic peptide on transcellular and paracellular transport pathways. *Am. J.*

Physiol. - Ren. Physiol. **274** (1998).

28. K. W. Beyenbach, H. Wieczorek, The V-type H$^+$ ATPase: molecular structure and

function, physiological roles and regulation. *J. Exp. Biol.* **209**, 577–589 (2006).

29. F. J. S. Novak, *et al.*, Primary structure of V-ATPase subunit B from *Manduca*

sexta midgut. BBA - Gene Struct. Expr. **1132**, 67–71 (1992).

30. Y. X. Pan, H. H. Gu, J. Xu, G. E. Dean, *Saccharomyces cerevisiae* expression of

exogenous vacuolar ATPase subunits B. *BBA - Biomembr.* **1151**, 175–185 (1993).

31. H. Wieczorek, D. Brown, S. Grinstein, J. Ehrenfeld, W. R. Harvey, Animal plasma

membrane energization by proton-motive V-ATPases. *BioEssays* **21**, 637–648

(1999).

32.   J. Zhang, Y. Feng, M. Forgac, Proton conduction and bafilomycin binding by the

V0 domain of the coated vesicle V-ATPase. *J. Biol. Chem.* **269**, 23518–23523

(1994).

33.   W. J. A. Dschida, B. J. Bowman, The vacuolar ATPase: sulfite stabilization and

the mechanism of nitrate inactivation. *J. Biol. Chem.* **270**, 1557–1563 (1995).

34.   J. Rein, M. Voss, W. Blenau, B. Walz, O. Baumann, Hormone-induced assembly

and activation of V-ATPase in blowfly salivary glands is mediated by protein

kinase A. *Am. J. Physiol. - Cell Physiol.* **294**, C56–C65 (2008).

35.   B. Zimmermann, P. Dames, B. Walz, O. Baumann, Distribution and serotonin-

induced activation of vacuolar-type H^+-ATPase in the salivary glands of the

blowfly *Calliphora vicina*. *J. Exp. Biol.* **206**, 1867–1876 (2003).

36.   H. Wieczorek, G. Grüber, W. R. Harvey, M. Huss, H. Merzendorfer, The plasma

membrane H^+-V-ATPase from tobacco hornworm midgut. *J. Bioenerg. Biomembr.*

31, 67–74 (1999).

37.   P. Rosay, *et al.*, Cell-type specific calcium signalling in a *Drosophila* epithelium.

J. Cell Sci. **110**, 1683–1692 (1997).

38.   M. C. Quinlan, N. J. Tublitz, M. J. O'Donnell, Anti-diuresis in the blood-feeding

insect *Rhodnius prolixus* Stal: The peptide CAP(2b) and cyclic GMP inhibit

Malpighian tubule fluid secretion. *J. Exp. Biol.* **200**, 2363–2367 (1997)..

39.   F. Tiburcy, K. W. Beyenbach, H. Wieczorek, Protein kinase A-dependent and -

independent activation of the V-ATPase in Malpighian tubules of *Aedes aegypti*. *J.*

Exp. Biol. **216**, 881–891 (2013).

40.   K. W. Beyenbach, P. M. Piermarini, Transcellular and paracellular pathways of

transepithelial fluid secretion in Malpighian (renal) tubules of the yellow fever

mosquito *Aedes aegypti. Acta Physiol. (Oxf).* **202**, 387–407 (2011).

41.   K. W. Beyenbach, Energizing epithelial transport with the vacuolar H+-ATPase.

News Physiol. Sci. **16**, 145–151 (2001).

42.   T. Nishi, M. Forgac, The vacuolar (H+)-ATPases - nature's most versatile proton

pumps. *Nat. Rev. Mol. Cell Biol.* **3**, 94–103 (2002).

43.   N. Nelson, W. R. Harvey, Vacuolar and plasma membrane proton-

adenosinetriphosphatases. *Physiol. Rev.* **79**, 361–385 (1999).

44.   K. W. Beyenbach, H. Skaer, J. A. T. Dow, The developmental, molecular, and

transport biology of malpighian tubules. *Annu. Rev. Entomol.* **55**, 351–374 (2010).

45.   P. M. Kane, Disassembly and reassembly of the yeast vacuolar H+-ATPase in vivo.

J. Biol. Chem. **270**, 17025–17032 (1995).

46.   M. Voss, O. Vitavska, B. Walz, H. Wieczorek, O. Baumann, Stimulus-induced

phosphorylation of vacuolar H+-ATPase by protein kinase A. *J. Biol. Chem.* **282**,

33735–33742 (2007).

47.   J. P. Sumner, *et al.*, Regulation of plasma membrane V-ATPase activity by

dissociation of peripheral subunits. *J. Biol. Chem.* **270**, 5649–5653 (1995).

48.   G. Coast, The endocrine control of salt balance in insects. *Gen. Comp. Endocrinol.*

152, 332–338 (2007).

49.   M. J. O'Donnell, J. H. Spring, Modes of control of insect Malpighian tubules:

synergism, antagonism, cooperation and autonomous regulation. *J. Insect Physiol.*

46, 107–117 (2000).

50.   J. C. Williams, H. H. Hagedorn, K. W. Beyenbach, Dynamic changes in flow rate

and composition of urine during the post-bloodmeal diuresis in *Aedes aegypti* (L.).

J. Comp. Physiol. **153**, 257–265 (1983).

51.   C. L. Jagge, P. V. Pietrantonio, Diuretic hormone 44 receptor in Malpighian

tubules of the mosquito *Aedes aegypti*: evidence for transcriptional regulation

paralleling urination. *Insect Mol. Biol.* **17**, 413–426 (2008).

52.   P. Gioino, B. G. Murray, J. P. Ianowski, Serotonin triggers cAMP and PKA-

mediated intracellular calcium waves in Malpighian tubules of *Rhodnius prolixus*.

Am. J. Physiol. - Regul. Integr. Comp. Physiol. **307**, R828–R836 (2014).

53.   F. Sajadi, J. P. V. Paluzzi, Hormonal regulation and functional role of the 'renal'

tubules in the disease vector, *Aedes aegypti*. *Vitamins and Hormones*, (Academic

Press, 2021), pp. 189–225.

54.   P. V. Pietrantonio, C. Jagge, C. McDowell, Cloning and expression analysis of a

5HT7-like serotonin receptor cDNA from mosquito *Aedes aegypti* female

excretory and respiratory systems. *Insect Mol. Biol.* **10**, 357–369 (2001).

55.   D. W. Lee, P. V. Pietrantonio, In vitro expression and pharmacology of the 5-HT7-

like receptor present in the mosquito *Aedes aegypti* tracheolar cells and hindgut-

associated nerves. *Insect Mol. Biol.* **12**, 561–569 (2003).

56. A. Petrova, D. F. Moffett, Comprehensive immunolocalization studies of a putative serotonin receptor from the alimentary canal of *Aedes aegypti* larvae suggest its diverse roles in digestion and homeostasis. *PLoS One* **11**, e0146587 (2016).

57. K. Clark, T M, Hayes, T K., Holman, G.M., Beyenbach, The concentration-dependence of CRF-like diuretic peptide: mechanisms of action. *J. Exp. Biol.* **201**, 1753–1762 (1998).

58. D. H. Petzel, H. H. Hagedorn, K. W. Beyenbach, Preliminary isolation of mosquito natriuretic factor. *Am. J. Physiol. - Regul. Integr. Comp. Physiol.* **18** (1985).

59. J. A. Dow, *et al.*, The Malpighian tubules of *Drosophila melanogaster*: a novel phenotype for studies of fluid secretion and its control. *J. Exp. Biol.* **197**, 421–428 (1994).

60. D. H. Petzel, M. M. Berg, K. W. Beyenbach, Hormone-controlled cAMP-mediated fluid secretion in yellow-fever mosquito. *Am. J. Physiol. - Regul. Integr. Comp. Physiol.* **253** (1987).

61. H. Kwon, H. L. Lu, M. T. Longnecker, P. V. Pietrantonio, Role in diuresis of a calcitonin receptor (GPRCAL1) expressed in a distal-proximal gradient in renal organs of the mosquito *Aedes aegypti* (L.). *PLoS One* **7**, e50374 (2012).

62. W. Kang'ethe, K. G. Aimanova, A. K. Pullikuth, S. S. Gill, NHE8 mediates

amiloride-sensitive Na^+/H^+ exchange across mosquito Malpighian tubules and

catalyzes Na^+ and K^+ transport in reconstituted proteoliposomes. *Am. J. Physiol. -*

Ren. Physiol. **292**, 1501–1512 (2007).

63.   A. K. Pullikuth, K. Aimanova, W. Kang'ethe, H. R. Sanders, S. S. Gill, Molecular

characterization of sodium/proton exchanger 3 (NHE3) from the yellow fever

vector, *Aedes aegypti. J. Exp. Biol.* **209**, 3529–3544 (2006).

64.   P. M. Piermarini, D. Weihrauch, H. Meyer, M. Huss, K. W. Beyenbach, NHE8 is

an intracellular cation/H^+ exchanger in renal tubules of the yellow fever mosquito

Aedes aegypti. Am. J. Physiol. - Ren. Physiol. **296**, 730–750 (2009).

65.   H. Merzendorfer, M. Huss, R. Schmid, W. R. Harvey, H. Wieczorek, A novel

insect V-ATPase subunit M9.7 is glycosylated extensively. *J. Biol. Chem.* **274**,

17372–17378 (1999).

66.   M. C. Quinlan, M. J. O'Donnell, Anti-diuresis in the blood-feeding insect

Rhodnius prolixus Stal: antagonistic actions of cAMP and cGMP and the role of

organic acid transport. *J. Insect Physiol.* **44**, 561–568 (1998).

67.   S. Seino, T. Shibasaki, PKA-dependent and PKA-independent pathways for

cAMP-regulated exocytosis. *Physiol. Rev.* **85**, 1303–1342 (2005).

68.   M. Efetova, *et al.*, Separate roles of PKA and EPAC in renal function unraveled by

the optogenetic control of cAMP levels in vivo. *J. Cell Sci.* **126**, 778–788 (2013).

69.   K. W. Beyenbach, A. Oviedo, D. J. Aneshansley, Malpighian tubules of *Aedes*

aegypti: five tubules, one function. *J. Insect Physiol.* **39**, 639–648 (1993).

70.   J. A. Veenstra, Effects of 5-hydroxytryptamine on the Malpighian tubules of *Aedes*

aegypti. J. Insect Physiol. **34**, 299–304 (1988).

71.   J. P. V. Paluzzi, W. Naikkhwah, M. J. O'Donnell, Natriuresis and diuretic

hormone synergism in *R. prolixus* upper Malpighian tubules is inhibited by the

anti-diuretic hormone, *Rhopr*CAPA-α2. *J. Insect Physiol.* **58**, 534–542 (2012).

72.   D. A. Rocco, D. H. Kim, J. P. V. Paluzzi, Immunohistochemical mapping and

transcript expression of the GPA2/GPB5 receptor in tissues of the adult mosquito,

Aedes aegypti. Cell Tissue Res. **369**, 313–330 (2017).

73.   A. Lajevardi, F. Sajadi, A. Donini, J. P. V. Paluzzi, Studying the activity of

neuropeptides and other regulators of the excretory system in the adult mosquito.

J. Vis. Exp. **2021** (2021).

74.   J. P. Ianowski, J. P. Paluzzi, V. A. Te Brugge, I. Orchard, The antidiuretic

neurohormone *Rhopr*CAPA-2 downregulates fluid transport across the anterior

midgut in the blood-feeding insect *Rhodnius prolixus. Am. J. Physiol. Integr.*

Comp. Physiol. **298**, R548–R557 (2010).

75.   M. Vanderveken, M. J. O'Donnell, Effects of diuretic hormone 31, *droso*kinin, and

allatostatin A on transepithelial K^+ transport and contraction frequency in the

midgut and hindgut of larval *Drosophila melanogaster. Arch. Insect Biochem.*

Physiol. **85**, 76–93 (2014).

76.   T. M. Clark, T. J. Bradley, Additive effects of 5-HT and diuretic peptide on *Aedes*

malpighian tubule fluid secretion. *Comp. Biochem. Physiol. - A Mol. Integr.*

879   *Physiol.* **119**, 599–605 (1998).

880   77.   V. Te Brugge, J. P. Paluzzi, D. A. Schooley, I. Orchard, Identification of the

881   elusive peptidergic diuretic hormone in the blood-feeding bug *Rhodnius prolixus*: a

882   CRF-related peptide. *J. Exp. Biol.* **214**, 371–381 (2011).

883   78.   H. R. Lee, M. Zandawala, A. B. Lange, I. Orchard, Isolation and characterization

884   of the corticotropin-releasing factor-related diuretic hormone receptor in *Rhodnius*

885   *prolixus*. *Cell. Signal.* **28**, 1152–1162 (2016).

886   79.   Overend, G., Cabrero, P., Halberg, K. A., Ranford-Cartwright, L. C., Woods, D. J.,

887   Davies, S. A., and Dow, J. A. T. (2015). A comprehensive transcriptomic view of

888   renal function in the malaria vector, *Anopheles gambiae*. *Insect Biochem. Mol.*

889   *Biol.* **67**, 47–58

890   80.   S. Jonusaite, S. P. Kelly, A. Donini, The physiological response of larval

891   *Chironomus riparius* (Meigen) to abrupt brackish water exposure. *J. Comp.*

892   *Physiol. B Biochem. Syst. Environ. Physiol.* **181**, 343–352 (2011).

893   81.   S. D. McCormick, Methods for nonlethal gill biopsy and measurement of Na^+, K^+-

894   ATPase activity. *Can. J. Fish. Aquat. Sci.* **50**, 656–658 (1993).

895   82.   H. Chasiotis, S. P. Kelly, Occludin immunolocalization and protein expression in

896   goldfish. *J. Exp. Biol.* **211**, 1524–1534 (2008).

Fig. 1. Effect of bafilomycin on fluid secretion rates and pH along with cyclic

nucleotide second messengers on adult *A. aegypti* MTs. Tubules were treated with

either (A) DH_{31} (B) 5HT or (C) DH_{44} and secreted droplets were measured at 10-min

intervals for 30 min. Immediately following measurement of the 30-min point fluid

droplet (solid arrow), MTs were treated with (A) DH_{31} (B) 5HT (C) or DH_{44} alone or in

combination with *Aedae*CAPA-1 or bafilomycin. (D) MTs were treated with DH_{31} alone

or in combination with *Aedae*CAPA-1, bafilomycin, or both *Aedae*CAPA-1 and

bafilomycin for 60 min. Secreted fluid pH was measured in tubules treated with either (E)

DH_{31} (F) 5HT or (G) DH_{44} before and after addition of *Aedae*CAPA-1 or bafilomycin,

along with unstimulated MTs (H). Production of (I) cGMP and (J) cAMP in DH_{31}-

stimulated MTs treated with *Aedae*CAPA-1. (A-C) Significant differences between

bafilomycin-treated and the corresponding time point controls and (E-H) significant

differences in secreted fluid pH between *Aedae*CAPA-1- and bafilomycin-treated and the

corresponding time point controls are denoted by an asterisk, as determined by a two-way

ANOVA and Bonferroni multiple comparison post-hoc test ($p < 0.05$). Data represent the

mean ± standard error (n=12-34), ns denotes no statistical significance. (D, I, J) Bars

labeled with different letters are significantly different from each other (mean± SEM;

one-way ANOVA with Bonferroni multiple comparison, $p < 0.05$, (D) n=7–17) (I, J) n=50

sets of MTs for all treatments (n=3 per treatment)). (K) Fluid secretion rates were

measured at 10 min intervals initially over a 30 min interval (unstimulated) and then over

a second 30 min interval after the addition (solid arrow) of 10^{-4} M cAMP or 10^{-8} M

cGMP. Significant differences between cAMP-treated MTs and the corresponding time

point controls (or cGMP- treated MTs) are denoted by an asterisk (mean± SEM; two-way

ANOVA with Bonferroni multiple comparison, $p < 0.05$, $n = 5$–6). (L) Significant

differences between cAMP-treated MTs and corresponding time point after addition at 30

min (downward arrow) of cGMP are denoted by an asterisk (similar with cGMP alone

and cAMP added at 30 min, mean± SEM; one-way ANOVA with Bonferroni multiple

comparison, $p < 0.05$, $n = 5$–9).

Fig 2. Effect of *Aedae*CAPA-1 and NOS/PKG inhibitors on VA and NKA activity in

diuretic stimulated *A. aegypti* MTs. MTs were incubated in *Aedes* saline, diuretics

(DH_{31}, 5HT, and DH_{44}) alone or in combination with *Aedae*CAPA-1 for 30 min before

collection to measure (A) VA and (B) NKA activity. (C) MTs were treated with

pharmacological blockers, NOS inhibitor (L-NAME) and PKG inhibitor (KT5823) in

combination with either *Aedae*CAPA-1 or cGMP. Bars labeled with different letters are

significantly different from each other (mean± SEM; one-way ANOVA with Bonferroni

multiple comparison, $p < 0.05$). N.S. denotes no statistical significance. For each

treatment, 50 sets of MTs were collected with $n = 3$ biological replicates per treatment.

Fig 3. Membrane and cytosolic protein abundance of the V_1 complex in MTs of *A.*

aegypti. The MTs ($n = 40$–50) were incubated in *Aedes* saline, DH_{31}, or DH_{31} +

*Aedae*CAPA-1 for one hour before collection. Protein abundance was measured in the

(A) 74 kDa band, A subunit (B) 56 kDa band, B subunit and (c) 32 kDa band, D subunit

of the V_1 complex. Individual band densities were normalized to total protein using

Coomassie staining, and graphed relative to saline-treated controls. Bars labeled with

different letters are significantly different from each other (mean± SEM; one-way

ANOVA with Bonferroni multiple comparison, $p<0.05$, n=3–4 replicates).

Fig 4. **Immunolocalization of the V_o and V_1 complexes in transverse sections of**

stimulated *A. aegypti* **MTs.** Representative paraffin-embedded sections of *A. aegypti*

MTs incubated in either (A–C) *Aedes* saline alone, (D–F) DH_{31} and (G–I) DH_{31} +

*Aedae*CAPA-1 for 30 min. Panels (A,D,G) show V_o staining (red), (B,E,H) show V_1

staining (green), and panels (C,F,I) show merged images with staining highly colocalized

in DH_{31} treatment but less evident in saline and *Aedae*CAPA-1 added treatments. Solid

white arrows denote apical VA staining, and empty arrows indicate cytosolic VA

staining. Where visible in sections, DAPI nuclear staining is shown in blue. Scale bar 100

μm, n = 4 biological replicates (SC = stellate cell).

Fig. 5. Immunolocalization of the V_o and V_1 complexes in blood-fed *A. aegypti*

females. Representative paraffin-embedded sections of whole-body non-blood-fed

females (A–C), blood-fed females isolated (D–F) 10 min (G–I) 30 min and (J–L) 3 hr

post blood-meal. Panels (A,D,G,J) show V_o staining (red), (B,E,H,K) show V_1 staining

(green), and panels (C,F,I,L) show merged immunoreactive staining. DAPI nuclear

staining is shown in blue. Scale bar 100 μm, n = 4 biological replicates.

Fig 6. Schematic diagram summarizing the signaling pathway of diuretic and anti-

diuretic control of adult *A. aegypti* MTs. The principal cells in *A. aegypti* MTs are

responsible for the transport of Na^+ and K^+ cations via secondary active transport,

energized by the V-type H^+-ATPase (VA), localized in the brush border of the apical

membrane. The movement of protons creates a gradient, driving the exchange of Na^+ and

K^+ through cation/H^+ antiporters (NHA). Neurohormone receptors for DH_{31}, 5HT, DH_{44},

and CAPA are localized to the basolateral membrane of the principal cells, while the

kinin receptor is localized exclusively in the stellate cells. The current results together

with previous data indicates that DH_{31} stimulates diuresis through activation and

assembly of the VA in the apical membrane, with no effect on the Na^+/K^+-ATPase

(NKA). The anti-diuretic effect of *Aedae*CAPA-1, facilitated by the NOS/cGMP/PKG

pathway, causes V_1 dissociation from the membrane, hindering activity, and thus

reducing fluid secretion. The biogenic amine, 5HT, has also been shown to stimulate

activation of the VA, however, to a lesser extent. DH_{44}-related peptide receptor activation

increases Ca^{2+} (as well as cAMP at higher doses) but its action was found to be

independent of PKA and VA. Lastly, it was shown earlier that cGMP inhibits kinin-

stimulated diuresis, suggesting an additional anti-diuretic factor may exist that acts

specifically on stellate cells.

| | Non-fed | 10min PBF | 30min PBF | 3hr PBF |