An *arginase 2* promoter transgenic illuminates anti-inflammatory signalling in

zebrafish

Ffion R. Hammond[1]*, Amy Lewis[1]*, Holly E. Anderson[1], Lewis G. Williams[1], Annemarie H.

Meijer[2], Geert F. Wiegertjes[3] and Philip M. Elks[1]

[1]The Bateson Centre, Department of Infection and Immunity and Cardiovascular Disease,

University of Sheffield, Western Bank, Sheffield, UK.

[2]Institute of Biology Leiden, Leiden University, Einsteinweg 55, 2333 CC, Leiden, The

Netherlands.

[3]Aquaculture and Fisheries Group, Department of Animal Sciences, Wageningen University

& Research, 6700 AH Wageningen, The Netherlands.

*These authors contributed equally.

Corresponding author:

Dr Philip M. Elks

The Bateson Centre,

University of Sheffield,

Firth Court,

Western Bank,

Sheffield,

South Yorkshire,

S10 2TN.

UK

Tel: +44 (0) 1142 223609

p.elks@sheffield.ac.uk

Abstract

The innate immune response to inflammatory stimuli must be finely balanced to

produce an appropriate pro-inflammatory response while allowing a subsequent return to

homeostasis. In recent years, *in vivo* transgenic zebrafish models have shed light on the

temporal regulation of the pro-inflammatory innate response to immune challenges.

However, until now, there have been no zebrafish transgenic models of anti-inflammatory

signalling. We compared existing expression data of arginase genes in zebrafish neutrophils

and macrophages, strong candidates for an anti-inflammatory marker, and identified that

arginase 2 is the most highly expressed Arginase in zebrafish immune cells. We developed

an *arginase 2* (*arg2*) bacterial artificial chromosome (BAC) transgenic line,

TgBAC(arg2:eGFP)sh571, driving GFP expression under the control of the *arg2* promoter.

We show that, under resting conditions, *arg2:GFP* is expressed in ionocytes, matching the *in*

situ hybridisation pattern. Upon immune challenge by injury, bacterial and fungal insults,

arg2:GFP is predominantly expressed in neutrophils at early timepoints post-insult. Later in

infections, *arg2:GFP* is expressed in cells associated with foci of infection (including

neutrophils and macrophages), alongside liver expression. Our data indicate that *arginase 2*

is predominantly expressed in neutrophils after immune challenge and suggest that anti-

inflammatory signals coincide with pro-inflammatory signals during early wound and infection

responses.

Introduction

Initial immune responses to inflammatory or infection stimuli are mediated by innate

immune cells, of which leukocytes are major players. Two important leukocyte cell types,

neutrophils and macrophages, work together to neutralise invading threats whilst promoting

tissue healing and restoration of homeostasis. Neutrophils and macrophages have evolved

together and co-exist in invertebrate and vertebrate species [1]. Neutrophils are often

observed to be the first immune cell type to respond to immune challenge, rapidly migrating

towards the stimuli and becoming activated to a pro-inflammatory and antimicrobial state,

clearing damaged cells and invading pathogens. Macrophages also rapidly respond to

challenge, with pro-inflammatory phenotypes emerging soon after immune challenge aiding

microbial clearance (often termed M1 or classical activation). Dogmatically, neutrophils are

considered a blunt first line of defense that once activated remain so until cleared from

tissues, either by apoptosis and subsequent efferocytosis by macrophages or by migration

away [2–4]. Persistence of neutrophils during disease can cause considerable bystander

damage to healthy surrounding tissues driving chronic pathologies. Macrophage

phenotypes, on the other hand, are well-characterised as being plastic throughout the

pathogenesis of inflammation and infections in a process termed macrophage polarisation.

Pro-inflammatory macrophage phenotypes are followed by anti-inflammatory phenotypes

that promote healing and restoration of homeostasis (often termed M2 or alternatively

activated). Many of the above observations come from *in vitro* approaches. *In vivo*

exploration indicates that macrophage polarisation is not binary but rather a spectrum of

phenotypes and behaviours, and that the neutrophil response is more plastic than previously

thought, with emerging roles for neutrophil subsets in tissue protection and healing [5–7].

Both neutrophils and macrophages are activated by danger and pathogen associated

molecular patterns (DAMPs and PAMPs), which trigger the production of pro-inflammatory

signals (e.g. IL-1β and TNF-α) after immune challenge [8,9]. These activate the production

of antimicrobial molecules, including reactive nitrogen species (RNS), via the enzyme

inducible nitric oxide synthase (iNOS) [10]. Immunomodulatory signals are required to limit

and eventually turn-off the pro-inflammatory response in order for tissues to regain

homeostasis. The best characterised of these anti-inflammatory signaling molecules are

cytokines including the interleukins (IL), IL-4, IL-13 and IL-10, alongside proteins that

dampen the pro-inflammatory response, including Arginase [11]. Despite their classification

as anti-inflammatory, these signals have also been identified as being upregulated in some

pro-inflammatory situations [12]. The production of antimicrobial RNS is negatively regulated

by the enzyme Arginase. Arginase competes with iNOS for a shared substrate, L-arginine, a

limited resource within the cell. The immunomodulatory properties of Arginase extend

beyond the regulation of RNS production. Arginase 2 is essential for IL-10 mediated

downregulation of pro-inflammatory factors making it a key anti-inflammatory enzyme [13].

Arginase and iNOS have been well-characterised in murine models as being strongly

expressed in macrophage subtypes, with Arginase being a wound-healing/anti-inflammatory

macrophage marker and iNOS a pro-inflammatory macrophage marker [14]. In human

macrophages the distinction between iNOS and Arginase expressing macrophage subtypes

is less well-defined, due, in part, to lower macrophage RNS in humans compared to mice.

Interestingly, human neutrophils constitutively express Arginase with levels increasing after

infection *in vitro*, but its roles are poorly understood *in vivo* [15]. Considering that Arginase

is not expressed highly by murine neutrophils [16,17], alternative *in vivo* models are required

to investigate the roles of Arginase in immunomodulation and human disease.

The zebrafish has become a powerful model organism to determine the molecular

mediators of immunity against injury and pathogenic challenges [18,19]. Advantages of the

zebrafish model include transparent larvae combined with fluorescent transgenic lines,

allowing detailed microscopy in an intact organism *in vivo*. A key turning-point in zebrafish

immunity research came with the development of transgenic lines marking neutrophil and

macrophage cell populations. Initially, transgenic lines were developed that labelled whole

populations of immune cells in zebrafish larvae, e.g., *TgBAC(mpx:GFP)i114* labelling the

total neutrophil population and *Tg(mpeg1:GFP)* labelling the total macrophage population

[20,21]. More recently, transgenic lines of important pro-inflammatory cytokines have

become available (e.g., *TgBAC(tnfa:GFP)pd1028* and *TgBAC(il-1beta)sh445*)) allowing in-

depth analysis of the cells producing these important signals following a variety of immune

challenges *in vivo* [22,23]. These transgenic lines utilised bacterial artificial chromosome

(BAC) technology, that allows tens of kilobases of promoter region to be used to drive the

fluorescent protein, ensuring that the expression of the transgene recapitulates endogenous

expression patterns as closely as possible. A key gap in this zebrafish "transgenic toolbox" is

an anti-inflammatory fluorescent transgenic line. We therefore set out to develop an *arginase*

2 promoter driven BAC transgenic line to understand the *arginase* response to immune

challenge *in vivo*.

Here, we compared existing expression data of *arginase* genes in zebrafish

neutrophils and macrophages and show that *arg2* is the most highly expressed *arginase* in

zebrafish immune cells. We developed a new BAC transgenic line that drives GFP under the

control of the *arginase 2* promoter and demonstrate that the *arg2:GFP* transgene is

expressed in ionocytes, a population of skin resident cells, in resting conditions. Following a

range of immune challenges, including tailfin transection (sterile injury), *Mycobacterium*

marinum (bacterial), and *Cryptococcus neoformans*/*Candida albicans* (fungal) infections,

arg2:GFP was predominantly upregulated in neutrophils. We identify a small population of

macrophages that express *arginase 2* after injury and infection suggesting the presence of

anti-inflammatory macrophages in zebrafish. The *arg2:GFP* transgenic line has the potential

to uncover new mechanisms behind innate immune regulation during *in vivo* immune

challenge.

Materials and Methods

Ethics statement and zebrafish

All zebrafish were raised in the Biological Services Unit (BSU) aquarium (University

of Sheffield, UK) and maintained according to standard protocols (zfin.org) in Home Office

approved facilities. All procedures were performed on embryos less than 5.2 dpf (days post

fertilisation) which were therefore outside of the Animals (Scientific Procedures) Act, to

standards set by the UK Home Office. Adult fish were maintained at 28°C with a 14/10 hour

light/dark cycle. To investigate expression in immune cells the *TgBAC(arg2:eGFP)sh571*

(hereon in termed *arg2:GFP*) reporter line was crossed with the macrophage reporter

Tg(mpeg1:mCherryCAAX)sh378 [24] (hereon in termed *mpeg:mCherry*) and the neutrophil

reporter *Tg(lyz:nfsB.mCherry)sh260* [25] (hereon in termed *lyz:mCherry*) to generate

embryos for experiments.

Generation of *TgBAC(arg2:eGFP)sh571* transgenic zebrafish

An eGFP SV40 polyadenylation cassette was inserted at the *arg2* ATG start site of the zebrafish bacterial artificial chromosome (BAC) CH-211-12d10 using established protocols [21]. Inverted Tol2 elements were inserted into the chloramphenicol coding sequence and the resulting modified BAC containing 115130 base pairs of the *arg2* promoter region was used. We identified two founder zebrafish (allele codes sh571 and sh572) and raised colonies. The embryos of both alleles had the same GFP expression pattern. The data generated in this manuscript is from the *TgBAC(arg2:eGFP)sh571* transgenic line (hereon in termed *arg2:GFP*) as this line had a higher fecundity than *TgBAC(arg2:eGFP)sh572*.

Tailfin transection

To induce an inflammatory stimulus, 2- or 3-days post fertilisation (dpf) zebrafish were anaesthetised in 0.168 mg/ml Tricaine (Sigma-Aldrich) in E3 media and visualised under a dissecting microscope. Using a scalpel blade (5mm depth, WPI) the tailfin was transected after the circulatory loop as previously described ensuring the circulation remained intact [26].

Pathogen strains and culture

Bacterial infection experiments were performed using *Mycobacterium marinum* strain M (ATCC #BAA-535), containing the pSMT3-Crimson vector, with liquid cultures prepared from bacterial plates [27]. Liquid cultures were washed and prepared in 2% polyvinylpyrrolidone40 (PVP40) solution as previously described for injection [28,29]. Injection inoculum was prepared to 100 colony forming units (cfu)/nl for all *Mm* experiments, which was injected into the circulation at 30hpf via the caudal vein.

Fungal infection experiments were performed using the *Candida albicans* strain TT21-*mCherry* [30]. Overnight liquid cultures were grown from fungal plates, then prepared

for injection as previously described [30]. Cultures were counted using a hemocytometer and

prepared in 10% PVP40 for 200cfu/nl injection dose which was injected into the circulation at

30hpf via the caudal vein.

Fungal infection experiments were also performed using the *Cryptococcus*

neoformans strain Kn99-*mCherry* [31]. Cryptococcal culture was performed as previously

described [24] and after counting on a hemocytometer, Kn99 was prepared in 10% PVP40

for 200cfu/nl injection dose which was injected into the circulation at 1-2dpf.

Microinjection of zebrafish larvae

Prior to injection, zebrafish were anaesthetised in 0.168 mg/ml Tricaine (MS-222,

Sigma-Aldrich) in E3 media and transferred onto 1% agarose in E3+methylene blue plates,

removing excess media. All pathogens were injected into the circulation to create systemic

infection, using a microinjection rig (WPI) attached to a dissecting microscope. A 10mm

graticule was used to measure 1nl droplets for consistency, and droplets were tested every

5-10 fish and recalibrated if necessary. Final injection volume of 1nl was injected to produce

doses calculated for each pathogen. After injection, zebrafish were transferred to fresh E3

media for recovery and maintained at 28°C.

Whole mount *in situ* hybridisation

RNA probes for zebrafish arginase type II (*arg2*, ENSDARG00000039269. Plasmid

obtained from Source Bioscience) were designed and synthesised after cloning into the

pCR™Blunt II-TOPO® vector according to manufacturer's instructions (ThermoFisher

Scientific). Plasmids were linearised and probes synthesised according to DIG RNA

Labeling Kit (SP6/T7) (Roche). Zebrafish larvae were fixed in 4% paraformaldehyde solution

(PFA, ThermoFisher Scientific) overnight at 4°C. Whole mount *in situ* hybridisation was

performed as previously described [32].

Confocal microscopy

Control, tailfin transected, and infected larvae were imaged using a Leica DMi8 SPE-

TCS microscope using a HCX PL APO 40x/1,10 water immersion lens. For confocal

microscopy larvae were anaesthetised in 0.168 mg/ml Tricaine and mounted in 1% low

melting agarose (Sigma) containing 0.168 mg/ml tricaine (Sigma) in 15µ-Slide 4 well glass

bottom slides (Ibidi).

Stereo microscopy

Zebrafish larvae were anaesthetised in 0.168 mg/ml Tricaine and transferred to a

50mm glass bottomed FluoroDish™ (Ibidi). Zebrafish were imaged using a Leica DMi8 SPE-

TCS microscope fitted with a Hamamatsu ORCA Flash 4.0 camera attachment using a HC

FL PLAN 2.5x/0.07 and HC PLAN APO 20x/0.70 dry lens. Both transgenic zebrafish and

whole mount *in situ* staining was imaged using a Leica MZ10F stereo microscope fitted with

a GXCAM-U3 series 5MP camera (GT Vision).

Lightsheet microscopy

2 and 3dpf larvae were imaged using a Zeiss Z1 lightsheet microscope with Plan-

Apochromat 20x/1.0 Corr nd=1.38 objective, dual-side illumination with online fusion and

activated Pivot Scan at 28°C chamber incubation. Zebrafish were anaesthetised in 0.168

mg/ml Tricaine and mounted vertically in 1% low melting agarose (Sigma) in a glass

capillary. Images were obtained using 16bit image depth, 1400x1400 pixel field of view and

GFP visualised with 488nm laser at 16% power, 49.94ms exposure and user-defined z-stack

depth (400-600 slices, 0.641µm slices).

Data analysis

Microscopy data was analysed using Leica LASX (Leica Microsystems) and Image J

software. Graphs were generated using Prism 8.0, GraphPad Software (San Diego, CA,

USA).

Results

***TgBAC(arg2:GFP)sh571* is expressed in ionocytes in resting conditions**

There are two isozymes of Arginase in most mammals and fish, Arginase 1 (ARG1)

and Arginase 2 (ARG2). In mice, ARG1 and ARG2 are both expressed by macrophages,

however it is ARG1 that is the most widely studied in macrophage polarisation, with

increased expression of cytosolic ARG1 protein depleting intracellular stores of arginine [33].

In fish, Arginases have been studied in common carp (*Cyprinus carpio*), a species

phylogenetically close to zebrafish, where *arg2* is the most highly expressed of the Arginase

family in immune cells [11,34]. Head-kidney derived macrophages of carp which have been

polarised towards anti-inflammatory phenotypes using cAMP have a 16-fold upregulation of

arg2 expression [11,35].

Zebrafish have orthologues of mammalian Arginase 1 and Arginase 2, which were

compared using existing transcriptomics datasets to identify innate immune cell expression.

FACS purified neutrophils (*Tg(mpx:GFP)i114*) and macrophages (*Tg(mpeg1:Gal4-*

VP16)gl24/(UAS-E1b:Kaede)s1999t) from unchallenged 5dpf larvae both expressed

arginase2 (*arg2*), while *arginase 1* (*arg1*) was not expressed at detectable levels

(Supplemental Figure 1A-B, using raw data from [36]). *arg2* expression was found at high

levels in the bulk, non-immune cell, population alongside being expressed in the immune cell

populations (Supplemental Figure 1C-D, data from [36]). *arg2* expression was approximately

1.5-fold higher in neutrophils than macrophages in unchallenged zebrafish larvae

(Supplemental Figure 1C-D, using raw data from [36]). In unchallenged adult zebrafish,

single-cell RNAseq identified *arg2* expression in a population of neutrophils, a smaller

population of monocytes/macrophages, with no detectable expression in other blood

lineages (thrombocytes/erythrocytes) (Supplemental Figure 1E-F, data from The Zebrafish

Blood Atlas, https://www.sanger.ac.uk/science/tools/basicz/basicz/, [37]). From the same

dataset *arg1* was not detected in any immune cell lineage (Supplemental Figure 1E-F).

Based on the predominant expression of *arg2* in fish innate immune cells, we chose

to develop an *arg2* transgenic zebrafish line to investigate its expression during immune

challenge *in vivo*. We adopted a BAC transgenesis approach, using a BAC (CH-211-12d10)

in which 11.5kb of the *arg2* promoter drives GFP expression, to generate two transgenic line

alleles with the same expression (*TgBAC(arg2:eGFP)sh571* and *TgBAC(arg2:eGFP)sh572*).

Due to the higher fecundity in the *sh571*, it was this line that was used in the following

studies (hereon in termed *arg2:GFP*).

In unchallenged *arg2:GFP* larvae, the transgene was expressed in cuboidal-shaped

cells in the skin, distributed over the yolk and caudal vein regions at 2 days post fertilisation

(dpf, Figure 1A) and 3dpf (Figure 1B-C). A subset of ionocytes, cells in the skin responsible

for the transport of sodium ions also known as H^+-ATPase-rich cells (HRCs), have been

shown to express high levels of *arginase 2* by *in situ* hybridisation in zebrafish [38–40].

arg2:GFP expression recapitulated the *arginase 2 in situ* hybridisation pattern, labelling

ionocytes (Figure 1D-E). To determine whether any of the cells over the yolk area

expressing *arg2:GFP* were leukocytes, we crossed the *arg2:GFP* line with a macrophage

transgenic line (*Tg(mpeg1:mCherry)sh378*, hereon in termed *mpeg:mCherry*) and a

neutrophil transgenic line (*Tg(lyz:nsfB.mCherry)sh260*, hereon in termed *lyz:mCherry*).

Under resting conditions there was no overlap between *mpeg:mCherry* positive

macrophages and *arg2:GFP* positive cells at 2dpf, 3dpf, 4dpf or 5dpf (Figure 1F-G). Nor was

there overlap between *lyz:mCherry* positive neutrophils (Figure 1H). These data indicate that

the *arg2:GFP* line is not expressed at detectable levels in immune cells in resting conditions

at these developmental timepoints, matching *in situ* hybridisation data. The zebrafish

arg2:GFP transgene and *in situ* hybridisation did not detect leukocyte *arginase 2* expression

in resting conditions, however *arginase 2* levels were detected by RNAseq of FACS purified

leukocyte populations (Supplemental Figure 1). This suggests that either the transgenic and

in situ hybridisation techniques were not as sensitive as RNAseq, or that techniques to purify

immune cells in RNAseq studies have led to upregulation of *arg2* not present in the intact

zebrafish.

280   ***arg2:GFP* is predominantly upregulated by neutrophils after tailfin transection**

To assess whether *arg2:GFP* expression is upregulated in innate immune cells

during an immune response, we challenged 2dpf larvae with a sterile tailfin wound. Using a

tailfin nick model we identified that highly mobile immune cells migrating towards the wound

expressed *arg2:GFP* within the first hour of timelapse microscopy (Figure 2A). We reasoned

that these cells were neutrophils, due to their size and amoeboid shape alongside their rapid

migration towards the wound. To investigate whether neutrophils expressed *arg2:GFP* after

injury, we crossed the *arg2:GFP* line with the neutrophil *lyz:mCherry* transgenic line.

Timelapse microscopy demonstrated that a subpopulation of neutrophils arriving at the tailfin

transection wound were *arg2:GFP* positive by 1 hour post wound (hpw) (Figure 2B) with

positive neutrophils at the site of injury present during the recruitment phase of inflammation

(1-6hpw) while expression was not observable in neutrophils away from the wound. There

was no expression of *arg2:GFP* observed in *mpeg:mCherry* positive macrophages in the

recruitment phase of inflammation between 1-6hpw (Figure 2C).

In order to observe potential anti-inflammatory macrophages, *mpeg:mCherry* larvae

crossed into the *arg2:GFP* line were imaged at a later timepoint, 3dpw (5dpf), by which time

the fin has partially regenerated. In uninjured larvae there were few *mpeg:mCherry* positive

macrophages in the tailfin fold at 5dpf and these did not express *arg2:GFP* (Figure 2D, top

panel). In tailfin transected larvae there were increased numbers of macrophages in the

2dpw and 3dpw healing/regenerating tailfin fold, but the majority of these were *arg2:GFP*

negative, while *mpeg:mCherry* negative cells with the morphology of neutrophils did express

arg2:GFP (Figure 2D, bottom two panels). Upon closer investigation using confocal

microscopy, *mpeg:mCherry* positive cells expressing *arg2:GFP* were identified at the wound

at 2dpw and 3dpw (Figure 2E), though these were greatly outnumbered by *arg2:GFP*

negative macrophages. *arg2:GFP* positive macrophages were not observed in uninjured

larvae. Together these data hint at the presence of *arginase 2* expressing anti-inflammatory

macrophages during tailfin regeneration.

Infection challenge upregulates neutrophil *arg2:GFP* expression

To investigate the expression of *arg2* after bacterial infection, *arg2:GFP* embryos were injected with *Mycobacterium marinum* (Mm) at 1dpf. We assessed the early response to Mm infection at 1dpi and found that a subpopulation of neutrophils express *arg2:GFP* early in infection (Figure 3A). Neutrophil *arg2:GFP* expression was not observed in mock-infected (PVP) controls (Figure 3B) with ionocyte expression confirming *arg2:GFP* positive larvae. *arg2:GFP* positive neutrophils were present in the vicinity of Mm infection (Figure 3A), with both infected and uninfected neutrophils expressing *arg2:GFP* (Figure 3C). A subset of neutrophils had *arg2:GFP* expression, with 31.7% (of n=123 neutrophils) of *lyz:mCherry* positive neutrophils in the region of infection expressing GFP, suggestive of differential immune responses between individual neutrophils (Figure 3D-E). At the same 1dpi timepoint, 6.1% (of n=98 macrophages) of *mpeg:mCherry* positive macrophages were positive for *arg2:GFP* (Figure 3E-G).

Fungal infections have been shown to modulate host arginine metabolism via Arginase [41], therefore we assessed *arg2:GFP* expression in two well-characterised fungal zebrafish infection models - *Candida albicans* [42] and *Cryptococcus neoformans* [24]. In both fungal infections, *arg2:GFP* was observed in a subpopulation of *lyz:mCherry* positive neutrophils at 1dpi (Figure 4A-B). As with Mm infection, subsets of neutrophils both with or without internalised pathogen were *arg2:GFP* positive in *Candida albicans* infection (Figure 4A). *arg2:GFP* positive *mpeg:mCherry* macrophages were also observed in *Cryptococcus neoformans* infection at 1dpi, however these were greatly outnumbered by negative macrophages and only 3 examples were identified (Supplemental Figure 2). In Cryptococcal infected larvae with a high fungal burden, *arg2:GFP* expression was observed in the liver (Figure 4C-D) at timepoints when this was not present in PVP injected larvae (Figure 4C). This was especially apparent in Cryptococcus infections at 2dpi (Figure 4D) and was confirmed by *in situ* hybridisation (Figure 4E). Arginase is a well characterised liver enzyme [43], yet in unchallenged embryos (Figure 1A-E) or PVP injected larvae (Figure 4C) there

was no visible liver expression of *arg2:GFP*. Liver expression was also present after Mm

infection, but was less pronounced (Supplemental Figure 3). Like Cryptococcal infected

larvae this occurred in individuals highly infected with Mm, but was not observed till later in

infection (at 4dpi), potentially reflecting the slower doubling time/pathogenesis of Mm

compared to Cryptococcus.

arg2:GFP is expressed in cells associated with developing granulomas

arg2:GFP expression was assessed at a later stage of Mm infection (4dpi), a stage

at which innate immune granulomas are forming and when it is likely that leukocyte

phenotypes are more diverse and polarised due to immune modulation by mycobacteria

[44]. PVP control injected larvae had minimal *arg2:GFP* expression, apart from ionocyte

expression and background green signal from the autofluorescence of pigment cells (Figure

5A). In Mm infected larvae, *arg2:GFP* was expressed in cells associated with developing

granulomas at 4dpi (Figure 5B). The brightest population of these cells were *lyz:mCherry*

positive, indicating a subset of *arg2:GFP* positive granuloma-associated neutrophils,

representing 37.5% of neutrophils imaged in granulomas (Figure 5C-E). There were also

granuloma-associated *arg2:GFP* expressing cells that were *lyz:mCherry* negative (Figure

5C), some containing phagocytosed bacteria (Figure 5C and F) and expressing macrophage

mpeg:mCherry (Figure 5F). *mpeg:mCherry* positive macrophages had lower *arg2:GFP*

expression in comparison to the *mpeg:mCherry* negative/*arg2:GFP* positive granuloma-

associated cells (Figure 5F-H). *arg2:GFP* expression was observed in both infected and

non-infected *mpeg:mCherry* positive macrophages (Figure 5F-H). Due to the dim and

variable expression of both *arg2:GFP* and *mpeg:mCherry*, numbers of positive granuloma-

associated macrophages were difficult to accurately quantitate, but *arg2:GFP* positive

mpeg:mCherry macrophages were outnumbered by *arg2:GFP* negative *mpeg:mCherry*

macrophages and *arg2:GFP* positive *lyz:mCherry* neutrophils in granulomas. Together,

these data suggest that Mm granulomas have *arginase* expressing neutrophils and that

other granuloma-associated cells, including potential anti-inflammatory macrophages, also

express *arginase*.

Discussion

Zebrafish transgenic lines have allowed *in vivo* exploration of the pro-inflammatory

immune response (including *il-1β* and *tnfa*) to a variety of stimuli, however, until now, there

has been a lack of similar tools for anti-inflammatory factors [22,23,45]. Here, we describe a

new transgenic line for *arginase 2* and show that this transgene is upregulated shortly after

immune challenge in neutrophils and a small subset of macrophages. These data correlate

with a growing body of evidence that suggest anti-inflammatory factors are expressed early

after immune challenge to suppress hyper-inflammation before a switch to an anti-

inflammatory environment is required at later stages for tissue healing and restoration of

homeostasis [46].

The *arg2:GFP* line shows that neutrophils are the predominant immune cell type that

express *arginase 2* early after immune challenge in zebrafish. The observation of *arginase 2*

expression in zebrafish neutrophils fits with our previous observations that zebrafish

neutrophils are the primary innate immune cell that produce antimicrobial reactive nitrogen

species (RNS) after Mm infection and suggests a neutrophil iNOS/Arginase balance [47].

Human neutrophils also produce high levels of RNS [12], differing from expression in mice

where it is primarily macrophages that produce RNS/Arginase [12,16,48,49]. Human

polymorphonuclear neutrophils (PMNs) produce high levels of Arginase at a transcript and

protein level following immune challenge and in resting states, that may act as a negative

regulator of the RNS response, as is the case in murine macrophages. [12,16,48,49].

The mechanisms balancing innate immune regulation during inflammatory and

infection responses are not well understood *in vivo*. Neutrophil *arg2:GFP* was observed at

timepoints that are considered to be pro-inflammatory stages of inflammation and infection.

arg2:GFP expression was reminiscent of our previous observations using the pro-

inflammatory *TgBAC(il-1 β:GFP)sh445* transgenic line, where neutrophil *il-1β* was found at

1hpw in the tailfin and 1dpi in Mm infections, the same timepoints that *arg2:GFP* was

observed [23]. This suggests that anti-inflammatory *arginase* expression coincides with pro-

inflammatory signals in neutrophils, evidence for a balanced response. The observation of

immune cells producing anti-inflammatory signals upon immune challenge has been

observed previously, but to date this has been mainly described in macrophages [46]. It is

becoming clear that a balanced response to immune challenge, including both pro- and anti-

inflammatory signals is beneficial in disease control [46]. In murine macrophages it has been

demonstrated *in vitro* that Arginase 1 is produced soon after infection and can have

immunomodulatory effects [16]. Our findings open up the possibility that a similar balance

may exist in neutrophils and add to recent evidence suggesting that neutrophil phenotypes

are more diverse and nuanced than previously appreciated [6,7].

Some pathogens have evolved to disrupt the pro- and anti-inflammatory response,

keeping pro-inflammatory factors low and increasing anti-inflammatory signals to allow for

immune cell evasion and survival. One such pathogen is *Mycobacterium tuberculosis* (Mtb),

that suppresses an initial pro-inflammatory response, in part by upregulation of macrophage

Arginase, to allow for intra-phagocyte survival and intracellular growth and proliferation to

form hallmark granuloma structures [16]. This makes Arginase a potential target for

therapeutic intervention during TB. In murine TB models, macrophage Arginase expression

is associated with decreased bacterial killing and is an immunomodulatory target of

Mycobacterium tuberculosis [16]. Similar observations have been described in fungal

infections, with mice infected with Cryptococcus (*Cryptococcus gattii*) having elevated levels

of Arginase expression in lung tissues [50]. In human-monocyte-derived macrophages,

Candida albicans infection induces Arginase that blocks host NO production as a fungal

survival mechanism via chitin exposure [41]. Our findings using the *arg2:GFP* line are

consistent with these mammalian observations, with early *arginase* expression observed in

innate immune cells after Mm and fungal infection. Further investigation is required to

understand the molecular mechanisms of this intriguing host-pathogen interaction.

Arginase has been described as an anti-inflammatory macrophage marker due, in

part, to high expression levels observed in mouse monocytes cultured with anti-inflammatory

stimuli such as IL-4 or IL-13 to make M2/anti-inflammatory macrophages and observations

in other murine macrophage models [51]. We observed macrophages expressing *arg2:GFP*

during infection and at wound-healing stages of tailfin transection. Our findings complement

studies on a zebrafish pro-inflammatory macrophage *tnfa* transgenic line, which suggest that

an anti-inflammatory population of macrophages exist based on the switching off of *tnfa:GFP*

in some macrophages during the wound-healing phase of tailfin transection, and are

consistent with anti-inflammatory macrophages being present in the developing zebrafish

larvae [45]. However in both inflammation and infection, *arginase*-expressing macrophages

were much less frequent than *arginase*-expressing neutrophils, and neutrophil *arginase*

expression predominated. It is important to note that the *mpeg1* promoter used to mark

macrophages in our study is downregulated by Mm infection [52] therefore it is possible that

our observations using the *mpeg1:mCherry* line is an underestimation of the population of

macrophages that express *arg2:GFP* during Mm infection. This seems likely as there are

many *arg2:GFP* positive granuloma-associated cells, some with phagocytosed bacteria, that

expressed *arg2:GFP* but were neutrophil-marker negative, indicative of macrophages that

lack visible *mpeg1:mCherry* marker. The granuloma-associated *arg2:GFP* observed may

also be expressed in epithelioid-like cells that make up a large proportion of the zebrafish

Mm granuloma structure, some of which are macrophage derived but may have lost

macrophage markers [44]. Here, we have identified *mpeg1:mCherry* positive granuloma

associated macrophages that express *arginase 2*, however it remains unclear as to how

many macrophages are polarised towards this potential anti-inflammatory phenotype in

zebrafish. Characterising the macrophage polarisation infection response fully will require

further study for which the *arg2:GFP* zebrafish line will be an important tool.

Outside of immune cells, the *arg2:GFP* line has also illuminated arginase expression

in the liver in highly infected individuals and in ionocytes, during both resting and

inflammatory states. Arginase is a well-characterised liver enzyme [53,54] therefore

expression here was not unexpected, though the function of the upregulated liver expression

in highly infected individuals remains unclear. Interestingly, ionocytes have recently been

described as a new airway epithelial cell type in human and mice and it is these cells that

most highly express cystic fibrosis transmembrane conductance regulator (CFTR) the anion

channel that is mutated in cystic fibrosis patients [55]. The *arg:GFP* line is a new tool that

could be used to investigate the roles of these intriguing cells *in vivo*.

Our data indicate that *arginase 2*, an important anti-inflammatory mediator, is

produced early after immune challenge, predominantly by neutrophils. The *arg2:GFP* line is

an exciting addition to the zebrafish transgenic toolbox with which to investigate innate

immunity during infections. It has the potential to be applied to multiple zebrafish disease

models of infection and inflammation and may also be relevant to any models with an

inflammatory component, from ageing to cancer.

Figures and legends

Figure 1

Figure 1: The *TgBAC(arg2:eGFP)sh571* line shows GFP expression in ionocytes but not in resting macrophages and neutrophils, recapitulating the *arginase 2 in situ hybridisation* expression pattern

(A-B) Lightsheet microscopy stereo micrographs of *TgBAC(arg2:eGFP)sh571* line shows ionocytes expression at 2 (A) and 3dpf (B).

(C) Enlarged image of section over yolk of (B).

(D) *arginase 2 in situ* hybridisation shows expression in cells over the yolk known as

ionocytes at 2 and 3 dpf in unchallenged zebrafish.

(E) Enlarged image of section over yolk of 3dpf larvae in (D).

(F) Stereo fluorescence micrographs of *arg2:GFP* crossed to *Tg(mpeg1:mCherry)sh378* at

2dpf and 5dpf with no overlap of *arg2:GFP* expression in ionocytes with magenta

macrophages.

(G) Stereo fluorescence micrograph of *arg2:GFP* crossed to *Tg(mpeg1:mCherry)sh378* at

2dpf, 3dpf, 4dpf and 5dpf with no overlap of *arg2:GFP* expression in ionocytes with magenta

macrophages.

(H) Stereo fluorescence micrograph of *arg2:GFP* crossed to *Tg(lyz:nfsB.mCherry)sh260* at

3dpf with no overlap of *arg2:GFP* expression in ionocytes with magenta neutrophils.

Figure 2

Figure 2: Neutrophils express *arg2:GFP* after wound challenge

(A) Fluorescence confocal timelapse micrographs of *TgBAC(arg2:eGFP)sh571* positive cells migrating towards a tailfin nick wound at early timepoints post injury. Dashed line indicates the edge of the fin and the asterisk marks the nick wound.

(B) Fluorescence confocal micrographs of *arg2:GFP* crossed to *Tg(lyz:nfsB.mCherry)sh260* showing overlap of *arg2:GFP* with neutrophils at a tailfin wound (dashed line). Arrowheads indicate *arg2:GFP* positive neutrophils migrating at the wound.

(C) Fluorescence confocal timelapse micrographs of *arg2:GFP* crossed to *Tg(mpeg1:mCherry)sh378* showing no overlap of macrophages with *arg2:GFP* expression early after injury (dashed line indicates the wound).

(D) Fluorescence widefield micrographs of *arg2:GFP* crossed to *Tg(mpeg1:mCherry)sh378*. The upper panels show an uninjured tailfin with few macrophages and no immune cell *arg2:GFP* expression at 5dpf. The middle panels show an injured tailfin at 2dpw (4dpf) showing no overlap between *mpeg:mCherry* and *arg2:GFP*, but some unlabelled cells expressing *arg2:GFP* with amoeboid, immune cell shape (white arrowheads). Dotted line indicates the edge of the tailfin fold. The lower panels show an injured tailfin at 3dpw (5dpf) showing no overlap between *mpeg:mCherry* and *arg2:GFP*, but some unlabelled cells expressing *arg2:GFP* with amoeboid, immune cell shape (white arrowheads). Dotted line indicates the edge of the tailfin fold.

(E) Fluorescence confocal micrographs of *arg2:GFP* crossed to *Tg(mpeg1:mCherry)sh378* at 2dpw (upper panels) and 3dpw (lower panels), showing examples of *mpeg:mCherry* positive *arg2:GFP* expressing cells in the proximity of the healing wound (arrowheads).

Figure 3

Figure 3: Neutrophils are the predominant immune cell that express *arg2:GFP* post Mm challenge

(A) Fluorescence confocal micrographs of *TgBAC(arg2:GFP)sh571* crossed to *Tg(lyz:nfsB.mCherry)sh260* after Mm infection at 1dpi showing GFP positive neutrophils (filled arrowheads) and GFP negative neutrophils (hollow arrowheads), around an area of high infection (asterisk).

(B) Fluorescence confocal micrographs of 1dpi *arg2:GFP* crossed to *lyz:mCherry* after PVP

control injection.

(C) Fluorescence confocal micrographs of *arg2:GFP* crossed to *lyz:mCherry* after Mm

infection at 1dpi showing that both infected (filled arrowhead) and non-infected (hollow

arrowhead) neutrophils can express *arg2:GFP*.

(D) Graph showing the number of *arg2:GFP* positive and negative neutrophils in a 40x

region of interest in the caudal vein region that contained Mm bacteria, post infection, in

individual larvae.

(E) Graph showing the percentage of *arg2:GFP* positive and negative neutrophils and

macrophages in a 40x region of interest around the infected caudal vein region. Data shown

are n = 98-123 cells accumulated over 3 independent experiments.

(F) Fluorescence confocal micrographs of *arg2:GFP* crossed to *Tg(mpeg1:mCherry)sh378*

after Mm infection at 1dpi showing a single GFP positive macrophage (filled arrowhead) in

this field of view. One of 6 instances observed.

(G) Graph showing the number of *arg2:GFP* positive and negative macrophages in a 40x

region of interest in the caudal vein region that contained Mm bacteria post infection in

individual larvae.

Figure 4

A *Candida albicans*

B *Cryptococcus neoformans*

C *PVP*

D *Cryptococcus neoformans*

E PVP

Figure 4: Fungal infections upregulate *arg2:GFP* in neutrophils and the liver

(A) Fluorescence confocal micrographs of *arg2:GFP* crossed to *lyz:mCherry* after *Candida*

albicans infection at 1dpi showing GFP positive neutrophils (filled arrowhead) and GFP

negative neutrophils (hollow arrowhead), with a zoomed in panel below showing a positive

non-infected neutrophil and a GFP negative infected neutrophil (GFP in this cell is

autofluorescence from the Candida, which in this instance has survived and formed a

hyphae).

(B) Fluorescence confocal micrographs of *arg2:GFP* crossed to *lyz:mCherry* after

Cryptococcus neoformans infection at 1dpi showing GFP positive neutrophils (filled

arrowheads). Asterisk indicates a Cryptococci that is auto fluorescent in both channels.

(C) Brightfield and fluorescence micrographs of *arg2:GFP* larvae at 2dpi and 3dpi after PVP

injection. The position of the *arg2:GFP* negative liver is shown by the arrowhead. The GFP

channel is turned up highly enough to show gut fluorescence (asterisk) in order to show that

the liver is *arg2:GFP* negative.

(D) Brightfield and fluorescence micrographs of *arg2:GFP* larvae at 2dpi after *Cryptococcus*

neoformans (Cn) infection showing *arg2:GFP* liver expression with heavy levels of infection.

(E) Brightfield stereo micrographs of *arg2* wholemount *in situ* hybridisation after PVP or

Cryptococcus neoformans (Cn) infection showing *arg2* liver expression (arrowhead) in an

infected individual, not present in the PVP injected larvae.

Figure 5: A subset of granuloma associated neutrophils and macrophages express *arg2:GFP*

(A) Fluorescence confocal micrographs of *TgBAC(arg2:GFP)sh571* crossed to *Tg(lyz:nfsB.mCherry)sh260* and *Tg(mpeg1:mCherry)sh378* after PVP control injection at 4dpi. Only ionocyte expression of *arg2:GFP* is present.

(B) Fluorescence confocal micrographs of *TgBAC(arg2:GFP)sh571* after Mm infection at 4dpi showing granuloma associated *arg2:GFP* positive cells.

(C) Fluorescence confocal micrographs of *arg2:GFP* crossed to *Tg(lyz:nfsB.mCherry)sh260* after Mm infection at 4dpi, showing neutrophils positive with *arg2:GFP* (filled arrowheads).

(D) Line analysis of a cross section through a granuloma showing fluorescence values of *arg2:GFP*, *lyz:GFP* and *Mm*.

(E) Graph showing the percentage of *arg2:GFP* positive and negative granuloma-associated neutrophils. Data shown are n = 95 cells accumulated over 3 independent experiments.

(F) Fluorescence confocal micrographs of *TgBAC(arg2:GFP)sh571* crossed to *Tg(mpeg1:mCherry)sh378* after Mm infection at 4dpi.

(G) Fluorescence confocal micrographs of *arg2:GFP* crossed to *mpeg:mCherry* after Mm infection at 4dpi showing a non-infected, *arg2:GFP* positive macrophage (filled arrowhead).

(H) Fluorescence confocal micrographs of *arg2:GFP* crossed to *mpeg:mCherry* after Mm infection at 4dpi showing an infected, *arg2:GFP* positive macrophage (filled arrowhead).

Acknowledgements

The authors would like to thank The BSU Aquarium Team for fish care and the IICD Technical Team for practical assistance (University of Sheffield). We would like to thank the Renshaw and Johnston groups for sharing their expertise in BAC transgenesis, especially Miss Catherine Loynes and Dr Stone Elworthy. Thanks to Dr Simon Johnston and Dr Stella Christou for invaluable help and expertise in fungal

infection models. Thanks to Prof Stephen Renshaw for his helpful comments and

discussions on the manuscript.

Funding

AL and PME are funded by a Sir Henry Dale Fellowship jointly funded by the

Wellcome Trust and the Royal Society (Grant Number 105570/Z/14/A) held by PME.

FRH is funded by a University of Sheffield PhD scholarship. Lightsheet imaging was

carried out in the Wolfson Light Microscopy Facility, supported by a BBSRC

ALERT14 award for light-sheet microscopy (BB/M012522/1).

Author Contributions

Conceived and designed the experiments: FRH, AHM, GFW, PME. Performed the

experiments: FRH, AL, HEA, LGW, PME. Generation of *TgBAC(arg2:eGFP)sh571*

transgenic zebrafish: AL. Analysed the data: FRH, AL, PME. Wrote and drafted the

manuscript: FRH, AL, AHM, GFW, and PME.

Conflicts of Interests

The authors declare that they have no conflicts of interest.

References

1 Hartenstein V (2006) Blood cells and blood cell development in the animal kingdom. *Annu*

Rev Cell Dev Biol **22**, 677–712.

2 Duffin R, Leitch AE, Fox S, Haslett C & Rossi AG (2010) Targeting granulocyte apoptosis:

mechanisms, models, and therapies. *Immunol Rev* **236**, 28–40.

3 Mathias JR, Perrin BJ, Liu T-X, Kanki J, Look AT & Huttenlocher A (2006) Resolution of

inflammation by retrograde chemotaxis of neutrophils in transgenic zebrafish. *J*

Leukoc Biol **80**, 1281–1288.

4 Elks PM, van Eeden FJ, Dixon G, Wang X, Reyes-Aldasoro CC, Ingham PW, Whyte MKB,

Walmsley SR & Renshaw SA (2011) Activation of hypoxia-inducible factor-1α (Hif-

1α) delays inflammation resolution by reducing neutrophil apoptosis and reverse

migration in a zebrafish inflammation model. *Blood* **118**, 712–722.

5 Phillipson M & Kubes P (2019) The Healing Power of Neutrophils. *Trends Immunol* **40**,

635–647.

6 Ballesteros I, Rubio-Ponce A, Genua M, Lusito E, Kwok I, Fernández-Calvo G, Khoyratty

TE, van Grinsven E, González-Hernández S, Nicolás-Ávila JÁ, Vicanolo T,

Maccataio A, Benguría A, Li JL, Adrover JM, Aroca-Crevillen A, Quintana JA, Martín-

Salamanca S, Mayo F, Ascher S, Barbiera G, Soehnlein O, Gunzer M, Ginhoux F,

Sánchez-Cabo F, Nistal-Villán E, Schulz C, Dopazo A, Reinhardt C, Udalova IA, Ng

LG, Ostuni R & Hidalgo A (2020) Co-option of Neutrophil Fates by Tissue

Environments. *Cell.*

7 Giese MA, Hind LE & Huttenlocher A (2019) Neutrophil plasticity in the tumor

microenvironment. *Blood* **133**, 2159–2167.

8 Kato T & Kitagawa S (2006) Regulation of neutrophil functions by proinflammatory

cytokines. *Int J Hematol* **84**, 205–209.

9 Shapouri-Moghaddam A, Mohammadian S, Vazini H, Taghadosi M, Esmaeili S-A, Mardani

F, Seifi B, Mohammadi A, Afshari JT & Sahebkar A (2018) Macrophage plasticity,

polarization, and function in health and disease. *J Cell Physiol* **233**, 6425–6440.

10 Stuehr DJ & Marletta MA (1985) Mammalian nitrate biosynthesis: mouse macrophages

produce nitrite and nitrate in response to Escherichia coli lipopolysaccharide. *Proc*

Natl Acad Sci USA **82**, 7738–7742.

11 Wiegertjes GF, Wentzel AS, Spaink HP, Elks PM & Fink IR (2016) Polarization of

immune responses in fish: The "macrophages first" point of view. *Mol Immunol* **69**,

146–156.

12 Munder M, Mollinedo F, Calafat J, Canchado J, Gil-Lamaignere C, Fuentes JM, Luckner

C, Doschko G, Soler G, Eichmann K, Müller F-M, Ho AD, Goerner M & Modolell M

(2005) Arginase I is constitutively expressed in human granulocytes and participates

in fungicidal activity. *Blood* **105**, 2549–2556.

13 Dowling JK, Afzal R, Gearing LJ, Cervantes-Silva MP, Annett S, Davis GM, De Santi C,

Assmann N, Dettmer K, Gough DJ, Bantug GR, Hamid FI, Nally FK, Duffy CP,

Gorman AL, Liddicoat AM, Lavelle EC, Hess C, Oefner PJ, Finlay DK, Davey GP,

Robson T, Curtis AM, Hertzog PJ, Williams BRG & McCoy CE (2021) Mitochondrial

arginase-2 is essential for IL-10 metabolic reprogramming of inflammatory

macrophages. *Nat Commun* **12**, 1460.

14 Munder M, Eichmann K & Modolell M (1998) Alternative metabolic states in murine

macrophages reflected by the nitric oxide synthase/arginase balance: competitive

regulation by CD4+ T cells correlates with Th1/Th2 phenotype. *J Immunol* **160**,

5347–5354.

15 Munder M (2009) Arginase: an emerging key player in the mammalian immune system.

Br J Pharmacol **158**, 638–651.

16 El Kasmi KC, Qualls JE, Pesce JT, Smith AM, Thompson RW, Henao-Tamayo M,

Basaraba RJ, König T, Schleicher U, Koo M-S, Kaplan G, Fitzgerald KA, Tuomanen

EI, Orme IM, Kanneganti T-D, Bogdan C, Wynn TA & Murray PJ (2008) Toll-like

receptor-induced arginase 1 in macrophages thwarts effective immunity against

intracellular pathogens. *Nat Immunol* **9**, 1399–1406.

17 MacMicking J, Xie QW & Nathan C (1997) Nitric oxide and macrophage function. *Annu*

Rev Immunol **15**, 323–350.

18 Yoshida N, Frickel E-M & Mostowy S (2017) Macrophage–Microbe Interactions: Lessons

from the Zebrafish Model. *Front Immunol* **8**.

19 Henry KM, Loynes CA, Whyte MKB & Renshaw SA (2013) Zebrafish as a model for the
study of neutrophil biology. *J Leukoc Biol* **94**, 633–642.

20 Ellett F, Pase L, Hayman JW, Andrianopoulos A & Lieschke GJ (2011) mpeg1 promoter
transgenes direct macrophage-lineage expression in zebrafish. *Blood* **117**, e49-56.

21 Renshaw SA, Loynes CA, Trushell DMI, Elworthy S, Ingham PW & Whyte MKB (2006) A
transgenic zebrafish model of neutrophilic inflammation. *Blood* **108**, 3976–3978.

22 Marjoram L, Alvers A, Deerhake ME, Bagwell J, Mankiewicz J, Cocchiaro JL, Beerman
RW, Willer J, Sumigray KD, Katsanis N, Tobin DM, Rawls JF, Goll MG & Bagnat M
(2015) Epigenetic control of intestinal barrier function and inflammation in zebrafish.
Proc Natl Acad Sci USA **112**, 2770–2775.

23 Ogryzko NV, Lewis A, Wilson HL, Meijer AH, Renshaw SA & Elks PM (2019) Hif-1α-
Induced Expression of Il-1β Protects against Mycobacterial Infection in Zebrafish. *J*
Immunol **202**, 494–502.

24 Bojarczuk A, Miller KA, Hotham R, Lewis A, Ogryzko NV, Kamuyango AA, Frost H,
Gibson RH, Stillman E, May RC, Renshaw SA & Johnston SA (2016) Cryptococcus
neoformans Intracellular Proliferation and Capsule Size Determines Early
Macrophage Control of Infection. *Sci Rep* **6**, 21489.

25 Buchan KD, Prajsnar TK, Ogryzko NV, de Jong NWM, van Gent M, Kolata J, Foster SJ,
van Strijp JAG & Renshaw SA (2019) A transgenic zebrafish line for in vivo
visualisation of neutrophil myeloperoxidase. *PLoS ONE* **14**, e0215592.

26 Elks PM, Loynes CA & Renshaw SA (2011) Measuring inflammatory cell migration in the
zebrafish. *Methods Mol Biol* **769**, 261–275.

27 van der Sar AM, Spaink HP, Zakrzewska A, Bitter W & Meijer AH (2009) Specificity of the
zebrafish host transcriptome response to acute and chronic mycobacterial infection
and the role of innate and adaptive immune components. *Mol Immunol* **46**, 2317–
2332.

28 Cui C, Benard EL, Kanwal Z, Stockhammer OW, van der Vaart M, Zakrzewska A, Spaink

HP & Meijer AH (2011) Infectious disease modeling and innate immune function in

zebrafish embryos. *Methods Cell Biol* **105**, 273–308.

29 Benard EL, van der Sar AM, Ellett F, Lieschke GJ, Spaink HP & Meijer AH (2012)

Infection of zebrafish embryos with intracellular bacterial pathogens. *J Vis Exp*.

30 Seman BG, Moore JL, Scherer AK, Blair BA, Manandhar S, Jones JM & Wheeler RT

(2018) Yeast and Filaments Have Specialized, Independent Activities in a Zebrafish

Model of Candida albicans Infection. *Infection and Immunity* **86**.

31 Gibson RH, Evans RJ, Hotham R, Bojarczuk A, Lewis A, Bielska E, May RC, Elks PM,

Renshaw SA & Johnston SA (2018) Mycophenolate mofetil increases susceptibility

to opportunistic fungal infection independent of lymphocytes. *bioRxiv*, 131540.

32 Thisse C & Thisse B (2008) High-resolution in situ hybridization to whole-mount zebrafish

embryos. *Nat Protoc* **3**, 59–69.

33 Rath M, Müller I, Kropf P, Closs EI & Munder M (2014) Metabolism via Arginase or Nitric

Oxide Synthase: Two Competing Arginine Pathways in Macrophages. *Front Immunol*

5, 532.

34 Wentzel AS, Petit J, van Veen WG, Fink IR, Scheer MH, Piazzon MC, Forlenza M,

Spaink HP & Wiegertjes GF (2020) Transcriptome sequencing supports a

conservation of macrophage polarization in fish. *Sci Rep* **10**, 13470.

35 Wentzel AS, Janssen JJE, de Boer VCJ, van Veen WG, Forlenza M & Wiegertjes GF

(2020) Fish Macrophages Show Distinct Metabolic Signatures Upon Polarization.

Front Immunol **11**.

36 Rougeot J, Torraca V, Zakrzewska A, Kanwal Z, Jansen HJ, Sommer F, Spaink HP &

Meijer AH (2019) RNAseq Profiling of Leukocyte Populations in Zebrafish Larvae

Reveals a cxcl11 Chemokine Gene as a Marker of Macrophage Polarization During

Mycobacterial Infection. *Front Immunol* **10**, 832.

37 Athanasiadis EI, Botthof JG, Andres H, Ferreira L, Lio P & Cvejic A (2017) Single-cell
RNA-sequencing uncovers transcriptional states and fate decisions in
haematopoiesis. *Nature Communications* **8**, 2045.

38 Hsiao C-D, You M-S, Guh Y-J, Ma M, Jiang Y-J & Hwang P-P (2007) A positive
regulatory loop between foxi3a and foxi3b is essential for specification and
differentiation of zebrafish epidermal ionocytes. *PLoS ONE* **2**, e302.

39 Eisenhoffer GT, Slattum G, Ruiz OE, Otsuna H, Bryan CD, Lopez J, Wagner DS,
Bonkowsky JL, Chien C-B, Dorsky RI & Rosenblatt J (2017) A toolbox to study
epidermal cell types in zebrafish. *J Cell Sci* **130**, 269–277.

40 Jänicke M, Carney TJ & Hammerschmidt M (2007) Foxi3 transcription factors and Notch
signaling control the formation of skin ionocytes from epidermal precursors of the
zebrafish embryo. *Dev Biol* **307**, 258–271.

41 Wagener J, MacCallum DM, Brown GD & Gow NAR (2017) Candida albicans Chitin
Increases Arginase-1 Activity in Human Macrophages, with an Impact on
Macrophage Antimicrobial Functions. *mBio* **8**.

42 Brothers KM & Wheeler RT (2012) Non-invasive imaging of disseminated candidiasis in
zebrafish larvae. *J Vis Exp.*

43 Wright PA, Campbell A, Morgan RL, Rosenberger AG & Murray BW (2004) Dogmas and
controversies in the handling of nitrogenous wastes: Expression of arginase Type I
and II genes in rainbow trout: influence of fasting on liver enzyme activity and mRNA
levels in juveniles. *Journal of Experimental Biology* **207**, 2033–2042.

44 Cronan MR, Hughes EJ, Brewer WJ, Viswanathan G, Hunt EG, Singh B, Mehra S,
Oehlers SH, Gregory SG, Kaushal D & Tobin DM (2021) A non-canonical type 2
immune response coordinates tuberculous granuloma formation and epithelialization.
Cell **184**, 1757-1774.e14.

45 Nguyen-Chi M, Laplace-Builhe B, Travnickova J, Luz-Crawford P, Tejedor G, Phan QT,
Duroux-Richard I, Levraud J-P, Kissa K, Lutfalla G, Jorgensen C & Djouad F (2015)
Identification of polarized macrophage subsets in zebrafish. *Elife* **4**, e07288.

46 Cicchese JM, Evans S, Hult C, Joslyn LR, Wessler T, Millar JA, Marino S, Cilfone NA,

Mattila JT, Linderman JJ & Kirschner DE (2018) Dynamic balance of pro- and anti-

inflammatory signals controls disease and limits pathology. *Immunol Rev* **285**, 147–

167.

47 Elks PM, Brizee S, van der Vaart M, Walmsley SR, van Eeden FJ, Renshaw SA & Meijer

AH (2013) Hypoxia inducible factor signaling modulates susceptibility to

mycobacterial infection via a nitric oxide dependent mechanism. *PLoS Pathog* **9**,

e1003789.

48 Abebe T, Takele Y, Weldegebreal T, Cloke T, Closs E, Corset C, Hailu A, Hailu W, Sisay

Y, Corware K, Corset M, Modolell M, Munder M, Tacchini-Cottier F, Müller I & Kropf

P (2013) Arginase activity - a marker of disease status in patients with visceral

leishmaniasis in ethiopia. *PLoS Negl Trop Dis* **7**, e2134.

49 Jacobsen LC, Theilgaard-Mönch K, Christensen EI & Borregaard N (2007) Arginase 1 is

expressed in myelocytes/metamyelocytes and localized in gelatinase granules of

human neutrophils. *Blood* **109**, 3084–3087.

50 Oliveira-Brito PKM, Rezende CP, Almeida F, Roque-Barreira MC & da Silva TA (2020)

iNOS/Arginase-1 expression in the pulmonary tissue over time during Cryptococcus

gattii infection. *Innate Immun* **26**, 117–129.

51 Mantovani A, Sica A, Sozzani S, Allavena P, Vecchi A & Locati M (2004) The chemokine

system in diverse forms of macrophage activation and polarization. *Trends Immunol*

25, 677–686.

52 Benard EL, Racz PI, Rougeot J, Nezhinsky AE, Verbeek FJ, Spaink HP & Meijer AH

(2015) Macrophage-expressed perforins mpeg1 and mpeg1.2 have an anti-bacterial

function in zebrafish. *J Innate Immun* **7**, 136–152.

53 Berüter J, Colombo JP & Bachmann C (1978) Purification and properties of arginase

from human liver and erythrocytes. *Biochem J* **175**, 449–454.

54 Haraguchi Y, Takiguchi M, Amaya Y, Kawamoto S, Matsuda I & Mori M (1987) Molecular

cloning and nucleotide sequence of cDNA for human liver arginase. *Proc Natl Acad*

Sci U S A **84**, 412–415.

55 Shah VS, Chivukula RR, Lin B, Waghray A & Rajagopal J (2022) Cystic Fibrosis and the

Cells of the Airway Epithelium: What Are Ionocytes and What Do They Do? *Annu*

Rev Pathol **17**, 23–46.

765