Behavioral signatures of Y-like neuronal responses

in human vision

Abbreviated title: Behavioral signatures of Y-like cells in humans

Ana L. Ramirez[1], Lowell W. Thompson[2], Ari Rosenberg[2], and Curtis L. Baker Jr.[1]

[1]Department of Ophthalmology
McGill University
Montreal, QC, H3G 1A4, Canada.

[2]Department of Neuroscience
School of Medicine and Public Health
University of Wisconsin – Madison
Madison, WI, 53705, USA.

Correspondence:
Curtis Baker
curtis.baker@mcgill.ca
McGill Vision Research
1650 Cedar Ave, Room L11.112
Montreal, H3G 1A4
Canada

Number of Pages: 23
Number of Figures: 7
Number of Tables: N/A
Number of Movies: N/A
Number of Words in the Abstract: 236
Number of Words in the Introduction: 643
Number of Words in the Discussion: 1459

Conflict of Interest: Based in part on these findings, the technology transfer office for UW-Madison (WARF) has filed a patent application (pending) with Ari Rosenberg, Curtis Baker, and Ana Ramirez Hernandez listed as inventors.

Acknowledgments
We thank Chang'an Zhan and Liang Zhen, who each provided invaluable help with technical issues. This work was supported by a Canadian NSERC grant OPG0001978 to CLB and a National Institutes of Health Grant (EY029438) to AR.

Abstract

Retinal ganglion cells (RGCs) initiating the magnocellular/Y-cell visual pathways are distinguished by responding nonlinearly to high spatial frequencies (SFs) and temporal frequencies (TFs). This nonlinearity is implicated in the processing of contrast modulation (CM) stimuli in cats and monkeys, but its contribution to human visual perception is not well understood. Here we formulated an approach to reveal the function of human Y-like RGCs by leveraging the Y-like carrier response properties of cortical neurons to CM stimuli consisting of a high SF contrast-reversing grating carrier whose contrast is modulated by a low SF drifting sinewave envelope. Subjects reported the direction of motion of CM envelopes or luminance modulation (LM) gratings, presented at different eccentricities. The dependence of task performance on SF (for LMs) or carrier SF (for CMs) was measured for different TFs (LMs) or carrier TFs (CMs). The best performance for LM stimuli was at lower TFs and SFs, which decreased systematically with eccentricity. However, performance with CM stimuli was bandpass with carrier SF, largely independent of carrier TF, and at the highest carrier TF (20 Hz) decreased minimally with eccentricity. Since the nonlinear subunits of Y-cells respond better at higher TFs compared to the linear response components, and respond best at higher SFs that are relatively independent of eccentricity, these results suggest that behavioral tasks employing CM stimuli with high spatiotemporal carrier frequencies can reveal the nonlinear contributions of retinal Y-like cells to human perception.

Significance Statement

Y-like retinal ganglion cells (RGCs), characterized by responding linearly at low spatial and temporal frequencies and nonlinearly at high spatiotemporal frequencies, have been well characterized neurophysiologically in cats and monkeys. However, a clear counterpart of Y-like RGCs has remained elusive in humans. Here, we leveraged these response properties to reveal a behavioral signature of Y-like cells in human observers. To drive linear or nonlinear response mechanisms, we presented drifting luminance modulation and contrast modulation stimuli respectively at multiple spatial and temporal frequencies and eccentricities. Our results suggest that humans exhibit behavioral response patterns to contrast modulation stimuli that are consistent with the nonlinear response properties of Y-like cells.

Introduction

Information about the visual world is provided to the brain through the responses of retinal ganglion cells (RGCs). There are many categorically distinct types of RGCs (Masland, 2001), and establishing their functional contributions to perception has been challenging. Early studies in the cat characterized two principal types of RGCs, X-beta cells and Y-alpha cells (Enroth-Cugell & Robson, 1966; Lennie, 1980; Hochstein & Shapley, 1976), which comprise key elements along the retino-geniculate pathway. Later studies in the primate similarly characterized two main types of RGCs, midget and parasol cells, which were the origin of the parvocellular (P) and magnocellular (M) geniculocortical pathways. Similar to primate midget cells, X-beta cells are compact neurons with small axons and dendritic fields. Primate parasol cells are like Y-alpha cells in having large receptive and dendritic fields. However, it was not immediately clear whether parasol cells possessed the characteristic frequency-doubling nonlinearity of Y-alpha cells, and it was therefore ambiguous whether primates possess a RGC counterpart of cat Y-alpha cells. Later research demonstrated that parasol and smooth/upsilon cells, which we collectively refer to here as Y-like cells, exhibit both anatomical and functional properties of cat Y-alpha cells including nonlinear responses to visual stimuli (Petrusca et al., 2007; Crook et al., 2008a, b). The functional role of Y-like cells' nonlinear responses is not well understood in human vision.

Classic experiments showed that selective lesions of the magnocellular pathway decrease contrast sensitivity for visual stimuli at high temporal frequencies (TFs) and low spatial frequencies (SFs) but do not affect color vision (Merigan & Manusell, 1990; Schiller et al., 1990; Merigan et al., 1991 a). However, those experiments did not investigate the contributions of the nonlinear responses of Y-like cells. Y-like cells are distinctive in that, at low SFs they show a linear first Fourier harmonic (F1) response at the temporal frequency of a drifting grating, but when contrast-reversing gratings at high SFs are presented, they display a prominent second Fourier harmonic (F2) nonlinear response. This pattern of SF-dependence of F1 and F2 responses, often referred to as the "Y-cell signature", is characteristic of the nonlinear summation of Y-cells (Hochstein & Shapley, 1976). The F1 response is dependent on the spatial phase of a contrast-reversing grating, whereas the F2 nonlinear response is phase-independent. This characteristic phase-invariance of the F2 response of Y-like cells reflects the activation of nonlinear subunits originating from a specialized class of bipolar cells (Demb et al., 1999; Crook

et al., 2008a, b). In neurophysiology experiments in primate retina, the estimated F1 linear center receptive fields clearly decrease systematically with eccentricity, as might be expected. In contrast, the F2 nonlinear center receptive fields maintain a relatively similar size across a large range of retinal eccentricities, although the differences have not been compared quantitatively (Crook et al., 2008b).

Contrast modulation (CM) stimuli generally consist of a carrier sine wave grating at a high SF, whose contrast is modulated by a sinewave envelope at a low SF. Neurophysiology studies have demonstrated that cortical neurons responding to CM stimuli exhibit selectivity for the carrier that is similar to the selectivity of nonlinear subunits of subcortical Y-cells (Gharat & Baker, 2017; Rosenberg et al., 2010; Rosenberg & Issa, 2011). These results support the idea that cortical CM responses are critically dependent on Y-cell inputs (Demb et al., 2001b; Rosenberg & Talebi, 2009; Rosenberg et al., 2010).

Here we leveraged the Y-like carrier response properties of cortical neurons to CM stimuli to create a novel approach capable of revealing the nonlinear contributions of Y-like cells to human perception. We employed CM stimuli with a carrier at high spatial and temporal frequencies and an envelope at low spatial and temporal frequencies. Judgments of the direction of motion for conventional luminance modulation (LM) gratings were consistent with conventionally described linear mechanisms, whereas results from envelope (CM) motion demonstrated a pattern of results consistent with Y-like nonlinear processing.

Materials and Methods

Subjects

Six healthy subjects (aged 23 to 35 years; 3 males, 3 females) participated in the study. Five were naive to the aims of the study (students from McGill University) and one was an author (ARH). All subjects had a monocular visual acuity or best-corrected acuity of at least 20/25 and reported no history of ophthalmological diseases or surgeries. All procedures were approved by the Research Ethics Board of the Research Institute of McGill University Health Center. All subjects gave written informed consent to participate.

Apparatus

Visual stimuli were produced on a Macintosh computer (MacPro 4.1, MacOS X 10.6.8, 2x2.8 GHz Quad-Core, 24 GB RAM) with custom software written in MATLAB (MathWorks, Inc.) using Psychophysics Toolbox, version 3.0.10 (Brainard, 1997; Kleiner et al., 2007; Pelli, 1997). The stimuli were presented on a cathode-ray tube (CRT) monitor (Iiyama, 39.5 x 29.5 cm, 120 Hz, 1024 x 768 pixels) at a viewing distance of 221 cm. To achieve linearization and high monochromatic luminance level resolution, we first measured the CRT gamma nonlinearity with a photometer (United Detector Technology S370), and then used the LOBES video switcher (Li et al., 2003) that combines blue and attenuated red outputs from the graphics card to generate 16 bits of voltage resolution.

Stimuli

The visual stimuli were presented at the center of the monitor's screen within a cosine-tapered circular window, on a uniform background at the mean luminance of the stimulus. We presented both first-order, luminance modulation (LM) gratings as well as second-order, contrast modulation (CM) gratings. A drifting LM grating was defined by:

$$I(x, y, t) = L_B \left\{ 1.0 + C_{LM} \cos(\omega_{\mathrm{SF}} \cdot [\cos(\theta) \cdot x + \sin(\theta) \cdot y] - \omega_{\mathrm{TF}} \cdot t) \right\} \qquad (1)$$

where $I(x, y, t)$ is the luminance intensity of a pixel at spatial location (x, y) at time t, L_B is the background (and mean) luminance, C_{LM} is the Michelson contrast of the luminance modulation, ω_{SF} is the spatial frequency, θ is the orientation, and ω_{TF} is the temporal frequency. The sign of ω_{TF} determines the direction of motion. In these experiments, the orientation was always vertical, so the motion was either leftwards or rightwards.

A CM grating was defined by the contrast modulation of a high spatial frequency carrier grating by a low spatial frequency envelope grating (**Fig. 1A**):

$$I(x, y, t) = L_B \{ 1.0 + Carr(x, y, t) \cdot [1 + Env(x, y, t)/2] \} \qquad (2)$$

Here, $Carr(x, y, t)$ is the carrier, which is a contrast-reversing grating with an orientation, spatial frequency, and temporal frequency that are defined independently of the envelope:

$$Carr(x, y, t) = C_c \cos(\omega_{\text{SFc}} \cdot [\cos(\theta_c) \cdot x + \sin(\theta_c) \cdot y]) \sin(\omega_{\text{TFc}} \cdot t) \quad (3)$$

where ω_{SFc} is the carrier SF, ω_{TFc} is the carrier TF, θ_c is the carrier orientation and C_c is the carrier contrast. In these experiments, the carrier orientation was always right-oblique, as in **Figure 1A**.

Likewise, $Env(x, y, t)$ is the envelope, which is a drifting grating with an orientation, spatial frequency, and temporal frequency (direction) that are defined independently of the carrier:

$$Env(x, y, t) = C_e \cos(\omega_{\text{SFe}} \cdot [\cos(\theta_e) \cdot x + \sin(\theta_e) \cdot y] - \omega_{\text{TFe}} \cdot t) \quad (4)$$

where ω_{SFe} is the envelope SF, ω_{TFe} is the envelope TF, θ_e is the envelope orientation and C_e is the envelope contrast, which henceforth we will refer to as the "modulation depth". In these experiments, the envelope orientation was always vertical, so the motion was either leftwards or rightwards, as in **Figure 1A**.

A grey cardboard surround, 81 x 102 cm (width x height), framed the monitor and approximately matched the mean luminance (L_B) of the stimuli on the CRT.

Design and experimental procedures

Subjects were instructed to look at a fixation target which was placed in the upper right quadrant of the screen (**Fig. 1B**), such that the stimulus would be presented at 4.3° of eccentricity in the subject's visual field, in the first experiment, and at varying eccentricities in the second experiment (see below). The fixation target and all visual stimuli were presented monocularly to the right eye for all subjects. On each trial, a stimulus was presented for 250 ms. The subject indicated the perceived direction of motion of the LM grating or CM envelope (**Fig. 1C**) by pressing a key, with no time limit, and with subsequent feedback (visual icon on the display screen) for incorrect responses. Performance was measured by the method of constant stimuli

with seven logarithmically spaced values of spatial frequency (0.7, 1, 1.5, 2.1, 2.9, 4.2, and 5.9 cycles per degree (cpd)) and four values of temporal frequency (5, 10, 15, and 20 Hz). All experiments had a fixed contrast (C_{LM} and C_c from Eq 1 and 3, respectively) of 5% for luminance modulation gratings and 80% for contrast modulation envelope and carrier components, chosen based on pilot results to avoid ceiling effects in psychophysical performance and to compensate for the very different contrast sensitivities for the two kinds of stimuli (Smith et al, 1994; Schofield & Georgeson, 1999). A minimum of three blocks of 140 trials (20 trials per condition) was tested for each condition to provide a total of at least 60 trials per condition. Subjects generally completed one block in 5 to 10 minutes. In a second experiment, stimuli were presented at a fixed temporal frequency (20 Hz), at 2.1°, 4.3°, 6.4°, or 8.5° of eccentricity from the fixation target (**Fig. 1B**).

Figure 1. Visual stimuli and behavioral task. **A)** Example contrast modulation (CM) stimulus, presented within a cosine-tapered circular window, consisting of a high spatial frequency contrast-reversing carrier grating (right-oblique) whose contrast is modulated by a low spatial frequency, vertical sinewave envelope. **B)** Example luminance modulation (LM) stimulus at the center of the screen, with different fixation targets (only one fixation target was visible on any given trial) that would place the stimulus at 2.1°, 4.3° (in a red square), 6.4°, and 8.5° of eccentricity. **C)** Depiction of the 2 alternative forced choice (2AFC) task in which the subject reported the direction of motion (left vs right) of a CM envelope (shown here) or LM grating. The stimulus images are shown at larger than true scale for illustration purposes.

Experimental Design and Statistical Analysis

Performance was determined by calculating each subject's percent correct responses for each unique experimental condition. Standard errors were calculated for each condition based on the variance of the proportion in a binomial distribution resulting from Bernoulli trials: var(p) = p*(1-p)/N, where p is the proportion correct and N is the number of trials. For mean responses from all subjects (N=6), standard error of the mean was calculated. We used 2-way ANOVAs to test whether direction discrimination performance was significantly impacted by the stimulus SF, TF (or eccentricity), and their interaction.

To determine the SF acuity, i.e., the highest SF supporting a criterion performance threshold, for each subject and experimental condition, plots of percent correct data were fit with a logistic function. To do so, only the last 5 values of the high-SF fall-off were included (1.5, 2.1, 2.9, 4.2, and 5.9 cpd) to provide a monotonically decreasing function. The logistic curve-fitting was performed using Psignifit 4 (Schütt et al., 2016). Threshold values of maximal SF (for LM) or carrier SF (for CM) were taken at the 75% correct level. Thresholds were included in an analysis if the 95% confidence interval did not include 0 (Experiment 1: 45/48 cases; Experiment 2: 41/48 cases). We used linear regression models to test whether the log SF acuity varied systematically as a function of TF (or eccentricity), stimulus type (LM or CM), and their interaction. We report the corresponding fixed-effect coefficients (b-values), F statistics, and p-values.

Results

Experiment 1: Spatial frequency dependence at different temporal frequencies

We hypothesized that direction discrimination performance for CM stimuli would remain relatively similar across a broad range of carrier TFs if the envelope motion was computed from the output of nonlinear Y-like cells rather than linear mechanisms. To test this prediction, we

assessed performance across a variety of carrier SFs and TFs at a fixed eccentricity. To provide a comparative context for contrasting the results with those reflecting linear processing, we made similar measurements for luminance gratings (LM stimuli) over the same ranges of SFs and TFs, and at the same eccentricity. In particular, direction discrimination performance for LM stimuli was predicted to decrease systematically as a function of TF, as expected if perception were mediated by linear mechanisms generally thought to underlie performance for moving gratings (Robson, 1966; Kulikowski, 1976; Watson & Ahumada, 2016).

In each stimulus presentation of 250 ms duration, an LM or CM stimulus was presented in the center of the screen while the subject fixated a target 4.3° away from the stimulus (**Fig. 1B**, red square). The task was to report the perceived direction of motion of the LM grating or of the CM envelope. Within each block of trials, the LM grating SF or CM carrier SF was varied using the method of constant stimuli at a fixed TF (LM gratings) or carrier TF (CM envelopes). In separate blocks of trials, the dependence of performance on SF was measured for different values of TF (LM gratings) or carrier TF (CM envelopes).

We first analyzed performance with the LM stimuli (vertical gratings, drifting leftwards or rightwards), which were presented with a Michelson contrast of 5%. The percent correct performance as a function of SF is plotted for one subject in **Figure 2A** for a series of values of TF. The ability to correctly report the direction of motion at relatively high SFs (1.5 to 3.0 cpd) was only possible at low TFs (5 to 10 Hz). Note that the performance fall-off with increasing SF was faster at successively higher TFs. **Figure 2B-F** plots the results for 5 additional subjects, who showed similar patterns. The average performance across all 6 subjects is plotted in **Figure 2G**. In general, performance fell off with increasing SF, and was more rapid for higher TF values (except at the two lowest TF values, at which performance was similar). These observations were confirmed with a 2-way ANOVA (**Fig. 2G**), which showed a significant main effect of SF, $F(6,168) = 117.02$, $p = 2.39 \times 10^{-57}$; a significant main effect of TF, $F(3,168) = 20.08$, $p = 3.59 \times 10^{-11}$; and a significant interaction effect, $F(18,168) = 3.3$, $p = 2.43 \times 10^{-5}$.

In general, the performance for LM stimuli at relatively high SFs was best at low TFs, and decreased with higher values of SF or TF. These findings replicate well-known results for LM stimuli as functions of SF and TF in central vision, where performance at high SFs is best at low TFs, and vice versa (Robson, 1966; Kulikowski, 1970; Watson & Ahumada, 2016).

Figure 2. Performance for luminance modulation (LM) stimuli (i.e., drifting sinusoidal gratings). The stimuli consisted of luminance gratings at different SFs and TFs. The task was to discriminate the direction of motion of a grating presented monocularly at 4.3° of eccentricity. Percent correct responses were measured while varying SF (cpd) within blocks for different values of TF: magenta, 5 Hz; green, 10 Hz; cyan, 15 Hz; and dark blue, 20 Hz. **A-F)** Percent correct responses for each subject. **G)** Mean percent correct responses of the six subjects. Error bars represent the binomial standard error (SE) of each condition for each subject (**A-F**) and the SEM (N=6 subjects) for the subject-averaged results (**G**).

Next, we examined performance with the CM stimuli which had a contrast-reversing grating carrier with right-oblique orientation (45°). The carrier contrast (C_c in Eq 3) was modulated by a vertical drifting sinewave grating envelope with a modulation depth (C_e in Eq 4) of 80%, SF of 0.25 cpd, and TF of 3 Hz. The carrier contrast was 80%, chosen based on pilot experiments to provide roughly similar ranges of performance as for the LM stimuli. Percent correct performance to report the direction of envelope motion is plotted as a function of carrier SF for one subject in **Figure 3A**, for a series of carrier TFs. Performance was generally best at mid-range carrier SFs (1.5 to 3.0 cpd) that were higher than the SFs giving best performance for LM stimuli (**Fig. 2**). Another difference from the LM results is that CM performance at higher SFs was more robust with increasing carrier TF. **Figure 3B-F** plots the results for the 5 other subjects, who showed largely similar results. The average performance across all 6 subjects (**Fig. 3G**) shows a bandpass dependence on carrier SF, which is very similar for the different carrier

TFs, with the curves almost overlapping. These observations were confirmed with a 2-way ANOVA, which showed a statistically significant effect of carrier SF, $F(6,168) = 70.27$, $p = 2.97 \times 10^{-43}$, but no significant effect of carrier TF, $F(3,168) = 2.02$, $p = 0.11$, or interaction of carrier SF and TF, $F(18,168) = 0.76$, $p = 0.75$.

Figure 3. Performance for contrast modulation (CM) stimuli. The stimuli consisted of a contrast-reversing grating (carrier) modulated by a drifting sinewave envelope with a SF of 0.25 cpd and a TF of 3 Hz. The task was to discriminate the direction of motion of the envelope of a CM stimulus presented monocularly at 4.3° of eccentricity. Percent correct responses were measured while varying carrier SF (cpd) within blocks for different values of carrier TF: magenta, 5 Hz; green, 10 Hz; light blue, 15 Hz; and dark blue, 20 Hz. **A-F**) Percent correct responses of the six subjects. **G**) Mean percent correct responses of the six subjects. Error bars represent the binomial SE of each condition for each subject (**A-F**) and the SEM (N=6 subjects) for the subject-averaged results (**G**).

As a whole, the performance for CM stimuli was better at higher carrier SFs and TFs than for LM stimuli. In particular, there was a large degree of independence between the effects of carrier SF and TF. These results would not be expected based on the linear mechanisms that detect LM stimuli, which support good performance when high SFs are combined with low TFs, or vice versa (Robson, 1966; Kulikowski, 1976; Watson & Ahumada, 2016). Importantly, the current results for CM stimuli (**Fig. 3**) are in sharp contrast to those for LM stimuli (**Fig. 2**) over

the same ranges of spatiotemporal frequencies, consistent with the idea that the processing of CM stimuli is fundamentally different from that of luminance stimuli.

In comparing the results for LM and CM stimuli, differences in the highest SFs giving good performance (i.e., thresholds describing SF acuity) provides a useful way to quantitatively compare LM and CM performance in a concise manner. To this end, we obtained measures of SF acuity (threshold) from the individual subject data in **Figures 2 & 3** by curve-fitting the high-SF fall-off and taking the value of SF corresponding to 75% correct as an estimate of SF acuity (see **Materials and Methods**). Thresholds were successfully obtained for 23/24 and 22/24 of the individual subject curves in **Figures 2 & 3** respectively. The results of this analysis show that SF acuity for LM stimuli (orange curve) declines systematically with TF whereas carrier SF acuity for CM stimuli (purple curve) remains relatively constant with carrier TF (**Fig. 4**). To quantify this assessment, we regressed the log SF acuity on the TF and stimulus type (LM or CM) as well as their interaction. A significant main effect of TF was observed, $F(1,41) = 10.30$, $p = 2.7 \times 10^{-3}$, as well as a significant interaction between TF and stimulus type, $F(1,41) = 6.85$, $p = 0.012$, such that SF acuity declined faster with increasing TF for LM stimuli (b = -0.03) than CM stimuli (b = -0.003).

Figure 4. Spatial frequency acuity (i.e., SF corresponding to 75% correct) obtained by fitting the high-SF fall-off of the individual subject data in **Figures 2** and **3** for LM (orange) and CM (purple) stimuli, respectively. Error bars indicate ± SEM (N=6 subjects). Data points for CM stimuli are shifted slightly relative to the actual TF values for visualization purposes.

Experiment 2: Spatial frequency dependence at different eccentricities

Based on previous retinal neurophysiology (Crook et al., 2008b), we further hypothesized that direction discrimination performance for CM stimuli would remain relatively constant as a function of stimulus eccentricity if the envelope motion was computed from the output of nonlinear Y-like cells rather than linear mechanisms. To test this prediction, we performed a second experiment in which direction discrimination performance for CM stimuli was measured as a function of carrier SF, with a fixed carrier TF, at different eccentricities. Similar to Experiment 1, we compared the results to those with LM stimuli over the same values of SFs, TFs, and retinal eccentricities. For LM stimuli, previous findings would predict a systematic fall-off of performance for higher SFs at increasing eccentricities (Koenderink et al., 1977; Rovamo et al., 1978).

In each stimulus presentation of 250 ms duration, an LM or CM stimulus was presented in the center of the screen while the subject fixated a target that was 2.1°, 4.3°, 6.4°, or 8.5° away from the stimulus (**Fig. 1B**). The task was to report the direction of motion of the LM grating or of the CM envelope. Within each block, the SF of the LM grating or the carrier SF of the CM stimulus was varied using the method of constant stimuli with a fixed TF (for LM stimuli) or carrier TF (for CM stimuli) of 20 Hz, respectively. This high value of TF or carrier TF was specifically chosen to minimize the contribution of linear (luminance-based) mechanisms, based on the results of the previous section. Different eccentricities were tested in separate trial blocks.

As in Experiment 1, the LM stimuli (vertical gratings, drifting leftwards or rightwards) were presented with a Michelson contrast of 5%. Percent correct performance to report the direction of motion of the grating as a function of SF is plotted for one subject in **Figure 5A** for each of a series of eccentricities. Note that the performance declined with SF, and the high-SF fall-off decreased with stimulus eccentricity. **Figure 5B-F** plots the results for the 5 other subjects, who showed largely similar results. The average performance across all 6 subjects is plotted in **Figure 5G**, where a systematic fall-off in performance at successively higher SFs and eccentricities is evident. These interpretations were confirmed with a 2-way ANOVA (**Fig. 5G**), showing a significant main effect for SF, $F(6,168) = 132.27$, $p = 4.58 \times 10^{-61}$, a significant main

effect for eccentricity $F(3,168) = 15.16$, $p = 8.88 \times 10^{-9}$, and a significant interaction effect $F(18,168) = 2.28$, $p = 0.003$.

Figure 5. Performance for luminance modulation (LM) stimuli at different retinal eccentricities and a fixed TF of 20 Hz. The stimuli consisted of a luminance grating at different SFs and a fixed TF. The task was to discriminate the direction of motion of the grating when presented at different retinal eccentricities. Percent correct responses were measured while varying SF within blocks for different eccentricities: dark purple, fixation at 2.1°; dark blue, 4.3°; pink, 6.4°; and light blue, 8.5°. **A-F)** Percent correct responses for each subject. **G)** Mean percent correct responses of the six subjects. Error bars represent the binomial SE of each condition for each subject (**A-F**) and the SEM (N=6 subjects) for subject-averaged results (**G**).

Across subjects the performance for LM stimuli decreased systematically with higher SF and greater eccentricity. These results are consistent with what is expected for LM stimuli, where the dependence of contrast sensitivity as a function of SF shifts to lower SFs as eccentricity increases (Koenderink et al., 1977; Rovamo et al., 1978).

Following Experiment 1, the CM stimuli had a contrast-reversing grating carrier (20 Hz) with right-oblique orientation (45°) and a carrier contrast of 80%. The carrier contrast was modulated by a vertical drifting sinewave grating envelope with a modulation depth of 80%, SF of 0.25 cpd, and TF of 3 Hz. Percent correct performance to report the direction of envelope motion as a function of carrier SF at a fixed carrier TF (20 Hz) is plotted for one subject in

Figure 6A for each of a series of eccentricities. The best performance was at mid-range carrier SFs (1.5 to 3.0 cpd) and was relatively independent of eccentricity compared to the results for LM stimuli (**Fig. 5**). **Figure 6B-F** plot the results for the 5 other subjects who generally showed a bandpass dependence on carrier SF. The average performance across all 6 subjects (**Fig. 6G**) shows this bandpass dependence on carrier SF, with the curves for the different eccentricities so similar as to be largely overlapping. These impressions were confirmed with a 2-way ANOVA (**Fig. 6G**), which showed a significant main effect for the carrier SF, $F(6,168) = 55.08$, $p = 3.43 \times 10^{-37}$, but no significant main effect for eccentricity $F(3,168) = 0.84$, $p = 0.48$, and no significant interaction effect $F(18,168) = 0.92$, $p = 0.56$.

Figure 6. Performance for contrast modulation (CM) stimuli at different retinal eccentricities and a fixed TF of 20 Hz. The stimuli consisted of a contrast-reversing grating (carrier) modulated by a drifting sinewave envelope with a SF of 0.25 cpd and a TF of 3 Hz. The task was to discriminate the direction of motion of the envelope when presented at different retinal eccentricities. Percent correct responses were measured while varying carrier SF within blocks for different eccentricities: dark purple, fixation at 2.1°; dark blue, 4.3°; pink, 6.4°; and light blue, 8.5°. **A-F)** Percent correct responses for each subject. **G)** Mean percent correct responses of the six subjects. Error bars represent the binomial SE of each condition for each subject (**A-F**) and the SEM (N=6 subjects) for the subject-averaged results (**G**).

Notably, the performance with CM stimuli exhibited relatively little dependence on eccentricity (**Fig. 6**), whereas the results with LM stimuli showed a clear, systematic eccentricity-dependence (**Fig. 5**). Only the latter is expected based on a linear mechanism, since

the SF-dependence of contrast sensitivity shifts to lower SFs with increasing eccentricity (Koenderink et al., 1977; Rovamo et al., 1978). Thus, these results are consistent with the idea that the processing of CM stimuli is fundamentally different (nonlinear) from that of LM stimuli.

Similar to Experiment 1, a useful way to compare LM and CM performance in a concise manner was to measure the differences in highest SFs giving good performance. Consequently, we obtained measures of SF acuity (threshold) from the individual subject data in **Figures 5 & 6** by curve-fitting the high-SF fall-off and taking the value of SF corresponding to 75% correct as an estimate of SF acuity (see **Materials and Methods**). Thresholds were successfully obtained for 18/24 and 23/24 of the individual subject curves in **Figures 5 & 6** respectively. The results of this analysis show that SF acuity for LM stimuli (orange) declines systematically and substantially with eccentricity, whereas the carrier SF acuity for CM stimuli (purple) remains relatively constant, or declines very little, with eccentricity (**Fig. 7**). To quantify this assessment, we regressed the log SF acuity on the eccentricity and stimulus type (LM or CM), as well as their interaction. Consistent with our observations, the effect of eccentricity on SF acuity was significant, $F(1,37) = 28.36$, $p = 5.14\text{x}10^{-6}$, and was twice as large for LM stimuli (b = -0.10) compared to CM stimuli (b = -0.05), although the interaction did not reach statistical significance, $F(1,37) = 2.65$, $p = 0.11$.

Figure 7. Spatial frequency acuity (i.e., SF corresponding to 75% correct) obtained by curve-fitting the high-SF fall-off of the individual subject data in **Figures 5** and **6** for LM (orange) and CM (purple) stimuli, respectively. Error bars indicate ± SEM (N=6 subjects). Data points for CM stimuli are shifted slightly relative to the actual eccentricity values for visualization purposes.

Discussion

The encoding of "second-order" stimuli, such as the CM patterns used here, are of particular interest in the context of neural mechanisms underlying perception, since figure-ground boundaries in natural images often entail modulations in texture properties such as contrast (Johnson & Baker, 2004). Here we introduced a way to assess the subcortical inputs to CM processing in human perception, which was previously possible only in animal models.

CM stimuli have been found to drive early visual cortex neurons in cats (Mareschal and Baker, 1998) and macaque monkeys (Li et al., 2014), with selectivity to the carrier spatial frequency. The responses to CM stimuli in both cat LGN (Rosenberg et al., 2010) and area 18 (Rosenberg & Issa 2011; Gharat & Baker, 2012) occur at surprisingly high carrier TFs, suggesting a subcortical origin, likely retinal Y-cells. Using CM stimuli at high carrier spatiotemporal frequencies, here we were able to translate this evidence from previous neurophysiology experiments into a human behavioral approach. The results of these experiments suggest that human motion perception of CM stimuli is mediated by carrier processing with nonlinear subunits of Y-like subcortical cells (Rosenberg et al., 2010; Rosenberg & Issa, 2011), rather than by conventional linear mechanisms thought to detect LM stimuli.

Y-like cells, such as retinal parasol cells, are characterized by their ability to respond linearly to low SF drifting gratings and nonlinearly to contrast-reversing gratings at high spatiotemporal frequencies (e.g., Hochstein & Shapley, 1976; Crook et al., 2008a). To drive linear mechanism responses, here we employed drifting LM stimuli at different SFs and TFs, which are known to elicit the best behavioral response at combinations of low TF and high SF, or vice versa (Robson, 1966; Kulikowski, 1970; Watson & Ahumada, 2016). Not surprisingly, the performance that we found for LM stimuli was consistent with this idea (**Figs. 2**, **4**). However, CM stimuli are thought to be specifically processed by nonlinear mechanisms that arise from retinal bipolar cell inputs to the Y-like retinal ganglion cells that project to the LGN (Demb et al., 2001a, b), whose responses are then relayed to the visual cortex. The nonlinear subunits of LGN Y-cells have been demonstrated to respond to relatively higher TFs compared to linear cortical mechanisms (Rosenberg et al., 2010; Rosenberg & Issa, 2011). Therefore, nonlinear (carrier) responses of cortical neurons to CM stimuli should be able to produce good behavioral performance at higher carrier spatiotemporal frequencies. This prediction is consistent with the results in our experiments with CM stimuli, where performance was best at relatively high

carrier SFs and high carrier TFs (**Figs. 3**, **4**). In addition, the demonstration of a bandpass dependence on carrier SF (**Fig. 3**) is consistent with similar findings for CM-responsive single neurons in early visual cortex of cats (Zhou & Baker, 1994, 1996; Tanaka & Ohzawa, 2006; Rosenberg et al., 2010) and macaque monkeys (Li et al., 2014).

Previous human behavioral studies using luminance gratings demonstrated that the SF-dependence on contrast sensitivity shifts to lower SFs with increasing eccentricity (Koenderink et al., 1977; Rovamo et al., 1978), much as we found in our experiments with LM stimuli (**Fig. 5**). However, for CM stimuli, performance as a function of carrier SF changed relatively little with eccentricity. This relative invariance to eccentricity for the carrier SF dependence of CM stimuli might reflect the organization of the nonlinear (F2, second harmonic) receptive fields of the Y-like cells. Indeed, previous primate neurophysiology demonstrated that estimated sizes of the center F2 receptive fields of Y-like cells varied minimally with eccentricity, while estimates of the size of the linear (F1, first harmonic) center receptive fields decrease substantially with eccentricity (Crook et al., 2008b).

Notably, in the experiments of Crook et al. (2008b), the eccentricities in the primate retina used to measure F1 and F2 center receptive fields (7°-40°) were mostly higher than the eccentricities (ca 2°-8°) examined here. While it can be challenging for inexperienced subjects to perform perceptual judgments in more peripheral vision, it nevertheless might be worthwhile in future studies to test a higher range of eccentricities, to make further comparisons to the results of Crook et al. (2008b). Based on our results here, and our conclusion that behavioral responses to CM stimuli are mediated by the nonlinear subunits of Y-like cells, we would expect performance to continue declining at larger retinal eccentricities for the LM stimuli at a greater rate than for the CM stimuli.

One can conceivably argue that for our results with CM stimuli, the bandpass carrier SF dependence, in particular the fall-off at low carrier SFs (**Figs. 3**, **6**), might be secondary to a lack of a constant ratio between the carrier and envelope SFs, because the envelope SF was kept at a fixed value. In this view, if the carrier and envelope SFs become more similar, it might interfere with the ability to judge the direction of envelope motion as distinct from the carrier flicker, or even that there might be insufficient cycles of the carrier within each envelope cycle to provide a genuine CM stimulus. However, if we had kept a constant ratio, at the lowest eccentricity the visual stimulus would be so large that part of the stimulus would be overlapping the fovea and in

peripheral vision it would be too small to make a reasonable judgement of envelope motion. Although in our experiments the ratio was not constant, we were careful to maintain a ratio of at least 3.5:1 at the lowest carrier SF, which would provide a reasonable number of carrier cycles within each cycle of the envelope to enable judgement of the direction of envelope motion.

A possible future research direction might be to employ a task in which subjects identify the orientation of the envelope rather than its direction of motion. Because cortical neurons that encode CM stimuli are selective for the orientation as well as the direction of motion of the envelope (Li et al., 2014), this approach should also be similarly specific for downstream, cortical processing of Y-like input.

A natural concern when using CM stimuli is that an early nonlinearity in the display screen, or in the photoreceptors, might act to provide an artifactual luminance signal. We performed a careful measurement of the CRT display nonlinearity and incorporated that into the gamma correction of the video attenuator. Then we used the photometer to verify the linearity of luminance with intensity of the corrected video signal. The visual stimuli were always displayed in the center of the CRT (where calibration for gamma correction was performed), with eccentricity being varied by placement of the fixation target rather than the stimulus. After gamma correction, we also employed a diffusing sheet that acts as a spatial low-pass filter to verify that no luminance artifact was visible, thus indicating the calibration was successful in preventing nonlinear distortions (Scott-Samuel & Georgeson, 1999). To prevent possible luminance artifacts from the "adjacent pixel nonlinearity" (Schofield & Georgeson, 1999; Sukumar & Waugh, 2007), we used sinusoidal carrier stimuli that changed luminance smoothly along the line scan of the CRT display. Furthermore, if the responses to CM stimuli were due to either CRT or photoreceptor nonlinearities, one would not expect a bandpass dependence on carrier spatial frequency (**Figs. 3, 6**). In particular, the fall-off at low carrier SFs demonstrates that the mechanism underlying the CM responses is not secondary to simple luminance artifacts.

The results of this study suggest that tasks employing CM stimuli with high spatiotemporal carrier frequencies can reveal the nonlinear contributions of retinal Y-like cells to human perception. With this approach, we can reveal the SF-selective nonlinear behavior of Y-like cells, separate from the contributions to visual processing from other RGCs. To our knowledge, this is the first human study to employ CM stimuli at different eccentricities with the aim of isolating the nonlinear response function of Y-like RGCs.

Previous studies suggest that "first-order" stimuli, which are based on variations in luminance, and "second-order" stimuli, which are based on differences in local contrast and texture (such as a CM stimulus) (Cavanagh & Mather, 1989), engage distinct processing mechanisms (Cavanagh & Mather,1989; Manahilov et al., 2003; Schofield & Georgeson, 1999). Our results suggest that CM processing likely arises from Y-like cells, and is consistent with neurophysiological evidence from the responses of neurons in the LGN and early visual cortex to CM stimuli (Gharat & Baker, 2017; Rosenberg et al., 2010; Rosenberg & Issa, 2011). Since Y-like cells carry a mixture of first- and second-order information, an important future research direction will be to explore the relationship between the mechanisms underlying the results here and the various other behavioral findings that have supported either common or separate processing for first- and second-order stimuli (Allard & Faubert, 2008, 2013; Holliday & Anderson, 1994; Scott-Samuel & Georgeson, 1999; Smith & Ledgeway, 1997).

References

Allard R, Faubert J (2008) First- and second-order motion mechanisms are distinct at low but common at high temporal frequencies. J Vis 8:12.

Allard R, Faubert J (2012) No second-order motion system sensitive to high temporal frequencies. J Vis 12:773–773.

Baker CL Jr (1999) Central neural mechanisms for detecting second-order motion. Curr Opin Neurobiol 9:461–466.

Brainard DH (1997) The Psychophysics Toolbox. Spat Vis 10:433–436.

Cavanagh P, Mather G (1989) Motion: the long and short of it. Spat Vis 4:103-29.

Crook JD, Peterson BB, Packer OS, Robinson FR, Gamlin PD, Troy JB, Dacey DM (2008a). The smooth monostratified ganglion cell: evidence for spatial diversity in the Y-cell pathway to the lateral geniculate nucleus and superior colliculus in the macaque monkey. J Neurosci 28:12654-71.

Crook JD, Peterson BB, Packer OS, Robinson FR, Troy JB, Dacey DM (2008b) Y-cell receptive field and collicular projection of parasol ganglion cells in macaque monkey retina. J Neurosci 28:11277–11291.

Demb JB, Haarsma L, Freed MA, Sterling P (1999) Functional circuitry of the retinal ganglion cell's nonlinear receptive field. J Neurosci 19:9756–9767.

Demb JB, Zaghloul K, Haarsma L, Sterling P (2001a) Bipolar cells contribute to nonlinear spatial summation in the brisk-transient (Y) ganglion cell in mammalian retina. J Neurosci 21:7447–7454.

Demb JB, Zaghloul K, Sterling P (2001b) Cellular basis for the response to second-order motion cues in Y retinal ganglion cells. Neuron 32:711–721.

Efron B, Tibshirani RJ (1993) An Introduction to the Bootstrap. J R Stat Soc Series B Stat Methodol.

Enroth-Cugell C, Robson JG (1966) The contrast sensitivity of retinal ganglion cells of the cat. J Physiol 187:517–552.

Gharat A, Baker CL Jr (2017) Nonlinear Y-Like Receptive Fields in the Early Visual Cortex: An Intermediate Stage for Building Cue-Invariant Receptive Fields from Subcortical Y Cells. J Neurosci 37:998–1013.

Hochstein S, Shapley RM (1976) Linear and nonlinear spatial subunits in Y cat retinal ganglion cells. J Physiol 262:265–284.

Holliday IE, Anderson SJ (1994) Different processes underlie the detection of second-order motion at low and high temporal frequencies. Proc R Soc Lond B 257:165–173.

Johnson AP, Baker CL Jr (2004) First- and second-order information in natural images: a filter-based approach to image statistics. J Opt Soc Am A Opt Image Sci Vis 21:913-925.

Kleiner M, Brainard D, Pelli D, Ingling A, Murray R, Broussard C (2007) What's new in Psychtoolbox-3? Perception 36:1-16.

Koenderink JJ, Bouman MA, de Mesquita AEB, Slappendel S (1978) Perimetry of contrast detection thresholds of moving spatial sine wave patterns. I. The near peripheral visual field (eccentricity 0--8). JOSA 68:845–849.

Kulikowski JJ (1976) Effective contrast constancy and linearity of contrast sensation. Vision Res 16:1419–1431.

Lennie P (1980) Perceptual signs of parallel pathways. Philos Trans R Soc Lond B Biol Sci 290:23–37.

Li G, Yao Z, Wang Z, Yuan N, Talebi V, Tan J, Wang Y, Zhou Y, Baker CL Jr (2014) Form-cue invariant second-order neuronal responses to contrast modulation in primate area V2. J Neurosci 34:12081–12092.

Li X, Lu Z-L, Xu P, Jin J, Zhou Y (2003) Generating high gray-level resolution monochrome displays with conventional computer graphics cards and color monitors. J Neurosci Methods 130:9–18.

Livingstone M, Hubel D (1988) Segregation of form, color, movement, and depth: anatomy, physiology, and perception. Science 240:740–749.

Manahilov V, Simpson WA, Calvert J (2005) Why is second-order vision less efficient than first-order vision? Vision Res 45:2759–2772.

Merigan WH, Maunsell JH (1990) Macaque vision after magnocellular lateral geniculate lesions. Vis Neurosci 5:347–352.

Merigan WH, Byrne CE, Maunsell JH (1991) Does primate motion perception depend on the magnocellular pathway? J Neurosci 11:3422–3429.

Pelli DG (1997) The VideoToolbox software for visual psychophysics: transforming numbers into movies. Spat Vis 10:437–442.

Petrusca D, Grivich MI, Sher A, Field GD, Gauthier JL, Greschner M, Shlens J, Chichilnisky EJ, Litke AM (2007) Identification and Characterization of a Y-Like Primate Retinal Ganglion Cell Type. J Neurosci 27:11019–11027.

Robson JG (1966) Spatial and Temporal Contrast-Sensitivity Functions of the Visual System. J Opt Soc Am 56:1141.

Rosenberg A, Talebi V (2009) The primate retina contains distinct types of Y-like ganglion cells. J Neurosci 29:5048–5050.

Rosenberg A, Husson TR, Issa NP (2010) Subcortical Representation of Non-Fourier Image Features. J Neurosci 30:1985–1993.

Rosenberg A, Issa NP (2011) The Y Cell Visual Pathway Implements a Demodulating Nonlinearity. Neuron 71:348–361.

Rovamo J, Virsu V, Näsänen R (1978) Cortical magnification factor predicts the photopic contrast sensitivity of peripheral vision. Nature 271:54–56.

Schiller PH, Logothetis NK, Charles ER (1990) Role of the color-opponent and broad-band channels in vision. Vis Neurosci 5:321–346.

Schofield AJ, Georgeson MA (1999) Sensitivity to modulations of luminance and contrast in visual white noise: separate mechanisms with similar behaviour. Vision Res 39:2697–2716.

Schütt HH, Harmeling S, Macke JH, Wichmann FA (2016) Painfree and accurate Bayesian estimation of psychometric functions for (potentially) overdispersed data. Vision Res 122:105–123.

Shapley RM, Victor JD (1981) How the contrast gain control modifies the frequency responses of cat retinal ganglion cells. J Physiol 318:161–179.

Smith AT, Hess RF, Baker CL Jr (1994) Direction identification thresholds for second-order motion in central and peripheral vision. J Opt Soc Am A 11:506-514.

Smith AT, Ledgeway T (1997) Separate Detection of Moving Luminance and Contrast Modulations: Fact or Artifact? Vision Res 37:45–62.

Sukumar S, Waugh SJ (2007) Separate first- and second-order processing is supported by spatial summation estimates at the fovea and eccentrically. Vision Res 47:581–596.

Tanaka H, Ohzawa I (2006) Neural Basis for Stereopsis from Second-Order Contrast Cues. J Neurosci 26:4370–4382.

Watson AB, Ahumada AJ (2016) The pyramid of visibility. Electronic Imaging 2016:1–6.

Zhou YX, Baker CL Jr (1994) Envelope-responsive neurons in areas 17 and 18 of cat. J Neurophysiol 72:2134–2150.

Zhou, YX, Baker, CL Jr (1996). Spatial properties of envelope-responsive cells in area 17 and 18 neurons of the cat. J Neurophysiol, 75:1038–1050.