Brain endothelial STING1 activation by *Plasmodium*-sequestered heme promotes cerebral malaria via type I IFN response.

Teresa F. Pais[1]*, Hajrabibi Ali[1], Joana Moreira da Silva[1,2], Nádia Duarte[1], Rita Neres[1], Chintan Chhatbar[3,4], Rita C. Acúrcio[5], Rita C. Guedes[5], Maria Carolina Strano Moraes[6], Bruno Costa Silva[6], Ulrich Kalinke[3] and Carlos Penha-Gonçalves[1]*

Affiliations:

[1]Instituto Gulbenkian de Ciência; Oeiras, Portugal.

[2]CIISA, Faculdade de Medicina Veterinária, Universidade de Lisboa; Lisbon, Portugal.

[3] Institute for Experimental Infection Research, TWINCORE, Centre for Experimental and Clinical Infection Research, a joint venture between the Helmholtz Centre for Infection Research and the Hannover Medical School; Hannover, Germany.

[4]Institute of Neuropathology, Medical Faculty, University of Freiburg; Freiburg, Germany.

[5]Research Institute for Medicines (iMed.ULisboa), Faculty of Pharmacy, Universidade de Lisboa; Lisbon, Portugal.

[6]Systems Oncology Group, Champalimaud Research, Champalimaud Centre for the Unknown; Lisbon, Portugal

*Corresponding author. Email: mtpais@igc.gulbenkian.pt; cpenha@igc.gulbenkian.pt

Keywords: cerebral malaria, brain inflammation, brain endothelium**,** type I IFN, STING1

Abstract

Cerebral malaria (CM) is a life-threatening form of *Plasmodium falciparum* infection caused by brain inflammation. Brain endothelium dysfunction is a hallmark of CM pathology, which is also associated with the activation of the type I interferon (IFN) inflammatory pathway. The molecular triggers and sensors eliciting brain type I IFN cellular responses during CM remain largely unknown. We herein identified the stimulator of interferon response cGAMP interactor 1 (STING1) as the key innate immune sensor that induces *Ifnβ1* transcription in the brain of mice infected with *Plasmodium berghei* (*Pba*). This STING1/IFNβ-mediated response increases brain CXCL10 governing the extent of brain leucocyte infiltration and blood-brain barrier (BBB) breakdown, and determining CM lethality. The critical role of brain endothelial cells (BECs) in fueling type I IFN-driven brain inflammation was demonstrated in brain endothelial-specific IFNβ-reporter and STING1-deficient *Pba*-infected mice, which are significantly protected from CM lethality. Moreover, extracellular particles (EPs) released from *Pba*-infected erythrocytes activated STING1-dependent type I IFN response in BECs, a response requiring intracellular acidification. Fractionation of the EPs enabled us to identify a defined fraction carrying hemoglobin degradation remnants that activates STING1/IFNβ in the brain endothelium, a process correlated with heme content. Notably, stimulation of STING1-deficient BECs with heme, docking experiments and *in vitro* binding assays unveiled that heme is a putative STING1 ligand. This work shows that heme resultant from the parasite heterotrophic activity operates as an alarmin triggering brain endothelial inflammatory responses via STING1/IFNβ/CXCL10 axis crucial to CM pathogenesis and lethality.

Significance

CM results from loss of blood-brain endothelial barrier function caused by unrestrained inflammatory response in the natural course of infection by *Plasmodium* parasites. However, the role of brain endothelium in triggering inflammatory mechanisms is still undetermined. We found that the innate immune sensor STING1 is crucial for production of IFNβ in brain endothelial cells in *Plasmodium*-infected mice. This in turn stimulates CXCL10-mediated recruitment of leukocytes and subsequent brain inflammation and tissue damage. We identified within extracellular particles released from *Plasmodium*-infected erythrocytes, a fraction containing products of hemoglobin degradation, namely heme, which we show can bind STING1. Our results unravel a new angle of CM pathogenesis: heme contained in particles triggers the STING/IFNβ/CXCL10 axis in brain endothelial cells.

Introduction

Malaria, the disease caused by Plasmodium infection, accounted for over 400 thousand deaths worldwide in 2019 (World Malaria Report 2019). Despite the available malarial treatments, children under 5 years of age can develop cerebral malaria (CM) upon infection with *Plasmodium falciparum*, facing the highest risk of severe malaria morbidity and mortality (15-20%) (1). Brain swelling with increased compression of respiration centers in the brain stem predicts a fatal CM outcome (2). Children who survive CM are often affected by long-term neurological sequelae (1).

Infected erythrocytes (IE), parasite components released during the blood-stage infection (e.g. malarial hemozoin and labile heme), and systemic inflammatory factors (eg. CXCL10, TNF, IL1β and IFNγ) elicit an unfettered brain endothelial cell response that compromises regular endothelium function (reviewed in (3)). In particular, BBB integrity is lost, which results in leakage of plasma proteins and water into the cerebral parenchyma and consequent intracranial hypertension (reviewed in (4)). However, whether brain endothelial cells (BECs) are bystander activated by systemic inflammatory factors or the actual source of brain inflammation and pathology remains largely unexplored in CM.

Over the last years, the type I interferons, IFNα and IFNβ, emerged as key mediators in innate immune responses to malaria, promoting brain immunopathology (reviewed in (5)). In children with uncomplicated malaria, activation of the type I IFN response was associated with protective immunity (6). However, in children with CM, polymorphism in regulatory regions of type I IFN receptor (IFNAR1 subunit) gene were associated with increased type I IFN signaling and CM development (7). In the experimental mouse model of CM, despite the protective role of type I IFN in early infection (8, 9), several reports show that ablation of IFNAR1 signaling prevents BBB disruption and pathogenesis of experimental CM driven by parasite-specific cytotoxic CD8[+] T cells (10, 11). In addition, mice deficient in type I IFN signaling molecules—like transcription

factors interferon regulatory factor 3 (IRF3) and 7 (IRF7) and the upstream TAK-binding kinase 1 (TBK1)—are protected from CM (12). Moreover, a point mutation in the ubiquitin-specific protease 15 (USP 15) deters the type I IFN response in the mouse brain, which dampens brain inflammation and increases survival to CM (13).

Several pattern recognition receptors (PRRs) underlie inflammatory responses during *Plasmodium* infection by binding to *Plasmodium*-derived components (14), such as GPI anchors, plasmodial DNA and RNA, and hemozoin—a byproduct of hemoglobin digested by the parasite, consisting of crystals made of polymerized heme (15). Labile heme, a byproduct of hemolysis during *Plasmodium* infection, can also signal via PRR (16) and contribute to the pathogenesis of CM (17, 18).

The cGAS-STING1 pathway is a major sensor of cytosolic DNA that can recognize AT-rich motifs in plasmodial DNA and induce IFNβ transcription in human monocytes and mouse macrophages (12, 19, 20). Interestingly, extracellular vesicles derived from *Plasmodium*-IE (19) and complexes of plasmodial DNA with hemozoin (12) were proposed as carriers of parasite DNA into the cytosol of human monocytes.

BECs can also capture and cross-present malarial antigens to parasite-specific CD8$^+$ T cells (21). Two mechanisms by which malarial antigens could reach the endothelial cell endosomal and cytosolic compartments were proposed, one through membrane fusion with infected-erythrocytes, and another by uptake of merozoites and digestive vacuoles (21, 22).

Ubiquitous expression of type I IFN and IFNAR1 poses a significant challenge to the identification of specific cellular and molecular contributors to type I IFN signaling activation in CM development. Although BECs are within the first cells to sense the IE, their role in type I IFN-mediated innate immune responses during CM remains elusive. Here, we demonstrate that in the brain, specifically in BECs, STING1 activation by

112   parasite-sequestered heme present in particles released from Plasmodium-IE is a critical

113   mechanism underlying CM pathogenesis.

Results

Brain IFNβ production via STING 1 activation correlates with CM

There is indirect evidence connecting brain type I IFN responses with tissue pathology associated with CM in *Pba*-infected mice (23). We used an IFNβ-luciferase reporter mouse line (24) to directly test whether IFNβ induction in the brain was triggered by *Pba* infection. We detected strong luminescence (fig. S1A) and significantly increased luciferase activity in brain tissue as well as in the lung of mice developing CM (Fig. 1A). In contrast, mice infected with *PbNK65,* a parasite strain that does not cause CM, induced weak or no luminescence signal (fig. S1B). The IFNβ induction in the brain correlated with a type I response spread in three main brain regions—olfactory bulb, cerebrum and cerebellum—as indicated by the expression of four type I IFN response interferon-stimulated genes (*Irf1*, *Irf7*, *Ifit1* and *Cxcl10*) (fig. S1C). We then assessed the contribution of IFNβ to CM pathogenesis. BBB integrity was measured by EB quantification in the brain after perfusion at day 6 post infection (PI). BBB leakage of EB was lower in the brains of IFNβ KO than in the brains of wild type (WT) mice (Fig. 1B and fig. S1D). Accordingly, CM was less lethal in IFNβ KO mice than in WT controls (Fig 1D). This decrease in lethality was not attributable to differences in parasite burden between IFNβ KO and WT mice, and was not affected by sex (Fig. S1E).

Several PRRs sense *Plasmodium* infection and initiate downstream signaling converging at the phosphorylation and activation of type I interferon regulatory family of transcription factors (IRFs)(14). Pba-IE, *Plasmodium* AT-rich DNA motifs and parasite DNA in EPs that are released from *P. falciparum*-IE were shown to activate the STING1 pathway of type I IFN induction in mouse macrophages and human monocytes in culture (12, 19). However, the role of STING1 activation in the development of CM has not been addressed. We found that *Sting1* deletion significantly delayed death by CM and

conferred approximately 50% protection against CM lethality without affecting the

parasitemia (Fig. 1D). Although we cannot exclude minor contributions of the other

PRRs, we did not detect a significant difference in CM lethality when STING1 KO mice

were compared with quadruple KO mice for *Myd88*, *Trif*, *Mavs* and *Sting1* (MyTrMaSt

KO), which are impaired in all known type I response induction pathways (fig. 1F). Lower

levels of IFNβ induction, measured by luciferase activity, were also found in the brains

and lungs of IFNβ-reporter mice deficient for STING1 with CM (Fig. 1E). These results

indicate that STING1 is a major innate immune sensor of *Pba* infection ensuing IFNβ

production in the brain and subsequent BBB disruption and CM development.

STING1 activation induces *Cxcl10* gene expression and leukocyte recruitment to the

brain

In *Pba*-infected mice, brain innate immune responses associated occur before leukocyte

recruitment to this organ (13, 25). We used IFNβ-reporter mice in both alleles to measure

induction of type I IFN at early stages of infection. We detected a significant increase of

luciferase activity in the brain of mice as early as day 4 PI (Fig. 2A). Next, we tested

whether IFNβ induction by STING1 in the brain could drive expression of *Cxcl10*, a

known mediator of leukocyte recruitment downstream of IFN signaling. Deletion of *Ifnβ1*

or *Sting1* significantly decreased the levels of *Cxcl10* mRNA in the brain, at day 4 PI

(Fig. 2B).

Plasmodium infection induces CXCL10 in the brain, which in turn recruits CXCR3-

expressing antigen-specific pathogenic CD8[+] T cells (26–28). We analyzed whether brain

leukocyte infiltration was impaired in IFNβ KO and STING1 KO mice and found lower

percentage (Fig. 2C) and numbers of leukocytes (CD45[high]), namely CD8[+] T cells

(CD45$^+$TCRβ$^+$CD8$^+$) and monocytes (CD45^{high} Mac-1$^+$), in the brains of STING1 KO

mice (Fig. 2D) and of IFNβ KO mice (fig. S2A) than in those of WT mice. However, the

expression of CD8$^+$ T cell surface activation markers (CD44^{high}/ CD62L^{low}) and of

Granzyme B and IFNγ was similar in WT, STING1 KO (Fig. 2D) and IFNβ KO mice

(fig. S2A), which indicates that CD8$^+$ T cell activation was equally effective. The low

numbers of CD8$^+$ T cells in the brain of infected STING1 KO mice could not be attributed

to decreased local proliferation (25) as the frequency of BrdU$^+$ cells amongst CD8$^+$ T

cells was 44.5% ± 7.74 in STING1 KO mice similar to the 33.7% ± 6.95 found in WT

mice. Likewise, microglia activation assessed by MHC class I (25) expression was similar

in WT and KO mice (Fig. 2D and fig. S2A).

Notably, the decrease in the number of brain-infiltrating leukocytes in the brain was not

related to a decrease in the activation of splenic T cells or monocytes in STING1 KO or

IFNβ KO mice (fig. S2B). This suggests that innate immune activation of STING1

leading to IFNβ production and *Cxcl10* gene up-regulation in the brain contributes to

leukocyte recruitment into the brain and immunopathology in CM.

180 STING1 activation and IFNβ production by brain endothelial cells contributes to

181 CM

It is unclear whether specific brain cell types are responsible for the induction of IFNβ

response to blood-borne parasites. We investigated IFNβ production in different brain

cells during CM by using cell type-specific IFNβ-reporter mice (IFNβ$^{floxβ-luc/floxβ-luc}$ mice

bred to mice with Cre expression driven by cell-type specific promoters) (29). In these

mice one *Ifnβ1* allele is kept wild-type while the other allele drives luciferase activity by

the *Ifnβ1* floxed allele in cells that undergo Cre recombination. We found no significant

induction of luciferase activity in astrocytes (GFAPCre$^+$ mice) upon CM induction while

a clear response was obtained 24 hours after Poly I:C injection (Fig. 3A). On the other

hand, LysMCre$^+$ brain cells, including different types of myeloid cells (25), showed *Ifnβ1*

transcription activation upon infection as detected by increased luciferase activity in the

brain (Fig. 3B).

To determine whether IFNβ is induced in microglia, the brain resident macrophage

population, during CM, we used CX3CR1CreERT2$^{+/-}$ mice (30). After Cre induction with

tamoxifen, the monocyte compartment will be replenished with bone-marrow derived

cells without flox excision, while the self-renewing microglia will maintain the *Ifnβ1*-luc

reporter gene active and give a luciferase signal if triggered. Under these conditions, we

detected luciferase activity indicating *Ifnβ1* gene transcription in the microglial cell

compartment during CM (Fig. 3C).

Induction of *Ifnβ1* in the endothelial compartment was assessed in reporter mice

expressing Cre in endothelial and hematopietic cells under the Tie2 promoter (Fig. 3D),

and in reporter mice carrying inducible Cre under the Slco1c1 promoter activated

specifically in brain endothelial/ependymal cells (31) (Fig. 3E). Induction of luciferase

activity in both mouse lines showed that brain endothelial cells contribute to local brain

production of IFNβ in the brain during CM.

To determine the impact of cell-specific expression of STING1 or IFNβ in CM

development, we specifically deleted these genes in different cell types by using cell-

lineage specific Cre expression in mice carrying two floxed alleles or one inactivated

allele and one floxed allele for the target gene. We found that absence of STING1 or IFNβ

in LysM$^+$ or CX3CR1$^+$ cells did not prevent death by CM (Fig. 3F and fig. S3A). In

contrast, ablation of STING1 or IFNβ in endothelial and hematopoietic cells in mice

expressing Cre under Tie2 promoter (Fig. 3G and fig. S3B) or ablation of STING1

specifically in brain endothelial cells in Slco1c1CreERT[+/-] mice significantly protected

from CM lethality (Fig. 3H). In addition health/neurologic score on day 6 PI was

improved (fig. S3C) while parasitemia was unaffected (fig. S3D). Furthermore, mice with

Tie2[+]-induced STING1 deletion that survived CM showed clinical improvement after day

8 PI (fig. S3E), while parasite burden was not reduced (fig. S3F).

Overall, these results unveil that STING1-dependent activation of IFNβ production in

brain endothelial cells starting early in infection is a critical determinant of brain

immunopathology during CM development.

IE-derived EPs induce IFNβ via STING1 after intracellular acidification in brain

endothelial cells

The generation of extracellular particles (100-1000 nm) by erythrocytes has been

associated with the pathogenesis of human and rodent CM (32, 33). Also, vesicles

released from *P. falciparum*-IE activate expression of pro-inflammatory cytokines (i.e

TNF and IFNβ) in innate immune cells (19, 34). We asked whether extracellular particles

(EPs) obtained from the serum of *P. berghei*-infected mice or from IE cell cultures can

induce type I IFN signaling in primary cultures of brain endothelial cells (BECs).

Nanoparticle tracking analysis of EPs collected from mouse serum showed significantly

increased mean particle size in *Pba*-infected, CM-sick mice than in non-infected mice

(Fig. 4A and fig. S4A). Negative-stain transmission electron microscopy (TEM) (Fig. 4B)

confirmed the presence of large particles with more than 200 nm in the serum of infected

mice, while EPs isolated from non-infected animals were around 100 nm, the exosome

size range (Fig. 4A). Moreover, the concentration of particles within the range of 100.5-

265.5 nm increased approximately 3-fold in the blood of infected mice compared to non-infected controls (fig. 4B). The particles isolated from blood of *Pba*-infected mice that differ from those found in non-infected mice were mostly from leukocyte (CD45$^+$) and erythrocytic origin (TER119$^+$) as revealed by antibody staining for cell-type surface markers (Fig. 4C).

In BECs cultures, particles isolated from the blood of mice with CM activated IFNβ gene transcription (Fig 4D) and protein expression (fig. S4C), but not those from non-infected mice. Similarly, EPs obtained from IE cultures activated *Ifnβ1* transcription measured by the luciferase activity in IFNβ-reporter BECs (Fig. 4E). In contrast, neither erythrocytes from non-infected mice (NIE) nor erythrocytes from *Pba*-infected mice (IE) induced luciferase activity (Fig. 4E). To determine whether endosomal acidification was required for endothelial response, cells were pretreated with ammonium chloride, which elevates the intralysosomal pH (35). Remarkably, induction of IFNβ in response to EPs was abolished in BECs pre-exposed to ammonium chloride (Fig. 4F). Furthermore, luciferase activity was lower in IFNβ-reporter STING1 KO BECs stimulated with IE-derived EPs (Fig. 4G). This result indicates that *Sting1* deletion in BECs reduces induction of IFNβ by IE-derived EPs. In sum, EPs derived from infected erythrocytes are processed in acidic compartments in BECs, activate STING1 and induce IFNβ.

Particles emanating from the parasite's digestive-vacuole induce *Ifnβ1* and *Cxcl10* in brain endothelial cells

To identify bioactive factors associated with EPs that trigger type I IFN response in BECs, we established a protocol to fractionate bulk EPs obtained from *P. berghei*-IE cultures on basis of a combination of differential centrifugation, paramagnetic separation and

presence of host erythrocytic membrane (Fig. 5A). We found that compared to total EPs, the fraction (Fr1) pulled down at lower centrifugation (5630 g) induced higher *Ifnβ1* transcription measured by luciferase activity (fig. S5A). Therefore, we further fractionated Fr1 using a magnetic LS column to separate a fraction enriched in particles containing iron and hemozoin (Fr2) from the flow through (Fr3) (Fig. 5A). To isolate particles with erythrocytic membrane (TER119[+]), Fr3 was incubated with biotinylated anti-TER119 antibody and streptavidin magnetic beads (Fig. 5A). The retained fraction in immuno-magnetic separation originated the Fr4, and the flow through originated Fr5. Flow cytometry analysis (Fig. 5B and fig. S5B) showed that compared to Fr1, Fr2 and Fr5 are enriched in TER119 negative particles. *Pba*-GFP parasites were used to generate IE cultures, thus, the GFP signal in Fr5 indicate the presence of parasite-derived material. TEM of fixed samples confirmed that Fr2 contained vesicles with hemozoin crystals likely derived from parasite's digestive vacuoles (Fig. 5C). Among all fractions, the hemozoin-enriched Fr2, was the strongest inducer of luciferase activity in BECs. Compared to total EPs ($2x10^6$ EPs), a 10-fold lower amount ($2x10^5$ EPs) of Fr2 induced luciferase activity (Fig. 5D). In BECs, Fr2 was also a potent inducer of *Cxcl10* (Fig. 5E) and *Irf7* (fig. S5C) mRNA. Importantly, Fr2-induced expression of these genes was about two-fold lower in STING1 KO BECs than in WT BECs and was abolished in IFNβ KO and in quadruple MyTrMaSt KO BECs (Fig. 5E and fig. S5C). Nevertheless, these KO BECs expressed *Cxcl10* and *Irf7* in response to direct stimulation with IFNβ, which indicates that the IFNβ-IFNAR axis was preserved (fig. S5D). These results indicate that Fr2, the IE-derived particle fraction containing digestive vacuole remnants, strongly induces a type I IFN response in BECs, which is at least partially dependent on STING1.

Labile heme originated by the parasite activates STING1 in brain endothelial cells

We then addressed the role of hemozoin in STING1 activation by inhibiting the hemozoin

synthesis pathway in IE cultures with chloroquine, an anti-malarial drug described to

increase the levels of labile heme in the parasite (15). Unexpectedly, EPs obtained from

chloroquine-treated IE cell cultures induce stronger IFNβ expression than EPs from

untreated IE (Fig. 6A). Both hemozoin and labile heme can activate PRRs and contribute

to inflammatory storm and BBB dysfunction in CM (14, 16, 17). We stimulated BECs

directly with different amounts of heme and hemozoin. While hemozoin was unable to

stimulate BECs, 20 μM of heme induced significant luciferase activity in WT but not in

STING1 KO IFNβ-reporter BECs (Fig. 6B and fig. S5E). Moreover, heme induced

Cxcl10 gene expression in WT BECs, as previously described (36), but not in STING1

KO BECs (Fig. 6C).

Following these results, we tested a putative interaction of heme with STING1 by

performing docking calculations. GOLD software was used to dock heme into the

structure of STING1 and search for the optimal pose (orientation, conformation and

binding energy) between them. We found that the heme is positioned within the STING1

binding site for its ligand c-di-GMP, mostly overlapping with it (Fig. 6D). This shows

that heme can position itself in the pocket in a propitious way for interaction with the

STING1. Although well positioned, heme has less affinity to STING1 than c-di-GMP

(ChemPLP score of 61 for heme compared to 104 for c-di-GMP). We next tested the

ability of STING1 to bind heme by examining the peroxidase-like activity and migration

shift of recombinant human STING1 upon incubation with heme. We detected peroxidase

activity of STING1-heme complexes at the molar ratio of 1:150, while protein shift was

detected at a molar ratio of 1:60 (Fig 6E), values that are similar to those reported for

other heme-binding proteins in this assay (37).

Taken together, the results indicate that heme-containing EPs derived from IE are able to

induce type I IFN responses in BECs in a STING1-dependent manner, possible involving

direct heme binding to STING1.

Discussion

We identified STING1 activation in brain endothelial cells as a molecular pathway that mediates neuroinflammation and brain pathology underlying the pathogenesis of CM. We demonstrated that, early in infection, the brain endothelium uses STING1 to sense *Plasmodium* blood-stage infection and initiate a type I IFN response through activation of *Ifnβ1* and *Cxcl10* gene transcription. This is associated with leukocyte recruitment and subsequent loss of BBB function, a hallmark of CM. We found that extracellular particles derived from IE, containing hemozoin and enriched for labile heme, trigger STING1-dependent IFNβ responses in brain endothelial cells (Fig. 6F).

Our study identifies IFNβ as a critical trigger of IFNAR signaling and a determinant of BBB disruption and lethality in CM. Whole and cell-specific IFNβ-reporter mice enabled us to identify BECs, as well as brain monocytes/microglia, as main producers of IFNβ, and STING1 as the key PRR upstream of IFNβ induction. STING1—a main activator of *Ifnβ1*transcription—also controlled early *Cxcl10* gene induction in the brain of *Pba*-infected mice. CXCL10 is known to promote chemotaxis of CXCR3 expressing CD8[+] T cells to the central nervous system during *Toxoplasma* infection (28) and has been strongly linked to CM severity both in humans (38, 39) and in rodents (27). We observed that both IFNβ KO and STING1 KO have lower numbers of CD8[+] T cells and monocytes in the brain infiltrates. Thus, our data support a role for CXCL10 in inducing chemotactic leukocyte migration into the brain.

We provide the first demonstration that activation of STING1 in BECs promotes CM pathology. Infected mice whose brain endothelial/ependymal cells are made deficient in STING1 (Slco1c1CreERT STING1 floxed mice) displayed milder neurological deficits and increased survival. Several studies reported that plasmodial DNA activates STING1 in innate immune cells such as monocytes and macrophages (12, 19, 20). However, we

found that specific *Ifnβ1* or *Sting1* deletion in myeloid and microglia cells did not affect

CM development in mice. Instead, our data indicate that BECs are the main cell drivers

of the STING1-IFNβ axis in the brain during CM.

STING1 is localized in the ER and does not bind DNA directly, but interacts with cyclic

dinucleotides, namely cGAMP. cGAMP is produced by cyclic-GMP-AMP synthase

(cGAS) upon dsDNA sensing in the cytosol. After translocation into the Golgi apparatus,

STING1 and TBK1 phosphorylate IRF3 that promotes transcription of type I IFN genes

(40). Interestingly, a population of exosomes isolated from *P. falciparum* (Pf) IE cultures

at the ring stage, could transfer Pf-DNA to PBMCs and activate the cGAS-

STING1pathway with consequent induction of *Ifn*β1/α gene expression (19). Our study

further highlights the likely important role for IE-derived particles in the activation of

innate immune mechanisms during malaria. We found that IE-derived particles larger

than exosomes and enriched in vesicles containing hemozoin are able to activate *Ifnβ* and

Cxcl10 transcription in BECs. This activation depends on intracellular organelle

acidification suggesting a requirement for endosome maturation and/or acidic hydrolase

activity (35). IE-derived particles containing hemozoin could represent digestive vacuoles

that accumulate heme and/or hemoglobin. Although plasmodial DNA-bound to hemozoin

has been shown to activate cyclic GMP-AMP (cGAS) upstream STING1 in macrophages

(12), hemozoin itself does not seem to be required for STING1 activation in BECs. In

fact, exposure of IE to chloroquine, which reduces hemozoin and increases labile heme

bioavailability, enhances IFNβ in BECs. This prompted us to hypothesize that heme could

be involved in STING1 activation mediated by IE-derived EPs. Indeed, we observed

STING1-dependent up-regulation of *Ifnβ* and *Cxcl10* in the presence of labile heme at

concentrations within the ranges found in the plasma of children with CM (41) and known

to activate *Cxcl10* transcription in endothelial cells (36). Recently, hemolysis in sickle

cell disease and injection of hemin in mice were shown to activate and recruit monocytes

to the liver through activation of type I IFN signaling and independently of TLR4 (42).

Our docking experiments and protein binding assay indicate that heme can bind STING1

in vitro. However, further evidence of direct binding of heme to STING1 in the ER is

required to clarify whether heme is a direct or indirect activator of STING1 in BECs.

Interestingly, late endosomes tightly contact with the ER (43) indicating that heme-

STING1 interaction may occur during the association between acidified mature

endosomes containing heme and the ER (Fig. 6f).

Overall, we demonstrate that STING1 in the brain endothelium senses molecular by

products of erythrocyte infection by *Plasmodium*, such as labile heme, initiating a type I

IFN response mediated by IFNβ with induction of CXCL10 in early phases of infection.

This, in turn, triggers a cascade of events that promote brain inflammation, including

leukocyte recruitment, and may also affect neuronal responses. This study suggests that

type I IFN targeted strategies may minimize brain immunopathology, thus preventing

fatal CM and subsequent neurological sequels in survivors.

Material and Methods

Mice and treatments

All procedures involving laboratory mice were in accordance with national (Portaria

1005/92) and European regulations (European Directive 86/609/CEE) on animal

experimentation and were approved by the Instituto Gulbenkian de Ciência Ethics

Committee and the Direcção-Geral de Alimentação e Veterinária, the national authority

for animal welfare.

Mice were housed and bred in the facilities of Instituto Gulbenkian de Ciência. $Myd88^{-/-}$

(MyD88 KO) mice were originally a gift from Prof. Shizuo Akira (Osaka University).

$Mda5^{-/-}$ mice were obtained from Dr. Maria Mota (IMM, Portugal). Transgenic mice

expressing Cre under the GFAP promoter (GFAPCre) in C57BL/6 background were

obtained from Dr. Henrique Veiga Fernandes (Champalimaud Foundation, Portugal).

$Ifn\beta1^{-/-}$ (IFNβ KO), $Ifn\beta1^{+/\Delta\beta\text{-}luc}$ (IFNβ-reporter mice), $Ifn\beta1^{flox\beta\text{-}luc/flox\beta\text{-}luc}$ (IFNβ $^{flox\beta\text{-}}$

$^{luc/flox\beta\text{-}luc}$), $Trif^{-/-}$ (TRIF KO), B6(Cg)-$Tmem173^{tm1.2Camb}$/J ($Tmem173^{-/-}$, STING1 KO),

Tg(Slco1c1-icre/ERT2)1Mrks (Slco1c1CreERT2) and $Myd88^{-/-}Trif^{-/-}Mavs^{-/-}Sting1^{-/-}$

(MyTrMaSt KO) mice were obtained from the TWINCORE, Centre for Experimental and

Clinical Infection Research, Hanover, Germany.

C57BL/6J, B6;129-$Mavs^{tm1Zjc}$/J (MAVS KO), B6;SJL-$Sting1^{tm1.1Camb}$/J (STING1$^{flox/flox}$) ,

B6.Cg-Tg(Tek-cre)1Ywa/J (Tie2Cre), B6J.B6N(Cg)-$Cx3cr1^{tm1.1(cre)Jung}$/J (CX3CR1Cre),

B6.129P2(Cg)-$Cx3cr1^{tm2.1(cre/ERT2)Litt}$/WganJ (CX3CR1CreERT2) and B6.129P2-

$Lyz2^{tm1(cre)Ifo}$/J mice (LysMCre) were originally purchased from The Jackson Laboratory

(Bar Harbor, ME) (Table S1). MAVS KO mice were backcrossed to C57BL/6J mice up

to 96.82% of recipient genome. IFNβ$^{flox\beta\text{-}luc/flox\beta\text{-}luc}$ and STING1$^{flox/flox}$ were crossed with

the different Cre lines to generate cell lineage conditional IFNβ-reporter, IFNβ KO mice

or STING1 KO mice. Experimental animals carrying one loxP-flanked and one KO allele

of $Sting1$ were generated by crossing Slco1c1Cre^{ERT2}STING1$^{flox/flox}$ or CX3CR1Cre^{ERT2}

STING1$^{flox/flox}$ mice with STING1 KO mice. The efficiency of the conditional *Sting1*

deletion was expected to be improved by having just one floxed allele. Crossing of

STING1$^{flox/flox}$ mice with CX3CR1CreSTING1$^{+/+}$ or Tie2Cre STING1$^{+/+}$ mice generated

a deleted loxP-flanking Sting1 allele in the germ line. These mice were then crossed with

STING1$^{flox/flox}$ and originated Cre^{+} mice with one deleted *Sting1* allele while another

allele remained loxP-flanked and was conditionally deleted. Mice with either a deletion

of the floxed allele or a KO allele but also expressing a floxed *Sting1* allele were

designated STING1$^{flox/-}$.

Wild type, deleted and floxed alleles of *Ifnβ1* and *Sting1* were identified in experimental

animals by genotyping using a 3-primer strategy (Table S2). For induction of *Sting1*

deletion, 4-6 week old male and female mice were treated with tamoxifen (2 mg/10 g

body weight) in corn oil administered by gavage every other day, totaling five doses. Mice

were allowed to recover from tamoxifen treatment for 4 weeks.

For *in vivo* labelling of proliferating cells, mice received four intraperitoneal injections of

BrdU (50 mg/kg,) every two hours. Brains were collected for FACS analysis two hours

after the last injection and intracardiac perfusion of the mice with PBS.

As a positive control for luciferase activity in IFNβ-reporter mice in GFAP^{+} cells, mice

received an intraperitoneal injection (i.p.) of 0.4 mg of Poly I:C.

Parasites and infection

Mice were infected with 10^{6} IE with the following parasite strains: *Plasmodium berghei*

ANKA (*Pba*) or *Pba*-GFP (44) to induce CM or with *Pb* NK65, a strain that proliferates

in mice but does not induce neurological signs of CM.

Frozen IE stocks were expanded in C57BL/6 mice prior to infection. Parasitemia in mice

infected with non-GFP parasites (*Pba* and *Pb* NK65) and GFP-*Pba* was determined by

flow cytometry analysis as the percentage of DRAQ5$^+$ erythrocytes (blood samples

incubated with DRAQ5™, 3 uM) and GFP$^+$ erythrocytes, respectively (LSRFortessa™

X-20 cell analyser, BD Biosciences and FACSDiVa software version 6.2).

Evaluation of CM

Neurological score

Wild-type mice developed severe signs of CM such as head deviation, paralysis, and

convulsions at day 6–7 post infection (PI) and died within 4–6 h of severe disease

development. In different genetic backgrounds death by CM occurred up to 11 days PI

("CM time window"). Mice resistant to CM died around day 20 with hyperparasitemia

(~60%) without signs of severe neurological dysfunction.

Disease severity was evaluated in mice before and after infection following an adapted

health score for CM (45). From day 4 PI mice were scored based on eleven parameters:

gait/body posture, grooming, motor performance, balance, hindlimb clasping reflex, limb

strength, toe pinch, aggressiveness, head deviation, paralysis and convulsions. For each

parameter a maximum score of two was given to good response or health condition

(normal gait, full body extension, clean and sheen hair, exploring of 3-4 corners in 30 sec,

extending front feet on the wall of the cage, normal plantar reaction, active pull away, toe

pinch reaction, normal aggressiveness, no convulsions, no paralysis or no head deviation).

Clear signs of CM (convulsions, paralysis and head deviation) with no strength, no

balance and no toe pinch reaction scored zero. Healthy mice scored 22 and mice dying of

CM could score between 3-16 points at day 6 PI.

Blood–brain barrier integrity assay

At day 6 PI, when C57BL/6J WT mice show signs of CM, the different mouse groups

were killed 1 hour after retro-orbital injection of 100 μl of 2% Evans blue (EB) dye

(Sigma) and were perfused intracardially with 15 ml of PBS. Brains were dissected,

weighed, and pictured. EB retained in the brain tissue was extracted by immersion in 2

ml formamide (Merck) at 37°C in the dark for 48 h and measured by spectrophotometry

at 620 and 740 nm (Thermo Scientific™ Multiskan™ GO Microplate

Spectrophotometer). Amounts of EB were calculated using an EB standard and expressed

as mg EB/g of brain tissue.

Immune cells analysis

Spleen (day 5 PI) and brain (day 6 PI) single cell suspensions of infected and non-infected

C57BL/6J and KO mice were immuno-stained for T cell and monocyte/macrophage

activation cell surface markers. Brain single-cell suspensions were prepared as described

before (25). Briefly, mice were perfused with 20 ml of PBS via the left ventricle of the

heart. Brains were removed and homogenized in HBSS containing collagenase VIII (0.2

mg/ml) (Sigma). After incubation for 45 min at 37 °C, the digested tissue was minced,

forced through a nylon cell strainer of 100 μm (BD Falcon™) and centrifuged. The pellet

was resuspended in a gradient of 30% Percoll (Amersham Biosciences) in PBS and

centrifuged at 520 g for 30 min at RT to obtain a pellet with brain mononuclear cells.

Spleen and brain mononuclear cells were incubated with anti-mouse Fc block in FACS

buffer (2% FCS and 0.01% NaN3 in PBS) for 30 min at 4 °C to block irrelevant binding

of the antibodies to the Fc receptor of myeloid cells. After washing, cells were stained

with defined cocktails of fluorochrome-conjugated antibodies (Table S3). For

intracellular staining, cells were incubated for 5h in RPMI complete medium containing

eBioscience™ Cell Stimulation Cocktail (Invitrogen). Cells were then washed and fixed

and permeablized with BD Cytofix/Cytoperm™ (BD) before staining with anti-mouse

Granzyme B-efluor 450 and anti-mouse IFNγ-PE (BioLegend). Brain cells isolated from

Pba-infected mice injected with BrdU were stained using the FITC BD BrdU Flow Kit

(BD Pharmingen™) according to the manufacturer's instructions. Stained cells were

resuspended in FACS buffer with 5,000 of 10-µm counting beads (SureCount™

Standards, Bang Laboratories) and analyzed by flow cytometry (LSR Fortessa X20™,

BD). Data were analyzed with FlowJo v10 software and cell types identified according

to the gate strategy (Fig. S6)

Isolation of extracellular particles (EPs)

All the solutions were filtered (0.22 µm) before use. Blood EPs were obtained from mouse

blood collected by cardiac puncture with a syringe containing 100 µl of EDTA, 0.5 M.

Plasma was prepared by two successive centrifugations at 1500 g for 15 minutes at 4ºC.

Harvested plasma was further centrifuged at 20 000g for 25 minutes and the pellet

containing EPs was ressuspended in 0.2% BSA in PBS (46).

IE-derived EPs were prepared from the supernatant of IE cultures. Whole blood of *Pba*

infected mice (day 5-6 PI, 10-20% parasitaemia) was collected by cardiac puncture with

a syringe containing heparin and suspended in 5 ml of RPMI 1640 medium (Biowest),

followed by centrifugation at 450*g* for 10 min. The pellet was resuspended in RPMI 1640

medium containing Glutamax and Neomycin (0.05 mg/ml), supplemented with 20% FBS

(Biowest) without or with chloroquine, 8 µM, and then incubated in culture flasks

overnight in an orbital shaker at 36.5ºC and 50 rpm. EPs were isolated adapting a

previously described procedure (34) (Fig. 5a). Briefly, the cell suspension was first

centrifuged at 600g to remove IE and non-IE and then centrifuged at 1600g for 15min to

remove cell debris. The supernatant of this centrifugation was then sequentially

centrifuged at 5630g and 20 000g for 20 min (total EPs). The pellet resulting from

centrifugation at 5630g was designated fraction 1 (Fr1) and the supernatant was further

centrifuged at 20 000g producing a pellet, "–Fr1". Fr1 was resuspended in PBS and

passed through a LS magnetic exclusion column according to the manufacturer's

instructions (MACS Milteny Biotec). Particles retained in this column were designated

Fr2. The flow through (Fr3) was centrifuged at 10 000g for 10 min and the obtained pellet

incubated with biotinylted anti-mouse TER119 antibody (1 μg/ml) (Invitrogen). The

pellet was then washed twice and resuspended in 80 μl of labeling buffer (PBS, 2 mM

EDTA) containing 20 μl of Anti-Biotin MicroBeads (Milteny Biotec). After incubation

for 15 min at 4ºC and washing, the pellet was resuspended in separation buffer (0.5%

BSA, 2 mM EDTA in PBS) followed by magnetic separation through a second LS

column. Particles retained in this column and flow through were designated Fr4 and Fr5,

respectively.

Characterization of EPs

NanoSight

Blood-derived MPs were analyzed for particle concentration and size distribution by the

NS300 Nanoparticle Tracking Analysis (NTA) system with a red laser (638 nm)

(NanoSight –Malvern Panalytical, United Kingdom). Samples were pre-diluted in PBS

to a concentration range of $5x10^7$-$1x10^9$ particles/ml and 10-50 particles/frame for

optimal NTA analysis. Video acquisitions were performed using camera level 16 and

threshold 5. Five videos of 30s each were captured per sample. Analysis of particle

concentration per milliliter and size distribution were performed using the NTA software

v3.4.

Flow cytometry

Blood MPs or IE-derived EPs were pre-incubated with Fc-block and then stained with PE-conjugated anti-mouse Ter119, CD41, CD45 and CD105 antibodies (1:100) (Life Technologies) during 20 min. After staining, the samples were washed in PBS (10 000g, 10 min, 4ºC) and the pellet was resuspended in PBS and acquired with flow cytometer LSRFortessa™ (BD Biosciences, run with FACSDiVa software version 6.2) using side scatter of blue violet laser (SSC-BV) and FSC-PMT tuned to detect small particles. PBS was used to set background signal and unstained samples used to define PE-negative particles. Total EPs numbers were calculated using 5000 counting beads (10 µm) in each sample.

Transmission electron microscopy

For negative staining, 5 µl of the pellet of serum MPs in HEPES buffer (10 mM) was adhered to glow discharged Mesh100 grids coated with 2% formvar in chloroform and carbon for 2 min. Following attachment, samples were washed with distilled H_2O and stained with 2% uranyl acetate for 2 min.

EPs samples were fixed with 2% formaldehyde and 2.5% glutaraldehyde in 0.1 M phosphate buffer (PB) overnight at 4ºC. After washing three times, samples were embedded for support in 2% low melting point agarose. After solidifying, samples were post-fixed with 1% osmium tetroxide in 0.1 M PB for 30 min at 4ºC. Samples were then washed in PB and distilled water and pre-stained with 1% tannic acid for 20 min at 4ºC and with 0.5% uranyl acetate for 1h at RT in the dark followed by dehydration using increasing concentrations of ethanol. After dehydration, samples were infiltrated and embedded using a graded series of Embed-812 epoxy resin polymerized at 60ºC overnight. Ultrathin sections (70 nm) were obtained with a Leica UC7 Ultramicrotome

and collected in 2% formvar in chloroform-coated slot grids that were post-stained with

1% uranyl acetate and Reynold's Lead Citrate for 5 min each.

Electron microscopy images were acquired on a FEI Tecnai G2 Spirit BioTWIN transmission electron microscope operating at 120 keV and equipped with an Olympus-SIS Veleta CCD Camera.

Brain endothelial cell cultures

Mouse brains were removed from the skull and kept in a petri dish with HBSS medium (Biowest) on ice. The brains were transferred to a freezer block covered with gauze and the meninges removed by rolling a cotton swab over each brain. Brain blood vessels were then isolated adapting a previously described procedure (47). Briefly, brains were minced with a blade in 1 ml of HBSS medium and aspirated with a syringe. The tissue was then homogenized by pushing it five times through a 23G needle. This homogenate was mixed with equal volume of cold 30% (wt/vol) dextran (MW ~60 kDa) (Sigma) in PBS and then centrifuged at 10 000g for 15 min. After centrifugation the myelin top layer was removed and the pellet containing the vessels was suspended in DMEM and passed through a 40-µm cell strainer. Five milliliters of medium was added to wash away other cells and myelin debris. The strainer was back-flushed with 5 ml of MCDB 131 to collect the brain vessels. These were digested by supplementing the medium with 1mg/ml of collagenase type IV (Millipore), 10 µg/ml DNAse (Roche) and 2% FBS, followed by incubation for 2-3 hours at 36.5 °C in an orbital shaker. The digested vessels were then centrifuged and suspended in EGM™-2 Endothelial Cell Growth Medium-2 BulletKit™ (LONZA) supplemented with GlutaMax (ThermoFisher), 10% of FBS and 4 µg/ml puromycin, and cultured in 24 well-plates. Medium was changed for fresh medium without puromycin every four days.

When cultures reached 50-80% confluence (~14 days, 95% of the cells are CD31$^+$ by

flow cytometry analysis), cells were exposed for 18-20h to different amounts (2x10^5-

2x10^6) of IE and NIE, EPs fractions (Total EPs, -Fr1 and Fr1 to Fr5), MPs (present in 600

μl of serum), or different activators of IFNβ signaling (IFNβ, 1000 U/ml; LPS, 1μg/ml;

Poly I:C, 1-5 μg/ml). To inhibit endosome acidification, BECs were pre-incubated for 20

min with 20 mM of NH4Cl before adding the stimuli.

Heme was added as ferric chloride heme (hemin, SIGMA) to BECs at a final

concentration range of 10-30 μM. A freshly prepared stock solution was obtained by

dissolving hemin in NaOH 0.1 M for 1h at 37°C and then adding 1M Tris, pH 7.5 to a

final concentration of 0.1 M. HCl was added to the solution to bring the pH back to 7.5.

Molecular docking

STING1-c-di-GMP crystal structure (PDB ID: 6S26) was retrieved from Protein Data

Bank (PDB). The crystal structure was then prepared using the Molecular Operating

Environment (MOE v2019.01) structure preparation wizard by stripping out the water

molecules. Hydrogen atoms were added, and protonation states were assigned using

Protonate-3D tool at pH=7.4 of the MOE software package. The co-crystallized ligand

(c-di-GMP) was also prepared using the MOE software package. The EHT10 force field

was used in all minimization steps. Again, hydrogen atoms were added using the

Protonate-3D tool at pH=7.4.

C-di-GMP was then removed and re-docked into the corresponding STING1 crystal

structure. Self-docking was performed in GOLD v5.7 software using ChemPLP as

scoring function. The docking binding site was centered at threonine T263 (Fig. S5G), a

crucial residue for c-di-GMP and STING1 interaction, and a search radius of 15 Å was

set. Upon self-docking successfully validated the developed protocol, molecular docking

of heme was performed, 500 GA runs. Heme followed the same ligand preparation

procedure as c-di-GMP.

Peroxidase activity and gel migration shift assay

Recombinant human STING1 protein (Gentaur) (0.3 μg, 1μM) in 20 mM Tris-HCl, 0.15

M NaCl, pH=8.0 (Tris buffer) was incubated for 15 min at room temperature (RT) with

3000, 600, 300, 150, 60 and 6 μM of heme. After adding Tris sample buffer (pH=6.8) for

native PAGE (no SDS or reducing agents), the samples were loaded on 12% Criterion^{TM}

TGX^{TM} Stain-Free^{TM} precast gels for PAGE (Bio-Rad). Heme-STING1 complexes were

separated by gel electrophoresis in non-denaturing running buffer. Peroxidase-like

activity of heme bound to STING1 was detected on the gel after incubation with TMB

Substrate Reagent Set (BD Biosciences). After developing a blue-colored band, the gel

was washed with 0.5 M sodium acetate (pH=5) and isopropanol (30%). Migration shift

of protein-heme complexes (37) was examined after washing the gel in Milli-Q water and

staining with Commassie Brilliant Blue R250 (0.05% in 50% of methanol and 10% of

acetic acid in H_2O) for 2 hours at RT, followed by incubation with destaining solution

(10% etanol and 7.5% acetic acid in H_2O). STING1 without incubation with heme and

heme alone were used as negative controls for peroxidase-like activity. Serum bovine

albumin (3.3 μg, 5μM) incubated with 600 μM of heme was used as a positive control

(48).

Gene expression

RNA was prepared from total mouse brains of mice and from BECs using Trizol and

Cells-to-CT kit (Thermo Fisher Scientific), respectively. One microgram of total brain

RNA was converted to cDNA (Transcriptor High Fidelity cDNA Synthesis Kit; Roche).

A final volume of 20 µl of the cDNA reaction was diluted 1:3 in RNase-free water to be

used in qRT-PCR. *Ifnβ1*, *Cxcl10*, *Irf7*, *Irf1* and *Ifit1* mRNA were quantified using

Mm00439552_s1, Mm00445235_m1, Mm00516788_m1, Mm01288580_m1 and

Mm00515153_m1 TaqMan Gene Expression Assays from ABI, respectively.

IFNβ-luciferase reporter assay

Tissues were lysed in Glo Lysis Buffer (Promega) followed by centrifugation at 20,000g

for 20 min. BECs were gently washed with PBS before lysis. 50 µl of tissue or cell lysates

were mixed with 50µl of Bright-Glo™ Luciferase Assay System (Promega) in a 96-well

luminometer plate. The luminescence was immediately measured with GloMax®

Explorer (Promega) (excitation, 405nm; emission, 500-550nm; integration time, 5).

Protein was quantified by Bradford Assay (BioRad) and luciferase activity expressed as

RFU per mg of protein.

Satistical Analysis

Statistical analysis was performed using Prism v6.01 software (GraphPad Software, Inc.).

Comparisons between treated and non-treated conditions used two-tailed unpaired t-test,

comparisons between multiple groups used one-way ANOVA, comparisons between

treated cells of different genotypes used two-way ANOVA as indicated in figure legends.

Bars in figures represent the mean (±SD) and each symbol represents one mouse or well

as indicated in the figures. Sample size and number of the independent experiments

performed can be found in the figures and figure legends.

Acknowledgments

We gratefully acknowledge the Electron Microscopy Unit at IGC, in particular Ana

Laura Sousa for technical support. We thank M. Soares and S. Gomes for discussions

and critical reading of the manuscript. This work was supported by ERA-NET

NEURON.

References

1.   R. Idro, K. Marsh, C. C. John, C. R. J. Newton, Cerebral malaria: Mechanisms of
brain injury and strategies for improved neurocognitive outcome. *Pediatr. Res.*
68, 267–274 (2010).

2.   K. B. Seydel, *et al.*, Brain Swelling and Death in Children with Cerebral Malaria.
N. Engl. J. Med. **372**, 1126–1137 (2015).

3.   J. Dunst, F. Kamena, K. Matuschewski, Cytokines and Chemokines in Cerebral
Malaria Pathogenesis. *Front. Cell. Infect. Microbiol.* **7** (2017).

4.   U. Frevert, A. Nacer, Fatal cerebral malaria: a venous efflux problem. *Front.*
Cell. Infect. Microbiol. **4**, 1–7 (2014).

5.   I. Sebina, A. Haque, Effects of type I interferons in malaria. *Immunology* **155**,
176–185 (2018).

6.   M. Krupka, *et al.*, Mild Plasmodium falciparum malaria following an episode of
severe malaria is associated with induction of the interferon pathway in
Malawian children. *Infect. Immun.* **80**, 1150–1155 (2012).

7.   C. M. Feintuch, *et al.*, Type i interferon receptor variants in gene regulatory
regions are associated with susceptibility to Cerebral Malaria in Malawi. *Am. J.*
Trop. Med. Hyg. **98**, 1692–1698 (2018).

8.   A. M. Vigário, *et al.*, Recombinant Human IFN-α Inhibits Cerebral Malaria and
Reduces Parasite Burden in Mice. *J. Immunol.* **178**, 6416–6425 (2007).

9.   A. Hassan, *et al.*, A virus hosted in malaria-infected blood protects against T cell-
mediated inflammatory diseases by impairing DC function in a type I IFN-
dependent manner. *MBio* **11** (2020).

10.   E. A. Ball, *et al.*, IFNAR1 controls progression to cerebral malaria in children
and CD8+ T cell brain pathology in Plasmodium berghei-infected mice. *J.*
Immunol. **190**, 5118–27 (2013).

11.   C. L. Edwards, *et al.*, Spatiotemporal requirements for IRF7 in mediating type I
IFN-dependent susceptibility to blood-stage Plasmodium infection. *Eur. J.*
Immunol. **45**, 130–141 (2015).

12.   S. Sharma, *et al.*, Innate Immune Recognition of an AT-Rich Stem-Loop DNA
Motif in the Plasmodium falciparum Genome. *Immunity* **35**, 194–207 (2011).

13.   S. Torre, *et al.*, USP15 regulates type i interferon response and is required for
pathogenesis of neuroinflammation. *Nat. Immunol.* **18**, 54–63 (2017).

14.   R. T. Gazzinelli, P. Kalantari, K. A. Fitzgerald, D. T. Golenbock, Innate sensing
of malaria parasites. *Nat. Rev. Immunol.* **14**, 744–757 (2014).

15.   S. E. Francis, D. J. Sullivan, D. E. Goldberg, Hemoglobin metabolism in the
malaria parasite Plasmodium falciparium. *Annu. Rev. Microbiol.* **51**, 97–123
(1997).

16.   M. P. Soares, M. T. Bozza, Red alert: Labile heme is an alarmin. *Curr. Opin.*
Immunol. **38**, 94–100 (2016).

17.   A. Pamplona, *et al.*, Heme oxygenase-1 and carbon monoxide suppress the
pathogenesis of experimental cerebral malaria. *Nat. Med.* **13**, 703–710 (2007).

18.   A. Ferreira, *et al.*, Sickle hemoglobin confers tolerance to plasmodium infection.
Cell **145**, 398–409 (2011).

19.   X. Sisquella, *et al.*, Malaria parasite DNA-harbouring vesicles activate cytosolic
immune sensors. *Nat. Commun.* **8** (2017).

20.   C. Gallego-Marin, *et al.*, Cyclic GMP–AMP Synthase Is the Cytosolic Sensor of
Plasmodium falciparum Genomic DNA and Activates Type I IFN in Malaria . *J.*
Immunol. **200**, 768–774 (2018).

21.   S. W. Howland, C. M. Poh, L. Rénia, Activated Brain Endothelial Cells Cross-
Present Malaria Antigen. *PLOS Pathog.* **11**, e1004963 (2015).

22.   R. Jambou, *et al.*, Plasmodium falciparum adhesion on human brain
microvascular endothelial cells involves transmigration-like cup formation and
induces opening of intercellular junctions. *PLoS Pathog.* **6**, 1–13 (2010).

23.   F. E. Lovegrove, *et al.*, Expression microarray analysis implicates apoptosis and
interferon-responsive mechanisms in susceptibility to experimental cerebral
malaria. *Am. J. Pathol.* **171**, 1894–1903 (2007).

24.   S. Lienenklaus, *et al.*, Novel Reporter Mouse Reveals Constitutive and
Inflammatory Expression of IFN-β In Vivo. *J. Immunol.* **183**, 3229–3236 (2009).

25.   T. F. Pais, S. Chatterjee, Brain macrophage activation in murine cerebral malaria
precedes accumulation of leukocytes and CD8+ T cell proliferation. *J.*
Neuroimmunol. **163**, 73–83 (2005).

26.   E. W. Sorensen, *et al.*, CXCL10 stabilizes T cell-brain endothelial cell adhesion
leading to the induction of cerebral malaria. *JCI insight* **3** (2018).

27.   G. S. V Campanella, *et al.*, Chemokine receptor CXCR3 and its ligands CXCL9
and CXCL10 are required for the development of murine cerebral malaria. *Proc.*
Natl. Acad. Sci. U. S. A. **105**, 4814–9 (2008).

28.   T. H. Harris, *et al.*, Generalized Lévy walks and the role of chemokines in
migration of effector CD8 + T cells. *Nature* **486**, 545–548 (2012).

29.   E. Solodova, J. Jablonska, S. Weiss, S. Lienenklaus, Production of IFN-β during
730   listeria monocytogenes infection is restricted to monocyte/macrophage lineage.
731   *PLoS One* **6**, 2–11 (2011).

732   30.   S. Yona, *et al.*, Fate Mapping Reveals Origins and Dynamics of Monocytes and
733   Tissue Macrophages under Homeostasis. *Immunity* **38**, 79–91 (2013).

734   31.   D. A. Ridder, *et al.*, TAK1 in brain endothelial cells mediates fever and lethargy.
735   *J. Exp. Med.* **208**, 2615–2623 (2011).

736   32.   J. B. Pankoui Mfonkeu, *et al.*, Elevated Cell-Specific Microparticles Are a
737   Biological Marker for Cerebral Dysfunctions in Human Severe Malaria. *PLoS*
738   *One* **5**, 13415 (2010).

739   33.   F. El-Assaad, J. Wheway, N. H. Hunt, G. E. R. Grau, V. Combes, Production,
740   Fate and Pathogenicity of Plasma Microparticles in Murine Cerebral Malaria.
741   *PLoS Pathog.* **10** (2014).

742   34.   P. Y. Mantel, *et al.*, Malaria-infected erythrocyte-derived microvesicles mediate
743   cellular communication within the parasite population and with the host immune
744   system. *Cell Host Microbe* **13**, 521–534 (2013).

745   35.   I. Mellman, R. Fuchs, A. Helenius, ACIDIFICATION OF THE ENDOCYTIC
746   AND EXOCYTIC PATHWAYS. *Annu. Rev. Biochem.* **55**, 663–700 (1986).

747   36.   C. M. Dickinson-Copeland, *et al.*, Heme-mediated induction of CXCL10 and
748   depletion of CD34+ progenitor cells is toll-like receptor 4 dependent. *PLoS One*
749   **10** (2015).

750   37.   E. Karnaukhova, *et al.*, Characterization of heme binding to recombinant Î±1-
751   microglobulin. *Front. Physiol.* **5**, 465 (2014).

752   38.   H. B. Armah, *et al.*, Cerebrospinal fluid and serum biomarkers of cerebral
753   malaria mortality in Ghanaian children. *Malar. J.* **6** (2007).

754   39.   V. Jain, *et al.*, Plasma IP-10, apoptotic and angiogenic factors associated with
755   fatal cerebral malaria in India. *Malar. J.* **7** (2008).

756   40.   D. Wan, W. Jiang, J. Hao, Research Advances in How the cGAS-STING
757   Pathway Controls the Cellular Inflammatory Response. *Front. Immunol.* **11**, 615
758   (2020).

759   41.   R. E. Elphinstone, *et al.*, Dysregulation of the haem-haemopexin axis is
760   associated with severe malaria in a case-control study of Ugandan children.
761   *Malar. J.* **14**, 511 (2015).

762   42.   Y. Liu, *et al.*, Type I interferon is induced by hemolysis and drives antibody-
763   mediated erythrophagocytosis in sickle cell disease. *Blood* **138**, 1162–1171
764   (2021).

43.   J. R. Friedman, J. R. DiBenedetto, M. West, A. A. Rowland, G. K. Voeltz,
Endoplasmic reticulum-endosome contact increases as endosomes traffic and
mature. *Mol. Biol. Cell* **24**, 1030–1040 (2013).

44.   B. Franke-Fayard, *et al.*, A Plasmodium berghei reference line that constitutively
expresses GFP at a high level throughout the complete life cycle. *Mol. Biochem.*
Parasitol. **137**, 23–33 (2004).

45.   R. W. Carroll, *et al.*, A rapid murine coma and behavior scale for quantitative
assessment of murine cerebral malaria. *PLoS One* **5**, 1–12 (2010).

46.   S. Dinkla, R. Brock, I. Joosten, G. J. Bosman, Gateway to understanding
microparticles: Standardized isolation and identification of plasma membrane-
derived vesicles. *Nanomedicine* **8**, 1657–1668 (2013).

47.   S. W. Howland, S. Y. Gun, C. Claser, C. M. Poh, L. Rénia, Measuring antigen
presentation in mouse brain endothelial cells ex vivo and in vitro. *Nat. Protoc.*
10, 2016–2026 (2015).

48.   F. A. Englert, *et al.*, Labile heme impairs hepatic microcirculation and promotes
hepatic injury. *Arch. Biochem. Biophys.* **672**, 108075 (2019).

 Figure legends

 Fig. 1. Cerebral malaria pathogenesis is associated to type I IFN responses induced

 by STING1 activation in *Plasmodium* infection.

(A) *Ifnβ1* gene expression in the indicated organs of IFNβ-Luc reporter (IFNβ$^{+/Δβ\text{-luc}}$)

mice infected with *Pba* (day 6 PI) or non-infected was ascertained by *ex vivo* luciferase

activity normalized for tissue protein contents. The present data are representative of three

independent experiments. **(B)** Blood-brain barrier integrity was evaluated by Evans Blue

(EB) quantification in the brain after mouse perfusion. Data were normalized for brain

weight in WT and IFNβ KO mice at day 6 PI. **(C)** Survival curve of WT and IFNβ KO

mice infected with 10^6 *Pba*-IE. **(D)** Survival and blood parasitemia curves in mice

deficient for STING1. **(E)** Induction of *Ifnβ1* gene expression in the indicated organs of

IFNβ-Luc reporter (IFNβ$^{+/Δβ\text{-luc}}$STING1$^{+/-}$) and in IFNβ-Luc reporter lacking STING

(IFNβ$^{+/Δβ\text{-luc}}$STING1$^{-/-}$) mice infected with *Pba* (day 6 PI) was ascertained by *ex vivo*

luciferase activity normalized for tissue protein amounts.

Dashed line in **(C)** and **(D)** delimits the time-window of death by CM. Data points in **(A)**,

(B) and **(E)** represent individual mice. Data in **(A)**, **(B)** and **(E)** are analyzed by one-way

ANOVA, Tukey's multiple comparisons test. Data in **(C)** and **(D)** comprise two

independent infection experiments and are compared between mouse strains with Gehan-

Breslow-Wilcoxon test.

Fig. 2. Leukocyte brain infiltration is mediated by STING1 induction of *Ifnβ* and

***Cxcl10* in the brain.**

(A) *Ifnβ1* gene expression in the brain of mice with two reporter alleles for *Ifnβ1* (IFNβ$^{Δβ\text{-}}$

$^{\text{luc}/Δβ\text{-luc}}$) at day 4 of infection was ascertained by *ex vivo* luciferase activity normalized for

tissue protein amounts. **(B)** *Cxcl10* gene expression (qPCR) in the brain of WT and IFNβ

and STING1 KO mice at day 4 of infection. **(C)** Representative flow cytometry dot blots

showing the percentage of CD45^{high} cells within brain infiltrates of WT and STING1 KO

mice. **(D)** Flow cytometry analysis of brain leukocytes isolated from non-infected or *Pba*-

infected (day 6 PI) WT and STING1 KO mice. Extracellular staining against CD45 in

combination with staining for TCRβ, CD8 and Mac-1 identified CD8$^+$T cells (CD45$^+$

TCRβ$^+$ CD8$^+$), monocytes (CD45^{high} Mac-1$^+$), and microglia (CD45^{int}Mac-1$^+$). Activated

CD8$^+$T cells were identified by extracellular expression of CD62L and CD44 and

intracellular staining for Granzyme B and IFNγ. MHC I expression identified activated

microglial cells. Experimental groups were compared using one-way ANOVA with

Tukey's multiple comparisons test.

Data points represent individual animals in **(A)**, **(B)** and **(D)**. Data are analyzed by

unpaired, two-tailed *t*-test in **(A)** and **(B)**.

Fig 3. **STING1 expression on brain endothelial cells is a determinant of IFNβ**

induction and CM lethality.

Ifnβ1 gene expression was ascertained by *ex vivo* luciferase activity in mice infected with

Pba (day 6 PI) and non-infected mice carrying a IFNβ-Luc reporter conditionally

activated in cells expressing **(A)** GFAP (glial cells), **(B)** LysM (myeloid cells), **(C)**

CX3CR1 (microglia cells), **(D)** Tie-2 (endothelial and hematopoietic cells) or **(E)** Slco1c1

(brain endothelial cells). I.p. injection of Poly I:C was used as positive response control

of *Ifnβ1* gene induction. Results comprise two independent infection experiments and

data are analyzed with unpaired, two-tailed *t*-test.

Survival curves after *Plasmodium* infection in mice carrying conditional or induced

STING deletion driven by **(F)** CX3CR1Cre and CX3CR1CreERT

(monocytes/microglia), **(G)** Tie2Cre (endothelial cells) or **(H)** Slco1c1CreERT (brain

endothelial). Results represent at least three independent infection experiments for each

mouse strain. Survival curves were compared by Gehan-Breslow-Wilcoxon test.

Fig. 4. EPs isolated from blood and IE induce IFNβ in brain endothelial cells after

acidic processing and in a STING1-dependent manner.

(A) NanoSight analysis of mean particle size after serum centrifugation (20 000g) of

symptomatic *Pba*-infected mice (day 6 PI) and non-infected control mice. Results

represent two independent infection experiments. Non-infected mice were compared with

Pba-infected mice by unpaired, two-tailed *t*-test. **(B)** TEM image of negative stained EPs

obtained from the serum of mice with CM or non-infected animals. **(C)** Representative

flow cytometry histograms of serum EPs pooled from 3-5 animals and stained with the

indicated cell lineage membrane markers. EPs were collected after 20 000g centrifugation

from non-infected (full histogram) and infected (empty histogram) mice. Data shown are

representative of three independent experiments. **(D)** *Ifnβ1* RNA quantification in BECs

stimulated with MPs isolated from the serum (600 μl) of symptomatic *Pba*-infected mice

(day 6 PI) (EPs-CM, n=6 independent isolation experiments) and non-infected control

mice (EPs-NI, n=3 independent isolation experiments). Exposure to LPS (1 μg/ml) and

Poly I:C (5 μg/ml) were used as positive response controls. **e**, Measurement of luciferase

activity in IFNβ-reporter BECs exposed to two different doses ($2x10^5$ and $2x10^6$) of

erythrocytes isolated from the blood of non-infected (NIE) or *Pba*-infected mice (IE), or

with EPs obtained from IE-cultures centrifuged at 20 000g. One representative

experiment of two performed in duplicate or triplicate. **(F)** Activation of *Ifnβ1* gene

expression measured by luciferase activity in IFNβ-reporter BECs treated or not with

NH4CL (20 mM) and then exposed to EPs collected from *Pba*-IE cell cultures. One

representative experiment of three performed in triplicate is shown. **(G)** *Ifnβ1* gene

expression measured by luciferase activity in IFNβ-reporter WT or STING1 KO BECs

stimulated with EPs. Results comprise data of two independent experiments done in

triplicate. Treatments are compared between the two genotypes by two-way ANOVA

using Tukey's multiple comparisons test.

Data are presented as mean values ± SD in **(D-G)** and analyzed by one-way ANOVA

using Tukey's multiple comparisons test in **(D-F)**.

Fig. 5. Particles derived from the parasite's digestive-vacuole activate type I IFN

response in brain endothelial cells.

(A) Experimental workflow for EP fractionation from *Pba*-IE cultures combining multi-

step centrifugation with magnetic and immunomagnetic separation to obtain five fractions

(Fr1-5). Fr2 is isolated by magnetic field capture of iron-enriched EPs. Fr4 is obtained by

immuno-separation of EPs expressing the erythrocyte surface marker Ter119. **(B)** Flow

cytometry analysis of EPs staining for TER119 and/or GFP-positive. Data comprising

four independent experiments were compared to Fr1 by two-way ANOVA, Dunnett's

multiple comparisons test. ****$p<0.0001$, **$p<0.01$. **c**, TEM imaging of Fr2 (fixed

sample) showing membrane structures surrounding hemozoin crystals. **(D)** Measurement

of luciferase activity in IFNβ-reporter BECs stimulated with each one of the 5 different

EPs fractions; at $2x10^5$ EPs/fraction. Data comprise two (Fr4), four (Fr1 and Fr3), six

(Fr5) and ten (Fr2) independent isolation experiments and each spot represents a cell

culture well. One-way ANOVA, Tukey's multiple comparisons test. **(E)** *Cxcl10* gene

expression in BECs stimulated with blood of *Pba*-infected mice containing 2×10^5 IE and

with 2×10^5 EPs of different fractions. Data comprise 3-4 experiments of different BECs

and EPs isolations and are compared between genotypes by two-way ANOVA using

Tukey's multiple comparisons test, $*p < 0.05$, $**** p < 0.0001$. Each spot represents a cell

culture well.

Fig. 6. Heme sequestered by the parasite induce *Ifnβ1* and *Cxcl10* gene expression

in BECs through STING1 activation

(A) Luciferase activity in IFNβ-reporter BECs exposed to 2×10^6 EPs from *Pba*-IE

cultured in the presence or absence of chloroquine (8 μM). One representative experiment

of two performed in triplicate is shown. **(B)** Luciferase activity in IFNβ-reporter BECs

stimulated with hemin or hemozoin at the indicated concentrations or with IFNβ (1,000

U/ml). Data comprise 2-3 independent experiments and each point represents a cell

culture well. **(C)** *Cxcl10* gene expression in WT or STING1 KO BECs stimulated with

different concentrations of hemin for 20h. Data comprise two independent experiments.

Comparison between genotypes were performed by two-way ANOVA, Tukey's multiple

comparisons test, $* p < 0.05$. **(D)** STING1-Heme molecular docking results. Close-up view

of STING1 pocket with superposition of best-scoring docked pose of c-di-GMP and

Heme. c-di-GMP and Heme are colored in yellow and cyan sticks, respectively.

Molecular docking was performed using GOLD software and ChemPLP as scoring

function. **(E)** Detection of peroxidase activity and migration shift assay of STING1 (1μM)

incubated with increasing doses of heme. Albumin (5 μM) incubated with heme was used

as positive control for peroxidase activity and heme alone was used as negative control.

(F) Schematic model of STING1 activation in brain endothelium by parasite-derived

heme-containing particles and its contribution to the pathogenesis of cerebral malaria.

Hb, hemoglobin; DV, digestive vacuole; *Pba*, *Plasmodium berghei*; RBC, red blood cell;

BEC, brain endothelial cell; MΦ, monocyte; Mic., microglia; TBK1, TANK binding

kinase 1; IRF3; interferon regulatory factor 3; ISRE; interferon-sensitive response

element.

Data are presented as mean values ± SD in **(A-C)** and analyzed by one-way ANOVA

using Tukey's multiple comparisons test in **(A)** and **(B).**

Fig. 1

Fig. 2

Fig. 3, Pais et al

A Size

Mean particle size (nm)

p=0.0007

○ NI (n=6)
● INF (n=8)

NI   INF

B

NI

178 nm

112 nm

500 nm

CM

633 nm

514 nm

500 nm

C

CD41
(Platelets)

CD45
(Leukocytes)

CD105
(Endothelial)

TER119
(RBCs)

D IFNβ mRNA expression

Fold change

p=0.0066

p=0,042

CTR   Poly I:C   LPS   EPs_NI   EPs_CM

E

Fold change (RFU)

□ 2x10⁵
■ 2x10⁶

p=0.0035

CTR   NIE   IE   EPs

F

Fold change (RFU/mg)

p<0.0001

CTR   EPs   NH4Cl   NH4Cl+EPs

G

Fold change (RFU)

■ WT
■ STING1 KO

p<0.001

CTR   EPs   IFNβ

Fig. 4

Fig. 5

Fig. 6

Supplementary Information for

Brain endothelial STING1 activation by *Plasmodium*-sequestered heme promotes cerebral malaria via type I IFN response.

Pais TF *et al.*

Corresponding author: Teresa F. Pais, mtpais@igc.gulbenkian.pt; Carlos Penha-Gonçalves, cpenha@igc.gulbenkian.pt

The PDF includes:

Tables S1 to S3
Figures S1 to S6

Strain	Source	JAX mice database stock #	Reference
$B6;Ifn\beta1^{-/-}$	Dr. Kalinke TWINCORE		(49)
$B6;Ifn\beta1^{+/\Delta\beta\text{-luc}}$	Dr. Kalinke TWINCORE		(50)
$B6;Ifn\beta1^{\text{flox}\beta\text{-luc/flox}\beta\text{-luc}}$	Dr. Kalinke TWINCORE		(29)
$B6;Trif^{/-}$	Dr. Kalinke TWINCORE	005037	
B6(Cg)-$Tmem173^{tm1.2Camb}$/J ($Tmem173^{-/-}$, STING KO)	Dr. Kalinke TWINCORE	025805	
$B6;Myd88^{-/-}Trif^{-/-}Mavs^{-/-}Tmem173^{-/-}$ (MyTrMaSt KO)	Dr. Kalinke TWINCORE		(51)
$B6;Myd88^{-/-}$	Dr. Akira Osaka University		(52)
$B6;Mda5^{-/-}$	Dr. Mota IMM	015812	
B6;129-$Mavs^{tm1Zjc}$/J	JAX	008634	
B6;SJL-$Sting1^{tm1.1Camb}$/J (STING1$^{\text{flox/flox}}$)	JAX	031670	
Tg(Slco1c1-icre/ERT2)1Mrks (Slco1c1CreERT2)	Dr. Kalinke TWINCORE		(31)
GFAPCre in C57BL/6 background	Dr. Veiga Fernandes Champalimaud Foundation	004600	
B6.Cg-Tg(Tek-cre)1Ywa/J (Tie2Cre)	JAX	008863	
B6J.B6N(Cg)-$Cx3cr1^{tm1.1(cre)Jung}$/J (CX3CR1Cre)	JAX	025524	
B6.129P2(Cg)-$Cx3cr1^{tm2.1(cre/ERT2)Litt}$/$WganJ$ (CX3CR1CreERT2)	JAX	021160	
B6.129P2-$Lyz2^{tm1(cre)Ifo}$/J mice (LysMCre)	JAX	004781	
GFAPCre$^{+/-}$IFNβ$^{+/\text{ flox}\beta\text{-luc}}$	This study		
LysMCre$^{+/-}$IFNβ$^{+/\text{ flox}\beta\text{-luc}}$	This study		
CX3CR1CreERT$^{+/-}$IFNβ$^{+/\text{ flox}\beta\text{-luc}}$	This study		
Tie2Cre$^{+/-}$IFNβ$^{+/\text{ flox}\beta\text{-luc}}$	This study		
Slco1c1CreERT2$^{+/-}$IFNβ$^{+/\text{ flox}\beta\text{-luc}}$	This study		
LysMCre$^{+/-}$IFNβ$^{\text{flox}\beta\text{-luc / flox}\beta\text{-luc}}$	This study		
CX3CR1Cre$^{+/-}$IFNβ$^{\text{flox}\beta\text{-luc / }\Delta\text{-luc}}$	This study		
Tie2Cre$^{+/-}$IFNβ$^{\text{flox}\beta\text{-luc / flox}\beta\text{-luc}}$	This study		
CX3CR1CreERT$^{+/-}$STING1$^{\text{flox/-}}$	This study		
CX3CR1Cre$^{+/-}$STING1$^{\text{flox/}\Delta}$	This study		
TieCre$^{+/-}$STING1$^{\text{flox/}\Delta}$	This study		
Slco1c1CreERT$^{+/-}$STING1$^{\text{flox/-}}$	This study		

Table S1. Strains used in this study.

Oligonulceotides	Genotyping
5'-CACAGGCCATGAAGGAAGAT-3'	*Ifnb1* wt, floxed and deleted alleles-
5'-TTGGAGCTGGAGCTGCTTAT-3'	
5'-GCCTTATGCAGTTGCTCTCC-3'	
5'-GCGCACACACTAAAAACTG-3'	*Sting1* wt, floxed and deleted alelles
5'-TTGCTAGTAGCTACTGAAAGGC-3	
5'-TTTCATCTGCCTTCCAGGTC-3'	

Table S2. Oligonucleotides used to genotyping *Ifnβ1* and *Sting1* modified alleles.

Antibody	Supplier	Clone
Anti-mouse-Fc-block/CD16/32	IGC (antibody facility)	2.4G2
Anti-mouse FITC CD45.2	IGC (antibody facility)	104.2
Anti-mouse PE CD45.2	IGC (antibody facility)	104.2
Anti-mouse PE CD41	eBiosciences	MWReg30
Anti-mouse PE CD45	Life Technologies	30-F11
Anti-mouse PE CD105	Life Technologies	MJ7/18
Anti-mouse PE TER119	Life Technologies	TER119
Anti-mouse efluor 450 CD45.2	eBiosciences	104
Anti-mouse biotinylated TCR beta	IGC (antibody facility)	H57-597
Anti-mouse Alexa 648 CD8	IGC (antibody facility)	YTS169.4
Anti-mouse Pacific Blue CD8	IGC (antibody facility)	YTS169.4
Anti-mouse biotinylated CD4	IGC (antibody facility)	GK1.5
Anti-mouse eFluor 450 CD44	IGC (antibody facility)	IM7
Anti-mouse PE CD69	IGC (antibody facility)	H1.2F3
Anti-mouse PE CD62L	IGC (antibody facility)	MEL-14
Anti-mouse Alexa 647 CD25	IGC (antibody facility)	PC61
Anti-mouse biotinylated CD11b/Mac-1	IGC (antibody facility)	M1/70
Anti-mouse Alexa 488 MHC-I/H-2Kb	IGC (antibody facility)	AF6-88.5
Anti-mouse efluor 450 Granzyme B	BioLegend	N6ZB
Anti-mouse PE IFNγ	BioLegend	XMG1.2

Table S3. Antibodies used for flow cytometry.

Fig. S1

Fig. S1. **Role of IFNβ signaling in CM development.**

(A) Ex vivo luminescence (upper panel) and corresponding light imaging (lower panel) of spleen (Sp) lung (Lu), liver (Liv), abdominal fat (Fat) and brain (Br) of IFNβ-reporter mice infected with *Pba* (day 5, 20 % parasitaemia). **(B)** *In vivo* luminescence imaging of IFNβ-reporter mice infected with *Pba* (10-20% parasitaemia with CM), injected with a IFNβ inducer (Poly I:C) or with a strain of *P. berghei* that does not induce CM (PbNK65) (day 18, 13% parasitaemia). **(C)** Interferon response signature detected in the designated brain regions was evaluated by measuring interferon-stimulated gene expression by quantitative PCR (qPCR). Data points represent individual mice from two independent infection experiments. Symptomatic *Pba*-infected mice (day 6 PI) were compared to non-infected mice using unpaired, two-tailed *t*-test (p values on the figure). **(D)** Photograph of brains collected from WT and IFNβ KO mice infected with *Pba* (day 6 PI) after Evans blue i.v. and perfusion. **(E)** Survival and parasitemia curves of WT and IFNβ KO male and female mice, infected with the indicated amounts of *Pba*-IE. **(F)** Survival and parasitemia curves of WT, STING1 KO and quadruple KO mice for IFNβ critical components of IFNβ induction (*MyD88$^{-/-}$*, *Trif$^{-/-}$*, *Mavs$^{-/-}$* and *Sting1$^{-/-}$*) infected with *Pba*. Protection from CM development in quadruple KO mice, is recapitulated in the STING1 KO alone.

Fig. S2

Fig. S2. **Decreased leukocyte infiltration in the brain is not associated with deficient cell activation.** **(A)** Flow cytometry analysis of brain leukocyte infiltration in WT and IFNβ KO mice infected with *Pba*. The number of infiltrating CD8$^+$ T cells and monocytes are lower in IFNβ KO while cell activation profiles are maintained. **(B)** Flow cytometry analysis of splenocytes in *Pba*-infected WT, IFNβ KO and STING1 KO mice at day 5 PI. The number of peripheral CD8$^+$ T cells, CD4$^+$ T cells and monocyte are increased in STING KO mice but the activation profile is not affected. Non-infected (NI) and *Pba*-infected mice of the different genotypes were compared using two-way ANOVA (* p<0.05, ** p<0.01 and **** p<0.0001).

Fig. S3

Fig. S3. **Development of CM in mice with *Ifnβ1* or *Sting1* conditional deletion in myeloid and endothelial compartments**.

(A) Survival of *Pba*-infected mice with conditional IFNβ deletion targeting macrophages (LysM-Cre), microglia (CX3CR1-Cre) and **(B)** endothelial cells (Tie2-Cre). Gehan-Breslow-Wilcoxon test, **p<0.01 **(C)** Health/neurological score of mice with STING-specific deletion in the different cellular compartments. Non-infected animals have a score of 22 that decreases in mice with CM at day 6 PI. Data were compared by unpaired, two-tailed *t*-test. **p<0.01, ***p<0.001 by Mann Whitney test. **(D)** Parasitemia at day 4 and day 6 in conditional STING1 KO mice. **(E)** Kinetics of the health/neurological score, and **f**, parasitemia in Tie2⁺ cell-specific conditional STING1 KO mice.

Fig. S4

Fig. S4. **Characterization of extracellular particles obtained from serum of CM affected mice and from *Pba*-IE cultures**.

(A) Representative histogram of the size and concentration of EPs obtained from non-infected and CM wild-type mice. **(B)** NanoSight analysis of concentration of particles

within the range of 100.5-265.5 nm in EPs collected from symptomatic *Pba*-infected mice (day 6 PI) and non-infected control mice. **(C)** In vitro IFNβ secretion in brain endothelial cells stimulated with particles obtained from the serum of *Pba*-infected animals (EPs-CM) or *Pba*-IE cultures (EPs-IE), measured by ELISA in cell culture supernatants. Data were compared by unpaired, two-tailed *t*-test.

Fig. S5

Fig. S5. Identification of IFNβ-inducing factors in fractions derived from EPs collected from *Pba*-IE cultures.

(**A**) IFNβ gene expression measured by luciferase activity in wild IFNβ-reporter BECs stimulated with 2×10^6 of total EPs or of Fr1. Results are representative of two independent experiments. One-way ANOVA, Tukey's multiple comparisons test. (**B**) Flow cytometry profiles of EPs fractions obtained from *PbaGFP*-IE cultures and stained for the erythrocytic marker TER119. (**C**) Induction of *Irf7* expression in BEC of the indicated genotypes upon stimulation with IE and EPs fractions. 4XKO refers to quadruple KO (*Myd88*[-/-], *Trif*[-/-], *Mavs*[-/-] and *Sting1*[-/-]). *Irf7* expression is induced by Fr2 in WT cells but is ablated in absence of endogenous IFNβ expression. Data comprise 3-4 experiments. Comparison between genotypes were performed by two-way ANOVA using Tukey's multiple comparisons test, $*p < 0.05$, $****\ p < 0.0001$. Each spot represents a cell culture well. (**D**) Induction of *Cxcl10* and *Irf7* expression in BEC of indicated genotypes stimulated with IFNβ (1000 U/ml) showing response to IFNAR signaling in the different genotypes. (**E**) Luciferase activity of STING1 KO IFNβ-reporter BECs exposed to different concentrations of hemin for 20 hours. Data comprise two different experiments. (**F**) Surface view of c-di-GMP pocket. Close-up view of specific recognition of c-di-GMP by STING1 and detailed interactions between ribose-phosphate of GMP and water, as well as c-di-GMP and T263 from STING1 monomer. Hydrogen bonds between c-di-GMP and water are shown as yellow dashed lines and red spheres, respectively. π-π interaction of c-di-GMP and Y167 is shown as a green dashed line. Residues from STING1 monomer A that interact with c-di-GMP are shown as orange sticks. c-di-GMP is shown as yellow sticks.

Gate strategy_Brain

Fig. S6. Flow cytometry gating strategy for the identification of immune cells.

Brain cells were initially gated on forward scatter (FSC)-A vs side scatter (SSC)-A and then **(A)** gated as CD45$^+$ cells and identified as TCRβ$^+$CD8$^+$. **(B)** TCRβ$^+$CD8$^+$ gated cells were examined using cell surface markers for activation (CD44 and CD62L). **(C)** Monocytes and microglia, distinguished based on CD45 expression and Mac-1, were characterized as CD45^{high} Mac-1$^+$ and CD45^{int} Mac-1$^+$, respectively. **(D)** Within microglial cell population, cellular activation was examined based on MHC class I expression. **(E)** Activation of CD8$^+$ T cells was further studied using positivity for IFNγ and Granzyme B. **(F)** Analysis of cell activation profile in splenic CD8$^+$ and CD4$^+$ T cells and **(G)** monocytes (Mac-1$^+$) of experimental mice.