Pilotins are mobile T3SS components involved in assembly and substrate specificity of the bacterial type III secretion system

Stephan Wimmi*, Moritz Fleck*, Carlos Helbig, Corentin Brianceau, Katja Langenfeld, Witold G. Szymanski, Georgia Angelidou, Timo Glatter, Andreas Diepold

Max Planck Institute for Terrestrial Microbiology, Department of Ecophysiology,

Karl-von-Frisch-Str. 10, 35043 Marburg, Germany

Corresponding author: Andreas Diepold, andreas.diepold@mpi-marburg.mpg.de,

+49-6421-178302

*: These authors contributed equally to this work

Short title: Pilotins contribute to T3SS assembly and substrate specificity

Abstract

In animal pathogens, assembly of the type III secretion system injectisome requires the presence of so-called pilotins, small lipoproteins that assist the formation of the secretin ring in the outer membrane. Using a combination of functional assays, interaction studies, proteomics and live-cell microscopy, we determined the contribution of the pilotin to assembly, function and substrate selectivity of the T3SS and identified potential new downstream roles of pilotin proteins. In the absence of its pilotin SctG, *Yersinia enterocolitica* forms few, largely polar injectisome sorting platforms and needles. In line, the majority of export apparatus subcomplexes is mobile in these strains, suggesting the absence of fully assembled injectisomes. Remarkably, the export of late T3SS substrates including the virulence effectors is hardly affected in these bacteria, whereas early T3SS substrates, such as the needle subunits, are only exported to a very low degree. We found that pilotins transiently interact with the secretin and the large export apparatus component SctV, but mostly localize throughout the bacterial membrane, where they form transient mobile clusters, which do not colocalize with assembled injectisomes. In addition, pilotins interact with non-T3SS components, including sugar transporters. Pilotins therefore have additional functions downstream injectisome assembly, which include the regulation of type III secretion and a potential new link to the cellular energy metabolism.

Introduction

Bacteria that live in contact to eukaryotic cells greatly benefit from being able to manipulate those cells. The bacterial type III secretion system (T3SS) allows such manipulation by injecting effector proteins directly from the bacterial cytosol into the host cell cytoplasm [1–5]. Although the T3SS is also used by symbionts and commensals, it is best known for its essential role in infections of important human pathogens such as *Salmonella*, *Shigella* and *Yersinia*. In these pathogens, the T3SS machinery, also called injectisome[1], is often assembled upon entry into the host organism. Once the basal body (membrane rings and export apparatus, Fig. 1a) and the cytosolic components are assembled (reviewed in Diepold and Wagner, 2014), the T3SS starts to secrete its own distal components. Amongst those so-called early secretion substrates are the subunits of the needle and proteins required for its correct assembly, as well as early regulatory proteins. When the required needle length is reached [7,8], middle secretion substrates – the translocon proteins forming the needle tip and two hydrophobic interactors that build a pore in the host membrane – are secreted. At this point, the injectisome is ready for secretion of the late secretion substrates, the virulence effectors, which is initiated by host cell contact.

Injectisomes are assembled at the bacterial surface according to the needs of the bacteria in a given situation. While in some cases, one to few injectisomes are sufficient (such as for the intracellular *Salmonella* SPI-2 T3SS [9]), bacteria that need to actively make contact with the host cells usually build more injectisomes [10,11]. The distribution of injectisomes across the bacterial surface is non-random [12–14], but it is currently unclear what determines where new injectisomes are built.

A prime candidate to establish the site of assembly for new injectisomes is the pilotin, a lipoprotein anchored in the inner leaflet of the outer membrane (OM). Pilotins are required for the proper integration and oligomerization of the secretin protein, which forms a large channel in the OM in the T3SS of animal pathogens, but also type II secretion systems (T2SS) and type IV pili (T4P) [15–27]. Since the secretin ring is one of the nucleation structures of the injectisome [6,28] and anchors the T3SS at its final position in the peptidoglycan mesh [29] (Fig. 1a), the localization of the pilotin may determine the place of secretin ring formation and in consequence the distribution of injectisomes.

---

[1] The T3SS is at the core of both injectisomes and bacterial flagella [11,96]. In this manuscript, "T3SS" refers to the injectisome-type, virulence-associated system.

Pilotins and their cognate secretin proteins in the T2SS and T3SS have been purified and structurally

analyzed in 1:1 complexes [23,30,31]. As a lipoprotein, the pilotin also interacts with the localization

of lipoproteins (Lol) protein export machinery [31,32]. Based on these properties, different models

for the export of the pilotin itself and its role in secretin ring assembly have been proposed [30,31].

Whether the pilotin stays attached to the secretin after secretin ring formation in the OM and the

subsequent assembly of the injectisome is debated and might differ between organisms. For T2SS

secretins, such attachment was initially proposed, but later rejected [23,33]; recent analyses of

purified secretin rings found pilotins attached stoichiometrically outside the main ring [34,35]. *In*

situ structures of the T3SS showed electron densities around the secretin, possibly corresponding

to pilotin proteins, in *Shigella flexneri* [36], but not in *Salmonella enterica* [37]. In spite of the

importance of pilotins for the localization of the central secretin ring structure, protein secretion by

the T3SS was found to be strongly impaired, but not completely prevented in absence of the pilotin

in some organisms [15,19,25] – the structural and mechanistic basis for this phenotype is currently

unclear.

Despite their central role, many key characteristics of pilotins are poorly understood or debated:

Why are pilotins usually not encoded with all other structural components of the T3SS, but upstream

the main transcription factor of the system? What determines their localization (and eventually that

of the injectisome)? Do pilotins remain part of the injectisome after the assembly and do they adapt

a functional role in type III secretion at this point?

To shed light on these questions and the role of pilotins in assembly and function of the T3SS, we

studied the role of the *Yersinia enterocolitica* pilotin SctG[2] with functional assays, fluorescence

microscopy, proteomics, as well as protein interaction and dynamics studies. We found that,

unexpectedly, SctG interacts with the secretin SctC only temporarily, and mostly localizes in dynamic

patches that are dispersed throughout the bacterial membrane. In absence of the pilotin, only a

small subset of bacteria assemble single, mostly polar needles which are not stably attached to fully

assembled injectisomes, as indicated by the lack of colocalization with the cytosolic sorting platform

component SctL. To our surprise, presence of the pilotin not only influenced injectisome assembly

and overall secretion levels, but specifically altered the secretion pattern of the T3SS. While effector

proteins and other late substrates were efficiently secreted in the absence of the pilotin, the export

of early substrates, such as the needle components, was greatly impaired. Based on specific

---

[2] This manuscript uses the common nomenclature for T3SS components [97,98]; the species-specific nomenclature of the *Yersinia* pilotin is YscW; other names can be found in Table S1.

95 interaction studies, we provide possible explanations for this unique phenotype and an extended

96 model for the function of the pilotin.

Results

Deletion of the pilotin specifically reduces secretion of early export substrates

To define the overall role of the pilotin SctG in type III secretion, we tested the effects of a complete

sctG deletion on the virulence plasmid in *Y. enterocolitica* in a secretion assay. Absence of SctG

distinctly altered the secretion pattern. While secretion of effector proteins was partially reduced,

but still clearly present, early secretion substrates such as the needle subunit SctF, the ruler protein

SctP and translocators SctA, B, E were much more severely affected in secretion or not secreted at

all. *In trans* expression of SctG complemented the deletion (Fig. 1b, Fig. S1a).

In order to determine the role of SctG in secretion more precisely, we analyzed the secretome of

the wild-type and *ΔsctG* strains in minimal medium by label-free quantitative mass spectrometry.

The results show that T3SS substrates cluster according to their nature, rather than their overall

secretion level (Fig. 1c, Table 1, Fig. S1a): The late substrates, effector proteins YopH, O, P, E, M, T

as well as the gatekeeper SctW (YopN) and the negative regulators YscM1/2 (marked green in Fig.

1bc), were exported efficiently in absence of SctG. In clear contrast, export of all early and middle

substrates of the T3SS (marked magenta in Fig. 1bc) was strongly reduced in *ΔsctG*. These T3SS

substrates build up the needle (SctF) and the inner rod (SctI) connecting it to the export apparatus,

control needle length (SctP) and form its tip structure (SctA) as well as a pore in the host membrane

(SctB, E). YscX is an early T3SS substrate specific to the Ysc subgroup of T3SS [38,39], which is

required for the export of any other protein by the T3SS of the Ysc subgroup [40,41].

Since most pilotins, including SctG, are expressed in an operon with the main transcription factor of

the T3SS, VirF [42], we tested if the observed phenotype is caused by impaired VirF expression in

the absence of *sctG*. Expression of VirF from plasmid in a *ΔsctG* strain indeed slightly increased the

secretion of effectors, but not of early substrates (Fig. S1b), indicating that the specific loss of

secretion of these early substrates is not caused by an influence on VirF expression.

Fig. 1 – SctG is required for the secretion of early export substrates, but not of effectors.

a) Schematic illustration of the injectisome (adapted from Diepold and Wagner, 2014). Main substructures indicated on the left; components analyzed in this study, including the pilotin SctG and the secretin SctC, are indicated on the right and shown in color. **b**) Secretion assay showing the export of native T3SS substrates into the culture supernatant in the listed strains (*n*=3). Left, molecular weight in kDa; right, assignment of exported proteins. **c**) Label-free mass spectrometry quantification of secreted proteins in *ΔsctG* and wild-type (WT), intensity ratio plotted against mean intensity (*n*=3). In b) and c), early/middle T3SS export substrates are marked in magenta; late substrates (effectors) are marked in green; substrate with unknown time point of export marked in yellow.

135   **Table 1 – Influence of SctG on T3SS secretion substrates**

136   Label-free quantitative mass spectrometry of known T3SS substrates in the culture supernatant of

137   different strains, as indicated. Magenta shade, early and middle substrates; green shade, late substrates;

138   yellow shade, unknown. # pept., number of total detected peptides. Hits sorted by ratio of

139   *ΔsctG*/wild-type (WT). *n*=3, individual measurements listed in Table S2.

T3SS substrate	# pept.	Intensity ratio *ΔsctG* / WT	Mean intensity		
			WT	*ΔsctG*	*ΔsctC*
Needle component SctF	7	0.0547	3.95E+09	2.16E+08	1.65E+07
Early substrate / regulator YscX	4	0.1366	3.55E+08	4.85E+07	8.80E+06
Tip protein SctA (LcrV)	24	0.1580	8.72E+09	1.38E+09	1.07E+08
Rod / washer protein SctI	3	0.1621	3.56E+08	5.78E+07	2.07E+07
Needle length regulator / ruler SctP	23	0.2359	1.53E+09	3.60E+08	4.84E+07
Translocator SctE (YopB)	19	0.3213	2.73E+09	8.77E+08	3.44E+08
Translocator SctB (YopD)	46	0.4502	3.26E+10	1.47E+10	9.06E+08
Secreted protein YscH/YopR	4	0.5604	2.90E+08	1.62E+08	1.15E+07
Effector YopQ	16	0.6357	1.90E+09	1.21E+09	6.51E+06
Effector YopO	37	0.9833	1.72E+09	1.69E+09	6.58E+07
Effector YopH	64	1.1095	3.11E+10	3.45E+10	1.23E+09
Effector YopE	10	1.5746	1.16E+10	1.82E+10	2.67E+09
Effector YopP	25	2.1810	7.21E+08	1.57E+09	5.24E+06
Gatekeeper SctW (YopN)	17	2.5811	3.65E+09	9.42E+09	3.18E+07
Effector YopM	6	2.6303	4.81E+09	1.26E+10	7.99E+07
Effector YopT	7	4.1605	1.15E+07	4.77E+07	1.28E+07
Regulator YscM1	12	5.5968	4.84E+08	2.71E+09	9.81E+06
Regulator YscM2	7	10.2833	4.52E+07	4.65E+08	5.43E+06

The pilotin is essential for stable localization of the export apparatus and the cytosolic

T3SS components, but few needles can form in its absence

We next investigated the specific role of the pilotin in the assembly of the injectisome, which may

provide a clue to its function. To this aim, we deleted *sctG* in strains expressing fluorescently labeled

variants of the secretin (SctC-mCherry) [28], the large export apparatus component (SctV-EGFP)

[29], the cytosolic sorting platform component SctL (mCherry-SctL) [43], and a strain expressing

SctF$_{S5C}$, a variant of the needle protein that can be labeled by maleimide dyes [44]. Except for SctF$_{S5C}$,

which is expressed from plasmid, all other labeled proteins are expressed from their native genetic

environment. As expected, fluorescence microscopy of the respective bacteria showed an

important role of SctG in the assembly of the secretin ring. Far from its usual assembly in >10 mostly

lateral foci in the wild-type background, SctC mostly localized in large, often polar clusters in the

absence of the pilotin (Fig. 2a, left, Dataset S1). These SctC foci mostly do not correspond to fully

assembled injectisomes: SctV foci were motile in the membrane in the absence of SctG, like in the

absence of the secretin SctC (Fig. 2b, Fig. S2a) [29]. Moreover, we found that the cytosolic

component SctL formed very few foci in the absence of SctG (Fig. 2, center; Dataset S1).

Nevertheless, a small subset of bacteria lacking SctG did display needles (Fig. 2, right; Dataset S1).

These needles, however, did not colocalize with the sorting platform protein SctL (Fig. S2b). *In trans*

complementation of SctG restored wild-type localization of all tested components (Fig. S3). Notably,

levels of SctG significantly lower than in the wild-type were already sufficient for complementation

of secretion and injectisome assembly (Fig. S3).

The results above indicate that SctG is essential for proper assembly and anchoring of all major

subcomplexes of the injectisome, but that a subset of cells forms mostly polar needles in its absence.

We therefore next analyzed SctG itself to figure out how it conveys its function.

Fig. 2 – SctG is required for normal localization of SctC and stable assembly of the export apparatus and the sorting platform component SctL, but not fully essential for the formation of needles.

a) Fluorescence micrographs of *Y. enterocolitica* expressing the proteins indicated from their native location on the virulence plasmid (SctC-mCherry and mCherry-SctL) or *in trans* (SctF$_{S5C}$, stained with CF 488A maleimide stain) in an effector-less strain background. Top row displays native localization in wild-type (WT) background, bottom row localization in respective Δ*sctG* strain background. Scale bar, 2 µm. The fraction of bacteria with foci is listed for the respective Δ*sctG* strains. See Suppl. Dataset 1 for details. *n*=3-6. **b**) Red/green/blue (RGB) overlay and minimum intensity projection showing the movement of SctV-EGFP over time in the indicated effector-less strain backgrounds in live *Y. enterocolitica*. Red/green/blue channel: micrograph at t=0/15/30 s; white foci indicate stable protein localization, whereas colors indicate movement of foci over time. Minimum projection: micrographs at 0/15/30/45/60/75/90 s; foci indicate stable foci over this time period. *n*=3.

Pilotin expression is controlled by temperature, similarly to structural T3SS components

Pilotin proteins are abundant but not omnipresent in secretion systems. Despite their consistent reported role for assembly of the highly conserved secretin, pilotins show low sequence conservation (Fig. S4a). Accordingly, we found that even the pilotin from the closely related T3SS of *P. aeruginosa* could not complement a *sctG* deletion in *Y. enterocolitica* (Fig. S4b).

Despite this low sequence conservation, pilotins have a conserved unique genetic location within the virulence plasmids or genetic islands encoding the T3SS. While most other structural and regulatory components of the injectisome are under the control of a main transcriptional regulator (VirF in *Y. enterocolitica*, called LcrF in other *Yersinia* species), pilotins are located upstream of the transcriptional regulator itself (genetic neighborhood analysis, Fig. 3a), indicating a specific role in the T3SS with a potentially distinct expression pattern.

To unravel the expression kinetics, as well as the subcellular localization of SctG, we introduced a *sctG-sfGFP* fusion in the native *sctG* locus on the *Y. enterocolitica* virulence plasmid by homologous recombination [45]. The SctG-sfGFP fusion protein was stable and functional (Fig. S5). The expression of SctG, VirF and, in consequence, other T3SS components was moderately increased (1.5-2-fold) in the strain. Additionally, few non-T3SS proteins were affected by the fusions (Table S3).

SctG-sfGFP was expressed at a level below the detection limit prior to the shift to 37°C, which activates the VirF-dependent expression of other T3SS components. After the temperature shift, SctG expression increased steadily over the next two hours (Fig. 3bc). Somewhat surprisingly, given their different genetic regulation, the SctG-sfGFP expression profile did not differ significantly from mCherry-SctD, a basal body protein labeled in the same strain (Fig. 3c).

Fig. 3 – Despite its unique genetic location, SctG expression is regulated similar to other T3SS components.

a) Genetic neighborhood analysis of pilotin proteins (green) of the different T3SS indicated on the left side. Main T3SS transcription factors (orange), secretins (blue) and other T3SS components (gray) are indicated. The *ysrRS* phosphorelay system activating transcription of the Ysa system [46] is marked in yellow. **b**) Expression of SctG-sfGFP (expected molecular weight 42.2 kDa) over time under non-secreting conditions. Immunoblot using antibodies directed against GFP. Control: Expression of GFP from plasmid. Numbers indicate time in min relative to temperature shift to 37°C, which triggers the expression of T3SS components. Right, molecular weight in kDa. *n*=2. **c**) Expression of SctG-sfGFP and the IM ring component mCherry-SctD over time before and after temperature shift to 37°C in a double-labeled strain under non-secreting conditions. Quantification of fluorescence per bacterium, normalized by the respective 120 min value. Dots indicate individual biological replicates with a total of 52, 100, 128, 128, 217, 185, 143 bacteria (from left to right).

Pilotins localize dynamically within the membrane and do not stably colocalize with other injectisome components

The functional SctG-sfGFP fusion allowed us to assay the cellular localization of SctG in live *Y. enterocolitica*. Stable stoichiometric association with the secretin SctC would result in the formation of membrane foci colocalizing with other injectisome components [28,43]. However, we detected a disperse membrane localization of SctG-sfGFP which increased over time similar to overall protein levels in the cell, in addition to cytosolic background fluorescence (Fig. 4a). While a certain clustering of SctG-sfGFP in the membrane was visible, these clusters did not significantly colocalize with the mCherry-labeled T3SS components SctC, SctD, or SctL, in contrast to the clear colocalization observed for mCherry-SctD and EGFP-SctQ used as positive control (Fig. 4b, Fig. S6).

Fig. 4 – SctG predominantly localizes in the membrane, but does not significantly colocalize with other injectisome components.

a) Time course microscopy of SctG-sfGFP in *Y. enterocolitica* under non-secreting conditions. Time with reference to the temperature shift to 37°C. Bottom, intracellular localization of SctG-sfGFP at the respective time points. Line scans across single bacteria, approximate average positions of bacterial membranes (m., dark grey) and cytosol (beige) indicated. *n*=78, 90, 81, 90 cells (left to right) from three independent biological replicates. **b**) Colocalization of labeled T3SS components (all expressed from native genetic locus on virulence plasmid) in double-labeled strains as indicated, 120 min after temperature shift to 37°C. *n*=3, except for colocalization of mCherry-SctD and EGFP-SctQ, which was reported earlier [43] and included here as a positive control (*n*=1). Scale bars = 2 μm.

To analyze the mobility and cluster formation of SctG in more detail, we tracked SctG-sfGFP in

individual bacteria. Still images and overlays (Fig. 5a), as well as movies (Suppl. Movie 1) showed

that the localization of SctG and its clusters in the membrane strongly fluctuates, indicating diffusion

within the membrane. We tested this hypothesis by fluorescence recovery after photobleaching

(FRAP) and found that indeed, SctG-sfGFP is highly mobile within the membrane with an average

half time of recovery at the bleached pole of 21.5 s (+/- 13.0s) (Fig. 5b).

Fig. 5 – Pilotin proteins move dynamically within the membrane, forming transient patches.

Fluorescence microscopy of a strain expressing SctG-sfGFP and mCherry-SctD from their native genetic

locations on the virulence plasmid. **a**) Left, single micrographs of SctG-sfGFP taken with 1 s difference.

Right, red/green/blue (RGB) overlay of those micrographs for SctG-sfGFP and mCherry-SctD (mCh-SctD)

(*n*=3). **b**) Fluorescence recovery after photobleaching for SctG-sfGFP. Left, representative individual

micrographs before and at the indicated time points after the bleach event. Bleach spot indicated by red

circle. Right, quantification of fluorescence recovery after photobleaching (*n*=21 cells from 3

independent experiments, displayed in different shades of grey). The recovery half-time $t_{1/2}$ was

calculated for each individual recovery curve (see Material and Methods for details) and is displayed as

average with standard deviation.

To investigate if SctG localization requires any other T3SS component or non-T3SS-related *Y. enterocolitica* protein, SctG or SctG-sfGFP were expressed from plasmid in a *Y. enterocolitica* strain cured of the virulence plasmid (pYV⁻), as well as *Escherichia coli* Top10 (which has no T3SS) (Fig. S7a). While SctG-sfGFP displayed a membrane localization in *Y. enterocolitica* pYV⁻, similar to its native localization, it localized predominantly to the cytosol in *E. coli* (Fig. S7ab). Notably, expression of SctG negatively influenced the growth of *E. coli*. Strains expressing SctG or SctG-sfGFP only grew to a low OD and microscopy experiments revealed severe lesions and plasmolysis in the respective bacteria (Fig. S7ac), indicating that the pilotin localizes independently of other T3SS components, but can be toxic upon ectopic expression. Taken together, SctG follows a similar expression pattern as other VirF-controlled T3SS components despite its unique genetic location. It localizes predominantly throughout the membrane, independently from other T3SS components, and forms small transient patches.

Proteome analysis and specific interactions of the pilotin to T3SS components and proteins not directly linked to the T3SS

The unexpected localization of SctG and the finding that amounts below the wild-type level are sufficient for its function in injectisome assembly (Fig. S3) prompted us to consider additional roles of SctG. To identify such roles in an unbiased way, we first compared the *Y. enterocolitica* proteome in presence and absence of SctG under non-secreting conditions. In *ΔsctG*, the amount of the main transcriptional activator VirF was reduced to about one third. Consequently, most other T3SS components were also present at lower concentrations (Table S4). Absence of SctG had little impact beyond the T3SS, with the notable exception of the OM protein OmpW, whose amount was reduced by a factor of eight (Table S4).

To better understand the function of the pilotin, both with respect to the T3SS and other pathways, we screened for interaction partners by co-immunoprecipitation with SctG-sfGFP, analyzed by label-free mass spectrometry. As expected, SctG interacted with the secretin SctC, but also with other T3SS components, most prominently the effector YopM and the large export apparatus protein SctV (Fig. 6). Most other T3SS components were enriched to a lower degree (Fig. 6, Table S5).

Interestingly, a number of proteins not directly linked to the T3SS were enriched highly significantly. Most prominently, the phosphoenolpyruvate (PEP)-dependent phosphotransferase system (PTS) trehalose transporter subunit EIIBC (PTS-IIBC) consistently interacted very specifically with SctG (Fig. 6), despite low overall expression levels (Table S4). A closer look at the results also showed a significant enrichment of several components of the Lol protein export pathway for the insertion of lipoproteins into the OM, namely LolD and LolE, as well as other non-T3SS proteins, such as Lon protease (termed endopeptidase La in *Y. enterocolitica*) and the prokaryotic tubulin homolog FtsZ, a major player in cell division (Fig. 6, Table S5).

Fig. 6 – SctG interacts with T3SS components and additional non-T3SS proteins.

Co-immunoprecipitation experiments with SctG-sfGFP as bait, compared to an unlabeled control strain expressing sfGFP from plasmid at comparable levels. Interactors were quantified with label-free mass spectrometry (*n*=3). The bait protein SctG is marked in orange, other T3SS components in dark red, interactors discussed in the main text in green. Greek letters denote subunits of the F1 subcomplex of the ATP synthase. A full list of all interactors with *p*<0.001 and T3SS components can be found in Table S5.

Interestingly, PTS sugar transporters were shown to have a role in the regulation of the T3SS earlier [47]. We therefore constructed a clean *Δpts-IIBC* deletion mutant and tested its impact on assembly and function of the T3SS. PTS-IIBC was not required for assembly of injectisomes or effector secretion (Fig. 7a) and overexpression did not significantly impact T3SS localization or secretion itself (Fig. 7b).

Fig. 7 – The SctG interactor PTS trehalose transporter subunit EIIBC does not affect assembly or function of the T3SS in rich medium.

a) Left: Micrographs of SctG-sfGFP and mCherry-SctL in double-labeled *Y. enterocolitica* strains otherwise wild-type (WT) or with a deletion of the PTS trehalose transporter subunit EIIBC (Δ*pts*). The experiment was performed under non-secreting conditions (*n*=3). Right: Secretion assay of the PTS trehalose transporter subunit EIIBC (Δ*pts*) deletion with WT and Δ*sctD* strains serving as positive and negative controls, under secreting conditions (*n*=2), molecular weight in kDa indicated. **b)** Left: Micrographs of EGFP-SctQ expressed from its native location on the virulence plasmid in *Y. enterocolitica* otherwise WT or overexpressing the PTS trehalose transporter subunit EIIBC (+PTS). Experiments performed under non-secreting and secreting conditions as indicated (*n*=3 each). Right: corresponding secretion assay, molecular weight in kDa indicated. Scale bar for all micrographs, 2 µm.

Discussion

Pilotin proteins are conserved components of T3SS in many bacteria including most animal pathogenic variants. They have a unique genetic location and a central role for the assembly of the T3SS injectisome. However, little was known about their behavior and role beyond the assistance in the proper localization and oligomerization of the secretin ring of the injectisome in the OM. In this study, we confirm this role, but uncover an additional effect on T3SS export substrate selection. Unexpectedly, we found that pilotins are mobile membrane components which may link the T3SS to other cellular pathways, including sugar metabolism.

Pilotins differentially affect major steps in T3SS assembly

Pilotins have long been known to assist the assembly of the SctC secretin ring in the OM [18–22,24,25]. Our data confirm these reports: In the absence of the pilotin, secretins localized in large foci at one of the bacterial poles (Fig. 2a). The localization of other T3SS components in absence of SctG differed from that of the secretin: SctL, a component of the cytosolic sorting platform of the T3SS, only formed single, mostly polar, foci. Similarly, we observed few, also mostly polar, needles for a subset of bacteria. However, these needles and the sorting platform protein SctL did not colocalize (Fig. 2a, Fig. S2b), indicating an absence of stably assembled injectisomes. Consistent this, the nonameric large export apparatus component SctV [48] assembled, but moved within the membrane in the absence of SctG (Fig. 2b, Fig. S2a), indicating a lack of anchoring by the ring-forming SctCDJ proteins [29]. Since all cytosolic T3SS components, including SctL, are required for protein export under normal conditions, we speculate that transient interactions of SctL and the other cytosolic T3SS components with basal body components at the poles might allow the limited export of the needle subunit, which would then stay stably associated with the bacteria despite the disassembly of the cytosolic sorting platform. In agreement with this hypothesis, we had found earlier that the cytosolic T3SS sorting platform components do exchange between the injectisome-bound sorting platform and a cytosolic pool [43,49] and that the sorting platform components temporarily dissociate from the remaining injectisomes under certain conditions, such as low external pH [50]. The presence of a low number of needles had previously been reported for the closely related *P. aeruginosa* T3SS [19]. Notably, the fraction of bacteria with associated needles and the overall secretion efficiency of the pilotin deletion showed an unusually high variation between individual replication experiments, suggesting an influence of external factors. However, despite our best efforts, we were unable to identify these factors.

Role of pilotins in T3SS substrate specificity

Analysis of the T3SS secretome by SDS-PAGE and quantitative mass spectrometry revealed a specific

effect of the presence of the pilotin on export of the so-called early substrates, required for

formation of the needle and its tip structure (Fig. 1, Table 1). Notably, while deletion or

modifications of *sctG* have an effect on the expression of VirF and its targets (Table S3, 4), most

likely due to an influence on the formation of hairpin structures in the *sctG-virF* mRNA that regulate

VirF transcription as part of the thermoregulation of T3SS expression [42,51], the role of SctG in

secretion substrate is independent from VirF levels (Fig. S1b). What could then be the reason for

the striking change of substrate specificity in absence of the pilotin, where early export substrate

are much more strongly affected by the absence of SctG than the late substrates, the virulence

effectors? Similar phenotypes had been found, although not always explicitly noted, in earlier

studies in *Y. enterocolitica* [15,25] and *P. aeruginosa* [19]. Taking into account the limited, probably

transient interaction of the pilotin with the secretin that we observed, it is tempting to speculate

that presence of the pilotin at the injectisome favors the secretion of early substrates. Subsequent

dissociation of the pilotin would then contribute to a switch in export cargo specificity towards the

effectors. The specific (albeit not necessarily direct) interaction of SctG with the large export

apparatus component SctV (Fig. 6), which is a key player in substrate selection [52–54], supports

this hypothesis.

Alternative explanations for the change in export substrate specificity might be (i) the polar

localization of the secretin (and consequently the hypothetical transient T3SS), which might change

the accessibility for certain substrates, or (ii) a two-step export pathway, with a first transport step

across the IM, facilitated by the T3SS export apparatus (which is assembled, albeit not anchored in

the peptidoglycan, in the absence of the pilotin (Fig. 2b)) and a subsequent export step across the

OM, facilitated by the secretin (without which no export is observed (Fig. S1)). The two-step

transport might prevent the export of early substrates, many of which polymerize outside the

cytosol. Alternatively, (iii) strains lacking the pilotin might be able to provide less energy due to the

missing link to the PTS sugar transporters or membrane stress as a consequence of secretin

misassembly. If effector export is less energy costly than the export of early substrates, this might

explain the observed bias in export in absence of SctG. However, in the absence of direct evidence

for these alternative explanations, we consider a direct influence of SctG on the substrate specificity

through interactions with other T3SS components to be the most likely explanation.

Substrate specificity switching is a central, but incompletely understood, feature type III secretion.
Our data suggest that pilotin proteins play a key role in this process, together with the export
apparatus components SctV and SctU [55], the gatekeeper complex [56,57] and the sorting platform
[58]. If and how the interaction of the pilotin with the effector YopM (Fig. 6) contributes to this
process is unclear at the moment.

Additional interactions of the pilotin after its dissociation from the injectisome and
possible functions

An unexpected finding of our study was that pilotin concentrations significantly below wild-type
levels are sufficient to restore normal T3SS assembly and function (Fig. S3). Although it is one of the
few T3SS machinery components encoded outside the VirF-regulated operons, native SctG
expression develops very similarly to other T3SS components (Fig. 3), consistent with the co-
transcription of *sctG* and *virF* in one operon [42]. However, the findings that low pilotin levels
already were sufficient for injectisome assembly and secretion and that SctG does not significantly
colocalize with other T3SS components (Fig. 4), but is highly mobile in the bacterial membrane (Fig.
5) suggests that the pilotin might have additional functions beyond its role in secretin assembly,
possibly after its release from the secretin. We therefore screened for additional interactors of SctG
by co-immunoprecipitation. Notably, while the pilotin mostly fractionates with the OM in
membrane preparations [18,25,59], many of the proteins found to interact with the labeled pilotin
in the co-immunoprecipitation assay are integral IM proteins or IM-associated proteins. The used
SctG-sfGFP fusion is expressed from its native genetic location, supports native secretion (Fig. S5),
and interacts with both known interactors, the secretin SctC and proteins of the Lol pathway, in our
assays (Fig. 6). Despite these indications for correct localization and interactions, it is possible that
transport of the fusion protein to the OM is impaired and that this at least enhances the fraction of
time the pilotin can contact these IM(-associated) proteins. Alternatively, the interactions may be
caused by natively soluble pilotin proteins, possibly bound to other proteins such as LolB [30,31].
Finally, the N-terminal lipid anchor of the pilotin might be removed after assembly of the secretin
ring or the injectisome, which would render the protein soluble [25] in the periplasm.

The most striking finding of the interaction studies was the highly specific enrichment of several PTS
sugar transporter subunits, most significantly the membrane-bound trehalose transporter subunit
EIIBC (Fig. 6, Table S5). PTS systems are used for sugar uptake in many bacteria. They create a
concentration gradient by transferring the PEP phosphoryl group onto the incoming sugar via

several intermediate phosphate carriers. The EIIC subunit, which can be fused to the EIIB subunit,

serves as channel in the IM. While many bacteria express several PTS transporters, PTS are often

unspecific and redundant [60]. Notably, PTS were repeatedly found to act as carbohydrate sensors

that control bacterial metabolism in response to the environment [61]. Transporters sensing host-

specific compounds can regulate expression and activity of virulence systems [62], indicating that

SctG and its main interactor, the PTS trehalose transporter subunit EIIBC, might serve to modulate

T3SS function according to the bacterial environment. The fact that so far, no pilotin homologs were

found in plant pathogens [3] is consistent with this hypothesis. Indeed, PTS systems were repeatedly

associated with bacterial virulence (reviewed in [60,63]): amongst others, presence of the PTS

glucose transporter subunit EIIBC conferred a competitive advantage to *Y. pestis* in serum-like

medium [64], a *Vibrio cholerae* PTS system was found to modulate virulence gene expression [65],

and *Salmonella* Typhimurium lacking a PTS EI subunit showed a nearly 1000-fold increase in LD_{50}

[66]. Specifically a PTS trehalose transporter system induces hypervirulence in *Clostridium difficile*

[67]. Notably, the PTS glucose transporter EIIA subunit directly interacts with cytosolic components

of the *Salmonella* SPI-2 T3SS and activates type III secretion [47]. Overall, the two most likely links

between PTS sugar transporters and virulence appear to be energy provision and/or host sensing.

Both possibilities are consistent with the lack of a phenotype of the *pts-IIbc* deletion in our

experiments (which were performed in rich medium in the absence of host cells, Fig. 7); however,

further experiments are required to describe the link in more detail.

Other significant interaction partners of the pilotin outside the T3SS are FtsZ, a prokaryotic tubulin

homolog with a key role in cell division, the protease Lon, components of the ATP synthase (subunits

α, β, γ), the Sec export pathway (SecF/A), and LolD/E, two components of the ABC transporter of

the Lol lipoprotein export pathway in the IM. FtsZ is an interesting interactor, as it might play a role

in the long-known, but still incompletely understood link between T3SS activity and cell growth and

division (actively secreting bacteria of many species cease growth and division and are often bigger

than T3SS-deficient or non-secreting bacteria [44,68–70]). Since to our knowledge, this is the first

time that a T3SS component was found to be associated with bacterial cytoskeleton proteins, this

might indicate an involvement of SctG in the non-random localization of T3SS around the cell

[12,13]. The Lon protease degrades unfolded or misfolded proteins, especially under stress

conditions [71], but can also have more specific roles in metabolism, cell cycle control and virulence

through proteolysis of specific regulatory proteins [72,73]. While it cannot be ruled out that the

specific interaction of SctG-sfGFP with Lon denotes degradation of partially misfolded SctG-sfGFP,

Lon proteases have been shown to influence T3SS activity, both positively via degradation of

histone-like proteins like *Yersinia* YmoA that suppress the transcription of T3SS components [74–

77] and negatively, most likely by directly degrading key components of the T3SS, mainly in plant

pathogens [78–81]. The presence of three components of the IM ATP synthase amongst the less

than forty specific interactors of the pilotin protein is remarkable and, in combination with the

strong interaction with the PTS sugar transporter, suggests a link with the bacterial energy

metabolism. While such a link is more than conceivable, given the high energy expense of type III

secretion [82], its details are unclear at the moment. As lipoproteins in the OM, pilotins are expected

to interact with components of both the Sec and the Lol pathway [30,32,83]. While most models

emphasize the interaction with LolA and LolB in the periplasm and OM, respectively, we detected

strong binding to LolD and LolE in the IM (Fig. 6, Table S5), but no significant binding to LolA or B. Of

the two models proposed for the role of pilotins in the assembly of the secretin ring by Majewski

and colleagues [31], our data therefore supports the second model where the pilotin is released

from LolB before interacting with the secretin.

Taken together, our data indicate that, besides their function in the assembly of the secretin ring of

the T3SS, pilotin proteins have additional roles linked to type III secretion. In their absence, assembly

of all T3SS subcomplexes and specifically the export of early secretion substrates is strongly

impeded, while effectors are exported efficiently. Pilotin protein interact with components of the

T3SS, most prominently the secretin SctC and the large export apparatus component SctV, but are

mobile in the bacterial membrane and do not stay attached to or even significantly colocalize with

the injectisome. In line with these findings, the pilotin also interacted with other cell wall-associated

proteins such as a PTS sugar transporter subunit.

A potential model integrating these findings is that during injectisome assembly, SctG interacts with

SctC, facilitates its oligomerization in the OM, and the formation of needles. At a later time point,

SctG dissociates from the fully assembled injectisome, which then favors the export of effectors

upon activation of the T3SS. The pilotin is mobile at this stage and may additionally support this

function by interacting with other partners, including PTS sugar transporters, which could provide

the necessary energy for secretion. Challenging this model in future studies may therefore lead to a

better understanding of the energy provision for the T3SS and its link to sugar import.

Material and methods

Bacterial strain generation and genetic constructs

A list of strains and plasmids used in this study can be found in Supplementary Table 6. All *Y. enterocolitica* strains used in this study are based on the *Y. enterocolitica* wild-type strain MRS40 or the strain IML421asd (ΔHOPEMTasd). In IML421asd all major virulence effector proteins (YopH,O,P,E,M,T) are absent. Furthermore, this strain harbors a deletion of the aspartate-beta-semialdehyde dehydrogenase gene which renders the strain auxotrophic for diaminopimelic acid (DAP), making it suitable for work in an biosafety class 1 environment [84].

Bacterial cultivation, secretion assays and protein analysis

Y. enterocolitica day cultures were inoculated from fresh stationary overnight cultures supplemented with the relevant antibiotics and DAP (80 mg/ml), where required, to an OD_{600} of 0.15 for secreting and 0.12 for non-secreting conditions, respectively. As growth medium, BHI (brain heart infusion broth) supplemented with nalidixic acid (35 mg/ml), DAP (80 mg/ml) where required, glycerol (0.4%) and $MgCl_2$ (20 mM) was used. Where required, ampicillin was added (0.2 mg/ml) to select for the maintenance of expression plasmids. For secreting conditions, 5 mM EGTA were added; for non-secreting conditions, 5 mM $CaCl_2$ were added to the pre-warmed (approx. 55°C) medium, which was filtered through a 0.45 μm filter before the addition of other supplements and cooled down to ambient temperature for the experiments. After inoculation, day cultures were incubated at 28°C for 90 min to reach exponential growth phase. Expression of the *yop* regulon was then induced by a rapid temperature shift to 37°C in a water bath. Where indicated, protein expression from plasmid (derivates of pBAD-His/B (Invitrogen) in all cases except expression of VirF from pCL5 [85]) was induced at this point by the addition of 0.02–1.0% L-arabinose as indicated in the respective experiments.

For protein secretion assays and analysis of total cellular proteins, bacteria were incubated for 2-3 h at 37°C, as indicated. 2 ml of the culture were collected at 21,000 *g* for 10 min. The supernatant was removed from the total cell pellet and proteins were precipitated by addition of a final concentration of 10% trichloroacetic acid (TCA) and incubation at 4°C for 1–8 h. Precipitated proteins were collected by centrifugation for 15 min at 21,000 *g* and 4°C, the pellet was washed once with 1 ml ice cold acetone and subsequently resuspended and normalized in SDS-PAGE loading buffer. Total cellular protein samples were normalized to 0.3 OD units (ODu, 1 ODu is equivalent to

1 ml of culture at on OD_{600} of 1, corresponding to approximately 5 x 10^8 *Y. enterocolitica*) and

supernatant samples to 0.6 ODu. Before loading, samples were incubated for 5 min at 99°C.

Separation was performed on 11%, 15% or gradient 12–20% SDS-PAGE gels, using BlueClassic

Prestained Marker (Jena Biosciences) as a size standard. For visualization, the SDS-PAGE gels were

stained with InstantBlue (Expedeon). For immunoblots, the separated proteins were transferred

from the SDS-PAGE gel onto a nitrocellulose membrane. Primary mouse antibodies against GFP

(1:4000) (ThermoFisher Proteintech 66002-1-Ig, 1:4000) were used in combination with secondary

anti-mouse antibodies conjugated to horseradish peroxidase (Sigma A8275, 1:5000). For

visualization, ECL chemiluminescence substrate (Pierce) was used in a LAS-4000 Luminescence

Image Analyzer.

Fluorescence microscopy

For fluorescence microscopy, bacteria were treated as described above. After 2–3 h at 37°C, 400 µl

of bacterial culture were collected by centrifugation (2400 × g, 2 min) and reconstituted in 200 µl

minimal medium (100 mM 2-[4-(2-Hydroxyethyl)piperazin-1-yl]ethane-1-sulfonic acid (HEPES) pH

7.2, 5 mM $(NH_4)_2SO_4$, 100 mM NaCl, 20 mM sodium glutamate, 10 mM $MgCl_2$, 5 mM K_2SO_4, 50 mM

2-(N-morpholino)ethanesulfonic acid (MES), 50 mM glycine). 2 µl of resuspension were spotted on

agarose pads (1.5% low melting agarose (Sigma-Aldrich) in minimal medium) in glass depression

slides (Marienfeld). Where required, agarose pads and media were supplemented with 80 mg/ml

DAP for ΔHOPEMTasd-based strains, 5 mM $CaCl_2$ for non-secreting conditions, or 5 mM EGTA for

secreting conditions.

Microscopy was performed on a Deltavision Elite Optical Sectioning Microscope equipped with an

UPlanSApo 100×/1.40 oil objective (Olympus), using an Evolve EMCCD camera (Photometrics).

Where applicable, the prepared samples were illuminated first in the mCherry = 0.4 s with a mCherry

filter set and afterwards in GFP = 0.2 sec with a GFP filter set. Depending on the experiment, z stacks

with 1-15 slices (Δz = 0.15 µm) per fluorescence channel were acquired. The acquired micrographs

where subsequently deconvolved using softWoRx 7.0.0 (standard "conservative" settings). Images

were further processed with FIJI (ImageJ 1.51f/1.52i/1.52n) [86]. Where necessary, a drift correction

with the StackReg Plugin [87] was performed. Fluorescence quantification was performed in FIJI. For

presentation of micrographs in a figure, representative fields of view were selected, and brightness

and contrast of the micrographs was adjusted identically within the compared image sets.

Maleimide-based needle (SctF) staining.

Maleimide-based staining to visualize the injectisome needles was performed as described in [50].

Briefly, expression of SctF$_{S5C}$ from the pBAD-His/B plasmid was induced by addition of 1.0% L-

arabinose at the time of the temperature shift in strains expressing the native SctF at the same time.

After 2-3 h, bacteria were collected and resuspended in 0.2 volumes of minimal medium

supplemented with 5 µM of CF 488A maleimide dye (Sigma-Aldrich, USA) at 37 °C for 5 min. After

staining, cells were washed once with 1500 µl of minimal medium and resuspended in 200 µl of

minimal medium. 2 µl of bacterial suspension were spotted on 1.5% agarose pads in minimal

medium and visualized under the microscope as described above.

Classification and quantification of fluorescent foci

To determine the number of stable YscV-EGFP foci, seven micrographs were taken in 15 s intervals

and the minimal intensity for each location was determined from the deconvoluted micrographs.

Foci were then identified in FIJI using an intensity threshold of 700 (determined in images of the

wild-type background strain) and the "analyze particles" function with a minimal size of 4 pixels and

a particle circularity of 0.5-1.0. To evaluate the fraction of bacteria with foci for SctC-mCherry,

mCherry-SctL and SctF$_{S5C}$ in the absence of SctG, foci were manually identified and classified as

polar, subpolar and lateral based on their subcellular localization in deconvoluted micrographs. In

case a cell displayed multiple foci in different localization classes, bacteria were classified as

containing a lateral, subpolar, polar focus in descending order (e.g. a bacterium with a lateral and a

polar focus was assigned "lateral"). The overall number of bacteria in a field of view was determined

by using the deep learning based network, DeepBacs [88], that was previously trained with

differential interferometry contrast (DIC) micrographs of *Y. enterocolitica*. The overall percentage

of cells with foci is displayed in the figure.

Fluorescence distribution analysis

For analysis of the fluorescence distribution across bacteria, line scan analysis was performed on

deconvoluted micrographs in FIJI. With the straight line freehand tool, measurements areas were

defined from one lateral membrane to the other at a 90° angle. Per condition, 3 fields of view with

18-30 cells were analyzed and the background was measured. The line scans were individually

corrected for background fluorescence, normalized (fluorescence fraction relative to overall

fluorescence over background) and centered at the position closest to center of the total

fluorescence over background.

Fluorescence recovery after photobleaching (FRAP)

Cells were grown in non-secreting medium as described above. Samples were taken after 3 h at

37°C. To minimize unnecessary bleaching, no z stacks were acquired. Three pre-bleach images were

acquired and locations near selected bacterial cell poles were photobleached by individual 30 ns

pulses of a 488 nm laser. Recovery of the bleached foci was followed by time-lapse microscopy.

Deconvolved micrographs were used for the fluorescence quantification.

The relative fluorescence of the bleached spot was calculated as follows:

$$\text{Relative flourescence} = \frac{(Intensity\ bleached\ cell\ pole - Intensity\ background)}{(Intensity\ total\ cell - Intensity\ background)}$$

The relative fluorescence was then normalized (average pre-bleach value set to 1; post-bleach value

set to 0) and an exponential recovery was fitted using the equation:

$$y = A(1 - e^{-bx})$$

where y stands for the relative fluorescence, x for the time, A for the overall ratio of fluorescence

recovery, and b for the inverse of the time constant $\tau_{1/2}$. For the fit, A was restricted to values

between 0.7 and 1.1. Each bleach curve was fitted individually, and fits with $R^2 < 0.7$ were excluded

from further analysis. The half-time of recovery was calculated from the time constant as follows:

$$t_{1/2} = \ln(2)\,\tau_{1/2}$$

Proteome and secretome analysis by shotgun proteomics based mass spectrometry

Strains were grown and prepared as described above. After 2.5 h at 37°, cultures were normalized

to an OD_{600} of 0.5 and 2 ml were harvested by centrifugation at 9,391 *g* for 2 min. Cell were washed

3 times with 1x phosphate buffered saline (PBS) (8 g/l NaCl, 0.2 g/l KCl, 1.78 g/l $Na_2HPO_4*2H_2O$, 0.24

g/l KH_2PO_4, pH 7.4). After washing, the cell pellet was resuspended in 300 µl lysis buffer (2% sodium

lauroyl sarcosinate (SLS), 100 mM ammonium bicarbonate (ABC)) and incubated at 90°C for 10 min.

Protein concentrations were measured using a BCA assay (Thermo Scientific) and following

reduction in 5 mM Tris(2-carboxyethyl) phosphin (TCEP, 15 min at 90 °C) and alkylation in 10 mM

iodoacetamid (30 min, 20°C in the dark), 50 µg protein was used for tryptic digestion (1 µg, Serva)

at 30°C overnight in presence of 0.5% SLS. After digestion, SLS was precipitated by adding a final

concentration of 1.5% trifluoroacetic acid (TFA).

Desalting of cleared peptide samples was performed in C18 solid phase extraction (SPE) cartridges

(Macherey-Nagel). First, cartridges were prepared by adding acetonitrile, followed by equilibration

using 0.1% TFA and loading of peptide samples. Cartridges were washed with buffer containing 5%

acetonitrile and 0.1% TFA, and finally eluted with 50% acetonitrile and 0.1% TFA into a fresh tube.

Eluted peptides were dried and reconstituted in 0.1% TFA for LC-MS analysis.

Purified peptides were analyzed by liquid chromatography-mass spectrometry (MS) carried out on

a Q-Exactive Plus instrument connected to an Ultimate 3000 rapid-separation liquid

chromatography (RSLC) nano instrument and a nanospray flex ion source (all Thermo Scientific).

Peptide separation was performed on a reverse-phase high-performance liquid chromatography

(HPLC) column (75 μm by 42 cm) packed in-house with C_{18} resin (2.4 μm; Dr. Maisch GmbH). The

peptides were first loaded onto a C18 precolumn (preconcentration set-up) and then eluted in

backflush mode with a gradient from 98 % solvent A (0.15 % formic acid) and 2 % solvent B (99.85

% acetonitrile, 0.15 % formic acid) to 25 % solvent B over 66 min, continued from 25 to 40 % of

solvent B up to 90 min. The flow rate was set to 300 nl/min. The data acquisition mode for label-

free quantification (LFQ) was set to obtain one high-resolution MS scan at a resolution of 70,000

(*m/z* 200) with scanning range from 375 to 1500 *m/z* followed by MS/MS scans of the 10 most

intense ions at a resolution of 17,500. To increase the efficiency of MS/MS shots, the charged state

screening modus was adjusted to exclude unassigned and singly charged ions. The dynamic

exclusion duration was set to 10 s. The ion accumulation time was set to 50 ms (both MS and

MS/MS). The automatic gain control (AGC) was set to 3×10^6 for MS survey scans and 1×10^5 for

MS/MS scans.

Due to an instrument upgrade, the QExactive Plus system was replaced by an Exploris 480 mass

spectrometer (Thermo Scientific). For total proteome analysis, purified peptides were analyzed

using identical LC settings, and the Exploris MS data acquisition settings were the following: one

high-resolution MS scan at a resolution of 60,000 (m/z 200) was obtained with scanning range from

350 to 1650 m/z followed by MS/MS scans of 2 sec cycle time at a resolution of 15,000. Charge state

inclusion was set between 2 and 6. The dynamic exclusion duration was set to 14 s. The ion

accumulation time was set to 25 ms for MS and AUTO for MS/MS. The automatic gain control (AGC)

was set to 300% for MS survey scans and 200% for MS/MS scans.

LFQ analysis was performed using MaxQuant [89] in standard settings using a *Y. enterolitica* protein

database containing proteins of the closely related *Y. enterocolitica* strain W22703 [90] and of the

pYVe227 virulence plasmid (GenBank entry AF102990.1). Statistical follow-up analysis of the

MaxQuant LFQ data was performed on an updated SafeQuant [91,92] R-script modified to routinely

process MaxQuant "protein groups" outputs. The missing value imputation was adapted from

Perseus [93] in default settings and implemented into the update SafeQuant script.   Unless

otherwise indicated, proteins with ≥3 detected peptides and a *p* value ≤ 0.01 of the sample versus

the control using Student's t-test were considered as enriched and displayed in the tables. Intensity

values were shaded according to their value (blue for high intensity; transparent for low intensity)

for easier visualization.

For secretome analysis and quantification of secreted effectors in the supernatant, 5 ml BHI cultures

were grown under secreting conditions in 15 ml tubes as otherwise described before. After two

hours of incubation at 37°C, the medium was changed from BHI to minimal medium supplemented

with nalidixic acid (35 mg/ml), diaminopimelic acid (80 mg/ml) where required, glycerol (0.4%),

$MgCl_2$ (20 mM) and EGTA (5 mM). Next, bacteria were again incubated at 37°C shaking for one hour.

Samples were OD-normalized in a 2 ml volume, and bacteria were removed by centrifugation (10

min at 21,000 *g*, 4°C). In the next step, the supernatant was removed and precipitated with a final

concentration of 10% TCA for 1–8 h at 4°C; proteins were collected by centrifugation for 15 min at

21,000 *g* at 4°C and washed once with 1 ml ice cold acetone. A detailed description of the

experimental procedure is shown in [94]. The remaining pellet was resuspended in 300 µl lysis buffer

and incubated at 90°C for 10 min. Following reduction and alkylation (see above), proteins were

digested using 1 µg trypsin. All further steps including C18-SPE, LC-MS analysis and LFQ data analysis

were identical to the ones described above. The LC separating gradient length was reduced to 45

min due to lower sample complexity.

Co-immunoprecipitation and shotgun proteomics

For co-immunoprepitation, 100 ml of non-secreting culture medium were inoculated to an OD_{600} of

~0.15 and treated as described above. After 2.5 h incubation at 37°C, the cultures were transferred

to 50 ml tubes and centrifuged at 2,000 *g* for 15 min at 4°C. Pellets were washed with 10 ml of 1x

PBS. The contents of two tubes were pooled and centrifuged at 2,000 *g* for 15 min at 4°C. The pellet

was resuspended in 2 ml HNN lysis buffer (50 mM HEPES pH 7.5, 150 mM NaCl, 50mM NaF, sterile

filtered; protease inhibitor (cOmplete Mini, EDTA free (Roche)) added before use), and incubated

for 30 min on a turning wheel at 4°C. Afterwards the bacteria were sonicated with 12 pulses à 30 s

in a Hielscher UP200ST ultrasonic homogenizer. The cells were again placed on a turning wheel for

30 min at 4°C and the sonification process and the subsequent incubation on a turning wheel were

repeated. Unbroken cells were removed by centrifugation (20,000 *g*, 20 min, 4°C). For affinity

purification, 10 µl of bead slurry (GFP-Trap Magnetic Agarose, Chromotek) were added to the lysate

and incubated shaking for 1 h at 4°C on a turning wheel. Beads were washed with 500 µl HNN lysis

buffer + 0.1% NP40, followed by seven consecutive washing steps with 700 µl 100 mM ABC. 200 µl

of elution buffer (100 mM ABC, 1 µg trypsin per sample) were added to the beads and incubated for

45 min on a thermomixer at 30°C at 1200 rpm. Beads were separated and the supernatant was

collected. Beads were washed twice using 80 µl of elution buffer 2 (1 M urea, 100 mM ABC, 5 mM

TCEP), combined with the first eluate fraction and incubated o/n at 30°C to complete proteolysis.

To monitor the experimental steps, samples for Western blot analysis were taken throughout the

whole procedure. After digest, 10 mM iodoacetamid were added and the samples were incubated

for 30 min in the dark. For desalting, C18 solid phase extraction cartridges samples were acidified

and the procedure was carried out as described above. The samples were analyzed on a QExactive

Plus mass spectrometer using a resolving LC gradient of 40 min. The analytical settings and data

analysis steps including MaxQuant and SafeQuant are described above. Unless otherwise indicated,

proteins with ≥3 detected peptides and a *p* value ≤ 0.001 of the sample versus the control using

Student's t-test were considered as enriched and displayed in the tables.

Sequence analysis

For the analysis of the genetic environment of T3SS pilotin proteins from model organisms, the

following GenBank sequences were used: AF102990.1 (*Yersinia enterocolitica* plasmid pYVe227,

complete sequence), NC_002516.2 (*Pseudomonas aeruginosa* PAO1, complete genome reference

sequence), NC_003197.2 (*Salmonella enterica* subsp. *enterica* serovar Typhimurium str. LT2,

complete genome reference sequence), NC_004851.1 (*Shigella flexneri* 2a str. 301 plasmid pCP301,

complete sequence), AM286415.1 (*Yersinia enterocolitica* subsp. *enterocolitica* 8081, complete

genome). Multiple sequence alignment of the *sctG* and *sctC* genes of the same systems was

performed using T-COFFEE 11.0 [95].

<u>**Acknowledgements**</u>

This work was supported by the Max Planck Society. The authors thank Christoph Spahn (Max Planck Institute for Terrestrial Microbiology, Marburg, Germany) for fruitful discussions about microscopy data analysis and his help with implementing the DeepBacs segmentation tool for *Y. enterocolitica.*

Author contributions

S.W. performed the majority of the experiments and data analysis and participated in study design and writing of the manuscript. M.F. performed experiments, implemented methods and generated key strains for this study. C.H., C.B. and K.L. assisted in experiments and strain generation. W.S., G.A. T.G. from the Proteomics facility provided this technology and assisted in data evaluation and statistical analysis. A.D. conceived and designed the study, performed data analysis and wrote the manuscript.

Competing interests

The authors declare no competing interests.

References

1.   Galán JE, Wolf-Watz H. Protein delivery into eukaryotic cells by type III secretion machines. Nature. 2006 Dec 1;444(7119):567–73.

2.   Cornelis GR. The type III secretion injectisome. Nat Rev Microbiol. 2006 Oct 17;4(11):811–25.

3.   Büttner D. Protein export according to schedule: architecture, assembly, and regulation of type III secretion systems from plant- and animal-pathogenic bacteria. Microbiol Mol Biol Rev. 2012 Jun 1;76(2):262–310.

4.   Burkinshaw BJ, Strynadka NC. Assembly and structure of the T3SS. Biochim Biophys Acta - Mol Cell Res. 2014 Feb 7;1843(8):1648–93.

5.   Wagner S, Grin I, Malmsheimer S, Singh N, Torres-Vargas CE, Westerhausen S. Bacterial type III secretion systems: a complex device for the delivery of bacterial effector proteins into eukaryotic host cells. FEMS Microbiol Lett. 2018 Aug 1;365(19).

6.   Diepold A, Wagner S. Assembly of the bacterial type III secretion machinery. FEMS Microbiol Rev. 2014 Jul 3;38(4):802–22.

7.   Journet L, Agrain C, Broz P, Cornelis GR. The needle length of bacterial injectisomes is determined by a molecular ruler. Science (80-). 2003 Dec 6;302(5651):1757–60.

8.   Wee DH, Hughes KT. Molecular ruler determines needle length for the Salmonella Spi-1 injectisome. Proc Natl Acad Sci. 2015 Mar 16;112(13):4098–103.

9.   Chakravortty D, Rohde M, Jäger L, Deiwick J, Hensel M. Formation of a novel surface structure encoded by Salmonella Pathogenicity Island 2. EMBO J. 2005 Jun 1;24(11):2043–52.

10.   Blocker AJ, Gounon P, Larquet E, Niebuhr K, Cabiaux V, Parsot C, et al. The tripartite type III secreton of Shigella flexneri inserts IpaB and IpaC into host membranes. J Cell Biol. 1999 Nov 1;147(3):683–93.

11.   Diepold A, Armitage JP. Type III secretion systems: the bacterial flagellum and the injectisome. Philos Trans R Soc B Biol Sci. 2015 Oct 5;370(1679):20150020.

12.   Kudryashev M, Diepold A, Amstutz M, Armitage JP, Stahlberg H, Cornelis GR. Yersinia enterocolitica type III secretion injectisomes form regularly spaced clusters, which incorporate new machines upon activation. Mol Microbiol. 2015 Dec 18;95(5):875–84.

13.   Zhang Y, Lara-Tejero M, Bewersdorf J, Galán JE. Visualization and characterization of individual type III protein secretion machines in live bacteria. Proc Natl Acad Sci U S A. 2017 Jun 6;114(23):6098–103.

14.   Burgess JL, Case HB, Burgess RA, Dickenson NE. Dominant negative effects by inactive Spa47 mutants inhibit T3SS function and Shigella virulence. Cascales E, editor. PLoS One. 2020 Jan 24;15(1):e0228227.

15.   Allaoui A, Scheen R, Lambert de Rouvroit C, Cornelis GR. VirG, a Yersinia enterocolitica lipoprotein involved in Ca2+ dependency, is related to exsB of Pseudomonas aeruginosa. J Bacteriol. 1995 Jul 31;177(15):4230–7.

16.   Hardie KR, Lory S, Pugsley AP. Insertion of an outer membrane protein in Escherichia coli requires a chaperone-like protein. EMBO J. 1996 Mar;15(5):978–88.

17.   Koo J, Burrows LL, Lynne Howell P. Decoding the roles of pilotins and accessory proteins in secretin escort services. FEMS Microbiol Lett. 2012 Mar 1;328(1):1–12.

18.   Rau R, Darwin AJ. Identification of YsaP, the pilotin of the Yersinia enterocolitica Ysa type III

secretion system. J Bacteriol. 2015;197(17):2770–9.

19.   Perdu C, Huber P, Bouillot S, Blocker AJ, Elsen S, Attree I, et al. ExsB is required for correct
assembly of Pseudomonas aeruginosa type III secretion apparatus in the bacterial membrane
and full virulence in vivo. Infect Immun. 2015 Feb 17;IAI.00048-15.

20.   Koster M, Bitter W, de Cock H, Allaoui A, Cornelis GR, Tommassen J. The outer membrane
component, YscC, of the Yop secretion machinery of Yersinia enterocolitica forms a ring-shaped
multimeric complex. Mol Microbiol. 1997 Nov;26(4):789–97.

21.   Daefler S, Russel M. The Salmonella typhimurium InvH protein is an outer membrane
lipoprotein required for the proper localization of InvG. Mol Microbiol. 1998 Jun 1;28(6):1367–
80.

22.   Crago A, Koronakis V. Salmonella InvG forms a ring-like multimer that requires the InvH
lipoprotein for outer membrane localization. Mol Microbiol. 1998 Oct 1;30(1):47–56.

23.   Nouwen N, Ranson N, Saibil HR, Wolpensinger B, Engel A, Ghazi A, et al. Secretin PulD:
association with pilot PulS, structure, and ion-conducting channel formation. Proc Natl Acad Sci
U S A. 1999 Jul 5;96(14):8173–7.

24.   Schuch R, Maurelli AT. MxiM and MxiJ, base elements of the Mxi-Spa type III secretion system
of Shigella, interact with and stabilize the MxiD secretin in the cell envelope. J Bacteriol. 2001
Nov 22;183(24):6991–8.

25.   Burghout P, Beckers F, de Wit E, van Boxtel R, Cornelis GR, Tommassen J, et al. Role of the pilot
protein YscW in the biogenesis of the YscC secretin in Yersinia enterocolitica. J Bacteriol. 2004
Aug 31;186(16):5366–75.

26.   Korotkov K V, Gonen T, Hol WGJ. Secretins: dynamic channels for protein transport across
membranes. Trends Biochem Sci. 2011 May 9;

27.   Gu S, Rehman S, Wang X, Shevchik VE, Pickersgill RW. Structural and functional insights into the
pilotin-secretin complex of the type II secretion system. PLoS Pathog. 2012;8(2).

28.   Diepold A, Amstutz M, Abel S, Sorg I, Jenal U, Cornelis GR. Deciphering the assembly of the
Yersinia type III secretion injectisome. EMBO J. 2010 Jun 2;29(11):1928–40.

29.   Diepold A, Wiesand U, Cornelis GR. The assembly of the export apparatus (YscR,S,T,U,V) of the
Yersinia type III secretion apparatus occurs independently of other structural components and
involves the formation of an YscV oligomer. Mol Microbiol. 2011 Oct 16;82(2):502–14.

30.   Okon M, Moraes TF, Lario PI, Creagh AL, Haynes CA, Strynadka NC, et al. Structural
Characterization of the Type-III Pilot-Secretin Complex from Shigella flexneri. Structure. 2008
Oct 8;16(10):1544–54.

31.   Majewski DD, Okon M, Heinkel F, Robb CS, Vuckovic M, McIntosh LP, et al. Characterization of
the Pilotin-Secretin Complex from the Salmonella enterica Type III Secretion System Using
Hybrid Structural Methods. Structure. 2021 Feb;29(2):125-138.e5.

32.   Collin S, Guilvout I, Nickerson NN, Pugsley AP. Sorting of an integral outer membrane protein via
the lipoprotein-specific Lol pathway and a dedicated lipoprotein pilotin. Mol Microbiol.
2011;80(3):655–65.

33.   Chami M, Guilvout I, Gregorini M, Rémigy HW, Müller SA, Valerio M, et al. Structural insights
into the secretin PulD and its trypsin-resistant core. J Biol Chem. 2005;280(45):37732–41.

34.   Yin M, Yan Z, Li X. Structural insight into the assembly of the type II secretion system pilotin–
secretin complex from enterotoxigenic Escherichia coli. Nat Microbiol. 2018 May 9;3(5):581–7.

35.   Chernyatina AA, Low HH. Core architecture of a bacterial type II secretion system. Nat Commun.

2019;10(1):1–10.

36.   Hu B, Morado DR, Margolin W, Rohde JR, Arizmendi O, Picking WL, et al. Visualization of the
type III secretion sorting platform of Shigella flexneri. Proc Natl Acad Sci. 2015 Jan
12;112(4):1047–52.

37.   Hu B, Lara-Tejero M, Kong Q, Galán JE, Liu J. In Situ Molecular Architecture of the Salmonella
Type III Secretion Machine. Cell. 2017 Mar 9;168(6):1065-1074.e10.

38.   Troisfontaines P, Cornelis GR. Type III Secretion: More Systems Than You Think. Physiology.
2005 Oct;20(5):326–39.

39.   Abby SS, Rocha EPC. The non-flagellar type III secretion system evolved from the bacterial
flagellum and diversified into host-cell adapted systems. PLoS Genet. 2012 Sep;8(9):e1002983.

40.   Iriarte M, Cornelis GR. Identification of SycN, YscX, and YscY, three new elements of the Yersinia
yop virulon. J Bacteriol. 1999;181(2):675–80.

41.   Diepold A, Wiesand U, Amstutz M, Cornelis GR. Assembly of the Yersinia injectisome: the
missing pieces. Mol Microbiol. 2012 Sep;85(5):878–92.

42.   Böhme K, Steinmann R, Kortmann J, Seekircher S, Heroven AK, Berger E, et al. Concerted actions
of a thermo-labile regulator and a unique intergenic RNA thermosensor control Yersinia
virulence. Isberg RR, editor. PLoS Pathog. 2012 Feb 16;8(2):e1002518.

43.   Diepold A, Sezgin E, Huseyin M, Mortimer T, Eggeling C, Armitage JP. A dynamic and adaptive
network of cytosolic interactions governs protein export by the T3SS injectisome. Nat Commun.
2017;8(May):15940.

44.   Milne-Davies B, Helbig C, Wimmi S, Cheng DWC, Paczia N, Diepold A. Life After Secretion—
Yersinia enterocolitica Rapidly Toggles Effector Secretion and Can Resume Cell Division in
Response to Changing External Conditions. Front Microbiol. 2019 Sep 13;10:2128.

45.   Kaniga K, Delor I, Cornelis GR. A wide-host-range suicide vector for improving reverse genetics
in Gram-negative bacteria: inactivation of the blaA gene of Yersinia enterocolitica. Gene. 1991
Dec 20;109(1):137–41.

46.   Walker KA, Miller VL. Synchronous Gene Expression of the Yersinia enterocolitica Ysa Type III
Secretion System and Its Effectors. J Bacteriol. 2009 Mar 15;191(6):1816–26.

47.   Mazé A, Glatter T, Bumann D. The central metabolism regulator EIIAGlc switches salmonella
from growth arrest to acute virulence through activation of virulence factor secretion. Cell Rep.
2014;7(5):1426–33.

48.   Abrusci P, Vergara-Irigaray M, Johnson S, Beeby MD, Hendrixson DR, Roversi P, et al.
Architecture of the major component of the type III secretion system export apparatus. Nat
Struct Mol Biol. 2013 Jan 1;20(1):99–104.

49.   Diepold A, Kudryashev M, Delalez NJ, Berry RM, Armitage JP. Composition, Formation, and
Regulation of the Cytosolic C-ring, a Dynamic Component of the Type III Secretion Injectisome.
Stock AM, editor. PLOS Biol. 2015 Jan 15;13(1):e1002039.

50.   Wimmi S, Balinovic A, Jeckel H, Selinger L, Lampaki D, Eisemann E, et al. Dynamic relocalization
of cytosolic type III secretion system components prevents premature protein secretion at low
external pH. Nat Commun. 2021 Dec 12;12(1):1625.

51.   Schwiesow L, Lam H, Dersch P, Auerbuch V. Yersinia Type III Secretion System Master Regulator
LcrF. Margolin W, editor. J Bacteriol. 2016 Feb 15;198(4):604–14.

52.   Bange G, Kümmerer N, Engel C, Bozkurt G, Wild K, Sinning I. FlhA provides the adaptor for
coordinated delivery of late flagella building blocks to the type III secretion system. Proc Natl

Acad Sci U S A. 2010 Jun 21;107(25):11295–300.

53.   Xing Q, Shi K, Portaliou AG, Rossi P, Economou A, Kalodimos CG. Structures of chaperone-
substrate complexes docked onto the export gate in a type III secretion system. Nat Commun.
2018 Dec 2;9(1):1773.

54.   Inoue Y, Kinoshita M, Kida M, Takekawa N, Namba K, Imada K, et al. The FlhA linker mediates
flagellar protein export switching during flagellar assembly. Commun Biol. 2021;4(1).

55.   Sorg I, Wagner S, Amstutz M, Müller SA, Broz P, Lussi Y, et al. YscU recognizes translocators as
export substrates of the Yersinia injectisome. EMBO J. 2007 Jun 19;26(12):3015–24.

56.   Day JB, Plano G V. A complex composed of SycN and YscB functions as a specific chaperone for
YopN in Yersinia pestis. Mol Microbiol. 1998 Nov 1;30(4):777–88.

57.   Botteaux A, Sory M-P, Biskri L, Parsot C, Allaoui A. MxiC is secreted by and controls the
substrate specificity of the Shigella flexneri type III secretion apparatus. Mol Microbiol.
2009;71(2):449–60.

58.   Lara-Tejero M, Kato J, Wagner S, Liu X, Galán JE. A Sorting Platform Determines the Order of
Protein Secretion in Bacterial Type III Systems. Science (80-). 2011 Feb 2;331(6021):1188–91.

59.   Schuch R, Maurelli AT. The Mxi-Spa type III secretory pathway of Shigella flexneri requires an
outer membrane lipoprotein, MxiM, for invasin translocation. Infect Immun. 1999;67(4):1982–
91.

60.   Jeckelmann JM, Erni B. Transporters of glucose and other carbohydrates in bacteria. Pflugers
Arch Eur J Physiol. 2020;472(9):1129–53.

61.   Pflüger-Grau K, Görke B. Regulatory roles of the bacterial nitrogen-related phosphotransferase
system. Trends Microbiol. 2010;18(5):205–14.

62.   Pacheco AR, Sperandio V. Enteric pathogens exploit the microbiota-generated nutritional
environment of the gut. Metab Bact Pathog. 2015;(10):279–96.

63.   Jeckelmann JM, Erni B. Carbohydrate Transport by Group Translocation: The Bacterial
Phosphoenolpyruvate: Sugar Phosphotransferase System. Vol. 92, Subcellular Biochemistry.
Springer International Publishing; 2019. 223–274 p.

64.   Palace SG, Proulx MK, Lu S, Baker RE, Goguen JD. Genome-wide mutant fitness profiling
identifies nutritional requirements for optimal growth of Yersinia pestis in deep tissue. MBio.
2014;5(4).

65.   Wang Q, Millet YA, Chao MC, Sasabe J, Davis BM, Waldor MK. A genome-wide screen reveals
that the Vibrio cholerae phosphoenolpyruvate phosphotransferase system modulates virulence
gene expression. Infect Immun. 2015;83(9):3381–95.

66.   Kok M, Bron G, Erni B, Mukhija S. Effect of enzyme I of the bacterial phosphoenolpyruvate :
Sugar phosphotransferase system (PTS) on virulence in a murine model. Microbiology.
2003;149(9):2645–52.

67.   Collins J, Robinson C, Danhof H, Knetsch CW, Van Leeuwen HC, Lawley TD, et al. Dietary
trehalose enhances virulence of epidemic Clostridium difficile. Nature. 2018;553(7688):291–4.

68.   Carter PB, Zahorchak RJ, Brubaker RR. Plague virulence antigens from Yersinia enterocolitica.
Infect Immun. 1980 May;28(2):638–40.

69.   Sasakawa C, Kamata K, Sakai T, Murayama SY, Makino S, Yoshikawa M. Molecular alteration of
the 140-megadalton plasmid associated with loss of virulence and Congo red binding activity in
Shigella flexneri. Infect Immun. 1986 Feb;51(2):470–5.

70. Sturm A, Heinemann M, Arnoldini M, Benecke A, Ackermann M, Benz M, et al. The cost of
virulence: retarded growth of Salmonella Typhimurium cells expressing type III secretion system
1. PLoS Pathog. 2011 Jul;7(7):e1002143.

71. Gur E, Sauer RT. Recognition of misfolded proteins by Lon, a AAA+ protease. Genes Dev.
2008;22(16):2267–77.

72. Tsilibaris V, Maenhaut-Michel G, Van Melderen L. Biological roles of the Lon ATP-dependent
protease. Res Microbiol. 2006;157(8):701–13.

73. Omnus DJ, Fink MJ, Szwedo K, Jonas K. The lon protease temporally restricts polar cell
differentiation events during the caulobacter cell cycle. Elife. 2021;10:1–24.

74. Cornelis GR, Sluiters C, Delor I, Geib D, Kaniga K, de Rouvroit CL, et al. ymoA, a Yersinia
enterocolitica chromosomal gene modulating the expression of virulence functions. Mol
Microbiol. 1991 May;5(5):1023–34.

75. Cornelis GR. Role of the Transcription Activator VirF and the Histone-like Protein YmoA in the
Thermoregulation of Virulence Functions in Yersiniae. Zentralblatt fur Bakteriol. 1993;278(2–
3):149–64.

76. Jackson MW, Silva-Herzog E, Plano G V. The ATP-dependent ClpXP and Lon proteases regulate
expression of the Yersinia pestis type III secretion system via regulated proteolysis of YmoA, a
small histone-like protein. Mol Microbiol. 2004;54(5):1364–78.

77. Breidenstein EBM, Janot L, Strehmel J, Fernandez L, Taylor PK, Kukavica-Ibrulj I, et al. The Lon
Protease Is Essential for Full Virulence in Pseudomonas aeruginosa. PLoS One. 2012;7(11).

78. Bretz J, Losada L, Lisboa K, Hutcheson SW. Lon protease functions as a negative regulator of
type III protein secretion in Pseudomonas syringae. Mol Microbiol. 2002;45(2):397–409.

79. Zhou X, Teper D, Andrade MO, Zhang T, Chen S, Song WY, et al. A phosphorylation switch on lon
protease regulates bacterial type iii secretion system in host. MBio. 2018;9(1).

80. Lee JH, Ancona V, Zhao Y. Lon protease modulates virulence traits in Erwinia amylovora by
direct monitoring of major regulators and indirectly through the Rcs and Gac-Csr regulatory
systems. Mol Plant Pathol. 2018;19(4):827–40.

81. Figaj D, Czaplewska P, Przepióra T, Ambroziak P, Potrykus M, Skorko-Glonek J. Lon protease is
important for growth under stressful conditions and pathogenicity of the phytopathogen,
bacterium Dickeya solani. Int J Mol Sci. 2020;21(10).

82. Renault TT, Guse A, Erhardt M. Export Mechanisms and Energy Transduction in Type-III
Secretion Machines. In: Current Topics in Microbiology and Immunology. Springer, Berlin,
Heidelberg; 2019. p. 1–17.

83. Konovalova A, Silhavy TJ. Outer membrane lipoprotein biogenesis: Lol is not the end. Philos
Trans R Soc B Biol Sci. 2015;370(1679).

84. Kudryashev M, Stenta M, Schmelz S, Amstutz M, Wiesand U, Castaño-Díez D, et al. In situ
structural analysis of the Yersinia enterocolitica injectisome. Elife. 2013 Jan;2(Im):e00792.

85. Lambert de Rouvroit C, Sluiters C, Cornelis GR. Role of the transcriptional activator, VirF, and
temperature in the expression of the pYV plasmid genes of Yersinia enterocolitica. Mol
Microbiol. 1992 Feb;6(3):395–409.

86. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-
source platform for biological-image analysis. Nat Methods. 2012 Jul 28;9(7):676–82.

87. Thevenaz P, Ruttimann UE, Unser M. A pyramid approach to subpixel registration based on
intensity. IEEE Trans Image Process. 1998;7(1):27–41.

88. Spahn C, Laine RF, Pereira P, Gómez-de-mariscal E. Supplementary information - DeepBacs :
Bacterial image analysis using open-source deep learning approaches. bioRxiv. 2021;1–15.

89. Cox J, Mann M. MaxQuant enables high peptide identification rates, individualized p.p.b.-range
mass accuracies and proteome-wide protein quantification. Nat Biotechnol. 2008 Dec
30;26(12):1367–72.

90. Fuchs TM, Brandt K, Starke M, Rattei T. Shotgun sequencing of Yersinia enterocolitica strain
W22703 (biotype 2, serotype O:9): Genomic evidence for oscillation between invertebrates and
mammals. BMC Genomics. 2011;12:1–14.

91. Glatter T, Ludwig C, Ahrné E, Aebersold R, Heck AJR, Schmidt A. Large-scale quantitative
assessment of different in-solution protein digestion protocols reveals superior cleavage
efficiency of tandem Lys-C/trypsin proteolysis over trypsin digestion. J Proteome Res.
2012;11(11).

92. Ahrné E, Glatter T, Viganò C, Von Schubert C, Nigg EA, Schmidt A. Evaluation and improvement
of quantification accuracy in isobaric mass tag-based protein quantification experiments. J
Proteome Res. 2016;15(8):2537–47.

93. Tyanova S, Temu T, Sinitcyn P, Carlson A, Hein MY, Geiger T, et al. The Perseus computational
platform for comprehensive analysis of (prote)omics data. Nat Methods. 2016;13(9):731–40.

94. Lampaki D, Diepold A, Glatter T. A Serial Sample Processing Strategy with Improved
Performance for in-Depth Quantitative Analysis of Type III Secretion Events in Pseudomonas
aeruginosa. J Proteome Res. 2020 Jan 3;19(1):543–53.

95. Wallace IM, O'Sullivan O, Higgins DG, Notredame C. M-Coffee: combining multiple sequence
alignment methods with T-Coffee. Nucleic Acids Res. 2006;34:1692–9.

96. Erhardt M, Namba K, Hughes KT. Bacterial nanomachines: the flagellum and type III injectisome.
Cold Spring Harb Perspect Biol. 2010 Nov 1;2(11):a000299.

97. Hueck CJ. Type III protein secretion systems in bacterial pathogens of animals and plants.
Microbiol Mol Biol Rev. 1998 Jun;62(2):379–433.

98. Wagner S, Diepold A. A Unified Nomenclature for Injectisome-Type Type III Secretion Systems.
In: Current topics in microbiology and immunology. 2020. p. 427: 1-10.

949