Variation in *Plasmodium falciparum* sexual commitment rates and responses
to environmental modification

Short title: Malaria parasite sexual commitment rate variation

Lindsay B. Stewart [1a], Aline Freville [1a], Till S. Voss [2], David A. Baker [1], Gordon A. Awandare [3],
David J. Conway [1*]

[1] Department of Infection Biology, Faculty of Infectious and Tropical Diseases, London
School of Hygiene and Tropical Medicine, London WC1E 7HT, UK

[2] Department of Medical Parasitology and Infection Biology, Swiss Tropical and Public Health
Institute, and University of Basel, Switzerland

[3] West African Centre for Cell Biology of Infectious Pathogens (WACCBIP), Department of
Biochemistry, Cell and Molecular Biology, University of Ghana, Accra, Ghana

[a] These authors contributed equally to the research

[*] Corresponding author: david.conway@lshtm.ac.uk

Abstract

Asexual blood stage malaria parasites must produce sexual progeny in order to infect

mosquitoes. It is therefore important to understand the scope of variation in sexual

commitment rates of the most important human parasite *Plasmodium falciparum*, and to

assess responses to conditions that may affect this commitment. First, using a genetically

modified line with inducible elevated sexual commitment we compared two different

methods of measuring the rates, either staining the Pfs16 marker in very early gametocytes

or counting gametocytes after further development. The methods showed correlated

estimates in evaluation across multiple independent replicate assays, with higher sensitivity

and precision being achieved by Pfs16 marker detection in early gametocytes, so this

method was used to survey a wide range of *P. falciparum* lines. A panel of six recently

culture-established clinical isolates from Ghana showed mean sexual commitment rates per

cycle ranging from 3.3% to 12.2%, with some significant differences among isolates as well

as variation among biological replicates of each. Comparing 13 long-term laboratory-

adapted lines of diverse origins, eight had sexual commitment rates similar to those of the

recent clinical isolates, with means across multiple biological replicates for each line ranging

from 4.7% to 14.5%, while four lines had significantly lower rates with means ranging from

0.3 to 1.6%, and one line with a non-functional *ap2-g* gene always showed zero sexual

commitment. Among a subset of parasite lines, adding choline to suppress commitment had

quantitatively variable effects, although these were significant in most assays of lines that

had relatively high rates. This study indicates the importance of multiple assay replicates

and comparisons of diverse isolates to understand natural and culture-induced variation in

this key reproductive trait, relevant to investigating potential effects of parasite adaptation

on transmission.

Author Summary

Only sexual malaria parasites are transmitted from humans to mosquitoes, so it is vital to

understand variation in sexual commitment rates of blood stage malaria parasites, and

responses to conditions that affect this. We compared two different methods of measuring

the rates in a *Plasmodium falciparum* line with engineered sexual commitment variation,

demonstrating higher sensitivity and precision by detection of an early differentiation

marker, and this method was then used to survey diverse *P. falciparum* lines in culture with

extensive biological replicate measurements. Recent clinical isolates from Ghana showed

mean sexual commitment rates per cycle ranging from 3% to 12%, with significant

differences among isolates as well as variation among biological replicates. Long-term

laboratory-adapted lines of diverse origins had a wide range, most having rates similar to

those of the clinical isolates, while a minority consistently had much lower or zero rates.

There was quantitative variation among lines in the effects of adding choline to suppress

commitment, although most assays of lines that had relatively high rates showed significant

effects. Performing multiple biological replicates and comparisons of recent parasite isolates

is vital to understand intra-specific variation in this important reproductive trait, and move

towards investigating direct effects on disease transmission.

Introduction

For malaria to be transmitted, asexual blood stage parasites must undergo differentiation

which leads to formation of male and female gametocytes that are required to infect

mosquitoes. This process involves an initial parasite commitment phase, when the switch to

sexual development is induced, followed by a longer developmental phase to produce the

mature gametocyte forms. From an evolutionary or ecological perspective, it is important to

understand what determines the rate of commitment to sexual forms [1], and how this may

vary within a species [2]. It is particularly vital for understanding *P. falciparum* transmission

as more effective control is needed to reduce the global malaria burden [3], which is

particularly challenging as chronic asymptomatic infections are often highly infective to

mosquitoes [4].

Research on malaria parasite sexual conversion has been long established [5], and progress

has been made in elucidating key aspects of the mechanism. An ApiAP2-family transcription

factor AP2-G is essential for sexual conversion, controlling expression of genes critical to

early gametocytogenesis [6-9]. The *ap2-g* locus is clonally variably expressed, with histone 2

lysine 9 trimethylation (H3K9me3) marks of heterochromatin silencing initially indicating

that its expression could be epigenetically regulated [10], and conditional depletion of

heterochromatin protein 1 (HP1) from the promoter was shown to increase *ap2-g*

transcription and sexual conversion [11]. Furthermore, in *P. falciparum* a gametocyte

development protein GDV1, that was previously linked with sexual commitment [12], is

involved in removal of HP1 from the *ap2-g* locus so that over-expression of *gdv1* causes

elevated sexual commitment rates [13].

The classic model is that sexual commitment occurs in the intra-erythrocytic cycle prior to

the cycle in which parasite sexual differentiation occurs, based on a finding that in static

culture individual schizonts of the 3D7 parasite clone yield spatially-clustered intra-

erythrocytic progeny that tend to differentiate similarly, either sexually or asexually [14].

However, there is recent evidence that some sexual commitment may also occur in the

same cycle as sexual differentiation, if *ap2-g* expression is induced to start very early after

erythrocyte invasion during the ring stage [15], although it is not known how commonly this

occurs in parasites under normal conditions of infection or laboratory culture. It is difficult to

study *P. falciparum* gametocyte conversion rates precisely in natural infections, as most

stages of developing gametocytes do not occur in the peripheral circulation and are

therefore not detected in blood samples, although circulating ring stage parasites that are

sexually committed may become detectable by developing assays for early markers such as

AP2-G or GEXP02 [16, 17]. Correlations between *P. falciparum* gametocyte numbers and

other measured variables in human hosts, such as age or hemoglobin levels, may not be

causally related to varying gametocyte conversion rates and are not well replicated across

different studies [18]. Experiments with other malaria parasite species in laboratory mice

indicate sexual commitment to be a plastic phenotype that can respond to the *in vivo*

environment [19]. There are few data relating to *P. falciparum* gametocyte conversion in

experimentally induced human infections, but modelling of results from human

experimental infections with the parasite clone 3D7 has indicated that the conversion rate

per cycle is usually less than 1% [20], and older data on induced malaria infections with

strains that no longer exist suggest broadly similar average rates [21].

Early studies of *P. falciparum* in culture indicated that gametocyte production varied among

parasite clones [22, 23], and also varied with culture conditions, although precise estimates

of conversion rates per cycle were not established. Subsequent studies estimated

gametocyte conversion rates of several laboratory-adapted *P. falciparum* clones to be

usually above 1% per cycle when grown at low parasitaemia [6, 24], with higher rates if

parasites were stressed at high parasitaemia or by antimalarial drug treatment [14, 24].

Parasites appear to respond to metabolic challenges during growth, although most cellular

mechanisms remain to be identified [25]. An increase in gametocytogenesis may be

triggered by cellular stress involving redox perturbation [26, 27], by adding low doses of the

antimalarial artemisinin at the early trophozoite growth stage [28], or by adding parasite-

conditioned culture medium [29] or extracellular vesicles from such medium [30]. Notably,

gametocyte conversion rates in cultured lines are suppressed by lysophosphatidylcholine

(LysoPC) within serum, or by adding choline as a supplement in serum-free medium [31, 32].

In a study of parasite development during the initial *ex vivo* cycle cultured from clinical

samples in Ghana, the proportions of parasites showing conversion to gametocytes

correlated positively with level of parasitaemia and negatively with concentration of LysoPC

in plasma of patients, supporting a hypothesis that developmental responses to metabolic

stress may occur during natural infections [33].

Despite these advances, the quantitative scope of sexual commitment rate variation in *P.*

falciparum has not yet been clearly defined. Estimates of commitment rates have previously

varied due to methodological differences or to biological differences between nominally

similar parasite lines, and there are few controlled comparisons of different isolates

employing biological replicate measurements. Performing multiple independent replicate

assays is important, because some individual lines may have temporally variable phenotypes

and because precision is generally increased with repeated measurements. In this study, we

first compared different methods of measuring sexual commitment rates, and then

employed the method with greater quantitative sensitivity and accuracy to conduct multiple

biological replicate assays on recently adapted clinical isolates from Ghana as well as a broad

selection of long-term laboratory adapted lines representing most of those that are

commonly cultured globally. Although each clinical isolate showed variability among

repeated assays, overall there are significant differences among isolates, indicating

potentially adaptive variation in sexual commitment within this important parasite species.

Separate assays of selected lines on the suppressive effect on sexual commitment by adding

choline to serum-free culture media identified significant quantitative variation among lines,

which further encourages research to test for relevance to disease transmission.

Results

Comparison of sensitivity and precision of two different assays to measure gametocyte

conversion rates

Two different assay methods were compared to quantify sexual commitment rates in

multiple independent experiments, using a genetically engineered *P. falciparum* clone

(3D7/iGP_D9) to over-express GDV1 with a destabilising domain (DD) in the presence of the

stabilising reagent Shield-1 which leads to elevated gametocyte conversion [34]. Method 1

involved differential counting of stage I gametocytes detected by anti-Pfs16

immunofluorescence staining as a proportion of all parasites developing beyond ring stages

in the next cycle, while Method 2 involved Giemsa-stained slide microscopy to compare the

next cycle ring stage parasitaemia (referred to as day 0 in this assay, D0) with the

parasitaemia of morphologically-defined stage II gametocytes (at day 4, D4) (details given in

the Materials and Methods section). In each of multiple biological replicate experiments,

both methods clearly show that the gametocyte conversion rate is very significantly higher

when GDV1 is overexpressed and stabilised with Shield-1 (GDV1-DD-ON) compared to when

GDV1 overexpression is disabled by the absence of Shield-1 (GDV1-DD-OFF) (Figure 1,

Supplementary Table S1).

In all biological replicate assays, Method 1 showed a marked difference in gametocyte

conversion rate between the GDV1-DD ON and OFF conditions, with a mean conversion rate

of 36.9 % under ON and 6.7 % under OFF conditions (Figure 1). In comparison, Method 2

yielded an estimated mean conversion rate of 15.9 % under ON and 4.1 % under OFF

conditions, with the rates for Method 2 being lower than for Method 1 in all replicates for

which both were tested in parallel. The precision of the estimates for each of the biological

replicates was higher for Method 1, with relatively narrow confidence intervals for each

experimental replicate as this method is based on direct counts of developmentally discrete

parasites. In comparison, Method 2 yielded wider confidence intervals as it is based on a

ratio-of-ratio proportion of two temporally sequential counts of parasitaemia which involves

more statistical sampling error (Supplementary Table S1), and this method also depends on

successful parasite development over a longer time in culture.

Correlation between gametocyte conversion rate estimates from different assay methods

To test for overall correlation between the estimates from the two assay methods, analysis

was performed on data from a broader range of paired assays. In addition to the eight

measurements from paired assays performed in the first comparison (Figure 1) nine

additional measurements of the 3D7/iGP_D9 clone were performed with different

concentrations of the Shield-1 reagent, to yield a total of 17 paired measurements with both

methods for analysis (Supplementary Table S2). Overall, this showed a high and significant

nonparametric rank correlation between the results from each assay method (Spearman's

rho = 0.70, P = 0.003), with Method 1 generally giving higher conversion rate estimates

(Figure 2) with narrower 95% confidence intervals (Supplementary Table S2). While this

supports both assays as separately giving informative relative estimates of gametocyte

conversion, Method 1 was chosen for subsequent quantitative analyses of additional lines

and culture conditions given its significantly higher sensitivity and tighter precision (Figures 1

and 2, Supplementary Tables S1 and S2).

Variation in sexual commitment rates of recently culture-established Ghanaian clinical isolates

Recent culture-established clinical isolates from a highly endemic area in Ghana were assayed to measure gametocyte conversion rates. Six isolates were selected from a larger panel for analysis as they represent unrelated genetically unmixed infections [35], so that biological replicate assays correspond to a single parasite genotype in each case, and after culture for several months they did not have any known loss of function mutations detected in their genome sequences that would affect sexual commitment [35]. These isolates show mean gametocyte conversion rates that range from 3.3% to 12.2% (Figure 3, Supplementary Table S3), and there were significant differences in pairwise comparisons (Mann-Whitney tests on results of a minimum of six biological replicate assays performed for each isolate, Supplementary Table S4). For example, isolate 296 had a significantly higher conversion rate than three of the others (isolates 272, 289 and 292), and isolate 289 had a significantly lower rate than three of the others (292, 293 and 296)(Supplementary Table S4).

There was also significant variation in gametocyte conversion rates among individual biological replicate assays of each isolate, with non-overlapping 95% confidence intervals of some replicates (Figure 3A). The variation among biological replicate assays for each line was not correlated with the culture parasitaemia, which was always between 0.5 and 2.5% (Figure 3B and Supplementary Table S3). Overall, there was no significant evidence of reduction in gametocyte conversion rates over time in culture (Supplementary Table S3), and variation among replicates did not correspond with the use of different erythrocyte batches for culture (Supplementary Table S3).

Range of gametocyte conversion rates among diverse long-term cultured laboratory lines

To compare with these recent clinical isolates, a panel of 13 long-term culture-adapted *P. falciparum* lines of diverse origins was tested, with multiple biological replicate assays (Figure 4, Supplementary Table S5). The clone F12 (containing a nonsense mutation in *ap2-g*)[6] here showed zero conversion in all replicate assays (Figure 4A, Supplementary Table S5). Four of the other lines showed consistently low levels of gametocyte conversion, with means ranging between 0.3% and 1.6%, significantly lower than the conversion rates of the remaining lines (Mann-Whitney tests on pairwise comparisons, Supplementary Table S6).

The lines with the lowest conversion rates (D10 and T9/96) were previously described as

having multiple gene deletions in a sub-telomeric region of chromosome 9 including the

gdv1 gene [36]. The 3D7 line may contain undetected mutants within bulk culture, as 3D7

has been elsewhere shown to be a source of parasite subclones with loss of function

mutations (including the *ap2-g* nonsense mutation in parasite subclone F12 [6], deletion of

gdv1 in subclone 3D7.G$_{def}$ [12] and truncation of *gdv1* in two other subclones [37]). The Palo

Alto line was previously shown to have undergone sub-telomeric gene deletions on different

chromosomes [36, 38] but not to have lost the *gdv1* locus on chromosome 9 [36, 39],

although the possibility of a loss-of-function mutation has not been excluded.

For the remaining eight long-term culture-adapted *P. falciparum* lines, the gametocyte

conversion rates for each showed means ranging from 4.7% (for D6) to 14.5% (for FCC2)

(Figure 4A). This range is similar to that seen for the recently culture-established clinical

isolates, with no significant differences between these groups (Mann-Whitney test, P =

0.47). Similar to the clinical isolates, the individual long-term culture-adapted parasite lines

also showed significant variation in gametocyte conversion rates among biological replicate

assays (Figure 4A), with non-overlapping 95% confidence intervals for some replicates

(Figure 4B). This shows that high and variable sexual commitment rates have been

maintained in a majority of the diverse cultured *P. falciparum* lines, in contrast to a minority

of lines that have markedly reduced rates or zero commitment.

Choline in culture medium suppresses sexual commitment rate with varying levels of

effect among *P. falciparum* lines

As it has been previously shown that parasite sexual commitment responds to the presence

or absence of choline in culture medium [13, 28, 31], we investigated five long-term culture-

adapted lines (T9/96, 3D7, Dd2, NF54 and HB3) for quantitative effects of 2 mM choline

compared to no added choline, performing between six and eight biological replicate assay

comparisons for each parasite line. Significant suppression of sexual commitment in the

presence of choline occurred in most lines (Figure 5, Supplementary Table S7). No significant

effect was seen in the *P. falciparum* clone T9/96 which has a deletion of *gdv1* and several

adjacent genes deleted on Chromosome 9 [36], and which has a very low rate of sexual

commitment either in the presence or absence of choline, consistent with the importance of

gdv1 in responding to environmental induction [13]. The responses of the other lines to

choline varied, with statistically significant suppression in sexual commitment in most of the

replicate assays for NF54 and Dd2, in half of the assays for HB3, and in a quarter of the

assays for 3D7 (Figure 5). Some individual assay replicates showed more than 10 fold

suppression of sexual commitment rate in the presence of choline, but most showed more

moderate rates (Supplementary Table S7).

The mean response of each parasite line to choline was estimated by the Mantel-Haenszel

rate ratio (RR_{M-H}), which analyses data across replicates under a simple fixed-effects model.

With the exception of T9/96 which showed no response as noted above, the remaining four

lines show overall increases in gametocyte conversion rates, but the mean responses varied

among the different lines (Supplementary Table S7). The greatest mean response to choline

was shown by line Dd2 (3.5 fold, 95% CI 3.0-4.1 fold), while 3D7 showed the lowest level of

effect (1.5 fold, 95% CI 1.2-1.9 fold), and the other lines showed intermediate levels (for

NF54 = 2.4 fold, 95% CI 2.0-3.0 fold; for HB3 = 2.0 fold, 95% CI 1.6-2.5 fold).

Discussion

Analyses of multiple biological replicate assays are informative to characterise the range of

sexual commitment rates in *P. falciparum*. A controlled comparison of two alternative

methods showed that phenotyping was more sensitive and precise with an assay that

measures proportions of sexually differentiating parasites by immunofluorescence staining

of the Pfs16 marker expressed in early stage I gametocytes [15]. Although the sensitivity and

precision of the methods differ, they give correlated measurements, and the lower

proportions measured using a method that does not employ Pfs16 staining may be

expected, given the requirement for parasites to successfully develop to stage II

gametocytes over four days in culture before they are counted by morphological inspection

of Giemsa-stained slides [33].

The recently culture-established Ghanaian clinical isolates analysed here had gametocyte

conversion rates of between 3.3% and 12.2% per cycle, measured as means of results from

at least six biological replicate assays for each isolate. These rates are higher than those

previously estimated for clinical isolates in the first cycle *ex vivo* in Ghana with an assay

which involves counting of stage II gametocytes, which indicated median rates less than 1%

and only approximately a quarter of isolates having rates above 3% [33]. The lower *ex vivo*

rates may be partly due to lower assay sensitivity compared to the Pfs16 staining method, as

well as effects of plasma components such as variable lysophosphatidylcholine levels in

some individuals inhibiting parasite sexual commitment [33]. It is notable that the isolates in

the present study were cultured consistently in serum-free medium supplemented by

Albumax-II for several months, demonstrating that use of serum in culture is not required to

preserve the intrinsic capacity of parasites to show robust rates of sexual commitment over

this period. Nevertheless, parasites with very high rates of sexual commitment might have

been selected against, while eventual loss of sexual commitment in some cultures may occur

over longer periods of culture.

Most of the diverse long-term laboratory-adapted lines had mean gametocyte conversion

rates between 4.7% and 14.5% per cycle, and showed variation in conversion rate among

different biological replicates. The only line that always had zero sexual conversion was F12

with a previously described non-functional *ap2-g* gene due to a nonsense mutation [6]. All

other lines showed some detectable sexual conversion, although four lines had much lower

rates (means of 0.3 to 1.6%) than the rest, likely due to loss-of-function mutations acquired

in culture including deletion of the *gdv1* gene in lines T9/96 and D10 [36] which showed the

lowest conversion rates. Loss of *gdv1* in two other laboratory lines was reportedly

associated with no detectable gametocytes [12, 37], and it is possible that the low level of

conversion in T9/96 and D10 reflects a small amount of commitment to gametocyte

development post-invasion due to stochastic activation of *ap2-g*, independent of the main

pathway involving *gdv1* [15].

The multiple replicate measurements of each *P. falciparum* line enable the identification of

significant variation among recently derived clinical isolates from the same local endemic

population, as well as among long-term laboratory-adapted lines that have diverse

geographical origins. Given that modelling of data from experimental infections has

indicated much lower rates of approximately 1% or less per cycle [20, 21], there may be a

significant loss *in vivo* during the long period of development to circulating late stage

gametocytes, or suppression of gametocyte conversion in these infections compared to

culture.

Variation in gametocyte conversion rates among biological replicate assays was similar for

recently established clinical isolates and most of the long-term laboratory adapted lines.

Measured differences among replicates may reflect intrinsic parasite variation, stochastic or

due to determinants yet unidentified, or could be due to assay conditions. Previous studies

on a several different culture-adapted *P. falciparum* clones did not show as much variation

between experimental replicates [6, 13, 28, 31], which may be due to tighter parasite

synchronisation in those studies, whereas a broad window of synchronisation in the present

study was intended to be similar to what may be sampled in peripheral blood (in which ring

stage parasites may range in ages up to approximately 24 hours). The inter-replicate assay

variation in this study might be partly due to the possibility that a proportion of parasites in

some replicates were too young to stain with the Pf16 marker, although this was aimed to

be minimised by inspecting parasite development to determine the exact time of assay

completion. Although most parasites assayed would be mature enough for Pfs16 staining,

some replicates may have parasites at least 36 hours post invasion when all stage I

gametocytes would be Pfs16-positive [15, 17, 40], while other replicates may have parasites

24 hours post invasion when a majority but not all stage I gametocytes may already be

Pfs16-positive [15]. Exact standardisation of parasite ages in culture of diverse isolates is not

feasible, as isolates intrinsically vary in developmental cycle times [41, 42], but it is likely

that tighter synchronisation of parasite cultures would reduce variation among replicates for

any individual isolate. Assay precision may also be increased if detection assays are

developed to enable antibody staining of sexually committed parasites before they express

Pfs16, which may become possible as particular gene transcripts are expressed in committed

ring stages [43], and at least one protein (GEXP02) is expressed earlier than Pfs16 [17].

Although a majority of the diverse laboratory-adapted parasite lines tested in this study

retain an intact mechanism for gametocyte production, there were differences in responses

to choline. The presence or absence of choline had no effect in any of eight replicate

experiments on the T9/96 clone that has a culture-acquired deletion of *gdv1*, consistent

with the requirement of this locus for mediating a response to induce enhanced levels of

sexual commitment [12, 13]. In contrast, choline had a significant but variable effect on

sexual commitment in each of the other parasite lines tested, each of which also showed

substantial variation in responsiveness among biological replicates. Although a few

individual replicate assays showed greater than 10 fold effects, mean suppressive effects of

choline in each parasite line ranged from 1.5 fold (for 3D7) to 3.5 fold (for Dd2), more

moderate than described in previous studies of parasite clones maintained in medium

containing choline or serum before being switched to choline-free medium [13, 17, 31].

Gametocyte conversion rates of >30% have occasionally been reported elsewhere [28],

which were not seen in the non-genetically engineered lines here, but only with

overexpression of GDV1 in an engineered line. As all parasite lines including clinical isolates

in this study were maintained in a standard medium with Albumax without choline

supplementation, it is possible that higher gametocyte conversion rates were selected

against, as they might not be sustained in continuous culture without using conditions to

suppress conversion during long-term maintenance [31].

The results support the utility of choline in medium to suppress parasite sexual commitment

under controlled environmental conditions (10, 20, 21), and notably show significant

variation in the responsiveness of different parasite lines. It will be important to understand

if there is intraspecific variation in natural regulation of GDV1 expression, particularly the

control of gene-silencing antisense transcripts [13], and investigate the potential influence

of naturally-occurring cis-variation at the sub-telomeric locus on chromosome 9 [33, 39, 44]

distinct from the deletions disrupting function in some cultured lines [12, 36, 37, 45]. An

important new insight into the effect of available choline has just been reported, indicating

that sexual commitment is affected by competition between histone methyltransferases

(involved in *ap2-g* gene silencing) and phosphoethanolamine methyltransferase (involved in

de novo synthesis of phosphatidylcholine when choline availability is low) for the methyl

donor S-adenosylmethionine, which could have implications for future studies on inter-

isolate variation in rates of sexual commitment [32]. Investigation of parasites in natural

populations requires assay precision and experimental replication to be considered where

possible alongside statistical power from increased numbers of clinical samples, whether for

candidate gene investigations or larger scale scanning for associations with unlinked

genomic variation in parasite populations [46].

Materials and Methods

Ethics Approval

Approval for sampling of clinical isolates for culture experimental analysis was granted by

the Ethics committees of the Ghana Health Service, the Noguchi Memorial Institute for

Medical Research at the University of Ghana, the Navrongo Health Research Centre and the

London School of Hygiene and Tropical Medicine, as previously described for the collection

[35] from which specific isolates were selected in this analysis.

Plasmodium falciparum laboratory-adapted clones and clinical isolates

The genetically modified *P. falciparum* cloned line 3D7/iGP_D9 was used to compare the

sensitivity and precision of gametocyte conversion rate assays, as artificially elevated sexual

commitment is selectively inducible in this line by overexpression of GDV1 fused to a

destabilization domain (DD) which maintains activity in the presence of Shield-1 reagent

[34].

Thirteen different *P. falciparum* long-term laboratory-adapted lines that have not

undergone targeted genetic modification were studied. Each of these was either previously

derived by cloning or is a line that has an apparently monoclonal single-genome sequence,

and a putative origin based on the geographical location of the original isolation: NF54 (a

strain of 'airport malaria' from The Netherlands with genome sequence similarity to African

isolates), 3D7 (the *P. falciparum* reference genome strain originally cloned from NF54 and

maintained in culture for many years) [47], F12 (derived by cloning from cultured 3D7 and

having a nonsense mutant *ap2-g* gene) [6], D10 (Papua New Guinea), D6 (Sierra Leone), Dd2

(Southeast Asia), FCC2 (China), HB3 (Honduras), R033 (Ghana), Palo Alto (Uganda), 7G8

(Brazil), GB4 (Ghana) and T9/96 (Thailand). All lines were thawed from pre-existing frozen

stocks with distinct identities verified by previous targeted genotyping and sequencing [39,

48] and were not re-cloned prior to this study, and each line was cultured for at least one

week after thawing prior to any assay, all culture being conducted using a commercially-

prepared serum-free medium supplemented with lipid-rich bovine serum albumen (Gibco

RPMI with 0.5% AlbuMAX™ II lot number 2177881). This consistent medium preparation had

very minimal choline (less than 0.02 mM choline chloride) which would not suppress

parasite sexual commitment, unlike high concentrations of choline (2 mM as used for some

assays described in a separate section below) or serum which has variable components

between batches.

Six *P. falciparum* clinical isolates from Ghana with single-genome unmixed infections were

selected from a larger panel of infections sampled from children with clinical malaria in

Navrongo (in the Upper East Region of the country), in which parasite asexual multiplication

rate variation was previously studied [35]. These isolates were cultured with the same batch

of commercially-prepared serum-free medium with 0.5% AlbuMAX™ II as for the laboratory-

adapted parasite lines, ensuring the same batch was used throughout all experiments so

that no biological variation should be due to media components, and these clinical isolates

were not cultured with added serum at any time as they had been cultured with 0.5%

AlbuMAX™ II since isolation [35]. Clinical isolates with mixed genome-infections were

purposefully not selected for inclusion in this study, as their sexual commitment rates would

be complex to interpret due to potential changes in mixed genomic composition over time

[35].

Quantification of sexual commitment rates by measuring gametocyte conversion

Multiple biological replicate preparations of *P. falciparum* long-term laboratory-adapted

lines and more recently culture-established clinical isolates were each cultured in human

erythrocytes at 3% hematocrit in RPMI 1640 media supplemented with AlbuMAX™ II and

incubated at 37°C in atmospheric air with 5% CO_2. Prior to assay, each asynchronous

parasite culture was synchronized to within a 24-hour window of intra-erythrocytic asexual

stage development, intended to correspond approximately to the parasite age range that

would be typically sampled in peripheral blood of patients, using the following two-step

technique. Parasites containing haemozoin, predominantly schizont and late trophozoite

stages, were first positively selected on MACS® LD magnetic columns [49], and after culture

for 24 hours a second MACS purification step was performed from which the flow through

erythrocytes (containing predominantly ring stage parasites) were collected and put back

into culture, which then contained exclusively ring stage parasites as any gametocytes or

residual mature asexual parasites were removed on the column. This broad synchronization

scheme is illustrated in Supplementary Figure S1. For the 3D7/iGP parasite line the effect of

GDV1-DD overexpression was tested by adding Shield-1 reagent to this preparation of ring

stages (1 µM compared to none, with intermediate concentration treatments of 0.05, 0.15

and 0.5 µM Shield-1 also being tested in some assays), with Shield-1 being retained in the

culture throughout the cycle.

Two methods were used for measuring the gametocyte conversion rate:

Method 1. Counting of the proportion of stage I gametocytes compared to all parasites

developing in the first cycle post-invasion using anti-Pfs16 staining.

This method involves distinguishing very early gametocytes from asexual parasites by

detection of the gametocyte marker Pfs16 protein [15, 40, 50]. After two rounds of MACS

column purification to obtain ring stages followed by a further 24 hours of growth, when a

culture with a broad synchronicity containing schizonts was obtained, parasites were

allowed to invade erythrocytes and were cultured for a period of between 38 and 46 hours,

the exact time of harvesting of each culture being decided on the basis of microscopical

examination of Giemsa stained slides indicating that most post-invasion parasites were

clearly beyond the ring stage of development and therefore should be either trophozoites or

stage I gametocytes. The harvested post-invasion parasites were examined by fluorescence

microscopy by Pfs16 staining of early gametocytes and DAPI staining of all parasites. Stage I

gametocytes were counted as those staining positive for Pfs16 and showing typical round

morphology similar to trophozoites (whereas any Pfs16-positive parasites with

differentiated morphology of stage II gametocytes potentially residual from induction prior

to the previous cycle were not counted). The assay enables most sexually-developing

parasites to be counted in comparison to the total of post-invasion parasites, on the basis

that a majority of early stage I gametocytes express Pfs16 after 24 hours post invasion [15]

as a proportion of all gametocytes which eventually express [40], with only a minority of

sexual parasites likely not to be counted due to being too young. Parasite cultures harvested

for this purpose were washed in phosphate buffered saline (PBS) with 3% bovine serum

albumen (BSA), resuspended to 2% hematocrit, spotted onto multiwell slides (Hendley,

Essex, UK), air-dried and stored at -80°C. Prior to immunofluorescence staining, slides were

fixed with 4% paraformaldehyde in PBS for 30 min and permeabilized with 0.1% Triton X-100

in PBS for 10 min. The slides were incubated for 30 minutes at room temperature with a

monoclonal α-Pfs16 murine antibody 93A3A2 [50], diluted 1:2000 in PBS with 3% BSA. Alexa

Fluor® 594-conjugated anti-mouse IgG (Invitrogen, A11032) was used as secondary

antibody, diluted 1:1000 in PBS with 3% BSA and incubated for 30 minutes at room

temperature. Slides were mounted with coverslips using DAPI-containing Prolong™

Diamond antifade mountant (ThermoFisher Scientific), and parasite counting was performed

using a Zeiss CCD fluorescence microscope, with identical settings used for each experiment.

An average of approximately 1000 parasites were counted to determine proportions

expressing Pfs16 in each measurement, and where the density of parasites did not allow this

a quality control was performed to exclude any assay data that did not have a minimum of

at least 300 parasites counted.

Method 2. Counting of developing gametocytemia compared to all parasites at the ring

stage culture.

This method estimates the proportion of parasites that produce gametocytes developing to

morphological stage II, by comparing with the ring parasitaemia counted 4 days previously

at the beginning of the parasite developmental cycle. The technique is based on a protocol

that uses Giemsa staining and does not require immunofluorescence staining of an early

gametocyte marker [33]. To initiate this assay, after two MACS column purifications and

when a broadly synchronous culture containing a majority of schizonts were obtained,

parasites were allowed to re-invade fresh erythrocytes and cultured until ring stage

development was observed. Then, ring stage parasitaemia was assessed by visual

examination of thin blood smears of each culture to determine the proportion of

erythrocytes containing parasites (referred to as D0 parasitaemia), ensuring that at least

1,000 erythrocytes were counted. The cultures were then treated with 50 mM N-acetyl-D-

glucosamine (GlcNAc) to prevent further development of asexual parasites [51]. After 4

days, the stage II gametocytemia was assessed by counting of Giemsa-stained thin blood

smears (referred to as D4 gametocytemia), ensuring that at least 10,000 erythrocytes were

counted. The gametocyte conversion rate was calculated by dividing D4 stage II

gametocytemia (as a proportion of erythrocytes infected) by D0 ring parasitaemia (as a

proportion of erythrocytes infected), yielding a 'ratio-of-ratio' proportion with 95%

confidence intervals.

For both methods, all assays were set up at 3% haematocrit in a culture volume of 5 ml

(static in 6-well culture plates), with parasitaemia levels at the start of each replicate assay

being between 0.5 - 2.5% erythrocytes infected, to allow sufficient numbers of parasites for

high precision counting while minimising stress on parasites that may be induced if cultures

have higher parasitaemia.

Quantification of gametocyte conversion rate using culture media without choline

Parasites were prepared for assays in the same manner as described in the previous section,

except that they were split after the second MACS column purification (at ring stage in the

cycle before gametocyte counting) into separate wells with culture media (RPMI 1640 with

25 mM HEPES, 100 µM Hypoxanthine, 1 mM L-Glutamine, 0.39% fatty acid-free BSA, 30 mM

Palmitic acid, 30 mM Oleic acid) either supplemented with 2 mM choline or lacking choline,

based on a previously described protocol [31]. Parasites were allowed to grow until schizont

stage and to invade fresh erythrocytes, without changing the culture medium. After 38-46

hours post invasion, parasites were collected and fixed for analysis of gametocyte

conversion by differential counting using Pfs16 staining by immunofluorescence microscopy

as well as DAPI staining as described above for Method 1, with an average of approximately

1000 parasites being counted for each individual measurement. Between six and eight

replicate assays were performed for each parasite line, the effect within each replicate

calculated as the rate ratio (with 95% confidence intervals) of the gametocyte conversion

rate in choline-free medium compared with the conversion rate in medium containing 2 mM

choline.

Statistical analyses

Spearman's ρ (rho) coefficient was used to evaluate the strength and significance of the rank

correlation between results from the two different gametocyte conversion rate assays

performed in parallel. Statistical comparisons of gametocyte conversion rate variation

between different *P. falciparum* lines were performed using the Mann-Whitney test on the

multiple replicate assay measurements of each line. Each different parasite line was assayed

with a mean of more than six biological replicate cultures, most individual lines having at

least six and some up to nine biological replicates performed. Each individual biological

replicate data point had 95% confidence intervals calculated for the proportions of parasites

showing sexual commitment on the basis of the raw numerical counts performed, enabling

accurate estimation of within-assay sampling variation, and all count data are provided for

verification and to enable any future meta-analyses. For each parasite line, the mean effect

of modified culture media conditions was estimated by calculating the Mantel-Haenszel rate

ratio (RR_{M-H}) across biological replicate assays, based on a simple fixed-effects model.

Statistical analyses were performed using Prism version 9 and Epi-Info version 7 software.

Acknowledgements

This study was funded by Project Grant MR/S009760/1 from the UK Medical Research Council to DJC and DAB. Original sampling of clinical isolates was supported by a Royal Society Africa Award AA110050 to GAA and DJC. TSV received funding from the Swiss National Science Foundation grant BSCGI0_157729. We are grateful to all colleagues who supported the research environment, including sharing of laboratory facilities and equipment, enabling continuous parasite culture and experimentation as well as storage and shipment of samples and reagents.

Figure Legends

Figure 1. Comparison of the sensitivity and precision of two different methods for measuring *P. falciparum* sexual commitment in terms of gametocyte conversion rates. Multiple paired replicate assays were performed with the parasite clone 3D7/iGP_D9 [34] which allows inducible overexpression of GDV1 fused to a destabilization domain (DD) stabilized in the presence of Shield-1 reagent to provide controlled elevation of sexual commitment (closed circles indicate GDV1–ON, open circles GDV1-OFF). Both methods show the expected significant elevation in gametocyte conversion in the presence of Shield-1 reagent in all individual assays (asterisks indicate P<0.0001 for each of the individual replicate assays). In each panel, the green dotted line denotes the mean for GDV1-ON and the red dotted line for GDV1-OFF. **A.** The gametocyte conversion rate (with 95% CI) for each of seven independent experiments using Method 1 involving staining of the gametocyte marker Pfs16, yielding a mean conversion rate of 38.3 % under GDV1-ON and 6.3% under GDV1-OFF conditions **B.** The gametocyte conversion rate (with 95% CI) for each of four experiments using Method 2 which involves a comparison of Day 4 (D4) gametocyte parasitaemia with D0 ring-stage parasitaemia (assays paired with the first four independent experiments shown for Method 1), yielding a mean conversion rate of 15.9 % under GDV1-ON and 4.1 % under GDV1-OFF conditions. Exact parasite and erythrocyte counts and statistical analyses for all assays are detailed in Supplementary Table S1.

Figure 2. Significant correlation in the estimates of gametocyte conversion rate obtained by two different assay methods. Paired assays were performed using Method 1 (involving Pfs16 staining to discriminate sexually from asexually-developing parasites) and Method 2 (D0D4, involving counts of parasites 4 days apart) on multiple biological replicates of 3D7/iGP_D9 with a range of concentrations of Shield-1 reagent (N=17 paired assays in total, details in Supplementary Table S2). This highly significant correlation confirms that either method gives valid measurements if used alone, whereas Method 1 is more sensitive (as indicted here and in Figure 1). Method 1 is also more precise as it involves calculation of a single numerical ratio, rather than a ratio-of-ratios based on two different counts performed 4 days apart which is highly vulnerable to effects of small numbers in either count. For visual clarity, the individual assay 95% confidence intervals are not shown on the plot, but these and numerical data of all assays are given in Supplementary Table S2.

Figure 3. Gametocyte conversion rates of six recently culture-established Ghanaian clinical

isolates. These were selected for study as they are unrelated single genotype infections,

chosen from a larger number of samples from a highly endemic population that contained

many multiple genotype infections [35]. **A.** For each isolate, a minimum of six biological

replicate assays was performed, and the estimate from each replicate with 95% confidence

intervals (based on numbers of parasites counted) is plotted, while the red dotted line

represents the mean across replicate assays. There were significant differences in pairwise

comparisons (Mann-Whitney tests, Supplementary Table S4), isolate 296 having a

significantly higher conversion rate than three others (isolates 272, 289 and 292), and isolate

289 having a significantly lower rate than three others (292, 293 and 296)(Supplementary

Table S4). There were no significant correlations with length of time in culture, or batches of

erythrocytes used for the different replicate assays (these details and all numerical data

values are given in Supplementary Table S3). **B.** For each isolate the gametocyte conversion

rate of individual replicates is plotted in relation to the culture parasitaemia in the previous

cycle (between 0.5 and 2.5% for all experimental replicates). 95% confidence intervals

individual conversion rate points are not shown as they are the same as in panel A. There

was no systematic trend towards positive or negative correlation, except a marginally

significant positive correlation for isolate 289 ($P = 0.03$) which may indicate slight

enhancement at higher parasitaemia or a chance finding due to multiple comparisons.

Figure 4. Gametocyte conversion rates of 13 long-term culture adapted *P. falciparum* lines

of diverse geographical origins. Each different parasite line was tested with multiple

independent biological replicate assays, most individual lines having at least six replicates.

All counts and calculated values for all replicates are given in full in Supplementary Table S5.

A. For each parasite line, the conversion rates are shown for each of the biological replicates

and the horizontal line shows the mean across all biological replicates, with a spectrum of

variation among the different lines, means ranging from 0% to 14.5%. The background with

different shades of grey show that the conversion rates separate into three different groups,

based on pairwise comparisons of all replicates between the parasite lines (Mann-Whitney

tests, Supplementary Table S6): One line (F12) showed no conversion, four lines (D10, T9/96,

Palo Alto, and 3D7) showed consistently low conversion (means between 0.3% and 1.6%),

and eight lines (D6, GB4, Dd2, HB3, NF54, RO33, 7G8, and FCC2) showed higher levels of

conversion (means between 4.7% and 14.5%). **B.** Variation in the gametocyte conversion

rates among independent replicate assays for each line is shown in more detail by including

95% CI of the counts for each assay of each line, except F12 for which there was zero

conversion in all assays. Identical methods were used in assaying these parasite lines as for

the clinical isolates shown in Figure 3, assays being conducted in parallel under the same

conditions in the same laboratory. Erythrocytes from different erythrocyte donors used at

various times throughout the series of experiments did not determine the variation between

assay replicates (Supplementary Table S5).

Figure 5. Significant variation in sensitivity of different parasite lines to choline in media to

suppress sexual commitment. Closed circles represent biological replicates cultured in the

presence of 2 mM choline, open circles represent biological replicates grown in the absence

of choline. T9/96 shows no response to choline in any of seven biological replicate assays,

whereas each of the other cultured lines show responses that vary among biological

replicates. Significant differences between paired measurements with and without choline

in individual replicate assays for 3D7, Dd2, HB3 and NF54 are highlighted with asterisks

(Fisher's Exact Test * P < 0.05, ** P <0.01, *** P<0.001). Across all replicates, estimated by

Mantel-Haenszel rate ratios the mean effects of choline were greatest for Dd2 (3.5 fold, 95%

CI 3.0-4.1 fold), intermediate for NF54 (2.4 fold, 95% CI 2.0-3.0 fold) and HB3 (2.0 fold, 95%

CI 1.6-2.5 fold), and least for 3D7 (1.5 fold, 95% CI 1.2-1.9 fold). All individual counts and

assay values for all biological replicates are given in Supplementary Table S7.

References

1.   McKenzie FE, Bossert WH. The optimal production of gametocytes by *Plasmodium falciparum*. J Theor Biol. 1998;193:419-28. Epub 1998/09/15. doi: 10.1006/jtbi.1998.0710. PMID: 9735270.

2.   Schneider P, Greischar MA, Birget PLG, Repton C, Mideo N, Reece SE. Adaptive plasticity in the gametocyte conversion rate of malaria parasites. PLoS Pathog. 2018;14:e1007371. Epub 2018/11/15. doi: 10.1371/journal.ppat.1007371. PMID: 30427935.

3.   WHO. World Malaria Report 2020. Geneva: 2020.

4.   Barry A, Bradley J, Stone W, Guelbeogo MW, Lanke K, Ouedraogo A, et al. Higher gametocyte production and mosquito infectivity in chronic compared to incident Plasmodium falciparum infections. Nat Commun. 2021;12:2443. Epub 2021/04/28. doi: 10.1038/s41467-021-22573-7. PMID: 33903595.

5.   Alano P, Carter R. Sexual differentiation in malaria parasites. Annu Rev Microbiol. 1990;44:429-49. doi: 10.1146/annurev.mi.44.100190.002241. PMID: 2252389.

6.   Kafsack BF, Rovira-Graells N, Clark TG, Bancells C, Crowley VM, Campino SG, et al. A transcriptional switch underlies commitment to sexual development in malaria parasites. Nature. 2014;507:248-52. doi: 10.1038/nature12920. PMID: 24572369.

7.   Kent RS, Modrzynska KK, Cameron R, Philip N, Billker O, Waters AP. Inducible developmental reprogramming redefines commitment to sexual development in the malaria parasite Plasmodium berghei. Nat Microbiol. 2018;3:1206-13. Epub 2018/09/05. doi: 10.1038/s41564-018-0223-6. PMID: 30177743.

8.   Sinha A, Hughes KR, Modrzynska KK, Otto TD, Pfander C, Dickens NJ, et al. A cascade of DNA-binding proteins for sexual commitment and development in *Plasmodium*. Nature. 2014;507:253-7. doi: 10.1038/nature12970. PMID: 24572359.

9.   Josling GA, Russell TJ, Venezia J, Orchard L, van Biljon R, Painter HJ, et al. Dissecting the role of PfAP2-G in malaria gametocytogenesis. Nat Commun. 2020;11:1503. Epub 2020/03/22. doi: 10.1038/s41467-020-15026-0. PMID: 32198457.

10.   Rovira-Graells N, Gupta AP, Planet E, Crowley VM, Mok S, Ribas de Pouplana L, et al. Transcriptional variation in the malaria parasite *Plasmodium falciparum*. Genome Res. 2012;22:925-38. Epub 2012/03/15. doi: 10.1101/gr.129692.111. PMID: 22415456.

11.   Brancucci NMB, Bertschi NL, Zhu L, Niederwieser I, Chin WH, Wampfler R, et al. Heterochromatin protein 1 secures survival and transmission of malaria parasites. Cell Host Microbe. 2014;16:165-76. doi: 10.1016/j.chom.2014.07.004. PMID: 25121746.

12.   Eksi S, Morahan BJ, Haile Y, Furuya T, Jiang H, Ali O, et al. *Plasmodium falciparum* gametocyte development 1 (Pfgdv1) and gametocytogenesis early gene identification and commitment to sexual development. PLoS Pathog. 2012;8:e1002964. Epub 2012/10/25. doi: 10.1371/journal.ppat.1002964. PMID: 23093935.

13.   Filarsky M, Fraschka SA, Niederwieser I, Brancucci NMB, Carrington E, Carrio E, et al. GDV1 induces sexual commitment of malaria parasites by antagonizing HP1-dependent gene silencing. Science. 2018;359:1259-63. doi: 10.1126/science.aan6042. PMID: 29590075.

14.   Bruce MC, Alano P, Duthie S, Carter R. Commitment of the malaria parasite *Plasmodium falciparum* to sexual and asexual development. Parasitology. 1990;100 Pt 2:191-200. Epub 1990/04/01. doi: 10.1017/s0031182000061199. PMID: 2189114.

15.   Bancells C, Llora-Batlle O, Poran A, Notzel C, Rovira-Graells N, Elemento O, et al. Revisiting the initial steps of sexual development in the malaria parasite *Plasmodium falciparum*. Nat Microbiol. 2019;4:144-54. Epub 2018/11/28. doi: 10.1038/s41564-018-0291-7. PMID: 30478286.

16.   Llora-Batlle O, Michel-Todo L, Witmer K, Toda H, Fernandez-Becerra C, Baum J, et al. Conditional expression of PfAP2-G for controlled massive sexual conversion in *Plasmodium falciparum*. Sci Adv. 2020;6:eaaz5057. Epub 20200610. doi: 10.1126/sciadv.aaz5057. PMID: 32577509.

17.   Portugaliza HP, Llora-Batlle O, Rosanas-Urgell A, Cortes A. Reporter lines based on the gexp02 promoter enable early quantification of sexual conversion rates in the malaria parasite

Plasmodium falciparum. Sci Rep. 2019;9:14595. Epub 2019/10/12. doi: 10.1038/s41598-019-50768-
y. PMID: 31601834.
18.   Koepfli C, Yan G. *Plasmodium* gametocytes in field studies: Do we measure commitment to
transmission or detectability? Trends Parasitol. 2018;34:378-87. Epub 2018/03/17. doi:
10.1016/j.pt.2018.02.009. PMID: 29544966.
19.   Carter LM, Kafsack BF, Llinas M, Mideo N, Pollitt LC, Reece SE. Stress and sex in malaria
parasites: Why does commitment vary? Evolution, Medicine, and Public Health. 2013;2013:135-47.
doi: 10.1093/emph/eot011. PMID: 24481194.
20.   Cao P, Collins KA, Zaloumis S, Wattanakul T, Tarning J, Simpson JA, et al. Modeling the
dynamics of *Plasmodium falciparum* gametocytes in humans during malaria infection. Elife. 2019;8.
Epub 2019/10/30. doi: 10.7554/eLife.49058. PMID: 31658944.
21.   Eichner M, Diebner HH, Molineaux L, Collins WE, Jeffery GM, Dietz K. Genesis, sequestration
and survival of *Plasmodium falciparum* gametocytes: parameter estimates from fitting a model to
malariatherapy data. Trans R Soc Trop Med Hyg. 2001;95:497-501. Epub 2001/11/15. doi:
10.1016/s0035-9203(01)90016-1. PMID: 11706658.
22.   Carter R, Miller LH. Evidence for environmental modulation of gametocytogenesis in
Plasmodium falciparum in continuous culture. Bull World Health Organ. 1979;57 Suppl 1:37-52. Epub
1979/01/01. PMID: 397008.
23.   Graves PM, Carter R, McNeill KM. Gametocyte production in cloned lines of *Plasmodium*
falciparum. Am J Trop Med Hyg. 1984;33:1045-50. doi: 10.4269/ajtmh.1984.33.1045. PMID:
6391217.
24.   Buckling A, Ranford-Cartwright LC, Miles A, Read AF. Chloroquine increases *Plasmodium*
falciparum gametocytogenesis in vitro. Parasitology. 1999;118:339-46. Epub 1999/05/26. doi:
10.1017/s0031182099003960. PMID: 10340323.
25.   Neveu G, Beri D, Kafsack BF. Metabolic regulation of sexual commitment in *Plasmodium*
falciparum. Curr Opin Microbiol. 2020;58:93-8. Epub 2020/10/15. doi: 10.1016/j.mib.2020.09.004.
PMID: 33053503.
26.   Beri D, Balan B, Chaubey S, Subramaniam S, Surendra B, Tatu U. A disrupted
transsulphuration pathway results in accumulation of redox metabolites and induction of
gametocytogenesis in malaria. Sci Rep. 2017;7:40213. Epub 2017/01/17. doi: 10.1038/srep40213.
PMID: 28091526.
27.   Chaubey S, Grover M, Tatu U. Endoplasmic reticulum stress triggers gametocytogenesis in
the malaria parasite. J Biol Chem. 2014;289:16662-74. Epub 2014/04/24. doi:
10.1074/jbc.M114.551549. PMID: 24755215.
28.   Portugaliza HP, Miyazaki S, Geurten FJ, Pell C, Rosanas-Urgell A, Janse CJ, et al. Artemisinin
exposure at the ring or trophozoite stage impacts *Plasmodium falciparum* sexual conversion
differently. Elife. 2020;9. Epub 2020/10/22. doi: 10.7554/eLife.60058. PMID: 33084568.
29.   Fivelman QL, McRobert L, Sharp S, Taylor CJ, Saeed M, Swales CA, et al. Improved
synchronous production of *Plasmodium falciparum* gametocytes in vitro. Mol Biochem Parasitol.
2007;154:119-23. Epub 20070420. doi: 10.1016/j.molbiopara.2007.04.008. PMID: 17521751.
30.   Mantel PY, Hoang AN, Goldowitz I, Potashnikova D, Hamza B, Vorobjev I, et al. Malaria-
infected erythrocyte-derived microvesicles mediate cellular communication within the parasite
population and with the host immune system. Cell Host Microbe. 2013;13:521-34. Epub 2013/05/21.
doi: 10.1016/j.chom.2013.04.009. PMID: 23684304.
31.   Brancucci NMB, Gerdt JP, Wang C, De Niz M, Philip N, Adapa SR, et al.
Lysophosphatidylcholine regulates sexual stage differentiation in the human malaria parasite
Plasmodium falciparum. Cell. 2017;171:1532-44 e15. Epub 2017/11/14. doi:
10.1016/j.cell.2017.10.020. PMID: 29129376.
32.   Harris CT, Tong X, Campelo R, Vanheer LN, Marreiros IM, Nahiyaan N, et al. Metabolic
competition between lipid metabolism and histone methylation regulates sexual differentiation in
human malaria parasites. BioRxiv. 2022. doi: 10.1101/2022.01.18.476397.
33.   Usui M, Prajapati SK, Ayanful-Torgby R, Acquah FK, Cudjoe E, Kakaney C, et al. *Plasmodium*
falciparum sexual differentiation in malaria patients is associated with host factors and GDV1-

dependent genes. Nat Commun. 2019;10:2140. Epub 2019/05/16. doi: 10.1038/s41467-019-10172-
6. PMID: 31086187.
34. Boltryk SD, Passecker A, Alder A, Carrington E, van de Vegte-Bolmer M, van Gemert GJ, et al.
CRISPR/Cas9-engineered inducible gametocyte producer lines as a valuable tool for Plasmodium
falciparum malaria transmission research. Nat Commun. 2021;12:4806. Epub 2021/08/12. doi:
10.1038/s41467-021-24954-4. PMID: 34376675.
35. Stewart LB, Diaz-Ingelmo O, Claessens A, Abugri J, Pearson RD, Goncalves S, et al. Intrinsic
multiplication rate variation and plasticity of human blood stage malaria parasites. Commun Biol.
2020;3:624. Epub 2020/10/30. doi: 10.1038/s42003-020-01349-7. PMID: 33116247.
36. Cheeseman IH, Gomez-Escobar N, Carret CK, Ivens A, Stewart LB, Tetteh KK, et al. Gene copy
number variation throughout the *Plasmodium falciparum* genome. BMC Genomics. 2009;10:353.
PMID: 19653891.
37. Tiburcio M, Hitz E, Niederwieser I, Kelly G, Davies H, Doerig C, et al. A 39-amino-acid C-
Terminal truncation of GDV1 disrupts sexual commitment in *Plasmodium falciparum*. Msphere.
2021;6. doi: 10.1128/mSphere.01093-20. PMID: WOS:000692590000015.
38. Scherf A, Mattei D. Cloning and characterization of chromosome breakpoints of *Plasmodium
falciparum*: breakage and new telomere formation occurs frequently and randomly in subtelomeric
genes. Nucleic Acids Res. 1992;20:1491-6. Epub 1992/04/11. doi: 10.1093/nar/20.7.1491. PMID:
1579440.
39. Duffy CW, Amambua-Ngwa A, Ahouidi AD, Diakite M, Awandare GA, Ba H, et al. Multi-
population genomic analysis of malaria parasites indicates local selection and differentiation at the
gdv1 locus regulating sexual development. Sci Rep. 2018;8:15763. Epub 2018/10/27. doi:
10.1038/s41598-018-34078-3. PMID: 30361631.
40. Bruce MC, Carter RN, Nakamura K, Aikawa M, Carter R. Cellular location and temporal
expression of the *Plasmodium falciparum* sexual stage antigen Pfs16. Mol Biochem Parasitol.
1994;65:11-22. Epub 1994/05/01. doi: 10.1016/0166-6851(94)90111-2. PMID: 7935618.
41. Reilly Ayala HB, Wacker MA, Siwo G, Ferdig MT. Quantitative trait loci mapping reveals
candidate pathways regulating cell cycle duration in *Plasmodium falciparum*. BMC Genomics.
2010;11:577. doi: 10.1186/1471-2164-11-577. PMID: 20955606.
42. Smith LM, Motta FC, Chopra G, Moch JK, Nerem RR, Cummins B, et al. An intrinsic oscillator
drives the blood stage cycle of the malaria parasite Plasmodium falciparum. Science. 2020;368:754-
9. Epub 2020/05/16. doi: 10.1126/science.aba4357. PMID: 32409472.
43. Prajapati SK, Ayanful-Torgby R, Pava Z, Barbeau MC, Acquah FK, Cudjoe E, et al. The
transcriptome of circulating sexually committed *Plasmodium falciparum* ring stage parasites
forecasts malaria transmission potential. Nat Commun. 2020;11:6159. Epub 2020/12/04. doi:
10.1038/s41467-020-19988-z. PMID: 33268801.
44. Rono MK, Nyonda MA, Simam JJ, Ngoi JM, Mok S, Kortok MM, et al. Adaptation of
Plasmodium falciparum to its transmission environment. Nat Ecol Evol. 2018;2:377-87. doi:
10.1038/s41559-017-0419-9. PMID: 29255304.
45. Day KP, Karamalis F, Thompson J, Barnes DA, Peterson C, Brown H, et al. Genes necessary for
expression of a virulence determinant and for transmission of *Plasmodium falciparum* are located on
a 0.3-megabase region of chromosome 9. Proc Natl Acad Sci U S A. 1993;90:8292-6. PMID: 8367496.
46. MalariaGen, Ahouidi A, Ali M, Almagro-Garcia J, Amambua-Ngwa A, Amaratunga C, et al. An
open dataset of *Plasmodium falciparum* genome variation in 7,000 worldwide samples. Wellcome
Open Res. 2021;6:42. Epub 2021/07/30. doi: 10.12688/wellcomeopenres.16168.2. PMID: 33824913.
47. Bohme U, Otto TD, Sanders M, Newbold CI, Berriman M. Progression of the canonical
reference malaria parasite genome from 2002-2019. Wellcome Open Res. 2019;4:58. Epub
2019/06/06. doi: 10.12688/wellcomeopenres.15194.2. PMID: 31080894.
48. Tetteh KK, Stewart LB, Ochola LI, Amambua-Ngwa A, Thomas AW, Marsh K, et al. Prospective
identification of malaria parasite genes under balancing selection. PLoS One. 2009;4:e5568. PMID:
19440377.

49.   Ribaut C, Berry A, Chevalley S, Reybier K, Morlais I, Parzy D, et al. Concentration and
purification by magnetic separation of the erythrocytic stages of all human *Plasmodium* species.
Malar J. 2008;7:45. PMID: 18321384.
50.   Baker DA, Daramola O, McCrossan MV, Harmer J, Targett GA. Subcellular localization of
Pfs16, a *Plasmodium falciparum* gametocyte antigen. Parasitology. 1994;108 129-37. Epub
1994/02/01. doi: 10.1017/s0031182000068219. PMID: 8159458.
51.   Gupta SK, Schulman S, Vanderberg JP. Stage-dependent toxicity of N-acetyl-glucosamine to
Plasmodium falciparum. J Protozool. 1985;32:91-5. Epub 1985/02/01. doi: 10.1111/j.1550-
7408.1985.tb03020.x. PMID: 3886901.

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Supplementary Table S1. Gametocyte conversion rate (GCR) estimates (with 95%
confidence intervals) and all numerical count data for two different methods with multiple
experimental replicates applied to *P. falciparum* line 3D7/iGP under elevated GDV1
expression with Shield-1 reagent (+SHLD) and control (-SHLD) conditions.

Replicate	Condition	Method 1			Method 2		
		Pfs16 +ve	DAPI +ve	Pfs16 GCR (95%CI)	D0 % Parasit.	D4 % Parasit.	D0D4 GCR (95% CI)
1	-SHLD	84	921	9.12 (7.4 - 11.2)	10.1 (101/999)	0.3 (26/8262)	3.4 (2.2 -5.2)
	+SHLD	474	826	57.4 (54.0 - 60.7)	9.8 (98/999)	1.9 (117/6124)	20.9 (16.0 - 27.3)
2	-SHLD	104	1171	8.9 (7.4 - 10.7)	7.2 (67/932)	0.2 (15/8210)	2.7 (1.6 - 4.7)
	+SHLD	262	978	26.8 (24.1 - 29.7)	3.0 (29/970)	0.4 (49/13415)	12.5 (7.9 - 19.8)
3	-SHLD	41	926	4.4 (3.3 - 6.0)	5.6 (53/946)	0.1 (8/9305)	1.6 (2.8 - 3.4)
	+SHLD	150	595	25.2 (21.9 - 28.9)	4.4 (42/957)	0.4 (36/8625)	9.9 (6.3 - 15.4)
4	-SHLD	47	1043	4.5 (3.4 - 6.0)	5.0 (48/951)	0.4 (30/7278)	8.5 (5.4 - 13.5)
	+SHLD	267	701	38.1 (34.6 - 41.7)	4.4 (42/952)	1.0 (77/8070)	20.1 (14.0 - 28.8)

Supplementary Table S2. Estimates of gametocyte conversion rate (GCR) for 17 assays of
the *P. falciparum* inducible 3D7/iGP_D9 line in which two different methods were
performed in parallel for comparison (as detailed in Materials and Methods). Varying
concentrations of Shield-1 (0.055 – 1.0 µM, and none) were used for the assays. Data are
plotted and analysed in Figure 2, showing an overall correlation between the methods, with
Method 1 being generally more sensitive (and more precise with narrower confidence
intervals as shown here).

Independent assays of *P. falciparum* line	Method 1 Pfs16 GCR (95%CI)	Method 2 D0/D4 GCR (95%CI)
iGP - SHLD	9.1 (7.4 - 11.2)	3.4 (2.2 - 5.3)
iGP +SHLD (1.0)	57.4 (54.0 - 60.7)	20.9 (16.0 - 27.31)
iGP - SHLD	8.9 (7.4 - 10.7)	2.7 (1.6 - 4.8)
iGP + SHLD (0.055)	14.2 (12.3 - 16.4)	4.0 (2.2 - 7.2)
iGP + SHLD (0.15)	16.8 (14.8 - 19.1)	2.4 (1.2 (4.9)
iGP + SHLD (0.5)	17.4 (15.3 - 19.7)	2.6 (1.3 - 5.2)
iGP +SHLD (1.0)	26.8 (24.1 - 30.0)	12.5 (8.0 - 19.8)
iGP - SHLD	4.4 (3.4 - 7.0)	1.6 (0.8 - 3.4)
iGP + SHLD (0.055)	20.5 (17.9 - 23.4)	7.2 (4.6 - 11.3)
iGP + SHLD (0.15)	16.1 (13.8 - 18.7)	4.7 (3.2 - 6.9)
iGP + SHLD (0.5)	26.5 (23.5 - 29.6)	5.1 (3.2 - 8.2)
iGP +SHLD (1.0)	25.2 (21.9 - 28.9)	9.9 (6.4 - 15.4)
iGP - SHLD	4.5 (3.4 - 6.9)	8.5 (5.4 - 13.5)
iGP + SHLD (0.055)	8.9 (7.2 - 11.2)	10.7 (7.3 - 15.7)
iGP + SHLD (0.15)	22.3 (19.2 - 25.8)	8.9 (6.1 - 12.8)
iGP + SHLD (0.5)	37.4 (34.0 - 41.0)	17.5 (12.9 - 23.8)
iGP +SHLD (1.0)	38.1 (34.6 - 41.7)	20.1 (14.0 - 28.8)

808 **Supplementary Table S3.** Multiple biological replicate assays of gametocyte conversion
809 rates (GCR) for each of 6 cultured *P. falciparum* clinical isolates from Navrongo, Ghana.

Isolate	Replicate	RBC Batch	Days in culture	% Parasit.	Pfs16 +ve	DAPI +ve	GCR (95%CI)
	1	12	92	1.4	38	961	4.0 (2.9 - 5.4)
	2	12	99	0.7	9	870	1.0 (0.6 - 2.0)
289	3	12	99	0.9	23	740	3.1 (2.1 - 4.6)
	4	13	106	0.9	53	946	5.6 (4.3 - 7.3)
	5	13	106	0.8	28	971	2.9 (2.0 - 4.1)
	6	15	125	0.8	28	939	3.0 (2.1 - 4.3)
	Mean (SD)						*3.3 (SD = 1.5)*
	1	12	83	1.0	48	951	5.0 (3.8 - 6.6)
	2	12	83	0.8	45	954	4.7 (3.5 - 6.3)
	3	13	91	1.2	34	965	3.5 (2.5 - 4.9)
	4	13	91	0.6	26	973	2.7 (1.8 - 3.9)
	5	15	110	0.6	69	999	6.9 (5.5 - 8.7)
272	6	15	110	0.6	51	880	5.8 (4.4 - 7.5)
	Mean (SD)						*4.8 (SD = 1.5)*
	1	1	67	0.8	66	677	9.7 (7.7 - 12.2)
	2	12	62	0.9	34	965	3.5 (2.5 - 4.9)
	3	12	62	0.7	58	882	6.6 (5.1 - 8.4)
292	4	13	69	2.3	63	936	6.7 (5.3 - 8.5)
	5	13	69	1.1	46	953	4.8 (3.6 - 6.4)
	6	15	88	0.5	49	939	5.2 (4.0 - 6.8)
	7	15	88	0.8	19	564	3.4 (2.2 - 5.2)
	Mean (SD)						*5.7 (SD = 2.2)*
	1	1	67	1.7	109	1841	5.9 (4.9 - 7.1)
	2	12	55	0.8	120	878	13.7 (11.6 - 16.1)
293	3	13	69	0.9	118	881	13.4 (11.3 - 15.8)
	4	13	69	1.3	118	883	13.4 (11.3 - 15.8)
	5	13	76	1.0	39	962	4.1 (3.0 - 5.5)
	6	13	76	1.0	41	617	6.6 (4.9 - 8.9)
	Mean (SD)						*9.5 (SD = 4.4)*
	1	12	48	1.5	116	883	13.1 (11.1 - 15.5)
	2	12	55	1.0	31	968	3.2 (2.3 - 4.5)
	3	12	55	0.6	65	632	10.3 (8.2 - 12.9)
296	4	13	62	1.2	94	905	10.4 (8.6 - 12.5)
	5	13	62	1.1	84	915	9.2 (7.5 - 11.2)
	6	13	69	1.0	93	906	10.3 (8.5 - 12.4)
	7	13	69	1.0	118	887	13.3 (11.2 - 15.7)
	Mean (SD)						*10.0 (SD = 3.4)*
	1	1	55	1.0	156	1406	11.1 (9.6 - 12.9)
	2	2	64	2.2	103	360	28.6 (24.2 - 33.5)
	3	2	64	1.5	209	794	26.3 (23.5 - 29.6)
	4	3	71	0.7	131	907	14.4 (12.3 - 16.9)
280	5	3	71	1.0	109	787	13.9 (11.6 - 16.4)
	6	3	78	1.4	27	973	2.8 (1.9 - 4.0)
	7	3	78	0.6	49	884	5.5 (4.2 - 7.3)
	8	4	85	1.4	12	998	1.2 (0.7 - 2.1)
	9	4	85	0.6	56	943	5.9 (4.6 - 7.6)
	Mean (SD)						*12.2 (SD = 9.8)*

The table shows exact counts and 95% confidence intervals of the GCR estimates using
Method 1 (with Pfs16 staining) as plotted and analysed in Figure 3. The number of days in
culture is shown as well as the % parasitaemia measured in the cycle before. The assay
replicates are presented in the order in which they were performed (the duration of time in
culture occasionally does not exactly follow the same sequential temporal order for Isolates
292 and 293 as an intermediate cryopreservation and thawing of the isolates was
performed). Erythrocytes from 15 different erythrocyte donors used at various times
throughout the series of experiments shown here and in Supplementary Table S5 are
numbered (RBC Batch), and inspection of the data indicates that these did not determine
the variation between assay replicates.

Supplementary Table S4. Pairwise tests for significant differences in gametocyte conversion rates between different Ghanaian *P. falciparum* clinical isolates.

Clinical Isolates	272	280	289	292	293	296
272						
280	0.1447					
289	0.1797	0.0663				
292	0.5594	0.2523	0.0350*			
293	0.0649	0.7756	0.0043**	0.1375		
296	0.0221*	0.7577	0.0047**	0.0379*	0.7308	

P values are from Mann-Whitney comparisons of all of the individual biological replicate measures for each of the six isolates (as shown in Supplementary Table S3). Statistically significant comparisons are highlighted in green.

826 **Supplementary Table S5.** Multiple biological replicate assays of gametocyte conversion
827 rates (GCR) for each of 13 long-term culture adapted *P. falciparum* lines of diverse origins.

Line	Replicate	RBC batch	Pfs16 +ve	DAPI +ve	GCR (95% CI)
	1	3	0	1000	0 (0)
	2	3	0	1000	0 (0)
	3	4	0	1023	0 (0)
F12	4	10	0	394	0 (0)
	5	11	0	600	0 (0)
	6	11	0	711	0 (0)
	7	11	0	380	0 (0)
	Mean (SD)				0 (SD = 0)
	1	2	0	515	0 (0)
	2	2	1	999	0.1 (0 - 5.6)
	3	2	10	1177	0.8 (0.5 - 1.6)
D10	4	3	2	997	0.2 (0.1 - 0.7)
	5	3	2	1000	0.2 (0.1 -0.7)
	6	9	3	997	0.3 (0.1 - 0.9)
	7	9	2	997	0.2 (01 - 0.7)
	Mean (SD)				0.26 (SD = 0.27)
	1	12	6	993	0.6 (0.3 - 1.3)
	2	13	2	733	0.3 (0.1 - 1.0)
	3	13	3	992	0.3 (0.1 - 0.9)
T996	4	13	2	703	0.3 (0.1 - 1.0)
	5	14	2	997	0.2 (0 - 0.7)
	6	14	3	433	0.7 (0.2 - 2.0)
	Mean (SD)				0.39 (SD = 0.20)
	1	6	12	987	1.2 (0.7 - 1.2)
	2	7	14	514	2.7 (1.6 - 4.5)
	3	7	5	321	1.6 (0.7 - 3.6)
Palo Alto	4	9	1	998	0.1 (0 - 0.6)
	5	9	3	996	0.3 (0.1 - 0.9)
	6	9	1	998	0.1 (0 - 0.6)
	Mean (SD)				1.00 (SD = 1.04)
	1	2	20	980	2.0 (1.3 - 3.1)
	2	3	3	994	0.3 (0.1 - 0.9)
	3	3	12	979	1.2 (0.7 -2.1)
3D7	4	4	20	977	2.0 (1.3 -3.1)
	5	4	17	916	1.9 (1.2 - 3.0)
	6	12	20	885	2.3 (1.5 - 3.5)
	7	12	16	983	1.6 (1.0 - 2.6)
	Mean (SD)				1.62 (SD = 0.67)

Line	Replicate	RBC batch	Pfs16 +ve	DAPI +ve	GCR (95% CI)
	1	5	83	920	9.0 (7.3 -11.1)
	2	5	16	983	1.6 (1.0 - 2.6)
	3	5	6	885	0.7 (0.3 - 1.5)
D6	4	9	28	559	5.0 (3.5 - 7.1)
	5	9	86	905	9.5 (7.8 - 11.6)
	6	9	26	973	2.7 (1.8 - 3.9)
	7	9	50	950	5.3 (4.0 - 6.9)
	Mean (SD)				4.70 (SD = 3.36)
	1	6	22	978	2.2 (1.5 - 3.4)
	2	6	85	914	9.3 (7.6 -11.4)
	3	6	92	917	10.0 (8.3 - 12.2)
GB4	4	6	37	964	3.8 (2.8 - 5.3)
	5	7	59	940	6.3 (4.9 - 8.0)
	6	7	16	983	1.6 (1.0 - 2.6)
	7	9	34	969	3.5 (2.5 - 4.9)
	Mean (SD)				5.26 (3.35)
	1	2	130	902	14.4 (12.3 - 16.9)
	2	3	4	995	0.4 (0.2 - 1.0)
	3	3	9	986	0.9 (0.5 - 1.7)
Dd2	4	4	46	953	4.8 (3.6 - 6.4)
	5	4	84	916	9.2 (7.5 - 11.2)
	6	12	23	532	4.3 (2.9 -6.4)
	7	12	35	885	4.0 (2.9 - 5.5)
	Mean (SD)				5.42 (SD = 4.89)
	1	2	50	691	7.2 (5.5 - 9.4)
	2	2	103	1000	10.3 (8.6 - 12.3)
HB3	3	9	57	942	6.1 (4.7 -7.8)
	4	9	56	944	5.9 (4.6 -7.6)
	Mean (SD)				6.87 (SD = 2.32)
	1	2	225	1022	22.0 (19.6 - 24.7)
	2	2	182	1013	18.0 (15.7 - 20.5)
	3	3	73	926	7.9 (6.3 - 9.8)
NF54	4	3	54	925	5.8 (4.5 - 7.5)
	5	13	41	619	6.6 (4.9 - 8.9)
	6	13	11	975	1.1 (0.6 - 2.0)
	7	13	27	975	2.8 (1.9 - 4.0)
	Mean (SD)				9.18 (SD = 7.83)

Line	Replicate	RBC Batch	Pfs16 +ve	DAPI +ve	GCR (95% CI)
RO33	1	5	153	861	17.8 (15.4 - 20.5)
	2	5	68	943	7.2 (5.7 - 9.0)
	3	7	70	932	7.5 (6.0 - 9.4)
	4	7	92	910	10.1 (8.3 - 12.2)
	5	10	72	921	7.8 (6.3 - 9.7)
	6	10	124	875	14.2 (12.0 - 16.6)
	Mean (SD)				10.77 (SD = 4.31)
FCC2	1	5	211	785	26.9 (23.9 - 30.1)
	4	5	147	835	17.6 (15.2 - 20.3)
	5	5	16	484	3.3 (2.0 - 5.3)
	6	5	23	503	4.6 (3.1 - 6.8)
	7	10	69	935	7.4 (5.9 - 9.2)
	8	10	47	952	4.9 (3.7 - 6.5)
	Mean (SD)				10.78 (SD = 9.44)
7G8	1	6	159	856	18.6 (16.1 - 21.3)
	2	6	199	803	24.8 (21.9 - 27.9)
	3	10	135	452	29.9 (25.8 - 34.2)
	4	10	24	339	7.1 (4.8 - 10.3)
	5	10	9	456	2.0 (1.0 - 3.7)
	6	11	46	953	4.8 (3.6 - 6.4)
	Mean (SD)				14.52 (SD = 11.52)

Exact counts and 95% confidence intervals of the GCR estimates were performed using
Method 1 (with Pfs16 staining) as plotted and analysed in Figure 4. Erythrocytes from fifteen
different erythrocyte donors used at various times throughout the series of experiments
shown here and in Supplementary Table S3 are numbered (RBC Batch), and inspection of the
data shows that these did not determine the variation between assay replicates.

Supplementary Table S6. Pairwise tests for significant differences in gametocyte conversion rates between different *P. falciparum* laboratory
lines. P values are from Mann-Whitney comparisons of all of the individual biological replicate measures for each of the 13 lines (as shown in
Supplementary Table S5). Statistically significant comparisons are highlighted in green.

Lab Isolates	F12	D10	T9/96	Palo Alto	3D7	D6	GB4	Dd2	HB3	NF54	RO33	7G8	FCC2
F12													
D10	0.0047**												
T9/96	0.0022**	0.1696											
Palo Alto	0.0022**	0.2459	0.6082										
3D7	0.0012**	0.0017**	0.0052**	0.1521									
D6	0.0012**	0.0012**	0.0023**	0.0216*	0.0781								
GB4	0.0012**	0.0006***	0.0012**	0.0047**	0.0117*	0.6457							
Dd2	0.0012**	0.0012**	0.0047**	0.0338*	0.1282	>0.9999	>0.9999						
HB3	0.0048**	0.0030**	0.0095**	0.0061**	0.0012**	0.2303	0.4121	0.3124					
NF54	0.0012**	0.0006***	0.0012**	0.0082**	0.0175*	0.3176	0.535	0.3176	0.9273				
RO33	0.0022**	0.0012**	0.0022**	0.0022**	0.0012**	0.035*	0.035*	0.0734	0.026*	0.4452			
7G8	0.0022**	0.0012**	0.0022**	0.0043**	0.0082**	0.1807	0.1807	0.1101	0.3939	0.4452	>0.9999		
FCC2	0.0022**	0.0012**	0.0022**	0.0022**	0.0012**	0.336	0.2949	0.2343	0.9143	0.9452	0.3939	0.6991	

Supplementary Table S7. Numerical counts and calculated values of assays measuring the effect of choline-free medium on gametocyte conversion rates (GCR) of five different *P. falciparum* lines. The effect of each biological replicate assay is shown as a Rate Ratio with 95% confidence intervals, based on the GCR with and without choline. Mantel-Haenszel adjusted mean Rate Ratios show the overall effects across all replicates with 95% confidence intervals.

Biological Replicate Assays	Condition	Pfs16 +ve	Pfs16 -ve	Rate Ratio (± 95%CI)	GCR (± 95%CI)	Adjusted mean Rate Ratio (± 95%CI)
P. falciparum line HB3						
R1	-choline	16	967	2.03 (0.93 - 5.46)	1.6 (1.0 - 2.6)	
	+ choline	7	864		0.8 (0.4 - 1.6)	
R2	-choline	29	943	3.69 (1.69 - 8.04)	3.0 (2.1 - 4.3)	
	+ choline	8	983		0.8 (0.4 - 1.6)	
R3	-choline	34	902	12.05 (3.71 - 39.13)	3.6 (2.6 -5.0)	
	+ choline	3	993		0.3 (0.1 - 0.9)	
R4	-choline	36	929	2.01 (1.15 - 3.51)	3.7 (2.7 - 5.1)	
	+ choline	18	950		1.9 (1.2 - 2.9)	1.98 (1.55 - 2.52)
R5	-choline	24	954	1.26 (0.70 - 2.30)	2.5 (1.7 - 3.6)	
	+ choline	19	961		1.9 (1.2 - 3.0)	
R6	-choline	14	970	1.00 (0.48 - 2.09)	1.4 (0.8 - 2.4)	
	+ choline	14	971		1.4 (0.8 - 2.4)	
R7	-choline	20	959	0.95 (0.52 - 1.74)	2.0 (1.3 - 3.1)	
	+ choline	21	957		2.1 (1.4 - 3.3)	
R8	-choline	19	962	2.40 1.06 - 5.45)	1.9 (1.2-3.0)	
	+ choline	8	983		0.8 (0.4 - 1.6)	
P. falciparum line Dd2						
R1	-choline	194	619	5.58 (4.03 - 7.71)	23.9 (21.1 - 26.9)	
	+ choline	41	917		4.3 (3.2 - 5.8)	
R2	-choline	42	1867	0.95 (0.61 - 1.47)	2.2 (1.6 - 3.0)	
	+ choline	37	1562		2.31 (1.7 - 3.2)	
R3	-choline	51	912	1.21 (0.81 - 1.80)	5.3 (4.1 - 6.9)	
	+ choline	42	915		4.4 (3.3- 5.9)	3.51 (3.00 - 4.11)
R4	-choline	105	799	16.56 (7.75 - 35.40)	11.6 (9.7 - 13.9)	
	+ choline	7	991		0.7 (0.3 - 1.4)	
R5	-choline	141	847	12.82 (6.98 - 23.53)	14.3 (12.2 - 16.6)	
	+ choline	11	977		1.1 (0.6 - 2.0)	
R6	-choline	93	824	2.07 (1.47 - 2.90)	10.1 (8.4 - 12.3)	
	+ choline	47	910		4.9 (3.7 - 6.5)	

Biological Replicate Assays	Condition	Pfs16 +ve	Pfs16 -ve	Rate Ratio (± 95%CI)	GCR (± 95%CI)	Adjusted mean Rate Ratio (± 95%CI)
			P. falciparum line 3D7			
R1	-choline	17	965	1.22 (0.60 - 2.45)	1.7(1.1 - 2.8)	
	+ choline	14	969		1.4 (0.9 - 2.4)	
R2	-choline	1	997	0.20 (0.02- 1.70)	0.1 (0.0 - 0.6)	
	+ choline	5	989		0.5 (0.2 - 1.2)	
R3	-choline	6	987	3.01 (0.61 - 14.88)	0.6 (0.3 - 1.3)	
	+ choline	2	995		0.2 (0.0 - 0.7)	
R4	-choline	34	931	3.45 (1.72 - 6.95)	3.5 (2.5 - 4.9)	
	+ choline	10	970		1.0 (0.6 - 1.9)	1.50 (1.16 - 1.94)
R5	-choline	4	940	0.82 (0.22 - 3.04)	0.4 (0.2 - 1.1)	
	+ choline	5	961		0.5 (0.2 - 1.2)	
R6	-choline	6	987	0.65 (0.23 - 1.82)	0.6 (0.3 - 1.3)	
	+ choline	9	961		0.9 (0.5 - 1.8)	
R7	-choline	35	931	1.86 (1.08 - 3.24)	3.6 (2.6 - 5.0)	
	+ choline	19	961		1.9 (1.2 - 3.0)	
R8	-choline	43	913	1.28 (0.83 - 1.99)	4.5 (3.4 - 6.0)	
	+ choline	34	938		3.5 (2.5 - 4.8)	
			P. falciparum line T9/96			
R1	-choline	7	985	1.00 (0.35 - 2.84)	0.7 (0.3 - 1.4)	
	+ choline	7	985		0.7 (0.3 - 1.4)	
R2	-choline	2	995	0.57 (0.10 - 3.10)	0.2 (0.0 - 0.7)	
	+ choline	4	1131		0.4 (0.1 - 0.9)	
R3	-choline	2	995	no estimate	0.2 (0.0 - 0.7)	
	+ choline	0	999		0 (0)	
R4	-choline	2	934	0.26 (0.06 - 1.24)	0.2 (0.0 - 0.8)	0.8 (0.51 - 1.19)
	+ choline	8	983		0.8 (0.4 - 1.6)	
R5	-choline	5	643	0.83 (0.23 - 3.09)	0.8 (0.3 - 1.8)	
	+ choline	4	429		0.9 (0.4 - 2.4)	
R6	-choline	11	978	0.84 (0.38 - 1.88)	1.1 (0.6 - 2.0)	
	+ choline	13	973		1.3 (0.8 - 2.2)	
R7	-choline	9	739	0.79 (0.34 - 1.87)	1.2 (0.6 - 2.3)	
	+ choline	12	780		1.5 (0.9 -2.6)	

Biological Replicate Assays	Condition	Pfs16 +ve	Pfs16 -ve	Rate Ratio (± 95%CI)	GCR (± 95%CI)	Adjusted Risk Ratio (± 95%CI)
P.falciparum line NF54						
R1	-choline	27	945	2.50	2.8 (1.9 - 4.0)	
	+ choline	11	978	(1.25 - 5.01)	1.1 (0.6 - 2.0)	
R2	-choline	30	943	5.11	3.1 (2.2- 4.4)	
	+ choline	6	990	(2.13 - 12.24)	0.6 (0.3 - 1.3)	
R3	-choline	40	819	3.53	4.7 (3.4- 6.3)	
	+ choline	13	973	(1.90 - 6.56)	1.3 (0.8 - 2.2)	
R4	-choline	43	913	5.57	4.5 (3.4 - 6.0)	**2.41 (1.97 - 2.96)**
	+ choline	8	983	(2.63 - 11.79)	0.8 (0.4 - 1.6)	
R5	-choline	70	861	1.52	7.5 (6.0 - 9.4)	
	+ choline	47	905	(1.06 - 2.18)	4.9 (3.7 - 6.5)	
R6	-choline	60	880	1.86	6.4 (5.0 - 8.1)	
	+ choline	33	933	(1.23 - 2.83)	3.4 (2.4 - 4.8)	
R7	-choline	8	983	1.60	0.8 (0.4 - 1.6)	
	+ choline	5	986	(0.53 - 4.87)	0.5 (0.2 - 1.2)	
R8	-choline	15	970	5.06	1.5 (0.9 - 2.5)	
	+ choline	3	993	(1.47 - 17.41)	0.3 (0.1 - 0.9)	

848 **Supplementary Figure S1.** Scheme outlining the timing of processes undertaken for two different
849 methods to assay parasite gametocyte conversion rates of different *P. falciparum* lines in culture
850 (details given in Materials and Methods section).

851