Molecular dynamics simulations of the Spike trimeric ectodomain of the SARS-CoV-2 Omicron variant: structural relationships with infectivity, evasion to immune system and transmissibility

Vitor Martins de Freitas Amorim[1], Robson Francisco de Souza[1], Anacleto Silva de Souza[1,*], Cristiane Rodrigues Guzzo[1,*]

[1]Department of Microbiology, Institute of Biomedical Sciences, University of São Paulo, São Paulo, Brazil.

* Corresponding author
E-mail: crisguzzo@usp.br and crisguzzo@gmail.com
E-mail: anacletosilvadesouza@usp.br and anacletosilvadesouza@gmail.com

Running title: Structural impacts of the SARS-CoV-2 Omicron variant Spike

To whom correspondence should be addressed:

Cristiane R. Guzzo, Ph.D, Department of Microbiology, Institute of Biomedical Sciences, University of São Paulo, Av. Prof. Lineu Prestes, 1374, Cidade Universitária, 5508-900, São Paulo/SP, Brazil, +55 11 3091-7298; E-mail: crisguzzo@usp.br

Anacleto Silva de Souza, Ph.D, Department of Microbiology, Institute of Biomedical Sciences, University of São Paulo, Av. Prof. Lineu Prestes, 1374, Cidade Universitária, 5508-900, São Paulo/SP, Brazil, +55 11 3091-7298; E-mail: anacletosilvadesouza@usp.br

Keywords: Omicron variant of SARS-CoV-2, full spike protein, principal component analysis, molecular dynamics simulations, infectivity, immune system escape, transmission.

Graphical abstract

Abstract

The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) Omicron variant is replacing Delta, the most prevalent variant worldwide from the beginning of 2021 until early 2022. The Omicron variant is highly transmissible and responsible for a new worldwide COVID-19 wave. Herein, we calculated molecular dynamics simulations of the SARS-CoV-2 trimeric spike protein of Wuhan-Hu-1 strain (wild type, WT) and the Omicron variant of concern. Structural analyses reveal that the Spike[Omicron] presents more conformational flexibility than Spike[WT], mainly in the N-terminal domain (NTD) and receptor-binding domain (RBD). Such flexibility results in a broader spectrum of different conformations for Spike[Omicron], whereby the RBD can more easily visit an up-conformational state. We reported how the mutations in this variant may influence the intra- and inter-protomer contacts caused by conformational flexibility. Based on our analysis, we suggest that the differences in conformational flexibility between Spike[Omicron] and Spike[WT] may explain the observed gains in infectivity, immune system evasion and transmissibility in this novel variant.

Despite efforts in vaccination in several countries, more than 400 million cases and 5.8 million deaths by Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) have been reported since the beginning of the Coronavirus diseases 2019 (COVID-19) pandemic [1]. The viral genome encodes four structural proteins such as Spike protein (or S protein), nucleocapsid (N), membrane (M) and envelope (E). Furthermore, it contains 16 non-structural proteins [5] (**Figure 1a**). The trimeric Spike ectodomain has 1,273 amino acids containing two subunits, S1 and S2 (**Figure 1b-c**). The S1 subunit contains the N-terminal domain (NTD), a receptor-binding domain (RBD), and SD1 and SD2 domains compose the S1 subunit. The S2 subunit is composed by the fusion peptide (FP), the heptapeptide repeat sequences 1 (HR1) and 2 (HR2), a transmembrane domain (TM) and C-terminal domain (CT) [8,9]. The functional role of S1 is related with the NTD and RBD, which the first acts as a "wedge", playing a key role in the structural changes of the trimeric Spike protein [7] while RBD interacts with human angiotensin-converting enzyme 2 (hACE2) (**Figure 1d**). The S2 subunit favors fusion between viral envelope and host cell membrane, which is mediated by domains HR1, HR2, TM and CT.

Since the start of the pandemic, several SARS-CoV-2 variants of concern (VOC) emerged and some important lineages propagated worldwide, such as the Alpha (B.1.1.7), Beta (B.1.351), Gamma (P.1), Delta (B.1.617.2) and Omicron (B.1.1.529), all showing higher transmission rates than the original Wuhan-Hu-1 strain (wild type, WT). In particular, the Omicron variant was found to include a much higher number of mutations than other VOCs, including both insertions (EPE at position 214) and deletions (Δ69-70, Δ142-144, Δ211) [2,3]. The SARS-CoV-2 Omicron variant carries over 50 mutations spread throughout its polyprotein when compared to the WT, 31 of which are located in the Spike protein [4]. These mutations are A67V, Δ69-70, T95I, Δ142-144, Y145D, Δ211, L212I, insertion of EPE at position 214, G339D, S371L, S373P, S375F, K417N, N440K, G446S, S477N, T478K, E484A, Q493R, G496S, Q498R, N501Y, Y505H, T547K, D614G, H655Y, N679K, P681H, N764K, D796Y, N856K, Q954H, N969K and L981F [10]. In SARS-CoV-2 VOCs, the mutations in the Spike protein often cause an increase in the binding affinity to hACE2 [11,12]. However, when these binding affinity values are compared between VOCs, they are within the same order of magnitude [13–16], thus suggesting that transmissibility, viral load and lethality may also be related to other molecular factors,

such as: (1) an increase of RBD up dispositions in Spike [14,15,17]; (2) immune system evasion [18–22]; and (3) level of exposure of the furin cleavage site to proteases [23–25].

The loss of sensibility to neutralizing antibodies against Omicron may be explained due to the Δ144-145 in the NTD [16]. We hypothesize that insertions and deletions observed in this domain could also be associated with further structural changes between the protomers within the trimeric Spike protein. This hypothesis is supported by the observation that structural changes in the adjacent loops 246─260 and nearby loops 144─155 may decrease recognition of neutralizing antibodies [24]. In addition, P681H mutations near the furin cleavage site (residues 682-685) may slightly increase S1/S2 cleavage [25], probably because this substitution enhances exposure of this site to proteases as it confers greater structural flexibility to this region and probably decrease thermostability of proteins. Proline is the most rigid residue and substitution of this residue in the protein may cause a gain in local flexibility [26]. The D614G mutation may also increase the cleavage by furin, by changing the Spike ectodomain structure due to an allosteric effect [27].

In order to investigate how the mutations found in the Omicron variant affect its trimeric structure in relation to the Spike of Wuhan-Hu-1 strain (sequences in **Figures S1** and **S2**), we performed molecular dynamics simulations of the trimeric ectodomains of the Spike^{WT} and Spike^{Omicron} for 100 ns. The root-mean-square deviation (RMSD) of the Spike^{Omicron} and Spike^{WT} protomers are shown in **Figure S3**. In general, Spike^{Omicron} presents more conformational changes) than Spike^{WT} (**Figures S1, S4** and **S5**). These structural changes may be related with high flexibility of residues localized mainly in the S1 subunit (NTD and RBD) (**Figure 2**). We observed a decrease in flexibility in the furin cleavage region in the Omicron variant compared to the WT (**Figure 2**). However, the structural model of the Spike^{WT} for the MD analysis was based on the structure with PDB ID 7CAB that has a "GSAS" instead of "RRAR" at the furin cleavage site [28]. The root-mean-square fluctuation (RMSF) of Spike^{WT} presented high fluctuations in the NTD (residues 69-77, 144-155 and, 246-258), in the RBD (residues 470-490) and furin cleavage region (residues 676-690). The Spike^{Omicron} showed flexibility in the NTD (residues 70-258), in the RBD (residues 466-498), in SD2 (residues 615-638) and, in the furin cleavage region (residues 673-687). In the Spike^{Omicron}, the NTD mutations, deletions and insertions (A67V, Δ69-70, T95I, Δ142-144, Y145D, Δ211, L212I, insertion of EPE at position 214) may favor the up-conformation of the RBD as well as in the

increasement of flexibility for this domain in relation to SARS-CoV-2[WT]. Indeed, the structure of Spike[Omicron] by cryo-EM revealed that the NTD region is more flexible when compared with the remaining protein [16].

The RBD domain of the Spike protein binds human angiotensin-converting enzyme (hACE2) by two regions, E1 and E2 [29] (**Figure 1**). E1 binds hACE2 mainly through hydrophobic interactions. In contrast, the E2 region, made up of residues 444−454 and 493−505, mediates mostly polar interactions with hACE2 (**Figure 1d**). Curiously, we did not observe significant changes in RMSF in the residues affected by subtitutions G339D, S371L, S373P, S375F, K417N, N440K, G446S, N501Y, Y505H and T547K. However, the mutations S477N, T478K, E484A, Q493R, G496S, Q498R, localized in the RBD, correspond to the E1 region, which has intercalated hydrophobic and hydrophilic residues in WT [29]. The E1 region may be the first to be recognized by hACE2 and presents high RMSF values in the residues 470-490 [29].

Previous study has shown that the D614G mutation plays a key role in SARS-CoV-2's evolution because it is involved with the disruption of the interprotomer contact [30], modulates structural conformations that promote membrane fusion [23], increases virion spike density and infectivity [31,32], and increases the cleavage rate by furin after causing conformational changes in the Spike ectodomain [27]. Accordingly, we also verified that the G614 residue influences the conformational space of the Spike[Omicron] protein. Analysis of the trajectories of the Spike[WT] residues shows that stable interactions between the SD2 subdomain of one protomer and the closest S2 subunit of a neighboring protomer are mediated by hydrogen bonding between residues Asp614─Met835 (hydrogen bonding occupancy of 41.8%), Asp614─Tyr837 (hydrogen bonding occupancy of 84.0%), and Asp614─Arg854 (hydrogen bonding occupancy of 99.8%) (**Figure S6**). Such interactions confer some degree of structural rigidity to the Spike[WT] ectodomain. On the other hand, G614 does not interact with any amino acid residues in Spike[Omicron] and the loss of the aforementioned interactions increases mobility of the SD2 domain and of the link between the FP and HR1, which contains the S2' cleavage site, making the Spike[Omicron] ectodomain more flexible. Such greater flexibility probably favors structural rearrangements that contribute to the greater efficiency of infection observed for this variant [33] probably makes its recognition by the immune system difficult, contributing to the evasion of neutralizing antibodies. Although the G614 residue is distant from the RBD, this substitution may also affect the RBD up/down

dispositions [27]. The Spike S1 subunit is involved in recognition of the host cell through receptor binding and the S2 subunit participates in membrane fusion [8,9]. Mutations near the S1/S2 boundary (residues 682-685) may be associated with increased efficiency of cleavage by furin in VOCs. Spikes of D614G, Alpha, Beta, and Delta variants are efficiently cleaved by furin between residues 682-685 [23–25]. In agreement with cryo-EM data for Spike ectodomain from different SARS-CoV-2 VOCs, our molecular dynamics data reveal that this region is highly flexible in Spike^{Omicron} (**Figures 1**).

To gain insights into the differences between the MD trajectories of Spike^{WT} and Spike^{Omicron}, we performed a principal component analysis of the covariance matrix for positional deviations of backbone atoms of the Spike trimer [34]. From the diagonalization of the correlation matrices, we obtained eigenvalues and their corresponding orthogonal eigenvectors (**Table S1**). Projection of the original covariances onto the orthogonal space defined by the eigenvalues allows calculation of the eigenvectors. The highest eigenvectors are called principal components and represent the most important correlations between residues pairs. We evaluate the first and second eigenvectors, which represent the most relevant motions of the Spike protein detected by molecular dynamics trajectories. The PCA score plot reveals that the Spike protein visits two main sets of states throughout the simulation (**Figure 4**). Using the first eigenvector (PC1, first principal component), we observed that the most prominent movement of the Spike^{WT} is the well known repositioning of the RBD from the up- to the down-conformation (**Figure S7a** and **supplementary movies S1** and **S2**). The second eigenvector (PC2, second principal component), however, showed that the second most important movement of Spike^{WT} RBD is a reverse process, from down- to up-conformation state (**Figure S7b** and **supplementary movies S3** and **S4**). Therefore, PCA-projected trajectory suggests that Spike^{WT} RBD tends to visit more down than up conformation, exposing the residues 470-490 (E1 region) for interactions with hACE2.

As expected, the PCA score plot also showed that Spike^{Omicron} RBD populates the up and down conformations but, according to PC1-projected trajectory, it visits preferably from down- to up-conformation state (**Figure S8a** and **supplementary movies S5** and **S6**). The reverse process to PC1-projected trajectory also was observed, with the Spike^{Omicron} RBD changing from the up to down conformation (**Figure S8b** and **supplementary movies S7** and **S8).** Interestingly, the NTD is more

flexible in Spike^{Omicron} than Spike^{WT}, probably because its domain suffered mutations and also insertions and deletions that probably caused significant structural rearrangement allowing it to increase the RBD up dispositions, as observed by previous studies [27]. In agreement with RMSF analysis (**Figure 1**) and hydrogen bonding occupancies, our PCA data suggest that the D614G is fundamental for improving the RBD up dispositions, contributing to increase the viral infection rate of the Omicron variant.

In order to investigate the effect of the mutations in the Spike^{OMICRON} variant , we also calculated the C_α correlation matrix of the trimeric Spike^{WT} and Spike^{Omicron}. **Figure 4** shows the heat map of the correlation matrix, considering internal correlations and also the long-range interactions between each protomer in both proteins. In general, the anti-correlations may be associated with the movements observed in the PC1-projected trajectory, indicating that the RBD up dispositions on Spike^{Omicron} also may be governed by NTD movements. Mutations in NTD seem to play a functional role fundamental in the Spike protein, making it more flexible and efficient to infect the host cell than wild type.

Overall, the SARS-CoV-2 VOCs evade the neutralizing antibodies contained in plasma from convalescent or/and immunized patients [35] and both NTD and RBD carry important epitopes for recognition by neutralizing antibodies, the later one being the major target for these interactions [36]. Deletion of the residues L242, L243 and A244 in Spike^{Beta} causes structural changes in the adjacent loops 246─260 and nearby loops 144─155, decreasing recognition by neutralizing antibodies [24]. Since neutralizing antibodies show more sensitivity to recognize SARS-CoV-2^{WT} than SARS-CoV-2^{Omicron} and that the antibodies mainly recognize the NTD region and the RBD, we suggested that the significant conformational changes of the NTD and RBD of Spike^{Omicron} caused not only by deletions and insertions, but also by the D614G mutation, may be related to the loss of sensitivity of neutralizing antibodies. Mutations alone are sufficient to decrease antibody binding efficiency, together with structural changes and flexibility can synergistically significantly decrease the neutralization efficiency of antibodies acquired by prior infection by a previous variant or by vaccination. In addition, movements in the NTD domain, as observed by the Omicron variant, may be difficult, by steric hindrance, the binding of antibodies to the NTD and also to the RBD. The loss of sensitivity of neutralizing antibodies for SARS-CoV-2 Omicron variant probably contributes to higher virus transmissibility

than WT. Finally, since D614G may cause an effect in the cleavage site of Spike^{Omicron}, the decreasing flexibility in relation to Spike^{WT}, may lead to higher viral infectivity due to better exposition of the furin cleavage site in this variant. As D614G mutation is present in all VOCs, the decreasing flexibility may explain the increased efficiency of Spike cleavage by furin observed for many VOCs [23–25]. In summary, by using molecular dynamics simulations, we have detected various structural changes in the Spike^{Omicron} ectodomain that help explain features such as the increase in viral infectivity of this variant, while revealing aspects that may be relevant for immune system evasion. Further simulations exploring details of the interaction of this variant with hACE2 and both computational prediction and experimental scanning of epitopes will help clarify the impact of such structural changes in the interactions of the virus and its host.

Acknowledgment

The authors acknowledge the National Council for Scientific and Technological Development (CNPq), the Coordination for the Improvement of Higher Education Personnel (CAPES, grant 88887.374931/2019-00, Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Finance Code 01), and the São Paulo Research Foundation (FAPESP, grants 2019/00195-2, 2020/04680-0 and 2016/09047-8), Brazil, for financial support.

Conflicts of Interest

The authors declare no conflict of interest.

Figure 1. representative schematic of the amino acid sequence and the three-dimensional structure of the Spike protein. a) Polyprotein of SARS-CoV-2. **b)** Spike sequence showing the S1 and S2 subunits, NTD, RBD, SD1, SD2, FP, HR1, HR2, TM and CD. c) Tridimensional structure of the trimeric Spike ectodomain (PDB ID 7CAB[28]). d) RBD interacting with hACE2 (PDB ID 6M0J[37]). The painel d) describes two regions, E1 and E2. E1 region is delimited by K417, L455, F456 and residues 470−490[29]. The E2 region corresponds to residues 446−453 and 493−505 residues[29].

Figure 2. Structural aspects obtained by molecular dynamics simulations of Spike^{WT} and Spike^{Omicron}. Backbone root-mean-square fluctuation (RMSF) of **a)** Spike^{WT} and **b)** Spike^{Omicron}. The Spike^{WT} presented high flexibility in NTD (residues 69-77, 144-155, 246-258), in RBD (residues 470-490) and S1/S2 region (residues 676-690). The Spike^{Omicron} has more flexibility in NTD (residues 70-258), in RBD (residues 466-498), in SD2 (residues 615-638) and S1/S2 interface (residues 673-687). Mutations (A67V, Δ69-70, T95I, Δ142-144, Y145D, Δ211, L212I, insertion of EPE at position 214) in the NTD contribute to RBD to adopt an up conformation in Spike^{Omicron}.

Figure 3. Principal component analysis (PCA) of the molecular dynamics trajectories. From the average structure of Spike backbone (WT or Omicron variant), the correlation matrix (fitting non-mass weighted) was calculated. After diagonalizing the correlation matrix, the eigenvalues were determined (**Table S1**) and used to extract the orthogonal eigenvectors (or principal components). The principal components represent the maximum correlation between residue pairs along the MD simulation. MD trajectories were projected in the eigenvector 1 (or PC1, principal component 1) to obtain the main movements of the Spike protein. Similarly, MD trajectories were projected in the eigenvector 2 (or PC2, principal component 2) to obtain the second main movements of the Spike protein. The PC2 movements are less significant than PC1 movements. From projected MD trajectories in the two first principal components, PCA scores were calculated. The PCA score as a function of time (left) and PCA score plot (right) were performed from the backbone of **a)** Spike^{WT} and **b)** Spike^{Omicron}.

Figure 4. Trimeric Spike correlation matrix of the wild type and Omicron variant. The correlations were calculated to find correlated motions between internal residues and interprotomer residues in **a)** WT and **b)** Omicron. The gray circles indicate how the movement of residues influences the mobility of the Spike protein. The red and blue dots show the positively and negatively correlated pairs, respectively. Note that the correlations are more intenses in Spike[Omicron] than Spike[WT]. In Spike[WT], the negative and positive correlations are observed between pairs S2[protomer 1]/NTD[protomer 3] and RBD[protomer 3]/S2[protomer 3], respectively. In Spike[WT], all protomers were positively correlated between their domains (main diagonal matrix) but notably only protomer 3 presented anti-correlation between NTD/S2. In contrast, Spike[Omicron] showed more anti-correlated residue pairs between protomers than Spike[WT]. The internal anti-correlations between their domains are more than WT. In Spike[Omicron], the anti-correlation are observed between pairs NTD[protomer 1]/S2[protomer1], NTD[protomer 1]/NTD[protomer 3], RBD[protomer 1]/RBD[protomer 2], RBD[protomer 1]/RBD[protomer 3], S2[protomer 1]/S2[protomer 2], S2[protomer 1]/RBD[protomer 3], S2[protomer 1]/S2[protomer 3], NTD[protomer 2]/RBD[protomer 3], NTD[protomer 2]/S2[protomer 3] and S2[protomer 2]/NTD[protomer 3]. Conversely, only the pairs S2[protomer 1]/NTD[protomer 3] and S2[protomer 2]/S2[protomer 3] were positively correlated.

References

(1) WHO Coronavirus (COVID-19) Dashboard https://covid19.who.int (accessed 2022 -01 -17).

(2) Classification of Omicron (B.1.1.529): SARS-CoV-2 Variant of Concern https://www.who.int/news/item/26-11-2021-classification-of-omicron-(b.1.1.529)-sars-cov -2-variant-of-concern (accessed 2022 -01 -25).

(3) CDC. Potential Rapid Increase of Omicron Variant Infections in the United States https://www.cdc.gov/coronavirus/2019-ncov/science/forecasting/mathematical-modeling-outbreak.html (accessed 2022 -02 -09).

(4) outbreak.info https://outbreak.info/ (accessed 2022 -01 -25).

(5) Wu, F.; Zhao, S.; Yu, B.; Chen, Y.-M.; Wang, W.; Song, Z.-G.; Hu, Y.; Tao, Z.-W.; Tian, J.-H.; Pei, Y.-Y.; Yuan, M.-L.; Zhang, Y.-L.; Dai, F.-H.; Liu, Y.; Wang, Q.-M.; Zheng, J.-J.; Xu, L.; Holmes, E. C.; Zhang, Y.-Z. A New Coronavirus Associated with Human Respiratory Disease in China. *Nature* **2020**, *579* (7798), 265–269.

(6) Harvey, W. T.; Carabelli, A. M.; Jackson, B.; Gupta, R. K.; Thomson, E. C.; Harrison, E. M.; Ludden, C.; Reeve, R.; Rambaut, A.; COVID-19 Genomics UK (COG-UK) Consortium; Peacock, S. J.; Robertson, D. L. SARS-CoV-2 Variants, Spike Mutations and Immune Escape. *Nat. Rev. Microbiol.* **2021**, *19* (7), 409–424.

(7) Li, Y.; Wang, T.; Zhang, J.; Shao, B.; Gong, H.; Wang, Y.; He, X.; Liu, S.; Liu, T.-Y. Exploring the Regulatory Function of the -Terminal Domain of SARS-CoV-2 Spike Protein through Molecular Dynamics Simulation. *Adv Theory Simul* **2021**, *4* (10), 2100152.

(8) Wrapp, D.; Wang, N.; Corbett, K. S.; Goldsmith, J. A.; Hsieh, C.-L.; Abiona, O.; Graham, B. S.; McLellan, J. S. Cryo-EM Structure of the 2019-nCoV Spike in the Prefusion Conformation. *Science* **2020**, *367* (6483), 1260–1263.

(9) Walls, A. C.; Park, Y.-J.; Tortorici, M. A.; Wall, A.; McGuire, A. T.; Veesler, D. Structure, Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. *Cell* **2020**, *183* (6), 1735.

(10) GISAID - hCov19 Variants https://www.gisaid.org/hcov19-variants/ (accessed 2022 -02 -03).

(11) Barton, M. I.; MacGowan, S.; Kutuzov, M.; Dushek, O.; Barton, G. J.; van der Merwe, P. A. Effects of Common Mutations in the SARS-CoV-2 Spike RBD Domain and Its Ligand the Human ACE2 Receptor on Binding Affinity and Kinetics. https://doi.org/10.1101/2021.05.18.444646.

(12) Spratt, A. N.; Kannan, S. R.; Woods, L. T.; Weisman, G. A.; Quinn, T. P.; Lorson, C. L.; Sönnerborg, A.; Byrareddy, S. N.; Singh, K. Factors Associated with Emerging and Re-Emerging of SARS-CoV-2 Variants. *bioRxiv* **2021**. https://doi.org/10.1101/2021.03.24.436850.

(13) de Souza, A. S.; de Freitas Amorim, V. M.; Guardia, G. D. A.; dos Santos, F. R. C.; dos Santos, F. F.; de Souza, R. F.; de Araujo Juvenal, G.; Huang, Y.; Ge, P.; Jiang, Y.; Paudel, P.; Ulrich, H.; Galante, P. A. F.; Guzzo, C. R. Molecular Dynamics Analysis of Fast-Spreading Severe Acute Respiratory Syndrome Coronavirus 2 Variants and Their Effects in the Interaction with Human Angiotensin-Converting Enzyme 2. *bioRxiv*, 2021, 2021.06.14.448436. https://doi.org/10.1101/2021.06.14.448436.

(14) Zhang, J.; Xiao, T.; Cai, Y.; Lavine, C. L.; Peng, H.; Zhu, H.; Anand, K.; Tong, P.; Gautam, A.; Mayer, M. L.; Walsh, R. M., Jr; Rits-Volloch, S.; Wesemann, D. R.; Yang, W.; Seaman, M. S.; Lu, J.; Chen, B. Membrane Fusion and Immune Evasion by the Spike Protein of SARS-CoV-2 Delta Variant. *Science* **2021**, *374* (6573), 1353–1360.

(15) Wang, Y.; Xu, C.; Wang, Y.; Hong, Q.; Zhang, C.; Li, Z.; Xu, S.; Zuo, Q.; Liu, C.; Huang, Z.; Cong, Y. Conformational Dynamics of the Beta and Kappa SARS-CoV-2 Spike Proteins and Their Complexes with ACE2 Receptor Revealed by Cryo-EM. *Nat. Commun.* **2021**, *12* (1), 7345.

(16) Mannar, D.; Saville, J. W.; Zhu, X.; Srivastava, S. S.; Berezuk, A. M.; Tuttle, K. S.;

Marquez, A. C.; Sekirov, I.; Subramaniam, S. SARS-CoV-2 Omicron Variant: Antibody Evasion and Cryo-EM Structure of Spike Protein-ACE2 Complex. *Science* **2022**, eabn7760.

(17) Xia, S.; Wen, Z.; Wang, L.; Lan, Q.; Jiao, F.; Tai, L.; Wang, Q.; Sun, F.; Jiang, S.; Lu, L.; Zhu, Y. Structure-Based Evidence for the Enhanced Transmissibility of the Dominant SARS-CoV-2 B.1.1.7 Variant (Alpha). *Cell Discov* **2021**, *7* (1), 109.

(18) Li, Q.; Wu, J.; Nie, J.; Zhang, L.; Hao, H.; Liu, S.; Zhao, C.; Zhang, Q.; Liu, H.; Nie, L.; Qin, H.; Wang, M.; Lu, Q.; Li, X.; Sun, Q.; Liu, J.; Zhang, L.; Li, X.; Huang, W.; Wang, Y. The Impact of Mutations in SARS-CoV-2 Spike on Viral Infectivity and Antigenicity. *Cell* **2020**, *182* (5), 1284–1294.e9.

(19) Banoun, H. Evolution of SARS-CoV-2: Review of Mutations, Role of the Host Immune System. *Nephron* **2021**, *145* (4), 392–403.

(20) Chen, J.; Wang, R.; Wang, M.; Wei, G.-W. Mutations Strengthened SARS-CoV-2 Infectivity. *J. Mol. Biol.* **2020**, *432* (19), 5212–5226.

(21) Ku, Z.; Xie, X.; Davidson, E.; Ye, X.; Su, H.; Menachery, V. D.; Li, Y.; Yuan, Z.; Zhang, X.; Muruato, A. E.; I Escuer, A. G.; Tyrell, B.; Doolan, K.; Doranz, B. J.; Wrapp, D.; Bates, P. F.; McLellan, J. S.; Weiss, S. R.; Zhang, N.; Shi, P.-Y.; An, Z. Author Correction: Molecular Determinants and Mechanism for Antibody Cocktail Preventing SARS-CoV-2 Escape. *Nat. Commun.* **2021**, *12* (1), 4177.

(22) Xie, X.; Liu, Y.; Liu, J.; Zhang, X.; Zou, J.; Fontes-Garfias, C. R.; Xia, H.; Swanson, K. A.; Cutler, M.; Cooper, D.; Menachery, V. D.; Weaver, S. C.; Dormitzer, P. R.; Shi, P.-Y. Neutralization of SARS-CoV-2 Spike 69/70 Deletion, E484K and N501Y Variants by BNT162b2 Vaccine-Elicited Sera. *Nat. Med.* **2021**, *27* (4), 620–621.

(23) Zhang, J.; Cai, Y.; Xiao, T.; Lu, J.; Peng, H.; Sterling, S. M.; Walsh, R. M., Jr; Rits-Volloch, S.; Zhu, H.; Woosley, A. N.; Yang, W.; Sliz, P.; Chen, B. Structural Impact on SARS-CoV-2 Spike Protein by D614G Substitution. *Science* **2021**, *372* (6541), 525–530.

(24) Cai, Y.; Zhang, J.; Xiao, T.; Lavine, C. L.; Rawson, S.; Peng, H.; Zhu, H.; Anand, K.; Tong, P.; Gautam, A.; Lu, S.; Sterling, S. M.; Walsh, R. M., Jr; Rits-Volloch, S.; Lu, J.; Wesemann, D. R.; Yang, W.; Seaman, M. S.; Chen, B. Structural Basis for Enhanced Infectivity and Immune Evasion of SARS-CoV-2 Variants. *Science* **2021**. https://doi.org/10.1126/science.abi9745.

(25) Lubinski, B.; Fernandes, M. H. V.; Frazier, L.; Tang, T.; Daniel, S.; Diel, D. G.; Jaimes, J. A.; Whittaker, G. R. Functional Evaluation of the P681H Mutation on the Proteolytic Activation of the SARS-CoV-2 Variant B.1.1.7 (Alpha) Spike. *iScience* **2022**, *25* (1), 103589.

(26) Yu, H.; Zhao, Y.; Guo, C.; Gan, Y.; Huang, H. The Role of Proline Substitutions within Flexible Regions on Thermostability of Luciferase. *Biochim. Biophys. Acta* **2015**, *1854* (1), 65–72.

(27) Gobeil, S. M.-C.; Janowska, K.; McDowell, S.; Mansouri, K.; Parks, R.; Manne, K.; Stalls, V.; Kopp, M. F.; Henderson, R.; Edwards, R. J.; Haynes, B. F.; Acharya, P. D614G Mutation Alters SARS-CoV-2 Spike Conformation and Enhances Protease Cleavage at the S1/S2 Junction. *Cell Rep.* **2021**, *34* (2), 108630.

(28) Lv, Z.; Deng, Y.-Q.; Ye, Q.; Cao, L.; Sun, C.-Y.; Fan, C.; Huang, W.; Sun, S.; Sun, Y.; Zhu, L.; Chen, Q.; Wang, N.; Nie, J.; Cui, Z.; Zhu, D.; Shaw, N.; Li, X.-F.; Li, Q.; Xie, L.; Wang, Y.; Rao, Z.; Qin, C.-F.; Wang, X. Structural Basis for Neutralization of SARS-CoV-2 and SARS-CoV by a Potent Therapeutic Antibody. *Science* **2020**, *369* (6510), 1505–1509.

(29) Silva de Souza, A.; Rivera, J. D.; Almeida, V. M.; Ge, P.; de Souza, R. F.; Farah, C. S.; Ulrich, H.; Marana, S. R.; Salinas, R. K.; Guzzo, C. R. Molecular Dynamics Reveals Complex Compensatory Effects of Ionic Strength on the Severe Acute Respiratory Syndrome Coronavirus 2 Spike/Human Angiotensin-Converting Enzyme 2 Interaction. *J. Phys. Chem. Lett.* **2020**, *11* (24), 10446–10453.

(30) Mansbach, R. A.; Chakraborty, S.; Nguyen, K.; Montefiori, D. C.; Korber, B.; Gnanakaran, S. The SARS-CoV-2 Spike Variant D614G Favors an Open

Conformational State. *Sci Adv* **2021**, *7* (16). https://doi.org/10.1126/sciadv.abf3671.

(31)   Yurkovetskiy, L.; Wang, X.; Pascal, K. E.; Tomkins-Tinch, C.; Nyalile, T. P.; Wang, Y.; Baum, A.; Diehl, W. E.; Dauphin, A.; Carbone, C.; Veinotte, K.; Egri, S. B.; Schaffner, S. F.; Lemieux, J. E.; Munro, J. B.; Rafique, A.; Barve, A.; Sabeti, P. C.; Kyratsous, C. A.; Dudkina, N. V.; Shen, K.; Luban, J. Structural and Functional Analysis of the D614G SARS-CoV-2 Spike Protein Variant. *Cell* **2020**, *183* (3), 739–751.e8.

(32)   Zhang, L.; Jackson, C. B.; Mou, H.; Ojha, A.; Peng, H.; Quinlan, B. D.; Rangarajan, E. S.; Pan, A.; Vanderheiden, A.; Suthar, M. S.; Li, W.; Izard, T.; Rader, C.; Farzan, M.; Choe, H. SARS-CoV-2 Spike-Protein D614G Mutation Increases Virion Spike Density and Infectivity. *Nat. Commun.* **2020**, *11* (1), 6013.

(33)   Zhang, X.; Wu, S.; Wu, B.; Yang, Q.; Chen, A.; Li, Y.; Zhang, Y.; Pan, T.; Zhang, H.; He, X. SARS-CoV-2 Omicron Strain Exhibits Potent Capabilities for Immune Evasion and Viral Entrance. *Signal Transduct Target Ther* **2021**, *6* (1), 430.

(34)   Amadei, A.; Linssen, A. B. M.; Berendsen, H. J. C. Essential Dynamics of Proteins. *Proteins: Structure, Function, and Genetics*. 1993, pp 412–425. https://doi.org/10.1002/prot.340170408.

(35)   Casadevall, A.; Henderson, J. P.; Joyner, M. J.; Pirofski, L.-A. SARS-CoV-2 Variants and Convalescent Plasma: Reality, Fallacies, and Opportunities. *J. Clin. Invest.* **2021**, *131* (7). https://doi.org/10.1172/JCI148832.

(36)   Yang, Y.; Du, L. SARS-CoV-2 Spike Protein: A Key Target for Eliciting Persistent Neutralizing Antibodies. *Signal Transduct Target Ther* **2021**, *6* (1), 95.

(37)   Lan, J.; Ge, J.; Yu, J.; Shan, S.; Zhou, H.; Fan, S.; Zhang, Q.; Shi, X.; Wang, Q.; Zhang, L.; Wang, X. Structure of the SARS-CoV-2 Spike Receptor-Binding Domain Bound to the ACE2 Receptor. *Nature* **2020**, *581* (7807), 215–220.