Constitutively active STAT5b feminizes mouse liver gene expression

Dana Lau-Corona[1], Hong Ma[1], Cameron Vergato[1], Andre Sarmento-Cabral[2],
Mercedes del Rio-Moreno[2], Rhonda D Kineman[2], David J Waxman[1]

[1] Department of Biology and Bioinformatics Program
Boston University, Boston, MA

[2] Department of Medicine, Section of Endocrinology, Diabetes, and Metabolism, University of
Illinois at Chicago and Research and Development Division, Jesse Brown VA Medical Center,
Chicago, IL

Key words: Liver sex differences, STAT5 knockout, pituitary GH secretion profiles, JAK-STAT

Running title: Feminization of liver by STAT5b$_{CA}$

Supported in part by NIH grant DK121998 (to DJW) and grants NIH DK114326 and VA Merit
BX005382 (to RK)

Address correspondence to:

Dr. David J. Waxman
Dept. of Biology
Boston University
5 Cummington Mall
Boston, MA 02115 USA
Email: djw@bu.edu

Abstract

STAT5 is an essential transcriptional regulator of the sex-biased actions of growth hormone
(GH) in the liver. Delivery of constitutively active STAT5 (STAT5$_{CA}$) to male mouse liver using an
engineered adeno-associated virus with high tropism for the liver is shown to induce widespread
feminization of the liver, with extensive induction of female-biased genes and repression of
male-biased genes, largely mimicking results obtained when male mice are given GH as a
continuous infusion. Many of the gene expression changes observed were associated with
STAT5 binding to liver chromatin, supporting the proposed direct role of persistently active
STAT5 in continuous GH-induced liver feminization. The feminizing effects of STAT5$_{CA}$ were
dose-dependent; moreover, at higher levels, overexpression of STAT5$_{CA}$ resulted in some
histopathology not seen in continuous GH-infused male liver, including hepatocyte hyperplasia
and distorted liver architecture. These findings establish that the persistent activation of STAT5
by GH that characterizes female liver is by itself sufficient to account for the female-biased
expression of a majority of female-biased genes. Moreover, histological changes seen when
STAT5$_{CA}$ is overexpressed highlight the importance of carefully evaluating such effects before
considering such STAT5 derivatives for therapeutic use in treating liver disease.

Introduction

Sex-differences in pituitary growth hormone (GH) secretory patterns control the sex-biased expression of hundreds of genes in mouse liver (1) and contribute to the sexual dimorphism of liver physiology and pathology (2,3). In males, GH is secreted as a series of intermittent pulses followed by extended periods when circulating GH is undetectable, whereas in females, GH secretion is near continuous (1,4). Sex differences in liver gene expression are largely abolished when pituitary GH secretion is ablated by hypophysectomy, and can be substantially restored when exogenous GH is administered as a series of pulses (male circulating GH pattern), which confers a male pattern of expression, or as a continuous infusion (female GH pattern), which results in a female pattern of expression (5,6).

GH binding to its cell surface receptor activates the receptor-bound tyrosine kinase JAK2, which in turn phosphorylates and thereby activates the latent cytoplasmic transcription factor STAT5 (7,8). GH receptor/JAK2-activated STAT5 dimerizes and translocates to the nucleus, where it binds to thousands of sites in liver chromatin and induces transcription of many target genes (9,10). STAT5b, the major liver STAT5 form, plays a crucial role in liver metabolism and is an essential transcriptional regulator of the sex-dependent actions of GH in the liver (11,12). In male liver, repeated pulses of hepatic STAT5 activity, nuclear localization and transcriptional activity are observed and closely track the successive pulses of male pituitary gland GH release (13-15), whereas in female liver, STAT5 is persistently activated by the near-continuous female plasma GH profile (9,16). In whole body STAT5b knockout male mice, 90% of hepatic male-biased genes are repressed and 60% of hepatic female-biased genes are de-repressed (i.e., induced) (11). Further, STAT5 binds in a sex-biased manner to several thousand sites in mouse liver chromatin, with the set of male-biased STAT5 binding sites enriched nearby genes showing male-biased expression and the set of female-biased binding sites enriched nearby female-biased genes (9,17). Recent studies in a hepatocyte-specific STAT5a/STAT5b-KO mouse model (18) confirm these findings (19), supporting the proposal that the loss of hepatic STAT5, rather than an indirect feedback response to the loss of STAT5 in the hypothalamus or the pituitary gland, is responsible for the loss of liver sexual dimorphism seen in global STAT5b knockout mice.

Continuous infusion of GH in male mice (cGH treatment) substantially feminizes liver gene expression (20,21), most likely by a combination of two mechanisms. First, cGH overrides the endogenous male pulsatile plasma GH profile. This leads to widespread repression of male-biased genes that are positively regulated by male plasma GH pulses (class I male-biased genes) and to de-repression of female-biased genes that are negatively regulated by male GH pulses (class II female-biased genes) (22). Second, cGH treatment imposes a persistent, female-like pattern of plasma GH stimulation on male liver, which leads to the induction of female-biased genes that are positively regulated by the endogenous female GH pattern (class I female-biased genes) and represses male-biased genes that are negatively regulated by the female GH pattern (class II male-biased genes) (15,22). GH-activated STAT5 is a strong transcriptional activator of its well established direct target genes, including *Igf1* and *Socs2* (10,23), and similarly, can directly activate at least some class I male-biased genes when it is activated by a plasma GH pulse in male mouse liver (15). However, it is uncertain whether STAT5 signaling directly regulates other classes of sex-biased genes, most notably class I female-biased genes, which require a female plasma GH pattern for expression, or alternatively, whether other GH-stimulated but STAT5-independent signaling pathways downstream of GH receptor and JAK2 (24-26) mediate their sex-dependent expression. The requirement of STAT5 for sex-biased hepatic expression of both class I and class II sex-biased genes (21) does not entirely resolve this question, given the apparent perturbation of feedback mechanisms leading

102 to changes in circulating GH patterns seen in both STAT5 knockout and GH receptor-deficient
103 mouse models (18,27).
104
Here, we address these questions using AAV8-STAT5$_{CA}$, a novel adeno-associated virus (AAV)
vector that we engineered to deliver STAT5b with a point mutation in its SH2 domain (STAT5b
Asn-642 to His) (28). This mutation renders the STAT5 protein constitutively active ('CA') and
thus able to support robust expression of hepatic *Igf1* in the absence of GH (29). AAV8-
STAT5$_{CA}$ is based on AAV serotype 8, which has high specificity for infection of hepatocytes
(30,31). Moreover, the STAT5b mutant is expressed from the hepatocyte-specific thyroxin-
binding globulin promoter, which confers additional specificity for persistent expression of the
constitutively active STAT5 in hepatocytes. We hypothesized that the persistent expression of
active STAT5 protein in male mouse liver mimics the persistent activation of liver STAT5 seen in
female liver and will feminize gene expression. Our findings reveal that the delivery of STAT5$_{CA}$
to male mouse liver leads to widespread induction of female-biased genes and repression of
male-biased genes, and thus feminization of the liver by persistent plasma GH stimulation can
largely be attributed to the persistent activation of liver STAT5b.
118
119
120 **Materials and Methods**
121
Preparation of AAV8-STAT5$_{CA}$ - Wild-type mouse STAT5b expression plasmid with an N-
terminal Flag tag was obtained from SinoBiological, Wayne PA (cat. # MG51116-NF). A point
mutation at base 1963 (A to C), which changes Asn642 to His, was introduced into the STAT5b
coding sequence by Genscript (Piscataway, NJ). This mutation confers constitutive activity (CA)
to STAT5b, including the ability to induce *Igf1* gene expression in livers of hypophysectomized
(GH-deficient) rats (28,29). The mutated plasmid was sent to Penn Vector Core (University of
Pennsylvania, Philadelphia, PA), where the N-Flag-mSTAT5b$_{CA}$ cDNA was cloned into the Penn
Vector Core plasmid pENN.AAV.TBG.PI.eGFP.WPRE.bGH, replacing eGFP by the mutated
and tagged STAT5b$_{CA}$. The resultant plasmid, pENN.AAV.TBG.PI.N-FLAG-
mSTAT5bCA.WPRE.bGH, was sequence verified then packaged in AAV8 virus, which was
purified, quantified with respect to genome copy (GC) number, and supplied by Penn Vector as
a ready to use vector, AAV8-TBGp-mStat5b$_{CA}$, referred to here as AAV8-STAT5$_{CA}$. The plasmid
pENN.AAV.TBG.PI.N-FLAG-mSTAT5bCA.WPRE.bGH, required to prepare AAV8-STAT5$_{CA}$
(see lab methods, below), is being deposited and will be made available through Addgene
(https://www.addgene.org/).
137
AAV8-STAT5$_{CA}$ expression in mouse liver. Initial studies to validate AAV8-STAT5$_{CA}$ were
performed in male C57Bl/6 mice on a GHR[fl/fl] background at Jesse Brown VA Medical Center
(Chicago, IL) with Institutional Animal Care and Use Committee approval (protocol number 19-
05). AAV8-TBGp-Null vector (Penn Vector Core), referred to here as AAV8-Null, was used as a
control. Male GHR[fl/fl] mice (10-12 wk, and maintained as an in-house breeding colony, originally
provided by Dr. John Kopchick, Ohio University (32)) were injected via the lateral tail vein with
AAV8-Null or AAV8-STAT5$_{CA}$ (1.5 x 10^{11} GC per mouse, diluted in sterile PBS). Seven days
later the mice were decapitated, and livers were collected. Nuclear protein was extracted using
NE-PER Nuclear Cytoplasmic Extraction Reagent (cat. # 78833, Pierce, Rockford, IL). Western
blots were performed to detect the Flag-tag sequence using rabbit monoclonal anti-DYKDDDDK
Tag antibody (Cell Signaling #14793). To test the efficacy of AAV8-STAT5$_{CA}$ to increase hepatic
expression and restore circulating levels of IGF1 to normal levels, we used a mouse model with
adult-onset hepatocyte-specific GH receptor knockdown (aHepGHRkd) (33), prepared as
follows. Male GHR[fl/fl] mice (10-12 wk old) were injected via lateral tail vein with 1.5 x 10^{11}
GC/mouse of either AAV8-Null, to obtain GH receptor-intact control mice, or AAV8 expressing

Cre recombinase (AAV8-TBGp-Cre, Penn Vector Core), to generate aHepGHRkd mice. The
above AAV8 injections done either alone or in combination with AAV8-STAT5$_{CA}$ (at 0.36 or 0.75
x 10^{11} GC), with the dose of AAV8-Null adjusted to equalize the total GC of AAV8 per mouse
across groups. Fourteen days later, blood and liver were collected to assay circulating IGF1
levels (Mouse/Rat IGF-1 ELISA, 22-IG1MS-E01, ALPCO) and liver IGF1 mRNA levels by qPCR
(33). In other studies, GH receptor-intact male mice (C57Bl/6 strain from Jackson Labs, 10 wk
old) were treated with AAV8-Null or AAV8-STAT5$_{CA}$ (0.75 x 10^{11} GC per mouse, iv) and 14 d
later livers were collected and used for RNA-seq analysis (see below).
161
AAV8-STAT5$_{CA}$ dose response studies. Studies to test the ability of AAV-STAT5$_{CA}$ to feminize
hepatic gene expression were performed, in compliance with procedures approved by the
Boston University Institutional Animal Care and Use Committee (protocol # PROTO201800698),
and in compliance with ARRIVE 2.0 Essential 10 guidelines (34), including study design, sample
size, randomization, experimental animals and procedures, and statistical methods. Male and
female CD1 mice (Crl:CD(ICR), strain code 022), 7-8 wk of age, were purchased from Charles
River Laboratories. Mice were housed in a temperature and humidity-controlled environment
with a 12-h light, 12-h dark cycle, fed standard rodent chow, and supplied with tap water. AAV8-
STAT5$_{CA}$ or AAV8-Luciferase (control) (both produced in-house using methods described
below) were administered to male mice by tail vein injection at doses ranging from 0.125 x 10^{11}
to 2 x 10^{11} GC per mouse. Virus was diluted into PBS + 35 mM NaCl + 5% glycerol (stabilizer)
to give a total injection volume of 150 uL. Mice were euthanized by cervical dislocation at 11 AM
(lights on 7:30 AM- 7:30 PM) to minimize the impact of circadian variations of gene expression
(35,36).
176
AAV8 production. AAV coding STAT5b$_{CA}$ and Luciferase (control) was prepared and quantified
by triple transfection of HEK293FT cells using methods adapted from protocols provided by
Addgene (www.addgene.org). HEK293FT is a fast growing and highly transfectable derivative
(cat. #R70007, ThermoFisher Scientific) of the SV40 large T-antigen-expressing HEK293T cell
line commonly used for AAV production. On Day 0, 80% confluent flasks of HEK293FT cells
were split 1:4 into DMEM culture medium [10% Fetal Bovine Serum (cat. #10437-028, Gibco),
1% PenStrep (cat. #15140-122, Gibco)] and then grown overnight to achieve 50-60%
confluence at the time of transfection on Day 1. T182.5 flasks were each transfected with a total
of 80 µg of plasmid DNA in a 1:1:1 (molar ratio) of three plasmids: pAAV-Helper helper plasmid,
which encodes adenovirus genes E2A, E4 and (Cell Biolabs, VA); pAAV2/8 Rep/Cap plasmid
(Addgene, plasmid #112864; replication protein derived from AAV-2 serotype, and Capsid
protein from AAV-8); and either pENN.AAV.TBG.PI.N-FLAG-tag mSTAT5bCA.WPRE.bGH (see
above) or pENN.AAV.TBG.PI.ffLuciferase.RBG (firefly luciferase expressed from TBG promoter;
Addgene # 105538). Plasmids were mixed in a 3:1 weight ratio of total plasmid DNA:PEI in 4
mL of either OptiMeM™ (Gibco cat. # 31985-062) or serum-free DMEM per flask and then
shaken vigorously for 30 sec. The mixture was incubated for 15 min at room temperature in a
tissue culture hood, after which 4 ml of the mixture was added to each flask of ~85% confluent
HEK293FT cells. After 24 h, the culture medium was replaced by DMEM containing 2% FBS
and 1% PenStrep, and the cells were returned to the CO_2 incubator for 5 days.
196
Virus was harvested on day 6 after transfection. A cell scraper was used to detach the cells
from the flask, followed by collection of the media with cells in sterile 50 mL conical tubes. The
tubes were centrifuged at 1000 x g for 10 min at 4°C, and the supernatant was decanted into a
sterile bottle. Pellets were pooled and transferred to smaller tubes with 1x PBS + 200 mM NaCl
+ 0.001% Pluronic F68 (Poloxamer 188 Solution, cat. #P5556-100ML, Sigma) and pelleted
again. Cell pellets were frozen at -80 °C for future use in purifying virus. The supernatant (which
contains the majority of virus) was filtered through a 0.2 µM PES filter (VWR, cat. #97066-214).

25 mL of 40% PEG8000 (JT Baker, cat. #JTU222-09) was then added for every 100 mL of
media supernatant, followed by overnight stirring at 4°C. The next day, the supernatant was
incubated for 1 h at 4°C without stirring, to allow for full precipitation of virus before spinning at
2,818 x g for 15 min at 4°C. The supernatant was discarded and the PEG pellet, containing
concentrated virus, was resuspended in 10 mL of 1x PBS + 0.001% Pluronic F68 + 200 mM
NaCl by pipetting back and forth until the pellet was completely in solution.
210
The following buffers were prepared for iodixanol purification of the PEG-pelleted virus: (1) 1 M
NaCl/PBS-MK Buffer (5.84 g NaCl, 26.3 mg $MgCl_2$ and 14.91 mg KCl dissolved in 1x PBS to
give a final volume of 100 mL), sterile filtered through a 0.2 μm PES filter; and (2) 1x PBS-MK
Buffer (26.3 mg $MgCl_2$ and 14.91 mg KCl dissolved in 1x PBS to a final volume of 100 mL and
then filter-sterilized by passage through a 0.2 μm filter. Iodixanol gradients were prepared as
follows: 15% Layer (4.5 mL of 60% Iodixanol (OptiPrep Density Gradient Medium, cat. # D1556-
250ML, Sigma) + 13.5 mL of 1M NaCl/PBS-MK Buffer); 25% Layer (5 mL of 60% Iodixanol, 7
mL of 1x PBS-MK Buffer + 30 µL of Phenol Red (cat. #P0290-100ML, Sigma); 40% Layer (6.7
mL of Iodixanol + 3.3 mL of 1x PBS-MK Buffer; 60% layer (10 mL of 60% Iodixanol + 45 µL
phenol red). A 10 mL syringe with an 18 g needle and a glass Pasteur pipet were used to layer
the Iodixanol gradient in QuickSeal™ tubes (cat. # 344326, Beckman Coulter), from the bottom
to the top, starting with 8 mL of the 15% layer, then 6 mL of the 25% layer, 5 mL of the 40%
layer and 5 mL of the 60% layer (two tubes per AAV preparation from 10-12 T182.5 flasks).
Resuspended virus-PEG pellet solution (5 mL) was added to the top of each gradient and 1x
PBS was used to top off each tube before heat sealing. The tubes were spun for 90 min at
350,000 x g in a T70i Rotor at 10°C. The tubes were removed carefully (without disturbing the
gradient) and secured to a clamp stand using a clamp. Twenty 1.5 mL Eppendorf tubes were
opened and placed in a tube rack beneath the gradient in a biosafety cabinet. The QuickSeal™
tube was punctured at the 40-60% Iodixanol interface with an 18 g needle, using the bevel
pointing up, and then the top of the tube was punctured with a 16 g needle. Factions of 1 mL
were collected until the 25-45% interface was reached. 5 µL was removed from each fraction for
analysis by qPCR to locate AAV-rich fractions before storing the remainder of each fraction at
4°C overnight. Each 5 µL fraction sample was treated with DNase I (cat #M6101, Promega) for
1 h at 37°C and then diluted 1:20 in 50 µg/mL yeast tRNA (cat #AM7119, Invitrogen). A known
concentration of plasmid DNA used to prepare the virus was also diluted 1:200 in 50 µg/mL
yeast tRNA and processed in parallel. AAV-containing fractions were pooled, typically giving a
volume of 4-10 ml.
238
To concentrate the pooled fractions via buffer exchange, 12 mL of 0.1% Pluronic F68 in 1x PBS
was added to a new Amicon Ultra-15 100K Centrifugal Filter Unit (cat. #UFC910024, EMD
Millipore) and allowed to incubate at room temperature for 10 min. The Pluronic F68 in PBS was
removed from the filtration unit and 12 mL of 0.01% Pluronic F68 in 1x PBS was added to the
filtration unit before spinning at 3,000 RPM for 5 min at 4°C. The flow through was discarded
and 12 mL of 0.001% Pluronic F68 in 1x PBS + 200 mM NaCl was added and spun at 3000
RPM for 5 min at 4 °C. The flow-through was discarded and up to 12 mL of sample was added
to the column and spun at 3500 RPM for 8 min at 4°C. The flow through was discarded and
more sample was added up to 12 mL, mixed well with the concentrated remaining sample and
spun again. This continued until the sample was concentrated to a total final volume of 0.5-1
mL. The sample was washed with 10 mL of 0.001% Pluronic F68 in 1x PBS + 200 mM NaCl,
using a pipette to mix thoroughly, and then spun again. The sample was concentrated to about
0.5 mL, collected using a pipette, and the walls washed with a small volume of 0.001% Pluronic
F68 in 1x PBS + 200 mM NaCl to ensure all virus was collected. The concentrated virus was
stored at 4°C for up to 2 wk, or at -80°C for long term storage.
254

Using 5 µL of the concentrated virus, purified AAV was treated with DNase I for 30-60 min at
37°C to eliminate any contaminating plasmid DNA remaining from the viral purification process.
The DNase-treated viral samples were serially diluted 1:20 a total of five times in dilution buffer
(50 µg/mL Yeast tRNA, 0.05% Pluronic F68) and the reference plasmid (used to prepare the
virus) was serially diluted 1:10 a total of five times in dilution buffer. qPCR was performed as
indicated above, using each plasmid dilution series to construct a calibration curve used to
determine the titer of the virus.
262
Liver RNA extraction and qPCR analysis. Total liver RNA was extracted from frozen mouse
liver with TRIzol reagent (Invitrogen Life Technologies Inc., Carlsbad, CA), and nuclear RNA
was purified from nuclei isolated from frozen liver tissue and then extracted with TRIzol LS as
detailed elsewhere (37). RNA (1 µg) was treated with DNase I (cat. #M6101, Promega) to
remove DNA contamination. cDNA was then synthesized using High-Capacity cDNA Reverse
Transcription Kit (cat. #4368814, Applied Biosystems). qPCR was performed using primers
specific to each RNA, designed using Primer Express or Primer3 software
(http://bioinfo.ut.ee/primer3-0.4.0/), as follows: Sult3a1: 5'-
CATTGTCACATATCCAAAGTCTGGT-3' (forward), 5'-GAAAGGATCTGCTGGGTCCA-3'
(reverse) (oligonucleotides# 5735-5736); A1bg: 5'-GAACCCTCTGAGCCCAGTGA-3' (forward),
5'- GAGTGGGTGGAGCCTGTGAG-3' (reverse) (oligonucleotides# 5723-5724); Cux2: 5'-
CCTCAAGACGAACACCGTCAT-3' (forward), 5'-GCGCATCCTGGACCTGTAGT-3' (reverse)
(oligonucleotides# 1382-1383). Quantitative real-time PCR was carried out on a CFX384 Touch
Real-Time PCR Detection System (Bio-Rad) using Power SYBR Green PCR Master Mix
(ThermoFisher). Normalized linear Ct numbers based on 18S RNA as control were computed to
determine the relative expression level of each gene across treatments.
279
RNA-seq analysis. Polyadenylated mRNA was isolated from 1 µg of total liver RNA or from 1
µg of liver nuclear RNA from each treatment group (see below) using NEBNext Poly(A) mRNA
Magnetic Isolation Module (cat. #E7490, New England Biolabs). The resulting polyA-selected
RNA was used to prepare RNA-seq libraries using the NEBNext Ultra™ Directional RNA Library
Prep Kit for Illumina (cat. #E7420L, New England Biolabs), NEBNext Multiplex Oligos for
Illumina® Dual Index Primers (cat. #E7600, New England Biolabs), and AMPure XP Beads (cat.
#A63881, Beckman Coulter Inc., Indianapolis, IN). RNA-seq libraries were prepared from two
independent studies: 1) Sample series #G169: livers from 10-12 wk old male C57Bl/6 mice
treated with 0.75 x 10^{11} GC of AAV8-STAT5$_{CA}$ or AAV8-Null control then excised 2 wk later, with
two sequencing libraries prepared for each treatment group, each derived from polyA-selected
total liver RNA pooled from n = 2 or 3 individual livers (biological replicates); and 2) Sample
series #G186: livers from 7-8 wk old male CD1 mice treated with 2 x 10^{11} GC of AAV8-STAT5$_{CA}$
or control then excised 4-6 wk later, with a total of 9 sequencing libraries prepared, each one
from an individual liver (n=5 AAV8-STAT5$_{CA}$ libraries, n=4 control libraries) and derived from
polyA-selected nuclear RNA prepared from nuclei extracted from frozen liver tissue, to increase
the sensitivity for detection of the large fraction of lncRNAs that are nuclear-enriched and tightly
bound to chromatin in mouse liver (37). Libraries were multiplexed and sequenced by
Novogene, Inc (Sacramento, CA) to an average depth of 18-25 million 150 nt paired-end
sequence reads on an Illumina HiSeq instrument. RNA-seq data was analyzed using a modified
version of the custom pipeline described earlier (15). Briefly, sequence reads were aligned to
mouse genome build mm9 (NCBI 37) using STAR aligner (38) and FeatureCounts (39) was
used to count sequence reads mapping to the union of the exonic regions in all isoforms of a
given gene (collapsed exon counting) based on an mm9 Gene Transfer Format file comprised
of 75,798 mouse genes: n = 20,884 RefSeq protein coding genes, n=48,360 mouse liver-
expressed lncRNAs genes (a much more complete listing than the set described previously in
(40)), n=2,061 RefSeq non-coding genes (NR accession numbers) that do not overlap the set of

48,360 lncRNAs, n =4,490 Ensembl non-coding lncRNAs that do not overlap either the RefSeq
NR gene set or the 48,360 lncRNA gene set, and n=3 other lncRNAs with interesting liver
functions (lnc-LFAR1, LeXis, Lnclgr) (41) and in the gene annotations provided in **Table S1**.
Raw sequencing files and processed data files are available at GEO
(https://www.ncbi.nlm.nih.gov/geo/) under accession numbers GSE196014 and GSE196015.

Differential expression analysis. EdgeR (42) was used to determine significant differential
expression of RNA-seq data for the following comparisons: Males + AAV8-STAT5$_{CA}$ at 2×10^{11}
GC vs. Control males, and Males + AAV8-STAT5$_{CA}$ at 0.75×10^{11} GC vs. AAV8-null males, with
full datasets shown in **Tables S1-S3**. Further, a set of 475 sex-biased liver-expressed genes
was defined as genes with male liver vs female liver differential expression significant at FDR <
0.01 (**Table S2**). A set of 8,246 liver-expressed genes whose differential expression is
stringently sex-independent was defined based on |fold-change| for sex-difference < 1.2 and
FDR > 0.1 (**Table S3**). All analyses reported here for sex-independent genes used this list of
8,246 genes. Finally, for female-biased genes, a percent feminization value was calculated
based on each gene's response to each of the following treatments: AAV8-STAT5$_{CA}$ 2×10^{11}
GC, AAV8-STAT5$_{CA}$ 0.75×10^{11} GC, and continuous infusion of male mice with GH (cGH) for 7
or 14 d (20), as follows: % feminization = 100% [FPKM (treated male) – FPKM (control
male)]/[FPKM (control female) – FPKM (control male)] (**Table S2**).

Raw RNA-seq data obtained from livers of hypophysectomized mice (15) (GEO accession
number GSE66003) was re-analyzed to obtain expression data for the full set of 75,798 mouse
genes, including 48,360 liver-expressed lncRNAs, described above. Two distinct classes of sex-
biased genes were identified based on differential expression between livers of control and
hypophysectomized male and female mice at FDR <0.05 (22): class I male-biased genes are
genes that show significantly decreased expression in male liver after hypophysectomy due to
their dependence on the male plasma GH pulsatile pattern for expression; and class II male-
biased genes are genes that are repressed by the female pituitary profile and, consequently,
they are significantly induced in female liver following hypophysectomy. We also identified
corresponding sets of class I female-biased genes as those whose expression in female liver
decreases significantly after hypophysectomy due to a requirement for the female, near
continuous GH pattern for expression, and class II female-biased genes whose expression is
repressed by the male pituitary profile and consequently show significant de-repression in
hypophysectomized male liver (see Fig. 1 of (43)). The enrichment of STAT5$_{CA}$-responsive
genes in each set of hypophysectomy-responsive sex-biased genes was calculated compared
to that of a background set of the corresponding class of STAT5$_{CA}$-unresponsive sex-biased
genes. Statistical significance was evaluated using Fisher exact test.

Enrichment of STAT5 ChIP-seq binding sites at STAT5$_{CA}$-responsive genes. Male-enriched,
female-enriched and sex-independent STAT5 binding sites (1,765, 1,790, and 11,531 sites,
respectively) were obtained from our published ChIP-seq data for male and female mouse liver
(9) (GEO accession number GSE31578). Each STAT5 binding site was mapped to the closest
gene using the command bedtools -closest (44). All genes with a STAT5 binding site within 50
kb of the gene body were considered to be a STAT5 target. The enrichment of STAT5$_{CA}$-
responsive genes in each set of STAT5 binding site target genes was calculated by comparison
to a background set of STAT5$_{CA}$-unresponsive genes. Statistical significance was evaluated
using Fisher exact test.

Histology. Fresh liver tissue was submerged into 4% PFA and snap frozen. A portion of each
liver was then placed in phosphate-buffered saline with 0.05% NaN$_3$ for frozen tissue sectioning,
and the remainder was transferred to 70% ethanol for paraffin embedding and H&E staining.

Results
358
Constitutively active STAT5b feminizes liver gene expression. GH-activated JAK2
catalyzes tyrosine phosphorylation linked to the activation and nuclear translocation of STAT5,
which enables transcriptional activation of STAT5 target genes. GH-activated liver STAT5 is
known to be essential for sex-biased liver gene expression; however, GH also activates other,
STAT5-independent signaling pathways (24-26), which could contribute to the regulation of sex-
biased gene expression. To determine whether the control of sex-biased liver gene expression
by plasma GH pulses (male GH pattern) vs persistent GH stimulation (female GH pattern) is
due to the downstream effects of pulsatile vs persistent STAT5 activation *per se*, we expressed
a constitutively active form of STAT5 (STAT5$_{CA}$) in male mouse liver to mimic the pattern of
persistent STAT5 activation that occurs in female mouse liver (9). STAT5$_{CA}$ cDNA expressed
from the hepatocyte-specific thyroxin binding globulin promoter (45,46) was delivered using an
AAV serotype 8 vector, which has high intrinsic tropism for the liver (30,31). We validated the
AAV8-STAT5$_{CA}$ viral vector by its ability to target STAT5 protein to the nucleus (**Fig. 1A**), which
is a key feature of activated STAT5, and by its functional ability to increase hepatic *Igf1* RNA
levels and circulating IGF1 in mice deficient in GH receptor signaling in hepatocytes (**Fig. 1B,**
aHepGHRkd mouse liver).
375
Next, we investigated the impact of STAT5$_{CA}$ on sex-biased gene expression in the liver. We
delivered AAV8-STAT5$_{CA}$ (2 x 10^{11} GC/mouse) or a control virus (AAV8-Luciferase) to young
adult male mice and excised the livers 4-6 wk later. Liver gene expression was analyzed by
nuclear RNA-seq, which enabled us to determine the effects of STAT5$_{CA}$ on the nuclear
transcriptome, including lncRNA genes, which are highly enriched in the nucleus owing to their
tight binding to chromatin (37) (**Table S1**). AAV8-STAT5$_{CA}$ substantially feminized the
expression of liver sex-biased genes. Thus, 154 (59%) of 260 female-biased genes were
induced and only 2 female-biased genes were repressed by AAV8-STAT5$_{CA}$ in male liver.
Furthermore, 114 (53%) of 215 male-biased genes were repressed and only 5 male-biased
genes were induced by AAV8-STAT5$_{CA}$ (**Fig. 2A**, **Table S2**). In contrast, only 1.1% of stringent
sex-independent genes (**Table S3**) were induced and 0.7% were repressed by AAV8-STAT5$_{CA}$.
These trends, presented for the CD1 mouse liver model used extensively in our studies on liver
sex differences (15,19,20,47), were confirmed in an independent study in C57BL6 mice, which
were treated with AAV8-STAT5$_{CA}$ at a lower dose and for a shorter duration (0.75 x 10^{11} GC for
2 wk) (**Fig. 2A,** bottom; **Table S2**). Feminization of the liver transcriptome by AAV8-STAT5$_{CA}$
was dose-dependent, with partial feminization observed at virus doses as low as 0.125 x 10^{11}
GC, as shown for three highly female-biased genes (**Fig. 2B**). Sex-biased genes responsive to
AAV8-STAT5$_{CA}$ were enriched for established sex-biased metabolic pathways, whereas the set
of sex-independent genes responsive to AAV8-STAT5$_{CA}$ was enriched for extracellular
matrix/adhesion and glycoprotein/signal peptide genes (**Fig. 2C, Fig. 2D**).
396
Role of STAT5 in liver feminization induced by continuous GH infusion (cGH). cGH
treatment overrides the endogenous male, pulsatile GH secretion pattern and progressively
feminizes liver gene expression over a period of days (20). To understand the role of STAT5 in
this process, we compared the changes in gene expression following AAV8-STAT5$_{CA}$ treatment
(2 x 10^{11} GC) to the changes seen in livers of male mice given cGH infusion for 7 or 14 days.
AAV8-STAT5$_{CA}$ and cGH infusion induced very similar changes in the expression of sex-biased
genes in male liver (**Fig. 3A, Table S2**). Thus, 90% of the female-biased genes induced by
AAV8-STAT5$_{CA}$ were also induced by cGH (139/154 genes) and 81% of the male-biased genes
repressed by AAV8-STAT5$_{CA}$ were repressed by cGH (92/114 genes) (**Fig. 3B**). Overall, AAV8-
STAT5$_{CA}$ and cGH were both highly effective in feminizing liver gene expression, with median
feminization values of 87% and 116%, respectively (**Fig. 3C, Table S2**). Female-biased genes

408 that are highly induced (> 100-fold) by cGH infusion or by constitutive activation of STAT5
409 include the cytochrome P450 genes *Cyp3a16, Cyp3a41b, Cyp2c69,* and *Cyp2a4,* and other
410 metabolic genes, such as *Fmo3* and *Hao2.* Of note, 62 female-biased genes were induced in
411 male liver by cGH infusion but not by $STAT5_{CA}$ and 62 male-biased genes were repressed in
412 male liver by cGH infusion but not by $STAT5_{CA}$ (**Fig. 3B, Table S2**). These genes may be
413 regulated by GH signaling pathways independent of STAT5 (25). Finally, four male-biased
414 genes that were repressed by cGH were unexpectedly induced by AAV8-$STAT5_{CA}$ (*Chrna2,*
415 *Fmn2, Nuggc, lnc5159*; **Fig. 3A**, **Table S2**).

417 **$STAT5_{CA}$ regulates many direct STAT5 gene targets** - We identified an initial list of potential
418 STAT5 target genes, defined as genes that met the following three criteria: 1) gene expression
419 shows a significant change upon loss of STAT5, as seen in either male or female STAT5-
420 deficient mouse liver (12,19); 2) gene expression is induced or repressed in either male or
421 female liver following pituitary hormone ablation by hypophysectomy, which abolishes GH-
422 dependent STAT5 activation (15); and 3) gene expression is restored when hypophysectomized
423 mice are given an exogenous pulse of GH, which restores pulsatile STAT5 signaling in male
424 liver (15). We filtered out genes showing an inconsistent response to GH treatment or STAT5
425 deficiency to obtain a final list of 227 putative STAT5 target genes, of which 115 showed sex-
426 biased expression (**Table S4**). Importantly, 40% (90 of 227) of the predicted STAT5 target
427 genes and 45% (52 of 115) of the sex-biased predicted STAT5 targets were responsive to
428 AAV8-$STAT5_{CA}$ (**Table S4**, column AC). These $STAT5_{CA}$-responsive STAT5 targets both sex-
429 biased genes and well-established sex-independent STAT5 gene targets, such as *Igf1, Socs2,*
430 *Cish* and *Onecut1*, whose expression was increased in AAV8-$STAT5_{CA}$-treated male mouse
431 liver.

433 Next, we investigated whether responsiveness to AAV8-$STAT5_{CA}$ is associated with local
434 STAT5 binding to liver chromatin. We mapped liver STAT5 binding sites identified by chromatin
435 immunoprecipitation (ChIP-seq) (9) to their putative target genes (nearest gene within 50 kb;
436 **Table S2, Table S3**). These sites include many genomic regions where STAT5 binding to liver
437 chromatin is significantly stronger in males than in females, and vice versa (male-enriched and
438 female-enriched STAT5 binding sites, respectively) (9). For each gene class of interest, we
439 determined the enrichment of STAT5 binding in the set of genes responsive to AAV8-$STAT5_{CA}$
440 compared to the set of genes that were unresponsive to AAV8-$STAT5_{CA}$. We observed a strong,
441 significant enrichment of female-biased STAT5 binding at the set of female-biased genes
442 induced by $STAT5_{CA}$ as compared to the corresponding background set of female-biased genes
443 not induced by $STAT5_{CA}$ (**Fig. 4A**, **Table S5**). Male-biased genes repressed by $STAT5_{CA}$ were
444 depleted of STAT5 binding, but not significantly. The strong enrichment of female-biased STAT5
445 binding at female-biased genes induced by $STAT5_{CA}$ indicates that it is the persistence of
446 STAT5 binding, acting in a positive regulatory manner, that induces these female-biased genes
447 in male liver. By contrast, we observed strong enrichment of sex-independent STAT5 binding at
448 sex-independent genes that are repressed by $STAT5_{CA}$, but not at sex-independent genes
449 induced by $STAT5_{CA}$ (**Fig. 4A**). Thus, in the genomic context of these sex-independent genes,
450 STAT5 can act as a repressor, consistent with studies in GH-deficient rat liver (48). Finally, sex-
451 independent genes induced by $STAT5_{CA}$ were enriched for male-biased STAT5 binding,
452 consistent with the positive regulatory potential of those sites evident from our earlier work (9).

454 **$STAT5_{CA}$ preferentially induces class I female-biased genes.** In principle, $STAT5_{CA}$ may act
455 by either of two distinct mechanisms. By mimicking the actions of a persistent, female-like
456 plasma GH profile, $STAT5_{CA}$ may induce expression of female-biased genes that are positively
457 regulated by the female GH pattern, i.e., class I female-biased genes. Alternatively, by
458 eliminating the pulsatile nature of STAT5 signaling that is ongoing in male liver (15), $STAT5_{CA}$

may effectively override the inhibitory activity that endogenous male plasma GH pulses exert on
class II female-biased genes and thereby de-repress their expression (22). To distinguish these
two models, we compared the hypophysectomy response class frequency of the female-biased
genes that are induced by STAT5$_{CA}$ to those that are not induced by STAT5$_{CA}$ (**Table S2**, **Table
S5**). We found that the set of STAT5$_{CA}$-induced female-biased genes is significantly enriched in
class I female-biased genes; furthermore, it is significantly depleted of class II female-biased
genes (**Fig. 4B**, left). No significant hypophysectomy class enrichment was seen for STAT5$_{CA}$-
repressed male-biased genes. Sex-independent genes induced by STAT5$_{CA}$ showed a highly
significant 7-fold depletion of genes whose expression was induced when GH was ablated by
hypophysectomy, whereas sex-independent genes repressed by STAT5$_{CA}$ showed a tendency
toward enrichment for genes induced by hypophysectomy (**Fig. 4B,** right), both consistent with
the GH mimic nature of STAT5$_{CA}$.
471
Dose-dependent effects of AAV8-STAT5$_{CA}$ on liver histology and *Igf1* expression. There
were no consistent changes in hepatocyte morphology when AAV8-STAT5$_{CA}$ was delivered at
doses up to 0.75×10^{11} GC or in livers from mice given cGH infusion. However, when the dose
of AAV8-STAT5$_{CA}$ was increased to 2×10^{11} GC/mouse, we observed architectural distortion,
hepatocyte hyperplasia, binuclear hepatocytes, karyomegaly and nuclear atypia (**Fig. 5**). To
ascertain whether these responses are associated with STAT5 overexpression, we examined
the expression levels of two classical STAT target genes in liver, *Igf1* and *Socs2.* Whereas
AAV8-STAT5$_{CA}$ induced the expression of *Igf1* moderately (2-2.5-fold increase above control
liver levels at both doses), we observed strong induction of *Socs2,* which reached 8.5-fold at the
lower dose and 14.4-fold above baseline at the higher dose (2×10^{11} GC), consistent with a
functional overexpression of STAT5 activity at the higher AAV8-STAT5$_{CA}$ dose (**Fig. 6**).
483
484   **Discussion**
485
The sex-dependent patterns of pituitary GH secretion – pulsatile GH release in males and
persistent (near-continuous) GH release in females – activate hepatic GH receptor signaling to
STAT5 in a manner that mirrors circulating GH profiles, i.e., pulsatile STAT5 activation and
transcriptional activity in male liver, and persistent STAT5 activation in female liver. Here we
show that persistent expression of activated STAT5, achieved using a liver-specific AAV8 viral
vector to deliver the activated STAT5b mutant STAT5$_{CA}$ to mouse hepatocytes *in vivo*, has
extensive effects on sex-biased gene expression in the liver, with large numbers of male-biased
genes repressed and female-biased genes induced. These effects are highly selective for sex-
biased genes, insofar as fewer than 2% of genes showing sex-independent expression were
impacted by STAT5$_{CA}$ delivery. Many of the sex-independent gene responses are likely to be
specific, in view of their enrichment for specific biological pathways related to extracellular
matrix/cell adhesion and Wnt signaling.
498
Continuous infusion of GH in male mice (cGH treatment) mimics the endogenous female pattern
of persistent circulating GH and overrides the endogenous pulsatile male secretion pattern,
leading to substantial feminization of liver gene expression within 7 days (20,21). Here, we
report a high overlap between gene responses to STAT5$_{CA}$ expression and cGH infusion,
providing strong evidence that persistent STAT5 activity alone is sufficient to account for a large
fraction of the feminized genes. Nevertheless, some genes responded in a unique manner to
either STAT5$_{CA}$ expression or cGH infusion. While some of the differences in response to these
two feminizing treatments may be due to cutoff thresholds, others clearly are not, for example
Cyp2b10 and *Cyp2c37*, whose expression in male liver was induced, and substantially
feminized, by cGH infusion but which were unresponsive to STAT5$_{CA}$ (**Table S2**). Differences
between these two models for feminizing male liver gene expression may in part be due to

STAT5-independent signaling pathways that are initiated by continuous GH, such as the GH
receptor-dependent activation of PI-3 kinase, ERK signaling and Src family kinases (25).
512
STAT5$_{CA}$ and cGH infusion both increased the expression of some female-biased genes to a
level that exceeds levels of expression seen in untreated female liver. This phenomenon was
somewhat more pronounced with cGH infusion (29 genes expressed at >2-fold the level of
female liver after 14 days of cGH infusion vs. 18 genes exceeded that level following STAT5$_{CA}$
treatment; **Table S2**). There are several possible reasons for the excessive feminization
observed for these genes, and for the incomplete feminization observed for many other sex-
biased genes. First, levels of STAT5 activity reached in AAV8-STAT5$_{CA}$-infected hepatocytes
may exceed the physiological levels of activated STAT5 in female liver, in particular at the
higher dose of AAV8-STAT5$_{CA}$ used in our study (c.f., overexpression of *Socs2*; **Fig. 6**). Related
to this, both treatments used in this study, cGH infusion and expression of a persistently
activated STAT5, differ somewhat from the physiological patterns in female liver, where there
are short GH-off time periods, albeit not as prolonged or as frequent as the GH-off inter-pulse
intervals that characterize males. Second, although STAT5$_{CA}$ and cGH may both mimic female
liver with respect to STAT5 activity and GH stimulation, they do not recapitulate the female
hormonal environment with respect to other factors. One such factor is estrogen, which can also
impact sex-biased gene expression in the liver, albeit it affects only 4% of liver sex-biased
genes in a direct estrogen receptor-dependent manner (49). Indeed, it is remarkable how
effective STAT5$_{CA}$ and cGH are in feminizing male liver, despite the presence of a male gonadal
steroid environment. Finally, intrinsic chromosomal sex differences associated with liver-
expressed Y-chromosome-encoded epigenetic modifiers, such as *Uty* and *Kdm5d,* and partial
X-inactivation/incomplete gene dosage compensation remain (50) and could also contribute to
the observed differences between female liver and STAT5$_{CA}$-treated or cGH-infused male liver.
535
Female-biased genes induced by STAT5$_{CA}$ in male liver were significantly enriched for nearby
female-biased STAT5 binding sites in liver chromatin when compared to the set of female-
biased genes that were not induced by STAT5$_{CA}$. Moreover, the STAT5$_{CA}$-induced female-
biased genes were significantly enriched for class I female-biased genes, whose expression in
female liver is positively regulated by the female plasma GH pattern. Together, these findings
lend strong support to the proposal that the de-repression of these female-biased genes in male
liver involves direct binding of STAT5$_{CA}$ to these STAT5-binding regulatory sites in association
with induction of local, female-biased chromatin opening, which has been linked to female-
biased STAT5 binding and gene activation (9,17). Female-biased genes not induced by
STAT5$_{CA}$ could be regulated by other mechanisms, including GH signaling independent of
STAT5, as noted above. We did not observe a corresponding enrichment of STAT5 binding
sites nearby male-biased genes repressed by STAT5$_{CA}$, indicating that factors other than
differences in STAT5 binding determine whether an individual male-biased gene will be
repressed by STAT5$_{CA}$ expression in male liver. Finally, sex-independent genes repressed by
STAT5$_{CA}$ showed strong enrichment for local sex-independent STAT5 binding sites, supporting
the proposal that STAT5 can repress gene expression by a direct DNA binding mechanism.
552
We observed significant feminization of gene expression at AAV8-STAT5$_{CA}$ doses as low as
0.125 x 10^{11} GC per mouse, i.e., 10-fold or more lower than standard doses of 2 x 10^{11}
commonly used for AAV8-induced gene expression in mouse liver (31). This finding is in line
with earlier studies in a GH-deficient rat model, where infusion of GH at a dose that results in as
low as 3% of the nominal circulating female GH level restores the expression of certain female-
biased genes, while other genes require higher GH doses and show only partial induction (51).
559

It is well established that a reduction of GH signaling is associated with liver steatosis and development of non-alcoholic steatohepatitis, both in humans and in mouse models, and increasing GH signaling can improve fatty liver endpoints (52,53). However, we found that treatment with a high dose of AAV8-STAT5$_{CA}$ resulted in certain histopathological changes indicative of liver injury that cannot be attributed to the AAV8 infection *per se*, as they were not seen in AAV-Luciferase controls. Livers from mice given cGH infusion for 17 days did not show corresponding histopathological changes, despite the overall similarity of its effects on sex-biased gene expression. Conceivably, the histopathology associated with AAV8-STAT5$_{CA}$ at the high dose could be related to the propensity of this viral vector to integrate into the host genome in the context of liver injury (54,55) or could be a supraphysiologic effect of STAT5 overexpression. Nonetheless, these observations highlight the importance of carefully evaluating such effects before considering STAT5$_{CA}$ or other STAT5 derivatives for therapeutic use in treating liver disease.

Acknowledgements – The authors thank the following Waxman lab members: Kritika Karri for providing annotations for the 48,360 mouse liver-expressed lncRNAs genes examined in this study, Dr. Maxim Pyatkov for RNA-seq analysis pipeline development, and Dr. Ravi Sonkar for assistance with histology imaging.

Data availability. All data generated or analyzed during this study are included in this published article or in the associated Supplemental tables and the public data repository at GEO (https://www.ncbi.nlm.nih.gov/geo/) under accession numbers GSE196014 and GSE196015.

References

1. Waxman DJ, Holloway MG. Sex differences in the expression of hepatic drug metabolizing enzymes. *Mol Pharmacol*. 2009;76(2):215-228.

2. Kur P, Kolasa-Wolosiuk A, Misiakiewicz-Has K, Wiszniewska B. Sex Hormone-Dependent Physiology and Diseases of Liver. *Int J Environ Res Public Health*. 2020;17(8).

3. Lonardo A, Nascimbeni F, Ballestri S, Fairweather D, Win S, Than TA, Abdelmalek MF, Suzuki A. Sex Differences in Nonalcoholic Fatty Liver Disease: State of the Art and Identification of Research Gaps. *Hepatology*. 2019;70(4):1457-1469.

4. Brie B, Ramirez MC, De Winne C, Lopez Vicchi F, Villarruel L, Sorianello E, Catalano P, Ornstein AM, Becu-Villalobos D. Brain Control of Sexually Dimorphic Liver Function and Disease: The Endocrine Connection. *Cell Mol Neurobiol*. 2019;39(2):169-180.

5. Waxman DJ, Pampori NA, Ram PA, Agrawal AK, Shapiro BH. Interpulse interval in circulating growth hormone patterns regulates sexually dimorphic expression of hepatic cytochrome P450. *Proc Natl Acad Sci U S A*. 1991;88(15):6868-6872.

6. Mode A, Wiersma-Larsson E, Strom A, Zaphiropoulos PG, Gustafsson JA. A dual role of growth hormone as a feminizing and masculinizing factor in the control of sex-specific cytochrome P-450 isozymes in rat liver. *J Endocrinol*. 1989;120(2):311-317.

7. Waters MJ. The growth hormone receptor. *Growth Horm IGF Res*. 2016;28:6-10.

8. Piwien-Pilipuk G, Huo JS, Schwartz J. Growth hormone signal transduction. *J Pediatr*
Endocrinol Metab. 2002;15(6):771-786.

9. Zhang Y, Laz EV, Waxman DJ. Dynamic, sex-differential STAT5 and BCL6 binding to sex-
biased, growth hormone-regulated genes in adult mouse liver. *Mol Cell Biol*. 2012;32(4):880-
896.

10. Chia DJ. Minireview: mechanisms of growth hormone-mediated gene regulation. *Mol*
Endocrinol. 2014;28(7):1012-1025.

11. Clodfelter KH, Holloway MG, Hodor P, Park SH, Ray WJ, Waxman DJ. Sex-dependent liver
gene expression is extensive and largely dependent upon signal transducer and activator of
transcription 5b (STAT5b): STAT5b-dependent activation of male genes and repression of
female genes revealed by microarray analysis. *Mol Endocrinol*. 2006;20(6):1333-1351.

12. Holloway MG, Cui Y, Laz EV, Hosui A, Hennighausen L, Waxman DJ. Loss of sexually
dimorphic liver gene expression upon hepatocyte-specific deletion of Stat5a-Stat5b locus.
Endocrinology. 2007;148(5):1977-1986.

13. Choi HK, Waxman DJ. Plasma growth hormone pulse activation of hepatic JAK-STAT5
signaling: developmental regulation and role in male-specific liver gene expression.
Endocrinology. 2000;141(9):3245-3255.

14. Tannenbaum GS, Choi HK, Gurd W, Waxman DJ. Temporal relationship between the sexually
dimorphic spontaneous GH secretory profiles and hepatic STAT5 activity. *Endocrinology*.
2001;142(11):4599-4606.

15. Connerney J, Lau-Corona D, Rampersaud A, Waxman DJ. Activation of Male Liver Chromatin
Accessibility and STAT5-Dependent Gene Transcription by Plasma Growth Hormone Pulses.
Endocrinology. 2017;158(5):1386-1405.

16. Choi HK, Waxman DJ. Growth hormone, but not prolactin, maintains, low-level activation of
STAT5a and STAT5b in female rat liver. *Endocrinology*. 1999;140(11):5126-5135.

17. Sugathan A, Waxman DJ. Genome-wide analysis of chromatin states reveals distinct
mechanisms of sex-dependent gene regulation in male and female mouse liver. *Mol Cell Biol*.
2013;33(18):3594-3610.

18. Cui Y, Hosui A, Sun R, Shen K, Gavrilova O, Chen W, Cam MC, Gao B, Robinson GW,
Hennighausen L. Loss of signal transducer and activator of transcription 5 leads to
hepatosteatosis and impaired liver regeneration. *Hepatology*. 2007;46(2):504-513.

19. Hao P, Waxman DJ. STAT5 Regulation of Sex-Dependent Hepatic CpG Methylation at Distal
Regulatory Elements Mapping to Sex-Biased Genes. *Mol Cell Biol*. 2021;41(2).

20. Lau-Corona D, Suvorov A, Waxman DJ. Feminization of Male Mouse Liver by Persistent
Growth Hormone Stimulation: Activation of Sex-Biased Transcriptional Networks and
Dynamic Changes in Chromatin States. *Mol Cell Biol*. 2017;37(19).

21. Holloway MG, Laz EV, Waxman DJ. Codependence of growth hormone-responsive, sexually
dimorphic hepatic gene expression on signal transducer and activator of transcription 5b and
hepatic nuclear factor 4alpha. *Mol Endocrinol*. 2006;20(3):647-660.

22. Wauthier V, Sugathan A, Meyer RD, Dombkowski AA, Waxman DJ. Intrinsic sex differences
in the early growth hormone responsiveness of sex-specific genes in mouse liver. *Mol
Endocrinol*. 2010;24(3):667-678.

23. Rotwein P. Regulation of gene expression by growth hormone. *Mol Cell Endocrinol*.
2020;507:110788.

24. Chhabra Y, Lee CMM, Muller AF, Brooks AJ. GHR signalling: Receptor activation and
degradation mechanisms. *Mol Cell Endocrinol*. 2021;520:111075.

25. Frank SJ. Classical and novel GH receptor signaling pathways. *Mol Cell Endocrinol*.
2020;518:110999.

26. Carter-Su C, Schwartz J, Argetsinger LS. Growth hormone signaling pathways. *Growth Horm
IGF Res*. 2016;28:11-15.

27. List EO, Berryman DE, Funk K, Jara A, Kelder B, Wang F, Stout MB, Zhi X, Sun L, White TA,
LeBrasseur NK, Pirtskhalava T, Tchkonia T, Jensen EA, Zhang W, Masternak MM, Kirkland
JL, Miller RA, Bartke A, Kopchick JJ. Liver-specific GH receptor gene-disrupted (LiGHRKO)
mice have decreased endocrine IGF-I, increased local IGF-I, and altered body size, body
composition, and adipokine profiles. *Endocrinology*. 2014;155(5):1793-1805.

28. Ariyoshi K, Nosaka T, Yamada K, Onishi M, Oka Y, Miyajima A, Kitamura T. Constitutive
activation of STAT5 by a point mutation in the SH2 domain. *J Biol Chem*. 2000;275(32):24407-
24413.

29. Woelfle J, Billiard J, Rotwein P. Acute control of insulin-like growth factor-I gene transcription
by growth hormone through Stat5b. *J Biol Chem*. 2003;278(25):22696-22702.

30. Zincarelli C, Soltys S, Rengo G, Rabinowitz JE. Analysis of AAV serotypes 1-9 mediated gene
expression and tropism in mice after systemic injection. *Mol Ther*. 2008;16(6):1073-1080.

31. Sands MS. AAV-mediated liver-directed gene therapy. *Methods Mol Biol*. 2011;807:141-157.

32. List EO, Berryman DE, Funk K, Gosney ES, Jara A, Kelder B, Wang X, Kutz L, Troike K,
Lozier N, Mikula V, Lubbers ER, Zhang H, Vesel C, Junnila RK, Frank SJ, Masternak MM,
Bartke A, Kopchick JJ. The role of GH in adipose tissue: lessons from adipose-specific GH
receptor gene-disrupted mice. *Mol Endocrinol*. 2013;27(3):524-535.

33. Cordoba-Chacon J, Majumdar N, List EO, Diaz-Ruiz A, Frank SJ, Manzano A, Bartrons R,
Puchowicz M, Kopchick JJ, Kineman RD. Growth Hormone Inhibits Hepatic De Novo
Lipogenesis in Adult Mice. *Diabetes*. 2015;64(9):3093-3103.

34. Percie du Sert N, Ahluwalia A, Alam S, Avey MT, Baker M, Browne WJ, Clark A, Cuthill IC,
Dirnagl U, Emerson M, Garner P, Holgate ST, Howells DW, Hurst V, Karp NA, Lazic SE,
Lidster K, MacCallum CJ, Macleod M, Pearl EJ, Petersen OH, Rawle F, Reynolds P, Rooney

K, Sena ES, Silberberg SD, Steckler T, Würbel H. Reporting animal research: Explanation
and elaboration for the ARRIVE guidelines 2.0. *PLoS Biol*. 2020;18(7):e3000411.

35. Wang J, Symul L, Yeung J, Gobet C, Sobel J, Luck S, Westermark PO, Molina N, Naef F.
Circadian clock-dependent and -independent posttranscriptional regulation underlies
temporal mRNA accumulation in mouse liver. *Proc Natl Acad Sci U S A*. 2018;115(8):E1916-
E1925.

36. Menet JS, Rodriguez J, Abruzzi KC, Rosbash M. Nascent-Seq reveals novel features of
mouse circadian transcriptional regulation. *Elife*. 2012;1:e00011.

37. Goldfarb CN, Waxman DJ. Global analysis of expression, maturation and subcellular
localization of mouse liver transcriptome identifies novel sex-biased and TCPOBOP-
responsive long non-coding RNAs. *BMC Genomics*. 2021;22(1):212.

38. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M,
Gingeras TR. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*. 2013;29(1):15-21.

39. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning
sequence reads to genomic features. *Bioinformatics*. 2014;30(7):923-930.

40. Lodato NJ, Melia T, Rampersaud A, Waxman DJ. Sex-Differential Responses of Tumor
Promotion-Associated Genes and Dysregulation of Novel Long Noncoding RNAs in
Constitutive Androstane Receptor-Activated Mouse Liver. *Toxicol Sci*. 2017;159(1):25-41.

41. Goldfarb CN, Karri K, Pyatkov M, Waxman DJ. Interplay between murine sex-biased gene
expression and hepatic zonation revealed by single nucleus RNA sequencing. *bioRxiv*.
2022:2022.2001.2018.476791.

42. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. *Bioinformatics*. 2010;26(1):139-140.

43. Melia T, Waxman DJ. Sex-Biased lncRNAs Inversely Correlate With Sex-Opposite Gene
Coexpression Networks in Diversity Outbred Mouse Liver. *Endocrinology*. 2019;160(5):989-
1007.

44. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics. 2010;26(6):841-842.

45. Kiourtis C, Wilczynska A, Nixon C, Clark W, May S, Bird TG. Specificity and off-target effects
of AAV8-TBG viral vectors for the manipulation of hepatocellular gene expression in mice.
Biol Open. 2021;10(9).

46. Wolf Greenstein A, Majumdar N, Yang P, Subbaiah PV, Kineman RD, Cordoba-Chacon J.
Hepatocyte-specific, PPARgamma-regulated mechanisms to promote steatosis in adult mice.
J Endocrinol. 2017;232(1):107-121.

47. Melia T, Hao P, Yilmaz F, Waxman DJ. Hepatic Long Intergenic Noncoding RNAs: High
Promoter Conservation and Dynamic, Sex-Dependent Transcriptional Regulation by Growth
Hormone. *Mol Cell Biol*. 2016;36(1):50-69.

48. Ono M, Chia DJ, Merino-Martinez R, Flores-Morales A, Unterman TG, Rotwein P. Signal
transducer and activator of transcription (Stat) 5b-mediated inhibition of insulin-like growth
factor binding protein-1 gene transcription: a mechanism for repression of gene expression
by growth hormone. *Mol Endocrinol*. 2007;21(6):1443-1457.

49. Zheng D, Wang X, Antonson P, Gustafsson JA, Li Z. Genomics of sex hormone receptor
signaling in hepatic sexual dimorphism. *Mol Cell Endocrinol*. 2018;471:33-41.

50. Naqvi S, Godfrey AK, Hughes JF, Goodheart ML, Mitchell RN, Page DC. Conservation,
acquisition, and functional impact of sex-biased gene expression in mammals. *Science*.
2019;365(6450).

51. Pampori NA, Shapiro BH. Gender differences in the responsiveness of the sex-dependent
isoforms of hepatic P450 to the feminine plasma growth hormone profile. *Endocrinology*.
1999;140(3):1245-1254.

52. Vazquez-Borrego MC, Del Rio-Moreno M, Kineman RD. Towards Understanding the Direct
and Indirect Actions of Growth Hormone in Controlling Hepatocyte Carbohydrate and Lipid
Metabolism. *Cells*. 2021;10(10).

53. Kaltenecker D, Themanns M, Mueller KM, Spirk K, Suske T, Merkel O, Kenner L, Luis A,
Kozlov A, Haybaeck J, Muller M, Han X, Moriggl R. Hepatic growth hormone - JAK2 - STAT5
signalling: Metabolic function, non-alcoholic fatty liver disease and hepatocellular carcinoma
progression. *Cytokine*. 2019;124:154569.

54. Dalwadi DA, Torrens L, Abril-Fornaguera J, Pinyol R, Willoughby C, Posey J, Llovet JM,
Lanciault C, Russell DW, Grompe M, Naugler WE. Liver Injury Increases the Incidence of
HCC following AAV Gene Therapy in Mice. *Mol Ther*. 2021;29(2):680-690.

55. Deyle DR, Russell DW. Adeno-associated virus vector integration. *Curr Opin Mol Ther*.
2009;11(4):442-447.
737
738

Figure legends

Fig. 1. Functional validation of AAV8-STAT5$_{CA}$ *in vivo*. (A) Nuclear localization of Flag-Tag STAT5$_{CA}$ protein, detected by anti-Flag antibody on a Western blot of nuclear extracts from livers excised from mice 7 d after injection of AAV8-Null (control, lanes 1, 3) or AAV8-STAT5$_{CA}$ (1.5 x 10^{11} GC, lanes 2, 4). Extracts were prepared from four individual livers. (B) Plasma IGF1 levels and liver Igf1 and GH receptor (Ghr) mRNA levels in male mice, GH receptor floxed control (Null) male mice, or male mice with adult-onset GH receptor knockdown (aHepGHRkd mice), with or without AAV8-STAT5$_{CA}$ treatment at the doses indicated. Values shown are mean ± SEM for n = 8 mice per group. Significance was evaluated by one way ANOVA with Bonferroni multiple comparison correction, implemented in GraphPad Prism. For comparisons to aHepGHRkd (second bar in each set): p< 0.001, ***; p< 0.01, **; and p=0.0595, *. NS, p > 0.05 for comparisons to Null, indicating restoration of Igf1 RNA and plasma protein.

Fig. 2. **AAV8-STAT5$_{CA}$ responsive genes in male liver**. (A) Percentage of all genes that are either consistently sex biased at FDR < 0.05 (Table S2, n= 475 genes) or stringently sex-independent (SI) genes (Table S3, n = 8,246 genes) and whose expression in AAV8-STAT5$_{CA}$-treated liver (as indicated) is significantly induced or repressed compared to control liver (FDR < 0.05). Top panel: results for CD-1 male mice; bottom panel: results for C57BL/6 mice. GC, genome copies of AAV8 virus. Number of responding genes is shown above each bar. (B) Dose responses for AAV8-STAT5$_{CA}$ induction of three highly female-biased genes. Data based on RT-qPCR analysis of liver RNA, mean +/- SEM for n=2-6 livers per group, expressed as percentage of gene expression in control (untreated) female (F) mouse liver. M, male group. (C, D) DAVID pathway analysis of the sets of sex-biased genes (C) and sex-independent genes (D) that are responsive to STAT5$_{CA}$. Shown are the top five enriched clusters, with bars indicating -log10 P values (Benjamini-Hochberg corrected) for enrichment.

Fig. 3. **Effects of STAT5$_{CA}$ and cGH infusion on sex-biased gene expression**. (A) Heatmap showing impact of STAT5$_{CA}$ or cGH infusion on the expression of 275 responsive sex-biased genes (log2 Fold-change increases (red) or decreases (blue)) in male mouse liver. M, male; F, female. (B) Overlap between female-biased genes induced by AAV8-STAT5$_{CA}$ and those induced by cGH after either 7 or 14 days (left), and between male-biased genes repressed by AAV8-STAT5$_{CA}$ and those repressed by cGH after either 7 or 14 days (right). Excluded are 5 male-biased genes induced by AAV8-STAT5$_{CA}$ and 1 female-biased genes repressed by AAV8-STAT5$_{CA}$. (C) Boxplots showing the distribution of feminization values for 139 female-biased genes induced in male liver in both mouse models (see Table S2). Median value, horizontal line in each box.

Fig. 4. Enrichment of STAT5 binding sites (A) and hypophysectomy response class (B) for the sets of genes induced or repressed by STAT5$_{CA}$. (A) Data are shown for STAT5 binding sites that are male-biased (M; 1,765 sites), female-biased (F; 1,790 sites), and sex-independent (SI; 11,531 sites), as defined by ChIP-seq, and mapped to their target genes (nearest gene within 50 kb). Enrichments are expressed as the ratio of STAT5$_{CA}$-responsive genes compared to STAT5$_{CA}$-unresponsive genes that are associated with each of the 3 sets of STAT5 binding sites. (B) Enrichment of class I or class II sex-biased genes (left) and sex-independent genes (right) in the STAT5$_{CA}$-responsive gene sets, expressed as the ratio of the number of STAT5$_{CA}$-responsive genes compared to the number of STAT5$_{CA}$-unresponsive genes in each class (left) or that are induced or repressed by hypophysectomy (right). Significance was determined by Fishers exact test: *, p< 0.05, ***, P < 0.001. Blue asterisks, significant enrichment; red asterisks, significant depletion.

790   **Fig. 5**. **H&E staining of AAV8-STAT5$_{CA}$-infected liver**. Liver sections from mice treated with
791   AAV8-STAT5$_{CA}$ or AAV8-Luciferase at the indicated GC dose per mouse were stained with
792   H&E, revealing a STAT5$_{CA}$ dose-dependent histopathology, which was evident and most
793   consistent at the highest dose of AAV8 dose. No histopathology was evident in mice given cGH
794   infusion for 17 days. Images collected with an Olympus FSX-100 instrument at 16x.
795
796   **Fig. 6**. Expression of direct STAT5 target genes *Igf1* and *Socs2* in male liver: impact of cGH
797   infusion and AAV8-STAT5$_{CA}$. Shown are expression level determined by RNA-seq (fold-change
798   values relative to male control), indicating supraphysiological induction of *Socs2*, and to a lesser
799   extent *Igf1*, by AAV8-STAT5$_{CA}$. Significance was determined by edgeR: ***, adjusted p-value <
800   E-05.
801

A

Null	CA	Null	CA
1	2	3	4

B

Igf1 RNA

IGF1 mRNA (% to Null)

***

NS

*

Plasma IGF1

ng/ml

NS

**

**

Ghr RNA

GHR mRNA (% to Null)

***

☐ Null

■ aHepGHRkd

▨ aHepGHRkd + STAT5bCA 0.376x10^{11}GC

▨ aHepGHRkd + STAT5bCA 0.75x10^{11} GC

Fig. 2

A. STAT5 binding within 50 kb

Gene set / Response to STAT5$_{CA}$ in male liver

B. Hypophysectomy response class

Fig. 5

Fig. 6