Diverse innate immune factors protect yeast from lethal viral pathogenesis

Sabrina Chau[1], Jie Gao[1], Annette J. Diao[1], Shi Bo Cao[1], Alan R. Davidson[1,2], and Marc D. Meneghini[1*]

[1]Department of Molecular Genetics, University of Toronto, 661 University Ave, Toronto, ON, M5G

1M1, Canada.

[2]Department of Biochemistry, University of Toronto, 661 University Ave, Toronto, ON, M5G 1M1,

Canada.

[*]Correspondence: marc.meneghini@utoronto.ca

Summary

In recent years, newly characterized antiviral systems have been found to be remarkably conserved from

bacteria to mammals, revealing the potential to gain unique insight into these systems through the study

of microbial organisms[1]. Despite the enthusiasm generated by these findings, the key microbial model

organism *Saccharomyces cerevisiae* (budding yeast) has been minimally exploited for studies of viral

defense, primarily because it is not infected with exogenously transmitted viruses. However, most yeast

strains are infected with an endogenous double stranded RNA (dsRNA) virus called L-A, and previous

studies have identified conserved antiviral systems that attenuate L-A replication[2]. Although these

systems do not lead to eradication of L-A, we show here that they collaboratively prevent massive over-

proliferation of L-A that leads to proteostatic stress and lethality. Exploiting this new insight, we asked

if the powerful genetic screening methods available in yeast could be used to identify additional

conserved antiviral systems. Validating this approach, we discovered antiviral functions for the yeast

homologs of polyA-binding protein (PABPC1) and the La-domain containing protein Larp1, which are

both involved in viral innate immunity in humans[3,4]. We also identify new antiviral functions for the

RNA exonucleases *REX2* and *MYG1*, both of which have distinct but poorly characterized human and

bacterial homologs[5-8]. These findings highlight the potential of yeast as a powerful model system for

studies of the ubiquitous host-virus arms race.

Results and discussion

L-A belongs to the broadly dispersed *Totiviridae* family of double stranded RNA (dsRNA) viruses,

which has no extracellular phase and is transmitted vertically through cell division or fusion[2]. L-A

transcripts encode the capsid (Gag) proteins that comprise most of its viral particle as well as a Gag-pol

fusion protein produced at much lower levels that provides RNA dependent RNA polymerase activity

responsible for replication and transcription. Encapsidation of viral transcripts within nascent particles completes the L-A lifecycle[2]. L-A replication is maintained at a low level through the activities of the *SKI2*, *3*, and *8* genes, which encode subunits of a conserved ribosome-associated complex that opposes the translation of transcripts such as those encoded by L-A that lack polyA tails[9-11]. A separate pathway of L-A attenuation is provided by Xrn1, a 5'-3' exoribonuclease that degrades uncapped mRNAs[12,13]. Both of these systems have been associated with antiviral defense in mammals[14-17].

We showed that L-A was held in check during sporulation by the mitochondrial DNA/RNA endonuclease Nuc1, a homolog of endonuclease G (EndoG) found in all eukaryotes and many prokaryotes[18]. EndoG is involved in mammalian programmed cell death that frequently occurs in virus infected cells, and programmed cell death is intrinsic to yeast sporulation[19-22]. These and other observations have led to speculation that Nuc1/EndoG may reflect a prokaryotic antiviral origin of programmed cell death[23]. To query *NUC1* anti-viral function outside of sporulation and its associated programmed cell death, we investigated L-A attenuation in mitotic haploid cells in the reference BY4742 strain background. Visualization of L-A dsRNA using ethidium bromide staining of electrophoresed RNA revealed that while *nuc1Δ* and *ski3Δ* single mutants exhibited no effect, combining *nuc1Δ* and *ski3Δ* led to a large increase in L-A dsRNA **(Fig 1A)**. We corroborated these findings using immunofluorescence microscopy with a dsRNA antibody widely used to detect replicating RNA viruses[24,25]. These images revealed L-A dsRNA to accumulate in foci, reminiscent of "viral factory" sites of viral replication observed in human cells **(Fig 1B and S1)**[26]. Consistent with previous findings in other strain backgrounds, western blotting showed that Gag protein levels were elevated in *nuc1Δ* and *ski3Δ* mutants[12,18,27], and we determined that *nuc1Δ ski3Δ* double mutants accumulated massively elevated Gag levels, confirming that *NUC1* and *SKI3* participate in separate antiviral pathways **(Fig 2A)**.

We used spot test growth assays to determine if high L-A viral load had any fitness consequence and found that *nuc1Δ ski3Δ* double mutants grew normally at 30° C but exhibited conditional lethality at 37° C **(Fig 2B)**. Modest growth defects of *nuc1Δ* and *ski3Δ* single mutants were observed at high temperature when cells were grown with glycerol rather than glucose as the carbon source, a condition under which yeast relies on mitochondrial respiration **(Fig 2B)**. These growth phenotypes were completely reversed in isogenic strains cured of L-A (L-A⁰), implying that unrestricted L-A replication caused them **(Fig 2B)**.

As expected, a *NUC1*-expressing plasmid restored viability to a *nuc1Δ ski3Δ* strain grown at high temperature and elicited a corresponding decrease in Gag levels **(Fig 2A and 2B)**. We observed similar suppression of *nuc1Δ ski3Δ* using plasmid driven *XRN1* over-expression, suggesting that Xrn1 repressed

L-A via a parallel mechanism **(Fig 2A and 2B)**. Consistent with this model, *nuc1Δ xrn1Δ* mutants

accumulated greatly elevated levels of Gag and exhibited L-A-dependent conditional lethality at high

temperature **(Fig S2A and S2B)**, and *xrn1Δ ski3Δ* mutants are inviable[28]. We conclude that Nuc1, Ski3

and Xrn1 convergently oppose L-A, and that L-A causes lethal pathogenesis at high temperature when

unattenuated **(Fig 2C)**.

Since *XRN1* over-expression suppressed the growth defects of a *nuc1Δ ski3Δ* strain, we

hypothesized that over-expression of other antiviral factors would produce a similar effect, and that this

phenomenon could be used as a screen to identify new antiviral systems. To test this idea, we employed

a high-copy plasmid suppression screen to detect genes whose overexpression alleviated the conditional

lethality of *nuc1Δ ski3Δ*. Using this strategy, we identified *SRO9*, *SLF1*, and *PAB1* as high-copy

suppressors, all of which encode ribosome-associated RNA binding proteins **(Fig 3A)**[29,30]. Sro9 and Slf1

are paralogous lupus-autoantigen (La) domain containing proteins broadly found in eukaryotes.

Interestingly, the human Sro9/Slf1 ortholog Larp1 was recently found to bind to SARS-Cov-2 plus-

strand ssRNA and to oppose its replication in human cells through an unknown mechanism[3]. *PAB1*

encodes the highly conserved <u>P</u>oly<u>A</u>-<u>B</u>inding <u>P</u>rotein (PABPC1 in humans), which is a common target

of viral inhibition through diverse mechanisms[4]. *PAB1* and *SRO9* overexpression markedly reduced Gag

levels, explaining their rescuing phenotypes **(Fig 3B)**. Curiously, even though *SLF1* overexpression

rescued the *nuc1Δ ski3Δ* growth defect just as well as *SRO9*, it did not lead to any reduction in Gag

levels **(Fig 3A and 3B)**. These findings suggest that *SRO9* rescues cells by opposing L-A replication and

that *SLF1* protects cells from the pathogenic consequences of elevated viral replication **(Fig 2C)**.

To gain insight into the divergent mechanisms of Sro9 and Slf1, we considered further what

physiological consequences L-A pathogenesis has and how *SRO9*/*SLF1* might differentially impact

them. We noted a previous study in which deletions of *NUC1* or of *SKI*-complex genes led to weak

induction of a proteostatic stress GFP reporter gene controlled by Hsf1, a conserved transcription factor

that senses proteostatic stress and activates the gene expression response[31-34]. Using flow cytometry with

this reporter (HSE-GFP), we determined that a *nuc1Δ ski3Δ* double mutant exhibited strong activation of

HSE-GFP and that this activation was reverted in L-A^0 strains **(Fig 3C and S3)**. We hypothesized that

the massive accumulation of Gag observed in *nuc1Δ ski3Δ* mutants accounted for this proteostatic stress

response. Supporting this, HSE-GFP activation of a *nuc1Δ ski3Δ* double mutant was reverted by *PAB1*

or *SRO9* over-expression, but not by *SLF1* over-expression, mirroring these genes respective

consequences for Gag accumulation **(Fig 3B and 3C)**. The evolutionary divergence of *SRO9* and *SLF1*

has thus resulted in different viral-protective mechanisms, with *SRO9* opposing viral protein

accumulation and associated proteostatic stress, and *SLF1* possibly protecting cells from the toxic

consequences of viral induced proteostatic stress **(Fig 2C)**.

To identify new antiviral factors using a different approach, we searched a curated genetic interaction database for gene deletions that caused a synergistic growth defect when combined with *nuc1Δ* in at least two high-throughput screening studies. We found twenty-one such examples[35], two of which we confirmed as new viral attenuation factors. The first of these, *REX2*, is conserved from bacteria to humans[5]. Both Rex2 and its human homolog REXO2 are mitochondrial-localized 3'-5' RNA exonucleases that contain an EXOIII domain widely found in prokaryotic and eukaryotic proteins, including the interferon stimulated antiviral protein ISG20[6,7,36-38]. Mutant strains combining *rex2Δ* with *nuc1Δ* accumulated greatly increased levels of Gag and exhibited L-A-dependent lethality at high temperature **(Fig 4A and 4B)**. Remarkably, *nuc1Δ ski3Δ rex2Δ* and *nuc1Δ xrn1Δ rex2Δ* triple mutants were inviable under all growth conditions and these defects were reversed in L-A^0 strains **(Fig S4)**. These findings demonstrate the severe pathogenic potential of L-A and identify a new and highly conserved mitochondrial localized antiviral RNA nuclease **(Fig 2C)**.

The second gene identified in our survey was *MYG1*, the yeast homolog of human MelanocYte proliferation Gene 1, a 3'-5' RNA exonuclease associated with the human auto-immune disorder vitiligo[8,39]. Phylogenetic analysis has identified *MYG1* orthologs across all taxa[8]. A *myg1Δ nuc1Δ* double mutant strain displayed large increases of Gag accumulation compared to the single mutants and exhibited L-A-dependent lethality at high temperature **(Fig 4C, 4D, and S5)**. We were able to recover *nuc1Δ ski3Δ myg1Δ* triple mutants, though they were extremely slow growing at 30˚ C and accumulated even higher levels of Gag **(Fig 4C and 4D)**. These growth defects were also completely reversed in L-A^0 strains **(Fig S5)**. *MYG1* thus represents a new viral attenuation factor, acting in parallel to both *NUC1* and the *SKI* complex **(Fig 2C)**. Reflecting its underpinning as an auto-immune disease, Vitiligo is triggered by viral infection and is associated with mutations in the intracellular viral sensor MDA5[40]. An interesting possibility our findings thus suggest is that the elevated h*MYG1* expression associated with vitiligo may reflect an antiviral role for this protein[39]. Indeed, growth defects caused by *myg1Δ* are complemented by human Myg1, demonstrating a functional conservation to further investigate[8].

Although insightful studies of the L-A virus have been carried out, the L-A virus system has not yet been extensively exploited as a model for studying viral innate immunity, owing largely to its apparent benign nature. Here, we show that L-A is profoundly consequential for yeast fitness when its replication is uncontrolled, and that diverse innate immune pathways contribute to maintaining L-A replication at a harmless level. Importantly, our discovery that L-A is attenuated by homologs of known human antiviral factors (*SRO9/SLF1* and *PAB1*) supports the hypothesis that *REX2* and *MYG1* homologs may have viral innate immune functions in other organisms. Of note, Rex2 expands the arsenal of

mitochondrial localized antiviral nucleases beyond Nuc1, suggesting a key role of mitochondria in yeast viral attenuation. Among the many possible explanations for this, one scenario is that some aspect of the L-A replication cycle occurs in physical association with mitochondria similarly to as documented for the plus strand ssRNA flock house virus[41]. Whatever the case may be, our results highlight a potential general importance of mitochondria for viral innate immunity in eukaryotes, and position the yeast-L-A system as a powerful model for further studies of this topic[42].

Acknowledgements

We are grateful Dr. Charlie Boone and Dr. Brenda Andrews for generous sharing of strains and plasmid and to Angus McQuibban for critical comments on the manuscript. M.D.M. is funded by NSERC RGPIN-2019-07230. A.R.D. is funded by CIHR FDN-15427.

Author Contributions

S.C., J.G., A.D., S.B.C., and M.D.M. conceived, planned, and executed the experiments.

S.C., J.G., A.D., and M.D.M. analyzed the data. S.C., A.R.D., and M.D.M. wrote the manuscript.

Declaration of interests

We declare no competing financial interests.

STAR Methods

Key Resources Table.

REAGENT or RESOURCE	Source	Identifier
Antibodies		
HRP-conjugated horse anti-mouse	Cell Signaling Technology	Cat # 7075
HRP-conjugated goat anti-rabbit	Cell Signaling Technology	Cat # 7074
Anti-Pgk1	Abcam	Cat # ab113684
Anti-L-A Gag	Reed Wickner	
Anti-Sro9	Sandra Wolin	Sobel et al.[29]
Anti-Slf1	Sandra Wolin	Sobel et al.[29]
J2 anti-dsRNA	SCICONS	
Cyanine3 conjugated goat anti-mouse	Jackson ImmunoResearch Laboratories	
Experimental models: Organisms/strains*		
BY4741 *MAT*a L-A^0 *nuc1Δ::NatMX6 ski3Δ::HygMX6*	This study	jgy42
BY4741 *MAT*a L-A$^+$ p5476 *2μ LEU2*	This study	jgy156
BY4741 *MAT*a L-A$^+$ *nuc1Δ::NatMX6 ski3Δ::HygMX6* p5476 *2μ LEU2*	This study	jgy157

BY4741 *MAT*a L-A$^+$ *nuc1Δ::NatMX6* *ski3Δ::HygMX6* p5476 *PAB1 2μ LEU2*	This study	jgy158
BY4741 *MAT*a L-A$^+$ *nuc1Δ::NatMX6* *ski3Δ::HygMX6* p5476 *SRO9 2μ LEU2*	This study	jgy171
BY4741 *MAT*a L-A$^+$ *nuc1Δ::NatMX6* *ski3Δ::HygMX6* p5476 *SLF1 2μ LEU2*	This study	jgy173
BY4741 *MAT*α L-A$^+$ *nuc1Δ::NatMX6*	This study	mmy4076
BY4741 *MAT*α L-A$^+$	This study	mmy4699
BY4741 *MAT*a L-A$^+$	This study	mmy6042
BY4741 *MAT*a L-A$^+$ *ski3Δ::HygMX6*	This study	mmy6860
BY4741 *MAT*a L-A$^+$	This study	mmy7701
BY4741 *MAT*α L-A^0	This study	mmy8064
BY4741 *MAT*a L-A$^+$ *nuc1Δ::NatMX6* *ski3Δ::HygMX6*	This study	mmy8213
BY4741 *MAT*α L-A^0	This study	mmy8244
BY4741 *MAT*a L-A^0	This study	mmy8245
BY4741 *MAT*α L-A^0 *nuc1Δ::NatMX6*	This study	mmy8246
BY4741 *MAT*a L-A^0 *nuc1Δ::NatMX6*	This study	mmy8247
BY4741 *MAT*a L-A^0 *nuc1Δ::NatMX6*	This study	mmy8466
BY4741 *MAT*α L-A$^+$ *nuc1Δ::NatMX6* *ski3Δ::HygMX6* p5476 *2μ LEU2*	This study	mmy9292
BY4741 *MAT*α L-A$^+$ *nuc1Δ::NatMX6* *ski3Δ::HygMX6* p5476 *NUC1 2μ LEU2*	This study	mmy9293
BY4741 *MAT*α L-A$^+$ *nuc1Δ::NatMX6* p5476 *2μ LEU2*	This study	mmy9320
BY4741 *MAT*α L-A$^+$ p5476 *2μ LEU2*	This study	mmy9321
BY4741 *MAT*α L-A^0 *nuc1Δ::NatMX6* *ski3Δ::HygMX6* p5476 *2μ LEU2*	This study	mmy9322
BY4741 *MAT*α L-A$^+$ *ski3Δ::HygMX6* p5476 *2μ LEU2*	This study	mmy9323
BY4741 *MAT*a L-A$^+$ *nuc1Δ::NatMX6* *rex2Δ::KanMX6*	This study	mmy9342
BY4741 *MAT*a L-A$^+$ *rex2Δ::KanMX6*	This study	mmy9344
BY4741 *MAT*α L-A$^+$ *nuc1Δ::NatMX6* *ski3Δ::HygMX6* p5476 *XRN1 2μ LEU2*	This study	mmy9350
BY4741 *MAT*α L-A$^+$ *nuc1Δ::NatMX6* *xrn1Δ::KanMX6*	This study	mmy9351
BY4741 *MAT*α L-A$^+$ *xrn1Δ::KanMX6*	This study	mmy9353
BY4741 *MAT*α L-A^0 *xrn1Δ::NatMX6*	This study	mmy9609
BY4741 *MAT*α L-A^0 *xrn1Δ::NatMX6* *nuc1Δ::HygMX6*	This study	mmy9611
BY4741 *MAT*a L-A$^+$ *nuc1Δ::NatMX6* *rex2Δ::HygMX6 xrn1Δ::KanMX6* pRS316 *NUC1-FLAG URA3*	This study	mmy10001
BY4741 *MAT*α L-A^0 *xrn1Δ::NatMX6* *nuc1Δ::HygMX6 rex2Δ::KanMX6* pRS316 *NUC1-FLAG URA3*	This study	mmy10003

BY4741 *MAT*α L-A^0 *URA3::HSE-EmGFP nuc1Δ::NatMX6 ski3Δ::HygMX6*	This study	mmy10007
BY4741 *MAT*α L-A$^+$ *URA3::HSE-EmGFP nuc1Δ::NatMX6*	This study	mmy10073
BY4741 *MAT*α L-A$^+$ *URA3::HSE-EmGFP nuc1Δ::NatMX6 ski3Δ::HygMX6*	This study	mmy10074
BY4741 *MAT*a L-A$^+$ *URA3::HSE-EmGFP ski3Δ::HygMX6*	This study	mmy10076
BY4741 *MAT*α L-A$^+$ *URA3::HSE-EmGFP*	This study	mmy10077
BY4741 *MAT*a L-A$^+$ *nuc1Δ::NatMX6 ski3Δ::HygMX6* p5476 *2µ LEU2*	This study	mmy10461
BY4741 *MAT*a L-A$^+$ *nuc1Δ::NatMX6 ski3Δ::HygMX6* p5476 *SRO9 2µ LEU2*	This study	mmy10462
BY4741 *MAT*a L-A$^+$ *nuc1Δ::NatMX6 ski3Δ::HygMX6* p5476 *SLF1 2µ LEU2*	This study	mmy10463
BY4741 *MAT*a L-A$^+$ *nuc1Δ::NatMX6 ski3Δ::HygMX6* p5476 *PAB1 2µ LEU2*	This study	mmy10464
BY4741 *MAT*α L-A$^+$ *nuc1Δ::NatMX6 myg1Δ::KanMX6*	This study	mmy10845
BY4741 *MAT*a L-A$^+$ *ski3Δ::HygMX6 myg1Δ::KanMX6*	This study	mmy10846
BY4741 *MAT*α L-A$^+$ *myg1Δ::KanMX6*	This study	mmy10847
BY4741 *MAT*α L-A^0 *nuc1Δ::NatMX6 ski3Δ::HygMX6 myg1Δ::KanMX6*	This study	mmy10848
BY4741 *MAT*a L-A^0 *nuc1Δ::NatMX6 myg1Δ::KanMX6*	This study	mmy10849
BY4741 *MAT*α L-A$^+$ *nuc1Δ::NatMX6 ski3Δ::HygMX6 myg1Δ::KanMX6*	This study	mmy10852
BY4741 *MAT*a L-A^0 *rex2Δ::KanMX6*	This study	scy629
BY4741 *MAT*a L-A^0 *nuc1Δ::NatMX6 rex2Δ::KanMX6*	This study	scy630
BY4741 *MAT*a L-A$^+$ *nuc1Δ::NatMX6 ski3Δ::HygMX6 rex2Δ::KanMX6* pRS316 *NUC1 URA3*	This study	scy633
BY4741 *MAT*a L-A^0 *nuc1Δ::NatMX6 ski3Δ::HygMX6 rex2Δ::KanMX6* pRS316 *NUC1 URA3*	This study	scy635
BY4741 *MAT*a L-A$^+$ L-BC^0 *nuc1Δ::NatMX6 ski3Δ::HygMX6*	This study	scy838
BY4741 *MAT*a L-A$^+$ L-BC^0	This study	scy839
BY4741 *MAT*a L-A$^+$ L-BC^0 *ski3Δ::HygMX6*	This study	scy841
BY4741 *MAT*a L-A$^+$ L-BC^0 *nuc1Δ::NatMX6*	This study	scy842
BY4741 *MAT*a L-A^0 L-BC^0 *nuc1Δ::NatMX6 ski3Δ::HygMX6*	This study	scy894
BY4741 *MAT*a L-A$^+$ *URA3::HSE-EmGFP* p5476 *2µ LEU2*	This study	scy1100

BY4741 *MATα* L-A+ *URA3::HSE-EmGFP nuc1Δ::NatMX6 ski3Δ::HygMX6* p5476 *2μ LEU2*	This study	scy1104
BY4741 *MATα* L-A+ *URA3::HSE-EmGFP nuc1Δ::NatMX6 ski3Δ::HygMX6* p5476 *PAB1 2μ LEU2*	This study	scy1105
BY4741 *MATα* L-A+ *URA3::HSE-EmGFP nuc1Δ::NatMX6 ski3Δ::HygMX6* p5476 *SLF1 2μ LEU2*	This study	scy1106
BY4741 *MATα* L-A+ *URA3::HSE-EmGFP nuc1Δ::NatMX6 ski3Δ::HygMX6* p5476 *SRO9 2μ LEU2*	This study	scy1107
Software and algorithms		
ImageJ	N/A	https://imagej.nih.gov/ij/
Volocity	Volocity	https://www.volocity4d.com/
Flowing	Turku Bioscience	https://bioscience.fi/
Excel	Microsoft	https://www.microsoft.com/
R studio	Rstudio	https://www.rstudio.com/

154 *Strain background
155   BY4741 *his3 ura3 leu2*
156

Lead Contact and Materials Availability.

Further information and requests for yeast strains, resources, and reagents should be directed to and will

be fulfilled by the lead contact, Marc Meneghini (marc.meneghini@utoronto.ca).

Experimental Model and Subject Details.

Strains were grown at 30°C in either YPAD or synthetic complete media with appropriate amino acid

drop out for plasmid maintenance unless otherwise specified.

Method Details.

Strains, media, and plasmids.

Standard *S. cerevisiae* genetic and strain manipulation techniques were used for strain construction.

Refer to STAR Methods for strains and plasmids used in this paper. The yOB255 *URA3::HSE-EmGFP*

strain was obtained from Onn Brandman[43]. The p5476 *2μ LEU2*, p5476 *NUC1 2μ LEU2*, p5476 *XRN1*

2μ LEU2, p5476 *PAB1 2μ LEU2*, p5476 *SRO9 2μ LEU2*, p5476 *SLF1 2μ LEU2* MOBY plasmids were

obtained from Brenda Andrews[44]. The pVB3011 *L-BC GAG (P*$_{GAL1}$*)* plasmid was obtained from

Suzanne Sandmeyer[45].

Spot analysis.

Yeast strains were grown overnight at 30°C to saturation. Each strain was diluted to an $OD_{600} = 0.4$,

serial diluted four times by 10-fold, and spotted onto agar plates containing synthetic complete media or

drop out media supplemented with the indicated carbon source (2% glucose or 3% glycerol).

Protein extraction and Western blot.

5 OD of log-phase cells were harvested and permeabilized with 0.1N NaOH at room temperature for 5

minutes. The cells were then pelleted and resuspended in SDS/PAGE buffer before heating at 100°C for

10 minutes. The samples were centrifuged to isolate the soluble fraction for Western blotting. Protein

concentrations were determined with an RC/DC assay (BioRad 5000121). Equal amounts of protein

were electrophoresed on 10% SDS/PAGE and transferred to poly(vinylidene difluoride) membranes.

Blots were incubated in primary antibody overnight and probed with 1:3,000 horseradish peroxidase

(HRP)-conjugated horse anti-mouse (7075; Cell Signaling Technology) or goat anti-rabbit (7074; Cell

Signaling Technology) secondary antibody. The proteins were detected with Luminata Forte Western

HRP Substrate (EMD Millipore) or SuperSignal™ West Atto Ultimate Sensitivity Substrate (A38555,

Thermo Scientific). Blots were imaged with the Bio-Rad ChemiDoc XRS+ system, and images were

processed with the Image Lab software package (Bio-Rad). The primary antibodies and their dilutions

were 1:5,000 anti-Pgk1 (ab113687; Abcam), 1:2,000 anti-L-A Gag (obtained from Reed Wickner),

1:8000 anti-Hsp60 (received from Toshiya Endo)[46], 1:500 anti-Tom22[46], 1:2000 anti-Om45[46], 1:1000

anti-Cox2 (ab110271; Abcam), 1:1000 anti-Sro9, and 1:1000 anti-Slf1 (received from Sandra Wolin)[29].

Virus curing.

L-A^0 strains were generated similar to as we described previously[18]. Briefly, a progenitor L-A^0 strain

was isolated following treatment with 32 mM Anisomycin (BioShip ANS245) for 4 days in YPD media

and plating for single colonies. Isolates were tested for the presence of the L-A genome by RT-PCR.

This L-A^0 strain was then used for subsequent backcrossing with *mak3Δ* to generate other L-A^0 strain

background.

To evict L-BC, cells were transformed with the *L-BC GAG (P*$_{GAL1}$*)* plasmid and grown overnight in –

URA drop out media supplemented with 2% raffinose. The saturated culture was diluted and grown in

raffinose to log phase before treated with 2% galactose for 24 hours. The resulting culture was plated for

single colonies on an agar plate containing synthetic complete media supplemented with 5-FOA. Isolates

were tested for the presence of the L-BC genome by RT-PCR.

RNA extraction and reverse transcription.

10 OD of log phase cells was harvested and resuspended in phenol solution (P4682; Sigma-Aldrich),

SDS, and buffer AE (10 mM Tris·Cl and 0.5 mM EDTA, pH 9.0). Samples were incubated at 65°C for

30 minutes, and the phase-separated supernatant was washed with chloroform and precipitated in

isopropanol overnight. The precipitate was washed with 70% ethanol and dissolved in water. RNA

samples were then purified with the RNeasy Mini Kit (Qiagen). Four micrograms of RNA were reverse-

transcribed using random nonamers and Maxima H Minus Reverse Transcriptase (Thermo Fisher). The

cDNA product was isolated by alkaline hydrolysis and treated with RNase A.

Immunofluorescence.

Log phase cells were fixed with 4% formaldehyde and washed 2 times with 1x phosphate-buffered

saline (PBS). Cells were resuspended in sorbitol buffer (1.2 M sorbitol and 20 mM potassium phosphate

buffer, pH 7.2) and spheroplated with β-mercaptoethanol (M7522; Sigma-Aldrich) and zymolase at

30°C. Spheroplasted cells were added onto polylysine coated slides and incubated with primary antibody

1:1000 J2 anti-dsRNA (SCICONS) overnight. The cells were detected with 1:1000 cyanine3 conjugated

goat anti-mouse (Jackson ImmunoResearch Laboratories) and stained with DAPI. The images were

acquired on the Zeiss Axio Imager Z1 with Volocity and processed with ImageJ.

Flow cytometry.

Log phase cells were stained with propidium iodide, fixed with 4% formaldehyde, and washed 3 times

with 1x PBS. Fluorescence intensity was measured with Becton Dickinson LSR II flow cytometer and

analyzed with Flowing Software. The median GFP intensities of three biological replicates were plotted

using R studio ggplot2.

High copy suppression screen.

A plasmid library of random genomic inserts in the YEP24 plasmid backbone was transformed into a

nuc1Δ ski3Δ strain and spread onto -URA agar plate for single colonies[47]. Potential suppressors were

obtained by growing at 37°C supplemented with glycerol. To confirm that the suppression was plasmid

dependent, the potential suppressors were first grown on agar plate containing 0.1% 5-fluoroorotic acid

(5-FOA) for counterselection and then transferred to a fresh agar plate at 37°C supplemented with

glycerol. The plasmids of the candidates were isolated and sequenced to identify potential genes

responsible for the suppression of L-A pathogenesis when over-expressed. The candidate genes were

confirmed by spot analysis using p5476 MOBY plasmids.

Quantification and Statistical Analysis.

Data was tested for statistical significance using Excel and plotted with R studio ggplot2. The details of

the statistical tests are described in the figure legends.

Data and Code Availability.

This study did not generate new data/code.

References

1. Bernheim, A., and Sorek, R. (2020). The pan-immune system of bacteria: antiviral defence as a
community resource. Nat Rev Microbiol *18*, 113-119. 10.1038/s41579-019-0278-2.
2. Wickner, R.B., Fujimura, T., and Esteban, R. (2013). Viruses and prions of Saccharomyces
cerevisiae. Adv Virus Res *86*, 1-36. 10.1016/B978-0-12-394315-6.00001-5.
3. Schmidt, N., Lareau, C.A., Keshishian, H., Ganskih, S., Schneider, C., Hennig, T., Melanson, R.,
Werner, S., Wei, Y., Zimmer, M., et al. (2021). The SARS-CoV-2 RNA-protein interactome in
infected human cells. Nat Microbiol *6*, 339-353. 10.1038/s41564-020-00846-z.
4. Smith, R.W., and Gray, N.K. (2010). Poly(A)-binding protein (PABP): a common viral target.
Biochem J *426*, 1-12. 10.1042/BJ20091571.
5. Koonin, E.V. (1997). A conserved ancient domain joins the growing superfamily of 3'-5'
exonucleases. Curr Biol *7*, R604-606. 10.1016/s0960-9822(06)00311-3.
6. Zuo, Y., and Deutscher, M.P. (2001). Exoribonuclease superfamilies: structural analysis and
phylogenetic distribution. Nucleic Acids Res *29*, 1017-1026. 10.1093/nar/29.5.1017.
7. Szewczyk, M., Malik, D., Borowski, L.S., Czarnomska, S.D., Kotrys, A.V., Klosowska-Kosicka,
K., Nowotny, M., and Szczesny, R.J. (2020). Human REXO2 controls short mitochondrial RNAs
generated by mtRNA processing and decay machinery to prevent accumulation of double-
stranded RNA. Nucleic Acids Res *48*, 5572-5590. 10.1093/nar/gkaa302.
8. Grover, R., Burse, S.A., Shankrit, S., Aggarwal, A., Kirty, K., Narta, K., Srivastav, R., Ray,
A.K., Malik, G., Vats, A., et al. (2019). Myg1 exonuclease couples the nuclear and mitochondrial
translational programs through RNA processing. Nucleic Acids Res *47*, 5852-5866.
10.1093/nar/gkz371.
9. Brown, J.T., Bai, X., and Johnson, A.W. (2000). The yeast antiviral proteins Ski2p, Ski3p, and
Ski8p exist as a complex in vivo. RNA *6*, 449-457.
10. Benard, L., Carroll, K., Valle, R.C., Masison, D.C., and Wickner, R.B. (1999). The ski7 antiviral
protein is an EF1-alpha homolog that blocks expression of non-Poly(A) mRNA in
Saccharomyces cerevisiae. J Virol *73*, 2893-2900. 10.1128/JVI.73.4.2893-2900.1999.

11. Anderson, J.S., and Parker, R.P. (1998). The 3' to 5' degradation of yeast mRNAs is a general
mechanism for mRNA turnover that requires the SKI2 DEVH box protein and 3' to 5'
exonucleases of the exosome complex. EMBO J *17*, 1497-1506. 10.1093/emboj/17.5.1497.

12. Ball, S.G., Tirtiaux, C., and Wickner, R.B. (1984). Genetic Control of L-a and L-(Bc) Dsrna
Copy Number in Killer Systems of SACCHAROMYCES CEREVISIAE. Genetics *107*, 199-
217.

13. Rowley, P.A., Ho, B., Bushong, S., Johnson, A., and Sawyer, S.L. (2016). XRN1 Is a Species-
Specific Virus Restriction Factor in Yeasts. PLoS Pathog *12*, e1005890.
10.1371/journal.ppat.1005890.

14. Burgess, H.M., and Mohr, I. (2015). Cellular 5'-3' mRNA exonuclease Xrn1 controls double-
stranded RNA accumulation and anti-viral responses. Cell Host Microbe *17*, 332-344.
10.1016/j.chom.2015.02.003.

15. Ng, C.S., Kasumba, D.M., Fujita, T., and Luo, H. (2020). Spatio-temporal characterization of the
antiviral activity of the XRN1-DCP1/2 aggregation against cytoplasmic RNA viruses to prevent
cell death. Cell Death Differ *27*, 2363-2382. 10.1038/s41418-020-0509-0.

16. Eckard, S.C., Rice, G.I., Fabre, A., Badens, C., Gray, E.E., Hartley, J.L., Crow, Y.J., and Stetson,
D.B. (2014). The SKIV2L RNA exosome limits activation of the RIG-I-like receptors. Nat
Immunol *15*, 839-845. 10.1038/ni.2948.

17. Miyashita, M., Oshiumi, H., Matsumoto, M., and Seya, T. (2011). DDX60, a DEXD/H box
helicase, is a novel antiviral factor promoting RIG-I-like receptor-mediated signaling. Mol Cell
Biol *31*, 3802-3819. 10.1128/MCB.01368-10.

18. Gao, J., Chau, S., Chowdhury, F., Zhou, T., Hossain, S., McQuibban, G.A., and Meneghini,
M.D. (2019). Meiotic viral attenuation through an ancestral apoptotic pathway. Proc Natl Acad
Sci U S A *116*, 16454-16462. 10.1073/pnas.1900751116.

19. Li, L.Y., Luo, X., and Wang, X. (2001). Endonuclease G is an apoptotic DNase when released
from mitochondria. Nature *412*, 95-99. 10.1038/35083620.

20. Thomson, B.J. (2001). Viruses and apoptosis. Int J Exp Pathol *82*, 65-76. 10.1111/j.1365-
2613.2001.iep0082-0065-x.

21. Eastwood, M.D., Cheung, S.W., Lee, K.Y., Moffat, J., and Meneghini, M.D. (2012).
Developmentally programmed nuclear destruction during yeast gametogenesis. Dev Cell *23*, 35-
44. 10.1016/j.devcel.2012.05.005.

22. Eastwood, M.D., and Meneghini, M.D. (2015). Developmental Coordination of Gamete
Differentiation with Programmed Cell Death in Sporulating Yeast. Eukaryot Cell *14*, 858-867.
10.1128/EC.00068-15.

23. Koonin, E.V., and Krupovic, M. (2019). Origin of programmed cell death from antiviral
defense? Proc Natl Acad Sci U S A *116*, 16167-16169. 10.1073/pnas.1910303116.

24. Weber, F., Wagner, V., Rasmussen, S.B., Hartmann, R., and Paludan, S.R. (2006). Double-
stranded RNA is produced by positive-strand RNA viruses and DNA viruses but not in
detectable amounts by negative-strand RNA viruses. J Virol *80*, 5059-5064.
10.1128/JVI.80.10.5059-5064.2006.

25. Welsch, S., Miller, S., Romero-Brey, I., Merz, A., Bleck, C.K., Walther, P., Fuller, S.D., Antony,
C., Krijnse-Locker, J., and Bartenschlager, R. (2009). Composition and three-dimensional
architecture of the dengue virus replication and assembly sites. Cell Host Microbe *5*, 365-375.
10.1016/j.chom.2009.03.007.

26. Fernandez de Castro, I., Tenorio, R., and Risco, C. (2016). Virus assembly factories in a lipid
world. Curr Opin Virol *18*, 20-26. 10.1016/j.coviro.2016.02.009.

27. Liu, Y.X., and Dieckmann, C.L. (1989). Overproduction of yeast viruslike particles by strains
deficient in a mitochondrial nuclease. Mol Cell Biol *9*, 3323-3331.

28.   Johnson, A.W., and Kolodner, R.D. (1995). Synthetic lethality of sep1 (xrn1) ski2 and sep1
326   (xrn1) ski3 mutants of Saccharomyces cerevisiae is independent of killer virus and suggests a
327   general role for these genes in translation control. Mol Cell Biol *15*, 2719-2727.
328   29.   Sobel, S.G., and Wolin, S.L. (1999). Two yeast La motif-containing proteins are RNA-binding
329   proteins that associate with polyribosomes. Mol Biol Cell *10*, 3849-3862.
330   10.1091/mbc.10.11.3849.
331   30.   Proweller, A., and Butler, J.S. (1996). Ribosomal association of poly(A)-binding protein in
332   poly(A)-deficient Saccharomyces cerevisiae. J Biol Chem *271*, 10859-10865.
333   10.1074/jbc.271.18.10859.
334   31.   Brandman, O., Stewart-Ornstein, J., Wong, D., Larson, A., Williams, C.C., Li, G.W., Zhou, S.,
335   King, D., Shen, P.S., Weibezahn, J., et al. (2012). A ribosome-bound quality control complex
336   triggers degradation of nascent peptides and signals translation stress. Cell *151*, 1042-1054.
337   10.1016/j.cell.2012.10.044.
338   32.   Anckar, J., and Sistonen, L. (2011). Regulation of HSF1 function in the heat stress response:
339   implications in aging and disease. Annu Rev Biochem *80*, 1089-1115. 10.1146/annurev-
340   biochem-060809-095203.
341   33.   Sorger, P.K., and Pelham, H.R. (1987). Purification and characterization of a heat-shock element
342   binding protein from yeast. EMBO J *6*, 3035-3041.
343   34.   Li, J., Labbadia, J., and Morimoto, R.I. (2017). Rethinking HSF1 in Stress, Development, and
344   Organismal Health. Trends Cell Biol *27*, 895-905. 10.1016/j.tcb.2017.08.002.
345   35.   Stark, C., Breitkreutz, B.J., Reguly, T., Boucher, L., Breitkreutz, A., and Tyers, M. (2006).
346   BioGRID: a general repository for interaction datasets. Nucleic Acids Res *34*, D535-539.
347   10.1093/nar/gkj109.
348   36.   Espert, L., Degols, G., Gongora, C., Blondel, D., Williams, B.R., Silverman, R.H., and Mechti,
349   N. (2003). ISG20, a new interferon-induced RNase specific for single-stranded RNA, defines an
350   alternative antiviral pathway against RNA genomic viruses. J Biol Chem *278*, 16151-16158.
351   10.1074/jbc.M209628200.
352   37.   Hanekamp, T., and Thorsness, P.E. (1999). YNT20, a bypass suppressor of yme1 yme2, encodes
353   a putative 3'-5' exonuclease localized in mitochondria of Saccharomyces cerevisiae. Curr Genet
354   *34*, 438-448. 10.1007/s002940050418.
355   38.   van Hoof, A., Lennertz, P., and Parker, R. (2000). Three conserved members of the RNase D
356   family have unique and overlapping functions in the processing of 5S, 5.8S, U4, U5, RNase
357   MRP and RNase P RNAs in yeast. EMBO J *19*, 1357-1365. 10.1093/emboj/19.6.1357.
358   39.   Dwivedi, M., Laddha, N.C., and Begum, R. (2013). Correlation of increased MYG1 expression
359   and its promoter polymorphism with disease progression and higher susceptibility in vitiligo
360   patients. J Dermatol Sci *71*, 195-202. 10.1016/j.jdermsci.2013.04.026.
361   40.   Zhuang, T., Yi, X., Chen, J., Kang, P., Chen, X., Chen, J., Cui, T., Chang, Y., Ye, Z., Ni, Q., et
362   al. (2020). Intracellular virus sensor MDA5 exacerbates vitiligo by inducing the secretion of
363   chemokines in keratinocytes under virus invasion. Cell Death Dis *11*, 453. 10.1038/s41419-020-
364   2665-z.
365   41.   Kopek, B.G., Perkins, G., Miller, D.J., Ellisman, M.H., and Ahlquist, P. (2007). Three-
366   dimensional analysis of a viral RNA replication complex reveals a virus-induced mini-organelle.
367   PLoS Biol *5*, e220. 10.1371/journal.pbio.0050220.
368   42.   West, A.P., Shadel, G.S., and Ghosh, S. (2011). Mitochondria in innate immune responses. Nat
369   Rev Immunol *11*, 389-402. 10.1038/nri2975.
370   43.   Sitron, C.S., Park, J.H., Giafaglione, J.M., and Brandman, O. (2020). Aggregation of CAT tails
371   blocks their degradation and causes proteotoxicity in S. cerevisiae. PLoS One *15*, e0227841.
372   10.1371/journal.pone.0227841.

373   44.   Magtanong, L., Ho, C.H., Barker, S.L., Jiao, W., Baryshnikova, A., Bahr, S., Smith, A.M.,
374   Heisler, L.E., Choy, J.S., Kuzmin, E., et al. (2011). Dosage suppression genetic interaction
375   networks enhance functional wiring diagrams of the cell. Nat Biotechnol *29*, 505-511.
376   10.1038/nbt.1855.
377   45.   Bilanchone, V., Clemens, K., Kaake, R., Dawson, A.R., Matheos, D., Nagashima, K., Sitlani, P.,
378   Patterson, K., Chang, I., Huang, L., and Sandmeyer, S. (2015). Ty3 Retrotransposon Hijacks
379   Mating Yeast RNA Processing Bodies to Infect New Genomes. PLoS Genet *11*, e1005528.
380   10.1371/journal.pgen.1005528.
381   46.   Sakaue, H., Shiota, T., Ishizaka, N., Kawano, S., Tamura, Y., Tan, K.S., Imai, K., Motono, C.,
382   Hirokawa, T., Taki, K., et al. (2019). Porin Associates with Tom22 to Regulate the
383   Mitochondrial Protein Gate Assembly. Mol Cell *73*, 1044-1055 e1048.
384   10.1016/j.molcel.2019.01.003.
385   47.   Kamada, Y., Fujioka, Y., Suzuki, N.N., Inagaki, F., Wullschleger, S., Loewith, R., Hall, M.N.,
386   and Ohsumi, Y. (2005). Tor2 directly phosphorylates the AGC kinase Ypk2 to regulate actin
387   polarization. Mol Cell Biol *25*, 7239-7248. 10.1128/MCB.25.16.7239-7248.2005.
388

Figure 1. Loss of *NUC1* and *SKI3* displays a synergistic increase in L-A viral load. **(A)** An ethidium bromide-stained gel of total RNA prepared from the indicated strains is shown, with the 4.6 Kb L-A dsRNA band indicated with an arrow. **(B)** Immunofluorescence was used to visualize L-A dsRNA (red) in cells of the indicated genotypes. These strains were cured of the weakly abundant L-BC dsRNA virus to eliminate background staining (see Methods). DAPI staining of DNA is shown in blue. Scale bar, 1 μm.

Figure 2. L-A attenuation protects yeast from lethal pathogenesis. **(A)** Western blotting of L-A Gag and Pgk1 protein levels in the indicated strains is shown. Molecular weight markers are indicated on the right. **(B)** Spot test growth assays of the strains from 2A is shown. Strains were spotted on -Leu media containing either glucose or glycerol and grown at the indicated temperatures. **(C)** The mitochondrial proteins Nuc1 and Rex2 collaborate with the cytosolic proteins Xrn1, SkiC, and Myg1 to regulate L-A protein level and ensure cell fitness. The translation control factors Pab1 and Sro9 alleviate L-A pathogenesis by suppressing L-A level while Slf1, the paralog of Sro9, suppress L-A pathogenesis without affecting L-A level. Newly discovered L-A attenuation factors are indicated in red.

Figure 3. Overexpression of translation control factors alleviate L-A pathogenesis. **(A)** Spot test growth assays of the high copy suppressors *SRO9*, *SLF1*, and *PAB1* are shown. **(B)** Western blotting detection of L-A Gag, Pgk1, Sro9, and Slf1 protein levels in the strains from 3A is shown. Molecular weight markers are indicated on the right. **(C)** Flow cytometry was used to measure HSE-GFP expression in the indicated strains. Median GFP intensity is shown. n = 3. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. The p value was calculated using the unpaired student's t-test.

Figure 4. New antiviral factors are identified by exploiting L-A pathogenesis. **(A)** Spot analysis of strains defective in *NUC1* and *REX2* is shown. Strains are spotted on SC media containing either glucose or glycerol and grown at the indicated temperature. **(B)** Western blotting of L-A Gag and Pgk1 protein levels of strains in Figure 4A. Molecular weight markers are indicated on the right. **(C)** Spot analysis of strains defective in *MYG1*, *NUC1*, and *SKI3* is shown. Strains are spotted on SC media containing either glucose or glycerol and grown at the indicated temperature. **(D)** Western blotting detection of L-A Gag and Pgk1 protein levels in the indicated strains is shown. Molecular weight markers are indicated on the right.

Figure S1. Additional images of immunofluorescence of dsRNA (red) of strains from Figure 1B. Images of fields of cells of indicated genotype is shown. Cells are stained with DAPI to visualize nuclei. Scale bar, 30 μm.

A

	Glucose 30°C	Glucose 37°C	Glycerol 37°C

L-A⁺
- WT
- *nuc1Δ*
- *xrn1Δ*
- *nuc1Δ xrn1Δ*

L-A⁰
- WT
- *nuc1Δ*
- *xrn1Δ*
- *nuc1Δ xrn1Δ*

B

Figure S2. *XRN1*, *NUC1*, and *SKI3* work in parallel pathways to regulate L-A Gag level. **(A)** Spot analysis of strains defective in *NUC1* and *XRN1* is shown. Strains are spotted on SC media containing either glucose or glycerol and grown at the indicated temperature. **(B)** Western blotting of L-A Gag and Pgk1 protein levels of strains in Figure S2A. Molecular weight markers are indicated on the right.

Figure S3. L-A pathogenesis correlates with proteostatic stress. Flow cytometry was used to measure HSE-GFP expression in the indicated strains. Median GFP intensity is shown. $n = 3$. $*$ $p < 0.05$, $**$ $p < 0.01$. The p value was calculated using unpaired student's t-test.

Figure S4. The mitochondrial exonuclease *REX2* is an antiviral factor. Spot analysis of strains defective in 3 parallel antiviral pathways containing plasmid expressing *NUC1* is shown. Strains are spotted on -URA media or synthetic complete (SC) media supplemented with 0.1% 5-fluoroorotic acid (5-FOA).

Figure S5. *MYG1* collaborates with *NUC1* and *SKI3* to ensure cell fitness in a L-A dependent manner. Spot analysis of strains defective in *MYG1*, *NUC1*, and *SKI3* with and without L-A is shown. Strains are spotted on SC media containing either glucose or glycerol and grown at the indicated temperature.