**Diverse innate immune factors protect yeast from lethal viral pathogenesis**

Sabrina Chau[1], Jie Gao[1], Annette Diao[1], Michelle Cao[1], Alan R. Davidson[1,2], and Marc D. Meneghini[1*]

[1]Department of Molecular Genetics, University of Toronto, 661 University Ave, Toronto, ON, M5G

1M1, Canada.

[2]Department of Biochemistry, University of Toronto, 661 University Ave, Toronto, ON, M5G 1M1,

Canada.

[*]Corresponding Author

**Although the budding yeast *Saccharomyces cerevisiae* is chronically infected with a double-**

**stranded RNA virus called L-A, the lack of any known L-A fitness consequence has hindered use**

**of this model organism for the study of host-virus interactions. Here we show that L-A causes**

**lethal pathogenesis in cells lacking parallel-acting viral attenuation pathways. Leveraging this**

**discovery, we identify known and new antiviral factors exploiting unbiased genetic screens and**

**determine that rampant L-A proliferation causes proteostatic stress.**

In recent years, newly characterized antiviral systems have been found to be remarkably conserved from

bacteria to mammals, revealing the potential to gain unique insight into these systems through the study

of microbial organisms[1]. Despite the enthusiasm generated by these findings, the key microbial model

organism *Saccharomyces cerevisiae* (budding yeast) has been minimally exploited for studies of viral

defense, primarily because it is not infected with exogenously transmitted viruses. However, most yeast

strains are infected with an endogenous double stranded RNA (dsRNA) virus called L-A, and previous

studies have identified conserved antiviral systems that attenuate L-A replication[2]. Although these

systems do not lead to eradication of L-A, we show here that they prevent massive over-proliferation of

L-A that leads to proteostatic stress and lethality. Exploiting this new insight, we asked if the powerful

genetic screening methods available in yeast could be used to identify additional conserved antiviral

systems. Validating this approach, we discovered antiviral functions for the yeast homologs of polyA-

binding protein (PABPC1) and the La-domain containing protein Larp1, which are both involved in viral

innate immunity in humans[3,4]. We also identify new antiviral functions for the RNA exonucleases *REX2*

and *MYG1*, both of which have distinct but poorly characterized human and bacterial homologs. These

findings highlight the potential of yeast as a powerful model system for studies of the ubiquitous host-

virus arms race.

L-A belongs to the broadly dispersed *Totiviridae* family of double stranded RNA (dsRNA) viruses, which has no extracellular phase and is transmitted vertically through cell division or fusion[2]. L-A transcripts encode the capsid (Gag) proteins that comprise most of its viral particle as well as a Gag-pol fusion protein produced at much lower levels that provides RNA dependent RNA polymerase activity responsible for replication and transcription. Encapsidation of viral transcripts within nascent particles completes the L-A lifecycle[2]. L-A replication is maintained at a low level through the activities of the *SKI2*, *3*, and *8* genes, which encode subunits of a conserved ribosome-associated complex that opposes the translation of transcripts such as those encoded by L-A that lack polyA tails[5-7]. A separate pathway of L-A attenuation is provided by Xrn1, a 5'-3' exoribonuclease that degrades uncapped mRNAs[8,9]. Both of these systems have been associated with anti-viral defense in mammals[10-13].

We showed previously that L-A replication was held in check during sporulation by the mitochondrial DNA/RNA endonuclease Nuc1, a homolog of endonuclease G (EndoG) found in all eukaryotes and many prokaryotes[14]. EndoG is involved in mammalian programmed cell death that frequently occurs in virus infected cells, and programmed cell death is intrinsic to yeast sporulation[15-18]. These and other observations have led to speculation that Nuc1/EndoG may reflect a prokaryotic antiviral origin of programmed cell death[19]. To query *NUC1* anti-viral function outside of sporulation and its associated programmed cell death, we investigated L-A attenuation in mitotic haploid cells in the reference BY4742 strain background. Visualization of L-A dsRNA using ethidium bromide staining of electrophoresed RNA revealed that while *nuc1Δ* and *ski3Δ* single mutants exhibited no effect, combining *nuc1Δ* and *ski3Δ* led to a large increase in L-A dsRNA **(Fig 1A)**. We corroborated these findings using immunofluorescence microscopy with a dsRNA antibody widely used to detect replicating RNA viruses[20,21]. These images revealed L-A dsRNA to accumulate in foci, reminiscent of "viral factory" sites of viral replication observed in human cells **(Fig 1B and S1)**[22]. Consistent with previous findings in other strain backgrounds, western blotting showed that Gag protein levels were elevated in *nuc1Δ* and *ski3Δ* mutants[8,14,23], and we determined that *nuc1Δ ski3Δ* double mutants accumulated massively elevated Gag levels, confirming that *NUC1* and *SKI3* participate in separate antiviral pathways **(Fig 1C)**.

We used spot test growth assays to determine if high L-A viral load had any fitness consequence and found that *nuc1Δ ski3Δ* double mutants grew normally at 30° C but exhibited conditional lethality at 37° C **(Fig 1D)**. Modest growth defects of *nuc1Δ* and *ski3Δ* single mutants were observed at high temperature when cells were grown with glycerol rather than glucose as the carbon source, a condition under which yeast relies on mitochondrial respiration **(Fig 1D)**. These growth phenotypes were completely reversed in isogenic strains cured of L-A (L-A[0]), implying that unrestricted L-A replication

caused them (**Fig 1D**).

As expected, a *NUC1*-expressing plasmid restored viability to a *nuc1Δ ski3Δ* strain grown at high

temperature and elicited a corresponding decrease in Gag levels (**Fig 1C and 1D**). We observed similar

suppression of *nuc1Δ ski3Δ* using plasmid driven *XRN1* over-expression, suggesting that Xrn1 repressed

L-A via a parallel mechanism (**Fig 1C and 1D**). Consistent with this model, *nuc1Δ xrn1Δ* mutants

accumulated greatly elevated levels of Gag and exhibited L-A-dependent conditional lethality at high

temperature (**Fig S2**), and *xrn1Δ ski3Δ* mutants are inviable[24]. We conclude that Nuc1, Ski3 and Xrn1

convergently oppose L-A, and that L-A causes lethal pathogenesis at high temperature when

unattenuated (**Fig 1E**).

Since *XRN1* over-expression suppressed the growth defects of a *nuc1Δ ski3Δ* strain, we

hypothesized that over-expression of other antiviral factors would produce a similar effect, and that this

phenomenon could be used as a screen to identify new antiviral systems. To test this idea, we employed

a high-copy plasmid suppression screen to detect genes whose overexpression alleviated the conditional

lethality of *nuc1Δ ski3Δ*. Using this strategy, we identified *SRO9, SLF1*, and *PAB1* as high-copy

suppressors, all of which encode ribosome-associated RNA binding proteins (**Fig 2A**)[25,26]. Sro9 and Slf1

are paralogous lupus-autoantigen (La) domain containing proteins broadly found in eukaryotes.

Interestingly, the human Sro9/Slf1 ortholog Larp1 was recently found to bind to SARS-Cov-2 plus-

strand ssRNA and to oppose its replication in human cells through an unknown mechanism[3]. *PAB1*

encodes the highly conserved PolyA-Binding Protein (PABPC1 in humans), which is a common target

of viral inhibition through diverse mechanisms[4]. *PAB1* and *SRO9* overexpression markedly reduced Gag

levels, explaining their rescuing phenotypes (**Fig 2B**). Curiously, even though *SLF1* overexpression

rescued the *nuc1Δ ski3Δ* growth defect just as well as *SRO9*, it did not lead to any reduction in Gag

levels (**Fig 2A and 2B**). These findings suggest that *SRO9* rescues cells by opposing L-A replication and

that *SLF1* protects cells from the pathogenic consequences of elevated viral replication (**Fig 1E**).

To gain insight into the divergent mechanisms of Sro9 and Slf1, we considered further what

physiological consequences L-A pathogenesis has and how *SRO9/SLF1* might differentially impact

them. We noted a previous study in which deletions of *NUC1* or of *SKI*-complex genes led to weak

induction of a proteostatic stress GFP reporter gene controlled by Hsf1, a conserved transcription factor

that senses proteostatic stress and activates the gene expression response[27-30]. Using flow cytometry with

this reporter (HSE-GFP), we determined that a *nuc1Δ ski3Δ* double mutant exhibited strong activation of

HSE-GFP and that this activation was reverted in L-A$^0$ strains (**Fig 2C and S3**). We hypothesized that

the massive accumulation of Gag observed in *nuc1Δ ski3Δ* mutants accounted for this proteostatic stress

response. Supporting this, HSE-GFP activation of a *nuc1Δ ski3Δ* double mutant was reverted by *PAB1*

or *SRO9* over-expression, but not by *SLF1* over-expression, mirroring these genes respective consequences for Gag accumulation **(Fig 2B and 2C)**. The evolutionary divergence of *SRO9* and *SLF1* has thus resulted in different viral-protective mechanisms, with *SRO9* opposing viral protein accumulation and associated proteostatic stress, and *SLF1* possibly protecting cells from the toxic consequences of viral induced proteostatic stress **(Fig 1E)**.

To identify new antiviral factors using a different approach, we searched a curated genetic interaction database for gene deletions that caused a synergistic growth defect when combined with *nuc1Δ* in at least two high-throughput screening studies. We found fifteen such examples[31], two of which we confirmed as new viral attenuation factors. The first of these, *REX2*, is conserved from bacteria to humans[32]. Both Rex2 and its human homolog REXO2 are mitochondrial-localized 3'-5' RNA exonucleases that contain an EXOIII domain widely found in prokaryotic and eukaryotic proteins, including the interferon stimulated antiviral protein ISG20[33-37]. Mutant strains combining *rex2Δ* with *nuc1Δ* accumulated greatly increased levels of Gag and exhibited L-A-dependent lethality at high temperature **(Fig 2D and 2E)**. Remarkably, *nuc1Δ ski3Δ rex2Δ* and *nuc1Δ xrn1Δ rex2Δ* triple mutants were inviable under all growth conditions and these defects were reversed in L-A$^0$ strains **(Fig S4)**. These findings demonstrate the severe pathogenic potential of L-A and identify a new and highly conserved mitochondrial localized antiviral RNA nuclease **(Fig 1E)**.

The second gene identified in our survey was *MYG1*, the yeast homolog of human MelanocYte proliferation Gene 1, a 3'-5' RNA exonuclease associated with the human auto-immune disorder vitiligo[38,39]. Phylogenetic analysis has identified *MYG1* orthologs across all taxa[38]. A *myg1Δ nuc1Δ* double mutant strain displayed large increases of Gag accumulation compared to the single mutants and exhibited L-A-dependent lethality at high temperature **(Fig S5A, S5B, and S5C)**. We were able to recover *nuc1Δ ski3Δ myg1Δ* triple mutants, though they were extremely slow growing at 30˚C and accumulated even higher levels of Gag. These growth defects were also completely reversed in L-A$^0$ strains **(Fig S5C)**. *MYG1* thus represents a new viral attenuation factor, acting in parallel to both *NUC1* and the *SKI* complex **(Fig 1E)**. Reflecting its underpinning as an auto-immune disease, Vitiligo is associated with viral infection as well as with mutations in the intracellular viral sensor MDA5[40]. An interesting possibility our findings suggest is that the elevated h*MYG1* expression associated with vitiligo may reflect an antiviral role for this protein[39]. Indeed, growth defects caused by *myg1Δ* are complemented by human Myg1, demonstrating a functional conservation to further investigate[38].

Although insightful studies of the L-A virus have been carried out, the L-A virus system has not yet been extensively exploited as a model for studying innate antiviral immunity, owing largely to its apparent benign nature. Here, we show that L-A is profoundly consequential for yeast fitness when its

replication is uncontrolled, and that diverse innate immune pathways contribute to maintaining L-A

replication at a harmless level. Importantly, our discovery that L-A is attenuated by homologs of known

human antiviral factors (*SRO9/SLF1* and *PAB1*) supports the hypothesis that *REX2* and *MYG1* homologs

may have viral innate immune functions in other organisms. Of note, Rex2 expands the arsenal of

mitochondrial localized antiviral nucleases beyond Nuc1, suggesting a key role of mitochondria in yeast

viral attenuation. Among the many possible explanations for this, one scenario is that some aspect of the

L-A replication cycle occurs in physical association with mitochondria similarly to as documented for

the plus strand ssRNA flock house virus[41]. Whatever the case may be, our results highlight a potential

general importance of mitochondria for viral innate immunity in eukaryotes, and position the yeast-L-A

system as a powerful model for further studies of this topic[42].

**Methods and Materials**

**Strains, media, and plasmids.**

Standard *S. cerevisiae* genetic and strain manipulation techniques were used for strain construction.

Refer to Table S1 for strains and plasmids used in this paper. The yOB255 *URA3::HSE-EmGFP* strain

was obtained from Onn Brandman[43]. The p5476 *2μ LEU2*, p5476 *NUC1 2μ LEU2*, p5476 *XRN1 2μ*

*LEU2*, p5476 *PAB1 2μ LEU2*, p5476 *SRO9 2μ LEU2*, p5476 *SLF1 2μ LEU2* MOBY plasmids were

obtained from Brenda Andrews[44]. The pVB3011 *L-BC GAG (P_{GAL1})* plasmid was obtained from

Suzanne Sandmeyer[45].

**Spot analysis.**

Yeast strains were grown overnight at 30°C to saturation. Each strain was diluted to an $OD_{600}$ = 0.4,

serial diluted four times by 10-fold, and spotted onto agar plates containing synthetic complete media or

drop out media supplemented with the indicated carbon source (2% glucose or 3% glycerol).

**Protein extraction and Western blot.**

5 OD of log-phase cells were harvested and permeabilized with 0.1N NaOH at room temperature for 5

minutes. The cells were then pelleted and resuspended in SDS/PAGE buffer before heating at 100°C for

10 minutes. The samples were centrifuged to isolate the soluble fraction for Western blotting. Protein

concentrations were determined with an RC/DC assay (BioRad 5000121). Equal amounts of protein

were electrophoresed on 10% SDS/PAGE and transferred to poly(vinylidene difluoride) membranes.

Blots were incubated in primary antibody overnight and probed with 1:3,000 horseradish peroxidase

(HRP)-conjugated horse anti-mouse (7075; Cell Signaling Technology) or goat anti-rabbit (7074; Cell

Signaling Technology) secondary antibody. The proteins were detected with Luminata Forte Western

HRP Substrate (EMD Millipore) or SuperSignal™ West Atto Ultimate Sensitivity Substrate (A38555,

Thermo Scientific). Blots were imaged with the Bio-Rad ChemiDoc XRS+ system, and images were

processed with the Image Lab software package (Bio-Rad). The primary antibodies and their dilutions

were 1:5,000 anti-Pgk1 (ab113687; Abcam), 1:2,000 anti-L-A Gag (obtained from Reed Wickner),

1:8000 anti-Hsp60 (received from Toshiya Endo)[46], 1:500 anti-Tom22[46], 1:2000 anti-Om45[46]1:1000

anti-Cox2 (ab110271; Abcam), 1:1000 anti-Sro9, and 1:1000 anti-Slf1 (received from Sandra Wolin)[25].

**Virus curing.**

L-A$^0$ strains were generated similar to as we described previously[14]. Briefly, a progenitor L-A$^0$ strain

was isolated following treatment with 32 mM Anisomycin (BioShip ANS245) for 4 days in YPD media

and plating for single colonies. Isolates were tested for the presence of the L-A genome by RT-PCR.

This L-A$^0$ strain was then used for subsequent backcrossing with *mak3Δ* to generate other L-A$^0$ strain

background.

To evict L-BC, cells were transformed with the *L-BC GAG (P$_{GAL1}$)* plasmid and grown overnight in –

URA drop out media supplemented with 2% raffinose. The saturated culture was diluted and grown in

raffinose to log phase before treated with 2% galactose for 24 hours. The resulting culture was plated for

single colonies on an agar plate containing synthetic complete media supplemented with 5-FOA. Isolates

were tested for the presence of the L-BC genome by RT-PCR.

**RNA extraction and reverse transcription.**

10 OD of log phase cells was harvested and resuspended in phenol solution (P4682; Sigma-Aldrich),

SDS, and buffer AE (10 mM Tris·Cl and 0.5 mM EDTA, pH 9.0). Samples were incubated at 65°C for

30 minutes, and the phase-separated supernatant was washed with chloroform and precipitated in

isopropanol overnight. The precipitate was washed with 70% ethanol and dissolved in water. RNA

samples were then purified with the RNeasy Mini Kit (Qiagen). Four micrograms of RNA were reverse-

transcribed using random nonamers and Maxima H Minus Reverse Transcriptase (Thermo Fisher). The

cDNA product was isolated by alkaline hydrolysis and treated with RNase A.

**Immunofluorescence.**

Log phase cells were fixed with 4% formaldehyde and washed 2 times with 1x phosphate-buffered saline (PBS). Cells were resuspended in sorbitol buffer (1.2 M sorbitol and 20 mM potassium phosphate buffer, pH 7.2) and spheroplated with β-mercaptoethanol (M7522; Sigma-Aldrich) and zymolase at 30°C. Spheroplasted cells were added onto polylysine coated slides and incubated with primary antibody 1:1000 J2 anti-dsRNA (SCICONS) overnight. The cells were detected with 1:1000 cyanine3 conjugated goat anti-mouse (Jackson ImmunoResearch Laboratories) and stained with DAPI. The images were acquired with Zeiss Axio Imager Z1 and processed with ImageJ.

**Flow cytometry.**

Log phase cells were stained with propidium iodide, fixed with 4% formaldehyde, and washed 3 times with 1x PBS. Fluorescence intensity was measured with Becton Dickinson LSR II flow cytometer and analyzed with Flowing Software. The median GFP intensities of three biological replicates were plotted using R ggplot2.

**High copy suppression screen.**

A plasmid library of random genomic inserts in the YEP24 plasmid backbone was transformed into a *nuc1Δ ski3Δ* strain and spread onto -URA agar plate for single colonies[47]. Potential suppressors were obtained by growing at 37°C supplemented with glycerol. To confirm that the suppression was plasmid dependent, the potential suppressors were first grown on agar plate containing 0.1% 5-fluoroorotic acid (5-FOA) for counterselection and then transferred to a fresh agar plate at 37°C supplemented with glycerol. The plasmids of the candidates were isolated and sequenced to identify potential genes responsible for the suppression of L-A pathogenesis when over-expressed. The candidate genes were confirmed by spot analysis using p5476 MOBY plasmids.

**Acknowledgements**

We are grateful Dr. Charlie Boone and Dr. Brenda Andrews for generous sharing of strains and plasmid and to Angus McQuibban for critical comments on the manuscript.

**References**

1.   Bernheim, A., and Sorek, R. (2020). The pan-immune system of bacteria: antiviral defence as a
community resource. Nat Rev Microbiol *18*, 113-119. 10.1038/s41579-019-0278-2.
2.   Wickner, R.B., Fujimura, T., and Esteban, R. (2013). Viruses and prions of Saccharomyces
cerevisiae. Adv Virus Res *86*, 1-36. 10.1016/B978-0-12-394315-6.00001-5.
3.   Schmidt, N., Lareau, C.A., Keshishian, H., Ganskih, S., Schneider, C., Hennig, T., Melanson, R.,
Werner, S., Wei, Y., Zimmer, M., et al. (2021). The SARS-CoV-2 RNA-protein interactome in
infected human cells. Nat Microbiol *6*, 339-353. 10.1038/s41564-020-00846-z.
4.   Smith, R.W., and Gray, N.K. (2010). Poly(A)-binding protein (PABP): a common viral target.
Biochem J *426*, 1-12. 10.1042/BJ20091571.
5.   Brown, J.T., Bai, X., and Johnson, A.W. (2000). The yeast antiviral proteins Ski2p, Ski3p, and
Ski8p exist as a complex in vivo. RNA *6*, 449-457.
6.   Benard, L., Carroll, K., Valle, R.C., Masison, D.C., and Wickner, R.B. (1999). The ski7 antiviral
protein is an EF1-alpha homolog that blocks expression of non-Poly(A) mRNA in
Saccharomyces cerevisiae. J Virol *73*, 2893-2900. 10.1128/JVI.73.4.2893-2900.1999.
7.   Anderson, J.S., and Parker, R.P. (1998). The 3' to 5' degradation of yeast mRNAs is a general
mechanism for mRNA turnover that requires the SKI2 DEVH box protein and 3' to 5'
exonucleases of the exosome complex. EMBO J *17*, 1497-1506. 10.1093/emboj/17.5.1497.
8.   Ball, S.G., Tirtiaux, C., and Wickner, R.B. (1984). Genetic Control of L-a and L-(Bc) Dsrna
Copy Number in Killer Systems of SACCHAROMYCES CEREVISIAE. Genetics *107*, 199-
217.
9.   Rowley, P.A., Ho, B., Bushong, S., Johnson, A., and Sawyer, S.L. (2016). XRN1 Is a Species-
Specific Virus Restriction Factor in Yeasts. PLoS Pathog *12*, e1005890.
10.1371/journal.ppat.1005890.
10.  Burgess, H.M., and Mohr, I. (2015). Cellular 5'-3' mRNA exonuclease Xrn1 controls double-
stranded RNA accumulation and anti-viral responses. Cell Host Microbe *17*, 332-344.
10.1016/j.chom.2015.02.003.
11.  Ng, C.S., Kasumba, D.M., Fujita, T., and Luo, H. (2020). Spatio-temporal characterization of the
antiviral activity of the XRN1-DCP1/2 aggregation against cytoplasmic RNA viruses to prevent
cell death. Cell Death Differ *27*, 2363-2382. 10.1038/s41418-020-0509-0.
12.  Eckard, S.C., Rice, G.I., Fabre, A., Badens, C., Gray, E.E., Hartley, J.L., Crow, Y.J., and Stetson,
D.B. (2014). The SKIV2L RNA exosome limits activation of the RIG-I-like receptors. Nat
Immunol *15*, 839-845. 10.1038/ni.2948.
13.  Miyashita, M., Oshiumi, H., Matsumoto, M., and Seya, T. (2011). DDX60, a DEXD/H box
helicase, is a novel antiviral factor promoting RIG-I-like receptor-mediated signaling. Mol Cell
Biol *31*, 3802-3819. 10.1128/MCB.01368-10.
14.  Gao, J., Chau, S., Chowdhury, F., Zhou, T., Hossain, S., McQuibban, G.A., and Meneghini,
M.D. (2019). Meiotic viral attenuation through an ancestral apoptotic pathway. Proc Natl Acad
Sci U S A *116*, 16454-16462. 10.1073/pnas.1900751116.
15.  Li, L.Y., Luo, X., and Wang, X. (2001). Endonuclease G is an apoptotic DNase when released
from mitochondria. Nature *412*, 95-99. 10.1038/35083620.
16.  Thomson, B.J. (2001). Viruses and apoptosis. Int J Exp Pathol *82*, 65-76. 10.1111/j.1365-
2613.2001.iep0082-0065-x.
17.  Eastwood, M.D., Cheung, S.W., Lee, K.Y., Moffat, J., and Meneghini, M.D. (2012).
Developmentally programmed nuclear destruction during yeast gametogenesis. Dev Cell *23*, 35-
44. 10.1016/j.devcel.2012.05.005.
18.  Eastwood, M.D., and Meneghini, M.D. (2015). Developmental Coordination of Gamete
Differentiation with Programmed Cell Death in Sporulating Yeast. Eukaryot Cell *14*, 858-867.
10.1128/EC.00068-15.

19.    Koonin, E.V., and Krupovic, M. (2019). Origin of programmed cell death from antiviral
279           defense? Proc Natl Acad Sci U S A *116*, 16167-16169. 10.1073/pnas.1910303116.
280    20.    Weber, F., Wagner, V., Rasmussen, S.B., Hartmann, R., and Paludan, S.R. (2006). Double-
281           stranded RNA is produced by positive-strand RNA viruses and DNA viruses but not in
282           detectable amounts by negative-strand RNA viruses. J Virol *80*, 5059-5064.
283           10.1128/JVI.80.10.5059-5064.2006.
284    21.    Welsch, S., Miller, S., Romero-Brey, I., Merz, A., Bleck, C.K., Walther, P., Fuller, S.D., Antony,
285           C., Krijnse-Locker, J., and Bartenschlager, R. (2009). Composition and three-dimensional
286           architecture of the dengue virus replication and assembly sites. Cell Host Microbe *5*, 365-375.
287           10.1016/j.chom.2009.03.007.
288    22.    Fernandez de Castro, I., Tenorio, R., and Risco, C. (2016). Virus assembly factories in a lipid
289           world. Curr Opin Virol *18*, 20-26. 10.1016/j.coviro.2016.02.009.
290    23.    Liu, Y.X., and Dieckmann, C.L. (1989). Overproduction of yeast viruslike particles by strains
291           deficient in a mitochondrial nuclease. Mol Cell Biol *9*, 3323-3331.
292    24.    Johnson, A.W., and Kolodner, R.D. (1995). Synthetic lethality of sep1 (xrn1) ski2 and sep1
293           (xrn1) ski3 mutants of Saccharomyces cerevisiae is independent of killer virus and suggests a
294           general role for these genes in translation control. Mol Cell Biol *15*, 2719-2727.
295    25.    Sobel, S.G., and Wolin, S.L. (1999). Two yeast La motif-containing proteins are RNA-binding
296           proteins that associate with polyribosomes. Mol Biol Cell *10*, 3849-3862.
297           10.1091/mbc.10.11.3849.
298    26.    Proweller, A., and Butler, J.S. (1996). Ribosomal association of poly(A)-binding protein in
299           poly(A)-deficient Saccharomyces cerevisiae. J Biol Chem *271*, 10859-10865.
300           10.1074/jbc.271.18.10859.
301    27.    Brandman, O., Stewart-Ornstein, J., Wong, D., Larson, A., Williams, C.C., Li, G.W., Zhou, S.,
302           King, D., Shen, P.S., Weibezahn, J., et al. (2012). A ribosome-bound quality control complex
303           triggers degradation of nascent peptides and signals translation stress. Cell *151*, 1042-1054.
304           10.1016/j.cell.2012.10.044.
305    28.    Anckar, J., and Sistonen, L. (2011). Regulation of HSF1 function in the heat stress response:
306           implications in aging and disease. Annu Rev Biochem *80*, 1089-1115. 10.1146/annurev-
307           biochem-060809-095203.
308    29.    Sorger, P.K., and Pelham, H.R. (1987). Purification and characterization of a heat-shock element
309           binding protein from yeast. EMBO J *6*, 3035-3041.
310    30.    Li, J., Labbadia, J., and Morimoto, R.I. (2017). Rethinking HSF1 in Stress, Development, and
311           Organismal Health. Trends Cell Biol *27*, 895-905. 10.1016/j.tcb.2017.08.002.
312    31.    Stark, C., Breitkreutz, B.J., Reguly, T., Boucher, L., Breitkreutz, A., and Tyers, M. (2006).
313           BioGRID: a general repository for interaction datasets. Nucleic Acids Res *34*, D535-539.
314           10.1093/nar/gkj109.
315    32.    Koonin, E.V. (1997). A conserved ancient domain joins the growing superfamily of 3'-5'
316           exonucleases. Curr Biol *7*, R604-606. 10.1016/s0960-9822(06)00311-3.
317    33.    Espert, L., Degols, G., Gongora, C., Blondel, D., Williams, B.R., Silverman, R.H., and Mechti,
318           N. (2003). ISG20, a new interferon-induced RNase specific for single-stranded RNA, defines an
319           alternative antiviral pathway against RNA genomic viruses. J Biol Chem *278*, 16151-16158.
320           10.1074/jbc.M209628200.
321    34.    Zuo, Y., and Deutscher, M.P. (2001). Exoribonuclease superfamilies: structural analysis and
322           phylogenetic distribution. Nucleic Acids Res *29*, 1017-1026. 10.1093/nar/29.5.1017.
323    35.    Hanekamp, T., and Thorsness, P.E. (1999). YNT20, a bypass suppressor of yme1 yme2, encodes
324           a putative 3'-5' exonuclease localized in mitochondria of Saccharomyces cerevisiae. Curr Genet
325           *34*, 438-448. 10.1007/s002940050418.

36. van Hoof, A., Lennertz, P., and Parker, R. (2000). Three conserved members of the RNase D
family have unique and overlapping functions in the processing of 5S, 5.8S, U4, U5, RNase
MRP and RNase P RNAs in yeast. EMBO J *19*, 1357-1365. 10.1093/emboj/19.6.1357.
37. Szewczyk, M., Malik, D., Borowski, L.S., Czarnomska, S.D., Kotrys, A.V., Klosowska-Kosicka,
K., Nowotny, M., and Szczesny, R.J. (2020). Human REXO2 controls short mitochondrial RNAs
generated by mtRNA processing and decay machinery to prevent accumulation of double-
stranded RNA. Nucleic Acids Res *48*, 5572-5590. 10.1093/nar/gkaa302.
38. Grover, R., Burse, S.A., Shankrit, S., Aggarwal, A., Kirty, K., Narta, K., Srivastav, R., Ray,
A.K., Malik, G., Vats, A., et al. (2019). Myg1 exonuclease couples the nuclear and mitochondrial
translational programs through RNA processing. Nucleic Acids Res *47*, 5852-5866.
10.1093/nar/gkz371.
39. Dwivedi, M., Laddha, N.C., and Begum, R. (2013). Correlation of increased MYG1 expression
and its promoter polymorphism with disease progression and higher susceptibility in vitiligo
patients. J Dermatol Sci *71*, 195-202. 10.1016/j.jdermsci.2013.04.026.
40. Zhuang, T., Yi, X., Chen, J., Kang, P., Chen, X., Chen, J., Cui, T., Chang, Y., Ye, Z., Ni, Q., et
al. (2020). Intracellular virus sensor MDA5 exacerbates vitiligo by inducing the secretion of
chemokines in keratinocytes under virus invasion. Cell Death Dis *11*, 453. 10.1038/s41419-020-
2665-z.
41. Kopek, B.G., Perkins, G., Miller, D.J., Ellisman, M.H., and Ahlquist, P. (2007). Three-
dimensional analysis of a viral RNA replication complex reveals a virus-induced mini-organelle.
PLoS Biol *5*, e220. 10.1371/journal.pbio.0050220.
42. West, A.P., Shadel, G.S., and Ghosh, S. (2011). Mitochondria in innate immune responses. Nat
Rev Immunol *11*, 389-402. 10.1038/nri2975.
43. Sitron, C.S., Park, J.H., Giafaglione, J.M., and Brandman, O. (2020). Aggregation of CAT tails
blocks their degradation and causes proteotoxicity in S. cerevisiae. PLoS One *15*, e0227841.
10.1371/journal.pone.0227841.
44. Magtanong, L., Ho, C.H., Barker, S.L., Jiao, W., Baryshnikova, A., Bahr, S., Smith, A.M.,
Heisler, L.E., Choy, J.S., Kuzmin, E., et al. (2011). Dosage suppression genetic interaction
networks enhance functional wiring diagrams of the cell. Nat Biotechnol *29*, 505-511.
10.1038/nbt.1855.
45. Bilanchone, V., Clemens, K., Kaake, R., Dawson, A.R., Matheos, D., Nagashima, K., Sitlani, P.,
Patterson, K., Chang, I., Huang, L., and Sandmeyer, S. (2015). Ty3 Retrotransposon Hijacks
Mating Yeast RNA Processing Bodies to Infect New Genomes. PLoS Genet *11*, e1005528.
10.1371/journal.pgen.1005528.
46. Sakaue, H., Shiota, T., Ishizaka, N., Kawano, S., Tamura, Y., Tan, K.S., Imai, K., Motono, C.,
Hirokawa, T., Taki, K., et al. (2019). Porin Associates with Tom22 to Regulate the
Mitochondrial Protein Gate Assembly. Mol Cell *73*, 1044-1055 e1048.
10.1016/j.molcel.2019.01.003.
47. Kamada, Y., Fujioka, Y., Suzuki, N.N., Inagaki, F., Wullschleger, S., Loewith, R., Hall, M.N.,
and Ohsumi, Y. (2005). Tor2 directly phosphorylates the AGC kinase Ypk2 to regulate actin
polarization. Mol Cell Biol *25*, 7239-7248. 10.1128/MCB.25.16.7239-7248.2005.
367

**Figure 1**. L-A attenuation protects yeast from lethal pathogenesis. **(A)** An ethidium bromide-stained gel of total RNA prepared from the indicated strains is shown, with the 4.6 Kb L-A dsRNA band indicated with an arrow. **(B)** Immunofluorescence was used to visualize L-A dsRNA (red) in cells of the indicated genotypes. These strains were cured of the weakly abundant L-BC dsRNA virus to eliminate background staining (see Methods). DAPI staining of DNA is shown in blue. Scale bar, 1 μm. **(C)** Western blotting of L-A Gag and Pgk1 protein levels in the indicated strains is shown. Molecular weight markers are indicated on the right. **(D)** Spot test growth assays of the strains from 1C is shown. Strains were spotted on -Leu media containing either glucose or glycerol and grown at the indicated temperatures. **(E)** The mitochondrial proteins Nuc1 and Rex2 collaborate with the cytosolic proteins Xrn1, SkiC, and Myg1 to regulate L-A protein level and ensure cell fitness. The translation control factors Pab1 and Sro9 alleviate L-A pathogenesis by suppressing L-A level while Slf1, the paralog of Sro9, suppress L-A pathogenesis without affecting L-A level. Newly discovered L-A attenuation factors are indicated in red.

**Figure 2**. New antiviral factors are identified by exploiting the L-A pathogenesis. **(A)** Spot test growth assays of the high copy suppressors *SRO9*, *SLF1*, and *PAB1* are shown. **(B)** Western blotting detection of L-A Gag, Pgk1, Sro9, and Slf1 protein levels in the strains from 2A is shown. Molecular weight markers are indicated on the right. **(C)** Flow cytometry was used to measure HSE-GFP expression in the indicated strains. Median GFP intensity is shown. n = 3. * $p$ < 0.05, ** $p$ < 0.01, *** $p$ < 0.001. The $p$ value was calculated using the unpaired student's t-test. **(D)** Spot analysis of strains defective in *NUC1* and *REX2* is shown. Strains are spotted on SC media containing either glucose or glycerol and grown at the indicated temperature. **(E)** Western blotting of L-A Gag and Pgk1 protein levels of strains in Figure 2D. Molecular weight markers are indicated on the right.

**Figure S1**. Additional images of immunofluorescence of dsRNA (red) of strains from Figure 1B.
**(A)** Immunofluorescence of dsRNA (red) of indicated genotype is shown. Cells are stained with
DAPI to visualize nuclei. Scale bar, 1 μm. **(B)** Images of fields of cells of indicated genotype is
shown. Cells are stained with DAPI to visualize nuclei. Scale bar, 30 μm.

A

B

**Figure S2**. *XRN1*, *NUC1*, and *SKI3* work in parallel pathways to regulate L-A Gag level. **(A)** Spot analysis of strains defective in *NUC1* and *XRN1* is shown. Strains are spotted on SC media containing either glucose or glycerol and grown at the indicated temperature. **(B)** Western blotting of L-A Gag and Pgk1 protein levels of strains in Figure S2A. Molecular weight markers are indicated on the right.

**Figure S3**. L-A pathogenesis correlates with proteostatic stress. Flow cytometry was used to measure HSE-GFP expression in the indicated strains. Median GFP intensity is shown. n = 3. * $p$ < 0.05, ** $p$ < 0.01. The $p$ value was calculated using unpaired student's t-test.

**Figure S4**. The mitochondrial exonuclease *REX2* is an antiviral factor. Spot analysis of strains defective in 3 parallel antiviral pathways containing plasmid expressing *NUC1* is shown. Strains are spotted on -URA media or synthetic complete (SC) media supplemented with 0.1% 5-fluoroorotic acid (5-FOA).

A

B

C

**Figure S5**. *MYG1* collaborates with *NUC1* and *SKI3* to ensure cell fitness in a L-A dependent manner. **(A)** Spot analysis of strains defective in *MYG1*, *NUC1*, and *SKI3* is shown. Strains are spotted on SC media containing either glucose or glycerol and grown at the indicated temperature. **(B)** Western blotting detection of L-A Gag and Pgk1 protein levels in the indicated strains is shown. Molecular weight markers are indicated on the right. **(C)** Spot analysis of strains defective in *MYG1*, *NUC1*, and *SKI3* with and without L-A is shown. Strains are spotted on SC media containing either glucose or glycerol and grown at the indicated temperature.