Clinical phenotypes in acute and chronic infarction explained through human ventricular electromechanical modelling and simulations

Xin Zhou*[1], Zhinuo Jenny Wang*[1], Julia Camps[1], Jakub Tomek[2], Alfonso Santiago[3,4],

Adria Quintanas[3], Mariano Vazquez[3,4], Marmar Vaseghi[5,6], Blanca Rodriguez[1]

* co-first authors

1 Department of Computer Science, University of Oxford, UK

2 Department of Physiology, Anatomy and Genetics, University of Oxford, UK

3 Department of Computer Applications in Science and Engineering, Barcelona Supercomputing

Centre (BSC), Barcelona, Spain

4 ELEM Biotech, Barcelona, Spain

5 UCLA Cardiac Arrhythmia Center, University of California, Los Angeles, CA, USA

6 Neurocardiology Research Center of Excellence, University of California, Los Angeles, CA, USA

Author contact details:

Dr Xin Zhou: xin.zhou@cs.ox.ac.uk

Dr Zhinuo Jenny Wang: jenny.wang@cs.ox.ac.uk

Professor Blanca Rodriguez: blanca.rodriguez@cs.ox.ac.uk

Oxford address: Department of Computer Science, Wolfson Building, Parks Road, Oxford OX1

3QD, UK

Word count: 7913 excluding references only, 7589 excluding references and the abstract,

10,110 in total (excluding title page).

ABSTRACT

Aims:

Sudden death after myocardial infarction (MI) is associated with electrophysiological heterogeneities and ionic remodelling, which are reflected as variable phenotypes. Low ejection fraction (EF) is used in risk stratification, but its mechanistic links with the post-MI pro-arrhythmic heterogeneities are unknown. We aim to provide a mechanistic explanation of clinical phenotypes in acute and chronic MI, from ionic remodeling to ECG and EF, using human electromechanical modelling and simulation to augment experimental and clinical investigations.

Methods and Results:

A human ventricular electromechanical modelling and simulation framework is constructed and validated with rich experimental and clinical datasets. Abnormalities caused by scar and border zone ionic remodeling are introduced in varying degrees as reported in experimental data obtained in acute and chronic infarction. Simulations enabled reproducing and explaining clinical phenotypes post-MI, from ionic remodelling to ECGs and pressure-volume loops. In acute MI, T-wave inversion and Brugada phenocopy were explained by up to 57 ms of local APD prolongation and activation failure due to the inhibition of potassium, sodium and calcium channels in the border zone. In chronic MI, upright tall T-waves highlight large repolarisation dispersion caused by uneven potassium channel expression in border and remote zones, which promoted ectopic propagation at fast pacing. Post-MI ionic remodelling reduced EF by up to 10% through inhibition of calcium transient amplitude due to weaker calcium currents or SERCA activity, but the EF at resting heart rate was not

sensitive to the extent of repolarisation heterogeneity and the risk of repolarisation

abnormalities at fast pacing.

Conclusions:

Multi-scale modelling and simulation coherently integrates experimental and clinical data at subcellular, tissue, and organ scales to unravel electromechanical disease mechanisms in MI. In acute post-MI, ionic remodelling and its effect on refractoriness and propagation failure in the BZ have a strong impact on phenotypic ECG variability, whereas in chronic post-MI, the repolarisation dispersion across the BZ is crucial. T-wave and QT abnormalities are better indicators of repolarisation heterogeneities than EF in post-MI.

1 INTRODUCTION

Sudden cardiac death in post-myocardial infarction (post-MI) patients is due to lethal

arrhythmias in 50% of cases at both acute and chronic infarct stages (Vazquez *et al.*, 2009;

Elayi *et al.*, 2017). Risk stratification is currently based on low left ventricular ejection

fraction (LVEF) (Solomon *et al.*, 2005) to identify patients who need the implantation of

defibrillator device. However, only a very small subset of patients that suffer from sudden

cardiac death are identified by low LVEF, and a significant number of sudden deaths occur in

patients with relatively preserved LVEF (Vaduganathan *et al.*, 2018). The mechanistic link

between electrophysiological heterogeneities, ECG and LVEF phenotypes that underpin

arrhythmic events is not clear. A precision medicine approach using computational modelling

and simulations could help to elucidate disease mechanisms and provide an *in silico*

alternative for therapy evaluation and risk stratification. However, a key hurdle in clinical

adoption of such *in silico* tools is providing proof of model credibility through validation

studies. In this study, we set out to achieve the dual aim of demonstrating a multi-scale

approach for model validation and to identify important electrophysiological mechanisms for

post-MI risk stratifications.

Various electrocardiogram (ECG) characteristics were suggested by clinical studies to

be relevant to the arrhythmic risk of post-MI patients. Longer QT intervals have been

associated with increased mortality as well as ventricular tachycardia and fibrillation for both

the acute and chronic MI (Ahnve, 1985; Oikarinen *et al.*, 1998). However, QT prolongation

in single leads can be a reflection of reduced global repolarisation reserve, while the regional

heterogeneity of repolarisation in different post-MI zones can be more crucial (Cluitmans *et*

al., 2021) than the global reserve for the development of re-entrant waves. QT dispersion

between 12-lead ECGs was proposed as a potential marker for electrophysiological heterogeneity (Spargias *et al.*, 1999); however, it was found to have good specificity but low sensitivity (Spargias *et al.*, 1999). Prolonged T-peak to T-end interval (Tpe) (Shenthar *et al.*, 2015) and increased incidence of T-wave alternans (Bloomfield *et al.*, 2004) were also found to be useful risk predictors in post-MI patients. Other ECG metrics, such as QRST integral and spatial ventricular gradients have also shown some promise in improving SCD prediction (Waks *et al.*, 2016). However, despite the promising outcomes of those clinical studies, the ECG-based risk predictors are not widely applied in the clinical evaluation of the need for implantable cardioverter-defibrillators (Al-Khatib *et al.*, 2018).

A key factor that hinders the clinical utility of ECG biomarkers is the large variability in post-MI phenotypes, their progression from acute to chronic, and between different patients. Variability in QT prolongation (Ahnve, 1985) and post-MI ECG morphologies constrain the use of single biomarker thresholds as predictors. Furthermore, an important limitation of non-invasive ECG biomarkers is their inability to provide direct measurements of regional electrophysiological heterogeneity caused by scar and ionic remodelling, which is critical for the arrhythmic substrate.

Many experimental studies have investigated the underlying mechanisms of increased repolarisation dispersion in post-MI patients. Specifically, repolarisation dispersion increases after MI due to ionic differences between the border zone (BZ) surrounding the scar and the remote zone (RZ) myocardium (Mendonca Costa *et al.*, 2018). The BZ at the acute MI exhibits reduced sodium current (I_{Na}), L-type calcium current (I_{CaL}) (Aggarwal Rajesh & Boyden Penelope A., 1995), and rapid and slow delayed rectifier potassium currents (I_{Kr} and I_{Ks}) (Jiang *et al.*, 2000), as well as enhanced CaMKII activity (Hund *et al.*, 2008*a*) and gap

junction redistribution (Yao *et al.*, 2003). After a period of scar healing, ionic remodelling

may partially recover (Ursell *et al.*, 1985), but increased late sodium current (I_{NaL}), calcium-

activated potassium and chloride currents (I_{KCa} and I_{ClCa}) were observed in the chronic BZ and

RZ (Hegyi *et al.*, 2018). Post-MI ionic remodelling can cause conduction abnormalities,

repolarisation abnormalities, as well as variable alterations in the action potential duration

(APD), which act as substrates of ventricular tachycardia (Pogwizd *et al.*, 1992). It is,

however, unclear how various degrees of repolarisation dispersion affect ECG biomarkers

and LVEF used for risk stratification. Therefore, further studies are needed to bridge the gap

between cellular electrophysiological characteristics and variable patient phenotypes post-

MI. Furthermore, the wealth of cellular, tissue, and ECG data described in the literature

provides an excellent multi-scale dataset for model validation. Comparisons of simulated

action potential and ECG biomarkers with experimental and clinical data under acute and

chronic MI conditions help to improve the credibility of the computational model.

Human-based modelling and simulations of ventricular electrophysiology and

mechanics have demonstrated accurate prediction of post-MI arrhythmic risk and myocardial

stress and strain (Arevalo *et al.*, 2016; Sack *et al.*, 2016). However, previous work

concentrated on either electrophysiology or mechanics, while the crosstalk between the two,

and the relationship between LVEF and arrhythmic risk, is yet to be investigated.

The main goal of this study is, therefore, to quantify the contribution of varying

degrees of ionic remodelling to the phenotypic variability in ECG and LVEF biomarkers

observed in acute and chronic post-MI patients, and thereby create and validate models of

post-MI electromechanics. Using state-of-the-art electromechanical human biventricular

simulations, we aim to identify biomarkers that are most representative to the pro-arrhythmic

substrate for each state, thus facilitating post-MI risk stratifications to go beyond LVEF. We

hypothesise that different ECG abnormalities are explained by different degrees of ionic

remodelling leading to activation sequence abnormalities, dispersion of repolarisation, early

after depolarisations (EADs) and alternans, whereas LVEF is insensitive to ionic remodelling

that underpins ECG disease markers and reflects predominantly calcium transient and

structural abnormalities.

METHODS

1. Human multi-scale ventricular electromechanical modelling and simulation: from ionic remodelling to ECG and LVEF.

A human ventricular electromechanical modelling and simulation framework is constructed

and evaluated using experimental and clinical data to enable the investigations of variable

post-MI patient phenotypes, from ionic remodelling to body surface ECGs and pressure-

volume (PV) loops (Figure 1A). The cardiac magnetic resonance (CMR)-based biventricular

anatomical mesh (Figure 1B) was included in the torso for ECG simulations (Wang *et al.*,

2021). Electrical propagation was simulated using the monodomain equation with orthotropic

diffusion based on rule-based fields for fibre directions (Streeter *et al.*, 1970) with sheet

directions normal to the endocardial/epicardial surface (Levrero-Florencio *et al.*, 2020)

(Figure 1B). Transmural and apex-to-base heterogeneities (Mincholé *et al.*, 2019) were

introduced and a sensitivity analysis was performed to investigate their implications in ECG

biomarkers and LVEF (Figure 1B). Electrical stimulus via Purkinje-myocardial junctions

was simulated by an endocardial fast-activation layer with root node locations to achieve

realistic QRS complex morphologies simulated at clinically standard lead locations (Figure

1   1C) (Mincholé *et al.*, 2019). In healthy tissue (normal zone (NZ)), monodomain diffusivities

2   were calibrated to achieve experimentally measured orthotropic conduction velocities of 67

3   cm/s, 30 cm/s, and 17 cm/s (Caldwell *et al.*, 2009).

Figure 1: Human-based multi-scale modelling and simulation in acute and chronic

myocardial infarction. (A) Simulated clinical ECG and pressure volume biomarkers. (B)

Ventricular electrophysiology is simulated using a fast endocardial activation layer to

approximate Purkinje-myocardial junction, experimentally-informed transmural and apico-

basal heterogeneities in action potential duration, and transmurally varying myocyte

orientation. Mechanical pumping behaviour is modelled by coupling the intraventricular

pressures with a two-element Windkessel model of arterial haemodynamics with a fixed

basal plane. (C) An anterior 75% transmural infarction is modelled with acute and chronic

ionic remodelling embedded in the border zone and remote zones. Standard 12-lead ECG

was evaluated at standard body-surface locations. (D) Simulated action potentials using

populations of human ventricular models in healthy (baseline) and acute and chronic post-

MI conditions with different degrees of ionic remodelling . (E) Schematic representation of

ionic fluxes, calcium dynamics and actin/myosin contraction mechanisms in the human

ventricular electromechanically-coupled cellular model.

Since anterior infarction is very common and related to the worst prognosis (Stone *et al.*, 1988), 75% transmural extent of infarct from the endocardial surface was introduced to the anterior myocardial wall (Figure 1C) to match the clinical definition of transmural infarctions (Sato *et al.*, 2008), with scar and border zones (maximum width 0.5 cm) taking up 12.4% and 11.7% of ventricular volume (or 15.8% and 15.0% of left ventricular volume), respectively. Maximum border zone width was based on various reports of systolic strain dysfunction in < 1 cm proximity to the infarct (Gallagher *et al.*, 1986; Van Leuven *et al.*, 1994). Electrophysiological remodelling was implemented in the border zone (BZ) and the

chronic remote zone (RZ), which covered the entire non-infarcted and non-BZ region in both

ventricles. In BZ and infarct zones, the diffusivities were calibrated to reproduce conduction

slowing (one-third of the NZ conduction velocities). For each virtual cardiomyocyte, the

electromechanical single cell model considered the human-based ToR-ORd

electrophysiological model (Tomek *et al.*, 2019) (extensively validated in control, disease

and drug block conditions) coupled with human excitation-contraction and active tension

Land model (Land *et al.*, 2017; Levrero-Florencio *et al.*, 2020) (Figure 1E).

The human biventricular model incorporated strongly-coupled electromechanics with

orthotropic passive mechanical behaviour and balance of linear momentum with inertial

effects, as in our previous work (Levrero-Florencio *et al.*, 2020; Margara *et al.*, 2021; Wang

et al., 2021). In brief, firstly, intracellular calcium concentration drives crossbridge cycling

and force production through unblocking the crossbridge binding-site. Secondly, calcium

sensitivity and force production are a function of fibre stretch ratio. Thirdly, stretch rate

affects distortion-dependent cross-bridge unbinding (Land *et al.*, 2017) and the diffusivity

tensor is transformed using the deformation gradient tensor such that the prescribed values

describe conductivities in the deformed state (Levrero-Florencio *et al.*, 2020).

An elastic spring boundary condition was set to act perpendicularly to the epicardial

surface, to simulate pericardial constraint, and the basal plane was fixed in space to prevent

unphysiological tilting and expansion due to the unavailability of closed basal geometry.

Pressure boundary condition on the left and right endocardial surfaces were controlled using

a series of five equations (see SM2) that controls 1) active diastolic inflation followed by

electrical activation and 2) isovolumic contraction, 3) ejection coupled with two-element

Windkessel aortic haemodynamics model, 4) isovolumic relaxation, and 5) passive inflation and relaxation (Wang *et al.*, 2021).

The sheet active tension was set to 30% of fibre active tension to achieve sufficient LVEF in control conditions, based on the results from a sensitivity analysis in Supplementary Material (SM) section SM5. In brief, we evaluated the sensitivity of ECG morphology and LVEF to changes in the calcium sensitivity of troponin binding in excitation-contraction coupling and the active tension in the sheet direction as a percentage of that in the fibre direction.

The passive stiffness parameters were calibrated based on a previous sensitivity analysis (Wang *et al.*, 2021) to achieve 53% LVEF and a physiological pressure-volume (PV) loop in a control simulation (see SM Table S2 and S3 for a list of calibrated parameters).

The active tension was set to zero in the scar to represent the myocyte damage, and the chronic passive stiffness parameters of the infarct region were increased 10-fold to mimic fibrotic scar formation (Sun *et al.*, 2009). For each acute post-MI phenotype, a case of complete loss of contractile function (zero active tension) in the BZ was also simulated to evaluate the contribution of other non-ionic remodelling-related abnormalities on ejection fraction.

2. Experimentally-informed single cell populations of human post-MI electromechanical models.

To account for the inter-subject electrophysiological variability widely observed in clinical data, the baseline human cellular electromechanical ToR-Land model was extended to

populations of healthy cellular models. Then, post-MI ionic remodelling was applied to

generate populations of post-MI virtual cardiomyocytes (Figure 1D). In addition to the

baseline ToR-ORd model, several representative cellular models were selected from the

population and implemented into the biventricular electromechanical simulations.

An initial population of human ventricular cell models was constructed based on the

ToR-ORd model by varying the conductances or magnitudes of I_{Na}, I_{NaL}, I_{to}, I_{CaL}, I_{Kr}, I_{Ks}, I_{K1},

I_{NaCa}, I_{NaK}, J_{rel} and J_{up} by up to ±50% using Latin Hypercube Sampling (SM Figure S1 and S2).

As illustrated in previous studies, small populations of models with proarrhythmic ionic

remodelling achieved similar predictability as large populations with uniform current

variations (Zhou *et al.*, 2019). Therefore, we chose to start with a small healthy population

(n=500) and then introduce multiple combinations of ionic remodelling to mimic the large

post-MI variability. After calibration with human experimental data (SM Table S1) and

discarding those manifesting EADs at 1Hz, 245 sets of endocardial, midmyocardial and

epicardial models were accepted as the healthy population (SM Figure S2).

Several degrees of post-MI ionic remodelling were applied to the healthy population

to generate BZ and RZ populations for both acute and chronic post-MI, based on

experimental measurements, as described in SM Table S4. For acute post-MI (within a week

post-occlusion), three types of BZ remodelling (Acute BZ1-3) were considered based on

previous modelling work and experimental canine data collected within 5 days post-MI. For

chronic post-MI, ionic remodelling measured from minipigs 5 months post-MI with heart

failure were used to generate Chronic BZ (affecting only the BZ) and Chronic RZ1 (also

affecting the remote myocardium). Another type of remodelling, Chronic RZ2, was

established based on multiple experimental data from failing human cardiomyocytes. The RZ

covers the entire myocardium apart from the infarct and BZ in chronic MI simulations.

Furthermore, reduction of sodium current and SERCA, with enhanced CaMKII activity and

slower calcium release were also implemented in Chronic BZ, RZ1 and RZ2, as observed in

human failing cardiomyocytes (details in SM Table S4). Both human recordings and animal

data were used for model evaluation, given the scarcity of human tissue, summarised in SM

Table S4-S5. Action potential, calcium transient, and active tension characteristics of the

post-MI models are summarised in SM Table S6.

3. Simulation protocols and biomarker calculation.

Human virtual ventricular myocytes were paced at 1Hz for 500 beats to detect EAD

generation. For alternans generation, single cells were paced at cycle lengths (CLs) of

500 ms, 400 ms and 300 ms for 500 beats, and a ΔAPD greater than 3 ms between the last

two beats at steady state was defined as alternans.

Biventricular electromechanical simulations were performed at 800 ms CL (75 beats

per minute) with 100 ms allowed for active diastolic filling prior to endocardial activation for

each beat. For chronic post-MI, and baseline for comparison, an additional fast pacing

protocol was applied with 500 ms CL (120 beats per minute), with 50 ms allowed for active

diastolic filling for each beat. For these fast pacing chronic MI simulations, single cell

models with arrhythmic abnormalities were selected from the population of models for

simulation at the ventricular scale. Three beats were sufficient to achieve converged ECG

and PV characteristics at 800 ms CL, and six beats for 500 ms CL (see Figures S7~S8 for

ECGs for all beats).

Clinical biomarkers were quantified from the simulated ECG (including the QT interval, QRS duration, T-wave duration, T peak to T end duration, T onset to T peak duration, and QT dispersion, see definition and method of evaluation in SM2 and biomarker results in SM6 Table S8), and from the simulated PV loop (including end diastolic and end systolic volumes, peak systolic pressures and LV and RV ejection fractions), as well as wall thickening strain (see Table S8-S9 for ECG and PV biomarkers for all simulated beats).

4. Simulation software and computational framework.

Cellular electrophysiological simulations and Latin Hypercube Sampling were performed using bespoke MATLAB codes. Coupled cellular electromechanics, as well as biventricular electromechanics simulations, were performed using the high-performance numerical software, Alya, for complex coupled multi-physics and multi-scale problems (Santiago *et al.*, 2018) on the CSCS (Swiss National Supercomputing Centre) Piz Daint supercomputer multi-core clusters, granted through the PRACE (Partnership for Advanced Computing in Europe) project. The simulation input files and Alya executable required to replicate the simulated results are available upon request for scientific investigations.

5. Data availability

The simulation input files and Alya executable required to replicate the simulated results will be made available upon request. Python and MATLAB scripts for cellular populations of models simulations and post-processing of the biventricular simulations can be found at:

https://github.com/jennyhelyanwe/post_MI_postprocessing

RESULTS

1. Human modelling and simulation for ECG phenotypes in acute and chronic post-MI.

Figure 2 demonstrates the ability of human electromechanical simulations to reproduce a variety of clinically reported phenotypes in patients with acute and chronic infarction, in agreement with clinical measurements of ECG and pressure-volume biomarkers, as quantified in further detail in (SM Table S7, details of clinical database in SM4). When imposing acute post-MI remodelling, simulated ECGs reproduced fractionated QRS complexes, T-wave inversion, Brugada phenocopy ST-segment elevation and QT interval prolongation in the anterior leads (Figure 2A), which are common ECG phenotypes observed in acute post-MI patients. Simulations also recapitulated ECG morphology similar to healthy subjects with upright T-waves (Figure 2A, right), which can also be present in acute post-MI.

In chronic post-MI, simulated ECG displayed upright T-waves, and global prolongations of QT intervals in all precordial leads (Figure 2B, top row), which are characteristic of chronic patients with worse clinical outcomes (Ahnve, 1985; Oikarinen *et al.*, 1998). A range of T-wave durations were present, as can be found in chronic post-MI (Figure 2B, bottom row). SM4 summarises the clinical ECGs used for comparisons in Figure 2. The full 12-lead ECGs for acute and chronic post-MI simulations can be found in SM Figure S7 and S8.

Sensitivity analysis showed first that large changes in apex-to-base and transmural heterogeneities only altered T-wave amplitude but not its polarity and did not affect the ST-segment (SM Figure S3 and S4), and second that changes in mechanical parameters did not

1 affect the ECG morphology (SM Figure S5 and S6). This result supports the specificity of

2 post-MI signatures to underlying ionic remodelling.

4 *Figure 2: Agreement between simulated and clinical ECGs demonstrating variability in*

5 *clinical phenotypes in acute and chronic post-myocardial infarction (post-MI).* (A) In acute

MI, simulated ECGs show T-wave inversion (border zone model 1 (BZ1)), Brugada

phenocopy (BZ2), and normal phenotypes (BZ3), in accordance with phenotypes found in

clinical databases. (B) In chronic MI, simulated ECGs show prolonged QT and upright T-

waves with a range of amplitude and duration (remote zone model 1 and 2 (RZ1, RZ2))

comparable to those observed in clinical databases.

2. In acute MI, T-wave inversion and Brugada phenocopy can indicate reversed transmural repolarisation gradient and activation failure.

Analysis of simulation results enabled the uncovering of specific contributions of different

degrees of ionic remodelling to ECG phenotypes identified in acute versus chronic MI

(Figure 3 vs Figure 4). In acute MI, T-wave inversion phenotype was associated with a

reversed transmural repolarisation gradient (Figure 3A, repolarisation time map insets), due

to 57 ms of APD prolongation at the epicardial BZ compared with control (Figure 3B,

membrane potential). APD prolongation originated due to inhibition of multiple potassium

currents caused by BZ1 ionic remodellling (Figure 3C, the first column of I_{Kr}, and SM Table

S4 Acute BZ1).

Brugada phenocopy ECG phenotype was also observed in acute MI with BZ2

remodelling, causing regions of activation failure in the epicardial border zone (Figure 3A,

the activation and repolarisation time maps), as well as delayed repolarisation in the BZ near

the apex, caused by 29 ms of epicardial APD prolongation compared with control (Figure

3B, membrane potential). These were caused by strong inhibitions of sodium, calcium and

potassium ionic currents in the BZ (Figure 3C, the second column of I_{Na}, I_{CaL}, I_{Kr}, and SM

Table S4 Acute BZ2). In this simulation, electrical activation in the infarcted region was

preserved despite the conduction block in the BZ because of higher expressions of the L-type

calcium channel in the mid-myocardium (see SM Figure S11).

Acute MI with upright T-waves corresponded to a comparable transmural

repolarisation gradient as in control with BZ3 ionic remodelling (Figure 3A, the

repolarisation time map). In this case, the lack of potassium current inhibition accounted for

the lack of ECG manifestations (Figure 3C, the third column of I_{Kr}, and SM Table S4 Acute

BZ3).

Ionic bases of ECG abnormalities in acute MI

Figure 3: Multiscale explanation of ST and T-wave phenotypes in acute MI. *(A) Activation time maps reveal conduction delay in acute border zone in T-wave inversion and normal ST-T phenotypes, and conduction block in Brugada phenocopy, as well as large repolarisation dispersion and altered transmural repolarisation gradient in T-wave inversion and Brugada*

phenocopy. Red in activation map show regions of no activation (NA), green in

repolarisation map highlights regions of no repolarisation (NR). (B) Action potential

duration (APD) prolongation is present in T-wave inversion and Brugada phenocopy

cellular phenotypes, as well as slight shortening (normal ST-T) (red arrows), with decreased

calcium amplitude in all phenotypes, and a corresponding decrease in active tension

generation. (C) I_{Na}, I_{CaL}, and I_{Kr} remodelling underpin reduced conduction, reduced calcium

amplitude, and alterations in action potential duration, respectively, in all acute phenotypes.

3. In chronic post-MI, variable T-wave width can be explained by the extent of repolarisation dispersion between border zone and remote zone.

In chronic MI, global QT prolongation was due to APD prolongation in the remote

myocardium (Figure 4, repolarisation time maps and membrane potentials). Recovery of the

upright T-wave in the anterior leads (compared to acute MI) was due to a recovery of the

transmural repolarisation gradient (Figure 4A, repolarisation time maps, and SM Figure S10),

given the milder I_{Kr} inhibition in the border zone (Figure 4B, the first column of I_{Kr}, and SM

Table S4 Chronic BZ). Furthermore, T-wave duration in this stage was mainly determined by

the gradient between remote and border zone repolarisation times (Figure 4C, repolarisation

time maps), where more severe APD prolongation in the RZ led to larger repolarisation

gradients and, consequently, larger T-wave duration and amplitude (Figure 2B). Specifically,

there was an APD difference of 157 ms between remote and border zone cell models for the

large T-wave case versus only 12 ms for the slight QT-prolongation case, which accounts for

the differences in T-wave peak-to-end duration (162 ms vs 72 ms) and QT intervals (565 ms

vs 380 ms) between these two cases.

Ionic bases of ECG abnormalities in chronic MI

Figure 4: *Multiscale explanation of QT and T-wave phenotypes in chronic MI.* (A)

Conduction delay in chronic border zone causes slight QT prolongation and large T-wave

phenotypes, as well as large repolarisation dispersion in large T-wave. Red in activation

map show regions of no activation (NA), green in repolarisation map show regions of no

repolarisation (NR). (B) Varying degrees of action potential duration (ADP) prolongation in

the remote zone (RZ) corresponding to extent of QT prolongation (blue arrows), with

decreased calcium amplitude in remote and border zone of both phenotypes, and

corresponding decrease in active tension generation. (C) As in acute MI, I_{Na}, I_{CaL}, and I_{Kr}

remodelling underpin reduced conduction, reduced calcium amplitude, and degree of

prolongation in action potential duration, respectively, in both chronic phenotypes.

4. LVEF failed to indicate the extent of post-MI repolarisation dispersions.

In our simulations, acute MI ionic remodelling yielded mildly reduced LVEF (43% to 47%)

compared with control (53%) (Figure 5A). LVEF reductions were caused by contractile

dysfunction (Figure 5A, active tension) due to lowered calcium amplitude in BZ (Figure 3B,

intracellular calcium transient), which was directly caused by inhibitions of I_{CaL} in all acute

phenotypes (Figure 3C, I_{CaL}, and SM Table S4). A complete loss of contractile function in the

BZ resulted in a more severe reduction in LVEF in acute post-MI (to 40% for all acute

phenotypes in SM Figure S9). Stroke volumes of the left and right ventricles were well-

matched in control conditions (1 mL difference, see SM Table S9), and introducing

myocardial infarction caused a decrease of stroke volume in the left ventricles in both acute

and chronic MI (see SM Table S9).

Activation failure in Brugada phenocopy caused loss of contractility in the affected

regions in the BZ (Figure 3A, activation and repolarisation time maps) and resulted in a more

severely reduced LVEF to 43% with a substantial non-contracting region with significant

wall thinning (Figure 5A, systolic wall thinning). Acute MI with no observable ECG

abnormalities can still have reduced mechanical function, as measured by an LVEF of 47%

(Figure 5A, the last column). Comparison between this phenotype and the inverted T-wave

phenotype shows that the mechanical dysfunction can be dissociated from repolarisation

gradient and T wave abnormalities.

Chronic post-MI simulations showed mild reduction in peak systolic pressure (by

1 kPa) and some reduction in LVEF (Figure 5B), which were unaffected by differences in

repolarisation gradients. This is because there is a consistent reduction in calcium transient

amplitude in the remote zone that is independent of the extent of APD prolongation (Figure

4B, intracellular calcium transient and membrane voltage). Therefore, both the acute and

chronic electromechanical simulation results showed that the post-MI repolarisation

dispersion were not reflected by LVEF.

Cellular bases for mechanical dysfunction in MI

Figure 5: Reduced LVEF and heterogeneous systolic deformation caused by BZ ionic

remodeling in both acute and chronic post-myocardial infarction. (A) Reduced LVEF in all

acute phenotypes due to reduced active tension amplitude in the border zone (BZ1~3).

Brugada phenocopy shows the lowest LVEF due to activation block and loss of contractile

function in the border zone in addition to reduced active tension amplitude in the activated

border zone due to ionic remodelling. Reduced contractile function in infarct and border

zone results in infarct thinning and bulging in systole. Systolic cross section of control

simulation shown in black (CTL) with post-MI cross-sections superimposed. (B) Reduced

LVEF in both chronic phenotypes due to reduced active tension amplitude in remote zone

(RZ) and border zones, independent of the extent of QT prolongation (RZ1, RZ2). Scar

stiffening helped to reduce infarct bulging. Systolic cross-section of control simulation

shown in black (CTL) with post-MI cross-sections superimposed.

5. T-wave alternans and abnormal wave propagation are caused at fast pacing by cellular alternans and EADs, without reduced LVEF at resting heart rate.

Increased incidence of T-wave alternans is commonly observed in post-MI patients, and

abnormally propagating waves generated from post-MI electrophysiological heterogeneity

can trigger lethal arrhythmic events. T-wave alternans were reproduced in the ventricular

chronic MI simulations at fast pacing (Figure 6A), with RZ2 remodelling, and their

mechanisms were revealed through analysis of the high spatio-temporal resolution of

simulation data. Figure 6A shows upright T-wave morphology and preserved LVEF of 49%

at resting rate of 75 bpm (CL= 800 ms). However, at fast rates (120 bpm, CL = 500 ms),

significant beat-to-beat ST and T-wave morphology alterations were observed. This was due

to large alternans seen in mid-myocardial single cell simulations of the remote zone at CL of

500 ms (Figure 6C, green trace), with EAD-driven alternans. These results support the

importance of stress tests, since alternans in APD and T-wave can occur at fast heart rates with no sign of LVEF abnormalities at resting heart rate. This is consistent with reports that T-wave alternans under supine bicycle exercise testing was found to be predictive of arrhythmic event after acute post-MI (Ikeda *et al.*, 2000).

Simulations with the population of virtual cardiomyocytes models revealed that in addition to the EAD-driven alternans (SM Figure S14), classical calcium-driven alternans were also observed in the population of cell models (SM Figure S12 and S13). The key ionic remodelling underlying calcium-driven alternans include enhanced CaMKII activity and slower calcium release, as well as suppressed SERCA pump activity in the chronic MI, which are consistent with previous studies (Livshitz & Rudy, 2007; Zhou *et al.*, 2016; Tomek *et al.*, 2018) (SM Figure S15). I_{KCa} enhancement in the chronic MI suppressed alternans generation (detailed analysis provided in SM7, Figures S16 to S21, Table S10). The median of calcium amplitude was larger in the alternans models than in the non-alternating post-MI models (SM Figure S22), in agreement with the preserved LVEF in the simulations. We did not simulate the effect of this classical calcium-driven alternans on the ECG because the higher pacing rate at which this phenomenon occurs requires the model to include beta-adrenergic inotropic effects to preserve realistic systolic mechanical function.

Figure 6: T-wave alternans in simulations underpinned by APD and calcium alternans at

fast pacing (120 bpm), albeit with preserved left ventricular ejection fraction (LVEF=49%)

at rest (75 bpm) for the chronic MI phenotype (A). (B) Simulated APD and calcium traces

in midmyocardial population of models with remote zone 2 (RZ2) remodelling. (C) Large

action potential and calcium transient alternans were caused by EADs in simulations at 120

bpm with midmyocardial cells affected by RZ2 ionic remodelling (green traces,

representative example at 75 versus 120 bpm). A single cell model (in green) was selected

from the population of models (in grey) for embedding into the remote region for ventricular

simulations.

Another case of chronic MI simulation showed prolonged QT interval of 669 ms and

preserved LVEF of 49% at 75 bpm (Figure 7A). At 120bpm, in this case, ECG reflected

chaotic activity (Figure 7A, fast pacing simulation) and loss of coordinated mechanical

function. Abnormal electrotonic waves were caused by large repolarization dispersions.

Further investigations at ventricular level showed abnormal wave propagations at fast

pacing (Figure 7B). By the end of the first heartbeat (500 ms), the anterior epicardial BZ was

fully repolarised (the red trace), but the RZ showed EADs (green, purple and yellow traces).

This means that after the second beat stimulus, at 610 ms (Figure 7B), the long APD of the

RZ prevented full activation in the postero-lateral LV (Figure 7B, green and purple marker)

while the BZ APD was shorter than RZ (Figure 7B, red marker) and so could be fully

activated. This led to a significant membrane potential gradient between the BZ (red marker)

and RZ (green marker) at 610 ms, leading to ectopic wave generation due to injury current

caused by electronic effects at 710 ms in the boundary between BZ and RZ (Figure 7C,

compare red and green traces). This ectopic wave propagated in an anti-clockwise fashion

when viewed from the base from 810 to 910 ms (Figure 7B, basal view). In the right

ventricle at 610 ms, incomplete repolarisation caused a smaller action potential to be elicited

by the sinus stimulus (Figure 7C, yellow trace), thus allowing successful propagation of the

ectopic wave from the left to the right ventricle from 910 ms to 1010 ms (Figure 7B).

This episode illustrated that EADs present in the epicardial remote zone cell model

(Figure 7D) resulted in large APD dispersion between BZ and RZ, which functioned as the

trigger of ectopic wave propagation due to electrotonic gradients. Therefore, the prolonged

global QT interval with large T-wave duration and amplitude in leads facing the infarct can

be indicative of the risk of large repolarization dispersion, while the LVEF can be preserved

at rest.

Spontaneous EADs were frequently observed in the chronic MI cellular population of

models (Figure 7D population of models, SM Figure S23), due to less I_{CaL} inhibition in the

chronic MI (left, SM Table S4). The key underlying combination of ionic remodelling for

EAD generation include the inhibition of I_{Kr} and the enhancement of I_{NaL}, which facilitate

reactivation of I_{CaL} (SM Figure S24). Additional contribution of baseline I_{CaL}, I_{Kr}, and I_{NCX}

conductances are consistently observed in all chronic EAD populations (SM Figures S25

~S28, Table S11). Similarly, the cellular models that showed EADs also had larger calcium

amplitude (SM Figure S29), suggesting a preserved LVEF.

Figure 7: Prolonged QT and preserved LVEF at rest can manifest as severely abnormal

ECG at fast heart rates in chronic MI with RZ2 (A), due to electrotonically-triggered EADs

across the border zone (B). In (B), membrane potential changes for the first 1010 ms of fast

pacing simulation, showing ectopic wave generation driven by electrotonic gradient at

710 ms (arrow from red dot to green dot in lateral view), and anticlockwise propagation of

ectopic wave starting at 810 ms (anticlockwise arrow in basal view, and arrow from green

dot to purple dot in lateral view). Ectopic wave propagates towards the right ventricle via

the posterior side at 910 ms (arrow in basal view) and at 1010 ms (arrow in basal view). (C)

Local action potential at anterior (red), lateral (green), posterior (purple), and right

ventricular (yellow) sites. (D) A population of models demonstrating chronic remote zone 2

(RZ2) remodelling in promoting EADs. A representative example was selected from the

population of models that showed EAD and was embedded in ventricular simulations.

DISCUSSIONS

In this study, human electromechanical modelling and simulation enables quantification of

the contribution of electrophysiological abnormalities to clinical phenotypes in post-MI

patients, from ionic to whole-organ dynamics (summarised in Table 1). The credibility of the

human electromechanical models and simulation results is supported by their consistency

with experimental and clinical data from ionic dynamics to ECG and LVEF biomarkers in

healthy, acute and chronic post-MI conditions. Diverse clinical ECG phenotypes are

reproduced in the simulations with different degrees of experimentally reported ionic

remodelling for acute and chronic MI; their signature on the LVEF is however weak, with

only a small reduction observed. Key findings include:

 1) In acute MI, T-wave inversion, Brugada phenocopy, and QT prolongation were explained

by reversed transmural dispersion of repolarisation in the border zone and infarct, activation

failure in the border zone, and increased repolarisation time in the border zone, respectively.

2) In chronic MI, large T-wave duration and amplitude reflects large repolarisation

dispersion between remote and border zones, and global QT prolongation was caused by AP

prolongation in the remote zone.

3) Reduction in LVEF is ubiquitous across acute and chronic post-MI phenotypes and can be

explained by decreased intracellular calcium amplitude and activation failure (in the case of

Brugada phenocopy). This effect was independent of changes in the dispersion of

repolarisation.

4) Interestingly, fast pacing simulations with T-wave alternans or abnormal propagation

driven by cellular alternans or EADs both showed preserved LVEF at resting heart rate,

highlighting the fact that preserved LVEF at rest does not guarantee low arrhythmic risk.

Table 1: *Linking clinical ECG and left ventricular ejection fraction (LVEF) phenotypes to*

tissue heterogeneities and subcellular ionic remodelling in acute and chronic post-

myocardial infarction.

Clinical Phenotypes	Tissue or Cell Level phenomena	Corresponding Post Infarction Ionic Remodelling
Acute MI T-wave inversion in ECG	Reversed transmural repolarisation gradient due to delayed repolarization in the epicardial border zone	Inhibition of potassium currents in the border zone
Acute MI Brugada phenocopy in ECG	Delayed repolarisation, as well as a small region of activation failure in the epicardial border zone	Strong inhibitions of sodium, calcium and potassium ionic currents in the border zone
Chronic MI upright tall T-waves in ECG	Large repolarisation time gradient between remote and border zones caused by more severe delay of repolarization in the remote zone	More severe potassium channel suppression in the remote zone
Chronic MI T-wave alternans	Cellular repolarization alternans or early afterdepolarization	Suppressed SERCA and augmented CaMK activity for alternans; Enhanced late sodium current and suppressed hERG current for early

		afterdepolarization
Acute MI reduction in LVEF	Reduced calcium amplitude and/or regional conduction block	Inhibitions of calcium and sodium currents
Chronic MI reduction in LVEF	Reduced calcium amplitude	Decreased SERCA activity

1. Acute MI T-wave inversion and Brugada phenocopy are caused by reversed transmural repolarisation gradient and regional conduction abnormality.

Three distinct types of T-wave morphology were generated by our acute MI biventricular simulations: T-wave inversion, Brugada-phenocopy and normal upright T-wave. We obtained them by applying three types of acute BZs in ventricular simulations, considering both APD prolongation and shortening, as a reflection of the variable experimental results (Mendonca Costa *et al.*, 2018). Collectively, our results highlighted the importance of investigating the implications of the various degrees of experimentally-reported ionic remodelling to explain phenotypic variability of patients with MI. T-wave inversion is a commonly observed feature in the acute MI patients, and is commonly associated with arrhythmic risk (Tikkanen *et al.*, 2015). Here we showed the reversed transmural repolarisation gradient caused by APD prolongation in the epicardial BZ accounted for this phenotype. The link between transmural repolarisation gradient and T-wave polarity has been reported previously (Okada *et al.*, 2011) and is consistent with our results. Brugada-phenocopy was also observed in some acute MI patients (Anselm *et al.*, 2014), and our simulation results showed it could be a reflection of regional conduction abnormality combined with APD prolongation. Although some animal experiments showed acute MI BZ APD shortening (Mendonca Costa *et al.*, 2018), we showed the prolongation of BZ APD was underlying the QT prolongation, T-wave inversion and Brugada phenocopy in the leads

facing the infarct (Supplementary Material Figure S7 showing lead dispersions), which is

consistent with the QTc prolongation observed in the anterior leads of acute anterior

infarction patients (Guaricci *et al.*, 2018). Apart from the above, normal T-waves and QT

intervals were also commonly observed in patients post percutaneous coronary intervention,

which can be reproduced when the post-MI repolarisation dispersion was small between BZ

and NZ (BZ3). It is worth noting that, in addition to having a mild border zone remodelling

as shown in BZ3, a silent ECG signature can also be due to a reduced transmural extent of

the infarct, as has been shown in previous computational studies (Loewe *et al.*, 2018; Wang

et al., 2021). Therefore, T-wave inversion, Brugada phenocopy and QT prolongation occur in

the leads facing the infarct can be useful biomarkers indicating bigger repolarisation

dispersions and/or larger transmural extent in the acute MI.

2. Wide and tall T-wave is explained by large repolarisation dispersions between BZ

and RZ in healed post-MI hearts.

Our simulated chronic ECGs recapitulated the recovery of T-wave polarity observed in

patients after a period of healing. This was achieved through the recovery of the transmural

repolarisation gradient caused by the milder I_{Kr} inhibition in the chronic BZ (Hegyi *et al.*,

2018). Experimental studies in different species showed inconsistent results regarding the

chronic BZ APD (Mendonca Costa *et al.*, 2018). Our chronic BZ remodelling produced

slightly longer APD than the NZ, which is consistent with observations in healed human BZ

(Dangman *et al.*, 1982). However, in minipigs, these remodelling caused shorter BZ APD

than in NZ (Hegyi *et al.*, 2018). This interesting discrepancy between minipigs and human

may be due to the different balance of ionic currents across species, which showed the

benefits of human electrophysiology models in overcoming the inter-species differences.

The two types of T-wave morphologies in the simulated chronic ECGs corresponded

to different extents of RZ APD prolongations, which were commonly observed in healed RZ

of post-MI animals (Hegyi *et al.*, 2018), and in failing human myocytes (Li *et al.*, 2004). The

substantial RZ APD prolongation was reflected as global QT prolongation in all leads, and

the large APD dispersion between chronic BZ and RZ generated wide and tall T-waves in the

precordial leads facing the infarct (Supplementary Material Figure S8 for global and

dispersed ECG characteristics). Previous simulation studies also found the T-wave amplitude

and area were proportional to the dispersion of repolarisation (Arteyeva & Azarov, 2017).

Therefore, these results demonstrated that in patients with global QT prolongation, leads with

bigger T-wave amplitudes could reflect increased local heterogeneity in repolarisation.

3. T wave alternans and severely abnormal ECGs at fast pacing are caused by

alternans and EADs in chronic infarction.

Post-MI ionic remodelling promoted alternans generation, which resulted in T-wave

alternans in simulated ECGs, consistent with the higher incidence of T-wave alternans

reported in post-MI patients (Martin *et al.*, 2009). Two types of repolarisation abnormalities

were observed in our post-MI models: EADs and alternans. One crucial mechanism

promoting alternans behaviour at the cellular level is the increased activity of CaMKII,

observed in the acute MI BZ (Hund *et al.*, 2008*b*), as well as in the hypertrophied and failing

myocardium (Anderson *et al.*, 2011). Enhanced CaMKII phosphorylation may preserve the

contractility of the heart through the phosphorylation of phospholamban and the L-type

calcium channels, but increased RyR phosphorylation by CaMKII resulted in prolonged RyR

opening as well as the enhancement of spontaneous calcium sparks, which can contribute to

alternans and triggered arrhythmias (Maier & Bers, 2007).

Generation of EADs due to chronic post-MI ionic remodelling, such as I_{Kr} inhibition

and I_{NaL} enhancement, was consistent with previous studies (Coppini *et al.*, 2013). EADs in

the RZ can create large repolarisation dispersion in the ventricle, facilitating abnormal

electrotonic wave propagations. Similar re-entrant waves caused by electrotonic gradients

were also observed in previous studies of acute ischemia (Ridley *et al.*, 1992; Dutta *et al.*,

2016; Boukens *et al.*, 2021).

4. LVEF should be combined with QT and T-wave characteristics for arrhythmic risk stratification.

In this study, we observe non-structurally-induced reductions of LVEF in the both the acute

and the chronic post-MI stages. At both stages, ventricles with different extents of

repolarisation dispersion may have similar LVEF because they have similar degrees of

calcium reduction (acute MI T-wave inversion vs normal ST-T, and chronic MI two cases).

Models with inducibility of T-wave alternans and arrhythmia at fast pacing rates may present

with a preserved LVEF at resting heart rates (Figure 6-7). Our cellular level results also

showed models with inducibility of repolarisation abnormalities, such as alternans and

EADs, tended to have more preserved CaT magnitudes at rest rates. Therefore, these

phenomena all support the fact that preserved LVEF measured at rest does not guarantee low

arrhythmic risk.

A recent clinical study of post-MI patients with preserved LVEF showed defibrillators

are needed in those patients with electrophysiological risk factors, such as prolonged QTc,

increased T-wave alternans, to prevent sudden cardiac death (Gatzoulis *et al.*, 2019).

Consistently, we also found post-MI alternans and EADs can present as alternans of T-wave

morphology and prolonged QT intervals. In addition to the global ECG changes, our

simulation results also showed increased local repolarisation dispersion can be reflected in

the leads facing the infarct: inverted T-wave and prolonged QT in the acute MI, and wide and

tall T-wave in the chronic MI. Therefore, we suggest the consideration of these signs as

markers of high arrhythmic risk.

5. Limitations

The main goal of this study is to investigate phenotypic variability in ECG and LVEF

biomarkers arising from post-MI ionic remodelling. We therefore kept other sources of

variability constant, and specifically heart anatomy, location and timing of the early activates

sites, calcium sensitivity, and conduction velocity. These could all modulate quantitatively

the findings but we do not anticipate strong implications in the findings. LVEF reduction in

clinical cases are more significant than in our simulations due to factors other than ionic

remodelling: RZ structural remodelling and elevated myocardium stiffness, and

abnormalities in anatomy. However, these effects have been well-documented elsewhere and

this study serves to elucidate the non-structural mechanisms that underpin LVEF reduction

that is linked to electrophysiological remodelling and arrhythmic risk. The basal plane in our

simulation was fixed in space, which was necessary due to the segmented geometry from

clinical MRI and to prevent unphysiological motion at the truncated basal plane. Despite this

limitation, our conclusions regarding the relative comparisons of mechanical dysfunction are likely to still hold.

In addition, post-MI ionic remodelling can be modulated by other acute and chronic factors such as autonomic modulation (beta-adrenergic effects), inflammation, cell death, and metabolic remodelling, which can be explored in future work. The limitations in this study call for a need for personalised digital twins to be generated in the future to facilitate a better understanding of the interaction between structural remodelling and electrophysiological alterations.

6. CONCLUSIONS

Human-based electromechanical simulations reveal ionic mechanisms underlying T-wave inversion, Brugada phenocopy, and upright T-wave in acute post-MI, as well as the upright T-wave, QT prolongation and T-wave alternans in the healed chronic MI. In acute MI, while the potassium current reduction in the border zone was implicated for all phenotypes, the more severe ECG abnormalities in the Brugada phenocopy implicates additional remodelling for the sodium and calcium currents, which were also key factors in reduced mechanical function. In chronic MI, the degree of QT prolongation and the generation of pro-arrhythmic injury currents were directly related to the severity of potassium current remodelling in the remote region. In addition, late sodium current remodelling was also an important factor underpinning T-wave alternans in chronic MI. Our results show that T-wave inversion, wide and tall T-wave, and QT prolongation in the leads facing the infarct are indicative of local dispersion of repolarisation, which is independent from the reduction of LVEF. Our simulation results suggest the utilisation of T-wave morphology, T-wave alternans and QT

1   prolongation to improve risk stratification biomarkers even when the resting LVEF is

2   preserved.

REFERENCES

Aggarwal Rajesh & Boyden Penelope A. (1995). Diminished Ca2+ and Ba2+ Currents in Myocytes Surviving in the Epicardial Border Zone of the 5-Day Infarcted Canine Heart. *Circulation Research* **77,** 1180–1191.

Ahnve S (1985). QT interval prolongation in acute myocardial infarction. *Eur Heart J* **6 Suppl D,** 85–95.

Al-Khatib SM, Stevenson WG, Ackerman MJ, Bryant WJ, Callans DJ, Curtis AB, Deal BJ, Dickfeld T, Field ME, Fonarow GC, Gillis AM, Granger CB, Hammill SC, Hlatky MA, Joglar JA, Kay GN, Matlock DD, Myerburg RJ & Page RL (2018). 2017 AHA/ACC/HRS Guideline for Management of Patients With Ventricular Arrhythmias and the Prevention of Sudden Cardiac Death: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *Circulation* **138,** e272–e391.

Anderson ME, Brown JH & Bers DM (2011). CaMKII in myocardial hypertrophy and heart failure. *J Mol Cell Cardiol* **51,** 468–473.

Anselm DD, Evans JM & Baranchuk A (2014). Brugada phenocopy: A new electrocardiogram phenomenon. *World J Cardiol* **6,** 81–86.

Arevalo HJ, Vadakkumpadan F, Guallar E, Jebb A, Malamas P, Wu KC & Trayanova NA (2016). Arrhythmia risk stratification of patients after myocardial infarction using personalized heart models. *Nat Commun*; DOI: 10.1038/ncomms11437.

Arteyeva NV & Azarov JE (2017). Effect of action potential duration on Tpeak-Tend interval, T-wave area and T-wave amplitude as indices of dispersion of repolarization: Theoretical and simulation study in the rabbit heart. *J Electrocardiol* **50,** 919–924.

Bloomfield DM, Steinman RC, Namerow PB, Parides M, Davidenko J, Kaufman ES, Shinn T, Curtis A, Fontaine J, Holmes D, Russo A, Tang C & Bigger JT (2004). Microvolt T-wave alternans distinguishes between patients likely and patients not likely to benefit from implanted cardiac defibrillator therapy: a solution to the Multicenter Automatic Defibrillator Implantation Trial (MADIT) II conundrum. *Circulation* **110,** 1885–1889.

Boukens BJD, Dacey M, Meijborg VMF, Janse MJ, Hadaya J, Hanna P, Swid MA, Opthof T, Ardell JL, Shivkumar K & Coronel R (2021). Mechanism of ventricular premature beats elicited by left stellate ganglion stimulation during acute ischaemia of the anterior left ventricle. *Cardiovasc Res* **117,** 2083–2091.

Caldwell BJ, Trew ML, Sands GB, Hooks DA, LeGrice IJ & Smaill BH (2009). Three distinct directions of intramural activation reveal nonuniform side-to-side electrical coupling of ventricular myocytes. *Circ Arrhythm Electrophysiol* **2,** 433–440.

Cluitmans MJM et al. (2021). Noninvasive detection of spatiotemporal activation-repolarization interactions that prime idiopathic ventricular fibrillation. *Science Translational Medicine* **13,** eabi9317.

Coppini R, Ferrantini C, Yao L, Fan P, Del Lungo M, Stillitano F, Sartiani L, Tosi B, Suffredini S, Tesi C, Yacoub M, Olivotto I, Belardinelli L, Poggesi C, Cerbai E & Mugelli A (2013). Late sodium current inhibition reverses electromechanical dysfunction in human hypertrophic cardiomyopathy. *Circulation* **127,** 575–584.

Dangman KH, Danilo P, Hordof AJ, Mary-Rabine L, Reder RF & Rosen MR (1982). Electrophysiologic characteristics of human ventricular and Purkinje fibers. *Circulation* **65,** 362–368.

Dutta S, Mincholé A, Zacur E, Quinn TA, Taggart P & Rodriguez B (2016). Early afterdepolarizations promote transmural reentry in ischemic human ventricles with reduced repolarization reserve. *Prog Biophys Mol Biol* **120,** 236–248.

Elayi CS, Charnigo RJ, Heron PM, Lee BK & Olgin JE (2017). Primary Prevention of Sudden Cardiac Death Early Post-
Myocardial Infarction: Root Cause Analysis for Implantable Cardioverter-Defibrillator Failure and Currently
Available Options. *Circ Arrhythm Electrophysiol*; DOI: 10.1161/CIRCEP.117.005194.

Gallagher KP, Gerren RA, Stirling MC, Choy M, Dysko RC, McManimon SP & Dunham WR (1986). The distribution of
functional impairment across the lateral border of acutely ischemic myocardium. *Circulation Research* **58,**
570–583.

Gatzoulis KA et al. (2019). Arrhythmic risk stratification in post-myocardial infarction patients with preserved
ejection fraction: the PRESERVE EF study. *Eur Heart J* **40,** 2940–2949.

Guaricci AI, Carità P, Lorenzoni V, Casavecchia G, Rabbat M, Ieva R, Brunetti ND, Andreini D, Di Biase M, Marenzi G,
Bartorelli A, Pepi M & Pontone G (2018). QT-interval evaluation in primary percutaneous coronary
intervention of ST-segment elevation myocardial infarction for prediction of myocardial salvage index.
PLoS One; DOI: 10.1371/journal.pone.0192220.

Hegyi B et al. (2018). Complex electrophysiological remodeling in postinfarction ischemic heart failure. *Proc Natl*
Acad Sci USA **115,** E3036–E3044.

Hund TJ, Decker KF, Kanter E, Mohler PJ, Boyden PA, Schuessler RB, Yamada KA & Rudy Y (2008*a*). Role of activated
CaMKII in abnormal calcium homeostasis and I(Na) remodeling after myocardial infarction: insights from
mathematical modeling. *J Mol Cell Cardiol* **45,** 420–428.

Hund TJ, Decker KF, Kanter E, Mohler PJ, Boyden PA, Schuessler RB, Yamada KA & Rudy Y (2008*b*). Role of activated
CaMKII in abnormal calcium homeostasis and I(Na) remodeling after myocardial infarction: insights from
mathematical modeling. *J Mol Cell Cardiol* **45,** 420–428.

Ikeda T, Sakata T, Takami M, Kondo N, Tezuka N, Nakae T, Noro M, Enjoji Y, Abe R, Sugi K & Yamaguchi T (2000).
Combined assessment of T-wave alternans and late potentials used to predict arrhythmic events after
myocardial infarction. A prospective study. *J Am Coll Cardiol* **35,** 722–730.

Jiang M, Cabo C, Yao J, Boyden PA & Tseng G (2000). Delayed rectifier K currents have reduced amplitudes and
altered kinetics in myocytes from infarcted canine ventricle. *Cardiovasc Res* **48,** 34–43.

Land S, Park-Holohan S-J, Smith NP, dos Remedios CG, Kentish JC & Niederer SA (2017). A model of cardiac
contraction based on novel measurements of tension development in human cardiomyocytes. *Journal of*
Molecular and Cellular Cardiology **106,** 68–83.

Levrero-Florencio F, Margara F, Zacur E, Bueno-Orovio A, Wang ZJ, Santiago A, Aguado-Sierra J, Houzeaux G, Grau
V, Kay D, Vázquez M, Ruiz-Baier R & Rodriguez B (2020). Sensitivity analysis of a strongly-coupled human-
based electromechanical cardiac model: Effect of mechanical parameters on physiologically relevant
biomarkers. *Computer Methods in Applied Mechanics and Engineering* **361,** 112762.

Li G-R, Lau C-P, Leung T-K & Nattel S (2004). Ionic current abnormalities associated with prolonged action
potentials in cardiomyocytes from diseased human right ventricles. *Heart Rhythm* **1,** 460–468.

Livshitz LM & Rudy Y (2007). Regulation of Ca2+ and electrical alternans in cardiac myocytes: role of CAMKII and
repolarizing currents. *Am J Physiol Heart Circ Physiol* **292,** H2854-2866.

Loewe A, Wülfers EM & Seemann G (2018). Cardiac ischemia—insights from computational models. *Herzschr*
Elektrophys **29,** 48–56.

Maier LS & Bers DM (2007). Role of Ca2+/calmodulin-dependent protein kinase (CaMK) in excitation-contraction
coupling in the heart. *Cardiovasc Res* **73,** 631–640.

Margara F, Wang ZJ, Levrero-Florencio F, Santiago A, Vázquez M, Bueno-Orovio A & Rodriguez B (2021). In-silico human electro-mechanical ventricular modelling and simulation for drug-induced pro-arrhythmia and inotropic risk assessment. *Progress in Biophysics and Molecular Biology* **159,** 58–74.

Martin DT, Shoraki A, Nesto RW & Rutter MK (2009). Influence of diabetes and/or myocardial infarction on prevalence of abnormal T-wave alternans. *Ann Noninvasive Electrocardiol* **14,** 355–359.

Mendonca Costa C, Plank G, Rinaldi CA, Niederer SA & Bishop MJ (2018). Modeling the Electrophysiological Properties of the Infarct Border Zone. *Front Physiol*; DOI: 10.3389/fphys.2018.00356.

Mincholé A, Zacur E, Ariga R, Grau V & Rodriguez B (2019). MRI-Based Computational Torso/Biventricular Multiscale Models to Investigate the Impact of Anatomical Variability on the ECG QRS Complex. *Front Physiol* **10,** 1103.

Oikarinen L, Viitasalo M & Toivonen L (1998). Dispersions of the QT interval in postmyocardial infarction patients presenting with ventricular tachycardia or with ventricular fibrillation. *Am J Cardiol* **81,** 694–697.

Okada J, Washio T, Maehara A, Momomura S, Sugiura S & Hisada T (2011). Transmural and apicobasal gradients in repolarization contribute to T-wave genesis in human surface ECG. *American Journal of Physiology-Heart and Circulatory Physiology* **301,** H200–H208.

Pogwizd SM, Hoyt RH, Saffitz JE, Corr PB, Cox JL & Cain ME (1992). Reentrant and focal mechanisms underlying ventricular tachycardia in the human heart. *Circulation* **86,** 1872–1887.

Ridley PD, Yacoub MH & Curtis MJ (1992). A modified model of global ischaemia: application to the study of syncytial mechanisms of arrhythmogenesis. *Cardiovasc Res* **26,** 309–315.

Sack KL, Davies NH, Guccione JM & Franz T (2016). Personalised computational cardiology: Patient-specific modelling in cardiac mechanics and biomaterial injection therapies for myocardial infarction. *Heart Fail Rev* **21,** 815–826.

Santiago A, Aguado-Sierra J, Zavala-Aké M, Doste-Beltran R, Gómez S, Arís R, Cajas JC, Casoni E & Vázquez M (2018). Fully coupled fluid-electro-mechanical model of the human heart for supercomputers. *Int J Numer Method Biomed Eng* **34,** e3140.

Sato A, Hiroe M, Nozato T, Hikita H, Ito Y, Ohigashi H, Tamura M, Takahashi A, Isobe M & Aonuma K (2008). Early validation study of 64-slice multidetector computed tomography for the assessment of myocardial viability and the prediction of left ventricular remodelling after acute myocardial infarction. *Eur Heart J* **29,** 490–498.

Shenthar J, Deora S, Rai M & Nanjappa Manjunath C (2015). Prolonged Tpeak-end and Tpeak-end/QT ratio as predictors of malignant ventricular arrhythmias in the acute phase of ST-segment elevation myocardial infarction: a prospective case-control study. *Heart Rhythm* **12,** 484–489.

Solomon SD, Zelenkofske S, McMurray JJV, Finn PV, Velazquez E, Ertl G, Harsanyi A, Rouleau JL, Maggioni A, Kober L, White H, Van de Werf F, Pieper K, Califf RM, Pfeffer MA, & Valsartan in Acute Myocardial Infarction Trial (VALIANT) Investigators (2005). Sudden death in patients with myocardial infarction and left ventricular dysfunction, heart failure, or both. *N Engl J Med* **352,** 2581–2588.

Spargias KS, Lindsay SJ, Kawar GI, Greenwood DC, Cowan JC, Ball SG & Hall AS (1999). QT dispersion as a predictor of long-term mortality in patients with acute myocardial infarction and clinical evidence of heart failure. *Eur Heart J* **20,** 1158–1165.

Stone PH, Raabe DS, Jaffe AS, Gustafson N, Muller JE, Turi ZG, Rutherford JD, Poole WK, Passamani E & Willerson JT (1988). Prognostic significance of location and type of myocardial infarction: independent adverse outcome associated with anterior location. *J Am Coll Cardiol* **11,** 453–463.

Streeter DD, Vaishnav RN, Patel DJ, Spotnitz HM, Ross J & Sonnenblick EH (1970). Stress distribution in the canine left ventricle during diastole and systole. *Biophys J* **10,** 345–363.

Sun K, Stander N, Jhun C-S, Zhang Z, Suzuki T, Wang G-Y, Saeed M, Wallace AW, Tseng EE, Baker AJ, Saloner D, Einstein DR, Ratcliffe MB & Guccione JM (2009). A computationally efficient formal optimization of regional myocardial contractility in a sheep with left ventricular aneurysm. *J Biomech Eng* **131,** 111001.

Tikkanen JT, Kenttä T, Porthan K, Huikuri HV & Junttila MJ (2015). Electrocardiographic T Wave Abnormalities and the Risk of Sudden Cardiac Death: The Finnish Perspective. *Ann Noninvasive Electrocardiol* **20,** 526–533.

Tomek J, Bueno-Orovio A, Passini E, Zhou X, Minchole A, Britton O, Bartolucci C, Severi S, Shrier A, Virag L, Varro A & Rodriguez B (2019). Development, calibration, and validation of a novel human ventricular myocyte model in health, disease, and drug block ed. Faraldo-Gómez JD, Barkai N, Hund T & Maleckar M. *eLife* **8,** e48890.

Tomek J, Tomková M, Zhou X, Bub G & Rodriguez B (2018). Modulation of Cardiac Alternans by Altered Sarcoplasmic Reticulum Calcium Release: A Simulation Study. *Front Physiol*; DOI: 10.3389/fphys.2018.01306.

Ursell PC, Gardner PI, Albala A, Fenoglio JJ & Wit AL (1985). Structural and electrophysiological changes in the epicardial border zone of canine myocardial infarcts during infarct healing. *Circ Res* **56,** 436–451.

Vaduganathan M, Claggett BL, Chatterjee NA, Anand IS, Sweitzer NK, Fang JC, O'Meara E, Shah SJ, Hegde SM, Desai AS, Lewis EF, Rouleau J, Pitt B, Pfeffer MA & Solomon SD (2018). Sudden Death in Heart Failure With Preserved Ejection Fraction: A Competing Risks Analysis From the TOPCAT Trial. *JACC Heart Fail* **6,** 653–661.

Van Leuven SL, Waldman LK, McCulloch AD & Covell JW (1994). Gradients of epicardial strain across the perfusion boundary during acute myocardial ischemia. *American Journal of Physiology-Heart and Circulatory Physiology* **267,** H2348–H2362.

Vazquez R, Bayes-Genis A, Cygankiewicz I, Pascual-Figal D, Grigorian-Shamagian L, Pavon R, Gonzalez-Juanatey JR, Cubero JM, Pastor L, Ordonez-Llanos J, Cinca J, de Luna AB, & MUSIC Investigators (2009). The MUSIC Risk score: a simple method for predicting mortality in ambulatory patients with chronic heart failure. *Eur Heart J* **30,** 1088–1096.

Waks JW, Sitlani CM, Soliman EZ, Kabir M, Ghafoori E, Biggs ML, Henrikson CA, Sotoodehnia N, Biering-Sørensen T, Agarwal SK, Siscovick DS, Post WS, Solomon SD, Buxton AE, Josephson ME & Tereshchenko LG (2016). Global Electric Heterogeneity Risk Score for Prediction of Sudden Cardiac Death in the General Population. *Circulation* **133,** 2222–2234.

Wang ZJ, Santiago A, Zhou X, Wang L, Margara F, Levrero-Florencio F, Das A, Kelly C, Dall'Armellina E, Vazquez M & Rodriguez B (2021). Human biventricular electromechanical simulations on the progression of electrocardiographic and mechanical abnormalities in post-myocardial infarction. *Europace* **23,** i143–i152.

Yao J-A, Hussain W, Patel P, Peters NS, Boyden PA & Wit AL (2003). Remodeling of gap junctional channel function in epicardial border zone of healing canine infarcts. *Circ Res* **92,** 437–443.

Zhou X, Bueno-Orovio A, Orini M, Hanson B, Hayward M, Taggart P, Lambiase PD, Burrage K & Rodriguez B (2016). In Vivo and In Silico Investigation Into Mechanisms of Frequency Dependence of Repolarization Alternans in Human Ventricular Cardiomyocytes. *Circ Res* **118,** 266–278.

1 Zhou X, Qu Y, Passini E, Bueno-Orovio A, Liu Y, Vargas HM & Rodriguez B (2019). Blinded In Silico Drug Trial Reveals
2   the Minimum Set of Ion Channels for Torsades de Pointes Risk Assessment. *Front Pharmacol* **10,** 1643.

ADDITIONAL INFORMATION SECTION

Data availability statement

The simulation input files and Alya executable required to replicate the simulated results will

be made available upon request. Python and MATLAB scripts for cellular populations of

models simulations and post-processing of the biventricular simulations can be found at:

https://github.com/jennyhelyanwe/post_MI_postprocessing

Competing interests

The authors declare that they have no competing interests.

Funding

This work was funded in whole, or in part, by the Wellcome Trust (214290/Z/18/Z). For the

purpose of Open Access, the author has applied a CC BY public copyright licence to any

Author Accepted Manuscript version arising from this submission.

This work was supported by a Wellcome Trust Fellowship in Basic Biomedical Sciences to

B.R. (214290/Z/18/Z), an Oxford-Bristol Myers Squibb Fellowship to X.Z.

(R39207/CN063), the Personalised In-Silico Cardiology (PIC) project, the CompBioMed 1

and 2 Centre of Excellence in Computational Biomedicine (European Commission Horizon

2020 research and innovation programme, grant agreements No. 675451 and No. 823712), an

NC3Rs Infrastructure for Impact Award (NC/P001076/ 1), the TransQST project (Innovative

Medicines Initiative 2 Joint Undertaking under grant agreement No 116030, receiving

support from the European Union's Horizon 2020 research and innovation programme and

EFPIA), and the Oxford BHF Centre of Research Excellence (RE/13/1/30181), PRACE-ICEI

funding projects icp005, icp013, icp019.

Acknowledgements

The authors would like to acknowledge Dr Erica Dall'Armellina, Dr Arka Das, Dr Chris

Kelly, and Dr Lei Wang, for discussions on the study.

Preprint

This manuscript was first published as a preprint: Xin Zhou, Zhinuo Jenny Wang, Julia Camps,

Jakub Tomek, Alfonso Santiago, Adria Quintanas, Mariano Vazquez, Marmar Vaseghi, Blanca

Rodriguez (2022). [Clinical phenotypes in acute and chronic infarction explained through human

ventricular electromechanical modelling and simulations]. bioRxiv.

https://www.biorxiv.org/content/10.1101/2022.02.15.480392v2