Clinical phenotypes in acute and chronic infarction explained through human ventricular electromechanical modelling and simulations

Xin Zhou*[1], Zhinuo Jenny Wang*[1], Julia Camps[1], Jakub Tomek[2], Alfonso Santiago[3,4],

Adria Quintanas[3], Mariano Vazquez[3,4], Marmar Vaseghi[5,6], Blanca Rodriguez[1]

* co-first authors

1 Department of Computer Science, University of Oxford, UK

2 Department of Physiology, Anatomy and Genetics, University of Oxford, UK

3 Department of Computer Applications in Science and Engineering, Barcelona Supercomputing

Centre (BSC), Barcelona, Spain

4 ELEM Biotech, Barcelona, Spain

5 UCLA Cardiac Arrhythmia Center, University of California, Los Angeles, CA, USA

6 Neurocardiology Research Center of Excellence, University of California, Los Angeles, CA, USA

Author contact details:

Dr Xin Zhou: xin.zhou@cs.ox.ac.uk

Dr Zhinuo Jenny Wang: jenny.wang@cs.ox.ac.uk

Professor Blanca Rodriguez: blanca.rodriguez@cs.ox.ac.uk

Oxford address: Department of Computer Science, Wolfson Building, Parks Road, Oxford OX1

3QD, UK

Word count: 7913 excluding references only, 7589 excluding references and the abstract,

10,110 in total (excluding title page).

ABSTRACT

Aims:

Sudden death after myocardial infarction (MI) is associated with electrophysiological

heterogeneities and ionic remodelling, which are reflected as variable phenotypes. Low

ejection fraction (EF) is used in risk stratification, but its mechanistic links with the post-MI

pro-arrhythmic heterogeneities are unknown. We aim to provide a mechanistic explanation

of clinical phenotypes in acute and chronic MI, from ionic remodeling to ECG and EF, using

human electromechanical modelling and simulation to augment experimental and clinical

investigations.

Methods and Results:

A human ventricular electromechanical modelling and simulation framework is constructed

and validated with rich experimental and clinical datasets. Abnormalities caused by scar and

border zone ionic remodeling are introduced in varying degrees as reported in experimental

data obtained in acute and chronic infarction. Simulations enabled reproducing and

explaining clinical phenotypes post-MI, from ionic remodelling to ECGs and pressure-

volume loops. In acute MI, T-wave inversion and Brugada phenocopy were explained by up

to 57 ms of local APD prolongation and activation failure due to the inhibition of potassium,

sodium and calcium channels in the border zone. In chronic MI, upright tall T-waves

highlight large repolarisation dispersion caused by uneven potassium channel expression in

border and remote zones, which promoted ectopic propagation at fast pacing. Post-MI ionic

remodelling reduced EF by up to 10% through inhibition of calcium transient amplitude due

to weaker calcium currents or SERCA activity, but the EF at resting heart rate was not

sensitive to the extent of repolarisation heterogeneity and the risk of repolarisation

abnormalities at fast pacing.

Conclusions:

Multi-scale modelling and simulation coherently integrates experimental and clinical data at

subcellular, tissue, and organ scales to unravel electromechanical disease mechanisms in MI.

In acute post-MI, ionic remodelling and its effect on refractoriness and propagation failure in

the BZ have a strong impact on phenotypic ECG variability, whereas in chronic post-MI, the

repolarisation dispersion across the BZ is crucial. T-wave and QT abnormalities are better

indicators of repolarisation heterogeneities than EF in post-MI.

57
58
59

INTRODUCTION

Sudden cardiac death in post-myocardial infarction (post-MI) patients is due to lethal arrhythmias in 50% of cases at both acute and chronic infarct stages (Vazquez *et al.*, 2009; Elayi *et al.*, 2017). Risk stratification is currently based on low left ventricular ejection fraction (LVEF) (Solomon *et al.*, 2005) to identify patients who need the implantation of defibrillator device. However, only a very small subset of patients that suffer from sudden cardiac death are identified by low LVEF, and a significant number of sudden deaths occur in patients with relatively preserved LVEF (Vaduganathan *et al.*, 2018). The mechanistic link between electrophysiological heterogeneities, ECG and LVEF phenotypes that underpin arrhythmic events is not clear. A precision medicine approach using computational modelling and simulations could help to elucidate disease mechanisms and provide an *in silico* alternative for therapy evaluation and risk stratification. However, a key hurdle in clinical adoption of such *in silico* tools is providing proof of model credibility through validation studies. In this study, we set out to achieve the dual aim of demonstrating a multi-scale approach for model validation and to identify important electrophysiological mechanisms for post-MI risk stratifications.

Various electrocardiogram (ECG) characteristics were suggested by clinical studies to be relevant to the arrhythmic risk of post-MI patients. Longer QT intervals have been associated with increased mortality as well as ventricular tachycardia and fibrillation for both the acute and chronic MI (Ahnve, 1985; Oikarinen *et al.*, 1998). However, QT prolongation in single leads can be a reflection of reduced global repolarisation reserve, while the regional heterogeneity of repolarisation in different post-MI zones can be more crucial (Cluitmans *et al.*, 2021) than the global reserve for the development of re-entrant waves. QT dispersion

between 12-lead ECGs was proposed as a potential marker for electrophysiological

heterogeneity (Spargias *et al.*, 1999); however, it was found to have good specificity but low

sensitivity (Spargias *et al.*, 1999). Prolonged T-peak to T-end interval (Tpe) (Shenthar *et al.*,

2015) and increased incidence of T-wave alternans (Bloomfield *et al.*, 2004) were also found

to be useful risk predictors in post-MI patients. Other ECG metrics, such as QRST integral

and spatial ventricular gradients have also shown some promise in improving SCD prediction

(Waks *et al.*, 2016). However, despite the promising outcomes of those clinical studies, the

ECG-based risk predictors are not widely applied in the clinical evaluation of the need for

implantable cardioverter-defibrillators (Al-Khatib *et al.*, 2018).

A key factor that hinders the clinical utility of ECG biomarkers is the large variability

in post-MI phenotypes, their progression from acute to chronic, and between different

patients. Variability in QT prolongation (Ahnve, 1985) and post-MI ECG morphologies

constrain the use of single biomarker thresholds as predictors. Furthermore, an important

limitation of non-invasive ECG biomarkers is their inability to provide direct measurements

of regional electrophysiological heterogeneity caused by scar and ionic remodelling, which is

critical for the arrhythmic substrate.

Many experimental studies have investigated the underlying mechanisms of increased

repolarisation dispersion in post-MI patients. Specifically, repolarisation dispersion increases

after MI due to ionic differences between the border zone (BZ) surrounding the scar and the

remote zone (RZ) myocardium (Mendonca Costa *et al.*, 2018). The BZ at the acute MI

exhibits reduced sodium current (I_{Na}), L-type calcium current (I_{CaL}) (Aggarwal Rajesh &

Boyden Penelope A., 1995), and rapid and slow delayed rectifier potassium currents (I_{Kr} and

I_{Ks}) (Jiang *et al.*, 2000), as well as enhanced CaMKII activity (Hund *et al.*, 2008*a*) and gap

junction redistribution (Yao *et al.*, 2003). After a period of scar healing, ionic remodelling

may partially recover (Ursell *et al.*, 1985), but increased late sodium current (I_{NaL}), calcium-

activated potassium and chloride currents (I_{KCa} and I_{ClCa}) were observed in the chronic BZ

and RZ (Hegyi *et al.*, 2018). Post-MI ionic remodelling can cause conduction abnormalities,

repolarisation abnormalities, as well as variable alterations in the action potential duration

(APD), which act as substrates of ventricular tachycardia (Pogwizd *et al.*, 1992). It is,

however, unclear how various degrees of repolarisation dispersion affect ECG biomarkers

and LVEF used for risk stratification. Therefore, further studies are needed to bridge the gap

between cellular electrophysiological characteristics and variable patient phenotypes post-

MI. Furthermore, the wealth of cellular, tissue, and ECG data described in the literature

provides an excellent multi-scale dataset for model validation. Comparisons of simulated

action potential and ECG biomarkers with experimental and clinical data under acute and

chronic MI conditions help to improve the credibility of the computational model.

Human-based modelling and simulations of ventricular electrophysiology and

mechanics have demonstrated accurate prediction of post-MI arrhythmic risk and myocardial

stress and strain (Arevalo *et al.*, 2016; Sack *et al.*, 2016). However, previous work

concentrated on either electrophysiology or mechanics, while the crosstalk between the two,

and the relationship between LVEF and arrhythmic risk, is yet to be investigated.

The main goal of this study is, therefore, to quantify the contribution of varying

degrees of ionic remodelling to the phenotypic variability in ECG and LVEF biomarkers

observed in acute and chronic post-MI patients, and thereby create and validate models of

post-MI electromechanics. Using state-of-the-art electromechanical human biventricular

simulations, we aim to identify biomarkers that are most representative to the pro-arrhythmic

substrate for each state, thus facilitating post-MI risk stratifications to go beyond LVEF. We

hypothesise that different ECG abnormalities are explained by different degrees of ionic

remodelling leading to activation sequence abnormalities, dispersion of repolarisation, early

after depolarisations (EADs) and alternans, whereas LVEF is insensitive to ionic remodelling

that underpins ECG disease markers and reflects predominantly calcium transient and

structural abnormalities.

METHODS

1. Human multi-scale ventricular electromechanical modelling and simulation: from

ionic remodelling to ECG and LVEF.

A human ventricular electromechanical modelling and simulation framework is constructed

using a population of models approach and evaluated using experimental and clinical data to

enable the investigations of variable post-MI patient phenotypes, from ionic remodelling to

body surface ECGs and pressure-volume (PV) loops (Figure 1A). A cardiac magnetic

resonance (CMR)-based biventricular anatomical mesh (Figure 1B) with corresponding torso

geometry was used for all simulations in this study, with an anterior scar that is 75%

transmural (Wang *et al.*, 2021). Electrical propagation was simulated using the monodomain

equation with orthotropic diffusion based on rule-based fields for fibre directions (Streeter *et*

al., 1970) with sheet directions normal to the endocardial/epicardial surface (Levrero-

Florencio *et al.*, 2020) (Figure 1B). Transmural and apex-to-base heterogeneities (Mincholé

et al., 2019) were introduced and a sensitivity analysis was performed to investigate their

implications in ECG biomarkers and LVEF (Figure 1B). Electrical stimulus via Purkinje-

myocardial junctions was simulated by an endocardial fast-activation layer with root node

locations to achieve realistic QRS complex morphologies simulated at clinically standard

lead locations (Figure 1C) (Mincholé *et al.*, 2019). In healthy tissue (normal zone (NZ)),

monodomain diffusivities were calibrated to achieve experimentally measured orthotropic

conduction velocities of 67 cm/s, 30 cm/s, and 17 cm/s (Caldwell *et al.*, 2009).

Figure 1: Human-based multi-scale modelling and simulation in acute and chronic

myocardial infarction. *(A) Simulations using a population of ventricular models (n=17) to*

produce ECGs (light blue traces) and pressure-volume (PV) loops (grey traces)

superimposed with the baseline ventricular model (ECG in blue and PV in black).

Biomarkers are calculated from the baseline simulation of ECG and PV, as illustrated. (B)

Ventricular electrophysiology is simulated using a fast endocardial activation layer to

approximate Purkinje-myocardial junction, experimentally-informed transmural and apico-

basal heterogeneities in action potential duration, and transmurally varying myocyte

orientation. Mechanical pumping behaviour is modelled by coupling the intraventricular

pressures with a two-element Windkessel model of arterial haemodynamics with a fixed

basal plane. (C) An anterior 75% transmural infarction is modelled with acute and chronic

ionic remodelling embedded in the border zone and remote zones. Standard 12-lead ECG

was evaluated at standard body-surface locations. (D) Simulated action potentials using

populations of human ventricular models in healthy (baseline) and acute and chronic post-

MI conditions with different degrees of ionic remodelling. (E) Schematic representation of

ionic fluxes, calcium dynamics and actin/myosin contraction mechanisms in the human

ventricular electromechanically-coupled cellular model.

Since anterior infarction is very common and related to the worst prognosis (Stone *et*

al., 1988), 75% transmural extent of infarct from the endocardial surface was introduced to

the anterior myocardial wall (Figure 1C) to match the clinical definition of transmural

infarctions (Sato *et al.*, 2008), with scar and border zones (maximum width 0.5 cm) taking up

12.4% and 11.7% of ventricular volume (or 15.8% and 15.0% of left ventricular volume),

respectively. Maximum border zone width was based on various reports of systolic strain

dysfunction in < 1 cm proximity to the infarct (Gallagher *et al.*, 1986; Van Leuven *et al.*,

1994). Electrophysiological remodelling was implemented in the border zone (BZ) and the

chronic remote zone (RZ), which covered the entire non-infarcted and non-BZ region in both

ventricles. In BZ and infarct zones, the diffusivities were calibrated to reproduce conduction

slowing (one-third of the NZ conduction velocities). For each virtual cardiomyocyte, the

electromechanical single cell model considered the human-based ToR-ORd

electrophysiological model (Tomek *et al.*, 2019) (extensively validated in control, disease

and drug block conditions) coupled with human excitation-contraction and active tension

Land model (Land *et al.*, 2017; Levrero-Florencio *et al.*, 2020) (Figure 1E).

The human biventricular model incorporated strongly-coupled electromechanics with

orthotropic passive mechanical behaviour and balance of linear momentum with inertial

effects, as in our previous work (Levrero-Florencio *et al.*, 2020; Margara *et al.*, 2021; Wang

et al., 2021). In brief, firstly, intracellular calcium concentration drives crossbridge cycling

and force production through unblocking the crossbridge binding-site. Secondly, calcium

sensitivity and force production are a function of fibre stretch ratio. Thirdly, stretch rate

affects distortion-dependent cross-bridge unbinding (Land *et al.*, 2017) and the diffusivity

tensor is transformed using the deformation gradient tensor such that the prescribed values

describe conductivities in the deformed state (Levrero-Florencio *et al.*, 2020).

An elastic spring boundary condition was set to act perpendicularly to the epicardial

surface, to simulate pericardial constraint, and the basal plane was fixed in space to prevent

unphysiological tilting and expansion due to the unavailability of closed basal geometry.

Pressure boundary condition on the left and right endocardial surfaces were controlled using

a series of five equations (see SM2) that controls 1) active diastolic inflation followed by

electrical activation and 2) isovolumic contraction, 3) ejection coupled with two-element

Windkessel aortic haemodynamics model, 4) isovolumic relaxation, and 5) passive inflation

and relaxation (Wang *et al.*, 2021).

The sheet active tension was set to 30% of fibre active tension to achieve sufficient

LVEF in control conditions, based on the results from a sensitivity analysis in Supplementary

Material (SM) section SM5. In brief, we evaluated the sensitivity of ECG morphology and

LVEF to changes in the calcium sensitivity of troponin binding in excitation-contraction

coupling and the active tension in the sheet direction as a percentage of that in the fibre

direction.

The passive stiffness parameters were calibrated based on a previous sensitivity

analysis (Wang *et al.*, 2021) to achieve 53% LVEF and a physiological pressure-volume

(PV) loop in a control simulation (see SM Table S2 and S3 for a list of calibrated

parameters).

The active tension was set to zero in the scar to represent the myocyte damage, and

the chronic passive stiffness parameters of the infarct region were increased 10-fold to mimic

fibrotic scar formation (Sun *et al.*, 2009). For each acute post-MI phenotype, a case of

complete loss of contractile function (zero active tension) in the BZ was also simulated to

evaluate the contribution of other non-ionic remodelling-related abnormalities on ejection

fraction.

2. **Experimentally-informed single cell and ventricular populations of human post-MI**

   electromechanical models.

To account for the inter-subject electrophysiological variability widely observed in clinical

data, the baseline human cellular electromechanical ToR-Land model was extended to

populations of healthy cellular models. Then, post-MI ionic remodelling was applied to

generate populations of post-MI virtual cardiomyocytes (Figure 1D). In addition to the

baseline ToR-ORd model, several representative cellular models were selected from the

population and implemented into the biventricular electromechanical simulations.

An initial population of human ventricular cell models was constructed based on the

ToR-ORd model by varying the conductances or magnitudes of I_{Na}, I_{NaL}, I_{to}, I_{CaL}, I_{Kr}, I_{Ks}, I_{K1},

I_{NaCa}, I_{NaK}, J_{rel} and J_{up} by up to ±50% using Latin Hypercube Sampling (SM Figure S1 and

S2). As illustrated in previous studies, small populations of models with proarrhythmic ionic

remodelling achieved similar predictability as large populations with uniform current

variations (Zhou *et al.*, 2019). Therefore, we chose to start with a small healthy population

(n=500) and then introduce multiple combinations of ionic remodelling to mimic the large

post-MI variability. After calibration with human experimental data (SM Table S1) and

discarding those manifesting EADs at 1Hz, 245 sets of endocardial, midmyocardial and

epicardial models were accepted as the healthy population (SM Figure S2). From this

population, a total of 17 sets of cell models were randomly selected and uniformly embedded

in 17 ventricular models, to generate ventricular population of models that produce a variety

of ECGs (Figure 1A, light blue traces) and PV loops (Figure 1B, grey traces).

Several degrees of post-MI ionic remodelling were collated from a combination of

human and animal experimental data with variability in severity of disease to explore

whether such variabilities can explain variability in established clinical ECG phenotypes.

These remodellings have been applied to the healthy celllular model population (n=245) to

generate BZ and RZ populations for both acute and chronic post-MI. For acute post-MI

(within a week post-occlusion), three types of BZ remodelling (Acute BZ1-3) were

considered based on previous modelling work and experimental canine data collected within

5 days post-MI. The three models of acute border zone remodelling had in common strong

inhibition of I_{Na} (60~62%). The BZ2 model had more severe inhibition of I_{CaL} and I_{Kr} than

the BZ1 model, alongside other minor differences. The BZ3 model, while having less severe

potassium currents inhibition than BZ1, had additional remodellings in CaMKII dynamics,

RyR time constants, and I_{Cab}. For chronic post-MI, ionic remodelling measured from

minipigs 5 months post-MI with heart failure were used to generate Chronic BZ (affecting

only the BZ) and Chronic RZ1 (also affecting the remote myocardium). Another type of

remodelling, Chronic RZ2, was established based on multiple experimental data from failing

human cardiomyocytes. The RZ covers the entire myocardium apart from the infarct and BZ

in chronic MI simulations. Furthermore, reduction of sodium current and SERCA, with

enhanced CaMKII activity and slower calcium release induced by CaMKII activation were

also implemented in Chronic BZ, RZ1 and RZ2, as observed in human failing

cardiomyocytes (details in SM Table S4). Compared with RZ1, the RZ2 model had a

significantly stronger inhibition of potassium currents and a lower repolarisation reserve,

alongside other more minor differences. Both human recordings and animal data were used

for model evaluation, given the scarcity of human tissue, summarised in SM Table S4-S5.

Action potential, calcium transient, and active tension characteristics of the post-MI models

are summarised in SM Table S6. These post-MI ionic remodellings were then applied to the

ventricular population of models (n=17) to explain ECG and PV phenotypes while

considering physiological population variability in baseline ionic conductances. These

remodelled cell models are embedded uniformly within each region according to a prescribed

transmural heterogeneity of 30% endo, 40% mid-myocardial, and 30% epicardial cell types.

3. Simulation protocols and biomarker calculation.

Human virtual ventricular myocytes were paced at 1Hz for 500 beats to detect EAD

generation. For alternans generation, single cells were paced at cycle lengths (CLs) of

500 ms, 400 ms and 300 ms for 500 beats, and a ΔAPD greater than 3 ms between the last

two beats at steady state was defined as alternans.

Biventricular electromechanical simulations were performed at 800 ms CL (75 beats

per minute) with 100 ms allowed for active diastolic filling prior to endocardial activation for

each beat. Three beats were sufficient to achieve converged ECG and PV characteristics at

800 ms CL (see Figure S7 for ECGs for all beats).

For chronic post-MI, an additional fast pacing protocol was applied with 500 ms CL

(120 beats per minute), with 50 ms allowed for active diastolic filling for each beat. From

these fast pacing chronic MI simulations, two sets of cell models showing EAD and alternans

behaviours, respectively, were selected from the population of models and embedded

according to transmural heterogeneity in the remote zone for ventricular simulations, to test

whether the arrhythmic behaviours at the cellular level can result in arrhythmic behaviour at

ventricular level and manifest in the ECG. For this fast pacing protocol, six beats were

necessary to achieve converged ECG and PV characteristics at 500 ms CL (see Figure S8 for

simulated ECGs of all beats).

Clinical biomarkers were quantified from the simulated ECG (including the QT

interval, QRS duration, T-wave duration, T peak to T end duration, T onset to T peak

duration, and QT dispersion, see definition and method of evaluation in SM2 and biomarker

results in SM6 Table S8), and from the simulated PV loop (including end diastolic and end

systolic volumes, peak systolic pressures and LV and RV ejection fractions), as well as wall

thickening strain (see Table S8-S9 for ECG and PV biomarkers for all simulated beats).

4. Simulation software and computational framework.

Cellular electrophysiological simulations and Latin Hypercube Sampling were performed

using bespoke MATLAB codes. Coupled cellular electromechanics, as well as biventricular

electromechanics simulations, were performed using the high-performance numerical

software, Alya, for complex coupled multi-physics and multi-scale problems (Santiago *et al.*,

2018) on the CSCS (Swiss National Supercomputing Centre) Piz Daint supercomputer multi-

core clusters, granted through the PRACE (Partnership for Advanced Computing in Europe)

project. The simulation input files and Alya executable required to replicate the simulated

results are available upon request for scientific investigations.

5. Data availability

The simulation input files and Alya executable required to replicate the simulated results will

be made available upon request. Python and MATLAB scripts for cellular populations of

models simulations and post-processing of the biventricular simulations can be found at:

https://github.com/jennyhelyanwe/post_MI_postprocessing

RESULTS

1. Human modelling and simulation for ECG phenotypes in acute and chronic post-MI.

Figure 2 demonstrates the ability of human electromechanical simulations to reproduce a variety of clinically reported phenotypes in patients with acute and chronic infarction, in agreement with clinical measurements of ECG and pressure-volume biomarkers, as quantified in further detail in (SM Table S7, details of clinical database in SM4). When imposing acute post-MI remodelling, simulated ECGs reproduced fractionated QRS complexes, T-wave inversion, Brugada phenocopy ST-segment elevation and QT interval prolongation in the anterior leads (Figure 2A), which are common ECG phenotypes observed in acute post-MI patients. Simulations also recapitulated ECG morphology similar to healthy subjects with upright T-waves (Figure 2A, right), which can also be present in acute post-MI.

In chronic post-MI, simulated ECG displayed upright T-waves, and global prolongations of QT intervals in all precordial leads (Figure 2B, top row), which are characteristic of chronic patients with worse clinical outcomes (Ahnve, 1985; Oikarinen *et al.*, 1998). A range of T-wave durations were present, as can be found in chronic post-MI (Figure 2B, bottom row). SM4 summarises the clinical ECGs used for comparisons in Figure 2. The full 12-lead ECGs for acute and chronic post-MI simulations can be found in SM Figure S7 and S8.

Simulations using the ventricular population of models showed that the described ECG features of acute and chronic post-MI were mostly preserved across variations in ionic conductances (Figure 2C). Sensitivity analysis showed first that large changes in apex-to-base and transmural heterogeneities only altered T-wave amplitude but not its polarity and did not affect the ST-segment (SM Figure S3 and S4), and second that changes in mechanical parameters did not affect the ECG morphology (SM Figure S5 and S6). This result supports the specificity of post-MI signatures to underlying ionic remodelling.

Figure 2: Agreement between simulated and clinical ECGs demonstrating variability in

clinical phenotypes in acute and chronic post-myocardial infarction (post-MI). (A) In acute

MI, simulated ECGs show T-wave inversion (border zone model 1 (BZ1)), Brugada

phenocopy (BZ2), and normal phenotypes (BZ3), in accordance with phenotypes found in

clinical databases. (B) In chronic MI, simulated ECGs show prolonged QT and upright T-

waves with a range of amplitude and duration (remote zone model 1 and 2 (RZ1, RZ2))

comparable to those observed in clinical databases. (C) ECG simulations of control, and

acute and chronic post-MI considering ionic variability of the baseline ToR-ORd model. T

wave morphologies for acute and chronic post-MI are mostly preserved across ionic

variability.

2. In acute MI, T-wave inversion and Brugada phenocopy can indicate reversed

transmural repolarisation gradient and activation failure.

Analysis of simulation results enabled the uncovering of specific contributions of different

degrees of ionic remodelling to ECG phenotypes identified in acute versus chronic MI

(Figure 3 vs Figure 4). In acute MI, T-wave inversion phenotype was associated with a

reversed transmural repolarisation gradient (Figure 3A, repolarisation time map insets), due

to 57 ms of APD prolongation at the epicardial BZ compared with control (Figure 3B,

membrane potential). APD prolongation originated due to inhibition of multiple potassium

currents caused by BZ1 ionic remodellling (Figure 3C, the first column of I_{Kr}, and SM Table

S4 Acute BZ1).

Brugada phenocopy ECG phenotype was also observed in acute MI with BZ2

remodelling, causing regions of activation failure in the epicardial border zone (Figure 3A,

the activation and repolarisation time maps), as well as delayed repolarisation in the BZ near

the apex, caused by 29 ms of epicardial APD prolongation compared with control (Figure

3B, membrane potential). These were caused by strong inhibitions of sodium, calcium and

potassium ionic currents in the BZ (Figure 3C, the second column of I_{Na}, I_{CaL}, I_{Kr}, and SM

Table S4 Acute BZ2). In this simulation, electrical activation in the infarcted region was

preserved despite the conduction block in the BZ because of higher expressions of the L-type

calcium channel in the mid-myocardium (see SM Figure S11).

Acute MI with upright T-waves corresponded to a comparable transmural

repolarisation gradient as in control with BZ3 ionic remodelling (Figure 3A, the

repolarisation time map). In this case, the lack of potassium current inhibition accounted for

the lack of ECG manifestations (Figure 3C, the third column of I_{Kr}, and SM Table S4 Acute

BZ3).

Ionic bases of ECG abnormalities in acute MI

Figure 3: Multiscale explanation of ST and T-wave phenotypes in acute MI. *(A) Activation time maps reveal conduction delay in acute border zone in T-wave inversion and normal ST-T phenotypes, and conduction block in Brugada phenocopy, as well as large repolarisation dispersion and altered transmural repolarisation gradient in T-wave inversion and Brugada*

phenocopy. Red in activation map show regions of no activation (NA), green in

repolarisation map highlights regions of no repolarisation (NR). (B) Action potential

duration (APD) prolongation is present in T-wave inversion and Brugada phenocopy

cellular phenotypes, as well as slight shortening (normal ST-T) (red arrows), with decreased

calcium amplitude in all phenotypes, and a corresponding decrease in active tension

generation. (C) I_{Na}, I_{CaL}, and I_{Kr} remodelling underpin reduced conduction, reduced calcium

amplitude, and alterations in action potential duration, respectively, in all acute phenotypes.

3. In chronic post-MI, variable T-wave width can be explained by the extent of

repolarisation dispersion between border zone and remote zone.

In chronic MI, global QT prolongation was due to APD prolongation in the remote

myocardium (Figure 4, repolarisation time maps and membrane potentials). Recovery of the

upright T-wave in the anterior leads (compared to acute MI) was due to a recovery of the

transmural repolarisation gradient (Figure 4A, repolarisation time maps, and SM Figure

S10), given the milder I_{Kr} inhibition in the border zone (Figure 4B, the first column of I_{Kr},

and SM Table S4 Chronic BZ). Furthermore, T-wave duration in this stage was mainly

determined by the gradient between remote and border zone repolarisation times (Figure 4C,

repolarisation time maps), where more severe APD prolongation in the RZ led to larger

repolarisation gradients and, consequently, larger T-wave duration and amplitude (Figure

2B). Specifically, there was an APD difference of 157 ms between remote and border zone

cell models for the large T-wave case versus only 12 ms for the slight QT-prolongation case,

which accounts for the differences in T-wave peak-to-end duration (162 ms vs 72 ms) and

QT intervals (565 ms vs 380 ms) between these two cases.

Ionic bases of ECG abnormalities in chronic MI

Figure 4: Multiscale explanation of QT and T-wave phenotypes in chronic MI. (A)

Conduction delay in chronic border zone causes slight QT prolongation and large T-wave

phenotypes, as well as large repolarisation dispersion in large T-wave. Red in activation

map show regions of no activation (NA), green in repolarisation map show regions of no

repolarisation (NR). (B) Varying degrees of action potential duration (ADP) prolongation in

the remote zone (RZ) corresponding to extent of QT prolongation (blue arrows), with

decreased calcium amplitude in remote and border zone of both phenotypes, and

corresponding decrease in active tension generation. (C) As in acute MI, I_{Na}, I_{CaL}, and I_{Kr}

remodelling underpin reduced conduction, reduced calcium amplitude, and degree of

prolongation in action potential duration, respectively, in both chronic phenotypes.

4. LVEF failed to indicate the extent of post-MI repolarisation dispersions.

In our simulations, acute MI ionic remodelling yielded mildly reduced LVEF in the baseline

ventricular models (43% to 47%) compared with control (53%) (Figure 5A). LVEF

reductions were caused by contractile dysfunction (Figure 5A, active tension) due to lowered

calcium amplitude in BZ (Figure 3B, intracellular calcium transient), which was directly

caused by inhibitions of I_{CaL} in all acute phenotypes (Figure 3C, I_{CaL}, and SM Table S4). A

complete loss of contractile function in the BZ resulted in a more severe reduction in LVEF

in acute post-MI (to 40% for all acute phenotypes in SM Figure S9). Stroke volumes of the

left and right ventricles were well-matched in control conditions (1 mL difference, see SM

Table S9), and introducing myocardial infarction caused a decrease of stroke volume in the

left ventricles in both acute and chronic MI (see SM Table S9).

Activation failure in Brugada phenocopy caused loss of contractility in the affected

regions in the BZ (Figure 3A, activation and repolarisation time maps) and resulted in a more

severely reduced LVEF to 43% with a substantial non-contracting region with significant

wall thinning (Figure 5A, systolic wall thinning). Acute MI with no observable ECG

abnormalities can still have reduced mechanical function, as measured by an LVEF of 47%

(Figure 5A, the last column). Comparison between this phenotype and the inverted T-wave

phenotype shows that the mechanical dysfunction can be dissociated from repolarisation

gradient and T wave abnormalities.

Chronic post-MI simulations showed mild reduction in peak systolic pressure (by

1 kPa) and some reduction in LVEF (Figure 5B), which were unaffected by differences in

repolarisation gradients. This is because there is a consistent reduction in calcium transient

amplitude in the remote zone that is independent of the extent of APD prolongation (Figure

4B, intracellular calcium transient and membrane voltage). Therefore, both the acute and

chronic electromechanical simulation results showed that the post-MI repolarisation

dispersion were not reflected by LVEF.

For both acute and chronic post-MI, simulations done using the population of

ventricular models showed similar changes to the PV loop as the baseline model across

variabilities in ionic current conductances.

Figure 5: Reduced LVEF and heterogeneous systolic deformation caused by BZ ionic remodeling in both acute and chronic post-myocardial infarction. *Pressure-volume loops are shown in black (control) or red (post-MI) traces for the baseline model, and in gray*

(control) or pink (post-MI) traces for the population of models. (A) Reduced LVEF in all

acute phenotypes due to reduced active tension amplitude in the border zone (BZ1~3).

Brugada phenocopy shows the lowest LVEF due to activation block and loss of contractile

function in the border zone in addition to reduced active tension amplitude in the activated

border zone due to ionic remodelling. Reduced contractile function in infarct and border

zone results in infarct thinning and bulging in systole. Systolic cross section of control

simulation shown in black (CTL) with post-MI cross-sections superimposed. (B) Reduced

LVEF in both chronic phenotypes due to reduced active tension amplitude in remote zone

(RZ) and border zones, independent of the extent of QT prolongation (RZ1, RZ2). Scar

stiffening helped to reduce infarct bulging. Systolic cross-section of control simulation shown

in black (CTL) with post-MI cross-sections superimposed.

5. T-wave alternans and abnormal wave propagation are caused at fast pacing by

cellular alternans and EADs, without reduced LVEF at resting heart rate.

Increased incidence of T-wave alternans is commonly observed in post-MI patients, and

abnormally propagating waves generated from post-MI electrophysiological heterogeneity

can trigger lethal arrhythmic events. T-wave alternans were reproduced in the ventricular

chronic MI simulations at fast pacing (Figure 6A), with RZ2 remodelling, and their

mechanisms were revealed through analysis of the high spatio-temporal resolution of

simulation data. Figure 6A shows upright T-wave morphology and preserved LVEF of 49%

at resting rate of 75 bpm (CL= 800 ms). However, at fast rates (120 bpm, CL = 500 ms),

significant beat-to-beat ST and T-wave morphology alterations were observed. This was due

to large alternans seen in mid-myocardial single cell simulations of the remote zone at CL of

500 ms (Figure 6C, green trace), with EAD-driven alternans. These results support the

importance of stress tests, since alternans in APD and T-wave can occur at fast heart rates

with no sign of LVEF abnormalities at resting heart rate. This is consistent with reports that

T-wave alternans under supine bicycle exercise testing was found to be predictive of

arrhythmic event after acute post-MI (Ikeda *et al.*, 2000).

Simulations with the population of virtual cardiomyocytes models revealed that in

addition to the EAD-driven alternans (SM Figure S14), classical calcium-driven alternans

were also observed in the population of cell models (SM Figure S12 and S13). The key ionic

remodelling underlying calcium-driven alternans include enhanced CaMKII activity and

slower calcium release, as well as suppressed SERCA pump activity in the chronic MI,

which are consistent with previous studies (Livshitz & Rudy, 2007; Zhou *et al.*, 2016;

Tomek *et al.*, 2018) (SM Figure S15). I_{KCa} enhancement in the chronic MI suppressed

alternans generation (detailed analysis provided in SM7, Figures S16 to S17). The median of

calcium amplitude was larger in the alternans models than in the non-alternating post-MI

models (SM Figure S19), in agreement with the preserved LVEF in the simulations. We did

not simulate the effect of this classical calcium-driven alternans on the ECG because the

higher pacing rate at which this phenomenon occurs requires the model to include beta-

adrenergic inotropic effects to preserve realistic systolic mechanical function.

Figure 6: T-wave alternans in simulations underpinned by APD and calcium alternans at

fast pacing (120 bpm), albeit with preserved left ventricular ejection fraction (LVEF=49%)

at rest (75 bpm) for the chronic MI phenotype (A). (B) Simulated APD and calcium traces

in midmyocardial population of models with remote zone 2 (RZ2) remodelling. (C) Large

action potential and calcium transient alternans were caused by EADs in simulations at 120

bpm with midmyocardial cells affected by RZ2 ionic remodelling (green traces,

representative example at 75 versus 120 bpm). A single cell model (in green) was selected

from the population of models (in grey) for embedding into the remote region for ventricular

simulations.

Another case of chronic MI simulation showed prolonged QT interval of 669 ms and

preserved LVEF of 49% at 75 bpm (Figure 7A). At 120bpm, in this case, ECG reflected

chaotic activity (Figure 7A, fast pacing simulation) and loss of coordinated mechanical

function. Abnormal electrotonic waves were caused by large repolarization dispersions.

When isolated cells that showed EADs were embedded in the RZ of a ventricular

simulation at fast pacing, we saw ectopic wave propagation. This was because the EADs in

the RZ generated conduction block, which enabled a large repolarisation gradient to form

between the BZ and RZ, thereby leading to ectopy (Figure 7B). By the end of the first

heartbeat (500 ms), the anterior epicardial BZ was fully repolarised (the red trace), but the

RZ showed EADs (green, purple and yellow traces). This means that after the second beat

stimulus, at 610 ms (Figure 7B), the long APD of the RZ prevented full activation in the

postero-lateral LV (Figure 7B, green and purple marker) while the BZ APD was shorter than

RZ (Figure 7B, red marker) and so could be fully activated. This led to a significant

membrane potential gradient between the BZ (red marker) and RZ (green marker), leading to

ectopic wave generation due to injury current caused by electronic effects at 710 ms in the

boundary between BZ and RZ (Figure 7C, compare red and green traces)(Dutta *et al.*, 2016).

This ectopic wave propagated in an anti-clockwise fashion when viewed from the base from

810 to 910 ms (Figure 7B, basal view). In the right ventricle at 610 ms, incomplete

repolarisation caused a smaller action potential to be elicited by the sinus stimulus (Figure

7C, yellow trace), thus allowing successful propagation of the ectopic wave from the left to

the right ventricle from 910 ms to 1010 ms (Figure 7B).

This episode illustrated that EADs present in the epicardial remote zone cell model

(Figure 7D) resulted in large APD dispersion between BZ and RZ, which functioned as the

trigger of ectopic wave propagation due to electrotonic gradients. Therefore, the prolonged

global QT interval with large T-wave duration and amplitude in leads facing the infarct can

be indicative of the risk of large repolarization dispersion, while the LVEF can be preserved

at rest.

Spontaneous EADs were frequently observed in the chronic MI cellular population of

models (Figure 7D population of models, SM Figure S20), due to less I_{CaL} inhibition in the

chronic MI (left, SM Table S4). The key underlying combination of ionic remodelling for

EAD generation include the inhibition of I_{Kr} and the enhancement of I_{NaL}, which facilitate

reactivation of I_{CaL} (SM Figure S21). Additional contribution of baseline I_{CaL}, I_{Kr}, and I_{NCX}

conductances are consistently observed in all chronic EAD populations (Table S11).

Similarly, the cellular models that showed EADs also had larger calcium amplitude (SM

Figure S22), suggesting a preserved LVEF.

Abnormal wave propagation driven by cellular EADs in chronic MI

Figure 7: Prolonged QT and preserved LVEF at rest can manifest as severely abnormal

ECG at fast heart rates in chronic MI with RZ2 (A), *due to electrotonically-triggered EADs*

across the border zone (B). *In* (B), *membrane potential changes for the first 1010 ms of fast*

pacing simulation, showing ectopic wave generation driven by electrotonic gradient at

710 ms (arrow from red dot to green dot in lateral view), and anticlockwise propagation of

ectopic wave starting at 810 ms (anticlockwise arrow in basal view, and arrow from green

dot to purple dot in lateral view). Ectopic wave propagates towards the right ventricle via

the posterior side at 910 ms (arrow in basal view) and at 1010 ms (arrow in basal view). (C)

Local action potential at anterior (red), lateral (green), posterior (purple), and right

ventricular (yellow) sites. (D) A population of models demonstrating chronic remote zone 2

(RZ2) remodelling in promoting EADs. A representative example was selected from the

population of models that showed EAD and was embedded in ventricular simulations.

DISCUSSIONS

In this study, human electromechanical modelling and simulation enables quantification of

the contribution of electrophysiological abnormalities to clinical phenotypes in post-MI

patients, from ionic to whole-organ dynamics (summarised in Table 1). The credibility of the

human electromechanical models and simulation results is supported by their consistency

with experimental and clinical data from ionic dynamics to ECG and LVEF biomarkers in

healthy, acute and chronic post-MI conditions. Diverse clinical ECG phenotypes are

reproduced in the simulations with different degrees of experimentally reported ionic

remodelling for acute and chronic MI; their signature on the LVEF is however weak, with

only a small reduction observed. The simulated clinical ECG and LVEF phenotypes were

found to be consistent across physiological variabilities in ionic conductances in the baseline

electrophysiological model. Key findings include:

1) In acute MI, T-wave inversion, Brugada phenocopy, and QT prolongation were explained

by reversed transmural dispersion of repolarisation in the border zone and infarct, activation

failure in the border zone, and increased repolarisation time in the border zone, respectively.

2) In chronic MI, large T-wave duration and amplitude reflects large repolarisation

dispersion between remote and border zones, and global QT prolongation was caused by AP

prolongation in the remote zone.

3) Reduction in LVEF is ubiquitous across acute and chronic post-MI phenotypes and can be

explained by decreased intracellular calcium amplitude and activation failure (in the case of

Brugada phenocopy). This effect was independent of changes in the dispersion of

repolarisation.

4) Interestingly, fast pacing simulations with T-wave alternans or abnormal propagation

driven by cellular alternans or EADs both showed preserved LVEF at resting heart rate,

highlighting the fact that preserved LVEF at rest does not guarantee low arrhythmic risk.

Collectively, our results show the proarrhythmic post-MI electrophysiological dispersions

caused by cellular remodelling of ionic currents are reflected in QT and T wave morphology

biomarkers rather than in LVEF, which questions the use of LVEF as the dominant

biomarker in clinical risk stratifications.

Table 1: *Linking clinical ECG and left ventricular ejection fraction (LVEF) phenotypes to*

tissue heterogeneities and subcellular ionic remodelling in acute and chronic post-

myocardial infarction.

Clinical Phenotypes	Tissue or Cell Level phenomena	Corresponding Post Infarction Ionic Remodelling
Acute MI T-wave inversion in ECG	Reversed transmural repolarisation gradient due to delayed repolarization in the epicardial border zone	Inhibition of potassium currents in the border zone

Acute MI Brugada phenocopy in ECG	Delayed repolarisation, as well as a small region of activation failure in the epicardial border zone	Strong inhibitions of sodium, calcium and potassium ionic currents in the border zone
Chronic MI upright tall T-waves in ECG	Large repolarisation time gradient between remote and border zones caused by more severe delay of repolarization in the remote zone	More severe potassium channel suppression in the remote zone
Chronic MI T-wave alternans	Cellular repolarization alternans or early afterdepolarization	Suppressed SERCA and augmented CaMKII activity for alternans; Enhanced late sodium current and suppressed hERG current for early afterdepolarization
Acute MI reduction in LVEF	Reduced calcium amplitude and/or regional conduction block	Inhibitions of calcium and sodium currents
Chronic MI reduction in LVEF	Reduced calcium amplitude	Decreased SERCA activity

1. Acute MI T-wave inversion and Brugada phenocopy are caused by reversed

transmural repolarisation gradient and regional conduction abnormality.

Three distinct types of T-wave morphology were generated by our acute MI biventricular

simulations: T-wave inversion, Brugada-phenocopy and normal upright T-wave. We

obtained them by applying three types of acute BZs in ventricular simulations, considering

both APD prolongation and shortening, as a reflection of the variable experimental results

(Mendonca Costa *et al.*, 2018). Collectively, our results highlighted the importance of

investigating the implications of the various degrees of experimentally-reported ionic

remodelling to explain phenotypic variability of patients with MI. T-wave inversion is a

commonly observed feature in the acute MI patients, and is commonly associated with

arrhythmic risk (Tikkanen *et al.*, 2015). Here we showed the reversed transmural

repolarisation gradient caused by APD prolongation in the epicardial BZ accounted for this

phenotype. The link between transmural repolarisation gradient and T-wave polarity has been reported previously (Okada *et al.*, 2011) and is consistent with our results. Brugada-phenocopy was also observed in some acute MI patients (Anselm *et al.*, 2014), and our simulation results showed it could be a reflection of regional conduction abnormality combined with APD prolongation. Although some animal experiments showed acute MI BZ APD shortening (Mendonca Costa *et al.*, 2018), we showed the prolongation of BZ APD was underlying the QT prolongation, T-wave inversion and Brugada phenocopy in the leads facing the infarct (Supplementary Material Figure S7 showing lead dispersions), which is consistent with the QTc prolongation observed in the anterior leads of acute anterior infarction patients (Guaricci *et al.*, 2018). Apart from the above, normal T-waves and QT intervals were also commonly observed in patients post percutaneous coronary intervention, which can be reproduced when the post-MI repolarisation dispersion was small between BZ and NZ (BZ3). It is worth noting that, in addition to having a mild border zone remodelling as shown in BZ3, a silent ECG signature can also be due to a reduced transmural extent of the infarct, as has been shown in previous computational studies (Loewe *et al.*, 2018; Wang *et al.*, 2021). Therefore, T-wave inversion, Brugada phenocopy and QT prolongation occur in the leads facing the infarct can be useful biomarkers indicating bigger repolarisation dispersions and/or larger transmural extent in the acute MI.

2. Wide and tall T-wave is explained by large repolarisation dispersions between BZ and RZ in healed post-MI hearts.

Our simulated chronic ECGs recapitulated the recovery of T-wave polarity observed in patients after a period of healing. This was achieved through the recovery of the transmural

repolarisation gradient caused by the milder I_{Kr} inhibition in the chronic BZ (Hegyi *et al.*,

2018). Experimental studies in different species showed inconsistent results regarding the

chronic BZ APD (Mendonca Costa *et al.*, 2018). Our chronic BZ remodelling produced

slightly longer APD than the NZ, which is consistent with observations in healed human BZ

(Dangman *et al.*, 1982). However, in minipigs, these remodelling caused shorter BZ APD

than in NZ (Hegyi *et al.*, 2018). This interesting discrepancy between minipigs and human

may be due to the different balance of ionic currents across species, which showed the

benefits of human electrophysiology models in overcoming the inter-species differences.

631   The two types of T-wave morphologies in the simulated chronic ECGs corresponded

to different extents of RZ APD prolongations, which were commonly observed in healed RZ

of post-MI animals (Hegyi *et al.*, 2018), and in failing human myocytes (Li *et al.*, 2004). The

substantial RZ APD prolongation was reflected as global QT prolongation in all leads, and

the large APD dispersion between chronic BZ and RZ generated wide and tall T-waves in the

precordial leads facing the infarct (Supplementary Material Figure S8 for global and

dispersed ECG characteristics). Previous simulation studies also found the T-wave amplitude

and area were proportional to the dispersion of repolarisation (Arteyeva & Azarov, 2017).

Therefore, these results demonstrated that in patients with global QT prolongation, leads with

bigger T-wave amplitudes could reflect increased local heterogeneity in repolarisation.

3. T wave alternans and severely abnormal ECGs at fast pacing are caused by alternans

643   **and EADs in chronic infarction.**

Post-MI ionic remodelling promoted alternans generation, which resulted in T-wave

alternans in simulated ECGs, consistent with the higher incidence of T-wave alternans

reported in post-MI patients (Martin *et al.*, 2009). Two types of repolarisation abnormalities

were observed in our post-MI models: EADs and alternans. One crucial mechanism

promoting alternans behaviour at the cellular level is the increased activity of CaMKII,

observed in the acute MI BZ (Hund *et al.*, 2008*b*), as well as in the hypertrophied and failing

myocardium (Anderson *et al.*, 2011). Enhanced CaMKII phosphorylation may preserve the

contractility of the heart through the phosphorylation of phospholamban and the L-type

calcium channels, but increased RyR phosphorylation by CaMKII resulted in prolonged RyR

opening as well as the enhancement of spontaneous calcium sparks, which can contribute to

alternans and triggered arrhythmias (Maier & Bers, 2007).

Generation of EADs due to chronic post-MI ionic remodelling, such as I_{Kr} inhibition

and I_{NaL} enhancement, was consistent with previous studies (Coppini *et al.*, 2013). We also

found that both repolarisation reserve remodelling (I_{NaL} and I_{Kr}) and calcium system

remodelling (J_{up} and CaMKII) are important for the EAD-driven alternans (details provided

in SM9). EADs in the RZ can create large repolarisation dispersion in the ventricle,

facilitating abnormal electrotonic wave propagations. Similar re-entrant waves caused by

electrotonic gradients were also observed in previous studies of acute ischemia (Ridley *et al.*,

1992; Dutta *et al.*, 2016; Boukens *et al.*, 2021).

4. LVEF should be combined with QT and T-wave characteristics for arrhythmic risk

stratification.

In this study, we observe non-structurally-induced reductions of LVEF in the both the acute

and the chronic post-MI stages. At both stages, ventricles with different extents of

repolarisation dispersion may have similar LVEF because they have similar degrees of

calcium reduction (acute MI T-wave inversion vs normal ST-T, and chronic MI two cases).

Models with inducibility of T-wave alternans and arrhythmia at fast pacing rates may present

with a preserved LVEF at resting heart rates (Figure 6-7). Our cellular level results also

showed models with inducibility of repolarisation abnormalities, such as alternans and

EADs, tended to have more preserved CaT magnitudes at rest rates. Therefore, these

phenomena all support the fact that preserved LVEF measured at rest does not guarantee low

arrhythmic risk.

676   A recent clinical study of post-MI patients with preserved LVEF showed defibrillators

are needed in those patients with electrophysiological risk factors, such as prolonged QTc,

increased T-wave alternans, to prevent sudden cardiac death (Gatzoulis *et al.*, 2019).

Consistently, we also found post-MI alternans and EADs can present as alternans of T-wave

morphology and prolonged QT intervals. In addition to the global ECG changes, our

simulation results also showed increased local repolarisation dispersion can be reflected in

the leads facing the infarct: inverted T-wave and prolonged QT in the acute MI, and wide and

tall T-wave in the chronic MI. Therefore, we suggest the consideration of these signs as

markers of high arrhythmic risk.

5. Limitations

The main goal of this study is to investigate phenotypic variability in ECG and LVEF

biomarkers arising from post-MI ionic remodelling. We have shown that the relationship

between phenotypic variabilities and ionic remodelling remains consistent across

physiological ranges of variation of the ionic conductances in the baseline cell model. Other

sources of variability were not considered in this study including: heart anatomy, location

and timing of the early activates sites, calcium sensitivity, conduction velocity. These could

all modulate quantitatively the findings but we do not anticipate strong implications in the

findings. The effect of variability in location and size of the scar on the ECG has been

explored elsewhere(Li *et al.*, 2024). LVEF reduction in clinical cases are more significant

than in our simulations due to factors other than ionic remodelling: RZ structural remodelling

and elevated myocardium stiffness, and abnormalities in anatomy. However, these effects

have been well-documented elsewhere and this study serves to elucidate the non-structural

mechanisms that underpin LVEF reduction that is linked to electrophysiological remodelling

and arrhythmic risk. The basal plane in our simulation was fixed in space, which was

necessary due to the segmented geometry from clinical MRI and to prevent unphysiological

motion at the truncated basal plane. Despite this limitation, our conclusions regarding the

relative comparisons of mechanical dysfunction are likely to still hold.

In addition, post-MI ionic remodelling can be modulated by other acute and chronic

factors such as autonomic modulation (beta-adrenergic effects), inflammation, cell death, and

metabolic remodelling, which can be explored in future work. The limitations in this study

call for a need for personalised digital twins to be generated in the future to facilitate a better

understanding of the interaction between structural remodelling and electrophysiological

alterations.

6. CONCLUSIONS

Human-based electromechanical simulations reveal ionic mechanisms underlying T-wave

inversion, Brugada phenocopy, and upright T-wave in acute post-MI, as well as the upright

T-wave, QT prolongation and T-wave alternans in the healed chronic MI. In acute MI, while

the potassium current reduction in the border zone was implicated for all phenotypes, the

more severe ECG abnormalities in the Brugada phenocopy implicates additional remodelling

for the sodium and calcium currents, which were also key factors in reduced mechanical

function. In chronic MI, the degree of QT prolongation and the generation of pro-arrhythmic

injury currents were directly related to the severity of potassium current remodelling in the

remote region. In addition, late sodium current remodelling could be an important factor

underpinning T-wave alternans in chronic MI through the promotion of EAD-driven

alternans. Our results show that T-wave inversion, wide and tall T-wave, and QT

prolongation in the leads facing the infarct are indicative of local dispersion of repolarisation,

which is independent from the reduction of LVEF. Our simulation results suggest the

utilisation of T-wave morphology, T-wave alternans and QT prolongation to improve risk

stratification biomarkers even when the resting LVEF is preserved.

REFERENCES

Aggarwal Rajesh & Boyden Penelope A. (1995). Diminished Ca2+ and Ba2+ Currents in Myocytes Surviving in the Epicardial Border Zone of the 5-Day Infarcted Canine Heart. *Circulation Research* **77,** 1180–1191.

Ahnve S (1985). QT interval prolongation in acute myocardial infarction. *Eur Heart J* **6 Suppl D,** 85–95.

Al-Khatib SM, Stevenson WG, Ackerman MJ, Bryant WJ, Callans DJ, Curtis AB, Deal BJ, Dickfeld T, Field ME, Fonarow GC, Gillis AM, Granger CB, Hammill SC, Hlatky MA, Joglar JA, Kay GN, Matlock DD, Myerburg RJ & Page RL (2018). 2017 AHA/ACC/HRS Guideline for Management of Patients With Ventricular Arrhythmias and the Prevention of Sudden Cardiac Death: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *Circulation* **138,** e272–e391.

Anderson ME, Brown JH & Bers DM (2011). CaMKII in myocardial hypertrophy and heart failure. *J Mol Cell Cardiol* **51,** 468–473.

Anselm DD, Evans JM & Baranchuk A (2014). Brugada phenocopy: A new electrocardiogram phenomenon. *World J Cardiol* **6,** 81–86.

Arevalo HJ, Vadakkumpadan F, Guallar E, Jebb A, Malamas P, Wu KC & Trayanova NA (2016). Arrhythmia risk stratification of patients after myocardial infarction using personalized heart models. *Nat Commun*; DOI: 10.1038/ncomms11437.

Arteyeva NV & Azarov JE (2017). Effect of action potential duration on Tpeak-Tend interval, T-wave area and T-wave amplitude as indices of dispersion of repolarization: Theoretical and simulation study in the rabbit heart. *J Electrocardiol* **50,** 919–924.

Bloomfield DM, Steinman RC, Namerow PB, Parides M, Davidenko J, Kaufman ES, Shinn T, Curtis A, Fontaine J, Holmes D, Russo A, Tang C & Bigger JT (2004). Microvolt T-wave alternans distinguishes between patients likely and patients not likely to benefit from implanted cardiac defibrillator therapy: a solution to the Multicenter Automatic Defibrillator Implantation Trial (MADIT) II conundrum. *Circulation* **110,** 1885–1889.

Boukens BJD, Dacey M, Meijborg VMF, Janse MJ, Hadaya J, Hanna P, Swid MA, Opthof T, Ardell JL, Shivkumar K & Coronel R (2021). Mechanism of ventricular premature beats elicited by left stellate ganglion stimulation during acute ischaemia of the anterior left ventricle. *Cardiovasc Res* **117,** 2083–2091.

Caldwell BJ, Trew ML, Sands GB, Hooks DA, LeGrice IJ & Smaill BH (2009). Three distinct directions of intramural activation reveal nonuniform side-to-side electrical coupling of ventricular myocytes. *Circ Arrhythm Electrophysiol* **2,** 433–440.

Cluitmans MJM et al. (2021). Noninvasive detection of spatiotemporal activation-repolarization interactions that prime idiopathic ventricular fibrillation. *Science Translational Medicine* **13,** eabi9317.

Coppini R, Ferrantini C, Yao L, Fan P, Del Lungo M, Stillitano F, Sartiani L, Tosi B, Suffredini S, Tesi C, Yacoub M, Olivotto I, Belardinelli L, Poggesi C, Cerbai E & Mugelli A (2013). Late sodium current inhibition reverses electromechanical dysfunction in human hypertrophic cardiomyopathy. *Circulation* **127,** 575–584.

Dangman KH, Danilo P, Hordof AJ, Mary-Rabine L, Reder RF & Rosen MR (1982). Electrophysiologic characteristics of human ventricular and Purkinje fibers. *Circulation* **65,** 362–368.

Dutta S, Mincholé A, Zacur E, Quinn TA, Taggart P & Rodriguez B (2016). Early afterdepolarizations promote transmural reentry in ischemic human ventricles with reduced repolarization reserve. *Prog Biophys Mol Biol* **120,** 236–248.

Elayi CS, Charnigo RJ, Heron PM, Lee BK & Olgin JE (2017). Primary Prevention of Sudden Cardiac Death Early Post-
Myocardial Infarction: Root Cause Analysis for Implantable Cardioverter-Defibrillator Failure and Currently
Available Options. *Circ Arrhythm Electrophysiol*; DOI: 10.1161/CIRCEP.117.005194.

Gallagher KP, Gerren RA, Stirling MC, Choy M, Dysko RC, McManimon SP & Dunham WR (1986). The distribution of
functional impairment across the lateral border of acutely ischemic myocardium. *Circulation Research* **58,**
570–583.

Gatzoulis KA et al. (2019). Arrhythmic risk stratification in post-myocardial infarction patients with preserved
ejection fraction: the PRESERVE EF study. *Eur Heart J* **40,** 2940–2949.

Guaricci AI, Carità P, Lorenzoni V, Casavecchia G, Rabbat M, Ieva R, Brunetti ND, Andreini D, Di Biase M, Marenzi G,
Bartorelli A, Pepi M & Pontone G (2018). QT-interval evaluation in primary percutaneous coronary
intervention of ST-segment elevation myocardial infarction for prediction of myocardial salvage index. *PLoS*
One; DOI: 10.1371/journal.pone.0192220.

Hegyi B et al. (2018). Complex electrophysiological remodeling in postinfarction ischemic heart failure. *Proc Natl*
Acad Sci USA **115,** E3036–E3044.

Hund TJ, Decker KF, Kanter E, Mohler PJ, Boyden PA, Schuessler RB, Yamada KA & Rudy Y (2008*a*). Role of activated
CaMKII in abnormal calcium homeostasis and I(Na) remodeling after myocardial infarction: insights from
mathematical modeling. *J Mol Cell Cardiol* **45,** 420–428.

Hund TJ, Decker KF, Kanter E, Mohler PJ, Boyden PA, Schuessler RB, Yamada KA & Rudy Y (2008*b*). Role of activated
CaMKII in abnormal calcium homeostasis and I(Na) remodeling after myocardial infarction: insights from
mathematical modeling. *J Mol Cell Cardiol* **45,** 420–428.

Ikeda T, Sakata T, Takami M, Kondo N, Tezuka N, Nakae T, Noro M, Enjoji Y, Abe R, Sugi K & Yamaguchi T (2000).
Combined assessment of T-wave alternans and late potentials used to predict arrhythmic events after
myocardial infarction. A prospective study. *J Am Coll Cardiol* **35,** 722–730.

Jiang M, Cabo C, Yao J, Boyden PA & Tseng G (2000). Delayed rectifier K currents have reduced amplitudes and
altered kinetics in myocytes from infarcted canine ventricle. *Cardiovasc Res* **48,** 34–43.

Land S, Park-Holohan S-J, Smith NP, dos Remedios CG, Kentish JC & Niederer SA (2017). A model of cardiac
contraction based on novel measurements of tension development in human cardiomyocytes. *Journal of*
Molecular and Cellular Cardiology **106,** 68–83.

Levrero-Florencio F, Margara F, Zacur E, Bueno-Orovio A, Wang ZJ, Santiago A, Aguado-Sierra J, Houzeaux G, Grau V,
Kay D, Vázquez M, Ruiz-Baier R & Rodriguez B (2020). Sensitivity analysis of a strongly-coupled human-
based electromechanical cardiac model: Effect of mechanical parameters on physiologically relevant
biomarkers. *Computer Methods in Applied Mechanics and Engineering* **361,** 112762.

Li G-R, Lau C-P, Leung T-K & Nattel S (2004). Ionic current abnormalities associated with prolonged action potentials
in cardiomyocytes from diseased human right ventricles. *Heart Rhythm* **1,** 460–468.

Li L, Camps J, Jenny Wang Z, Beetz M, Banerjee A, Rodriguez B & Grau V (2024). Toward Enabling Cardiac Digital
Twins of Myocardial Infarction Using Deep Computational Models for Inverse Inference. *IEEE Trans Med*
Imaging **43,** 2466–2478.

Livshitz LM & Rudy Y (2007). Regulation of Ca2+ and electrical alternans in cardiac myocytes: role of CAMKII and
repolarizing currents. *Am J Physiol Heart Circ Physiol* **292,** H2854-2866.

Loewe A, Wülfers EM & Seemann G (2018). Cardiac ischemia—insights from computational models. *Herzschr*
Elektrophys **29,** 48–56.

Maier LS & Bers DM (2007). Role of Ca2+/calmodulin-dependent protein kinase (CaMK) in excitation-contraction
coupling in the heart. *Cardiovasc Res* **73,** 631–640.

Margara F, Wang ZJ, Levrero-Florencio F, Santiago A, Vázquez M, Bueno-Orovio A & Rodriguez B (2021). In-silico
human electro-mechanical ventricular modelling and simulation for drug-induced pro-arrhythmia and
inotropic risk assessment. *Progress in Biophysics and Molecular Biology* **159,** 58–74.

Martin DT, Shoraki A, Nesto RW & Rutter MK (2009). Influence of diabetes and/or myocardial infarction on
prevalence of abnormal T-wave alternans. *Ann Noninvasive Electrocardiol* **14,** 355–359.

Mendonca Costa C, Plank G, Rinaldi CA, Niederer SA & Bishop MJ (2018). Modeling the Electrophysiological
Properties of the Infarct Border Zone. *Front Physiol*; DOI: 10.3389/fphys.2018.00356.

Mincholé A, Zacur E, Ariga R, Grau V & Rodriguez B (2019). MRI-Based Computational Torso/Biventricular Multiscale
Models to Investigate the Impact of Anatomical Variability on the ECG QRS Complex. *Front Physiol* **10,** 1103.

Oikarinen L, Viitasalo M & Toivonen L (1998). Dispersions of the QT interval in postmyocardial infarction patients
presenting with ventricular tachycardia or with ventricular fibrillation. *Am J Cardiol* **81,** 694–697.

Okada J, Washio T, Maehara A, Momomura S, Sugiura S & Hisada T (2011). Transmural and apicobasal gradients in
repolarization contribute to T-wave genesis in human surface ECG. *American Journal of Physiology-Heart
and Circulatory Physiology* **301,** H200–H208.

Pogwizd SM, Hoyt RH, Saffitz JE, Corr PB, Cox JL & Cain ME (1992). Reentrant and focal mechanisms underlying
ventricular tachycardia in the human heart. *Circulation* **86,** 1872–1887.

Ridley PD, Yacoub MH & Curtis MJ (1992). A modified model of global ischaemia: application to the study of syncytial
mechanisms of arrhythmogenesis. *Cardiovasc Res* **26,** 309–315.

Sack KL, Davies NH, Guccione JM & Franz T (2016). Personalised computational cardiology: Patient-specific
modelling in cardiac mechanics and biomaterial injection therapies for myocardial infarction. *Heart Fail Rev*
21, 815–826.

Santiago A, Aguado-Sierra J, Zavala-Aké M, Doste-Beltran R, Gómez S, Arís R, Cajas JC, Casoni E & Vázquez M (2018).
Fully coupled fluid-electro-mechanical model of the human heart for supercomputers. *Int J Numer Method
Biomed Eng* **34,** e3140.

Sato A, Hiroe M, Nozato T, Hikita H, Ito Y, Ohigashi H, Tamura M, Takahashi A, Isobe M & Aonuma K (2008). Early
validation study of 64-slice multidetector computed tomography for the assessment of myocardial viability
and the prediction of left ventricular remodelling after acute myocardial infarction. *Eur Heart J* **29,** 490–
498.

Shenthar J, Deora S, Rai M & Nanjappa Manjunath C (2015). Prolonged Tpeak-end and Tpeak-end/QT ratio as
predictors of malignant ventricular arrhythmias in the acute phase of ST-segment elevation myocardial
infarction: a prospective case-control study. *Heart Rhythm* **12,** 484–489.

Solomon SD, Zelenkofske S, McMurray JJV, Finn PV, Velazquez E, Ertl G, Harsanyi A, Rouleau JL, Maggioni A, Kober L,
White H, Van de Werf F, Pieper K, Califf RM, Pfeffer MA, & Valsartan in Acute Myocardial Infarction Trial
(VALIANT) Investigators (2005). Sudden death in patients with myocardial infarction and left ventricular
dysfunction, heart failure, or both. *N Engl J Med* **352,** 2581–2588.

Spargias KS, Lindsay SJ, Kawar GI, Greenwood DC, Cowan JC, Ball SG & Hall AS (1999). QT dispersion as a predictor
of long-term mortality in patients with acute myocardial infarction and clinical evidence of heart failure.
Eur Heart J **20,** 1158–1165.

Stone PH, Raabe DS, Jaffe AS, Gustafson N, Muller JE, Turi ZG, Rutherford JD, Poole WK, Passamani E & Willerson JT
(1988). Prognostic significance of location and type of myocardial infarction: independent adverse outcome
associated with anterior location. *J Am Coll Cardiol* **11,** 453–463.

Streeter DD, Vaishnav RN, Patel DJ, Spotnitz HM, Ross J & Sonnenblick EH (1970). Stress distribution in the canine
left ventricle during diastole and systole. *Biophys J* **10,** 345–363.

Sun K, Stander N, Jhun C-S, Zhang Z, Suzuki T, Wang G-Y, Saeed M, Wallace AW, Tseng EE, Baker AJ, Saloner D, Einstein
DR, Ratcliffe MB & Guccione JM (2009). A computationally efficient formal optimization of regional
myocardial contractility in a sheep with left ventricular aneurysm. *J Biomech Eng* **131,** 111001.

Tikkanen JT, Kenttä T, Porthan K, Huikuri HV & Junttila MJ (2015). Electrocardiographic T Wave Abnormalities and
the Risk of Sudden Cardiac Death: The Finnish Perspective. *Ann Noninvasive Electrocardiol* **20,** 526–533.

Tomek J, Bueno-Orovio A, Passini E, Zhou X, Minchole A, Britton O, Bartolucci C, Severi S, Shrier A, Virag L, Varro A
& Rodriguez B (2019). Development, calibration, and validation of a novel human ventricular myocyte
model in health, disease, and drug block ed. Faraldo-Gómez JD, Barkai N, Hund T & Maleckar M. *eLife* **8,**
e48890.

Tomek J, Tomková M, Zhou X, Bub G & Rodriguez B (2018). Modulation of Cardiac Alternans by Altered Sarcoplasmic
Reticulum Calcium Release: A Simulation Study. *Front Physiol*; DOI: 10.3389/fphys.2018.01306.

Ursell PC, Gardner PI, Albala A, Fenoglio JJ & Wit AL (1985). Structural and electrophysiological changes in the
epicardial border zone of canine myocardial infarcts during infarct healing. *Circ Res* **56,** 436–451.

Vaduganathan M, Claggett BL, Chatterjee NA, Anand IS, Sweitzer NK, Fang JC, O'Meara E, Shah SJ, Hegde SM, Desai
AS, Lewis EF, Rouleau J, Pitt B, Pfeffer MA & Solomon SD (2018). Sudden Death in Heart Failure With
Preserved Ejection Fraction: A Competing Risks Analysis From the TOPCAT Trial. *JACC Heart Fail* **6,** 653–661.

Van Leuven SL, Waldman LK, McCulloch AD & Covell JW (1994). Gradients of epicardial strain across the perfusion
boundary during acute myocardial ischemia. *American Journal of Physiology-Heart and Circulatory*
Physiology **267,** H2348–H2362.

Vazquez R, Bayes-Genis A, Cygankiewicz I, Pascual-Figal D, Grigorian-Shamagian L, Pavon R, Gonzalez-Juanatey JR,
Cubero JM, Pastor L, Ordonez-Llanos J, Cinca J, de Luna AB, & MUSIC Investigators (2009). The MUSIC Risk
score: a simple method for predicting mortality in ambulatory patients with chronic heart failure. *Eur Heart*
J **30,** 1088–1096.

Waks JW, Sitlani CM, Soliman EZ, Kabir M, Ghafoori E, Biggs ML, Henrikson CA, Sotoodehnia N, Biering-Sørensen T,
Agarwal SK, Siscovick DS, Post WS, Solomon SD, Buxton AE, Josephson ME & Tereshchenko LG (2016). Global
Electric Heterogeneity Risk Score for Prediction of Sudden Cardiac Death in the General Population.
Circulation **133,** 2222–2234.

Wang ZJ, Santiago A, Zhou X, Wang L, Margara F, Levrero-Florencio F, Das A, Kelly C, Dall'Armellina E, Vazquez M &
Rodriguez B (2021). Human biventricular electromechanical simulations on the progression of
electrocardiographic and mechanical abnormalities in post-myocardial infarction. *Europace* **23,** i143–i152.

Yao J-A, Hussain W, Patel P, Peters NS, Boyden PA & Wit AL (2003). Remodeling of gap junctional channel function
in epicardial border zone of healing canine infarcts. *Circ Res* **92,** 437–443.

Zhou X, Bueno-Orovio A, Orini M, Hanson B, Hayward M, Taggart P, Lambiase PD, Burrage K & Rodriguez B (2016).
In Vivo and In Silico Investigation Into Mechanisms of Frequency Dependence of Repolarization Alternans
in Human Ventricular Cardiomyocytes. *Circ Res* **118,** 266–278.

Zhou X, Qu Y, Passini E, Bueno-Orovio A, Liu Y, Vargas HM & Rodriguez B (2019). Blinded In Silico Drug Trial Reveals
the Minimum Set of Ion Channels for Torsades de Pointes Risk Assessment. *Front Pharmacol* **10,** 1643.

ADDITIONAL INFORMATION SECTION

Data availability statement

The simulation input files and Alya executable required to replicate the simulated results will

be made available upon request. Python and MATLAB scripts for cellular populations of

models simulations and post-processing of the biventricular simulations can be found at:

https://github.com/jennyhelyanwe/post_MI_postprocessing

Competing interests

The authors declare that they have no competing interests.

Funding

This work was funded in whole, or in part, by the Wellcome Trust (214290/Z/18/Z). For the

purpose of Open Access, the author has applied a CC BY public copyright licence to any

Author Accepted Manuscript version arising from this submission.

This work was supported by a Wellcome Trust Fellowship in Basic Biomedical Sciences to

B.R. (214290/Z/18/Z), an Oxford-Bristol Myers Squibb Fellowship to X.Z.

(R39207/CN063), the Personalised In-Silico Cardiology (PIC) project, the CompBioMed 1

and 2 Centre of Excellence in Computational Biomedicine (European Commission Horizon

2020 research and innovation programme, grant agreements No. 675451 and No. 823712), an

NC3Rs Infrastructure for Impact Award (NC/P001076/ 1), the TransQST project (Innovative

914   Medicines Initiative 2 Joint Undertaking under grant agreement No 116030, receiving

915   support from the European Union's Horizon 2020 research and innovation programme and

916   EFPIA), and the Oxford BHF Centre of Research Excellence (RE/13/1/30181), PRACE-ICEI

917   funding projects icp005, icp013, icp019.

Acknowledgements

920   The authors would like to acknowledge Dr Erica Dall'Armellina, Dr Arka Das, Dr Chris

921   Kelly, and Dr Lei Wang, for discussions on the study.

922
Preprint

This manuscript was first published as a preprint: Xin Zhou, Zhinuo Jenny Wang, Julia Camps,

Jakub Tomek, Alfonso Santiago, Adria Quintanas, Mariano Vazquez, Marmar Vaseghi, Blanca

Rodriguez (2022). [Clinical phenotypes in acute and chronic infarction explained through human

ventricular electromechanical modelling and simulations]. bioRxiv.

https://www.biorxiv.org/content/10.1101/2022.02.15.480392v3