An electrophysiological and kinematic model of *Paramecium*, the "swimming neuron"

Irene Elices[1], Anirudh Kulkarni[1,2], Nicolas Escoubet[3], Léa-Laetitia Pontani[3], Alexis Michel Prevost[3], Romain Brette[1]

[1] Sorbonne Université, INSERM, CNRS, Institut de la Vision, 17 rue Moreau, F-75012 Paris, France

[2] Department of Bioengineering and Centre for Neurotechnology, Imperial College London, South Kensington Campus, SW7 2AZ London, United Kingdom

[3] Sorbonne Université, CNRS, Institut de Biologie Paris-Seine (IBPS), Laboratoire Jean Perrin (LJP), F-75005, Paris

Corresponding author: Romain Brette, romain.brette@inserm.fr

Abstract

Paramecium is a large unicellular organism that swims in fresh water using cilia. When stimulated by various means (mechanically, chemically, optically, thermally), it often swims backward then turns and swims forward again in a new direction: this is called the avoiding reaction. This reaction is triggered by a calcium-based action potential. For this reason, several authors have called *Paramecium* the "swimming neuron". Here we present an empirically constrained model of its action potential based on electrophysiology experiments on live immobilized paramecia, together with simultaneous measurement of ciliary beating using particle image velocimetry. Using these measurements and additional behavioral measurements of free swimming, we extend the electrophysiological model by coupling calcium concentration to kinematic parameters, turning it into a swimming model. In this way, we obtain a model of autonomously behaving *Paramecium*. Finally, we demonstrate how the modeled organism interacts with an environment, can follow gradients and display collective behavior. This work provides a modeling basis for investigating the physiological basis of autonomous behavior of *Paramecium* in ecological environments.

Author Summary

Behavior depends on a complex interaction between a variety of physiological processes, the body and the environment. We propose to examine this complex interaction in an organism consisting of a single excitable and motile cell, *Paramecium*. The behavior of *Paramecium* is based on trial and error: when it encounters an undesirable situation, it backs up and changes direction. This avoiding reaction is triggered by an action potential. Here we developed an empirically constrained biophysical model of *Paramecium*'s action potential, which we then coupled to its kinematics. We then demonstrate the potential of this model in investigating various types of autonomous behavior, such as obstacle avoidance, gradient-following and collective behavior.

Introduction

Behavior depends on a complex interaction between a variety of physiological processes, the body and the environment. This complexity makes it challenging to develop integrative models relating the different components. Thus, a strategy is to study model organisms that are experimentally accessible and structurally simpler than vertebrates. This strategy has been applied in particular to *C. Elegans*, with its 302 neurons and a known connectome (Cohen and Denham, 2019; Cohen and Sanders, 2014; Schafer, 2018). However, electrophysiology is difficult owing to the small size of its neurons (about 3 μm), and developing empirically valid neuromechanical models of *C. Elegans* remains challenging. More recently, other model organisms have been introduced: *Hydra*, with a few thousand neurons and the advantage of being transparent (Dupre and Yuste, 2017; Wang et al., 2020), and jellyfish *Aurelia aurita* (Pallasdies et al., 2019).

Here we propose to develop an integrative model of *Paramecium tetraurelia* (Brette, 2021), which is structurally much simpler than the abovementioned model organisms (Fig. 1A), since it is a unicellular organism, while being large enough to perform intracellular electrophysiology (about 120 μm long (Nagel and Machemer, 2000)). *Paramecium* is a common ciliate, which swims in fresh water by beating its ~4000 cilia (Fig. 1B) (Aubusson-Fleury et al., 2015; Iftode et al., 1989; Nagel and Machemer, 2000), and feeds on smaller microorganisms (bacteria, algae, yeast). It uses chemical signals to find food (Fig. 1C-D), avoids obstacles thanks to mechanosensitivity (Fig. 1E), displays collective behavior, adapts to changing environmental conditions and can even learn to respond to new stimuli (Hennessey et al., 1979).

Figure 1. *Presentation of Paramecium. A, Scanning electron microscopy image (left) (Valentine et al., 2012), and bright field microscopy image (right) of Paramecium tetraurelia (scale bars: 10 μm). B, Typical helicoidal swim of Paramecium (Herbert S. Jennings, 1899). C, Accumulation of paramecia in a drop of acid (Herbert S. Jennings, 1899). D, Trajectory of a single cell in the acid drop, showing directional changes at the boundary (Herbert S. Jennings, 1899). E, Avoiding reaction against an obstacle, showing ciliary reversal followed by reorientation (Jennings, 1906).*

More than a century ago, Jennings described the basis of its behavior as "trial-and-error" (Jennings, 1906). *Paramecium* normally swims in a helicoidal fashion at about 500-1000 μm/s, but when it encounters something undesirable (obstacle, hot region, noxious substance), it produces an *avoiding reaction* (Fig. 2A): it briefly swims backward, then turns and swims forward in a new direction. The avoiding reaction is triggered by a calcium-based graded action potential, which can be observed in an

immobilized cell in response to a current pulse (Fig. 2B). The calcium current is produced by L-type
calcium channels located in the cilia (Eckert, 1972), related to the Ca_V1 family found in neurons, heart
and muscles of mammals (Lodh et al., 2016). Genes for many ionic channels have been found in the
fully sequenced genome (Martinac et al., 2008), and a number have been electrophysiologically
identified (Eckert and Brehm, 1979). Many signaling pathways of neurons have been found in
Paramecium (Plattner and Verkhratsky, 2018). For this reason, *Paramecium* has been called a
"swimming neuron" (Kung and Saimi, 1985) and there is a vast amount of information about its
electrophysiology, from studies done mainly in the 1960-80s (Eckert, 1972; Eckert and Brehm, 1979).
However, there is no empirically based model of its action potential.

We developed a biophysical model of *Paramecium*'s action potential, based on electrophysiological
experiments on immobilized paramecia. We then augmented it by coupling ciliary calcium
concentration with kinematic variables, with a phenomenological model constrained by simultaneous
imaging of fluid motion induced by cilia beating and measurements of trajectories of freely swimming
cells. In this way, we obtain a model of a swimming *Paramecium*, which exhibits spiraling movements
and graded avoiding reactions. Finally, we show how the model can be used to investigate closed-loop
behavior, including chemotaxis and collective behavior.

Figure 2. *The avoiding reaction of Paramecium. A, Typical spontaneous avoiding reaction: the ciliate swims*
backward, then turns and eventually resumes forward swimming, while spinning around its main axis during
the entire movement. Images are separated by 150 ms, with intermediate shaded frames every 37 ms. The cell
was placed in 20 mM NaCl and 0.3 mM CaCl₂ to induce spontaneous avoiding reactions (Kung, 1971). B,
Intracellular recording of a voltage response (bottom right) to a square current pulse of amplitude 300 pA (top
right) in an immobilized cell (left; A: anterior end; P: posterior end), showing a small action potential (in the
standard extracellular solution, see Methods). The arrow points at a small upward inflexion due to the calcium
current. Inset: Paramecium immobilized on a filter (background) with two electrodes. C, Velocity field of the

fluid on a plane ~30 µm above the cell, calculated over the three shaded intervals shown in B. The blue arrow
indicates mean velocity in the whole field, represented twice larger for clarity. The red arrows highlight the
area neighboring the cell. C1, The fluid moves backward, which would make the cell swim forward. C2, The
fluid moves forward. C3, The flow direction reverts on the posterior end, but not on the anterior right end,
resulting in a swirling pattern.

Results

A brief overview of *Paramecium*'s action potential

In this section, we recapitulate known facts about *Paramecium's* electrophysiology, while illustrating
our experimental techniques. To perform intracellular electrophysiology (see Methods,
Electrophysiology), it is necessary to first immobilize the cell. To this end, we use a device we previously
developed (Kulkarni et al., 2020), which uses a transparent filter with holes smaller than the cell and
a peristaltic pump. The pump draws the extracellular solution (4 mM KCl and 1 mM CaCl$_2$) from an
outlet below the filter, immobilizing the cells against the filter (Fig. 2B, inset). Two high-resistance
electrodes are then inserted into the cell and the pump is stopped. The cell is then held by the
electrodes. One electrode is used to inject current, while the other is used to measure the membrane
potential. *Paramecium* is a large cell (about 120 µm long and 35 µm wide for *P. tetraurelia*), which
makes it isopotential (Eckert and Naitoh, 1970; Satow and Kung, 1979).

Depolarization opens voltage-gated calcium channels located in the ~4000 cilia (Aubusson-Fleury et
al., 2015; Iftode et al., 1989; Nagel and Machemer, 2000), similar to the L-type Cav1.2 family in
mammals (Lodh et al., 2016). This can be noticed on Fig. 2B as a small upward deflection before the
peak of the membrane potential (arrow). This calcium current, denoted as I_{Ca}, activates rapidly (a few
milliseconds), producing a current of up to about 4 nA (Oertel et al., 1977). Calcium entry then makes
cilia reorient, which makes them beat forward.

To observe the beating direction, we use 1 µm tracer particles and estimate their velocity with particle
image velocimetry (PIV, see Methods), on a plane about 30 µm above the cell (Fig. 2C1-C3, note that
fluid velocity is much smaller than swimming velocity as it is measured above the cell (Drescher et al.,
2010)). Before the stimulation, the cilia beat backward to the right (Fig. 2C1), which tends to make the
cell move forward, with a spiraling movement over to the left (Jennings, 1904; Machemer, 1972), with
the oral groove facing the spiral axis (Bullington, 1930; Herbert S. Jennings, 1899). When calcium
enters the cilia and calcium concentration reaches about 1 µM (Naitoh and Kaneko, 1972), cilia
reorient and beat forward, slightly to the left (Fig. 2C2), which makes the cell move backward. The
calcium channels inactivate rapidly (a few milliseconds) through calcium-mediated inactivation:
intraciliary calcium binds to calmodulin, which then inactivates the channels (Brehm et al., 1980;
Eckert and Brehm, 1979; Eckert and Chad, 1984). Calcium is then expelled by diffusion, buffering and
pumps, in particular plasma membrane calcium pumps (PMCA) identified in the cilia (Yano et al., 2015,
2013). After calcium has entered the cilia, voltage-gated K$^+$ channels located in the basal membrane
rapidly open, producing a delayed rectifier current I_{Kd} that damps the membrane potential (Fig. 2B,
just after the peak). Calcium also activates a smaller K$^+$ current $I_{K(Ca)}$, which can be seen in the
prolonged hyperpolarization after the stimulation (Saimi et al., 1983; Satow and Kung, 1980).

After the stimulation, when calcium concentration has decreased below ~1 µM (Naitoh and Kaneko,
1972), cilia revert to backward beating (Fig. 2C3). We notice a swirling pattern on Fig. 2C3, which can
be attributed to an asynchronous reversal of different groups of cilia. We will show how this relates to
the change in swimming direction seen on Fig. 2A.

141

Figure 3. *Summary of the model, showing a cilium attached to the base, and the movements of the two main*
ions, Ca^{2+} (red) and K^+ (blue). There is more Ca^{2+} outside than inside, and more K^+ inside than outside. Calcium
enters through ciliary voltage-gated channels as a current I_{Ca}. It then quickly inactivates these channels,
forming a negative feedback loop. Calcium activates motor proteins, triggering ciliary reversal, as well as a
ciliary K^+ channel, producing an outward K^+ current $I_{K(Ca)}$. The motor activation results in calcium
concentration $[Ca^{2+}]_I$ modulating kinematic parameters. Calcium is then expelled, in particular by plasma
membrane Ca^{2+} ATPases (PMCA). Depolarization opens voltage-gated K^+ channels in the basilar membrane,
presumably near the cilium, creating a current I_{Kd} (delayed rectifier). A linear leak current is also present.
Finally, an inward rectifier current I_{Kir} opens at very hyperpolarized voltages, which has little impact on the
avoiding reaction.

We used current-clamp recordings and PIV measurements of fluid motion to build a model of the
action potential together with electromotor coupling, summarized in Figure 3. We chose to use
current-clamp rather than voltage-clamp recordings because good control is difficult to achieve with
high resistance electrodes and several important processes are calcium-gated rather than voltage-
gated, making voltage-clamp less relevant. Our modeling strategy is outlined in Figure 4. We start by
determining passive properties (Fig. 5) by using responses to hyperpolarized pulses, which do not
trigger the calcium-based avoiding reaction. Then we isolate the delayed rectifier current I_{Kd}
mechanically by deciliating the cells, thereby removing the voltage-gated calcium channels (Fig. 6). We
use these findings to build a complete model of the action potential with calcium dynamics coupled
with ciliary reversal (as sketched in Fig. 3), which we fit to electrophysiological responses of ciliated
cells together with measurements of fluid motion (Fig. 7). To couple this electrophysiological model
with kinematics, we examine a simplified hydrodynamic model where cilia reverse with
heterogeneous calcium sensitivity, as suggested by our PIV measurements and previous studies (Fig.
8). This study allows us to infer the form of couplings between calcium concentration and kinematic
variables, which we partially constrain with behavioral measurements of freely swimming cells (Fig.
9). With these couplings, the electrophysiological model can then be used to simulate free swimming
and behavioral responses to stimuli (Fig. 9). Finally, we augment the model with elementary models
of transduction and show simulations of behavior in structured environments, such as obstacle
avoidance (Fig. 10) and gradient following (Fig. 11).

Figure 4. Outline of the modeling strategy. Each number indicates the corresponding figure. The left column
(5-7) is the electrophysiological modeling. The right column (8-9) is the modeling of couplings between
electrophysiology and kinematic variables. Put together, we obtain a model of a freely swimming Paramecium
(9), which is then augmented with elementary sensory transduction models to yield model simulations in
structured environments (10-11).

Passive properties

We start by estimating the passive properties (resistance, capacitance, reversal potentials) with model
fitting techniques (see Methods, *Electrophysiological modeling* and *Model optimization*), and we
compare with previous measurements in the literature. To this end, we use voltage responses to
hyperpolarizing current pulses (duration 100 ms, amplitude 0 to -4 nA in 300 pA increments; Fig. 5A,
top). Such stimuli are known to trigger different voltage-gated currents: a small inactivating calcium
current (Nakaoka and Iwatsuki, 1992; Preston et al., 1992a, 1992b), and a strong inward rectifier K^+
current I_{Kir}, i.e., which lets current pass mainly in the inward direction, below the reversal potential of
K^+ (Oertel et al., 1978; Preston et al., 1990). None of these currents are involved in the avoiding reaction
and the action potential, and therefore they will not be included in the final model. In particular, the
small calcium current is related to the escape reaction, an increase in swimming speed triggered by
hyperpolarizing stimuli (Machemer, 1974; Roesle, 1903). However, the inward rectifier current will
allow us to estimate E_K, the reversal potential of K^+.

When the pulse intensity is strong, an inward current can be seen to activate after ~15 ms. The inward rectifier current has the property to activate mainly below E_K (Lu, 2004). This can be seen by removing K^+ from the extracellular solution (making $E_K = -\infty$). With 4 mM extracellular KCl, hyperpolarization below about -60 mV activates a strong inward current, which is largely suppressed in 0 mM KCl (Fig. 5B). After the pulse, the K^+ current switches from inward to outward as it passes E_K. This results in a change in decay speed, which is noticeable at about -60 mV in Fig. 5A (top, arrow). We use this property to estimate E_K.

Figure 5. *Passive properties and inward rectifier current. A, Top: voltage responses of one cell to negative current pulses (I = 0 to -4 nA in 300 pA increments; dashed lines: start and end of pulses), in the standard extracellular solution (4 mM KCl and 1 mM CaCl₂). The arrow points at an inflexion due to the inward rectifier current I_{Kir}. Bottom: model responses fitted to the data, showing the inferred reversal potential of K^+ (E_K) and the half-activation voltage V_{Kir} of the inward rectifier current. B, Current-voltage relationship over all cells (mean $\pm$ standard deviation, measured at pulse end) in 4 mM KCl (grey) and 0 mM KCl (blue). Removing K^+ from the extracellular solution largely suppresses the inward current. C, Activation curve of the inward rectifier current in the fitted models. The current activates below E_K (E_L is leak reversal potential). The solid curve is the activation function with median parameters, the dashed curve is the activation function of the cell shown in A. D, Fitted parameters over n = 28 cells, grouped in passive parameters and inward rectifier parameters.*

To this end, we fit a biophysical model consisting of a linear leak current and an inward rectifier
current I_{Kir} (Fig. 5A, bottom) (see Methods, *Electrophysiological modeling*) using the Brian 2 simulator
(Stimberg et al., 2019) with the model fitting toolbox (Teska et al., 2020), which applies differential
evolution and gradient descent for least square estimation of model parameters.

We modeled the inward rectifier current as a non-inactivating current with Boltzmann activation, two
gates and a fixed time constant: $I_{Kir} = g_{Kir} n_{Kir}^2 (E_K - V)$ (equations (4-5); g_{Kir} is maximal conductance
and n_{Kir} is the gating variable). Figure 5C shows the activation curve $n_{Kir}^2(V)$ of the cell shown in Fig.
5A (dashed) and the curve with median parameters (solid), which confirms that the current activates
essentially below E_K.

We find that the resting potential V_0 is -24.5 mV $\pm$ 10.6 mV (n = 28; median -22.5 mV; Fig. 5D). Oertel
et al. (1977) previously reported about -23 mV with a slightly different extracellular solution.
Capacitance C is 289 $\pm$ 75 pF. By comparison, *P. caudatum*, which is larger, has a capacitance of about
700 pF. Since *P. caudatum* is 200 μm long and 46 μm wide (Machemer and Ogura, 1979) and *P.*
tetraurelia is 115 μm long and 34 μm wide (Nagel and Machemer, 2000), a simple scaling would predict
a capacitance (200 x 46) / (115 x 34) $\approx$ 2.35 times smaller for *P. tetraurelia,* i.e. about 300 pF, which is
consistent with our estimates.

Resistance at rest R (including the contribution from I_{Kir}) is 126 $\pm$ 62 MΩ. Finally, we find E_K = -48 $\pm$
10 mV, corresponding to an intracellular K^+ concentration $[K^+]_i$ = 29 $\pm$ 11 mM (based on the Nernst
equation and given that $[K^+]_o$ = 4 mM). This is consistent with estimates in the literature obtained with
various methods, varying between 18 and 34 mM (Hansma, 1974; Oertel et al., 1978; Ogura and
Machemer, 1980; Oka et al., 1986).

We briefly describe the parameter estimation results for I_{Kir}, even though these will not be further
used, as only E_K plays a role in depolarized responses. The estimates for total conductance g_{Kir} and half-
activation voltage V_{Kir} are variable across cells, presumably because these parameters are not well
constrained by the data (g_{Kir} and V_{Kir} cannot be estimated independently in the voltage region where
channels are mostly closed). Nonetheless, the results confirm that I_{Kir} activates essentially below E_K.
Activation slope (k_{Kir} = 32 $\pm$ 9 mV) and time constant (τ_{Kir} = 16 $\pm$ 4 ms) are better constrained. With
the estimated parameters, the inward rectifier current contributes about 14% of the resting
conductance (median; 16 $\pm$ 14 %).

A model of the deciliated cell

Figure 6. *The delayed rectifier current measured in deciliated cells. A, Current-voltage relationship in deciliated cells, showing a strong delayed rectifier current for depolarized voltages, and the inward rectifier current for hyperpolarized voltages. B, Top: voltage responses of one cell to positive current pulses (I = 0 to 4 nA in 300 pA increments). Bottom: responses of the two-gate Boltzmann model fitted to the data, showing the inferred half-activation voltage of the delayed rectifier current (dashed). C, Activation and time constant of the delayed rectifier current as a function of voltage in fitted models, with median parameters (solid) and for the cell shown in B (dashed). D, Statistics of fitted parameters (n = 16).*

Next, we analyzed the delayed rectifier current I_{Kd} responsible for repolarization. The ciliary calcium channel can be pharmacologically blocked with W-7, but this drug is toxic to *Paramecium* (Hennessey and Kung, 1984). Instead, we isolated the delayed rectifier current mechanically by removing the cilia with ethanol (Machemer and Ogura, 1979; Ogura, 1981) (see Methods, *Deciliation*). This procedure does not kill the cell, and cilia grow back after a few hours. It removes the voltage-gated calcium channels, which are located in the cilia (Lodh et al., 2016; Machemer and Ogura, 1979), and thereby also removes the calcium-activated K⁺ current $I_{K(Ca)}$. In addition, it is no longer necessary to use the immobilization device (Fig. 6A, inset). As can be seen on Fig. 3, this procedure should leave only the delayed rectifier current I_{Kd}, activated upon depolarization, and the inward rectifier current I_{Kir}, activated upon hyperpolarization. These two currents are indeed observed in electrophysiological measurements (Fig. 6A).

We fitted a Boltzmann model of the delayed rectifier current (see Methods, *Electrophysiological modeling*, equations (6-7) and (12-13)), $I_{Kd} = g_{Kd}n^2(E_K - V)$ (g_{Kd} is the maximal conductance, *n* is the gating variable), to responses to 100 ms depolarizing current pulses (0 to 4 nA in 300 pA increments).

This model turned out to fit the data as well as a Hodgkin-Huxley model, but with fewer parameters
(see Methods, *Model optimization*). Figure 6B (bottom) shows responses of this model, and Figure 6C
shows the activation curve and voltage-dependent time constant with median parameters and those
for the cell shown in Fig. 6B, with detailed statistics in Figure 6D. The delayed rectifier current activates
at a median value of $V_{Kd} \approx 21$ mV (30 ± 23 mV) with a slope $k_{Kd} \approx 7$ mV (9 ± 6 mV). The time constant
peaks at ~4.1 ms (4 ± 1.3 ms) at a voltage $V_\tau \approx 23$ mV (26 ± 48 mV), with a slope $k_\tau \approx 12$ mV (14 ± 8
mV).

Based on these results, we further simplified the model by enforcing $V_{Kd} = V_\tau$ and $k_\tau = 2k_{Kd}$. This
simplification slightly increases the fit error (1.82 vs. 1.8 mV; p = 0.009, two-tailed Wilcoxon test), but
reduces the number of parameters. We used this simplified model in the full model of ciliated cells
(leaving its parameters unconstrained).

The action potential

We now build a model of the action potential of ciliated cells, coupled with cilia reversal (Fig. 7A). Cilia
revert with very small depolarizations, of just a few mV (Machemer, 1974). For this reason, we used
two sets of pulses, large pulses from 0 to 5 nA in 300 pA increments, and small pulses from -100 to 500
pA in 25 pA increments (Fig. 7A, bottom; only positive currents are shown). Tip potentials could
fluctuate between these two sets, therefore we aligned the traces to the median resting potential of -
22 mV, and we fixed E_K at its median value of -48 mV. Simultaneously, we seeded the extracellular
medium with 1 μm tracer particles and imaged their motion at a frame rate of 30 Hz. Particle image
velocimetry was then used to calculate the fluid velocity field, giving an indication of the direction of
ciliary beating, as illustrated in Fig. 2C (See Methods, *Particle image velocimetry*). Figure 7B shows the
cosine of the mean angle α of the velocity field during stimulation, relative to the cell's anteroposterior
axis: 1 means that particles flow towards the posterior end, i.e., the cell is trying to swim forward; -1
means that the cell is trying to swim backward. Thus, cilia revert for a duration longer than the pulse
(120-200 ms in this cell), graded with pulse intensity.

Figure 7. *Fitting the action potential of Paramecium. A, Voltage responses of a cell (top) to two sets of current pulses (bottom), from 0 to 5 nA (in 300 pA increments) and from 0 to 300 pA (in 25 pA increments). B, Ciliary response to the same currents, measured as the cosine of the mean angle of the velocity field, relative to the anteroposterior axis. C, Close up of an action potential triggered by a 1.5 nA current pulse, with the model fit (dashed). The arrow points at an upward deflection due to the calcium current. D, Ionic current calculated by*

subtracting the estimated leak current from the capacitive current. The inward current (I<0, shaded)
corresponds to the calcium current. Integrating this current yields a calcium entry corresponding to a 10 μM
increase in intraciliary calcium concentration. E, Responses of the fitted model. E1, Voltage responses. E2,
Ciliary responses. E3, Voltage-gated calcium current I_{Ca} (top, negative traces), delayed rectifier K^+ current I_{Kd}
(bottom) and calcium-activated K^+ current $I_{K(Ca)}$ (top, positive traces) in the fitted model. Currents are shown
with the electrophysiological convention, i.e., I<0 is inward. E4, Intraciliary calcium concentration in the fitted
model. The dashed lines show the ciliary reversal threshold and the half-inactivation concentration. F, Ionic
currents inferred by the model for the action potential shown in C. G, Statistics of fitted parameters (n = 18).

In the electrophysiological responses, we notice a small upward deflection after stimulation as shown
in Fig. 7C (arrow). This deflection is due to an inward current, the Ca^{2+} current. This current can be
estimated by subtracting the estimated leak current from the capacitive current *C.dV/dt* (passive
properties estimated by model fitting, see below). With a pulse of intensity *I* = 1.5 nA, we find that the
inward part of that current peaks at about 2 nA (Fig. 7D). Note that in this Figure, currents are shown
with the electrophysiological convention so as to compare with previous voltage clamp studies (Eckert
and Brehm, 1979), i.e., I<0 is inward; this is the opposite of the modeling convention. The peak
amplitude shown in Fig. 7D is an underestimation of the calcium current since part of the inward
current may be masked by the K^+ current, but it is comparable to previous estimations in voltage-
clamp (Oertel et al., 1977). The calcium current is known to activate and inactivate quickly, within a
few ms (Brehm et al., 1980; Oertel et al., 1977), as can be seen on Fig. 7D.

We also observe small oscillations (Fig. 7A), due to the interplay between Ca^{2+} and K^+ currents, and a
pronounced hyperpolarization after the pulse. This hyperpolarization is due to a calcium-activated K^+
current $I_{K(Ca)}$. This current has been previously characterized electrophysiologically (Saimi et al., 1983;
Satow and Kung, 1980), as well as genetically and with immunochemistry (Valentine et al., 2012; Yano
et al., 2013).

Thus, we included the following currents in our model: a leak current I_L, a voltage-gated calcium
current I_{Ca}, with calcium-mediated inactivation, a delayed rectifier K^+ current I_{Kd}, and a calcium-
activated K^+ current $I_{K(Ca)}$ (see Methods, *Electrophysiological modeling*). The calcium current I_{Ca} is
produced by ciliary channels similar to L-type Cav1.2 channels (Lodh et al., 2016). We modeled it
similarly to (Eckert and Chad, 1984; Standen and Stanfield, 1982), but we allow for several inactivation
binding sites (equations (14-16)). In addition, the current uses the Goldman-Hodgkin-Katz equation,
which is more appropriate than the linear driving force (E_{Ca} - V) when intra- and extracellular
concentrations are very different (Hille, 2001). The calcium-activated K^+ current $I_{K(Ca)}$ is simply
modelled with a conductance increasing as a Hill function of calcium concentration (equations (17-
18)).

In addition, the model must include calcium dynamics (equation (19)). The decay of calcium
concentration after the action potential may be due to a combination of processes, including diffusion
towards the base, buffering (in particular by centrin), and pumps. We model this combination by a
simple linear model of the calcium flux. However, this is not sufficient because there is a large calcium
flux at rest through the voltage-gated calcium channels, which must also be expelled or buffered. Thus,
we postulate that the resting calcium concentration, about 0.1 μM (Iwadate, 2003; Klauke and Plattner,
1997), is maintained by a pump operating near that concentration, which is consistent with properties
of plasma membrane calcium pumps (PMCA), also present in the cilia (Yano et al., 2015, 2013). We
model this pump with Michaelis-Menten kinetics.

Finally, we couple calcium concentration with ciliary beating angle by a Hill function (equation (20)).

We then fitted this complete model simultaneously to electrophysiological and motor responses to 100
ms current pulses (n = 18), while ensuring that the resting calcium concentration was 0.1 µM (Fig. 7E).
Thus, calcium concentration is not directly measured, but indirectly constrained by several processes:
inactivation of I_{Ca}, activation of $I_{K(Ca)}$, and ciliary reversal.

Figures 7E1 and 7E2 show the fits for the cell shown in Figs. 7A and 7B. Consistently with previous
voltage-clamp measurements (Oertel et al., 1977), the inferred calcium current is transient and peaks
at about ~2.5 nA (-2.5 nA with the electrophysiology convention, Fig. 7E3). A residual current remains,
so that calcium concentration remains high in the model during stimulation (Fig. 7E4). This is
consistent with the fact that ciliary reversal can last for many seconds when the membrane is
depolarized (Machemer, 1974). With large currents, inferred calcium concentration raises to about 22
µM, similar to previous estimations.

The voltage-gated potassium current is delayed relative to the calcium current, and the calcium-
activated K⁺ current raises more slowly and is only dominant during repolarization (Fig. 7E3), with a
maximum of 0.65 nA. This is consistent with previous studies of that current (Satow and Kung, 1980).
Figure 7F shows the three different currents during the action potential shown in Fig. 7C. As previously
argued, the calcium-activated K⁺ current has a small contribution to the early current (Oertel et al.,
1977).

Over the *n* = 18 cells, we find in the fitted models that the calcium current has half-activation voltage
V_{Ca} = -1 mV (0 ± 6 mV), activation slope k = 4.3 mV (4.3 ± 1 mV) and time constant about 0.9 ms (0.9 ±
0.4 ms) (Fig. 7G). Estimated conductance is not well constrained and often very large. This is
presumably because the peak current is mainly determined by the inactivation properties, and
therefore the conductance parameter is not well constrained. Half-inactivation occurs at about K_{Ca} =
3.7 µM ($\log_{10}(K_{Ca}$ in M) = -5.4 ± 0.2). This is close to patch-clamp measurements on cardiac L-type
calcium channels (Höfer et al., 1997). The fitted models have about 4 binding sites (4.4 ± 1.7), larger
than previous models (Eckert and Chad, 1984; Standen and Stanfield, 1982) (which have a single site
but were not constrained by *Paramecium* data). Calcium decays in the model with a median time
constant of 130 ms and the pump operating near rest has a median maximum rate of 87 µM/s.

The delayed rectifier current has similar fitted parameters as in deciliated cells (median k_{Kd} = 4.9 mV,
τ_{Kd} = 4.5 ms), except half-activation tends to be lower (median V_{Kd} = 4 mV) – this might be because the
responses are essentially below V_{Kd}, in the unsaturated part of the activation curve. The calcium-
activated K⁺ current $I_{K(Ca)}$ has low affinity (K_{KCa} = 63 µM, $\log_{10}(K_{KCa}$ in M) = -4.2 ± 0.7). This is consistent
with the observation that in voltage-clamp, this current keeps on increasing for at least one second
(Satow and Kung, 1980). There are about n_{KCa} = 3 binding sites is (3.5 ± 2.3).

Finally, cilia revert when the inferred calcium concentration is about 2.4 µM ($\log_{10}(K_{motor}$ in M) = -5.9
± 0.8). This is close to measurements with triton-permeabilized cells, reporting about 1 µM (Naitoh
and Kaneko, 1972). We note that this and other concentration parameters depend on the estimation
of intraciliary volume, which is approximate.

Overall, parameters of the fitted models are compatible with known properties of the currents and of
ciliary reversal.

Swimming and turning

We now examine how *Paramecium* swims and turns, before coupling the electrophysiological model with swimming motion.

Figure 8. *Swimming and turning. A, Direction of fluid motion during forward swimming (blue) and backward swimming (red), relative to the anteroposterior axis. Averages are shown by arrows. B, Example of helicoidal motion of Paramecium, with the oral groove facing the axis. Highlighted frames are spaced by 750 ms. C, The translational velocity vector **v** is oriented along the anteroposterior axis. The rotation vector **ω** is in the dorsoventral plane (including the oral groove), making an angle θ with the anteroposterior axis. D, Rotating movement of the cell at the end of avoiding reactions of increasing strength (Jennings, 1904). E, Organization of ciliary basal bodies on the oral (ventral) side (Iftode et al., 1989, with permission). The oral apparatus (oa) is in the center (R: right; L: left; A: anterior; P: posterior; as: anterior suture; ps: posterior suture; cy: cytoproct). F, Calculation of kinematic parameters v, θ and ω in a spherical model of radius 60 μm, during successive phases of the avoiding reaction. First column: cilia beat to the rear and right, producing an axisymmetric force field pushing the organism forward while spinning around its axis. Local force amplitude is adjusted for a velocity of 500 μm/s. Second column: cilia revert and now beat to the front and right, pushing the organism backward. Third column: anterior left cilia revert back to the initial direction while anterior right cilia still beat towards the front, and posterior cilia partially revert, beating to the right. Translational velocity is now 0*

and the rotation axis tilts to about 34°. Spinning speed ω also increases by a factor four. Fourth column: all cilia revert back to the initial beating direction. G, Measurement of fluid velocity in a sample cell beyond the anterior end (top) and beyond the posterior end (bottom), in response to 100 ms positive current pulses (1-5 nA), relative to the anteroposterior axis. H, Over n = 9 cells, the direction of posterior motion reverts back about 30 ms after anterior fluid motion (dashed line: linear regression; solid line: identity). Reversal duration is calculated as the time when cos (α) crosses 0, relative to the pulse end time.

Before stimulation, the flow produced by the cilia is directed towards the posterior end, about 11° to the right (Fig. 8A, blue). This should produce a forward left spiraling movement, as documented from observations of free swimming (Jennings, 1904; Machemer, 1972). During a pulse that triggers an action potential, the flow is directed towards the anterior end, about 9° to the right (Fig. 8A, red). This would make the cell swim backward, also spiraling to the left. An example of this spiraling motion is shown on Fig. 8B: the organism swims forward while spinning around a tilted axis. The oral groove faces the spiral axis (Bullington, 1930; Herbert S. Jennings, 1899), which means that the rotational velocity vector $\boldsymbol{\omega}$ is tilted from the main axis towards the oral side by an angle θ (in the median plane; Fig. 8C). In freely swimming paramecia, we found that $\theta \approx 13°$ ($\pm 6.4°$) and the rotation speed $\|\boldsymbol{\omega}\|$ is about 1 cycle/s (1.03 ± 0.2 cycle/s) (see Methods, *Behavioral measurements* and *Behavioral analysis*).

How does *Paramecium* turn? A directional change can occur if the angle θ changes. According to Jennings (1904), θ increases during the avoiding reaction, in relation with stimulus strength (Fig. 8D). In the final model, we will directly couple the calcium concentration $[Ca^{2+}]$ to the three kinematic variables v, θ and ω. In order to postulate plausible calcium-kinematic couplings, and to understand the relationship between ciliary beating patterns and kinematic parameters, in particular θ, we now examine an idealized hydrodynamic model, consisting of a sphere of radius 60 μm, for which we can use analytical formula relating forces and motion (see Methods, *Hydrodynamic model*). Figure 8E shows the ciliature on the ventral (oral) side, which appears to be organized around the oral groove. In the model, we postulate that cilia on the left (L), right (R), anterior (A) and posterior (P) sides may beat in different directions. The fluid produces local forces opposite to the direction of ciliary beating, and at low Reynolds number, the total force and torque map linearly to the translational velocity vector **v** and the rotational velocity vector $\boldsymbol{\omega}$ in the cell coordinate system (Lauga and Powers, 2009).

At rest (low calcium concentration), cilia beat towards the rear, slightly to the right (Fig. 8F, first column), so that the fluid produces a force towards the front, slightly to the left. If the direction of ciliary beating is identical everywhere in spherical coordinates (that is, in terms of the cardinal directions North/South/East/West, Fig. 12), then the force field over the sphere is symmetrical with respect to the main axis. This makes both the total force and the total torque align with the main (antero-posterior) axis, and therefore **v** and $\boldsymbol{\omega}$ are also aligned with that axis, that is, $\theta = 0$. The organism then moves forward, with a spinning movement around the axis. We adjust the force so that the velocity is 500 μm/s, which makes the sphere spin at about 1.8 Hz. Upon stimulation, when calcium concentration is high, cilia revert and beat forward (Fig. 8F, second column), making the organism move backward.

Thus, the organism cannot turn unless there is some asymmetry in the ciliary beating pattern. Machemer (1969) and Párducz (1967) observed that during the turning phase, anterior and posterior cilia beat in different directions; Jennings (1904) observed that left and right anterior cilia beat in different directions, where "left" and "right" are relative to the oral groove. In Fig. 8F (third column), we examine what happens if cilia beat in a swirling pattern around the oral groove: the left anterior cilia beating backward, the right anterior cilia beating forward, and the posterior cilia beating to the right. This corresponds to what would happen near the calcium concentration threshold for global ciliary reversal, if cilia revert back first in the left anterior part, then in the posterior part, then in the right anterior part. The swirling pattern suggests that the cell is going to turn around an axis tilted

from the main axis, in the plane separating the left and right sides, and this can be confirmed
analytically. In this configuration, the sphere does not move along the main axis (v = 0 mm/s), because
the net force along that axis is null, but it turns along an axis tilted by $\theta \approx 34°$ from the main axis. The
sphere also spins about 4 times faster ($\omega/(2\pi) \approx 7.2$ Hz) – this is of course qualitative since
Paramecium is not a sphere. It returns to moving forward when all cilia revert back (Fig. 8F, fourth
column).

This pattern of ciliary reversal is suggested on Fig. 2C. However, given that the particle flow was
measured on a plane ~30 μm above the cell, that the cell could take different shapes and that the
position of the oral groove was often difficult to estimate, it was generally not possible to determine
the precise pattern of ciliary reversal empirically. Nevertheless, it is possible to demonstrate that cilia
revert asynchronously. We measured particle flow separately in two regions of the field, beyond the
anterior end, and beyond the posterior end (Fig. 8G). This was only possible for 9 cells, where video
quality was sufficient (see Methods, *Particle image velocimetry*). We measured the mean angle α of the
flow field during stimulation with current pulses between 1 nA and 5 nA; weak pulses were not
included because particle density tended to be lower due to sedimentation (weak pulses were
recorded after strong pulses).

Figure 8G shows in one cell that posterior cilia (bottom) revert back after anterior cilia (top). Thus,
before stimulation, when calcium concentration is low, anterior and posterior cilia beat in the same
direction. Beating direction is also spatially homogeneous during stimulation, when calcium
concentration is high. However, after stimulation, anterior and posterior cilia beat in different
directions for a short period. This was a reproducible finding across the 9 cells: the posterior side tends
to revert back slightly later than the anterior side (Fig. 8H) (p = 7.10[-17], one-tailed Wilcoxon test), with
a mean delay of 26 ms (s.d. 29 ms). This confirms Párducz' observations (Párducz, 1967), which were
based on electron microscopy of cells fixed during the avoiding reaction.

Modeling the avoiding reaction

This idealized hydrodynamic model shows that an asynchronous control of cilia across the membrane
by calcium can make the organism turn, by transiently increasing the rotation angle θ and the spinning
speed ω. The exact mechanism by which calcium might differentially activate the cilia is beyond the
scope of this paper (see Discussion). Here we simply postulate phenomenological couplings between
calcium concentration and kinematic parameters, as suggested by our analysis (note that the
hydrodynamic model is not explicitly included in our final model, as it was only used to explore the
relation between ciliary patterns and kinematics). As illustrated in Fig. 8C, we will assume that the
translational velocity vector **v** is aligned with the main axis, so that it is fully parameterized by the
velocity v, and that the rotation vector $\boldsymbol{\omega}$ lies in the plane of the oral groove, so that it is parameterized
by its angle θ relative to the main axis and the spinning speed ω. We assume that all three kinematic
parameters (v, θ, ω) are functions of intraciliary calcium concentration [Ca²⁺] (Fig. 7A-C) (See
Methods, *Electromotor coupling*, equations (21-23)).

We model velocity as a Hill function of [Ca²⁺], with threshold equal to the reversal threshold K_{Ca} and n
= 2 sites, linearly scaled to match the maximum positive and negative velocities measured empirically
(Fig. 7A; equation (21)). In freely swimming paramecia, we observed that the median velocity was 472
μm/s (521 ± 285 μm/s) for forward swimming and 370 μm/s (411 ± 200 μm/s) for backward
swimming. Thus, in the model, we simply set both forward and backward maximum velocity at v =
±500 μm/s. Figure 9A shows the resulting function for the cell shown in Fig. 7A-F.

We model both θ and ω as bell functions of $[Ca^{2+}]$ (Fig. 9B, C; equations (22-23); see Methods,
Electromotor coupling), peaking when $[Ca^{2+}]$ is near the global ciliary reversal threshold K_{motor} (2.4
µM), as suggested by our analysis of the spherical model (Fig. 8F). The minimum angle is taken from
measurements of trajectories of freely swimming paramecia ($\theta \approx 13°$). For the maximum angle, we
choose $\theta = 90°$ to account for the strongest avoiding reactions observed by Jennings (Fig. 8D),
corresponding to a rotation normal to the main axis. The minimum spinning speed is based on
measurements ($\omega \approx 1$ cycle/s), and the maximum spinning speed is set to 4 times the minimum, as in
the spherical model (Fig. 8D).

Figure 9. *Simulation of the avoiding reaction. A, Velocity v as a function of intraciliary calcium concentration*
$[Ca^{2+}]$ in the model. B, Angle θ of the rotation axis as a function of $[Ca^{2+}]$ in the model. C, Spinning speed ω as a
function of $[Ca^{2+}]$ in the model. Angle and spinning speed increase at intermediate Ca^{2+} concentration, as
implied by the spherical model in Fig. 8F. D, Simulated model trajectory with three 2 ms current pulse
stimulations of increasing amplitude. Images are shown at 400 ms intervals. Without stimulation, the organism

swims in spiral, with the oral groove facing the spiral axis. A very small stimulation deviates the trajectory.
Stronger stimulations produce avoiding reactions, with backward swimming and turning. E, Example of an
observed Paramecium trajectory showing a directional change without backward swimming (right), followed
by a full avoiding reaction (left). Images are shown at 400 ms intervals, starting on the right. F, Backward
swimming duration (F1) and reorientation angle (F2) as a function of current amplitude for 2 ms pulses. Red
and black curves show results for the same model but different initial positions of the oral groove, differing by
a quarter of a cycle. G, Backward swimming duration (G1) and reorientation angle (G2) as a function of current
pulse duration T with 100 pA amplitude. H, Reorientation angle vs. backward swimming duration in n = 1138
spontaneous avoiding reactions of Paramecium, showing a positive correlation (linear regression r = 0.2, p ≈
10^{-11}). About 15% of data points are not represented (larger angle or duration). Colors represent contour lines
of the distribution.

In this way, we obtain a model in which all kinematic variables are coupled to the electrophysiological model. We note that, in contrast with the electrophysiological part, this model is only loosely constrained by measurements (see Discussion). We can then calculate organism motion from these variables, and thereby simulate behavior in an environment (see Methods, *Kinematics*). In the following, the model of one particular cell is chosen for illustration, the same cell as in Fig. 7.

We first examine the trajectory of a model stimulated by 2 ms currents of varying amplitude (Fig. 9D) (see Methods, *Behavioral scenarios*). Without stimulation, the organism swims in a helicoidal path. With a small stimulation amplitude (0.3 nA), the organism changes direction without swimming backward. At larger amplitude (0.5 nA), the organism swims backward for a very short time then turns and swims forward. When the amplitude is increased (5 nA), backward swimming is more noticeable. Directional changes without backward swimming do occur in freely swimming paramecia, as illustrated on Fig. 9E: the organism first changes direction without swimming backward, then does an avoiding reaction.

In more detail (Fig. 9F1), we observe that the cell swims backward when current intensity exceeds a threshold (here 372 pA), then backward swimming duration tends to increase with intensity. The reorientation angle following the stimulation changes continuously with stimulation strength, but in a complex way (Fig. 9F2). In particular, small stimulations can trigger large turns without backward swimming. In addition, the directional change depends on the initial position of the oral groove: the red and black curves of Fig. 9F2 correspond to the same cell but an oral groove position (spinning angle) differing by a quarter of a cycle. This occurs because the observation plane is fixed while the organism spins around its main axis, and because movement is constrained in the plane.

The characteristics of the avoiding reaction also depend on stimulus duration (Fig. 9G). If the pulse amplitude is fixed (I = 0.1 nA) and its duration is increased, then the duration of backward swimming increases (Fig. 9G1), and reorientation angle tends to increase for large durations but is not monotonous near threshold (Fig. 9G2).

When we examine spontaneous avoiding reactions of freely swimming paramecia, we find that both backward swimming duration and reorientation angle vary broadly (156 ± 81 ms and 114 ± 66 °, respectively) (Fig. 9H), and there is a small although highly significant correlation (linear regression, r = 0.2, p = 10^{-11}). Thus, backward swimming duration and reorientation angle are variable and not deterministically related.

A closed-loop behavioral model of *Paramecium*

We now use the model in closed loop, to describe how the interaction between organism and
environment may give rise to behavior (see Methods, *Behavioral scenarios*), as opposed to motor
responses to predetermined input stimuli (open loop). The following scenarios are relatively abstract
and are meant mainly as illustrations of the possible uses of this integrative model to investigate the
relation between physiology and behavior.

Figure 10. *Interaction of a model Paramecium with a generic stimulus, modelled as a positive current*
proportional to the cell area within the stimulus area. A, Trajectory of the model doing several avoiding
reactions against the stimulus. B, Membrane potential (top), stimulus current (middle) and intraciliary
calcium concentration (bottom) at the first contact. Contact occurs at the boundary with the shaded region.
Orange curves indicate backward swimming. Several weak avoiding reactions occur in succession. C,
Trajectory of the model where sensory transduction has a 40 ms activation/deactivation time constant. In red,
the stimulus is placed 300 μm further away. D, Same as B, for the black trajectory in C. The stimulus current
lasts longer and peaks after the organism has started reacting, resulting in a stronger avoiding reaction.

First, we consider an organism swimming towards a generic object, which triggers a depolarizing
current when in contact with the membrane (for example a chemical substance, or hot water) (Fig.
10). Thus, we simply consider that the stimulus current is proportional to the surface area in contact
with the stimulus (see *Methods, Sensory transduction*). In contrast with previous situations, the
stimulus is not pre-determined but depends on behavior. As Dewey pointed out (1896), "*the motor*
response determines the stimulus, just as truly as sensory stimulus determines the movement.". When the
organism touches the object, a current is triggered, which depolarizes the membrane (Fig. 10A, 10B).
As the cell swims into the object, the current increases until an action potential is triggered. The cell
then swims backward, moves out from the object and the current stops. Thus, the sensory current is
necessarily small and short, because the organism withdraws as soon as the current reaches threshold.

This results in a small avoiding reaction, and the organism bumps again repetitively against the object
until it finally escapes (Movie 1).

Larger movements can be obtained if sensory transduction has slower kinetics (Figs. 10C, D and Movie
2). Here, the sensory current follows the stimulation with a time constant of 40 ms, modelled with first
order kinetics (to simplify, the spatial spread of channels is not modeled; equation (24)). In this case,
the sensory current keeps on increasing (slightly) after the organism has started swimming backward
and it lasts longer. This results in a larger avoiding reaction.

Although the model is deterministic, the directional change can be described as pseudo-random. If the
object is moved 300 μm further (Fig. 10C), then the organism escapes with a larger angle. This is
because the organism spins while swimming, so that its oral groove takes a different position when it
touches the object.

We now consider that the object is a disk in a square environment (Fig. 11A and Movie 4). To avoid
boundary effects, we consider that the environment has the topology of a torus (paramecia escaping
to the left reappear to the right). To account for spontaneous avoiding reactions (occurring at a rate of
about 0.2 Hz in our behavioral measurements), we added a noisy current to the membrane equation.
The paramecia are modeled as in Figs. 10C and D, with slow transduction. Fig. 11A shows 100
trajectories starting from random positions and simulated for 20 s. The proportion of paramecia inside
the disk decreases from 19 to 7%. Thus, the disk acts as a repelling stimulus. Occasionally, a
paramecium gives an avoiding reaction against the boundary, makes a large turn and swims backward
into the disk. In this case, it continues being stimulated and swims backward through the disk until it
escapes. This peculiar behavior might be avoided if currents of opposite polarity were triggered when
stimulating the rear, as is the case for thermal stimuli (Tominaga and Naitoh, 1992) and some chemical
substances (Oami, 1996). Traces shown in Fig. 11A show the membrane potential V, the stimulus
current I and the calcium concentration $[Ca^{2+}]$ for the trajectory shown in red, where we can
distinguish a number of spontaneous avoiding reactions and one avoiding reaction against the disk.

Next, we examine how an object can act as an attractive stimulus (Fig. 11B and Movie 5). This behavior
can be obtained if the cell responds to a hyperpolarizing stimulus with adaptation, so that a
depolarization is triggered when the stimulus stops. A simple model that exhibits this behavior is one
where the stimulus triggers currents through two pathways with different kinetics: a hyperpolarizing
current with fast kinetics, and a depolarizing current of equal magnitude with slow kinetics (equations
(25-27)). In this way, a transient hyperpolarizing current is triggered when the stimulus switches on,
and a transient depolarizing current is triggered when the stimulus switches off. This is shown on Fig.
11B. The proportion of paramecia inside the disk increases with time (from 15 % to 45 % in 20 s). It
can be seen that paramecia tend to aggregate inside the disk, mostly near the boundary. This is due to
the small avoiding reactions but also to the curvature of the disk.

Figure 11. *Closed-loop behavior of the Paramecium model. A, Top: trajectories of 100 models swimming for*
20 s in a torus with a depolarizing circular stimulus, modelled as in Fig. 10C. The proportion of cells in the disk
quickly decays (below). Membrane potential, stimulus current, and intraciliary calcium concentration are
shown for the trajectory highlighted in red, which does an avoiding reaction against the disk after a number
of spontaneous avoiding reactions. B, 100 model trajectories with a circular stimulus triggering an adapting
hyperpolarizing current. Organisms tend to make avoiding reactions on the inner boundary of the disk. The

proportion of cells in the disk increases over time. The highlighted trajectory enters the disk around t = 5 s with a large hyperpolarization, then displays several avoiding reactions against the boundary of the disk before exiting. C, Paramecia swimming in a linear stimulus gradient, modelled as in B. The position of 200 cells starting at position x = 5 mm is displayed every 5 s. D, Collective behavior in model paramecia induced by respiration and chemosensitivity. CO_2 produced by cells is displayed in shades of grey (normalized to the spatial peak), and diffusion is simulated. CO_2 concentration represents an attracting stimulus modelled as in B and C.

Using the same model, we then place the paramecia in a linear stimulus gradient (Fig. 11C); the environment is toric in the transversal dimension (up and down boundaries are glued). We place all paramecia at the same initial position, with random orientations. The population then rapidly ascends the gradient. This occurs because a depolarizing current is produced when the stimulus decreases, triggering an avoiding reaction. This behavior has been observed in *Paramecium* with thermal gradients (H. S. Jennings, 1906; Mendelssohn, 1895), and shares similarities with bacterial chemotaxis (Berg, 1975).

Finally, we show an example of collective behavior (Fig. 11D). *Paramecium* produces CO_2 by respiration, which acidifies its medium, and it is attracted by weak acids (Dryl, 1973; Houten, 1978; H. S. Jennings, 1899). As a result, it can form aggregates, for example around a source of food or at the bottom of a depression slide (H. S. Jennings, 1899). We simulate the production of CO_2 by paramecia and its diffusion in the medium (a square with torus topology), together with sensitivity to CO_2 modeled in the same way as in Fig. 9B and C (see Methods, *Collective behavior*). CO_2 concentration is represented in Fig. 11D as grey shades, after normalization. In this simulation, paramecia progressively form aggregates.

Discussion

Summary

We have built an integrative model of *Paramecium* that combines electrophysiology and motility. The model is informed by previous experimental literature and constrained by specific electrophysiology and trajectory measurements. It consists of an empirically constrained biophysical model of the action potential, with a coupling to kinematic parameters, which is more phenomenological. It can be simulated as a model of autonomous behavior in various environments.

The electrophysiological model was built by model fitting to current clamp data, with calcium-dependent properties indirectly constrained by ciliary reversal data. This method recovered properties of individual currents compatible with previous measurements obtained by different means. For example, the calcium threshold for ciliary reversal was estimated to be ~2 µM, the same order of magnitude as measured by varying extracellular calcium concentration in *Paramecium* with permeabilized membranes (Naitoh and Kaneko, 1972). This is notable since calcium was not measured but only inferred from electrophysiology (indirectly through the estimation of the calcium current by the fitting procedure). The fitting procedure also determined that the calcium-dependent K+ current is small during the action potential, as previously determined with voltage-clamp experiments (Oertel et al., 1977), but dominant after stimulation. The magnitude and time scale of calcium currents estimated by fitting were also compatible with voltage-clamp measurements (Oertel et al., 1977). Quantitative fitting allowed us to estimate additionally the calcium inactivation threshold (~3 µM) and the number of inactivation sites (~4).

By measuring ciliary induced flows during action potentials, we found that ciliary reversal is not synchronous across the cell, confirming previous observations obtained with electron microscopy

(Párducz, 1967). We showed with a simple hydrodynamic model that asynchronous ciliary reversal
allows the organism to turn, namely if the order of ciliary reversal follows a swirling pattern around
the oral groove. From these findings, we built a phenomenological model of the coupling between
calcium concentration and the main kinetic parameters, constrained with trajectory measurements
(speed, angle).

The integrated model shows helicoidal swimming with graded avoiding reactions, where backward
duration swimming and reorientation angle increase with stimulus strength or duration. As observed
in spontaneous behavior, the model can also slightly reorient without swimming backward.

Behavior of the autonomous model is more complex than stimulus-response experiments, because the
relation between sensory stimulus and motor response is circular. In particular, we noticed that the
interaction with an object (e.g. a chemical substance) critically depends on the properties of sensory
transduction. For example, efficient avoidance of the object requires persistent stimulation, e.g. with
slow sensory activation/deactivation. Furthermore, sensory adaptation to a hyperpolarizing stimulus
makes the stimulus attractive. Another possibility, which is non-stimulus-specific, is that excitability
adapts to voltage changes (Platkiewicz and Brette, 2010), for which there is some electrophysiological
evidence in *Paramecium* (Satow and Kung, 1979). This can allow the model to follow a stimulus
gradient. Finally, collective behavior can arise if organisms are sensitive to a substance that they
produce. In summary, relatively complex behavior can be generated by the interaction of this simple
"swimming neuron" with its environment. We note that we have not included any specific empirical
model of transduction, which would be required in any empirical investigation of behavior using our
model.

Limitations

This work has many limitations. First, ionic currents were measured simultaneously rather than in
isolation, although we could isolate I_{Kd} by deciliation. This was partly for technical reasons (one cannot
measure $I_{K(Ca)}$ while blocking the calcium current), and partly to ensure a global fit of the entire model
to the action potential. Nonetheless, current overlap may cause difficulties for model fitting. For this
reason, we strived to choose the simplest models that captured the phenomenology.

A second limitation is that calcium was not directly measured. Instead, it was indirectly constrained
by several observed phenomena: calcium-dependent inactivation of I_{Ca}, calcium-dependent activation
of $I_{K(Ca)}$, and ciliary reversal. Calcium imaging has been performed previously in *Paramecium* by
pressure injection of a calcium indicator, in other contexts (Iwadate et al., 1997; Iwadate and
Kikuyama, 2001; Klauke and Plattner, 1997). However, it is technically challenging to perform
quantitative time-resolved measurements of ciliary calcium, because the cilia represent a small
fraction of the total volume (2-3%) and beat at about 20 Hz.

Related to this limitation, our estimates of calcium-dependent parameters, for example the ciliary
reversal threshold, depend on an estimate of the effective ciliary volume. Compared to our estimate,
based on electron microscopy measurements, this effective volume may be reduced by crowding, or
increased by fast buffering. Changing this parameter results in proportional changes in concentration
parameters. However, the fact that the fitted ciliary reversal threshold is close to the threshold
measured on permeabilized *Paramecium* suggests that our estimate (also used in (Oertel et al., 1977))
was reasonable.

Another limitation is we did not measure ciliary beating directly, but rather its effect on the fluid. This
was motivated by the fact that we were interested primarily in the movement induced by ciliary
beating, as well as by technical reasons. The constraints of measuring fluid motion while the cell is
impaled with electrodes made it difficult to consistently achieve good spatial resolution over the entire
recordings. In future work, high speed imaging of ciliary beating could be used to determine the spatial
pattern of ciliary reversal with higher precision, although contractions of the cell may complicate the
analysis.

Because of these technical limitations, our model of electromotor coupling was highly simplified,
restricted to a phenomenological relation between calcium concentration and three kinetic
parameters. It could also be that this relation is not instantaneous, involving more indirect pathways.

Finally, we considered only generic rather than biophysical models of sensory transduction, and we
did not consider mechanical or hydrodynamic interactions with objects (Berke et al., 2008; Jana et al.,
2015; Ohmura et al., 2018). More generally, the behavioral repertoire of *Paramecium* includes other
aspects that we did not attempt to model, such as the escape reaction (Machemer, 1974; Roesle, 1903),
contractions (Nakaoka and Machemer, 1990), trichocyst discharge (Hamel et al., 2011; Knoll et al.,
1991) and gravitaxis (Jensen, 1893).

How *Paramecium* turns

Since our model directly couples calcium concentration to kinetic parameters, it is not tied to any
specific hypothesis about the ciliary beating pattern. However, turning is only possible if ciliary
reversal is asynchronous, leading to a strongly asymmetrical ciliary beating pattern, otherwise the
action potential would only trigger a back-and-forth movement in the direction of the main axis. We
have shown that one possibility, compatible with the observed movement, is that cilia beat in a swirling
pattern around the oral groove.

It is known that there is some structural and molecular heterogeneity between cilia, in particular
between locomotor and oral cilia (Aubusson-Fleury et al., 2015). Whereas basal bodies are regularly
placed on the dorsal side, they are spatially arranged on the ventral side with a characteristic pattern
around the oral groove (see Fig. 8E). The beating frequency during helicoidal swimming is also
spatially heterogeneous (Jung et al., 2014), and tail cilia are also known to be immobile (Machemer
and Machemer-Röhnisch, 1984). Ciliary heterogeneity is likely to be more complex than a distinction
between oral and locomotor cilia, because when *Paramecium* is cut in two pieces below the oral
groove, both pieces can turn in a similar way (Jennings and Jamieson, 1902). Such heterogeneity is
likely a general feature of motile microorganisms, some of which can exhibit complex gaits (Wan,
2020).

Machemer (1969) and Párducz (1967) described an asynchrony between anterior and posterior
ciliary reversal, with anterior cilia returning to their initial beating direction before posterior cilia,
which we confirmed with our PIV measurements. However, this is not sufficient to produce turning: if
both anterior and posterior beating patterns are axisymmetric, any combination of them would still
produce movement along the main axis. Jennings mentions that there is also a reversal asynchrony
between the anterior left and anterior right cilia, such that all anterior cilia transiently beat towards
the oral groove during the avoiding reaction (Jennings, 1904). More detailed investigation is necessary
to clarify this question.

Physiologically, asynchronous ciliary reversal can be due to differential calcium sensitivity, that is, the
calcium threshold for reversal might vary across cilia. This is the implicit assumption of our model. It
could also be that there are differences in calcium entry or removal across cilia (e.g. a gradient of
calcium channel expression, or calcium pumps, or calcium buffering molecules). Some studies suggest
that cyclic nucleotides may also differentially regulate the reversal threshold (Noguchi et al., 2000,
1991).

Previous models

We are aware of two previous attempts to model *Paramecium*'s action potential, neither of which was
based on quantitative measurements. Hook and Hildebrand (1979) used a calcium channel model with
instantaneous transitions, an ohmic current-voltage relation (instead of GHK), an inactivation state
accessible only from the closed state, and no voltage-dependent K^+ channel (only a model of $I_{K(Ca)}$,
which is not the major K^+ current). Kunita et al. (2014) used a Hodgkin-Huxley type model with
voltage-dependent inactivation of calcium channels, which is not the main inactivation mode of this
channel (Brehm et al., 1980), even though the phenomenon exists on a slow timescale (Hennessey and
Kung, 1985). The model included two calcium channels (fast and slow), for which there is no
electrophysiological support, and their relative activation was given as a function of time after stimulus
start (i.e., it is not modeled). The calcium-dependent K^+ current was not included. Neither
electrophysiological model was fitted to data, and neither was coupled to a kinematic model.

Perspectives

The model could be improved by addressing the technical limitations listed above. In particular, it
would be most enlightening to measure calcium concentration in the cilia at high temporal and spatial
resolution, although it might require new technical developments. Further investigations should be
carried out to understand in detail how *Paramecium* turns: to measure the spatial pattern of ciliary
responses and to determine how this heterogeneity is achieved physiologically.

We have addressed only the avoiding reaction of *Paramecium*. The modeling effort could be completed
by addressing other behavioral aspects, such as the escape reaction (increased ciliary beating speed
upon hyperpolarization), which involves distinct hyperpolarization-activated channels (Brette, 2021).
Paramecium is sensitive to many sensory modalities, including temperature, various chemical
substances, mechanical stimulation, light. Thus, the model should be completed by models of sensory
transduction, as well as of mechanical interaction with objects. This would allow us to use the model
to investigate the physiological basis of behavior of *Paramecium* in complex environments.

Finally, this work opens the perspective of addressing complex autonomous behavior in ecological
environments, including adaptation, learning and problem solving (Brette, 2021), with a systemic
modeling approach.

Materials and Methods

Paramecium culture and preparation

Cultures of *Paramecium tetraurelia* were obtained from Éric Meyer, Institut de Biologie, Ecole Normale
Supérieure, Paris, France. For electrophysiological experiments (at Institut de la Vision), paramecia
were co-cultured with *Klebsiella pneumoniae*, where each week 1 mL of culture was reinjected into 5
mL of Wheat Grass Powder (WGP) buffer supplemented with 1 µL of beta-sitosterol. Cultures were
kept at room temperature (about 20°C). Cells were harvested in the early stationary growth phase,
between 3 and 5 days after feeding them. To wash and concentrate cells for experiments, a droplet of
culture (approximately 600 µL) was placed in a narrow neck volumetric flask before adding
extracellular solution used for electrophysiology (see below). Due to negative gravitaxis (Naitoh and
Eckert, 1972), paramecia tend to accumulate at the top of the solution. Thus, after approximately 10
min, a concentrated population of cells were retrieved from the top of the flask and placed in a
microcentrifuge tube for at least 3h for adaptation (Machemer-Röhnisch and Machemer, 1989; Oka et
al., 1986). The tube was shaken before collecting cells to perform an experiment.

The culture method differed slightly for the behavioral measurements with freely swimming
paramecia, because these cultures were done in another lab (Laboratoire Jean Perrin). Instead, before
an experiment, bacteria were first grown in 5 mL of WGP for 24 h at 27°C, then paramecia were grown
by adding 1 mL of *Paramecium* culture and 1 µL of beta-sitosterol to the bacterized WGP, for 48 h at
27°C in the dark. About 0.4 mL of cell suspension were then pipetted from the top of the culture tube
into 4 mL of extracellular solution (see *Electrophysiology*), at least 20 minutes before an experiment.

Swimming pools

Freely swimming paramecia were imaged at room temperature (∼25°C) in square pools of side length
30 mm and depth 340 µm. These were obtained using micromilling and molding techniques. A
Plexiglas mold, consisting of a square trench, is first milled with a square end mill of diameter 1 mm
using a CNC micro-milling machine (Minitech, Machinary Corp., USA). Then a liquid mixture of Poly-
DiMethyl Siloxane (PDMS, Sylgard 184, Dow Corning, USA) and its crosslinker (10:1 mass ratio) is
poured onto the Plexiglas mold. It is immediately placed in a vacuum chamber for at least 1 h to remove
any air bubbles. Crosslinking of the mixture is then obtained by placing the whole in an oven at 65°C
for at least 4 h. Finally, the resulting transparent elastomer pool is gently peeled off the mold and put
on a microscope glass slide. Prior to any experiment, the pool is exposed to an oxygen plasma for about
1 min to render the PDMS surface hydrophilic and prevent the trapping of air bubbles.

Behavioral measurements

For all behavioral experiments, about 500 µL of the cell suspension is pipetted into the pool with a
concentration of 300 – 600 cells/mL. Trajectories are imaged at 50 Hz with a CMOS camera (Blackfly
S BFS-U3-51S5M-C, Flir, USA, 2448x2048 pixels^2, 10 bits), acquired with its dedicated acquisition
software (Spinview, Flir, USA). A high magnification variable zoom lens (MVL12X12Z, Thorlabs) is
used and yields a pixel size of 3.81 µm. The pool is uniformly illuminated with a dark field
configuration, by placing ∼10 cm beneath the pool a square LED panel (EFFI-SBL, Effilux, France), on
top of which a fully opaque mask is positioned, partially covering the LED panel (typically ¼ of its
surface). The LED panel produces a red light (wavelength λ = 625 nm) to minimize phototaxis
(Iwatsuki and Naitoh, 1983, 1982). Movies of the swimming paramecia are 200 s long (see Movie 7).

To limit hard drive space, images are stored without their background with lossless compression (TIFF
format). The background image is computed by taking for each of its pixels the minimum pixel
intensity over the first 100 frames. It is then subtracted from each frame, and pixels with an intensity

value below a threshold (automatically computed with the "triangle method", see e.g. (Zack et al.,
1977)) are set to 0.

Trajectories are extracted with the open source tracking software FastTrack (Gallois and Candelier,
2021), and manually inspected for corrections. Briefly, the software fits an ellipse to the cell's shape,
and disambiguates front and rear based on the asymmetry of the pixel histogram along the main axis.
Trajectories shorter than 1 s and sequences where the cell is immobile are discarded. Errors in
front/rear identification are automatically corrected as follows: when the cell turns by more than 20°
over two successive frames, it is considered an error and the angle is flipped.

Trajectories with circling motions are also discarded. To this end, we calculate the proportion of the
trajectory where the cell turns clockwise versus anti-clockwise (for trajectories longer than 4 s). These
proportions should be balanced (0.5) for helicoidal trajectories. The trajectory is eliminated if these
proportions differ by more than 0.1 from the expectation, with manual confirmation. In total, there
were n = 554 selected trajectories.

Behavioral analysis

Analysis of helicoidal trajectories

We manually selected 20 trajectories presenting clear helicoidal motion in the focal plane from 2
experiments, totaling 121 s. In each helicoidal trajectory, cell orientation $\gamma(t)$ varies periodically with
period T. We fitted $\gamma(t)$ to a sinusoidal signal. We found $T = 1.02 \pm 0.27\,s$ (mean $\pm$ s.d.),
corresponding to a spinning speed $\omega = \frac{1}{T} = 1.03 \pm 0.2$ cycle/s or about 2π/s in radians. The amplitude
was $\theta = 13 \pm 6.4°$, the angle relative to the spiral axis.

Analysis of avoiding reactions

An avoiding reaction is defined as a portion of trajectory during which the cell swims backward. This
backward swimming is detected when the instantaneous motion vector **m** and the orientation vector
o (posterior to anterior) point to opposite directions, i.e., $\boldsymbol{m} \cdot \boldsymbol{o} < 0$. Avoiding reactions consisting of a
single pair of frames were discarded. The mean frequency of spontaneous avoiding reactions was
calculated as the number of avoiding reactions across all trajectories, divided by the total duration,
yielding 0.18 Hz.

Only reorientation events involved in planar avoiding reactions were selected, based on
measurements of the eccentricity of the ellipse that best fits the shape of paramecia. Whenever this
eccentricity went below 0.8, the event was discarded. The total reorientation angle was obtained by
summing all successive instantaneous reorientation angles during the entire avoiding reaction.

In Fig. 7G, the 2D probability density of reorientation angle and backward duration was calculated with
Gaussian kernel density estimation.

Deciliation

Deciliated cells were obtained by adding 96% ethanol to a tube containing the previously washed and
adapted cells in the extracellular solution up to a final concentration of 5% (v/v) (Ogura, 1981). Then

the tube was shaken for 2 min and left to rest for 1 min. Deciliated cells were collected from the lower
half of the tube since they no longer accumulate at the top of the solution. Cilia start to grow back after
approximately 30 min. Thus, as described in (Ogura, 1981), in some experiments we blocked cilia
regrowth by adding 10 mM of colchicine to the extracellular solution.

Electrophysiology

The extracellular solution used in all experiments contains 1 mM $CaCl_2$, 4 mM KCl and 1 mM Tris-HCl
buffer with pH 7.2, except for Fig. 5B, where there was no KCl (blue curve). Microelectrodes of ~50 MΩ
resistance were pulled using a micropipette puller (P-1000, Sutter Instrument) from standard wall
borosilicate capillary glass with filament (o.d. 1 mm, i.d. 0.5 mm, Harvard Apparatus). They were filled
with a 1 M KCl solution using a MicroFil non-metallic syringe needle (MF 34G-5, World Precision
Instruments); a few recordings were done with 3 M KCl (no particular change was noticed).

We used an upright microscope (LNScope, Luigs & Newmann) with two objectives, a 20× air objective
(SLMPLN Plan Achromat, Olympus) to locate cells, and a 40× water immersion objective (LUMPLFLN,
Olympus) with DIC contrast enhancement for electrophysiology and imaging.

Paramecia were immobilized using the device described in (Kulkarni et al., 2020). Briefly, paramecia
are immobilized against a transparent filter (Whatman Cyclopore polycarbonate membranes;
diameter 25 mm, pore diameter 12 μm) thanks to a peristaltic pump (Gilson Minipulse 3) that
circulates the fluid from below the filter to above the device. Two microelectrodes are then lowered
into the cell, and the pump is stopped. The cell is then held in place by the electrodes.

Electrophysiology recordings were performed using an amplifier with capacitance neutralization
(Axoclamp 2B and Axoclamp 900 A, Molecular Device) and an analog–digital acquisition board
operating at a sampling frequency of 40 kHz (USB-6343, National Instruments). Custom Python
programs (https://github.com/romainbrette/clampy) were used to control the acquisition board.

Membrane potential was recorded with the reading electrode while 100 ms current pulses of various
amplitudes were injected through the second electrode, with at least 1 s between successive trials.

Particle image velocimetry

To measure the flows induced by cilia beating, the bath was seeded with 1 μm silica or polystyrene
particles (~0.2 mM) after paramecia were immobilized and the pump was stopped. Because of
sedimentation, particle density was typically higher at the beginning of the experiment. Images were
recorded at 30 Hz with a high-sensitivity CCD camera (Lumenera Infinity 3-6UR) over a 1392×1392
pixels region of interest surrounding the cell (8 bits depth, pixel width 0.178 μm). Frames were
synchronized with electrophysiology recordings using a digital trigger.

Frames were preprocessed by removing the background (average image) and band-pass filtering
(subtraction of two Gaussian filters with standard deviation 1 μm and 1.3 μm). Consecutive frames
were then analyzed with particle image velocimetry (PIV) using the OpenPIV Python package
(https://github.com/OpenPIV/openpiv-python.git), which calculates the velocity field using image
cross-correlation. We used 50 μm windows with 2/3 overlap.

In each frame, we calculated the mean angle of the velocity vector over the entire field, using circular
mean (argument of the mean complex unit vector; occasional missed frames were discarded). We then
subtracted the angle of the anteroposterior axis. The position of anterior and posterior ends was

measured manually. As the two ends can be visually ambiguous, they were automatically corrected (by swapping) when the flow measured before stimulation was directed towards the anterior end (indicating backward swimming).

In Figure 8A, for each cell we averaged the mean angle over all currents and over the 300 ms before stimulus (blue) or over the second half of the stimulus (red), for positive currents (<5 nA).

We also calculated the mean angle in the anterior and posterior regions as indicated in Fig. 8G. Each region is a half-plane orthogonal to the main axis, starting at one end. For this analysis, we selected n = 9 cells with high quality video recordings and clear cell positioning, indicated by an absence of missed frames and a pre-stimulus flow deviating by less than 45° from the main cell axis. The average was restricted to responses to large pulses (1 to 5 nA), because those were recorded before the small pulses and therefore had a higher density of particles (due to sedimentation).

Electrophysiological modeling

In this section, we describe the biophysical models. The parameter values are obtained by fitting the models to the data (section *Model fitting*).

Electrode model

All recordings were done with two electrodes, an injecting electrode and a reading electrode. Because of the capacitance and resistance of the injecting electrode, the current injected in the cell is a low-pass filtered version of the command current (Brette et al., 2008). To estimate this current, we model the injecting electrode as a simple RC circuit and estimate its parameters R_e and τ_e from responses to small pulses, assuming passive cell responses:

$$C\frac{dV}{dt} = -g_L(V - V_0) + I_e \tag{1}$$

$$\tau_e \frac{dV_2}{dt} = V - V_2 + R_e I + \Delta V \tag{2}$$

$$I_e = \frac{V_2 - V - \Delta V}{R_e} \tag{3}$$

where V is the membrane potential, assumed identical to the potential of the reading electrode, V_2 is the potential of the injecting electrode, I_e is the current injected in the cell, and ΔV accounts for a difference in tip potentials. The membrane equation (first equation) is a rough linear model of the cell, but only parameters R_e and τ_e are used subsequently, to estimate I_e from I according to the last two equations (ΔV has no impact on I_e and therefore can then be discarded). In the 29 ciliated cells analyzed for passive properties, we found R_e = 121 $\pm$ 8 MΩ and τ_e = 1 $\pm$ 0.9 ms.

Ionic currents

Paramecium electrophysiology is reviewed in (Eckert and Brehm, 1979) and updated in (Brette, 2021; Valentine and Van Houten, 2022). *Paramecium* in an isopotential cell (Eckert and Naitoh, 1970; Satow and Kung, 1979). Thus, we consider a single membrane equation:

$$C\frac{dV}{dt} = I_L + I_{Kir} + I_{Kd} + I_{Ca} + I_{K(Ca)} + I$$

where C is membrane capacitance, $I_L = g_L(E_L - V)$ is the leak current, I_{Kd} is the delayed rectifier K+ current responsible for repolarization, I_{Ca} is the ciliary voltage-dependent Ca^{2+} current, $I_{K(Ca)}$ is the calcium-activated K+ current, I_{Kir} is the inward rectifier K+ current and I is a stimulating current. We did not include a few other electrophysiologically identified currents that are less relevant for this study, namely: Na+ (Saimi, 1986; Saimi and Ling, 1990) and Mg^{2+} (Preston, 1998, 1990) currents (since our extracellular solution does not contain these two ion species), and hyperpolarization-activated calcium currents responsible for the escape reaction (Nakaoka and Iwatsuki, 1992; Preston et al., 1992a, 1992b), which we did not model.

The inward rectifier current I_{Kir} is a K+ current activated by hyperpolarization, most strongly below E_K (Oertel et al., 1978). It is modeled as follows:

$$I_{Kir} = g_{Kir}n_{Kir}^p(E_K - V) \tag{4}$$

$$\tau_{Kir}\frac{dn_{Kir}}{dt} = \frac{1}{1 + \exp\left(\frac{V - V_{Kir}}{k_{Kir}}\right)} - n_{Kir} \tag{5}$$

where $p = 1$ or 2 ($p = 2$ in the final version). We made this simple modeling choice because this current was only used as a way to infer the reversal potential E_K. In particular, we did not include inactivation (Preston et al., 1990). We also tested a version of the model where the linear driving force $(E_K - V)$ is replaced by the Goldman-Hodgkin-Katz expression (Hille, 2001), but it made no significant difference in fitting results.

For the delayed rectifier current I_{Kd}, we tested several models of the type:

$$I_{Kd} = g_{Kd}n^p(E_K - V) \tag{6}$$

$$\tau_{Kd}\frac{dn}{dt} = n_\infty(V) - n \tag{7}$$

We tested two classes of models. The Hodgkin-Huxley model is:

$$\alpha_n(V) = \frac{a_{Kd}}{\mathrm{exprel}\left(\frac{V_{Kd} - V}{k_{Kd}^a}\right)} \tag{8}$$

$$\beta_n(V) = b_{IK}\exp\left(\frac{V_{Kd} - V}{k_{Kd}^b}\right) \tag{9}$$

$$n_\infty(V) = \frac{\alpha_n(V)}{\alpha_n(V) + \beta_n(V)} \tag{10}$$

$$\tau_{Kd}(V) = \tau_{Kd}^{min} + \frac{1}{\alpha_n(V) + \beta_n(V)} \tag{11}$$

where

$$\mathrm{exprel}(x) = (e^x - 1)/x$$

For numerical stability (near $x = 0$), we use this special function in the code rather than the explicit expression.

The Boltzmann model is:

$$n_\infty(V) = \frac{1}{1 + \exp\left(\frac{V_{Kd} - V}{k_{Kd}}\right)} \tag{12}$$

$$\tau_{Kd}(V) = a_{Kd} + \frac{b_{Kd}}{\cosh\left(\frac{V - V_\tau^{Kd}}{k_\tau^{Kd}}\right)} \tag{13}$$

The voltage-gated calcium current I_{Ca} is a calcium-inactivated current located in the cilia (Brehm et al., 1980; Eckert and Brehm, 1979). The corresponding channels have been genetically identified; they are similar to the Cav1 mammalian family (L-type), with a putative calmodulin binding site (Lodh et al., 2016). We model it similarly to (Chad et al., 1984; Standen and Stanfield, 1982):

$$I_{Ca} = g_{Ca} m^2 h f_{GHK}(V) \tag{14}$$

where m is the activation gating variable, h is the inactivation gating variable, and $f_{GHK}(V)$ is the normalized current-voltage relation of the open channel, given by the Goldman-Hodgkin-Katz equation (Hille, 2001). In the Hodgkin-Huxley model, this relation is linear. However, with very different intracellular and extracellular calcium concentrations, a better model is the Goldman-Hodgkin-Katz equation. Resting intracellular concentration is about 50-200 nM (Iwadate, 2003; Klauke and Plattner, 1997), and rises to an estimated 20 µM during an action potential (Oertel et al., 1977). In contrast, extracellular concentration is 1mM in our experiments. Thus, we neglect intracellular concentration, which yields:

$$f_{GHK}(V) = \frac{1}{\text{exprel}(2FV/RT)} \tag{}$$

where F is the Faraday constant, R is the gas constant, and T = 293 K is temperature (20°C). Here, extracellular concentration has been lumped into g_{Ca}, which is now homogeneous to a current, while $f_{GHK}(V)$ is unitless and has no free parameter.

The activation gating variable is governed by:

$$\tau_m \frac{dm}{dt} = \frac{1}{1 + \exp\left(\frac{V_{Ca} - V}{k_{Ca}}\right)} - m \tag{15}$$

while the inactivation gating variable is a Hill function of intraciliary calcium concentration $[Ca^{2+}]$:

$$h([Ca^{2+}]) = \frac{1}{1 + \left(\frac{[Ca^{2+}]}{K_{Ca}}\right)^{n_{Ca}}} \tag{16}$$

This is similar to the model of (Standen and Stanfield, 1982), except that the number of sites n_{Ca} is allowed to be greater than 1, because we found that this was necessary to fit our data.

A calcium-activated K+ current has been identified by comparison with Pawn mutants lacking functional voltage-activated calcium currents (Satow and Kung, 1980). The current is largely reduced by EGTA. Genomic analysis indicates the presence of both BK and SK channels, with SK channels immunochemically identified in the cilia (Valentine et al., 2012; Yano et al., 2013). We model the current as follows, with activation as a Hill function of $[Ca^{2+}]$:

$$I_{K(Ca)} = g_{K(Ca)} m([Ca^{2+}])(E_K - V) \tag{17}$$

$$m([Ca^{2+}]) = \frac{1}{1 + \left(\frac{K_{K(Ca)}}{[Ca^{2+}]}\right)^{n_{K(Ca)}}} \tag{18}$$

Calcium dynamics

Resting intracellular calcium concentration $[Ca^{2+}]_0$ has been estimated between 50 and 200 nM (Iwadate, 2003; Klauke and Plattner, 1997). We chose $[Ca^{2+}]_0 = 100$ nM. Calcium enters the cilia when calcium channels open. The concentration increase is spatially uniform along the cilium (Iwadate and Suzaki, 2004). It then decreases by three mechanisms: buffering, pumps, and diffusion. Buffering can occur with a variety of calcium-binding proteins, an important one being centrin, located in the infraciliary lattice, at the base of cilia (Plattner, 2015; Plattner and Klauke, 2001). Plasma membrane calcium pumps (PMCA) have been identified in the basal membrane with low affinity, around 10^{-7} M (Wright and van Houten, 1990), and also in the cilia (Yano et al., 2015, 2013). Suppressing the ciliary PCMAs by RNA interference prolongs backward swimming, which means that they are involved in the removal of calcium after an action potential. In principle, calcium can also diffuse to the basal cytosol. However, this has not been observed (Husser et al., 2004). This might be because of cilia volume compared to the cell, or because calcium is buffered at the base of cilia. Both phenomena can be modeled by diffusion to the cilium base, with fixed resting concentration at the boundary.

We lump these diverse mechanisms into two simple processes: a linear process, with rate proportional to $([Ca^{2+}]-[Ca^{2+}]_0)$, modelling diffusion and low-affinity buffers, and a high-affinity process operating near rest, with rate given by a Hill function of $[Ca^{2+}]$, modelling PMCAs or high affinity buffers. This results in the following equation:

$$\frac{d[Ca^{2+}]}{dt} = \frac{I_{Ca}}{2Fv_{\text{cilia}}} - \lambda([Ca^{2+}] - [Ca^{2+}]_0) - \frac{J}{1 + \frac{[Ca^{2+}]_0}{[Ca^{2+}]}} \tag{19}$$

where v is the volume of cilia and F is the Faraday constant. It can be seen that the role of the high-affinity process in this model is to counteract the calcium flow at rest, namely $J = I_{rest}/Fv_{\text{cilia}}$, while the low-affinity process independently tunes the rate of calcium removal after an action potential.

There are 3000 – 4000 cilia in *P. tetraurelia* (Aubusson-Fleury et al., 2015; Iftode et al., 1989; Nagel and Machemer, 2000), with the upper estimates likely including oral cilia. Each cilium is 10-12 μm long (Eckert and Naitoh, 1970; Sedar and Porter, 1955). Each cilium is 270 nm wide but the fiber bundle is 200 nm wide (Aubusson-Fleury et al., 2015). This yields a total volume between 950 and 2750 μm³. We used the previous estimate $v_{\text{cilia}} = 1700$ μm³ from (Oertel et al., 1977), which is compatible with these bounds, but the uncertainty is large. In addition, the effective volume might be smaller because of crowding, or larger because of fast buffering. In practice, an error in the estimation of ciliary volume will translate into an equivalent change in all calcium binding constants (as well as λ and J). Binding constants and volume cannot be determined independently, because the (inverse) volume effectively acts as a unit for those constants.

For stability, the numerical implementation of calcium-dependent equations used equivalent versions written as a function of

$$p_{Ca} \equiv \log \frac{[Ca^{2+}]}{[Ca^{2+}]_0}$$

1041   where $[Ca]_0$ = 0.1 µM is the resting concentration. For example, inactivation is rewritten as:

$$h([Ca^{2+}]) = \frac{1}{1 + \exp\left(n_{Ca}(p_{Ca} - p_{K_{Ca}})\right)}$$

1043   This equivalent change of variables avoids numerical issues when $[Ca^{2+}]$ approaches 0. The calcium
1044   dynamics equation rewrites as follows:

$$\frac{dp_{Ca}}{dt} = \frac{I_{Ca}}{2F[Ca^{2+}]_0 v_{\text{cilia}}} e^{-pCa} + \lambda(e^{-pCa} - 1) - \frac{J}{1 + e^{pCa}}$$

1046

Electromotor coupling

1048   Cilia reorient when intraciliary calcium concentration reaches about 1 µM (Naitoh and Kaneko, 1972).
1049   We model the ciliary angle as a Hill function of $[Ca^{2+}]$:

$$\alpha = \alpha_0 + \frac{\Delta\alpha}{1 + \left(\frac{K_{\text{motor}}}{[Ca^{2+}]}\right)^{n_{\text{motor}}}} \tag{20}$$

1051   where angles are relative to the anteroposterior direction ($\alpha = 0$ when cilia beat to the rear), and
1052 K_{motor} is the reversal threshold.

1053   Velocity is modeled as an affine transformation of a Hill function with coefficient $n = 2$, changing sign
1054   at K_{motor}:

$$v = -v_{\text{max}} + \frac{2v_{\text{max}}}{1 + \left(\frac{[Ca^{2+}]}{K_{\text{motor}}}\right)^2} \tag{21}$$

1056   where v_{max} = 500 µm/s is maximum velocity (both backward and forward), according to our
1057   measurements.

1058   The angle θ of the rotation axis and the spinning speed ω are modeled as bell functions peaking at
1059 K_{motor}:

$$\theta = \theta_{\text{min}} + 2\frac{\theta_{\text{max}} - \theta_{\text{min}}}{\left(\frac{K_{\text{motor}}}{[Ca^{2+}]}\right)^2 + \left(\frac{[Ca^{2+}]}{K_{\text{motor}}}\right)^2} \tag{22}$$

$$\omega = \omega_{\text{min}} + 2\frac{\omega_{\text{max}} - \omega_{\text{min}}}{\left(\frac{K_{\text{motor}}}{[Ca^{2+}]}\right)^2 + \left(\frac{[Ca^{2+}]}{K_{\text{motor}}}\right)^2} \tag{23}$$

1062   We set θ_{min} = 13° based on our measurements, and θ_{max}= 90°, to allow for planar rotations as
1063   illustrated in Fig. 8D. We set $\omega_{\text{min}} = 2\pi/s$ (1 cycle/s) based on our measurements and $\omega_{\text{min}} = 8\pi/s$ (2
1064   cycles/s), based on the doubling found in the spherical model (Fig. 8F). Thus, no extra free parameter
1065   is introduced.

1066

 Sensory transduction

 Instantaneous transduction

1069 In Fig. 10, 10A and 10B where a well-delimited object (half-plane or disc) acts as a stimulus, we first
1070 calculate the proportion of the cell surface in contact with the object. To this end, cell shape is
1071 determined by the formula proposed by (Zhang et al., 2015):

1072
$$y(x) = \frac{b}{2}\left(\sqrt{1 - 4\frac{x^2}{a^2}} - \beta \sin\left(\frac{2\pi x}{a}\right)\right)$$

1073 where x is the position along the major axis and y the position along the minor axis, a = 120 μm is cell
1074 length, b = 35 μm is cell width and β = 0.15 is an asymmetry factor. We then simply calculate the
1075 intersection of cell and object shapes (as pixel images). The stimulus current is then $I = I_0 p$, where I_0
1076 is maximum current and p is the fraction of the cell image within the object.

1077

1078 *Delayed transduction*

1079 In Fig. 10B and 10A, we simply consider that the transduction current activates and deactivates with
1080 a time constant τ_I:

1081
$$\tau_I \frac{dI}{dt} = I_0 p - I \tag{24}$$

1082 This simple model corresponds to channels with finite opening and closing rates (namely, opening rate
1083 $\alpha = \tau_I s$ and closing rate $\beta = \tau_I(1 - s)$, where s is proportional to the stimulus $I_0 p$) but does not take
1084 into account the spatial recruitment of channels.

1085

1086 *Adaptation*

1087 In Fig. 11B, C and D, we consider that there are two pathways with opposite polarity, a fast pathway
1088 and a slower pathway:

1089
$$\tau_{\text{fast}} dI_{\text{fast}} = s - I_{\text{fast}} \tag{25}$$

1090
$$\tau_{\text{slow}} dI_{\text{slow}} = s - I_{\text{slow}} \tag{26}$$

1091
$$I = I_{\text{slow}} - I_{\text{fast}} \tag{27}$$

1092 where τ_{fast} = 40 ms and τ_{slow} = 200 ms. Thus, for a constant stimulus s, the stationary current is 0. In
1093 Fig. 11B (disc stimulus), the stimulus is $s = I_0 p$. In Fig. 11C and D, where the environment is spatially
1094 continuous, the stimulus is simply the value at the center of the cell.

1095

1096 **Model optimization**

1097 Model parameters are estimated with the model fitting toolbox of the Brian simulator (Stimberg et al.,
1098 2019; Teska et al., 2020) (https://github.com/brian-team/brian2modelfitting). Briefly, the software
1099 performs least square optimization using a combination of global optimization algorithms (we used

differential evolution) and gradient descent, where the gradient is calculated symbolically from the model equations. Optimization with multiple objectives is done by adding the errors associated to the different objectives. Compiled code is automatically produced by code generation. Each fitting procedure took up to a few hours, and fitting scripts were run in parallel on different cells, using a small cluster of 3 PC with 8 cores each.

After model fitting, cells were discarded if passive properties were abnormal, indicating a bad recording (C > 500 pF or R<30 MΩ or R>500 MΩ), or if $E_K > E_L$ (which is biophysically impossible), indicating a fitting problem.

Electrode fitting

First, for each cell we estimated electrode resistance R_e and time constant τ_e from responses to small 100 ms pulses, both hyperpolarizing and depolarizing (|I|<0.5 nA), using equations (1-3). The error criterion was the sum of quadratic errors on both electrode potentials, measured from 100 ms before to 100 ms after the pulse. The estimated parameters were then used in subsequent fits.

Hyperpolarized fits

For Fig. 5, we fitted the models described above with I_L and I_{Kir}, (equations (4-5)), with least square minimization of the error on the reading electrode potential, taken from pulse start to 50 ms after the pulse. The stimuli were 100 ms pulses with amplitude between -4 and 0 nA in 300 pA increments.

We fitted the model with p =1 and with p = 2 for I_{Kir}. Using two gates (n^2) gave better fits than using one (n = 40; median error 2.3 vs. 2.6 mV; p = 8.10^{-5}, one-tailed Wilcoxon test). We also fitted the n^2 model with the ohmic driving force replaced by a Goldman-Hodgkin-Katz model, but it did not yield any significant improvement (p = 0.27, two-tailed Wilcoxon test). Selection criteria (see above) were passed by n = 28 cells. Statistics of fitted parameters are shown in Table 1.

	Mean	Median	s.d.	s.e.m.
C (pF)	288.87	278.82	75.32	18.83
E$_K$ (mV)	-48.27	-47.92	9.54	2.38
E$_L$ (mV)	-20.32	-19.03	11.32	2.83
V$_{Kir}$ (mV)	-130.27	-120.75	35.36	8.84
g$_L$ (nS)	9.41	7.11	6.93	1.73
g$_{Kir}$ (nS)	1375.38	873.36	1206.00	301.50
k$_{Kir}$ (mV)	31.53	30.23	8.89	2.22
τ_{Kir} (ms)	15.44	15.10	3.69	0.92
1/R (nS)	1.21	1.14	0.72	0.18
V$_0$ (mV)	-24.56	-22.51	10.64	2.66

Table 1. *Statistics of fitted parameters (n = 28) for hyperpolarized responses (model with two gates).*

Deciliated fits

For Fig. 6, we fitted the models described above with I_L and I_{Kd}, with least square minimization of the error on the reading electrode potential, taken from pulse start to 50 ms after the pulse. The stimuli were 100 ms pulses with amplitude between 0 and 4 nA in 300 pA increments. Parameters E_K and C were taken from the previous fit to hyperpolarized responses. To make sure the short onset is well captured, the time interval is split in two windows: the first 30 ms and the rest of the response, and each window is equally weighted (meaning that a data point in the first window contributes more than a data point in the second window).

We tested the Hodgkin-Huxley (HH) type model (equations (6-11)) and the Boltzmann model (equations (6-7) and (12-13)). The two models performed similarly (n = 21; median 1.78 mV vs. 1.69 mV; p = 0.59, two-tailed Wilcoxon test). The median number of gates was 1.13 in the HH model (1.1-1.8, 25-75% interval) and 1.7 (1.3-2.7) in the Boltzmann model. In both models, the minimal time constant was very small (median 0.1 vs. 0.6 ms). Therefore, we chose a model with 2 gates (n^2) and a nearly null minimal time constant (100 μs for numerical reasons). A Boltzmann n^2 model gave similar results to a HH n^2 model (p = 0.96, two-tailed Wilcoxon test) and had fewer parameters, while performing similarly to the unconstrained model (p = 0.79, two-tailed Wilcoxon test). Therefore, we chose the Boltzmann n^2 model. Selection criteria (see above) were passed by n = 16 cells. Statistics of fitted parameters are shown in Table 2.

Fitting results motivated us to further simplify the model by enforcing $V_{Kd} = V_\tau$ and $k_\tau = 2k_{Kd}$. This corresponds to a simple biophysical model where opening and closing rates are of the form $e^{\pm V/k}$. This simplification slightly increases the fit error (1.82 vs. 1.8 mV; p = 0.009, two-tailed Wilcoxon test), but has the advantage of reducing the parameter set to a single kinetic parameter, the maximum time constant (median 3.2 ms).

	Mean	Median	s.d.	s.e.m.
$\mathbf{C}$ (pF)	158.59	145.61	48.76	12.19
$\mathbf{E_K}$ (mV)	-53.83	-56.38	12.35	3.09
$\mathbf{E_L}$ (mV)	-19.68	-18.33	6.48	1.62
$\mathbf{V_{Kd}}$ (mV)	29.77	21.31	22.86	5.71
$\mathbf{V_{\tau_{Kd}}}$ (mV)	26.32	23.09	48.37	12.09
$\mathbf{b_{Kd}}$ (ms)	9.21	4.40	12.69	3.17
$\mathbf{g_L}$ (nS)	10.62	9.72	5.73	1.43
$\mathbf{g_{Kd}}$ (nS)	1130.25	100.11	2656.07	664.02
$\mathbf{k_{Kd}}$ (mV)	8.96	6.61	6.29	1.57
$\mathbf{k_{\tau_{Kd}}}$ (mV)	13.54	11.74	8.08	2.02

Table 2. *Statistics of fitted parameters (n = 16) for deciliated cells (Boltzmann model with two gates).*

Ciliated fits

For Fig. 7, we fitted the complete model to electrophysiological and ciliary responses to two sets of 100 ms pulses, a set of pulses between 0 and 5 nA in 300 pA increments, and a set of pulses between -100 and 500 pA in 25 pA increments. The complete model consisted of I_L, the simplified n^2 Boltzmann model of I_{Kd} with $a_{Kd} = 0.1$ ms, $V_{K_d} = V_\tau^{Kd}$ and $k_\tau^{Kd} = 2k_{Kd}$ (equations (6-7) and (12-13)), I_{Ca}

(equations (14-16)), $I_{K(Ca)}$ (equations (17-18)), calcium dynamics (equation (19)) and electromotor
coupling (equation (20)).

To deal with possible shifts in tip potential between the two sets, we aligned all traces to a resting
potential of -22 mV (the median resting potential), and E_K was fixed at -48 mV (the median estimated
E_K).

The optimization error combined an error on the reading electrode potential, an error on the ciliary
angle (mean angle of the PIV analysis), and an error on resting calcium concentration. For the voltage
error, the response was divided in two equally weighted intervals: from pulse start to pulse end, and
from pulse end to 500 ms after the end (to capture the post-stimulus hyperpolarization). The angle
error was defined as the quadratic error on the corresponding unit vectors, which is equivalent to $E_\alpha =$
$\cos^2(\alpha - \hat{\alpha}) + \sin^2(\alpha - \hat{\alpha})$, and applied on the interval from 100 ms before the pulse to 500 ms after
it. Finally, we ensured that the resting [Ca^{2+}] was 0.1 μM by inserting an error on [Ca^{2+}] on the interval
from 100 ms before the pulse to the start of the pulse ($E_{Ca} = ([Ca^{2+}] - 0.1\ \mu M)^2$). This effectively
ensures $J = I_{rest}/Fv_{\text{cilia}}$.

Selection criteria (see above) were passed by n = 18 cells. Statistics of fitted parameters are shown in
Table 3.

1174

	Mean	Median	s.d.	s.e.m.
C (pF)	303.06	289.10	67.81	15.98
E$_L$ (mV)	-23.07	-23.30	1.18	0.28
J (1/s)	946.72	866.02	422.76	99.65
V$_{Ca}$ (mV)	0.47	-1.33	6.03	1.42
V$_{Kd}$ (mV)	9.55	4.19	12.48	2.94
λ (1/s)	19.43	7.79	32.31	7.62
b$_{Kd}$ (ms)	4.77	4.50	2.58	0.61
g$_{Ca}$ (nA)	434.81	226.63	413.79	97.53
g$_{K(Ca)}$ (nS)	3919.39	216.51	6499.27	1531.89
g$_L$ (nS)	10.38	9.52	4.78	1.13
g$_{Kd}$ (nS)	821.80	114.76	1469.93	346.47
k$_{Ca}$ (mV)	4.34	4.35	0.97	0.23
k$_{Kd}$ (mV)	5.74	4.89	2.97	0.70
n$_{Ca}$	4.41	4.25	1.66	0.39
n$_{K(Ca)}$	3.53	2.94	2.35	0.55
n$_{motor}$	7.37	7.02	5.70	1.34
pK$_{Ca}$	3.60	3.61	0.40	0.10
pK$_{K(Ca)}$	6.37	6.79	1.50	0.35
pK$_{motor}$	2.64	3.14	1.87	0.44
τ$_m$ (ms)	0.91	0.94	0.38	0.09

1175 **Table 3.** *Statistics of fitted parameters (n = 16) for depolarized ciliated cells.*

1176

Statistics

Fitting results obtained with different models were compared using the Wilcoxon test. Statistics are
given as mean $\pm$ standard deviation. In box plots, the box shows the first and third quartile with the
median value inside, and the whiskers are the minimum and maximum values excluding outliers,
which are shown as diamonds and defined as those at a distance exceeding 1.5 times the interquartile
range from the box.

1183

1184 **Hydrodynamic model**

1185 In Fig. 8F, we calculated the motion vectors from patterns of ciliary beating on a sphere of 60 μm radius.
1186 The velocity vector $\boldsymbol{U}$ and the rotation vector $\boldsymbol{\Omega}$ are given by:

1187
$$\begin{pmatrix} \boldsymbol{U} \\ \boldsymbol{\Omega} \end{pmatrix} = \begin{pmatrix} \boldsymbol{M} & \boldsymbol{N} \\ \boldsymbol{N^T} & \boldsymbol{O} \end{pmatrix} \begin{pmatrix} \boldsymbol{F} \\ \mathbf{L} \end{pmatrix}$$

1188 where $\mathbf{F}$ is the external force and $\mathbf{L}$ is the external torque (Lauga and Powers, 2009). The matrix is
1189 called the *mobility* matrix. For a sphere of radius r, the mobility matrix is diagonal:

1190
$$\boldsymbol{U} = (6\pi\eta r)^{-1}\boldsymbol{F}$$

1191
$$\boldsymbol{\Omega} = (8\pi\eta r^3)^{-1}\boldsymbol{L}$$

We consider that each patch of membrane is subjected to a force from the fluid, in the direction
opposite to the ciliary beating direction, and we calculate the total force and torque for different ciliary
beating patterns.

We use spherical coordinates (θ, ϕ) (see Fig. 12), where $\theta = 0$ corresponds to the North pole,
considered as the anterior end, and $\phi = 0$ is the meridian corresponding to the oral groove. Thus, a
surface element is:

1198
$$dS = r^2 \sin\theta \, d\theta d\phi$$

1199 The local force is tangent to the sphere and oriented with an angle α, where $\alpha = 0$ is the direction of
1200 the meridian, pointing South. Cartesian coordinates are chosen so that the x axis is dorso-ventral, the
1201 y axis is oriented left to right, and the z axis is posterior to anterior (i.e., South to North pole).

1202

1203 **Figure 12.** *Conventions on the spherical model, with spherical coordinates and a local force **F** on the surface*
1204 *of the sphere.*

1205 With these conventions, the local force expressed in Cartesian coordinates is:

1206
$$\boldsymbol{F} = A \begin{bmatrix} \cos\theta\cos\alpha\cos\phi - \sin\alpha\sin\phi \\ \cos\theta\cos\alpha\sin\phi + \sin\alpha\cos\phi \\ -\sin\theta\cos\alpha(\theta,\phi) \end{bmatrix}$$

1207 where A is the amplitude of the force per unit area and α is its angle (to obtain this result, start from
1208 the North pole, rotate along the y axis by θ, then rotate along the z axis by α). Therefore, the total force
1209 is:

1210
$$\boldsymbol{F}_{tot} = r^2 \int_{-\pi}^{\pi}\int_{0}^{\pi} \boldsymbol{F}(\theta,\phi)\sin\theta\, d\theta d\phi$$

1211 The local torque is $\boldsymbol{\tau} = \boldsymbol{r} \times \boldsymbol{F}$ where $\boldsymbol{r}$ is a radius. In Cartesian coordinates, we obtain:

1212
$$\boldsymbol{\tau} = rA \begin{bmatrix} -\cos\theta\sin\alpha\cos\phi - \cos\alpha\sin\phi \\ -\cos\theta\sin\alpha\sin\phi + \cos\alpha\cos\phi \\ \sin\theta\sin\alpha \end{bmatrix}$$

1213
$$\boldsymbol{\tau}_{tot} = r^2 \int_{-\pi}^{\pi}\int_{0}^{\pi} \boldsymbol{\tau}(\theta,\phi)\sin\theta\, d\theta d\phi$$

The reported velocity is $v = \|\boldsymbol{U}\|$, the angle of the rotation axis θ_{rotation} is calculated in the xz plane, and
the spinning speed is $\omega = \frac{\|\boldsymbol{\Omega}\|}{2\pi}$ in cycle/s. Local force amplitude is identical in all ciliary beating patterns,
and chosen so as to obtain a forward velocity of 500 µm/s.

Three different patterns are represented in Fig. 8F. In the forward pattern, the local angle is uniform:
$\alpha = $ -170°, corresponding to a beating direction of 10°, downward to the right. In the backward pattern,
the local angle is $\alpha = $ -10°, corresponding to a beating direction of 170°, upward to the right (obtained
by up/down symmetry). In both cases, the local force is axisymmetrical with respect to the

anteroposterior axis. It follows that both $\boldsymbol{F}_{tot}$ and $\boldsymbol{\tau}_{tot}$ are aligned with the z axis (anteroposterior).
This can be seen in the formulas above by integrating with respect to ϕ, which yields 0 for the x and y
coordinates.

In the turning pattern, the left anterior quarter ($\phi \in [-\pi, 0]$, $\theta \in [0, \frac{\pi}{2}]$) follows the forward pattern
(α = -170°), while the right anterior quarter ($\phi \in [0, \pi]$, $\theta \in [0, \frac{\pi}{2}]$) follows the backward pattern ($\alpha =$
-10°) and the posterior half ($\theta \in [\pi/2, \pi]$) has local forces pointing to the left ($\alpha =$ -90°), meaning cilia
beating to the right. The posterior half generates a rotating pattern around the main axis (by
axisymmetry), without translational movement. Each anterior quarter generates forces and torques

1229
$$\boldsymbol{F}_{tot} = \begin{bmatrix} 2\sin\alpha \\ -\cos\alpha \\ -\frac{\pi}{4}\cos\alpha \end{bmatrix}$$

1230
$$\boldsymbol{\tau}_{tot} = r \begin{bmatrix} 2\cos\alpha \\ \sin\alpha \\ \frac{\pi}{4}\sin\alpha \end{bmatrix}$$

1231   With $\alpha_L = \pi - \alpha_R$, we then find that the y and z components of the total torque vanish. Added to the
1232   torque generated by the posterior part, we obtain a torque vector in the xz plane, which is the plane of
1233   the oral groove, separating the cell in left and right parts.

1234

Kinematics

We consider that the organism is an object moving by rigid motion. The organism is characterized by
a position vector **x**, and an orientation matrix **R** defining the rotation of the reference frame (frame of
the organism), so that a point **y** on the reference frame is mapped to **Ry** in the observer frame. The
reference frame is chosen as in the spherical model above, so that $z>0$ points towards the anterior end
while the x axis the dorsoventral axis.

Translational velocity is assumed to be in the posterior-anterior direction only: $\boldsymbol{v} = [0, 0, v]$, so that

1242
$$\dot{\boldsymbol{x}} = \boldsymbol{R}\boldsymbol{v}$$

1243   Let $\boldsymbol{\omega}$ be the rotation vector in the reference frame. We assume it is tilted from the main axis by an
1244   angle θ in the xz plane:

1245
$$\boldsymbol{\omega} = -\omega[\sin(\theta), 0, \cos(\theta)]$$

1246   Thus, kinematics is determined by three variables v, ω and θ, which are functions of calcium
1247   concentration, as detailed in *Electrophysiological modeling (subsection Electromotor coupling)*.

1248   Over a time *dt*, the organism rotates by $\boldsymbol{\Omega}(dt) = \boldsymbol{I} + [\boldsymbol{\omega}].dt$, where

1249
$$[\boldsymbol{\omega}] = \omega \begin{bmatrix} 0 & -\cos(\theta) & 0 \\ \cos(\theta) & 0 & -\sin(\theta) \\ 0 & \sin(\theta) & 0 \end{bmatrix}$$

1250   is the infinitesimal rotation matrix, such that $[\boldsymbol{\omega}]\boldsymbol{y} = \boldsymbol{\omega} \times \boldsymbol{y}$. Therefore, the orientation matrix
1251   changes as $\boldsymbol{R}(t + dt) = \boldsymbol{R}\boldsymbol{\Omega}(dt)$, giving:

1252
$$\dot{R} = R[\boldsymbol{\omega}]$$

1253 For numerical reasons, we use quaternions instead of matrices (Graf, 2008), implemented with the
1254 Python packages *quaternion* and *pyquaternion*. Orientation is then represented by a unit quaternion q,
1255 and kinematic equations translate to:

1256
$$\dot{\boldsymbol{x}} = q\hat{\boldsymbol{v}}\bar{q}$$

1257 where $\hat{\boldsymbol{v}}$ is the pure quaternion with imaginary part **v**, and

1258
$$\dot{q} = \frac{1}{2}q\hat{\boldsymbol{\omega}}$$

1259

1260 *Confinement to a plane*

1261 We constrain the organism to move in a plane. This is done simply by rotating the orientation vector
1262 at each time step so that it lies in the plane. Concretely, we calculate the orientation vector in the
1263 observer frame:

1264
$$\hat{\boldsymbol{p}} = q(\widehat{0,0,1})\bar{q}$$

1265 Then we rotate the orientation vector around the axis **u** that is orthogonal to both the z axis and **p**:
1266 $\boldsymbol{u} = (0,0,1) \times \boldsymbol{p}$, by an angle $\theta = \sin^{-1}(\boldsymbol{p}_z/\|\boldsymbol{p}\|)$. The final orientation quaternion is then
1267 $q' = Q(\boldsymbol{u}, \theta)q$, where $Q(\boldsymbol{u}, \theta)$ is the rotation of axis **u** and angle θ.

1268

1269 **Behavioral scenarios**

1270 In all simulations, the electrophysiological model fitted to the cell shown in Fig. 7 is integrated with
1271 Euler method and a time step of 0.1 ms, using Brian 2 (Stimberg et al., 2019). Kinematics were
1272 integrated with a time step of 1 ms for simulations with stimuli with sharp boundaries (Figs. 7, 8, 9A
1273 and 8B) and 2 ms for simulations with spatially continuous stimuli (Figs. 9C and 9D).

1274

1275 *Avoiding reaction*

1276 In Fig. 9D, models are simulated in a plane with 2 ms pulse currents of amplitude 0.3, 0.5 and 5 nA, at
1277 1 second intervals. In Fig. 7F, the stimulus is a 2 ms current pulse of 0.01 to 10 nA amplitude. In Fig.
1278 7G, the stimulus is a 100 pA pulse of duration 0 to 100 ms. The reorientation angle is calculated as the
1279 change in angle before and after the stimulus, averaged over 1 second (the spinning period).

1280

Interaction with a stimulus

In Fig. 10 and all subsequent figures, trajectories are constrained to a plane. In Fig. 10A and B, the
stimulus is a half-plane. It produces an instantaneous depolarizing current, proportional to the fraction
of the cell shape inside the stimulus (see *Sensory transduction*), with maximum 5 nA. In Fig. 10C and D,
the stimulus additionally goes through a low-pass filter with time constant 40 ms, representing the
activation/deactivation rate of the receptors.

1287

Repelling and attracting discs

In Fig. 11A and 11B, the stimulus is a disc of radius 1 mm within a 4 mm torus. 100 trajectories are
simulated for 20 s, with random initial positions. A noisy current is added to the membrane equation
so as to produce spontaneous action potentials at the observed rate of 0.2 Hz. It is modeled as an
Ornstein-Uhlenbeck process:

$$\tau_n \frac{dI_n}{dt} = -I_n + \sigma_n \sqrt{\tau_n} \xi$$

with $\tau_n = 20$ ms and $\sigma_n = 9$ pA. Physiologically, this corresponds to the random opening of K^+
channels (Moolenaar et al., 1976). This noise is included in all subsequent simulations.

In Fig. 11A, the stimulus produces a depolarizing current with a 40 ms time constant, as in Fig. 8C and
D. In Fig. 9B, the stimulus produces an adapting hyperpolarizing current (see *Sensory transduction;*
τ_{fast} = 40 ms and τ_{slow} = 200 ms), with a maximum amplitude of 1 nA.

1299

1300 *Gradient following*

1301 In Fig. 11C, the environment has the topology of a cylindrical surface, i.e., circular in the small
1302 dimension (500 μm) and linear in the long dimension (25 mm). The stimulus is a linear gradient of 100
1303 pA/mm, with transduction modeled with adaptation as for the attracting disc (Fig. 9B).

1304

Collective behavior

In Fig. 11D, cells produce CO_2 by respiration, which then diffuses and acidifies the fluid. This is
modelled by the diffusion equation:

$$\frac{\partial S}{\partial t} = \alpha \cdot \mathbb{1}_{(x,y) \in \text{cell}} + D \Delta S$$

where S is the transduction current triggered by CO_2, α is the production rate and D is the diffusion
coefficient of CO_2. In water at 25°C, $D \approx 0.002$ mm^2/s but we accelerate it by a factor 5 to speed up the
simulation. The production rate is α =100 pA/s in a square pixel of width 20 μm.

1312

1313 **Code and data availability**

1318

1319 **Acknowledgments**

We thank Marcel Stimberg for technical assistance and Eric Meyer for providing us with specimens of *P. tetraurelia*.

This work was supported by Agence Nationale de la Recherche (ANR-20-CE30-0025-01 and ANR-21-CE16-0013-02), Programme Investissements d'Avenir IHU FOReSIGHT (Grant ANR-18-IAHU-01), Fondation Pour l'Audition Grant FPA RD-2017-2, CNRS (Défi Mécanobiologie, project PERCEE), and Sorbonne Université (Emergence, project NEUROSWIM).

Competing interests

The authors have no competing interests to declare.

References

Aubusson-Fleury A, Cohen J, Lemullois M. 2015. Ciliary heterogeneity within a single cell: the Paramecium model. *Methods Cell Biol* **127**:457–485. doi:10.1016/bs.mcb.2014.12.007

Berg HC. 1975. Bacterial behaviour. *Nature* **254**:389–392. doi:10.1038/254389a0

Berke AP, Turner L, Berg HC, Lauga E. 2008. Hydrodynamic attraction of swimming microorganisms by surfaces. *Phys Rev Lett* **101**:038102. doi:10.1103/PhysRevLett.101.038102

Brehm P, Eckert R, Tillotson D. 1980. Calcium-mediated inactivation of calcium current in Paramecium. *J Physiol* **306**:193–203.

Brette R. 2021. Integrative Neuroscience of Paramecium, a "Swimming Neuron." *eNeuro* **8**:ENEURO.0018-21.2021. doi:10.1523/ENEURO.0018-21.2021

Brette R, Piwkowska Z, Monier C, Rudolph-Lilith M, Fournier J, Levy M, Frégnac Y, Bal T, Destexhe A. 2008. High-resolution intracellular recordings using a real-time computational model of the electrode. *Neuron* **59**:379–91. doi:S0896-6273(08)00539-4

Bullington WE. 1930. A further study of spiraling in the ciliate Paramecium, with a note on morphology and taxonomy. *J Exp Zool* **56**:423–449. doi:10.1002/jez.1400560404

Chad J, Eckert R, Ewald D. 1984. Kinetics of calcium-dependent inactivation of calcium current in voltage-clamped neurones of Aplysia californica. *J Physiol* **347**:279–300.

Cohen N, Denham JE. 2019. Whole animal modeling: piecing together nematode locomotion. *Curr Opin Syst Biol*, • Systems biology of model organisms • Systems ecology and evolution **13**:150–160. doi:10.1016/j.coisb.2018.12.002

Cohen N, Sanders T. 2014. Nematode locomotion: dissecting the neuronal–environmental loop. *Curr Opin Neurobiol*, Theoretical and computational neuroscience **25**:99–106. doi:10.1016/j.conb.2013.12.003

Drescher K, Goldstein RE, Michel N, Polin M, Tuval I. 2010. Direct Measurement of the Flow Field around Swimming Microorganisms. *Phys Rev Lett* **105**:168101. doi:10.1103/PhysRevLett.105.168101

Dryl S. 1973. Chemotaxis in Ciliate ProtozoaBehaviour of Micro-Organisms. Springer US. pp. 16–30. doi:10.1007/978-1-4684-1962-7_2

Dupre C, Yuste R. 2017. Non-overlapping Neural Networks in Hydra vulgaris. *Curr Biol* **27**:1085–1097.

doi:10.1016/j.cub.2017.02.049

Eckert R. 1972. Bioelectric Control of Ciliary Activity. *Science* **176**:473–481.
doi:10.1126/science.176.4034.473

Eckert R, Brehm P. 1979. Ionic mechanisms of excitation in Paramecium. *Annu Rev Biophys Bioeng* **8**:353–
383. doi:10.1146/annurev.bb.08.060179.002033

Eckert R, Chad JE. 1984. Inactivation of Ca channels. *Prog Biophys Mol Biol* **44**:215–267.
doi:10.1016/0079-6107(84)90009-9

Eckert R, Naitoh Y. 1970. Passive electrical properties of Paramecium and problems of ciliary
coordination. *J Gen Physiol* **55**:467–483.

Gallois B, Candelier R. 2021. FastTrack: An open-source software for tracking varying numbers of
deformable objects. *PLOS Comput Biol* **17**:e1008697. doi:10.1371/journal.pcbi.1008697

Graf B. 2008. Quaternions and dynamics. *ArXiv08112889 Math-Ph*.

Hamel A, Fisch C, Combettes L, Dupuis-Williams P, Baroud CN. 2011. Transitions between three swimming
gaits in Paramecium escape. *Proc Natl Acad Sci U S A* **108**:7290–7295. doi:10.1073/pnas.1016687108

Hansma HG. 1974. Biochemical studies on the behavioral mutants of Paramecium aurelia: ion fluxes and
ciliary membrane proteins (PhD Thesis). University of California, Santa Barbara.

Hennessey TM, Kung C. 1985. Slow inactivation of the calcium current of Paramecium is dependent on
voltage and not internal calcium. *J Physiol* **365**:165–179. doi:10.1113/jphysiol.1985.sp015765

Hennessey TM, Kung C. 1984. An anticalmodulin drug, W-7, inhibits the voltage-dependent calcium
current in Paramecium caudatum. *J Exp Biol* **110**:169–181.

Hennessey TM, Rucker WB, McDiarmid CG. 1979. Classical conditioning in paramecia. *Anim Learn Behav*
7:417–423. doi:10.3758/BF03209695

Hille B. 2001. Ion Channels of Excitable Membranes. Sinauer Associates.

Höfer GF, Hohenthanner K, Baumgartner W, Groschner K, Klugbauer N, Hofmann F, Romanin C. 1997.
Intracellular Ca2+ inactivates L-type Ca2+ channels with a Hill coefficient of approximately 1 and an
inhibition constant of approximately 4 microM by reducing channel's open probability. *Biophys J*
73:1857–1865. doi:10.1016/S0006-3495(97)78216-X

Hook C, Hildebrand E. 1979. Excitation of paramecium. *J Math Biol* **8**:197–214. doi:10.1007/BF00279722

Houten JV. 1978. Two mechanisms of chemotaxis inParamecium. *J Comp Physiol* **127**:167–174.
doi:10.1007/BF01352301

Husser MR, Hardt M, Blanchard M-P, Hentschel J, Klauke N, Plattner H. 2004. One-way calcium spill-over
during signal transduction in Paramecium cells: from the cell cortex into cilia, but not in the reverse
direction. *Cell Calcium* **36**:349–358. doi:10.1016/j.ceca.2004.02.003

Iftode F, Cohen J, Ruiz F, Rueda AT, Chen-Shan L, Adoutte A, Beisson J. 1989. Development of surface
pattern during division in Paramecium. I. Mapping of duplication and reorganization of cortical
cytoskeletal structures in the wild type. *Development* **105**:191–211.

Iwadate Y. 2003. Photolysis of caged calcium in cilia induces ciliary reversal in Paramecium caudatum. *J*
Exp Biol **206**:1163–1170. doi:10.1242/jeb.00219

Iwadate Y, Katoh K, Asai H, Kikuyama M. 1997. Simultaneous recording of cytosolic Ca2+ levels
inDidinium andParamecium during aDidinium attack onParamecium. *Protoplasma* **200**:117–127.
doi:10.1007/BF01283288

Iwadate Y, Kikuyama M. 2001. Contribution of Calcium Influx on Trichocyst Discharge in Paramecium
caudatum. *Zoolog Sci* **18**:497–504. doi:10.2108/zsj.18.497

Iwadate Y, Suzaki T. 2004. Ciliary reorientation is evoked by a rise in calcium level over the entire cilium.
Cell Motil **57**:197–206. doi:10.1002/cm.10165

Iwatsuki K, Naitoh Y. 1983. Photobehavior in a Colorless Paramecium. *BioScience* **33**:714–715.
doi:10.2307/1309353

Iwatsuki K, Naitoh Y. 1982. Photoresponses in colorlessParamecium. *Experientia* **38**:1453–1454.
doi:10.1007/BF01955765

Jana S, Eddins A, Spoon C, Jung S. 2015. Somersault of Paramecium in extremely confined environments.
Sci Rep **5**. doi:10.1038/srep13148

Jennings. 1906. Behavior of the lower organisms. New York, The Columbia university press, The
Macmillan company, agents; [etc., etc.].

Jennings HS. 1906. Behavior of the lower organisms. New York, The Columbia university press, The
Macmillan company, agents; [etc., etc.].

Jennings HS. 1904. The behavior of paramecium. Additional features and general relations. *J Comp Neurol*
Psychol **14**:441–510. doi:10.1002/cne.920140602

Jennings Herbert S. 1899. Studies on reactions to stimuli in unicellular organisms. ii.—the mechanism of
the motor reactions of paramecium. *Am J Physiol-Leg Content* **2**:311–341.
doi:10.1152/ajplegacy.1899.2.4.311

Jennings H. S. 1899. Studies on reactions to stimuli in unicellular organisms. iv.—laws of chemotaxis in
paramecium. *Am J Physiol-Leg Content* **2**:355–379. doi:10.1152/ajplegacy.1899.2.4.355

Jennings HS, Jamieson C. 1902. Studies on Reactions to Stimuli in Unicellular Organisms. X. The
Movements and Reactions of Pieces of Ciliate Infusoria. *Biol Bull* **3**:225–234. doi:10.2307/1535876

Jensen P. 1893. Ueber den Geotropismus niederer Organismen. *Arch Für Gesamte Physiol Menschen Tiere*
53:428–480. doi:10.1007/BF01706283

Jung I, Powers TR, Valles JM. 2014. Evidence for two extremes of ciliary motor response in a single
swimming microorganism. *Biophys J* **106**:106–113. doi:10.1016/j.bpj.2013.11.3703

Klauke N, Plattner H. 1997. Imaging of Ca2+ transients induced in Paramecium cells by a polyamine
secretagogue. *J Cell Sci* **110**:975–983.

Knoll G, Haacke-Bell B, Plattner H. 1991. Local trichocyst exocytosis provides an efficient escape
mechanism for Paramecium cells. *Eur J Protistol* **27**:381–385. doi:10.1016/S0932-4739(11)80256-7

Kulkarni A, Elices I, Escoubet N, Pontani L-L, Prevost AM, Brette R. 2020. A simple device to immobilize
protists for electrophysiology and microinjection. *J Exp Biol* **223**. doi:10.1242/jeb.219253

Kung C. 1971. Genic mutants with altered system of excitation in Paramecium aurelia. II. Mutagenesis,
screening and genetic analysis of the mutants. *Genetics* **69**:29–45.

Kung C, Saimi Y. 1985. Chapter 3 Ca2+ Channels of Paramecium: A Multidisciplinary Study In: Bronner F,
Kleinzeller A, editors. Current Topics in Membranes and Transport, Genes and Membranes: Transport
Proteins and Receptors. Academic Press. pp. 45–66. doi:10.1016/S0070-2161(08)60149-5

Kunita I, Kuroda S, Ohki K, Nakagaki T. 2014. Attempts to retreat from a dead-ended long capillary by
backward swimming in Paramecium. *Front Microbiol* **5**. doi:10.3389/fmicb.2014.00270

Lauga E, Powers TR. 2009. The hydrodynamics of swimming microorganisms. *Rep Prog Phys* **72**:096601.
doi:10.1088/0034-4885/72/9/096601

Lodh S, Yano J, Valentine MS, Van Houten JL. 2016. Voltage-gated calcium channels of Paramecium cilia. *J*
Exp Biol **219**:3028–3038. doi:10.1242/jeb.141234

Lu Z. 2004. Mechanism of rectification in inward-rectifier K+ channels. *Annu Rev Physiol* **66**:103–129.
doi:10.1146/annurev.physiol.66.032102.150822

Machemer H. 1974. Frequency and directional responses of cilia to membrane potential changes
inParamecium. *J Comp Physiol* **92**:293–316. doi:10.1007/BF00696617

Machemer H. 1972. Ciliary Activity and the Origin of Metachrony in Paramecium: Effects of Increased
Viscosity. *J Exp Biol* **57**:239–259.

Machemer H. 1969. Regulation der Cilienmetachronie bei der "Fluchtreaktion" von Paramecium*. *J*
Protozool **16**:764–771. doi:10.1111/j.1550-7408.1969.tb02340.x

Machemer H, Machemer-Röhnisch S. 1984. Mechanical and electric correlates of mechanoreceptor
activation of the ciliated tail inParamecium. *J Comp Physiol A* **154**:273–278. doi:10.1007/BF00604993

Machemer H, Ogura A. 1979. Ionic conductances of membranes in ciliated and deciliated Paramecium. *J*
Physiol **296**:49–60.

Machemer-Röhnisch S, Machemer H. 1989. A Ca paradox: Electric and behavioural responses of
Paramecium following changes in external ion concentration. *Eur J Protistol* **25**:45–59.
doi:10.1016/S0932-4739(89)80077-X

Martinac B, Saimi Y, Kung C. 2008. Ion Channels in Microbes. *Physiol Rev* **88**:1449–1490.
doi:10.1152/physrev.00005.2008

Mendelssohn M. 1895. Ueber den Thermotropismus einzelliger Organismen. *Arch Für Gesamte Physiol*
Menschen Tiere **60**:1–27. doi:10.1007/BF01661667

Moolenaar WH, De Goede J, Verveen AA. 1976. Membrane noise in Paramecium. *Nature* **260**:344–346.
doi:10.1038/260344a0

Nagel U, Machemer H. 2000. Physical and physiological components of the graviresponses of wild-type
and mutant Paramecium Tetraurelia. *J Exp Biol* **203**:1059–1070.

Naitoh Y, Eckert R. 1972. Electrophysiology of Ciliate ProtozoaExp. in Physiol. and Biochem. pp. 17–31.

Naitoh Y, Kaneko H. 1972. Reactivated Triton-Extracted Models of Paramecium: Modification of Ciliary
Movement by Calcium Ions. *Science* **176**:523–524. doi:10.1126/science.176.4034.523

Nakaoka Y, Iwatsuki K. 1992. Hyperpolarization-activated inward current associated with the frequency
increase in ciliary beating of Paramecium. *J Comp Physiol A* **170**:723–727. doi:10.1007/BF00198983

Nakaoka Y, Machemer H. 1990. Effects of cyclic nucleotides and intracellular Ca on voltage-activated

ciliary beating in Paramecium. *J Comp Physiol A* **166**:401–406. doi:10.1007/BF00204813

Noguchi M, Nakamura Y, Okamoto K-I. 1991. Control of ciliary orientation in ciliated sheets from Paramecium–differential distribution of sensitivity to cyclic nucleotides. *Cell Motil* **20**:38–46. doi:https://doi.org/10.1002/cm.970200105

Noguchi M, Ogawa T, Taneyama T. 2000. Control of ciliary orientation through cAMP-dependent phosphorylation of axonemal proteins in Paramecium caudatum. *Cell Motil* **45**:263–271. doi:https://doi.org/10.1002/(SICI)1097-0169(200004)45:4<263::AID-CM2>3.0.CO;2-J

Oami K. 1996. Distribution of chemoreceptors to quinine on the cell surface of Paramecium caudatum. *J Comp Physiol A* **179**:345–352. doi:10.1007/BF00194988

Oertel D, Schein SJ, Kung C. 1978. A potassium conductance activated by hyperpolarization in paramecium. *J Membr Biol* **43**:169–185. doi:10.1007/BF01933477

Oertel D, Schein SJ, Kung C. 1977. Separation of membrane currents using a Paramecium mutant. *Nature* **268**:120–124. doi:10.1038/268120a0

Ogura A. 1981. Deciliation and Reciliation in Paramecium after Treatment with Ethanol. *Cell Struct Funct* **6**:43–50. doi:10.1247/csf.6.43

Ogura A, Machemer H. 1980. Distribution of mechanoreceptor channels in theParamecium surface membrane. *J Comp Physiol* **135**:233–242. doi:10.1007/BF00657251

Ohmura T, Nishigami Y, Taniguchi A, Nonaka S, Manabe J, Ishikawa T, Ichikawa M. 2018. Simple mechanosense and response of cilia motion reveal the intrinsic habits of ciliates. *Proc Natl Acad Sci* **115**:3231–3236. doi:10.1073/pnas.1718294115

Oka T, Nakaoka Y, Oosawa F. 1986. Changes in Membrane Potential during Adaptation to External Potassium Ions in Paramecium Caudatum. *J Exp Biol* **126**:111–117.

Pallasdies F, Goedeke S, Braun W, Memmesheimer R-M. 2019. From single neurons to behavior in the jellyfish Aurelia aurita. *eLife* **8**:e50084. doi:10.7554/eLife.50084

Párducz B. 1967. Ciliary Movement and Coordination in Ciliates In: Bourne GH, Danielli JF, editors. International Review of Cytology. Academic Press. pp. 91–128. doi:10.1016/S0074-7696(08)60812-8

Platkiewicz J, Brette R. 2010. A threshold equation for action potential initiation. *PLoS Comput Biol* **6**:e1000850. doi:10.1371/journal.pcbi.1000850

Plattner H. 2015. Molecular aspects of calcium signalling at the crossroads of unikont and bikont eukaryote evolution--the ciliated protozoan Paramecium in focus. *Cell Calcium* **57**:174–185. doi:10.1016/j.ceca.2014.12.002

Plattner H, Klauke N. 2001. Calcium in ciliated protozoa: sources, regulation, and calcium-regulated cell functions. *Int Rev Cytol* **201**:115–208. doi:10.1016/s0074-7696(01)01003-8

Plattner H, Verkhratsky A. 2018. The remembrance of the things past: Conserved signalling pathways link protozoa to mammalian nervous system. *Cell Calcium* **73**:25–39. doi:10.1016/j.ceca.2018.04.001

Preston RR. 1998. Transmembrane Mg2+ Currents and Intracellular Free Mg2+ Concentration in Paramecium tetraurelia. *J Membr Biol* **164**:11–24. doi:10.1007/s002329900389

Preston RR. 1990. A magnesium current in Paramecium. *Science* **250**:285–288. doi:10.1126/science.2218533

Preston RR, Saimi Y, Kung C. 1992a. Calcium current activated upon hyperpolarization of Paramecium
tetraurelia. *J Gen Physiol* **100**:233–251. doi:10.1085/jgp.100.2.233

Preston RR, Saimi Y, Kung C. 1992b. Calcium-dependent inactivation of the calcium current activated upon
hyperpolarization of Paramecium tetraurelia. *J Gen Physiol* **100**:253–268.

Preston RR, Saimi Y, Kung C. 1990. Evidence for two K+ currents activated upon hyperpolarization
ofParamecium tetraurelia. *J Membr Biol* **115**:41–50. doi:10.1007/BF01869104

Roesle E. 1903. Die Reaktion einiger Infusorien auf einzelne Induktionsschläge. *Z Für Allg Physiol* 139–
168.

Saimi Y. 1986. Calcium-dependent sodium currents inParamecium: Mutational manipulations and effects
of hyper- and depolarization. *J Membr Biol* **92**:227–236. doi:10.1007/BF01869391

Saimi Y, Hinrichsen RD, Forte M, Kung C. 1983. Mutant analysis shows that the Ca2+-induced K+ current
shuts off one type of excitation in Paramecium. *Proc Natl Acad Sci* **80**:5112–5116.
doi:10.1073/pnas.80.16.5112

Saimi Y, Ling KY. 1990. Calmodulin activation of calcium-dependent sodium channels in excised
membrane patches of Paramecium. *Science* **249**:1441–1444. doi:10.1126/science.2169650

Satow Y, Kung C. 1980. Ca-Induced K+-Outward Current in Paramecium Tetraurelia. *J Exp Biol* **88**:293–
304.

Satow Y, Kung C. 1979. Voltage Sensitive Ca-Channels and the Transient Inward Current in Paramecium
Tetraurelia. *J Exp Biol* **78**:149–161.

Schafer WR. 2018. The Worm Connectome: Back to the Future. *Trends Neurosci* **41**:763–765.
doi:10.1016/j.tins.2018.09.002

Sedar AW, Porter KR. 1955. The fine structure of cortical components of Paramecium
multimicronucleatum. *J Biophys Biochem Cytol* **1**:583–604. doi:10.1083/jcb.1.6.583

Standen NB, Stanfield PR. 1982. A Binding-Site Model for Calcium Channel Inactivation that Depends on
Calcium Entry. *Proc R Soc Lond B Biol Sci* **217**:101–110. doi:10.1098/rspb.1982.0097

Stimberg M, Brette R, Goodman DF. 2019. Brian 2, an intuitive and efficient neural simulator. *eLife*
8:e47314. doi:10.7554/eLife.47314

Teska A, Stimberg M, Brette R. 2020. brian2modelfitting. doi:10.5281/zenodo.4601961

Tominaga T, Naitoh Y. 1992. Membrane Potential Responses to Thermal Stimulation and the Control of
Thermoaccumulation in Paramecium Caudatum. *J Exp Biol* **164**:39–53.

Valentine MS, Rajendran A, Yano J, Weeraratne SD, Beisson J, Cohen J, Koll F, Van Houten J. 2012.
ParameciumBBS genes are key to presence of channels in Cilia. *Cilia* **1**:16. doi:10.1186/2046-2530-1-16

Valentine MS, Van Houten J. 2022. Ion Channels of Cilia: Paramecium as a Model. *J Eukaryot Microbiol*
e12884. doi:10.1111/jeu.12884

Wan KY. 2020. Synchrony and symmetry-breaking in active flagellar coordination. *Philos Trans R Soc Lond*
B Biol Sci **375**:20190393. doi:10.1098/rstb.2019.0393

Wang H, Swore JJ, Sharma S, Szymanski JR, Yuste R, Daniel TL, Regnier M, Bosma M, Fairhall AL. 2020.
From neuron to muscle to movement: a complete biomechanical model of Hydra contractile behaviors.

bioRxiv 2020.12.14.422784. doi:10.1101/2020.12.14.422784

Wright MV, van Houten JL. 1990. Characterization of a putative Ca2+-transporting Ca2+-ATPase in the
pellicles of Paramecium tetraurelia. *Biochim Biophys Acta BBA - Biomembr* **1029**:241–251.
doi:10.1016/0005-2736(90)90160-P

Yano J, Rajendran A, Valentine MS, Saha M, Ballif BA, Van Houten JL. 2013. Proteomic analysis of the cilia
membrane of Paramecium tetraurelia. *J Proteomics* **78**:113–122. doi:10.1016/j.jprot.2012.09.040

Yano J, Valentine MS, Van Houten JL. 2015. Novel Insights into the Development and Function of Cilia
Using the Advantages of the Paramecium Cell and Its Many Cilia. *Cells* **4**:297–314.
doi:10.3390/cells4030297

Zack GW, Rogers WE, Latt SA. 1977. Automatic measurement of sister chromatid exchange frequency. *J*
Histochem Cytochem **25**:741–753. doi:10.1177/25.7.70454

Zhang P, Jana S, Giarra M, Vlachos PP, Jung S. 2015. Paramecia swimming in viscous flow. *Eur Phys J Spec*
Top **224**:3199–3210. doi:10.1140/epjst/e2015-50078-x

1561