Strain-dependent induction of primary bile acid 7-dehydroxylation by cholic acid.

Authors

Eduard Vico-Oton[1], Colin Volet[1], Nicolas Jacquemin[1], Karin Lederballe Meibom[1], Rizlan Bernier-
Latmani[1*]

[1]Environmental Microbiology Laboratory, School of Architecture, Civil and Environmental Engineering,
École Polytechnique Fédérale de Lausanne, Lausanne, Switzerland.

*Corresponding author. E-mail address rizlan.bernier-latmani@epfl.ch. Postal address EPFL ENAC IIE
EML CH A1 375 (Bâtiment CH) Station 6 CH-1015 Lausanne Switzerland. Tel. +41 21 69 35001.

Abstract (250 words)

Bile acids (BAs) are steroid-derived molecules with important roles in digestion, the maintenance of
host metabolism and immunomodulation. Primary BAs are synthesised by the host and their
concentrations tightly regulated, while secondary BAs are produced by the gut microbiome through
transformation of the former. Regulation of microbial production of secondary BAs is not well
understood, particularly the production of 7-dehydroxylated BAs, which are the most potent agonists
for host BA receptors. The 7-dehydroxylation of cholic acid (CA) is well established and is linked to the
expression of a bile acid-inducible (*bai*) operon. However, little to no 7-dehydroxylation has been
reported for other host-derived BAs (e.g., chenodeoxycholic acid, CDCA or the mouse ursodeoxycholic
acid, UDCA) and it is unknown whether they induce *bai* gene expression. Here, we demonstrate that
the 7-dehydroxylation of CDCA and UDCA by *Clostridium scindens* is induced by CA suggesting that CA-
dependent transcriptional regulation of 7-dehydroxylation is generalisable to CDCA and UDCA.
Nevertheless, this effect is strain-specific and the murine isolate *Extibacter muris* did not respond to
CA exposure, suggesting that *bai* genes are regulated differently in this strain. Moreover, the accessory
gene *baiJ* was only upregulated in the *Clostridium scindens* ATCC 35704 strain, implying mechanistic
differences amongst isolates. Interestingly, the *C. scindens* strains were also capable of 7-DH-ing
muricholic acids (MCAs) to a limited extent. This study shows novel 7-dehydroxylation activity *in vitro*
as a result of CA-driven induction and suggests that it is likely that a different BA plays the same role
in the murine gut.

Keywords

7-dehydroxylation, *Clostridium scindens*, *Extibacter muris*, gut microbe, deoxycholic acid (DCA),
lithocholic acid (LCA), muricholic acid (MCA), CA co-induction, *bai* gene expression, conjugated bile
acids

Introduction

Primary bile acids (BAs) are metabolites synthesised from cholesterol by hepatocytes while secondary
BAs are produced by the gut microbiome through the transformation of primary BAs (Figure 1). In the
liver, the BA are conjugated to glycine or taurine. The three main microbial BA transformations are
deconjugation (loss of the amino acid group), oxidation (of one or several of the hydroxyl groups), and
7α-dehydroxylation (7-DH-ion), the loss of a hydroxyl group at the C7 position[1]. These microbial
transformations increase the diversity of the BA pool (Figure 1) and determine BA affinity to host
receptors. In particular, 7-DH-ion turns primary BAs such as cholic acid (CA) and chenodeoxycholic acid
(CDCA) into the 7-dehydroxylated (7-DH-ed) BAs deoxycholic acid (DCA) and lithocholic acid (LCA),
respectively[2].

BAs act as detergents to solubilise dietary fats, but also have important metabolic and
immunomodulatory roles through activation of their target receptors[3]. The two best-studied BA
receptors are the Farnesoid X Receptor (FXR), a nuclear receptor, and the G Protein-Coupled Bile Acid
Receptor, GPBAR1, also known as Takeda G-Protein Receptor 5 (TGR5), which is a membrane receptor.
FXR is activated through the binding of BA agonists, particularly the 7-DH-ed LCA but also DCA[4]. FXR
activation results in the inhibition of primary BA synthesis through repression of the cholesterol 7α-
hydroxylase CYP7A1. Regulation of BA production limits BA concentration and therefore, toxicity. On
the other hand, BA dysregulation can cause health issues such as cholestasis, Irritable Bowel Syndrome
(IBS), gallstone disease, or even the induction of colon cancer[5–7]. Besides BA homeostasis, FXR also has
focal roles in glucose and lipid homeostasis[8]. Similarly, TGR5 is a multifunctional regulator involved in
glucose homeostasis, energy expenditure, and the modulation of the inflammatory response[9,10]. LCA,
DCA and their tauro-conjugated forms TLCA and TDCA, are among the strongest agonists of TGR5[11,12],
highlighting the importance of microbial transformation, particularly 7-DH-ion, in TGR5 activation.
Moreover, DCA and LCA have protective properties against *Clostridium difficile* infection[13,14].

The study of BAs has traditionally been based on mouse models[15]. Besides CA and CDCA, mice (and
other rodents) also generate muricholic acids (MCAs) such as α-MCA and β-MCA (Figure 1) and re-
hydroxylate DCA and LCA in the liver[16]. Additionally, the mouse liver is capable of producing primary
ursodeoxycholic acid (UDCA)[16] although there is evidence that the gut microbiome is responsible for
a significant fraction of UDCA in the gut[14] (Figure 3 in manuscript provided for reviewers). On the other
hand, UDCA is exclusively a secondary BA in humans[7]. Because 7-DH-ion plays a major role in host
homeostasis, significant effort has been expended to study 7-dehydroxylating (7-DH-ing) bacteria.

Nonetheless, experimental evidence of 7-DH-ion is limited to a few species of the *Clostridiales* order.
One of the best characterised is the human isolate *Clostridium scindens* ATCC 35704, the type strain
of *C. scindens*[17].The *Extibacter muris* DSM 28560 (JM40) strain was recently isolated from mice and
identified as a 7-DH-ing organism[18,19]. The ability of *C. scindens* ATCC 35704 to 7-dehydroxylate both
in vivo and *in vitro* is well established[20]; *E. muris* DSM 28560 has also been shown to 7-dehydroxylate
in vivo, transforming primary BAs CA, CDCA, αMCA, βMCA and UDCA into their respective secondary
BAs DCA, LCA and MDCA (Figures 3 and 5 in manuscript provided for reviewers).

The biochemical machinery for 7-DH-ion is encoded in the *bai* (bile acid inducible) eight-gene operon
(*baiBCDEA2FGHI*)[21]. In addition, the *C. scindens* ATCC 35704 strain harbours the accessory gene *baiJ*
(HDCHBGLK_03451)[22] whereas the *E. muris* DSM 28650 genome includes a *baiJKL* pseudogene
cluster[23]. Most of the published work on the 7-DH-ion pathway has been performed with another *C.
scindens* strain, VPI 12708[24,25]. A recent publication by Funabashi *et al.* cloned the *bai* operon of *C.
scindens* VPI 12708 into *Clostridium sporogenes* which then showed *in vivo* 7-DH-ing activity[25].
Different strains exhibit varying efficiency in 7-DH-ing CA *in vitro*. The *C. scindens* ATCC 35704 and VPI
12708 strains show almost complete transformation to DCA while *E. muris* DSM 28560 has more
limited activity[19,20,26]. Other known 7-DH-ing strains such as *Clostridium hylemonae* and
Peptacetobacter hiranonis have been reported as harbouring weak and strong activity, respectively[27]
and a new strain of *P. hiranonis* recently isolated from dog faeces displayed *in vitro* 7-DH-ion at around
30% conversion of CA to DCA[28]. Notably, *in vitro* 7-DH-ion of other primary BAs has been reported to
be minor (CDCA) or non-existent (MCAs and UDCA)[20,23,29].

The limited *in vitro* 7-DH-ion of primary BAs other than CA (i.e., CDCA and MCAs) is striking considering
that secondary 7-DH-ed forms of these BAs are routinely detected at significant concentrations in the
host[30,31]. Most studies tackling *in vitro* primary bile acid 7-DH-ion consider each BA in isolation. In
addition, significant overexpression of the *bai* operon in response to CA has been reported[32,33] but

there is no information about the potential induction of this operon by other primary BAs (i.e., CDCA
and MCAs) and whether induction by CA also results in the transformation of the latter. We
hypothesise that CA-dependent induction of the *bai* operon promotes 7-DH-ing activity of other BAs
when they occur together with CA. Moreover, we posit that primary BAs other than CA cannot induce
their own transformation.

Here, the expression of *bai* genes was measured *in vitro* in the presence of CA, CDCA, αMCA, βMCA
and UDCA, with and without co-induction with CA to test whether the overexpression of *bai* genes
was exclusive to CA and whether CA-induced overexpression was sufficient to promote the 7-DH-ion
of other BAs. The experiments were performed with three strains, the human isolates *C. scindens* ATCC
35704 and VPI 12708 and the murine isolate *E. muris* DSM 28561 (SJ24). The results show that the
response to CA co-induction was strain-variable. It was highly effective for *C. scindens* strains and
sufficient to promote the transformation of other primary BAs*.

This work highlights the importance of the presence CA for the 7-DH-ion of other BAs. Moreover, the
lack of activity from *E. muris* SJ24 points at potential host-related differences whereby CA may not be
the key inducer for BA 7-DH-ion in the murine gut.

Results

In vitro bile acid transformation and impact of ^{13}C-CA co-induction

Two human isolates *C. scindens* ATCC 35704 and *C. scindens* VPI 12708 and one murine isolate *E. muris*
DSM 28561 (SJ24) were tested for their ability to 7-dehydroxylate human and mouse primary BAs *in*
vitro. Moreover, parallel experiments were performed by amending the cultures with ^{13}C-CA in order
to test whether the 7-DH-ion of CDCA, αMCA, βMCA and UDCA could be induced by CA
(Supplementary Table 1). As expected, all three strains 7-DH-ed CA but to varying extents (Figure 2).
C. scindens ATCC 35704 and *C. scindens* VPI 12708 showed strong 7-DH-ing activity with 97% and 80%
CA conversion to 7-DH-ed BAs after 48 hours, respectively. *E. muris* SJ24, on the other hand, only
converted 9% of the CA provided into 7-DH-ed forms (Figure 2). Similar results were obtained when
this experiment was repeated as part of the uninduced batch of conditions (Supplementary Figure 1).
C. scindens ATCC 35704 produced up to 52.26 μM DCA after 32 hours. Some of the DCA was
subsequently oxidised to 12-oxolithocholic acid (12-oxoLCA) (Figure 2A). In contrast, *C. scindens* VPI
12708 produced the highest amount of DCA after 48 hours (71.42 μM), with little to no oxidised DCA
forms (0.19 μM of 3-oxoDCA at 48 hours) (Figure 2B). The lack of 12-oxo forms from the *C. scindens*
VPI 12708 strain was expected since the 12α-hydroxysteroid dehydrogenase (12α-HSDH) required for
this process was not detected by PCR in this strain (the full genome is currently unavailable) (data not
shown). Finally, *E. muris* SJ24 only produced 8.1 μM of DCA after 48 hours with very low amounts of
oxidised forms of DCA (0.21 μM of 12-oxoLCA) (Figure 2C). It is important to highlight that *E. muris*
does not possess a 3α-HSDH encoded by *baiA2* which was recently identified as an important
component of the CA 7-DH-ion pathway[25]. This strain may perform hydroxyl group oxidations at the
C3 position using BaiA1/3 which has lower affinity to CA than BaiA2[34] and is present outside the *bai*
operon[35]. All three strains also showed a modicum of 7-oxidation activity (Figure 2), resulting in the
formation of 7-oxoDCA, which cannot be 7-DH-ed.

CDCA 7α-dehydroxylation is limited for all three strains. Indeed, *C. scindens* ATCC 35704 only produced
1.59 μM LCA and *C. scindens* VPI 12708 only 1.55 μM LCA (Figure 3AB), whereas for *E. muris*, no LCA
was detected. The latter is in line with previous reports[23]. The amendment of ^{13}C-CA significantly
increased the transformation of CDCA for both *C. scindens* strains (*p*-value < 0.001 two-way ANOVA)
but had no impact on *E. muris* SJ24 (Figure 3C). Indeed, the LCA yield increased to 9.77 μM for strain

ATCC 35704 and to 40.4 μM for strain VPI 12708 (Figure 3AB). No change was observed for *E. muris*
SJ24.

As for UDCA, 7α-dehydroxylation to LCA was observed only upon amendment with ^{13}C-CA (Figure 4).
None of the strains exhibited any detectable level of activity from cultures that included only UDCA.
14.92 μM of LCA as well as extremely low amounts of 3-oxoLCA (with a maximum of 0.12 μM at 32
hours) were detected when *C. scindens* ATCC 35704 was co-induced. An unknown oxidised form
labelled X-oxoUDCA was detected with a maximum concentration of 2.41 μM after 32 hours (Figure
4A). It is likely that this BA corresponds to 3-oxoUDCA (3-oxo-7β-Hydroxy-5β-cholan-24-oic acid) as
we can exclude 7-oxoLCA (the other product of oxidation of UDCA) (Figure 1). Another BA with the
same ionised mass as UDCA was detected at a maximum concentration of 4.24 μM after 24 hours. We
propose that this could be an isoform of UDCA with the hydroxyl group of the C3 carbon in β
conformation (3β,7β-Dihydroxy-5β-cholan-24-oic acid). However, the identity of these compounds
remains unconfirmed due to the lack of standards. Upon co-induction, the 7-DH-ing activity of *C.*
scindens VPI 12708 was comparable to that of the co-induced ATCC strain, with 13.61 μM of LCA and
0.68 μM of 3-oxoLCA after 48 hours. X-oxoUDCA was also detected at very small concentrations
around 0.5 μM from 12 hours until the end of the experiment. The potential isoform of UDCA was
detected at up to 9.92 μM at the 24-hour time point. Following the same trend observed with the
other primary BAs, *E. muris* SJ24 did not show any detectable activity with UDCA even after co-
induction. The chromatograms for the unknown bile acids can be found in Supplementary Figure 2.

As expected, neither *C. scindens* strain nor *E. muris* were capable of αMCA 7α-dehydroxylation in the
absence of ^{13}C-CA (Figure 5). *C. scindens* ATCC 35704 only produced minute amounts of an unknown
oxo form of αMCA (labelled Y-oxoαMCA) (0.85 μM at 32 hours). Once co-induced with ^{13}C-CA, 6-
oxoMDCA was detected at 2.7 μM after 48 hours in the *C. scindens* ATCC 35704 culture (Figure 5A).
This secondary bile acid has been 7α-DH-ed but also the hydroxyl at C6 oxidised. Moreover, several
intermediates for which standards are unavailable were also detected after 48 hours. These were
unknown oxidised forms of αMCA (labelled X- and Y- oxoαMCA) at concentrations not exceeding 5 μM
each. A third unknown BA was detected (albeit at very low concentrations, 0.41 μM at 32 hours) with
the same mass as 6-oxoMDCA, suggesting that it is an MCA species with one oxidation and one
dehydroxylation. This would indicate the production of another 7α-dehydroxylated form of αMCA *in*
vitro (Figure 1, Figure 5A). As for the ATCC 35704 strain, *C. scindens* VPI 12708 exhibited an increase
in the quantity of products from αMCA transformation in the presence of ^{13}C-CA relative to its absence
(Figure 5B). This includes the 7-DH-ed BA 6-oxoMDCA that reached a concentration of 8.18 μM after
48 hours and the X- and Y- αMCA forms that were detected at maximum concentrations of 4.22 μM
(4 hours) and 1.69 μM (32 hours), respectively. The aforementioned αMCA-derived bile acid with one
ketone group and one dehydroxylation was also detected at a maximum concentration of 3.49 μM
after 48 hours (Figure 5B). Surprisingly, *E. muris* SJ24 did not exhibit any observable 7-DH-ing activity
with or without ^{13}C-CA. Nevertheless, a small amount of X-oxoαMCA was detected at all time points,
with a stable concentration at around 2.4 μM without and 1.9 μM with ^{13}C-CA (Figure 5C). The results
for βMCA were very similar to those for αMCA and are discussed in further detail in the supplementary
information.

The concentration of ^{13}C-CA was also measured over time to ascertain that CA was being metabolised.
It was observed to decrease until disappearing after 48 hours in the *C. scindens* strains except in the
presence of CDCA, for which the concentration decreased slowly over time. We attribute this
observation to the toxicity of CDCA in that concentration[20]. On the other hand, the concentration of
^{13}C-CA in *E. muris* remained stable over time and in all conditions (Supplementary Figure 3).

Induction of bai genes expression in the presence of primary bile acids

In order to assess whether the amendment of [13]C-CA to the culture induced the *bai* operon as
hypothesised, the relative expression of *baiCD* and of *baiE* were measured. In addition, the expression
of *baiJ* (an accessory gene) was also monitored. Gene expression was normalised using at least three
reference genes and was calculated relative to the expression levels in a control group without BAs.
Both *E. muris* strains, that is JM40 (DSM 28560) and SJ 24 (DSM 28561), have a truncated *baiJ* gene[23].
However, while the *baiJ* pseudogene is interrupted by stop codons in strain JM40, it is not in strain
SJ24, making it worthwhile to investigate *baiJ* in the latter strain (as was done in this study).
Additionally, *baiO* was also analysed for *E. muris* SJ24 as an alternative accessory *bai* gene[36].

Results show that the expression of *bai* operon genes in both *C. scindens* strains was highly
upregulated as a response to exposure to CA or to CDCA but not to the other BAs (Figure 6). CA and
CDCA are also the only two primary BAs for which *in vitro* 7-DH-ion data are already available[20].
Moreover, it is worth highlighting that CDCA was tested using a concentration of 200 µM vs. 100 µM
for CA.

For *C. scindens* ATCC 35704, the three genes tested were highly upregulated when [13]C-CA was present
along with another BA (UDCA, αMCA, or βMCA) (Figure 6A). In the CDCA dataset, statistically
significant differences relative to the uninduced conditions were observed and all genes were slightly
more upregulated in the presence of [13]C-CA (p-value < 0.001 linear model), but *baiCD* was more so
than the other genes (Figure 6A). Most interestingly, none of the other primary BAs activated the
expression of *bai* genes on their own, consistent with the lack of 7-DH-ing activity with these BA
substrates alone. However, the expression was brought up to levels higher than those observed for
CA once co-induced (Figure 6A).

A similar pattern was observed for *C. scindens* VPI 12708 but with the significant difference being that
baiJ was not overexpressed under any conditions (Figure 6B) (p-value < 0.001 linear model). In the
CDCA dataset, the co-induction with [13]C-CA had an upregulatory effect if assessed with a paired
Wilcoxon test (Supplementary Figure 4) but this effect was not found to be significant when using the
linear statistical model displayed in Figure 6. It is worth noting that the expression data were obtained
from the mid- to late-log exponential phase (around 18 hours) when differences in activity between
uninduced and co-induced conditions were not very large (Figure 3B). As with *C. scindens* ATCC 35704,
co-induction had a dramatic effect on the expression levels of *baiCD* and *baiE* in the presence of UDCA,
αMCA, or βMCA, with upregulation reaching the expression levels observed with CA or CDCA alone
(Figure 6B).

E. muris SJ24 showed a slight upregulation of *baiCD* and *baiE* in the CA dataset but it was not significant
and did not translate to any of the other conditions (Figure 6C), mimicking its very poor 7-DH-ing
activity (Figure 2). As a matter of fact, the increased *baiCD* and *baiE* gene expression ratio observed in
the CA group was probably caused by one of the biological replicates which had a higher expression
level than the others.

Thus, CA had a large effect on *bai* expression which was strain specific. Genes of the *bai* operon in the
two *C. scindens* strains (ATCC 35704 and VPI 12708) exhibited a similar response to CA induction but
the accessory *baiJ* differed in its response. It was upregulated in strain ATCC 35704 but not in strain
VPI 12708. In contrast, CA had no significant effect on the expression of all the *bai* genes considered
in *E. muris* SJ24.

The *rhaS*_1 gene (HDCHBGLK_01429) is immediately upstream of the *bai* operon promoter on the
opposite strand and has also been proposed as <u>b</u>ile <u>a</u>cid-<u>r</u>egulatory (*bar*) due to its potential

implication in *bai* regulation[7]. The expression of *rhaS1* and *rhaS2* (a copy of *rhaS1* elsewhere in the
genome) was shown to have background levels across BAs (Supplementary Figure 5). This was tested
in *C. scindens* ATCC 35704 without the amendment of ^{13}C-CA. Results indicate that *rhaS* is not
upregulated by any of the tested BAs.

Discussion

Bile acid chemistry is a relevant field in human and veterinary medicine not only because of BA
detergent function during digestion but also for the wide range of roles related to host physiology and
homeostasis[37,38].

The *bai* operon was originally described in *C. scindens* VPI 12708 in 1990 and this strain has become
the reference for biochemical studies of the 7-DH-ion pathway[21,35,39]. However, genetic differences
occur in the *bai* operon amongst 7-DH-ing strains[35,40]. Moreover, the presence of accessory *bai* genes
with as-of-yet undefined roles varies across strains[35,39].

Out of all primary BAs, only CA has been consistently reported to be 7-DH-ed *in vitro*, with low levels
of 7-DH-ion for CDCA and none for other murine primary BAs (αMCA, βMCA or UDCA)[20,23,41–44]. We
confirm that the two *C. scindens* human isolates upregulated the expression of *bai* operon genes in
response to CA (Figure 6). However, we observed differences between the two strains in the response
of *baiJ*. This accessory gene is annotated as an urocanate reductase and a member of the
oxidoreductase family. Current pathway data indicate that the gene does not play a significant role in
7-DH-ion of CA in strain VPI 12708[25], which would align with the lack of expression observed from our
data. In contrast, the high levels of upregulation in the ATCC 35704 strain match previously published
data[32] and suggest that *baiJ* may play a role in the 7-DH-ion of CA or other BAs in that strain. On the
other hand, *E. muris* SJ24, the murine isolate, barely registered any response to CA, nor activity in
general, highlighting the potential differences in the 7-DH-ion pathway between human and mouse
isolates.

UDCA is the 7β isomer of CDCA and is reported to have significant therapeutic properties[2,7,45]. In
humans, UDCA is synthesised from CDCA by gut microbes containing 7α- and 7β-HSDHs[7,26,46,47] and is
thus, a secondary BA; in mice, it is produced by the liver, thus, it is a primary BA, and is used as a
precursor for βMCA[15,16]. Nevertheless, colonisation of mice lacking any 7-dehydroxylators with 7-DH-
ing bacteria increases UDCA levels, implying that the gut microbiome also plays a significant role in
the UDCA production of mice[14] (Studer *et al.,* and appended manuscript Figures 3 and 5). Despite its
importance, little is known about the capacity of bacteria to 7β-dehydroxylate UDCA *in vitro* and
previous data have shown no activity by *C. scindens* strain ATCC 35704[20]. In accordance with our
hypothesis, co-induction with ^{13}C-CA not only greatly upregulated *bai* gene expression (Figure 6B) but
also provided evidence of UDCA 7β-dehydroxylation as LCA was detected in the culture (Figure 4). The
absence of CDCA in the culture suggests that UDCA was not 7-epimerised to CDCA prior to its 7α-
dehydroxylation. Moreover, two unidentified compounds were detected, providing further evidence
of the 7β-dehydroxylation of UDCA and its associated unique set of intermediates. The oxidised
intermediate is very likely to be 3-oxoUDCA (3-oxo-7β-Hydroxy-5β-cholan-24-oic acid), as based on
the mass, the only alternative would have been 7-oxoLCA, which was quantified (Figure 4). The other
unknown compound had the same mass as UDCA albeit a different retention time, suggesting this
could be an iso- form of UDCA. It follows that this would be 3β-UDCA, tentatively named isoUDCA
(3β,7β-Dihydroxy-5β-cholan-24-oic acid). Isoforms are well known in the BA pool and particularly in
CDCA-related intermediates[20,25,33,48] so it is likely that UDCA follows a similar pattern. Despite the
production of unknown intermediates, the activity response to co-induction suggests that UDCA 7β-
dehydroxylation uses the same *bai* machinery as the 7-DH-ion of CDCA.

The BA pool of mice is significantly more diverse than that of humans due to the primary production
of muricholic acids (with a hydroxyl group at the C-6 position) and of UDCA. Their secondary bile acid
pool includes DCA and LCA but also MDCA and its 6α counterpart, hyodeoxycholic acid (HDCA).
Furthermore, mice can rehydroxylate TDCA back into TCA in the liver[15]. In general, the secondary BAs
derived from muricholic acids seem to be in low abundance in the gut, hinting at the difficulty in 7-
DH-ing these BAs. This is perhaps the reason why primary BAs such as βMCA are dominant in the bile
acid pool[49]. While we initially hypothesised that the lack of activity was due to the lack of expression,
our co-induced data show that *bai* gene expression in the human isolates is not sufficient for the
production of MDCA. [23]. On the other hand, *E. muris* showed no significant expression.

The human isolates showed marginal activity for αMCA and βMCA upon co-induction. 7-DH-ed forms
such as 6-oxoMDCA were detected (Figure 5 & Supplementary Figure 6) but the full 7-dehydroxylation
to MDCA was not observed. These results imply that other elements besides the *bai* operon (and *baiJ*)
might be needed to completely 7-dehydroxylate MCAs. A potential reason could be a missing gene
such as a 6β-HSDH required for complete 7-DH-ion.

The mouse isolate *E. muris* was a weak CA 7-dehydroxylator and had no activity with the other BAs.
This was coupled with a complete lack of gene upregulation in response to any BA. Nonetheless,
recently published *in vivo* data from Studer *et al.,* 2022 (appended manuscript Figure 3) quantified
MDCA as a direct result of *E. muris* colonisation, demonstrating that this strain is capable of 7-DH-ion
of MCA *in vivo*. To explain these differences, it is possible that an untested BA could be relevant for
inducing 7-DH-ion of MCAs. An option could be TβMCA, but conjugated forms were not tested here.
Both of these BAs are highly abundant in the BA pool of mice[16]. Finally, experimental conditions might
have not been optimal to stimulate the 7-DH-ing activity[50].

The novel activity for αMCA and βMCA yielded several intermediate BAs that we were not able to fully
characterise due to the lack of appropriate standards. All human and mouse BAs share a backbone of
four rings (Figure 1), this makes mass fractionation in the mass spectrometer unsuitable for
identification. Therefore, we currently rely on comparison of ionised mass and retention time to
standards. However, several assumptions can be made to speculate what these compounds could be.
MCAs oxidised at the C-3, C-6 or C-7 position (Figure 1)*.* A C-7 oxidation would yield 7-oxoMDCA
regardless of the primary MCA. Meanwhile, other oxidations would be differentiated by the α or β
conformation of the C-7 hydroxyl. It is possible that one of the intermediates that we detected was 7-
oxoMDCA but none shared retention times across MCAs, meaning that different intermediates were
produced for each of the two MCA substrates (Supplementary Figure 2). Considering that two oxidised
intermediates are detected from both MCAs, we hypothesise that those are the 3-oxo and 6-oxo forms
of αMCA and βMCA (3-oxoαMCA: 3-oxo-6β,7α-Dihydroxy-5β-cholan-24-oic acid; 6-oxoαMCA: 6-oxo-
3α,7α-Dihydroxy-5β-cholan-24-oic acid; 3-oxoβMCA: 3-oxo-6β,7β-Dihydroxy-5β-cholan-24-oic acid;
and 6-oxoβMCA: 6-oxo-3α,7β-Dihydroxy-5β-cholan-24-oic acid). A third intermediate was also
detected from both MCAs, with the ionised mass corresponding to secondary BAs with one
dehydroxylation and a ketone group (e.g., 7-oxoLCA) (Figure 5 & Supplementary Figure 6). Three
options are plausible: 1) A dehydroxylation at the C-3 position. This would yield a novel family of BAs
with a 6β- and 7α/β- hydroxyls which is highly unlikely as it would have been identified previously by
the multiple studies investigating the murine BA pool[15,16,51–54]. 2) An oxidation paired with a 6-
dehydroxylation would yield 3-oxoCDCA, 7-oxoLCA, or 3-oxoUDCA. 3-oxoCDCA and 7-oxoLCA were
included in our analysis (Supplementary Table 3) and would have been detected if present.
Meanwhile, the retention time of this compound is distinct from that of the compound proposed to
be 3-oxoUDCA from the transformation of UDCA (see above and Supplementary Figure 2). Thus, this
is not likely to be 3-oxoUDCA. 3) A 7-dehydroxylation could allow for ketone groups at the C-3 and C-

6 positions. 6-oxoMDCA was available as a standard but the other option, 3-oxoMDCA (3-oxo-6β-
Hydroxy-5β-cholan-24-oic acid), was not. It is therefore possible that this compound corresponds to
3-oxoMDCA, but this remains unconfirmed.

Conclusion

The findings presented here are threefold. First and foremost, we demonstrated that the previously
reported[32,33] strong upregulation of *bai* genes by CA increases the extent of 7-DH-ion of other primary
BAs. This was particularly true for UDCA which has been reported to be converted to LCA *in vitro* for
the first time. Secondly, the upregulation of *bai* genes exhibited strain-specific differences. *C. scindens*
ATCC 35704 upregulated the *bai* operon genes *baiCD* and *baiE* as well as the *bai* accessory gene *baiJ*.
While for *C. scindens* VPI 12708, it upregulated the *bai* operon genes, but *baiJ* expression was found
to be at a background level. This is consistent with the lack of involvement of *baiJ* in 7-DH-ion in strain
VPI 12708[25]. *E. muris* SJ24 was the third strain tested, a murine isolate with *in vivo* 7-DH-ing capabilities
(appended manuscript Figure 3 and 5). Our findings with the SJ24 strain showed weak 7-DH-ion of CA
and no upregulation of any of the *bai* genes tested. The lack of activity of this murine strain even in
the presence of murine primary BAs suggests that there is a missing factor, maybe an untested BA or
other co-factors, required for an efficient *bai* expression and 7-DH-ing activity. It is also possible that
7-DH-ion of MCAs occurs through a distinct pathway with different enzymatic requirements. Finally,
C. scindens human isolates can partially 7-dehydroxylate MCAs. Nevertheless, there might be missing
a key component such as a 6-βHSDH that would complete the reductive arm of 7-DH-ion to MDCA.

In conclusion, these data provide novel insights into the intricacies of 7-DH-ion and the significant
differences amongst 7-DH-ing bacteria. The CA-dependent co-induction can be attributed to the
abundance of this compound in the BA pool of humans and mice. However, the inducing factor for *E.*
muris activity remains elusive. Moreover, multiple novel BAs were observed and their identity
surmised based on mass spectrometry information. Further work is required to investigate the 7-DH-
ion pathway of CDCA, αMCA, βMCA and UDCA as well as exploring the strain-specific differences
regarding the 7-DH-ion pathway of CA.

Materials and Methods

Bacterial strains and growth conditions

The strains used were *Clostridium scindens* ATCC 35704, *Clostridium scindens* VPI 12708 and *Extibacter*
muris DSM 28561 (SJ24). Bacteria were grown in $\underline{B}$rain $\underline{H}$eart $\underline{I}$nfusion $\underline{S}$upplement – $\underline{S}$alts (BHI-S)
medium, consisting of 37g BHI, 1g L-cysteine, 5g yeast extract, 2.5g fructose, 50mL salts solution (0.2g
$CaCl_2$, 0.2 $MgSO_4$, 1g K_2HPO_4, 1g KH_2PO_4, 10g $NAHCO_3$ and 2g NaCl per L of ddH_2O) per L ddH_2O. The
salts solution and media were sterilised by autoclaving. Static growth was carried out at 37°C in an
anaerobic chamber (Coy Laboratory Products, 95% N_2, 5% H_2). A pre-inoculum was prepared from
glycerol stocks (using BHIS-S) before inoculating 25 mL of BHIS-S in Falcon tubes at a starting OD_{600} of
0.05. Results of the growth curves for the experiments can be found in the Supplementary
Information.

In vitro 7-dehydroxylation assays

Bacteria were grown in presence of BAs: CA (100 μM), CDCA (200 μM), αMCA (100 μM), βMCA (100
μM) and UDCA (100 μM) or the same volume of ethanol (solvent control). A sterile control (media
with ethanol solvent) was also included. Co-induction was performed by adding an additional 100 μM
of ^{13}C-CA to CDCA, UDCA, αMCA, or βMCA). ^{13}C-CA was chosen so its transformation products could
be separated from those of the other primary BAs during the quantification process. All uninduced

experiments were conducted at the same time while all the co-induced ones were run simultaneously at a different time. Both experiments included a condition consisting of the amendment of only 100 µM CA, which allows for comparison of the results from co-induced and uninduced experiments. Growth was monitored by periodically measuring the OD_{600}. During the main time points (0, 4, 8, 12, 24, 32 and 48 hours), 1mL samples were collected for BA extraction in a 2 mL bead-beating resistant tube. All conditions were performed in triplicates.

Bile acid extraction

Samples were vacuum dried overnight (ON). Approximately 450 mg of 0.5 mm zirconium beads were added to the dried samples as well as 500 µL ice-cold alkaline acetonitrile (acetonitrile – 25% ammonia 4:1 v/v) and 100 µL of ISTD solution (CA-d$_4$, CDCA-d$_4$, TCA-d$_4$, TUDCA-d$_4$, DCA-d$_4$ and LCA-d$_4$, each at 100 µM in methanol). Samples were homogenised in a Precellys 24 Tissue Homogenizer (Bertin Instruments, Montigny-le-Bretonneux, France) at 6500 rpm 3x 30" beat 30" rest. Samples were vortexed for 1 hour and centrifugated for 15 minutes at 16000 rcf at room temperature. Approximately 500 µL of suspension was carefully collected over the beads level and transferred into a new 1.5 mL epi tube which was then vacuum dried overnight. Finally, the samples were reconstituted in 1 mL of ammonium acetate [5mM] – methanol (50:50 v/v) and a 1:20 dilution with the same solvent was prepared in LC-MS glass vials, ready for injection.

RNA extraction and reverse transcription

1 mL of sample was collected in a 15 mL falcon tube during the mid-log to late-log phase for the RNA extraction. The sample was stored with RNAprotect following the manufacturer protocol (Protocol 5 from RNAprotect Bacteria Reagent Handbook 01/2020, Qiagen) at -80°C until processed. All conditions were performed in triplicates. Lysis and RNA purifications were done using the RNeasy Mini Kit (Qiagen, Hilden, Germany). Bacterial lysis was performed following Protocol 5: Enzymatic Lysis, Proteinase K Digestion and Mechanical Disruption of Bacteria (RNAprotect Bacteria Reagent Handbook 01/2020, Qiagen), with 20 µL of proteinase K for each sample and the required volumes for a number of bacteria <7.5 x 10^8. The cell lysis was performed using a Precellys 24 Tissue Homogenizer (Bertin Instruments, Montigny-le-Bretonneux, France) at 6500 rpm 3x 10 seconds beat 10 seconds rest. RNA purification was performed following Protocol 7: Purification of Total RNA from Bacterial Lysate using the RNeasy Mini Kit. Centrifugations were carried out at 15000 rcf except for the 2 min centrifugation which was done at 18000 rcf.

Purified RNA was further subject to a DNase treatment using the RQ DNase I (Promega, Madison, WI, USA) following the manufacturer protocol with small modifications: The final volume was adjusted to a 100 µL and incubation was extended to 1 hour at 37°C. The treated RNA was cleaned-up using the RNeasy Mini Kit (Qiagen, Hilden, Germany) following the RNA Clean-up protocol from the manufacturer (RNeasy Mini Handbook 10/2019) with the 2 min centrifugation done at 18000 rcf. Concentration and purity of RNA was measured with a NanoDrop One (Thermo Fisher Scientific, Waltham, MA, USA).

100 ng of RNA was reverse transcribed into cDNA using the GoScript™ Reverse Transcription Mix, Random Primers (Promega, Madison, WI, USA) following the manufacturer protocol. The process was done in duplicates with one group using water instead of the reaction buffer as a non-reverse transcription control (NRT).

Reverse transcription quantitative PCR (RT-qPCR)

RT-qPCRs were prepared using the Myra liquid handling system (Bio Molecular Systems, software
version 1.6.26) and performed using the Magnetic induction cycler (Mic) platform (Bio Molecular
Systems, Upper Coomera, QLD, Australia) with the micPCR software (v2.10.5).

The list of primers used can be found in Supplementary table 2. Samples were prepared with the
SensiFAST SYBR No-ROX Kit (Meridian Bioscience, Cincinnati, OH, USA) at a final volume of 10 µL. All
runs were performed with the following program, with small modifications: Initial hold at 95°C for 5
minutes with a cycle of 95°C for 5 seconds, 54.5°C for 20 seconds (54.1°C for *E. muris* SJ24) and 72°C
for 9 seconds. 40 cycles were done for *C. scindens* ATCC 35704 and 50 for *C. scindens* VPI 12708 and
E. muris SJ24. The melting curve, temperature control and acquisition settings were left as default.
The quantification was done using three or more reference genes (Supplementary Table 2) based on
their expression stability across conditions. NRTs as well as no template controls (NTCs) were included
to check for residual DNA or contaminations. Four technical replicates were done for each biological
triplicate, totalling at 12 data points per condition. Note that expression data presented in Figure 6 for
the CA-only condition (labelled CA) correspond to the pooled expression results (for the condition in
which only 100 µM Ca was added) from the two sets of experiments presented above (referred to as
the co-induced and the uninduced experiments, respectively).

Liquid chromatography – mass spectrometry (LC-MS)

The quantitative method was performed on an Agilent ultrahigh-performance liquid chromatography
1290 series coupled in tandem to an Agilent 6530 Accurate-Mass Q-TOF mass spectrometer. The
separation was done on a Zorbax Eclipse Plus C18 column (2.1 x 100mm, 1.8 µm) and a guard column
Zorbax Eclipse Plus C18 (2.1 x 5mm, 1.8 µm) both provided by Agilent technologies (Santa Clara, CA,
USA). The column compartment was kept heated at 50°C. Two different solutions were used as
eluents: ammonium acetate [5mM] in water as mobile phase A and pure acetonitrile as mobile phase
B. A constant flow of 0.4 mL/min was maintained over 26 minutes of run time with the following
gradient (expressed in eluent B percentage): 0-5.5 min, constant 21.5% B; 5.5-6 min, 21.5-24.5% B; 6-
10 min, 24.5-25% B; 10-10.5 min, 25-29% B; 10.5-14.5 min, isocratic 29% B; 14.5-15 min, 29-40% B;
15-18 min, 40-45% B; 18-20.5 min, 45-95% B; 20.5-23 min, constant 95% B; 23-23.1 min, 95-21.5% B;
23.10-26 min, isocratic 21.50% B. The system equilibration was implemented at the end of the
gradient for 3 minutes in initial conditions. The autosampler temperature was maintained at 10°C and
the injection volume was 5µL. The ionisation mode was operated in negative mode for the detection
using the Dual AJS Jet stream ESI Assembly. The QTOF acquisition settings were configured in 4GHz
high-resolution mode (resolution 17000 FWHM at m/z 1000), data storage in profile mode and the
high-resolution full MS chromatograms were acquired over the range of m/z 100-1700 at a rate of 3
spectra/s. The mass spectrometer was calibrated in negative mode using ESI-L solution from Agilent
technologies every 6 hours to maintain the best possible mass accuracy. Source parameters were
setup as follows: drying gas flow, 8 L/min; gas temperature, 300°C; nebulizer pressure, 35psi; capillary
voltage, 3500V; nozzle voltage, 1000V. Data were processed afterwards using the MassHunter
Quantitative software and MassHunter Qualitative software to control the mass accuracy for each
run. In the quantitative method, 42 bile acids were quantified by calibration curves (Supplementary
Table 3). The quantification was corrected by addition of internal standards in all samples and
calibration levels. Extracted ion chromatograms were generated using a retention time window of ±
1.5 min and a mass extraction window of ± 30ppm around the theoretical mass of the targeted bile
acid. Unknown BAs were identified when found within the retention time window of a standard with
the same ionised mass. Approximate quantification of these unknown BAs was done by using the
nearest standard (by retention time) with the same ionised mass.

Statistical analysis and data visualisation

Graphpad Prism 9.2.0 (GraphPad) was used to generate the figures shown in this paper and perform pairwise comparisons, the gene expression data was analysed with a linear model (LM) in R language v4.1.2[55] using RStudio[56]. The statistical significance boundary was stablished at a *p*-value < 0.05.

Conflicts of interest

The authors declare no conflicts of interest.

Data Availability Statement

The data used for this manuscript are publicly available in the following link:

 https://doi.org/10.5281/zenodo.6034320

References

1.   Monte MJ, Marin JJG, Antelo A, Vazquez-Tato J. Bile acids: Chemistry, physiology, and pathophysiology. World J Gastroenterol 2009; 15:804–16. doi: 10.3748/wjg.15.804

2.   Hanafi NI, Mohamed AS, Kadir SHSA, Othman MHD. Overview of bile acids signaling and perspective on the signal of ursodeoxycholic acid, the most hydrophilic bile acid, in the heart. Biomolecules2018; 8. doi: 10.3390/biom8040159

3.   Kuipers F, Bloks VW, Groen AK. Beyond intestinal soap - Bile acids in metabolic control. Nat Rev Endocrinol 2014; 10:488–98. doi: 10.1038/nrendo.2014.60

4.   Wang H, Chen J, Hollister K, Sowers LC, Forman BM. Endogenous bile acids are ligands for the nuclear receptor FXR/BAR. Mol Cell 1999; 3:543–53. doi: 10.1016/S1097-2765(00)80348-2

5.   Bernstein H, Bernstein C, Payne CM, Dvorakova K, Garewal H. Bile acids as carcinogens in human gastrointestinal cancers. Mutat Res - Rev Mutat Res 2005; 589:47–65. doi: 10.1016/j.mrrev.2004.08.001

6.   Joyce SA, Gahan CGM. Bile Acid Modifications at the Microbe-Host Interface: Potential for Nutraceutical and Pharmaceutical Interventions in Host Health. Annu Rev Food Sci Technol 2016; 7:313–33. doi: 10.1146/annurev-food-041715-033159

7.   Ridlon JM, Kang D-J, Hylemon PB, Ridlon JM, Kang D-J, Hylemon PB. Bile salt biotransformations by human intestinal bacteria. J Lipid Res 2006; 47:241–59. doi: 10.1194/jlr.R500013-JLR200

8.   Claudel T, Staels B, Kuipers F. The Farnesoid X receptor: A molecular link between bile acid and lipid and glucose metabolism. Arterioscler. Thromb. Vasc. Biol.2005; 25:2020–31. doi: 10.1161/01.ATV.0000178994.21828.a7

9.   Guo C, Chen WD, Wang YD. TGR5, not only a metabolic regulator. Front. Physiol.2016; 7. doi: 10.3389/fphys.2016.00646

10.   van Nierop FS, Scheltema MJ, Eggink HM, Pols TW, Sonne DP, Knop FK, Soeters MR. Clinical relevance of the bile acid receptor TGR5 in metabolism. Lancet Diabetes Endocrinol 2016; 5:224–33. doi: 10.1016/S2213-8587(16)30155-3

11.   Kawamata Y, Fujii R, Hosoya M, Harada M, Yoshida H, Miwa M, Fukusumi S, Habata Y, Itoh T, Shintani Y, et al. A G protein-coupled receptor responsive to bile acids. J Biol Chem 2003; 278:9435–40. doi: 10.1074/jbc.M209706200

12.   Brighton CA, Rievaj J, Kuhre RE, Glass LL, Schoonjans K, Holst JJ, Gribble FM, Reimann F. Bile acids trigger GLP-1 release predominantly by accessing basolaterally located G protein-coupled bile acid receptors. Endocrinology 2015; 156:3961–70. doi: 10.1210/en.2015-1321

13. Buffie CG, Bucci V, Stein RR, McKenney PT, Ling L, Gobourne A, No D, Liu H, Kinnebrew M, Viale
A, et al. Precision microbiome reconstitution restores bile acid mediated resistance to
Clostridium difficile. Nature 2015; 517:205–8. doi: 10.1038/nature13828

14. Studer N, Desharnais L, Beutler M, Brugiroux S, Terrazos MA, Menin L, Schürch CM, McCoy KD,
Kuehne SA, Minton NP, et al. Functional intestinal bile acid 7α-dehydroxylation by *Clostridium*
scindens associated with protection from *Clostridium difficile* infection in a gnotobiotic mouse
model. Front Cell Infect Microbiol 2016; 6. doi: 10.3389/fcimb.2016.00191

15. Li J, Dawson PA. Animal models to study bile acid metabolism. Biochim. Biophys. Acta - Mol.
Basis Dis.2019; 1865:895–911. doi: 10.1016/j.bbadis.2018.05.011

16. Honda A, Miyazaki T, Iwamoto J, Hirayama T, Morishita Y, Monma T, Ueda H, Mizuno S,
Sugiyama F, Takahashi S, et al. Regulation of bile acid metabolism in mouse models with
hydrophobic bile acid composition. J Lipid Res 2020; 61:54–69. doi: 10.1194/jlr.RA119000395

17. Morris GN, Winter J, Cato EP, Ritchie AE, Bokkenheuser VD. *Clostridium scindens* sp. nov., a
Human Intestinal Bacterium with Desmolytic Activity on Corticoids. Int J Syst Bacteriol 1985;
35:478–81. doi: 10.1099/00207713-35-4-478

18. Lagkouvardos I, Pukall R, Abt B, Foesel BU, Meier-Kolthoff JP, Kumar N, Bresciani A, Martínez I,
Just S, Ziegler C, et al. The Mouse Intestinal Bacterial Collection (miBC) provides host-specific
insight into cultured diversity and functional potential of the gut microbiota. Nat Microbiol
2016; 1:1–15. doi: 10.1038/nmicrobiol.2016.131

19. Streidl T, Kumar N, Suarez LN, Rohn S, Clavel T. *Extibacter*. In: Bergey's Manual of Systematics
of Archaea and Bacteria. American Cancer Society; 2019. page 1–7.doi:
10.1002/9781118960608.gbm01749

20. Marion S, Studer N, Desharnais L, Menin L, Escrig S, Meibom A, Hapfelmeier S, Bernier-Latmani
R. *In vitro* and *in vivo* characterization of *Clostridium scindens* bile acid transformations. Gut
Microbes 2019; 10:481–503. doi: 10.1080/19490976.2018.1549420

21. Mallonee DH, White WB, Hylemon PB. Cloning and sequencing of a bile acid-inducible operon
from *Eubacterium* sp. strain VPI 12708. J Bacteriol 1990; 172:7011–9.

22. Ridlon JM, Hylemon PB. Identification and characterization of two bile acid coenzyme A
transferases from *Clostridium scindens*, a bile acid 7α-dehydroxylating intestinal bacterium . J
Lipid Res 2012; 53:66–76. doi: 10.1194/jlr.m020313

23. Streidl T, Karkossa I, Segura Muñoz RR, Eberl C, Zaufel A, Plagge J, Schmaltz R, Schubert K, Basic
M, Schneider KM, et al. The gut bacterium *Extibacter muris* produces secondary bile acids and
influences liver physiology in gnotobiotic mice. Gut Microbes 2021; 13:1–21. doi:
10.1080/19490976.2020.1854008

24. Wells JE, Hylemon PB. Identification and Characterization of a Bile Acid 7α-Dehydroxylation
Operon in *Clostridium* sp. Strain TO-931, a Highly Active 7α-Dehydroxylating Strain Isolated
from Human Feces. Appl Environ Microbiol 2000; 66:1107–13. doi: 10.1128/AEM.66.3.1107-
1113.2000

25. Funabashi M, Grove TL, Wang M, Varma Y, McFadden ME, Brown LC, Guo C, Higginbottom S,
Almo SC, Fischbach MA. A metabolic pathway for bile acid dehydroxylation by the gut
microbiome. Nature 2020; :1–5. doi: 10.1038/s41586-020-2396-4

26. Macdonald IA, Hutchison DM. Epimerization versus dehydroxylation of the 7α-hydroxyl- group
of primary bile acids: Competitive studies with *Clostridium absonum* and 7α-dehydroxylating
bacteria (*Eubacterium* SP.). J Steroid Biochem 1982; 17:295–303. doi: 10.1016/0022-

4731(82)90203-5

27.   Doerner KC, Takamine F, LaVoie CP, Mallonee DH, Hylemon PB. Assessment of fecal bacteria
529   with bile acid 7α-dehydroxylating activity for the presence of *bai*-like genes. Appl Environ
530   Microbiol 1997; 63:1185–8.

531   28.   Vinithakumari AA, Hernandez BG, Ghimire S, Adams S, Stokes C, Jepsen I, Brezina C, Sahin O, Li
532   G, Tangudu C, et al. A model screening pipeline for bile acid converting anti-*Clostridioides
533   difficile* bacteria reveals unique biotherapeutic potential of *Peptacetobacter hiranonis*. bioRxiv
534   2021; :2021.09.29.462466. doi: 10.1101/2021.09.29.462466

535   29.   Ridlon JM. Enzymology and molecular biology of bile acid 7alpha- and 7beta- dehydroxylation
536   by the intestinal bacteria *Clostridium scindens* and *Clostridium hylemonae*. PhD Thesis Virginia
537   Commonw Univ 2008; :1–333. doi: https://doi.org/10.25772/BN62-H676

538   30.   Makino I, Nakagawa S. Changes in biliary lipid and biliary bile acid composition in patients after
539   administration of ursodeoxycholic acid. J Lipid Res 1978; 19:723–8. doi: 10.1016/s0022-
540   2275(20)41272-6

541   31.   Meir K, Kitsberg D, Alkalay I, Szafer F, Rosen H, Shpitzen S, Avi L Ben, Staels B, Fievet C, Meiner
542   V, et al. Human sterol 27-hydroxylase (CYP27) overexpressor transgenic mouse model.
543   Evidence against 27-hydroxycholesterol as a critical regulator of cholesterol homeostasis. J Biol
544   Chem 2002; 277:34036–41. doi: 10.1074/jbc.M201122200

545   32.   Devendran S, Shrestha R, Alves JMP, Wolf PG, Ly L, Hernandez AG, Méndez-García C, Inboden
546   A, Wiley J, Paul O, et al. *Clostridium scindens* ATCC 35704: Integration of nutritional
547   requirements, the complete genome sequence, and global transcriptional responses to bile
548   acids. Appl Environ Microbiol 2019; 85. doi: 10.1128/AEM.00052-19

549   33.   Ridlon JM, Devendran S, Alves JM, Doden H, Wolf PG, Pereira G V., Ly L, Volland A, Takei H,
550   Nittono H, et al. The '*in vivo* lifestyle' of bile acid 7α-dehydroxylating bacteria: comparative
551   genomics, metatranscriptomic, and bile acid metabolomics analysis of a defined microbial
552   community   in   gnotobiotic   mice.   Gut   Microbes   2019;   :1–24.   doi:
553   10.1080/19490976.2019.1618173

554   34.   Bhowmik S, Jones DH, Chiu HP, Park IH, Chiu HJ, Axelrod HL, Farr CL, Tien HJ, Agarwalla S, Lesley
555   SA. Structural and functional characterization of BaiA, an enzyme involved in secondary bile
556   acid synthesis in human gut microbe. Proteins Struct Funct Bioinforma 2014; 82:216–29. doi:
557   10.1002/prot.24353

558   35.   Ridlon JM, Harris SC, Bhowmik S, Kang D-J, Hylemon PB. Consequences of bile salt
559   biotransformations   by   intestinal   bacteria.   Gut   Microbes   2016;   7:22–39.   doi:
560   10.1080/19490976.2015.1127483

561   36.   Heinken A, Ravcheev DA, Baldini F, Heirendt L, Fleming RMTT, Thiele I. Systematic assessment
562   of secondary bile acid metabolism in gut microbes reveals distinct metabolic capabilities in
563   inflammatory bowel disease. Microbiome 2019; 7:75. doi: 10.1186/s40168-019-0689-3

564   37.   Hofmann AF, Hagey LR. Key discoveries in bile acid chemistry and biology and their clinical
565   applications: History of the last eight decades. J. Lipid Res.2014; 55:1553–95. doi:
566   10.1194/jlr.R049437

567   38.   Winston JA, Theriot CM. Diversification of host bile acids by members of the gut microbiota.
568   Gut Microbes2020; 11:158–71. doi: 10.1080/19490976.2019.1674124

569   39.   Ridlon JM, Kang DJ, Hylemon PB. Isolation and characterization of a bile acid inducible 7α-
570   dehydroxylating operon in *Clostridium hylemonae* TN271. Anaerobe 2010; 16:137–46. doi:

10.1016/j.anaerobe.2009.05.004

40.   Song I, Gotoh Y, Ogura Y, Hayashi T, Fukiya S, Yokota A. Comparative genomic and physiological
analysis against *Clostridium scindens* reveals *Eubacterium* sp. C-25 as an atypical deoxycholic
acid producer of the human gut microbiota. Microorganisms 2021; 9:2254. doi:
10.3390/microorganisms9112254

41.   Hylemon PB, Melone PD, Franklund C V., Lund E, Bjorkhem I. Mechanism of intestinal 7α-
dehydroxylation of cholic acid: Evidence that allo-deoxycholic acid is an inducible side-product.
J Lipid Res 1991; 32:89–96.

42.   Masuda N, Oda H, Hirano S, Tanaka H. Enhancement of the 7α-dehydroxylase activity of a
gram-positive intestinal anaerobe by flavins. Appl Environ Microbiol 1983; 45:308–9.

43.   Kitahara M, Takamine F, Imamura T, Benno Y. Assignment of *Eubacterium* sp. VPI 12708 and
related strains with high bile acid 7α-dehydroxylating activity to *Clostridium scindens* and
proposal of *Clostridium hylemonae* sp. nov., isolated from human faeces. Int J Syst Evol
Microbiol 2000; 50:971–8. doi: 10.1099/00207713-50-3-971

44.   Benno Y, Kitahara M, Takamine F, Imamura T. *Clostridium hiranonis* sp. nov., a human intestinal
bacterium with bile acid 7α-dehydroxylating activity. Int J Syst Evol Microbiol 2001; 51:39–44.
doi: 10.1099/00207713-51-1-39

45.   Abdallah E, Emile SH, Elfeki H, Fikry M, Abdelshafy M, Elshobaky A, Elgendy H, Thabet W,
Youssef M, Elghadban H, et al. Role of ursodeoxycholic acid in the prevention of gallstone
formation after laparoscopic sleeve gastrectomy. Surg Today 2017; 47:844–50. doi:
10.1007/s00595-016-1446-x

46.   Coleman JP, Hudson LL, Adams MJ. Characterization and regulation of the NADP-linked 7α-
hydroxysteroid dehydrogenase gene from *Clostridium sordellii*. J Bacteriol 1994; 176:4865–74.

47.   Ferrandi EE, Bertolesi GM, Polentini F, Negri A, Riva S, Monti D. In search of sustainable
chemical processes: Cloning, recombinant expression, and functional characterization of the
7α- and 7β-hydroxysteroid dehydrogenases from *Clostridium absonum*. Appl Microbiol
Biotechnol 2012; 95:1221–33. doi: 10.1007/s00253-011-3798-x

48.   Devlin AS, Fischbach MA. A biosynthetic pathway for a prominent class of microbiota-derived
bile acids. Nat Chem Biol 2015; 11:685–90. doi: 10.1038/nchembio.1864

49.   Sayin SI, Wahlström A, Felin J, Jäntti S, Marschall HU, Bamberg K, Angelin B, Hyötyläinen T,
Orešič M, Bäckhed F. Gut microbiota regulates bile acid metabolism by reducing the levels of
tauro-beta-muricholic acid, a naturally occurring FXR antagonist. Cell Metab 2013; 17:225–35.
doi: 10.1016/j.cmet.2013.01.003

50.   Stellwag EJ, Hylemon PB. 7α-Dehydroxylation of cholic acid and chenodeoxycholic acid by
Clostridium leptum. J Lipid Res 1979; 20:325–33.

51.   Narushima S, Itoh K, Miyamoto Y, Park SH, Nagata K, Kuruma K, Uchida K. Deoxycholic acid
formation in gnotobiotic mice associated with human intestinal bacteria. Lipids 2006; 41:835–
43. doi: 10.1007/s11745-006-5038-1

52.   Eyssen HJ, Parmentier GG, Mertens JA. Sulfated Bile Acids in Germ-Free and Conventional
Mice. Eur J Biochem 1976; 66:507–14. doi: 10.1111/j.1432-1033.1976.tb10576.x

53.   Han J, Liu Y, Wang R, Yang J, Ling V, Borchers CH. Metabolic profiling of bile acids in human and
mouse blood by LC-MS/MS in combination with phospholipid-depletion solid-phase extraction.
Anal Chem 2015; 87:1127–36. doi: 10.1021/ac503816u

54. Straniero S, Laskar A, Savva C, Härdfeldt J, Angelin B, Rudling M. Of mice and men: Murine bile
acids explain species differences in the regulation of bile acid and cholesterol metabolism. J
Lipid Res 2020; 61:480–91. doi: 10.1194/jlr.RA119000307

55. Team RC. R: A language and environment for statistical computing. 2021;

56. RStudio Team (2020). RStudio: Integrated Development for R. 2020;

Figure Legends

Figure 1. List of deconjugated human and rodent bile acids (BAs) discussed here. The characteristic that distinguishes BAs
is the presence of a hydroxyl group at the C-3, C-6, C-7 and/or C-12 position. The hydroxyl groups can be in α- or β-
conformation, oxidised into a ketone group, or fully removed (dehydroxylated BA). CA and CDCA are primary BAs of both
humans and rodents whereas MCAs are exclusively produced by rodents. (*) UDCA is a primary BA in rodents while it is a
secondary BA in humans where the gut microbes epimerise it from CDCA. (†) Bile acids that might have been detected in
this study but for which no standards were available. (‡) Also known as 6-oxolithocholic acid (6-oxoLCA). See
Supplementary Table 3 for full chemical names of the BAs.

Figure 2. *In vitro* 7-dehydroxylation of cholic acid. The 7-dehydroxylation of CA was tested in (A) *Clostridium scindens* ATCC
35704, (B) *C. scindens* VPI 12708 and (C) *Extibacter muris* DSM 28561 (SJ24) over time. All strains were grown anaerobically
in BHIS-S containing 100 µM CA. Bile acids were extracted from the suspended biomass. Error bars represent the standard
deviation of the mean of biological triplicates.

Figure 3. *In vitro* 7α-dehydroxylation of CDCA. The transformation of CDCA into secondary bile acids was tested with and
without co-induction with 100 µM of ¹³C-CA. 200 µM of CDCA were used based on previous experiments by Marion e*t al.*
(A) *Clostridium scindens* ATCC 35704, (B) *C. scindens* VPI 12708 and (C) *E. muris* DSM 28561 (SJ24). Bile acids were extracted
from the suspended biomass. Error bars represent the standard deviation of the mean of biological triplicates.

Figure 4. *In vitro* 7-dehydroxylation of UDCA. The transformation of 100 µM of UDCA into secondary bile acids was tested
with and without co-induction with 100 µM of ¹³C-CA*. (A) *Clostridium scindens* ATCC 35704, (B) *C. scindens* VPI 12708 and
(C) *Extibacter muris* DSM 28561 (SJ24) were grown anaerobically in BHIS-S. Bile acids were extracted from suspended
biomass. Two compounds were detected that could not be identified due to missing standards but their oxidative state
can be estimated based on their ionised mass. X-oxoUDCA had the same mass as other bile acids with one ketone group
and one hydroxyl group. The other unidentified compound had the same mass as UDCA and therefore it is likely to be an
isoform with a 3β conformation. The retention times for these compounds was unique and therefore could not be
identified further. Concentration values of the unknown BAs could only be estimated for the same reason. Error bars
represent the standard deviation of the mean of biological triplicates.

Figure 5. *In vitro* 7-dehydroxylation of αMCA. The transformation of 100 µM of αMCA into secondary bile acids was tested
with and without co-induction with 100 µM of ¹³C-CA*. (A) *Clostridium scindens* ATCC 35704, (B) *C. scindens* VPI 12708 and
(C) *Extibacter muris* DSM 28561 (SJ24) were grown anaerobically in BHIS-S. Bile acids were extracted from suspended
biomass. Several compounds were detected that could not be identified due to missing standards but their oxidative state
can be estimated based on their ionised mass. X- or Y- oxoαMCA had the same mass as other bile acids with one ketone
group and two hydroxyl groups. The other unidentified compound had the same mass as secondary bile acids that have
been dehydroxylated (-1 -OH), have one ketone group and one hydroxyl group. The retention times for these compounds
did not correspond to that of any known standard. Concentration values of the unknown BAs could only be semi-
quantitative for the same reason. Error bars represent the standard deviation of the mean of biological triplicates.

Figure 6. *bai* gene expression in the presence of CA (first panel) or in that of various primary BAs either uninduced or and
co-induced with ¹³C-CA. Data for the CA-only condition correspond to the pooled expression results from the two sets of
experiments (referred to as uninduced and co-induced) that both include this condition, as described in the text. The
expression (normalised to at least three reference genes) is relative to the no BA condition including an equivalent volume
of solvent (ethanol). (A) *Clostridium scindens* ATCC 35704, (B) *C. scindens* VPI 12708 and (C) *Extibacter muris* DSM 28561
(SJ24) gene expression of *baiCD*, *baiE* (part of the *bai* operon), and accessory genes *baiJ* and *baiO* (only in *E. muris* DSM
SJ24) was measured. CA, UDCA, αMCA, βMCA and the co-inducing ¹³C-CA were used at 100 µM, CDCA was used at 200
µM. A detailed view of the *C. scindens* VPI 12708 uninduced vs 13C-CA co-induced in the CDCA group is found in

Supplementary Figure 4. Coloured dots represent the average and error bars represent the standard deviation of 12
replicates. Some error bars may look elongated due to the logarithmic scale of the Y axis. (***) indicates a *p*-value < 0.001
in a linear model analysis comparing the uninduced vs ^{13}C-CA co-induced factor for each *bai* gene.

	Host	Bile Acid name	Abbreviations	C-3	C-6	C-7	C-12
Human and rodent	Primary	Cholic acid	CA	α -OH		α -OH	α -OH
		Chenodeoxycholic acid	CDCA	α -OH		α -OH	
		Ursodeoxycholic acid*	UDCA	α -OH		β -OH	
	secondary	Deoxycholic acid	DCA	α -OH			α -OH
		Lithocholic acid	LCA	α -OH			
		3-oxocholic acid	3-oxoCA	=O		α -OH	α -OH
		3-oxodeoxycholic acid	3-oxoDCA	=O			α -OH
		3-oxochenodeoxycholic acid	3-oxoCDCA	=O		α -OH	
		3-oxolithocholic acid	3-oxoLCA	=O			
		3-oxoursodeoxycholic acid†	3-oxoUDCA	=O		β -OH	
		7-oxodeoxycholic acid	7-oxoDCA	α -OH		=O	α -OH
		7-oxolithocholic acid	7-oxoLCA	α -OH		=O	
		12-oxochenodeoxycholic acid	12-oxoCDCA	α -OH			=O
		12-oxolithocholic acid	12-oxoLCA	α -OH			=O
		Isoursodeoxycholic acid†	isoUDCA	β -OH		β -OH	
Rodent	primary	α-Muricholic acid	αMCA	α -OH	β -OH	α -OH	
		B-Muricholic acid	βMCA	α -OH	β -OH	β -OH	
	secondary	Murideoxycholic acid	MDCA	α -OH	β -OH		
		3-oxo-α-muricholic acid†	3-oxoαMCA	=O	β -OH	α -OH	
		3-oxo-β-muricholic acid†	3-oxoβMCA	=O	β -OH	β -OH	
		3-oxomurideoxycholic acid†	3-oxoMDCA	=O	β -OH		

		6-oxo-α-muricholic acid†	6-oxoαMCA	α -OH	=O	α -OH	
		6-oxo-β-muricholic acid†	6-oxoβMCA	α -OH	=O	β -OH	
		6-oxomurideoxycholic acid‡	6-oxoMDCA	α -OH	=O		
		7-oxomurideoxycholic acid†	7-oxoMDCA	α -OH	β -OH	=O	
		ω-Muricholic acid†	ωMCA	α -OH	α -OH	β -OH	

C. scindens ATCC 35704 uninduced

C. scindens ATCC 35704 ^{13}C-CA co-induced

C. scindens VPI 12708 uninduced

C. scindens VPI 12708 ^{13}C-CA co-induced

E. muris SJ24 uninduced

E. muris SJ24 ^{13}C-CA co-induced

A)

B)

C)

A)

B)

C)