<u>Strain-dependent induction of primary bile acid 7-dehydroxylation by cholic acid.</u>

Authors

Eduard Vico-Oton[1], Colin Volet[1], Nicolas Jacquemin[1], Yuan Dong[2], Siegfried Hapfelmeier[2], Karin
Lederballe Meibom[1], Rizlan Bernier-Latmani[1*]

[1]Environmental Microbiology Laboratory, School of Architecture, Civil and Environmental Engineering,
École Polytechnique Fédérale de Lausanne, Lausanne, Switzerland.

[2]Institute for Infectious Diseases, University of Bern, Bern, Switzerland

*Corresponding author. E-mail address <u>rizlan.bernier-latmani@epfl.ch</u>. Postal address EPFL ENAC IIE
EML CH A1 375 (Bâtiment CH) Station 6 CH-1015 Lausanne Switzerland. Tel. +41 21 69 35001.

Abstract (250 words)

Bile acids (BAs) are steroid-derived molecules with important roles in digestion, the maintenance of
host metabolism and immunomodulation. Primary BAs are synthesised by the host, while secondary
BAs are produced by the gut microbiome through transformation of the former. Regulation of
microbial production of secondary BAs is not well understood, particularly the production of 7-
dehydroxylated BAs, which are the most potent agonists for host BA receptors. The 7-dehydroxylation
of cholic acid (CA) is well established and is linked to the expression of a bile acid-inducible (*bai*) operon
responsible for this process. However, little to no 7-dehydroxylation has been reported for other host-
derived BAs (e.g., chenodeoxycholic acid, CDCA or ursodeoxycholic acid, UDCA). Here, we
demonstrate that the 7-dehydroxylation of CDCA and UDCA by *Clostridium scindens* is induced by CA
suggesting that CA-dependent transcriptional regulation of 7-dehydroxylation is generalisable to
CDCA and UDCA. In contrast, the murine isolate *Extibacter muris* did not respond to CA exposure *in
vitro*, suggesting that *bai* genes are regulated differently in this strain. However, it could 7-
dehydroxylate *in vivo* and its *in vitro* activity was promoted by the addition of cecal content. The
accessory gene *baiJ* was only upregulated in the *Clostridium scindens* ATCC 35704 strain, implying
mechanistic differences amongst isolates. Interestingly, the human-derived *C. scindens* strains were
also capable of 7-dehydroxylating murine bile acids (muricholic acids) to a limited extent. This study
shows novel 7-dehydroxylation activity *in vitro* as a result of CA-driven induction and suggests distinct
bai gene induction mechanisms across bacterial species.

Keywords

7-dehydroxylation, *Clostridium scindens*, *Extibacter muris*, gut microbe, deoxycholic acid (DCA),
lithocholic acid (LCA), muricholic acid (MCA), ursodeoxycholic acid, CA co-induction, *bai* gene
expression, conjugated bile acids

Introduction

Primary bile acids (BAs) are metabolites synthesised from cholesterol by hepatocytes while secondary
BAs are produced by the gut microbiome through the transformation of primary BAs (Figure 1). In the
liver, the BA are conjugated to glycine or taurine. The three main microbial BA transformations are
deconjugation (loss of the amino acid group), oxidation (of one or several of the hydroxyl groups), and
7α-dehydroxylation (7-DH-ion), the loss of a hydroxyl group at the C7 position[1]. These microbial
transformations increase the diversity of the BA pool (Figure 1) and enhance BA affinity to host
receptors. In particular, 7-DH-ion turns primary BAs such as cholic acid (CA) and chenodeoxycholic acid

(CDCA) into the 7-dehydroxylated (7-DH-ed) BAs deoxycholic acid (DCA) and lithocholic acid (LCA),
respectively[2].

BAs act as detergents to solubilise dietary fats, but also have important metabolic and
immunomodulatory roles through activation of their target receptors[3]. The two best-studied BA
receptors are the Farnesoid X Receptor (FXR), a nuclear receptor, and the G Protein-Coupled Bile Acid
Receptor, GPBAR1, also known as Takeda G-Protein Receptor 5 (TGR5), which is a membrane receptor.
FXR is activated through the binding of BA agonists, particularly the 7-DH-ed LCA but also DCA[4]. FXR
activation results in the inhibition of primary BA synthesis through repression of the cholesterol 7α-
hydroxylase CYP7A1. Regulation of BA production limits BA concentration and therefore, toxicity. On
the other hand, BA dysregulation can cause health issues such as cholestasis, irritable bowel
syndrome, gallstone disease, or even the induction of colorectal cancer[5–7]. Besides BA homeostasis,
FXR also has focal roles in glucose and lipid homeostasis[8]. Similarly, TGR5 is a multifunctional regulator
involved in glucose homeostasis, energy expenditure, and the modulation of the inflammatory
response[9,10]. LCA, DCA and their tauro-conjugated forms TLCA and TDCA, are among the strongest
agonists of TGR5[11,12], highlighting the importance of microbial transformation, particularly 7-DH-ion,
in TGR5 activation. Moreover, DCA and LCA have protective properties against *Clostridium difficile*
infection[13,14].

The study of BAs has traditionally been based on mouse models[15]. Besides CA and CDCA, mice (and
other rodents) also generate muricholic acids (MCAs) such as α-MCA and β-MCA (Figure 1) and re-
hydroxylate DCA and LCA in the liver[16]. Additionally, the mouse liver is capable of producing primary
ursodeoxycholic acid (UDCA)[16] although there is evidence that the gut microbiome is responsible for
a significant fraction of UDCA in the gut[14]. On the other hand, UDCA is exclusively a secondary BA in
humans[7]. Because 7-DH-ion plays a major role in host homeostasis, significant effort has been
expended to study 7-dehydroxylating (7-DH-ing) bacteria.

Nonetheless, experimental evidence of 7-DH-ion is limited to a few species of the *Clostridiales* order.
One of the best characterised is the human isolate *Clostridium scindens* ATCC 35704, the type strain
of *C. scindens*[17].The *Extibacter muris* DSM 28560 (JM40) strain was recently isolated from mice and
identified as a 7-DH-ing organism[18,19]. The ability of *C. scindens* ATCC 35704 to 7-dehydroxylate both
in vivo and *in vitro* is well established[20]; *E. muris* has also been shown to 7-dehydroxylate *in vivo*,
transforming primary BAs CA, CDCA, αMCA, βMCA and UDCA into their respective secondary BAs DCA,
LCA and MDCA. Previous research had focused on *E. muris* strain DSM 28560[21] and in this study, we
demonstrate that *E. muris* strain DSM 28561 (SJ24) also has the ability to 7-dehydroxylate (7-DH-ate)
in vivo.

The biochemical machinery for 7-DH-ion is encoded in the *bai* (<u>b</u>ile <u>a</u>cid <u>i</u>nducible) eight-gene operon
(*baiBCDEA2FGHI*)[22]. In addition, the *C. scindens* ATCC 35704 strain harbours the accessory gene *baiJ*
(HDCHBGLK_03451)[23] whereas the *E. muris* DSM 28650 genome includes a *baiJKL* pseudogene
cluster[21]. Most of the published work on the 7-DH-ion pathway has been performed with another *C.*
scindens strain, VPI 12708[24,25]. A recent publication by Funabashi *et al.* cloned the *bai* operon of *C.*
scindens VPI 12708 into *Clostridium sporogenes* which then showed *in vivo* 7-DH-ing activity[25].
Different strains exhibit varying efficiency in 7-DH-ing CA *in vitro*. The *C. scindens* ATCC 35704 and VPI
12708 strains show rapid transformation to DCA while *E. muris* DSM 28560 has more limited
activity[19,20,26]. Other known 7-DH-ing strains such as *Clostridium hylemonae* and *Peptacetobacter*
hiranonis have been reported as harbouring weak and strong activity, respectively[27] and a new strain
of *P. hiranonis* recently isolated from dog faeces displayed *in vitro* 7-DH-ion at around 30% conversion

of CA to DCA[28]. Notably, *in vitro* 7-DH-ion of other primary BAs has been reported to be minor (CDCA)
or non-existent (MCAs and UDCA)[20,21,29].

The limited *in vitro* 7-DH-ion of primary BAs other than CA (i.e., CDCA and MCAs) is striking considering
that secondary 7-DH-ed forms of these BAs are routinely detected at significant concentrations in the
host[30,31]. Most studies tackling *in vitro* primary bile acid 7-DH-ion consider each BA in isolation. In
addition, significant overexpression of the *bai* operon in response to CA has been reported[32,33] but
there is no information about the potential induction of this operon by other primary BAs (i.e., CDCA
and MCAs) and whether induction by CA also results in the transformation of the latter. We
hypothesise that CA-dependent induction of the *bai* operon promotes 7-DH-ing activity of other BAs
when they occur together with CA. Moreover, we posit that primary BAs other than CA cannot induce
their own transformation.

Here, the expression of *bai* genes was measured *in vitro* in the presence of CA, CDCA, αMCA, βMCA
and UDCA, with and without co-induction with CA to test whether the overexpression of *bai* genes
was exclusive to CA and whether CA-induced overexpression was sufficient to promote the 7-DH-ion
of other BAs. The experiments were performed with three strains, the human isolates *C. scindens* ATCC
35704 and VPI 12708 and the murine isolate *E. muris* DSM 28561 (SJ24). The results show that the
response to CA co-induction was strain-variable. It was highly effective for *C. scindens* strains and
sufficient to promote the transformation of other primary BAs*.* For *E. muris*, none of the BAs tested
promoted 7-DH-ion, nevertheless, a positive effect was observed when the bacterium was co-cultured
with a small amount of faecal content from germ-free mice, suggesting that signaling from the host
may be responsible for the induction of 7-DH-ion in *E. muris* SJ24.

This work highlights the importance of the presence CA for the 7-DH-ion of other BAs. Moreover,
results from *E. muris* SJ24 point at host-related differences whereby BAs may not be the key inducers
for BA 7-DH-ion in the murine gut.

Results

In vitro bile acid transformation and impact of [13]C-CA co-induction

Two human isolates *C. scindens* ATCC 35704 and *C. scindens* VPI 12708 and one murine isolate *E. muris*
DSM 28561 (SJ24) were tested for their ability to 7-DH-ate human and mouse primary BAs *in vitro*. As
expected, all three strains 7-DH-ed CA but to varying extents (Figure 2). *C. scindens* ATCC 35704 and
C. scindens VPI 12708 showed strong 7-DH-ing activity with 97% and 80% CA conversion to 7-DH-ed
BAs after 48 hours, respectively. *E. muris* SJ24, on the other hand, only converted 9% of the CA
provided into 7-DH-ed forms (Figure 2). *C. scindens* ATCC 35704 produced up to 52.26 µM DCA after
32 hours with some of the DCA subsequently oxidised to 12-oxolithocholic acid (12-oxoLCA) (Figure
2A). In contrast, *C. scindens* VPI 12708 produced the highest amount of DCA after 48 hours (71.42 µM),
with little to no oxidised DCA forms (0.19 µM of 3-oxoDCA at 48 hours) (Figure 2B). The lack of 12-oxo
forms from the *C. scindens* VPI 12708 strain was expected since the 12α-hydroxysteroid
dehydrogenase (12α-HSDH) required for this process was not detected by PCR in this strain (the full
genome is currently unavailable) (data not shown). Finally, *E. muris* SJ24 only produced 8.1 µM of DCA
after 48 hours with very low amounts of oxidised forms of DCA (0.21 µM of 12-oxoLCA) (Figure 2C). It
is important to highlight that *E. muris* does not possess a 3α-HSDH encoded by *baiA2* which was
recently identified as an important component of the CA 7-DH-ion pathway[25]. BaiA1/3 has lower
affinity to CA than BaiA2[34] and is present outside the *bai* operon[35]. Finally, all three strains also showed
a modicum of 7-oxidation activity (Figure 2), resulting in the formation of 7-oxoDCA, which cannot be
7-DH-ed. Moreover, parallel experiments were performed by amending the cultures with [13]C-CA in

addition to other individual BAs in order to test whether the 7-DH-ion of CDCA, αMCA, βMCA and
UDCA could be induced by CA (Supplementary Table 1). Similar results to above were obtained for the
control (^{13}C-CA only) experiments (Supplementary Figure 1).

CDCA 7α-dehydroxylation is very limited for all three strains. Indeed, *C. scindens* ATCC 35704 only
produced 1.59 µM LCA and *C. scindens* VPI 12708 only 1.55 µM LCA (Figure 3AB), whereas for *E. muris*,
no LCA was detected. The latter is in line with previous reports[21]. The amendment of ^{13}C-CA
significantly increased the transformation of CDCA for both *C. scindens* strains (*p*-value < 0.001 two-
way ANOVA) but had no impact on *E. muris* SJ24 (Figure 3C). Indeed, the LCA yield increased to 9.77
µM for strain ATCC 35704 and to 40.4 µM for strain VPI 12708 (Figure 3AB). No change was observed
for *E. muris* SJ24.

As for UDCA, 7α-dehydroxylation to LCA was observed only upon amendment with ^{13}C-CA (Figure 4).
None of the strains exhibited any detectable level of activity from cultures that included only UDCA.
14.92 µM of LCA as well as extremely low amounts of 3-oxoLCA (with a maximum of 0.12 µM at 32
hours) were detected when *C. scindens* ATCC 35704 was co-induced. An unknown oxidised form
labelled X-oxoUDCA was detected with a maximum concentration of 2.41 µM after 32 hours (Figure
4A). It is likely that this BA corresponds to 3-oxoUDCA (3-oxo-7β-hydroxy-5β-cholan-24-oic acid) as we
can exclude 7-oxoLCA (the other product of oxidation of UDCA) (Figure 1). Another BA with the same
ionised mass as UDCA was detected at a maximum concentration of 4.24 µM after 24 hours. We
propose that this could be an isoform of UDCA with the hydroxyl group of the C3 carbon in the β
conformation (3β,7β-dihydroxy-5β-cholan-24-oic acid). However, the identity of these compounds
remains unconfirmed due to the lack of standards. Upon co-induction, the 7-DH-ing activity of *C.
scindens* VPI 12708 was comparable to that of the co-induced ATCC strain, with 13.61 µM of LCA and
0.68 µM of 3-oxoLCA after 48 hours (Figure 4B). X-oxoUDCA was also detected at very small
concentrations around 0.5 µM from 12 hours until the end of the experiment. The potential isoform
of UDCA was detected at up to 9.92 µM at the 24-hour time point. Following the same trend observed
with the other primary BAs, *E. muris* SJ24 did not show any detectable activity with UDCA even after
co-induction. The chromatograms for the unknown bile acids can be found in Supplementary Figure
2.

As expected, neither *C. scindens* strain nor *E. muris* were capable of αMCA 7α-dehydroxylation in the
absence of ^{13}C-CA (Figure 5). *C. scindens* ATCC 35704 only produced minute amounts of an unknown
oxo form of αMCA (labelled Y-oxoαMCA) (0.85 µM at 32 hours). Once co-induced with ^{13}C-CA, 6-
oxoMDCA was detected at 2.7 µM after 48 hours in the *C. scindens* ATCC 35704 culture (Figure 5A).
This secondary bile acid has been 7α-DH-ed but also the hydroxyl at C6 oxidised. Moreover, several
intermediates for which standards are unavailable were also detected after 48 hours. These were
unknown oxidised forms of αMCA (labelled X- and Y- oxoαMCA) at concentrations not exceeding 5 µM
each. A third unknown BA was detected (albeit at very low concentrations, 0.41 µM at 32 hours) with
the same mass as 6-oxoMDCA, suggesting that it is an MCA species with one oxidation and one
dehydroxylation. This would indicate the production of another 7α-dehydroxylated form of αMCA *in
vitro* (Figure 1, Figure 5A). As for the ATCC 35704 strain, *C. scindens* VPI 12708 exhibited an increase
in the quantity of products from αMCA transformation in the presence of ^{13}C-CA relative to its absence
(Figure 5B). This includes the 7-DH-ed BA 6-oxoMDCA that reached a concentration of 8.18 µM after
48 hours and the X- and Y- αMCA forms that were detected at maximum concentrations of 4.22 µM
(4 hours) and 1.69 µM (32 hours), respectively. The aforementioned αMCA-derived bile acid with one
ketone group and one dehydroxylation was also detected at a maximum concentration of 3.49 µM
after 48 hours (Figure 5B). Surprisingly, *E. muris* SJ24 did not exhibit any observable 7-DH-ing activity
with or without ^{13}C-CA. Nevertheless, a small amount of X-oxoαMCA was detected at all time points,

with a stable concentration at around 2.4 µM without and 1.9 µM with [13]C-CA (Figure 5C). The results
for βMCA were very similar to those for αMCA and are discussed in further detail in the supplementary
information.

The concentration of [13]C-CA was also measured over time to ascertain that CA was being metabolised.
It was observed to decrease until it disappeared after 48 hours in the *C. scindens* strains except in the
presence of CDCA, for which the concentration decreased slowly over time. We attribute this
observation to the toxicity of CDCA at that concentration[20]. On the other hand, the concentration of
[13]C-CA in *E. muris* remained stable over time and in all conditions (Supplementary Figure 3).

Induction of bai gene expression in the presence of bile acids

In order to assess whether the amendment of [13]C-CA to the culture induced the *bai* operon as
hypothesised, the relative expression of *baiCD* and of *baiE* were measured. In addition, the expression
of *baiJ* (an accessory gene) was also monitored. Gene expression was normalised using at least three
reference genes and was calculated relative to the expression levels in a control group without BAs.
Both *E. muris* strains, that is JM40 (DSM 28560) and SJ24 (DSM 28561), have a truncated *baiJ* gene[21].
However, while the *baiJ* pseudogene is interrupted by stop codons in strain JM40, it is not in strain
SJ24, making it worthwhile to investigate *baiJ* in the latter strain (as was done in this study).
Additionally, *baiO* was also analysed for *E. muris* SJ24 as an alternative accessory *bai* gene[36].

Results show that the expression of *bai* operon genes in both *C. scindens* strains was highly
upregulated as a response to exposure to CA or to CDCA but not to the other BAs (Figure 6). CA and
CDCA are also the only two primary BAs for which *in vitro* 7-DH-ion data are already available[20].
Moreover, it is worth highlighting that CDCA was tested using a concentration of 200 µM vs. 100 µM
for CA.

For *C. scindens* ATCC 35704, the three genes tested were highly upregulated when [13]C-CA was present
along with another BA (UDCA, αMCA, or βMCA) (Figure 6A). In the CDCA dataset, statistically
significant differences relative to the uninduced conditions were observed and all genes were slightly
more upregulated in the presence of [13]C-CA (p-value < 0.001 linear model), but *baiCD* was more so
than the other genes (Figure 6A). Most interestingly, none of the other primary BAs activated the
expression of *bai* genes on their own, consistent with the lack of 7-DH-ing activity with these BA
substrates alone. However, the expression was brought up to levels higher than those observed for
CA once co-induced (Figure 6A).

A similar pattern was observed for *C. scindens* VPI 12708 but with the significant difference being that
baiJ was not overexpressed under any conditions (Figure 6B) (p-value < 0.001 linear model). In the
CDCA dataset, the co-induction with [13]C-CA had an upregulatory effect if assessed with a paired
Wilcoxon test (Supplementary Figure 4) but this effect was not found to be significant when using the
linear statistical model displayed in Figure 6. It is worth noting that the expression data were obtained
from the mid- to late-log exponential phase (around 18 hours) when differences in activity between
uninduced and co-induced conditions are not very large (Figure 3B). Similarly to *C. scindens* ATCC
35704, co-induction had a dramatic effect on the expression levels of *baiCD* and *baiE* in the presence
of UDCA, αMCA, or βMCA, with upregulation reaching the expression levels observed with CA or CDCA
alone (Figure 6B).

E. muris SJ24 showed a slight upregulation of *baiCD* and *baiE* in the CA dataset but it was not significant
and did not translate to any of the other conditions (Figure 6C), consistent with its very poor 7-DH-ing

activity (Figure 2). As a matter of fact, the increased *baiCD* and *baiE* gene expression ratio observed in the CA group was probably caused by one of the biological replicates which had a higher expression level than the others.

Thus, CA had a large effect on *bai* expression which was strain specific. Genes of the *bai* operon in the two *C. scindens* strains (ATCC 35704 and VPI 12708) exhibited a similar response to CA induction but the accessory *baiJ* differed in its response. It was upregulated in strain ATCC 35704 but not in strain VPI 12708. In contrast, CA had no significant effect on the expression of all the *bai* genes considered in *E. muris* SJ24.

The *rhaS*_1 gene (HDCHBGLK_01429) is immediately upstream of the *bai* operon promoter on the opposite strand and has also been proposed as bile acid-regulatory A (*barA*) due to its potential implication in *bai* regulation[7]. The expression of *rhaS1* and *rhaS2* (a copy of *rhaS1* elsewhere in the genome) was shown to have background levels across BAs (Supplementary Figure 5). This was tested in *C. scindens* ATCC 35704 without the amendment of [13]C-CA. Results indicate that *rhaS* is not upregulated by any of the BAs tested.

Thus, the question remains about the conditions propitious for *bai* gene expression and robust 7-DH-ion in *E. muris* SJ24. We hypothesized that other mouse-specific BAs may be the key inducers

bai gene induction by other BAs in *E. muris* SJ24

Because the presence of [13]C-CA did not induce *bai* genes in *E. muris* SJ24, we tested four BA cocktails to probe whether other BAs commonly found in the BA pool could promote *bai* expression. The BA pool was divided into four cocktails: tauro-conjugated BAs, oxidised BAs, sulfonated BAs and ωMCA. The addition of these BAs did not yield the production of any detectable secondary BAs (Figure 7). A small CA concentration (<2 μM) was detected in the tauro-BA cocktail (Figure 7A) but this was likely the result of the presence of CA as an impurity in the TCA standard, as it was also detected at time 0. In the oxidised BA cocktail, 12-oxoCDCA was almost fully reduced to CA after 16 hours (Figure 7B). Small quantities of CDCA and βMCA were detected, while both are likely to be impurities from the standards used (detected at time 0), it is worth highlighting that the concentration of CDCA increased from an average of 2.74 μM (time 0) to 5.35 μM (time 24), meanwhile, the concentration of βMCA remained stable around 2 μM. No reduction of 3-oxo forms was detected, likely due to the absence of *baiA2*[25]. Finally, neither sulfonated BAs nor ωMCA were transformed by *E. muris* SJ24 in any way (Figure 7C-D). Therefore, we excluded the possibility that other BAs could induce *bai* expression and 7-DH-ing activity in *E. muris* SJ24.

The expression of *baiCD*, *baiE*, the pseudogene *baiJ* and *baiO* was also measured in the BA cocktail experiments and compared with a CA-only reference group. Given the lack of 7-DH-ion of the BAs within the cocktails, it is not surprising that no significant upregulation was observed in any of the BA cocktail groups when compared to the CA control. (Supplementary Figure 6A).

Bile acid 7-DH-ion by E. muris SJ24 in the presence of mouse cecal content

CA 7-DH-ion by *E. muris* SJ24 was investigated in the presence of cecal content from either germ-free mice or a stable gnotobiotic murine model, Oligo-Mouse-Microbiota (Oligo-MM12)[37] in order to further investigate potential non-BA triggers for 7-DH-ion. A significant fraction of CA was conjugated with Co-enzyme A (CoA) and therefore could not be detected, as there are no standards for CoA-forms. In the controls (no cecal content), the DCA concentration averaged 4.26 μM after 48 hours which corresponded to the transformation of approximately 7% of the initial CA (Figure 8A). The amendment of cecal content from germ-free mice increased the DCA produced to 7.6 μM which

corresponded to 12% of the initial CA (Figure 8B). Finally, the addition of cecal content from Oligo-
MM12 mice produced only 0.67 μM of DCA but 18.6 μM of 7-oxoDCA (Figure 8C). In all conditions,
DCA was detected after 12 hours of incubation and gradually increased. 12-oxoCDCA was detected in
all conditions, while 3-oxoCA was only found in the CA control and germ-free groups (Figure 8). The
control groups of cecal content without *E. muris* SJ24 showed no change in CA concentration other
than the potential conjugation with Co-A by the Oligo-MM12 mouse case (Supplementary Figure 7).

Despite the measurable impact on 7-DH-ion by the addition of germ-free mouse cecal content of CA
7-DH-ion, it was not sufficient to significantly upregulate *bai* expression when compared to the CA-
only reference group (Supplementary Figure 6B). In both assays, the gene expression ratio of *bai* genes
was never above 3.

E. muris SJ24 *In vivo* 7-DH-ion and *bai* gene expression

The ability of *E. muris* strain DSM 28560 (JM40) to 7-DH-ate *in vivo* has been previously documented[21].
Here, colonisation of Oligo-MM12 mice was performed with the DSM 28561 strain (*E. muris* SJ24) to
confirm 7-DH-ion *in vivo* and quantify *bai* gene expression. The bile acid composition confirms active
7-DH-ion *in vivo* in Oligo-MM12 mice. Indeed, DCA, LCA and MDCA, were exclusively identified in the
sDMDMm2 + *E. muris* SJ24 group (Supplementary Figure 8).

As for the aforementioned expression assays, expression of *baiCD*, *baiE,* the pseudogene *baiJ* and *baiO*
was quantified, normalised against at least three reference genes and calculated relative to the
background signal detected from non-specific amplification in the Oligo-MM12 mice (without *E. muris*
SJ24).

Results show the expression of *E.muris* SJ24 *bai* genes *in vivo* (Figure 9). Indeed, *baiE* was expressed
at significant levels above background (gene expression ratio of 35.9) while *baiCD* and the *baiJ*
pseudogene had low gene expression ratios of 1.78 and 1.38, respectively. Finally, the expression ratio
of the auxiliary oxidoreductase *baiO* was situated at 4.04.

Discussion

Bile acid chemistry is a relevant field in human and veterinary medicine not only because of BA
detergent function during digestion but also for the wide range of roles related to host physiology and
homeostasis[38,39].

The *bai* operon was originally described in *C. scindens* VPI 12708 in 1990 and this strain has become
the reference for biochemical studies of the 7-DH-ion pathway[22,35,40]. The operon (*baiBCDEFGHI*)
encodes one CoA ligase (*baiB*), two oxidoreductases (*baiCD* and *baiH*), a 7-dehydratase (*baiE*), a CoA
transferase (*baiF*), a transporter (*baiG*), and a putative ketosteroid isomerase (*baiI*), but not all genes
are required for 7-DH-ion[25]. Expectedly from such a complex operon, comparative genomics have
highlighted significant differences in the *bai* operon amongst 7-DH-ing strains[35,41]. Furthermore, a
novel genetic synteny has been recently described and proven to be capable of 7-DH-ion. This novel
synteny does not follow the traditional *bai* operon structure and instead breaks it down to smaller
operons that cluster together[41,42]. Moreover, accessory *bai* genes have also been described across
multiple strains[35,40]. These accessory genes often cluster in two operons *baiJKL* and *baiNO* which is not
always complete; *C. scindens* ATCC 35704 only has *baiJ* (urocanate reductase)[32] while the VPI 12708
strain has the full *baiJKL* set[35]. From the genome analysis, *E. muris* SJ24 has fragments of *baiJKL* as

pseudogenes, while its sister strain JM40 has similar pseudogenes but interrupted by stop codons. The
exact role of these genes has yet to be defined.

Out of all primary BAs, only CA has been consistently reported to be 7-DH-ed *in vitro*. In contrast, CDCA
is poorly 7-DH-ed and other murine primary BAs (αMCA, βMCA or UDCA) are not at all[20,21,43–46]. We
confirm that the two *C. scindens* human isolates upregulated the expression of *bai* operon genes in
response to CA (Figure 6), except for *baiJ*. Indeed, we observed differences in the response of *baiJ* to
CA in between the two strains. This accessory gene is annotated as an urocanate reductase and a
member of the oxidoreductase family. The current CA 7-DH-ion pathway would suggest that this gene
is not involved in that process in strain VPI 12708[25], which is consistent with the lack of expression
observed here (Figure 6B). Conversely, its high level of upregulation in the strain ATCC 35704 matches
previously published data[32] and suggests that *baiJ* may play a role in the 7-DH-ion of CA or other BAs
in that strain. Finally, *E. muris* SJ24, the murine isolate, barely registered any *bai* gene expression
response to CA or any other BA tested.

CDCA co-induction with [13]C-CA was particularly effective in both *C. scindens* strains. The significant *bai*
upregulation (Figure 6) was reflected in an increase on the 7-DH-ed products from 1 to 5% in *C.*
scindens ATCC 35704 and a dramatic increase from 1 to 23% in *C. scindens* VPI 12708 (Figure 3). These
data suggest that the 7-DH-ion pathway of CDCA uses the same *bai* machinery as CA and its regulation
is dependent on the CA pathway. Furthermore, strain ATCC 35704 shows lower 7-DH-ion of CDCA
(Figure 3) and less effective 7-DH-ion of CA (Figure 2 and Supplementary Figure 1) as compared to
strain VPI 12708 despite a significant upregulation in *baiJ* expression.

UDCA is the 7β isomer of CDCA and is reported to have significant therapeutic properties[2,7,47]. In
humans, UDCA is synthesised from CDCA by gut microbes containing 7β-HSDHs[7,26,48,49] and is thus, a
secondary BA; in mice, it is produced by the liver, thus, it is a primary BA, and is used as a precursor
for βMCA[15,16]. Nevertheless, colonisation of mice lacking 7-dehydroxylating bacteria with 7-DH-ing
bacteria increases UDCA levels, implying that the gut microbiome also plays a significant role in UDCA
production by mice[14]. Despite the importance of this BA for medical applications, little is known about
the capacity of bacteria to 7β-dehydroxylate UDCA *in vitro.* Previous work has reported no
transformation by *C. scindens* strain ATCC 35704[20]. In accordance with our hypothesis, co-induction
with [13]C-CA not only greatly upregulated *bai* gene expression (Figure 6B) but also provided evidence
of UDCA 7β-dehydroxylation as LCA was detected in the culture (Figure 4). Indeed, the amount of 7-
DH-ed products upon co-induction increased from 0 to 15 and 16% in *C. scindens* strains ATCC 35704
and VPI 12708, respectively. The absence of CDCA in the culture suggests that UDCA was not
epimerised to CDCA and subsequently 7α-dehydroxylated. Moreover, two unidentified compounds
consistent with 7β-dehydroxylation were detected, providing further evidence of the 7β-
dehydroxylation of UDCA and its associated unique set of intermediates. The oxidised intermediate is
very likely to be 3-oxoUDCA (3-oxo-7β-hydroxy-5β-cholan-24-oic acid), as based on the mass, the only
alternative would have been 7-oxoLCA, which can be excluded as it is one the standards within our
collection (Figure 4). The other unknown compound had the same mass as UDCA albeit a different
retention time, suggesting this could be an iso- form of UDCA. It follows that this would be 3β-UDCA,
tentatively named isoUDCA (3β,7β-dihydroxy-5β-cholan-24-oic acid). Isoforms are well known in the
BA pool and particularly in CDCA-related intermediates[20,25,33,50] so it is likely that UDCA follows a similar
pattern. Despite the production of unknown intermediates, the activity response to co-induction
suggests that UDCA 7β-dehydroxylation uses the same Bai machinery as CDCA 7-DH-ion.

7-DH-ion activity was uncovered for αMCA and βMCA, for the first time and it yielded several
intermediate BAs that we were not able to fully characterise due to the lack of appropriate standards.

All human and mouse BAs share a backbone of four rings (Figure 1), this makes mass fractionation in the mass spectrometer unsuitable for identification. Therefore, we currently rely on comparison of ionised mass and retention time to standards. However, several assumptions can be made to speculate what these compounds could be. MCAs could be oxidised at the C-3, C-6 or C-7 position (Figure 1). A C-7 oxidation would yield 7-oxoMDCA regardless of the primary MCA. Meanwhile, other oxidations could be differentiated by the α or β conformation of the C-7 hydroxyl. It is possible that one of the intermediates that we detected was 7-oxoMDCA but none shared retention times across MCAs, meaning that different intermediates were produced for each of the two MCA substrates (Supplementary Figure 2). Considering that distinct oxidised intermediates were detected for the two MCAs, we hypothesise that those are the 3-oxo and 6-oxo forms of αMCA and βMCA (3-oxoαMCA: 3-oxo-6β,7α-dihydroxy-5β-cholan-24-oic acid; 6-oxoαMCA: 6-oxo-3α,7α-dihydroxy-5β-cholan-24-oic acid; 3-oxoβMCA: 3-oxo-6β,7β-dihydroxy-5β-cholan-24-oic acid; and 6-oxoβMCA: 6-oxo-3α,7β-dihydroxy-5β-cholan-24-oic acid). A third intermediate was also detected from both MCAs, with the ionised mass corresponding to secondary BAs with one dehydroxylation and a ketone group (e.g., 7-oxoLCA) (Figure 5 & Supplementary Figure 9). Three options are plausible: 1) A dehydroxylation at the C-3 position. This would yield a novel family of BAs with a 6β- and 7α/β- hydroxyls which is highly unlikely as it would have been identified previously by the multiple studies investigating the murine BA pool[15,16,51–54]. 2) An oxidation paired with a 6-dehydroxylation would yield 3-oxoCDCA, 7-oxoLCA, or 3-oxoUDCA. 3-oxoCDCA and 7-oxoLCA were included as standards in our analysis (Supplementary Table 3) and would have been detected if present. Meanwhile, the retention time of this compound is distinct from that of the compound proposed to be 3-oxoUDCA from the transformation of UDCA (see above and Supplementary Figure 2). Thus, this is not likely to be 3-oxoUDCA. 3) A 7-dehydroxylation could allow for ketone groups at the C-3 and C-6 positions. 6-oxoMDCA was available as a standard but, the second option, 3-oxoMDCA (3-oxo-6β-hydroxy-5β-cholan-24-oic acid), was not. It is therefore possible that this compound corresponds to 3-oxoMDCA, but this remains to be confirmed.

The mouse BA pool is significantly more diverse than that of humans due to the primary production of muricholic acids (with a hydroxyl group at the C-6 position) and of UDCA. The murine secondary BA pool includes DCA and LCA but also MDCA and its 6α counterpart, hyodeoxycholic acid (HDCA). Furthermore, mice can rehydroxylate TDCA back into TCA in the liver[15] which magnifies the differences between mouse and human BA pools. In general, the secondary BAs derived from muricholic acids seem to be in low abundance in the gut, hinting at the difficulty in 7-DH-ing these BAs. This is perhaps the reason why primary BAs such as βMCA are highly abundant in the mouse BA pool[55]. While we initially hypothesised that the lack of 7-DH-ion activity for αMCA and βMCA was due to the lack of *bai* gene expression, our co-induced data show that even *bai* gene expression in the *C. scindens* strains is insufficient for the production of MDCA[21], and results in the detection of potential oxidized versions of that BA (Figure 5 and Supplementary Figure 9).

As reported above, the murine strain SJ24 has shown no significant *bai* upregulation nor 7-DH-ion *in vitro* either in the presence or absence of ^{13}C-CA. To investigate the underlying reasons for this lack of activity, we considered three additional conditions: (a) various BA mixtures, to determine whether *bai* gene expression was controlled by another (or several other) murine BAs; (b) *in vitro* in the presence of Germ-free or Oligo-MM12 cecal content to ascertain whether the presence of other gut bacteria or signalling molecules from the host itself induced 7-DH-ion; or (c) in the Oligo-MM12 environment, to confirm the activity of strain SJ24 *in vivo*.

First, the *in vivo* condition exhibited high upregulation of *baiE* relative to the background, non-colonised control (Figure 9) and 7-DH-ed BAs were detected in the BA pool from the same samples

(Supplementary Figure 8). On the other hand, *baiCD* did not show evidence of upregulation. It is
possible that the expression of *baiCD* by *E. muris* is constitutive while *baiE*, a 7α-dehydratase[56],
presents a higher expression ratio, perhaps due to the smaller size of the protein (169 aa in *E. muris*
SJ24), intra-operon regulatory elements, or differential mRNA half-life[57]. The expression of *baiO*, a
different oxidoreductase, was slightly above that of *baiCD*. Whilst the gene expression ratios remained
low in both instances, a role in 7-DH-ion by this protein cannot be ruled out. Thus, *E. muris* SJ24 is
capable of *in vivo* 7-DH-ion, although its *bai* machinery may require additional genes that have not
been identified in this study. However, the exact trigger for *in vivo* levels of 7-DH-ion from *E. muris*
still remained elusive at this point.

To elucidate that question, we tested all the BAs detected in the Oligo-MM12 environment, and found
no evidence of 7-DH-ing activity (Figure 7), excluding the possibility that non-CA BA triggered *bai* gene
expression in *E. muris*.

However, when SJ24 was grown in the presence of cecal content from germ-free mice, its 7-DH-ing
activity increased (Figure 8) despite *bai* gene expression not increasing significantly (Supplementary
Figure 6B). Indeed, the amendment of cecal content from germ-free mice (Figure 8B) resulted in a
significant increase from 7 to 13% of 7-DH-ed products, coupled with a decrease of the abundance of
the 12-oxoCDCA (3α,7α-dihydroxy-12-oxo-5β-cholanic acid) intermediate. Interestingly, the co-
cultivation of *E. muris* with the non-sterile cecal-content from Oligo-MM12 mice produced high
amounts of 7-oxoDCA (Figure 8C) while CA was not transformed by the same cecal content in the
absence of strain SJ24 (Supplementary Figure 7). This could suggest an interaction between *E. muris*
and the gnotobiotic community in which the former would promote the 7-oxidation of CA by the
latter, known to harbor 7-HSDHs[58]. This interaction appears to be exclusive to the *in vitro* environment
since the *in vivo* BA data show lower concentrations of 7-oxoDCA than DCA (Supplementary Figure 8).

The evidence presented in this study shows that the role of the host, presumably through signalling,
is a critical element for effective 7-DH-ion by *E. muris* and the regulatory mechanisms of this secondary
BA transformation is dramatically different amongst bacterial species. The results also highlight the
significant differences in 7-DH-ion between human and mouse isolates but also between the *in vitro*
and *in vivo* environments.

To further highlight the differences, the human isolates showed marginal activity for αMCA and βMCA
upon co-induction. 7-DH-ed forms such as 6-oxoMDCA were detected (Figure 5 & Supplementary
Figure 9) but the full 7-dehydroxylation to MDCA was not observed. These data add further evidence
that other elements besides the *bai* operon (and *baiJ*) might be needed to completely 7-dehydroxylate
MCAs. Perhaps, a missing 6β-HSDH gene would be required for complete 7-DH-ion. The apparent
simplicity of 7-DH-ion regulation from human isolates compared to that of *E. muris* could be due to
the more diverse diet of humans than mice. It has been observed that a less diverse diet can
overstimulate the BA pool in humans and increase the incidence of colorectal cancer[59]. The natural
mouse diet is less diverse than that of humans and their initial lactation period (a monotrophic diet)
plays a much stronger role in the mouse lifespan[60]. In these circumstances, a strong regulation of 7-
DH-ion might be an important mechanism to prevent BA pool unbalances. Nevertheless, much more
data on the 7-DH-ion mechanisms of various strains with particular focus on isolates from the mouse
and other animal models is required to investigate the potential differences in 7-DH-ion regulation.

Conclusion

The findings presented here are fourfold. First, we demonstrated that the previously reported[32,33]
strong upregulation of *bai* genes by CA increases the extent of 7-DH-ion of other primary BAs. This
was particularly true for UDCA which has been reported to be converted to LCA *in vitro* for the first
time, with important differences in the 7-DH-ing capabilities amongst strains.

Secondly, the upregulation of *bai* genes exhibited strain-specific differences. *C. scindens* ATCC 35704
upregulated the *bai* operon genes *baiCD* and *baiE* as well as the *bai* accessory gene *baiJ*. While for *C.*
scindens VPI 12708, it upregulated the *bai* operon genes, but *baiJ* expression was found to be at a
background level. This is consistent with the lack of involvement of *baiJ* in 7-DH-ion in strain VPI
12708[25]. *E. muris* SJ24 was the third bacterium tested, a murine isolate with *in vivo* 7-DH-ing
capabilities (Supplementary Figure 8). Strain SJ24 showed weak *in vitro* 7-DH-ion of CA and no
upregulation of any of the *bai* genes tested. The activity of this strain was promoted by the addition
of germ-free cecal content but not cecal content from Oligo-MM12 colonized mice. This result
suggests that host signalling is required for efficient 7-DH-ing activity by strain SJ24 but that the
presence of a minimum microbiome (12 strains) inhibits this activity potentially due to the promotion
of 7-HSDH activity from the microbiome *in vitro*. Unravelling the controls on BA 7-DH-ion by *E. muris*
requires further investigation.

Thirdly, *C. scindens* human isolates can partially 7-dehydroxylate MCAs, leading to the formation of
oxidized MDCA at the C-6 position. Therefore, an enzyme capable of reducing this compound, e.g., a
6β-HSDH, is required to achieve the end point of 7-DH-ion that is observed *in vivo*, namely MDCA. To
date, such protein has not been identified in any microorganism.

Finally, the mechanism of 7-DH-ion regulation differs significantly between murine- and human-
derived strains, which could be due to the nature of the host. Human isolates showcase a system
governed by CA, while murine isolates appear to utilise a BA-independent signal present in the lumen.

In conclusion, these data provide novel insights into the intricacies of 7-DH-ion and the significant
differences amongst 7-DH-ing bacteria. The CA-dependent co-induction can be attributed to the
abundance of this compound in the BA pool of humans. However, the inducing factor for *E. muris*
activity remains elusive despite evidence suggesting that it is host-derived. Moreover, multiple novel
BAs were observed and their identity surmised. Further work with these and other strains is required
to investigate the 7-DH-ion pathway of CDCA, αMCA, βMCA and UDCA as well as to explore the strain-
specific differences regarding the 7-DH-ion pathway of CA.

Materials and Methods

Bacterial strains and growth conditions

The strains used were *Clostridium scindens* ATCC 35704, *Clostridium scindens* VPI 12708 and *Extibacter*
muris DSM 28561 (SJ24), this strain was chosen instead of *E. muris* DSM 28560 (JM40) due to its ability
to grow faster *in vitro* (24h vs 48h, data not shown). Bacteria were grown in Brain Heart Infusion
Supplement – Salts (BHI-S) medium, consisting of 37g BHI, 1g L-cysteine, 5g yeast extract, 2.5g
fructose, 50mL salts solution (0.2g $CaCl_2$, 0.2 $MgSO_4$, 1g K_2HPO_4, 1g KH_2PO_4, 10g $NAHCO_3$ and 2g NaCl
per L of ddH_2O) per L ddH_2O. The salts solution and media were sterilised by autoclaving. Static growth
was carried out at 37°C in an anoxic chamber (Coy Laboratory Products, 95% N_2, 5% H_2). A pre-
inoculum was prepared from glycerol stocks (using BHIS-S) before inoculating 25 mL of BHIS-S in
Falcon tubes at a starting OD_{600} of 0.05. Results of the growth curves for the experiments can be found
in the Supplementary Information (Supplementary Figure 10).

In vitro 7-dehydroxylation assays

Bacteria were grown in presence of BAs: CA (100 µM), CDCA (200 µM), αMCA (100 µM), βMCA (100
µM) and UDCA (100 µM) or the same volume of ethanol (solvent control). A sterile control (media
with ethanol solvent) was also included. Co-induction was performed by adding an additional 100 µM
of ^{13}C-CA to CDCA, UDCA, αMCA, or βMCA). ^{13}C-CA was chosen so its transformation products could
be separated from those of the other primary BAs during the quantification process. All uninduced
experiments were conducted at the same time while all the co-induced ones were run simultaneously
at a different time. Both experiments included a condition consisting of the amendment of only 100
µM CA, which allows for comparison of the results from co-induced and uninduced experiments.
Growth was monitored by periodically measuring the OD_{600}. During the main time points (0, 4, 8, 12,
24, 32 and 48 hours), 1mL samples were collected for BA extraction in a 2 mL bead-beating resistant
tube. All conditions were performed in triplicates.

Four BA cocktails were prepared for the assays with *E. muris* SJ24 based on the BAs often present in a
meaningful amount within the BA pool of mice. All BAs within the cocktails were at 50 µM. The Tauro-
conjugated cocktail included TCA, TCDCA, TαMCA, TβMCA, TUDCA and THCA. The Oxo- cocktail
included 3-oxoCA, 3-oxoCDCA, 7-oxoDCA and 12-oxoCDCA. The Sulfo- cocktail included CA7S and
CDCA3S. Finally, the last group was only comprised of 50 µM of ωMCA. All BAs were diluted in ethanol
or methanol depending on their solubility. Three time points were collected (0, 16 and 24 hours).

Furthermore, *E. muris* SJ24 was also amended with cecal content to test its implications over 7-DH-
ion *in vitro*. For this assay, 25 mg of freeze-dried cecal content from germ-free mice or 25 mg of frozen
content with 5% glycerol (v/v) from sDMDMm2 mice were added to 25 mL of BHIS-S with 100 µM CA.
The control conditions for this experiment were a group with 100 µM CA but no additional cecal
content and two more groups with each respective type of cecal content but no *E. muris* SJ24. Seven
time points were taken (0, 4, 8, 12, 24, 32 and 48). RNA sample collection was particularly early for
this experiment (4-hour time point) due to a faster-than-usual growth (Supplementary Figure 10).

Bile acid extraction

Samples were vacuum dried overnight (ON). Approximately 450 mg of 0.5 mm zirconium beads were
added to the dried samples as well as 500 µL ice-cold alkaline acetonitrile (acetonitrile – 25% ammonia
4:1 v/v) and 100 µL of ISTD solution (CA-d$_4$, CDCA-d$_4$, TCA-d$_4$, TUDCA-d$_4$, DCA-d$_4$ and LCA-d$_4$, each at
100 µM in methanol). Samples were homogenised in a Precellys 24 Tissue Homogenizer (Bertin
Instruments, Montigny-le-Bretonneux, France) at 6500 rpm 3x 30" beat 30" rest. Samples were
vortexed for 1 hour and centrifugated for 15 minutes at 16000 rcf at room temperature.
Approximately 500 µL of suspension was carefully collected over the beads level and transferred into
a new 1.5 mL epi tube which was then vacuum dried overnight. Finally, the samples were reconstituted
in 1 mL of ammonium acetate [5mM] – methanol (50:50 v/v) and a 1:20 dilution with the same solvent
was prepared in LC-MS glass vials, ready for injection.

RNA extraction and reverse transcription

1 mL of sample was collected in a 15 mL falcon tube during the mid-log to late-log phase for the RNA
extraction. The sample was stored with RNAprotect following the manufacturer protocol (Protocol 5
from RNAprotect Bacteria Reagent Handbook 01/2020, Qiagen) at -80°C until processed. All conditions
were performed in triplicates. Lysis and RNA purifications were done using the RNeasy Mini Kit
(Qiagen, Hilden, Germany). Bacterial lysis was performed following Protocol 5: Enzymatic Lysis,
Proteinase K Digestion and Mechanical Disruption of Bacteria (RNAprotect Bacteria Reagent
Handbook 01/2020, Qiagen), with 20 µL of proteinase K for each sample and the required volumes for
a number of bacteria <7.5 x 10^8. The cell lysis was performed using a Precellys 24 Tissue Homogenizer

(Bertin Instruments, Montigny-le-Bretonneux, France) at 6500 rpm 3x 10 seconds beat 10 seconds
rest. RNA purification was performed following Protocol 7: Purification of Total RNA from Bacterial
Lysate using the RNeasy Mini Kit. Centrifugations were carried out at 15000 rcf except for the 2 min
centrifugation which was done at 18000 rcf.

Purified RNA was further subject to a DNase treatment using the RQ DNase I (Promega, Madison, WI,
USA) following the manufacturer protocol with small modifications: The final volume was adjusted to
a 100 µL and incubation was extended to 1 hour at 37°C. The treated RNA was cleaned-up using the
RNeasy Mini Kit (Qiagen, Hilden, Germany) following the RNA Clean-up protocol from the
manufacturer (RNeasy Mini Handbook 10/2019) with the 2 min centrifugation done at 18000 rcf.
Concentration and purity of RNA was measured with a NanoDrop One (Thermo Fisher Scientific,
Waltham, MA, USA).

100 ng of RNA was reverse transcribed into cDNA using the GoScript™ Reverse Transcription Mix,
Random Primers (Promega, Madison, WI, USA) following the manufacturer protocol. The process was
done in duplicates with one group using water instead of the reaction buffer as a non-reverse
transcription control (NRT).

Reverse transcription quantitative PCR (RT-qPCR)

RT-qPCRs were prepared using the Myra liquid handling system (Bio Molecular Systems, software
version 1.6.26) and performed using the Magnetic induction cycler (Mic) platform (Bio Molecular
Systems, Upper Coomera, QLD, Australia) with the micPCR software (v2.10.5).

The list of primers used can be found in Supplementary table 2. Samples were prepared with the
SensiFAST SYBR No-ROX Kit (Meridian Bioscience, Cincinnati, OH, USA) at a final volume of 10 µL. All
runs were performed with the following program, with small modifications: Initial hold at 95°C for 5
minutes with a cycle of 95°C for 5 seconds, 54.5°C for 20 seconds (54.1°C for *E. muris* SJ24) and 72°C
for 9 seconds. 40 cycles were done for *C. scindens* ATCC 35704 and 50 for *C. scindens* VPI 12708 and
E. muris SJ24. The melting curve, temperature control and acquisition settings were left as default.
The quantification was done using three or more reference genes (Supplementary Table 2) based on
their expression stability across conditions. NRTs as well as no template controls (NTCs) were included
to check for residual DNA or contaminations. Four technical replicates were done for each biological
replicate. Note that expression data presented in Figure 6 for the CA-only condition (labelled CA)
correspond to the pooled expression results (for the condition in which only 100 µM Ca was added)
from the two sets of experiments presented above (referred to as the co-induced and the uninduced
experiments, respectively). The *in vivo* expression data presented in Figure 9 was normalised against
the background signal detected from an uncolonised Oligo-MM12 control group.

Liquid chromatography – mass spectrometry (LC-MS)

The quantitative method was performed on an Agilent ultrahigh-performance liquid chromatography
1290 series coupled in tandem to an Agilent 6530 Accurate-Mass Q-TOF mass spectrometer. The
separation was done on a Zorbax Eclipse Plus C18 column (2.1 x 100mm, 1.8 µm) and a guard column
Zorbax Eclipse Plus C18 (2.1 x 5mm, 1.8 µm) both provided by Agilent technologies (Santa Clara, CA,
USA). The column compartment was kept heated at 50°C. Two different solutions were used as
eluents: ammonium acetate [5mM] in water as mobile phase A and pure acetonitrile as mobile phase
B. A constant flow of 0.4 mL/min was maintained over 26 minutes of run time with the following
gradient (expressed in eluent B percentage): 0-5.5 min, constant 21.5% B; 5.5-6 min, 21.5-24.5% B; 6-
10 min, 24.5-25% B; 10-10.5 min, 25-29% B; 10.5-14.5 min, isocratic 29% B; 14.5-15 min, 29-40% B;
15-18 min, 40-45% B; 18-20.5 min, 45-95% B; 20.5-23 min, constant 95% B; 23-23.1 min, 95-21.5% B;

23.10-26 min, isocratic 21.50% B. The system equilibration was implemented at the end of the
gradient for 3 minutes in initial conditions. The autosampler temperature was maintained at 10°C and
the injection volume was 5µL. The ionisation mode was operated in negative mode for the detection
using the Dual AJS Jet stream ESI Assembly. The QTOF acquisition settings were configured in 4GHz
high-resolution mode (resolution 17000 FWHM at m/z 1000), data storage in profile mode and the
high-resolution full MS chromatograms were acquired over the range of m/z 100-1700 at a rate of 3
spectra/s. The mass spectrometer was calibrated in negative mode using ESI-L solution from Agilent
technologies every 6 hours to maintain the best possible mass accuracy. Source parameters were
setup as follows: drying gas flow, 8 L/min; gas temperature, 300°C; nebulizer pressure, 35psi; capillary
voltage, 3500V; nozzle voltage, 1000V. Data were processed afterwards using the MassHunter
Quantitative software and MassHunter Qualitative software to control the mass accuracy for each
run. In the quantitative method, 42 bile acids were quantified by calibration curves (Supplementary
Table 3). The quantification was corrected by addition of internal standards in all samples and
calibration levels. Extracted ion chromatograms were generated using a retention time window of ±
1.5 min and a mass extraction window of ± 30ppm around the theoretical mass of the targeted bile
acid. Unknown BAs were identified when found within the retention time window of a standard with
the same ionised mass. Approximate quantification of these unknown BAs was done by using the
nearest standard (by retention time) with the same ionised mass.

Animals and Ethics Statement

sDMDMm2[61] mice were housed in the Clean Mouse Facility (CMF, Department of Clinical Research) of
the University of Bern. Animal experiments were performed in accordance with the Swiss Federal and
the Bernese Cantonal regulations and were approved by the Bernese Cantonal ethical committee for
animal experiments under the license number BE82/13.

In vivo colonisation with *E. muris* SJ24

A cohort of nine sDMDMm2[61] mice were used. Four mice were dedicated to an uncolonized control
group and the remaining five were colonised with *E. muris* DSM 28561 (SJ24). sDMDMm2 animals to
be colonised were imported from breeding isolators into small experimental isolators and
administrated orally with approximately 10^9 CFUs (in 200 µL). Control animals remained in the
breeding isolator during this period. After 11 days, all animals were exported into a laminar flow hood
and sacrificed. Cecal content was collected for bile acid measurement and RNA extractions. RNA was
extracted from 75-100 mg of cecal content obtained at the end of the experiment (11 days from the
initiation of the colonisation experiment). 20 to 50 mg (dry weight) of cecal content were used for BA
extraction. BA quantification and RT-qPCR assays were performed as described above.

Statistical analysis and data visualisation

Graphpad Prism 9.2.0 (GraphPad) was used to generate the figures shown in this paper and perform
pairwise comparisons, the gene expression data was analysed with a linear model (LM) in R language
v4.1.2[62] using RStudio[63] a 2-way ANOVA or a Welch's t-test. The statistical significance boundary was
stablished at a *p*-value < 0.05.**Conflicts of interest**

The authors declare no conflicts of interest.

Data Availability Statement

The data used for this manuscript are publicly available in the following link:

https://doi.org/10.5281/zenodo.6034320

Funding

The research hereby shown was funded by the Swiss National Science Foundation (Sinergia
CRSII5_180317)

References

1.   Monte MJ, Marin JJG, Antelo A, Vazquez-Tato J. Bile acids: Chemistry, physiology, and
617   pathophysiology. World J Gastroenterol 2009; 15:804–16. doi: 10.3748/wjg.15.804

618   2.   Hanafi NI, Mohamed AS, Kadir SHSA, Othman MHD. Overview of bile acids signaling and
619   perspective on the signal of ursodeoxycholic acid, the most hydrophilic bile acid, in the heart.
620   Biomolecules2018; 8. doi: 10.3390/biom8040159

621   3.   Kuipers F, Bloks VW, Groen AK. Beyond intestinal soap - Bile acids in metabolic control. Nat Rev
622   Endocrinol 2014; 10:488–98. doi: 10.1038/nrendo.2014.60

623   4.   Wang H, Chen J, Hollister K, Sowers LC, Forman BM. Endogenous bile acids are ligands for the
624   nuclear receptor FXR/BAR. Mol Cell 1999; 3:543–53. doi: 10.1016/S1097-2765(00)80348-2

625   5.   Bernstein H, Bernstein C, Payne CM, Dvorakova K, Garewal H. Bile acids as carcinogens in
626   human gastrointestinal cancers. Mutat Res - Rev Mutat Res 2005; 589:47–65. doi:
627   10.1016/j.mrrev.2004.08.001

628   6.   Joyce SA, Gahan CGM. Bile Acid Modifications at the Microbe-Host Interface: Potential for
629   Nutraceutical and Pharmaceutical Interventions in Host Health. Annu Rev Food Sci Technol
630   2016; 7:313–33. doi: 10.1146/annurev-food-041715-033159

631   7.   Ridlon JM, Kang D-JJ, Hylemon PB, Ridlon JM, Kang D-JJ, Hylemon PB. Bile salt
632   biotransformations by human intestinal bacteria. J Lipid Res 2006; 47:241–59. doi:
633   10.1194/jlr.R500013-JLR200

634   8.   Claudel T, Staels B, Kuipers F. The Farnesoid X receptor: A molecular link between bile acid and
635   lipid and glucose metabolism. Arterioscler. Thromb. Vasc. Biol.2005; 25:2020–31. doi:
636   10.1161/01.ATV.0000178994.21828.a7

637   9.   Guo C, Chen WD, Wang YD. TGR5, not only a metabolic regulator. Front. Physiol.2016; 7. doi:
638   10.3389/fphys.2016.00646

639   10.   van Nierop FS, Scheltema MJ, Eggink HM, Pols TW, Sonne DP, Knop FK, Soeters MR. Clinical
640   relevance of the bile acid receptor TGR5 in metabolism. Lancet Diabetes Endocrinol 2016;
641   5:224–33. doi: 10.1016/S2213-8587(16)30155-3

642   11.   Kawamata Y, Fujii R, Hosoya M, Harada M, Yoshida H, Miwa M, Fukusumi S, Habata Y, Itoh T,
643   Shintani Y, et al. A G protein-coupled receptor responsive to bile acids. J Biol Chem 2003;
644   278:9435–40. doi: 10.1074/jbc.M209706200

645   12.   Brighton CA, Rievaj J, Kuhre RE, Glass LL, Schoonjans K, Holst JJ, Gribble FM, Reimann F. Bile
646   acids trigger GLP-1 release predominantly by accessing basolaterally located G protein-coupled
647   bile acid receptors. Endocrinology 2015; 156:3961–70. doi: 10.1210/en.2015-1321

648   13.   Buffie CG, Bucci V, Stein RR, McKenney PT, Ling L, Gobourne A, No D, Liu H, Kinnebrew M, Viale
649   A, et al. Precision microbiome reconstitution restores bile acid mediated resistance to
650   Clostridium difficile. Nature 2015; 517:205–8. doi: 10.1038/nature13828

651   14.   Studer N, Desharnais L, Beutler M, Brugiroux S, Terrazos MA, Menin L, Schürch CM, McCoy KD,

Kuehne SA, Minton NP, et al. Functional intestinal bile acid 7α-dehydroxylation by *Clostridium scindens* associated with protection from *Clostridium difficile* infection in a gnotobiotic mouse model. Front Cell Infect Microbiol 2016; 6:191. doi: 10.3389/fcimb.2016.00191

15. Li J, Dawson PA. Animal models to study bile acid metabolism. Biochim. Biophys. Acta - Mol. Basis Dis.2019; 1865:895–911. doi: 10.1016/j.bbadis.2018.05.011

16. Honda A, Miyazaki T, Iwamoto J, Hirayama T, Morishita Y, Monma T, Ueda H, Mizuno S, Sugiyama F, Takahashi S, et al. Regulation of bile acid metabolism in mouse models with hydrophobic bile acid composition. J Lipid Res 2020; 61:54–69. doi: 10.1194/jlr.RA119000395

17. Morris GN, Winter J, Cato EP, Ritchie AE, Bokkenheuser VD. *Clostridium scindens* sp. nov., a Human Intestinal Bacterium with Desmolytic Activity on Corticoids. Int J Syst Bacteriol 1985; 35:478–81. doi: 10.1099/00207713-35-4-478

18. Lagkouvardos I, Pukall R, Abt B, Foesel BU, Meier-Kolthoff JP, Kumar N, Bresciani A, Martínez I, Just S, Ziegler C, et al. The Mouse Intestinal Bacterial Collection (miBC) provides host-specific insight into cultured diversity and functional potential of the gut microbiota. Nat Microbiol 2016; 1:1–15. doi: 10.1038/nmicrobiol.2016.131

19. Streidl T, Kumar N, Suarez LN, Rohn S, Clavel T. *Extibacter*. In: Bergey's Manual of Systematics of Archaea and Bacteria. American Cancer Society; 2019. page 1–7.doi: 10.1002/9781118960608.gbm01749

20. Marion S, Studer N, Desharnais L, Menin L, Escrig S, Meibom A, Hapfelmeier S, Bernier-Latmani R. *In vitro* and *in vivo* characterization of *Clostridium scindens* bile acid transformations. Gut Microbes 2019; 10:481–503. doi: 10.1080/19490976.2018.1549420

21. Streidl T, Karkossa I, Segura Muñoz RR, Eberl C, Zaufel A, Plagge J, Schmaltz R, Schubert K, Basic M, Schneider KM, et al. The gut bacterium *Extibacter muris* produces secondary bile acids and influences liver physiology in gnotobiotic mice. Gut Microbes 2021; 13:1–21. doi: 10.1080/19490976.2020.1854008

22. Mallonee DH, White WB, Hylemon PB. Cloning and sequencing of a bile acid-inducible operon from *Eubacterium* sp. strain VPI 12708. J Bacteriol 1990; 172:7011–9.

23. Ridlon JM, Hylemon PB. Identification and characterization of two bile acid coenzyme A transferases from *Clostridium scindens*, a bile acid 7α-dehydroxylating intestinal bacterium . J Lipid Res 2012; 53:66–76. doi: 10.1194/jlr.m020313

24. Wells JE, Hylemon PB. Identification and Characterization of a Bile Acid 7alpha - Dehydroxylation Operon in *Clostridium* sp. Strain TO-931, a Highly Active 7alpha - Dehydroxylating Strain Isolated from Human Feces. Appl Environ Microbiol 2000; 66:1107–13. doi: 10.1128/AEM.66.3.1107-1113.2000

25. Funabashi M, Grove TL, Wang M, Varma Y, McFadden ME, Brown LC, Guo C, Higginbottom S, Almo SC, Fischbach MA. A metabolic pathway for bile acid dehydroxylation by the gut microbiome. Nature 2020; :1–5. doi: 10.1038/s41586-020-2396-4

26. Macdonald IA, Hutchison DM. Epimerization versus dehydroxylation of the 7α-hydroxyl- group of primary bile acids: Competitive studies with *Clostridium absonum* and 7α-dehydroxylating bacteria (*Eubacterium* SP.). J Steroid Biochem 1982; 17:295–303. doi: 10.1016/0022-4731(82)90203-5

27. Doerner KC, Takamine F, LaVoie CP, Mallonee DH, Hylemon PB. Assessment of fecal bacteria with bile acid 7α-dehydroxylating activity for the presence of *bai*-like genes. Appl Environ Microbiol 1997; 63:1185–8.

28. Vinithakumari AA, Hernandez BG, Ghimire S, Adams S, Stokes C, Jepsen I, Brezina C, Sahin O, Li G, Tangudu C, et al. A model screening pipeline for bile acid converting anti-*Clostridioides difficile* bacteria reveals unique biotherapeutic potential of *Peptacetobacter hiranonis*. bioRxiv 2021; :2021.09.29.462466. doi: 10.1101/2021.09.29.462466

29. Ridlon JM. Enzymology and molecular biology of bile acid 7alpha- and 7beta- dehydroxylation by the intestinal bacteria *Clostridium scindens* and *Clostridium hylemonae*. PhD Thesis Virginia Commonw Univ 2008; :1–333. doi: https://doi.org/10.25772/BN62-H676

30. Makino I, Nakagawa S. Changes in biliary lipid and biliary bile acid composition in patients after administration of ursodeoxycholic acid. J Lipid Res 1978; 19:723–8. doi: 10.1016/s0022-2275(20)41272-6

31. Meir K, Kitsberg D, Alkalay I, Szafer F, Rosen H, Shpitzen S, Avi L Ben, Staels B, Fievet C, Meiner V, et al. Human sterol 27-hydroxylase (CYP27) overexpressor transgenic mouse model. Evidence against 27-hydroxycholesterol as a critical regulator of cholesterol homeostasis. J Biol Chem 2002; 277:34036–41. doi: 10.1074/jbc.M201122200

32. Devendran S, Shrestha R, Alves JMP, Wolf PG, Ly L, Hernandez AG, Méndez-García C, Inboden A, Wiley J, Paul O, et al. *Clostridium scindens* ATCC 35704: Integration of nutritional requirements, the complete genome sequence, and global transcriptional responses to bile acids. Appl Environ Microbiol 2019; 85. doi: 10.1128/AEM.00052-19

33. Ridlon JM, Devendran S, Alves JM, Doden H, Wolf PG, Pereira G V., Ly L, Volland A, Takei H, Nittono H, et al. The '*in vivo* lifestyle' of bile acid 7α-dehydroxylating bacteria: comparative genomics, metatranscriptomic, and bile acid metabolomics analysis of a defined microbial community in gnotobiotic mice. Gut Microbes 2019; :1–24. doi: 10.1080/19490976.2019.1618173

34. Bhowmik S, Jones DH, Chiu HP, Park IH, Chiu HJ, Axelrod HL, Farr CL, Tien HJ, Agarwalla S, Lesley SA. Structural and functional characterization of BaiA, an enzyme involved in secondary bile acid synthesis in human gut microbe. Proteins Struct Funct Bioinforma 2014; 82:216–29. doi: 10.1002/prot.24353

35. Ridlon JM, Harris SC, Bhowmik S, Kang D-JJ, Hylemon PB. Consequences of bile salt biotransformations by intestinal bacteria. Gut Microbes 2016; 7:22–39. doi: 10.1080/19490976.2015.1127483

36. Heinken A, Ravcheev DA, Baldini F, Heirendt L, Fleming RMTT, Thiele I. Systematic assessment of secondary bile acid metabolism in gut microbes reveals distinct metabolic capabilities in inflammatory bowel disease. Microbiome 2019; 7:75. doi: 10.1186/s40168-019-0689-3

37. Brugiroux S, Beutler M, Pfann C, Garzetti D, Ruscheweyh HJ, Ring D, Diehl M, Herp S, Lötscher Y, Hussain S, et al. Genome-guided design of a defined mouse microbiota that confers colonization resistance against *Salmonella enterica* serovar Typhimurium. Nat Microbiol 2016; 2:1–12. doi: 10.1038/nmicrobiol.2016.215

38. Hofmann AF, Hagey LR. Key discoveries in bile acid chemistry and biology and their clinical applications: History of the last eight decades. J. Lipid Res.2014; 55:1553–95. doi: 10.1194/jlr.R049437

39. Winston JA, Theriot CM. Diversification of host bile acids by members of the gut microbiota. Gut Microbes2020; 11:158–71. doi: 10.1080/19490976.2019.1674124

40. Ridlon JM, Kang DJ, Hylemon PB. Isolation and characterization of a bile acid inducible 7α-dehydroxylating operon in *Clostridium hylemonae* TN271. Anaerobe 2010; 16:137–46. doi:

10.1016/j.anaerobe.2009.05.004

41.   Song I, Gotoh Y, Ogura Y, Hayashi T, Fukiya S, Yokota A. Comparative genomic and physiological
analysis against *Clostridium scindens* reveals *Eubacterium* sp. C-25 as an atypical deoxycholic
acid producer of the human gut microbiota. Microorganisms 2021; 9:2254. doi:
10.3390/microorganisms9112254

42.   Bai Y, Zhao T, Gao M, Zou Y, Lei X. A Novel Gene Alignment in *Dorea* sp. AM58-8 Produces 7-
Dehydroxy-3β Bile Acids from Primary Bile Acids. Biochemistry 2022; doi:
10.1021/acs.biochem.2c00264

43.   Hylemon PB, Melone PD, Franklund C V., Lund E, Bjorkhem I. Mechanism of intestinal 7α-
dehydroxylation of cholic acid: Evidence that allo-deoxycholic acid is an inducible side-product.
J Lipid Res 1991; 32:89–96.

44.   Masuda N, Oda H, Hirano S, Tanaka H. Enhancement of the 7α-dehydroxylase activity of a
gram-positive intestinal anaerobe by flavins. Appl Environ Microbiol 1983; 45:308–9.

45.   Kitahara M, Takamine F, Imamura T, Benno Y. Assignment of *Eubacterium* sp. VPI 12708 and
related strains with high bile acid 7alpha-dehydroxylating activity to *Clostridium scindens* and
proposal of *Clostridium hylemonae* sp. nov., isolated from human faeces. Int J Syst Evol
Microbiol 2000; 50:971–8. doi: 10.1099/00207713-50-3-971

46.   Benno Y, Kitahara M, Takamine F, Imamura T. *Clostridium hiranonis* sp. nov., a human intestinal
bacterium with bile acid 7alpha-dehydroxylating activity. Int J Syst Evol Microbiol 2001; 51:39–
44. doi: 10.1099/00207713-51-1-39

47.   Abdallah E, Emile SH, Elfeki H, Fikry M, Abdelshafy M, Elshobaky A, Elgendy H, Thabet W,
Youssef M, Elghadban H, et al. Role of ursodeoxycholic acid in the prevention of gallstone
formation after laparoscopic sleeve gastrectomy. Surg Today 2017; 47:844–50. doi:
10.1007/s00595-016-1446-x

48.   Coleman JP, Hudson LL, Adams MJ. Characterization and regulation of the NADP-linked 7α-
hydroxysteroid dehydrogenase gene from *Clostridium sordellii*. J Bacteriol 1994; 176:4865–74.

49.   Ferrandi EE, Bertolesi GM, Polentini F, Negri A, Riva S, Monti D. In search of sustainable
chemical processes: Cloning, recombinant expression, and functional characterization of the
7α- and 7β-hydroxysteroid dehydrogenases from *Clostridium absonum*. Appl Microbiol
Biotechnol 2012; 95:1221–33. doi: 10.1007/s00253-011-3798-x

50.   Devlin AS, Fischbach MA. A biosynthetic pathway for a prominent class of microbiota-derived
bile acids. Nat Chem Biol 2015; 11:685–90. doi: 10.1038/nchembio.1864

51.   Narushima S, Itoh K, Miyamoto Y, Park SH, Nagata K, Kuruma K, Uchida K. Deoxycholic acid
formation in gnotobiotic mice associated with human intestinal bacteria. Lipids 2006; 41:835–
43. doi: 10.1007/s11745-006-5038-1

52.   Eyssen HJ, Parmentier GG, Mertens JA. Sulfated Bile Acids in Germ-Free and Conventional
Mice. Eur J Biochem 1976; 66:507–14. doi: 10.1111/j.1432-1033.1976.tb10576.x

53.   Han J, Liu Y, Wang R, Yang J, Ling V, Borchers CH. Metabolic profiling of bile acids in human and
mouse blood by LC-MS/MS in combination with phospholipid-depletion solid-phase extraction.
Anal Chem 2015; 87:1127–36. doi: 10.1021/ac503816u

54.   Straniero S, Laskar A, Savva C, Härdfeldt J, Angelin B, Rudling M. Of mice and men: Murine bile
acids explain species differences in the regulation of bile acid and cholesterol metabolism. J
Lipid Res 2020; 61:480–91. doi: 10.1194/jlr.RA119000307

55. Sayin SI, Wahlström A, Felin J, Jäntti S, Marschall HU, Bamberg K, Angelin B, Hyötyläinen T,
Orešič M, Bäckhed F. Gut microbiota regulates bile acid metabolism by reducing the levels of
tauro-beta-muricholic acid, a naturally occurring FXR antagonist. Cell Metab 2013; 17:225–35.
doi: 10.1016/j.cmet.2013.01.003

56. Dawson JA, Mallonee DH, Bjorkhem I, B P. Expression and characterization of a C24 bile acid
7a-dehydratase from *Eubacterium* sp. strain VPI 12708 in *Escherichia coli*. J Lipid Res 1996;
37:1258–67.

57. Laing E, Mersinias V, Smith CP, Hubbard SJ. Analysis of gene expression in operons of
Streptomyces coelicolor. Genome Biol 2006; 7:R46. doi: 10.1186/gb-2006-7-6-r46

58. Marion S, Desharnais L, Studer N, Dong Y, Notter MD, Poudel S, Menin L, Janowczyk A, Hettich
RL, Hapfelmeier S, et al. Biogeography of microbial bile acid transformations along the murine
gut. J Lipid Res 2020; 61:1450–63. doi: 10.1194/jlr.RA120001021

59. Wolf PG, Byrd DA, Cares K, Dai H, Odoms-Young A, Gaskins HR, Ridlon JM, Tussing-Humphreys
L. Bile Acids, Gut Microbes, and the Neighborhood Food Environment—a Potential Driver of
Colorectal Cancer Health Disparities. mSystems 2022; 7. doi: 10.1128/msystems.01174-21

60. National Research Council. Nutrient Requirements of the Mouse. Nutr Requir Lab Anim 1995;
:80–102.

61. Uchimura Y, Wyss M, Brugiroux S, Limenitakis JP, Stecher B, McCoy KD, Macpherson AJ.
Complete Genome Sequences of 12 Species of Stable Defined Moderately Diverse Mouse
Microbiota 2: TABLE 1 . Genome Announc 2016; 4:4–5. doi: 10.1128/genomea.00951-16

62. Team RC. R: A language and environment for statistical computing. 2023;

63. RStudio Team (2020). RStudio: Integrated Development for R. 2020;

Figures

	Host		Bile Acid name	Abbreviations	C-3	C-6	C-7	C-12
Human and rodent	Primary		Cholic acid	CA	α -OH		α -OH	α -OH
			Chenodeoxycholic acid	CDCA	α -OH		α -OH	
			Ursodeoxycholic acid*	UDCA	α -OH		β -OH	
	secondary		Deoxycholic acid	DCA	α -OH			α -OH
			Lithocholic acid	LCA	α -OH			
			3-oxocholic acid	3-oxoCA	=O		α -OH	α -OH
			3-oxodeoxycholic acid	3-oxoDCA	=O			α -OH

		Name	Abbreviation				
		3-oxochenodeoxycholic acid	3-oxoCDCA	=O		α -OH	
		3-oxolithocholic acid	3-oxoLCA	=O			
		3-oxoursodeoxycholic acid†	3-oxoUDCA	=O		β -OH	
		7-oxodeoxycholic acid	7-oxoDCA	α -OH		=O	α -OH
		7-oxolithocholic acid	7-oxoLCA	α -OH		=O	
		12-oxochenodeoxycholic acid	12-oxoCDCA	α -OH			=O
		12-oxolithocholic acid	12-oxoLCA	α -OH			=O
		Isoursodeoxycholic acid†	isoUDCA	β -OH		β -OH	
Rodent	primary	α-Muricholic acid	αMCA	α -OH	β -OH	α -OH	
		B-Muricholic acid	βMCA	α -OH	β -OH	β -OH	
	secondary	Murideoxycholic acid	MDCA	α -OH	β -OH		
		3-oxo-α-muricholic acid†	3-oxoαMCA	=O	β -OH	α -OH	
		3-oxo-β-muricholic acid†	3-oxoβMCA	=O	β -OH	β -OH	
		3-oxomurideoxycholic acid†	3-oxoMDCA	=O	β -OH		
		6-oxo-α-muricholic acid†	6-oxoαMCA	α -OH	=O	α -OH	
		6-oxo-β-muricholic acid†	6-oxoβMCA	α -OH	=O	β -OH	
		6-oxomurideoxycholic acid‡	6-oxoMDCA	α -OH	=O		
		7-oxomurideoxycholic acid†	7-oxoMDCA	α -OH	β -OH	=O	
		ω-Muricholic acid†	ωMCA	α -OH	α -OH	β -OH	

Figure 1. List of deconjugated human and rodent bile acids (BAs) discussed here. The characteristic that distinguishes BAs is the presence of a hydroxyl group at the C-3, C-6, C-7 and/or C-12 position. The hydroxyl groups can be in α- or β-conformation, oxidised into a ketone group, or fully removed (dehydroxylated BA). CA and CDCA are primary BAs of both humans and rodents whereas MCAs are exclusively produced by rodents. (*) UDCA is a primary BA in rodents while it is a secondary BA in humans where the gut microbes epimerise it from CDCA. (†) Bile acids that might have been detected in this study but for which no standards were available. (‡) Also known as 6-oxolithocholic acid (6-oxoLCA). See Supplementary Table 3 for full chemical names of the BAs.

815
816 **Figure 2.** *In vitro* **7-dehydroxylation of cholic acid. The 7-dehydroxylation of CA was tested in (A)** *Clostridium scindens* **ATCC**
817 **35704, (B)** *C. scindens* **VPI 12708 and (C)** *Extibacter muris* **DSM 28561 (SJ24) over time. All strains were grown anaerobically**
818 **in BHIS-S containing 100 µM CA. Bile acids were extracted from the suspended biomass. Error bars represent the standard**
819 **deviation of the mean of biological triplicates.**

Figure 3. *In vitro* 7α-dehydroxylation of CDCA. The transformation of CDCA into secondary bile acids was tested with and without co-induction with 100 µM of ^{13}C-CA. 200 µM of CDCA were used based on previous experiments by Marion e*t al.* (A) *Clostridium scindens* ATCC 35704, (B) *C. scindens* VPI 12708 and (C) *E. muris* DSM 28561 (SJ24). Bile acids were extracted from the suspended biomass. Error bars represent the standard deviation of the mean of biological triplicates.

828
Figure 4. *In vitro* **7-dehydroxylation of UDCA. The transformation of 100 µM of UDCA into secondary bile acids was tested**
with and without co-induction with 100 µM of [13]**C-CA. (A)** *Clostridium scindens* **ATCC 35704, (B)** *C. scindens* **VPI 12708 and**
(C) *Extibacter muris* **DSM 28561 (SJ24) were grown anaerobically in BHIS-S. Bile acids were extracted from suspended**
biomass. Two compounds were detected that could not be identified due to missing standards but their oxidative state
can be estimated based on their ionised mass. X-oxoUDCA had the same mass as other bile acids with one ketone group
and one hydroxyl group. The other unidentified compound had the same mass as UDCA and therefore it is likely to be an
isoform with a 3β conformation. The retention times for these compounds was unique and therefore could not be
identified further. Concentration values of the unknown BAs could only be estimated for the same reason. Error bars
represent the standard deviation of the mean of biological triplicates.
838

839
Figure 5. *In vitro* **7-dehydroxylation of αMCA. The transformation of 100 μM of αMCA into secondary bile acids was tested**
with and without co-induction with 100 μM of ^{13}**C-CA. (A)** *Clostridium scindens* **ATCC 35704, (B)** *C. scindens* **VPI 12708 and**
(C) *Extibacter muris* **DSM 28561 (SJ24) were grown anaerobically in BHIS-S. Bile acids were extracted from suspended**
biomass. Several compounds were detected that could not be identified due to missing standards but their oxidative state
can be estimated based on their ionised mass. X- or Y- oxoαMCA had the same mass as other bile acids with one ketone
group and two hydroxyl groups. The other unidentified compound had the same mass as secondary bile acids that have
been dehydroxylated (-1 -OH), have one ketone group and one hydroxyl group. The retention times for these compounds
did not correspond to that of any known standard. Concentration values of the unknown BAs could only be semi-
quantitative for the same reason. Error bars represent the standard deviation of the mean of biological triplicates.
849

Figure 6. *bai* gene expression in the presence of CA (first panel) or in that of various primary BAs either uninduced or and
co-induced with [13]C-CA. Data for the CA-only condition correspond to the pooled expression results from the two sets of
experiments (referred to as uninduced and co-induced) that both include this condition, as described in the text. The
expression (normalised to at least three reference genes) is relative to the no BA condition including an equivalent volume
of solvent (ethanol). (A) *Clostridium scindens* ATCC 35704, (B) *C. scindens* VPI 12708 and (C) *Extibacter muris* DSM 28561
(SJ24) gene expression of *baiCD*, *baiE* (part of the *bai* operon), and accessory genes *baiJ* and *baiO* (only in *E. muris* DSM
SJ24) was measured. CA, UDCA, αMCA, βMCA and the co-inducing [13]C-CA were used at 100 μM, CDCA was used at 200
μM. A detailed view of the *C. scindens* VPI 12708 uninduced vs 13C-CA co-induced in the CDCA group is found in

Supplementary Figure 4. Coloured dots represent the average and error bars represent the standard deviation of 12
replicates. Some error bars may look elongated due to the logarithmic scale of the Y axis. (***) indicates a *p*-value < 0.001
in a linear model analysis comparing the uninduced vs ¹³C-CA co-induced factor for each *bai* gene.

Figure 7. *In vitro* transformation of BA cocktails by *E. muris* DSM SJ24. The transformation of several BAs at 50 μM by *E.
muris* DSM 28561 (SJ24) was tested anaerobically in 25 mL of BHIS-S. Different cocktails were prepared based on their
similarity. Tauro-conjugated BAs (A). Oxidised BAs (B), Sulfonated BAs (C) or the secondary BA ωMCA (D) which is present
in the murine BA pool. Three time points were taken for BA extraction from suspended biomass. An additional sample of
1 mL was taken at 16 hours for RNA extraction and RT-qPCR analysis. Minute concentrations (<2 μM) of CA (A) and βMCA
(B) were detected, most likely as as impurities from the standards used as they were detected from time 0 and
concentrations remained stable. Error bars represent the standard deviation of the mean of biological replicates.

Figure 8. *In vitro* 7-DH-ion of CA by *E. muris* SJ24 amended with cecal content. The transformation of 100 μM of CA into
secondary BAs was tested with and without the amendment of cecal content. A control group with only CA (A) was used
as intra-assay reference vs (B) cecal content from germ-free mice and (C) cecal content from sDMDMm2 mice. 25 mg of
cecal content were added to 25 mL of BHIS-S. The mass imbalance between the added CA (100 μM) and the measured

(aprox. 80 µM) can be attributed to the presence of CoA- forms that cannot currently be quantified due to lack of
standards. Error bars represent the standard deviation of biological triplicates. Data from the control groups of this
experiment can be found in Supplementary Figure 7.

Figure 9. *In vivo E. muris* SJ24 *bai* expression. The expression (normalised to at least three reference genes) is relative to
a non-colonised sDMDMm2 control group. Each bar represents the average value from five biological replicates with four
technical replicates each. Error bars represent the standard deviation of the mean.