1   **Finding coarse and fine scale population structure in a coastal species: population**

2   **demographics meets genomics**

3
Running title: Geographic scale effect on population structure

Diana Catarino[1,2], Per Erik Jorde[3], Lauren Rogers[4], Jon Albretsen[3], Marlene Jahnke[5], Marte

Sodeland[1], Ida Mellerud[3], Carl Andre[5], Halvor Knutsen[1, 3]

[1]Centre for Coastal Research, Department of Natural Sciences, University of Agder, 4630

Kristiansand, Norway

[2] Okeanos – Institute of Marine Sciences, University of the Azores, Rua Professor Doutor

Frederico Machado 4, 9901-862 Horta, Portugal

[3]Institute of Marine Research, Nye Flødevigveien 20, 4817 His, Norway

[4]Alaska Fisheries Science Center, National Oceanic and Atmospheric Administration, 7600 Sand

Point Way NE, Seattle, WA 98115, USA

[5]Department of Marine Sciences – Tjärnö, University of Gothenburg, 45296 Strömstad, Sweden

*Correspondence:

e-mail: dscatarino@gmail.com; diana.catarino@uia.no

Abstract

Population genetic studies often focus on patterns at a regional scale and use spatially

aggregated samples to draw inferences about population structure and drivers, potentially

masking ecologically relevant population sub-structure and dynamics. In this study we use a

multidisciplinary approach combining genomic, demographic, and habitat data with an

oceanographic particle drift model, to unravel the patterns of genetic structure at different

scales in the black goby (*Gobius niger*) along the Norwegian coast. Using a high-density

sampling protocol, we observed restricted gene flow both at a surprisingly fine (kms) and large

(100s km) scale. Our results showed a pattern of isolation by distance related to the level of

exposure along the Skagerrak coast, where sheltered sampling stations had an overall level of

genetic divergence about three times higher (F_{ST} =0.0046) than levels observed among

exposed samples (F_{ST} =0.0015). These results were corroborated by demographic analyses

which showed that population-fluctuations decrease in synchrony with distance at much

smaller scales for sheltered samples (20 km) than for exposed sites (80 km), suggesting higher

population connectivity among exposed sites. We also found a pronounced genetic

discontinuity between populations along the Norwegian west and east coasts, with a sharp

"break" around the southern tip of Norway, likely driven both by lack of habitat and by

oceanographic features.

Keywords: connectivity; Isolation by distance; habitat discontinuity; oceanography; marine

fish; demographic synchrony

Introduction

Many marine species are broadcast spawners and disperse at early life stages by drifting with ocean currents. Traditionally, these species were expected to show little or no genetic population structure across small (kms) or even large geographic scales (ocean regions; i.e. Longmore et al., 2014). Although for some species this holds true, in the past decades this paradigm has considerably shifted (Palumbi, 2004; Hauser & Carvalho 2008), and it has been shown for many species that phenomena such as ocean currents, bathymetry, local adaptation, habitat fragmentation, or reduced migration, limit genetic connectivity across different spatial and temporal levels (i.e. Catarino et al., 2015; Kelley et al., 2016; Barth et al., 2019). In fact, many species with extensive geographic ranges may consist of genetically separated populations despite an apparent homogeneous distribution. Therefore, understanding and identifying the geographic scales of connectivity is a crucial step for basic and applied ecology of marine species (Cowen et al., 2006; Selkoe et al., 2016; Benestan et al. 2021), since this information is essential to improve conservation efforts such as the design of marine reserve networks at suitable scales (Almany et al., 2009; Mertens et al., 2018) and to define stock boundaries for fisheries management (Bernatchez et al., 2017; Quintela et al., 2020; Lindegren et al., 2022). However, detecting the magnitude and geographic scale of connectivity of fish in the marine environment is not always an easy task. Many studies may have missed existing population divergence because population genetic studies usually focus on patterns at a regional scale and use spatially aggregated samples to draw inferences about population structure and drivers, potentially masking ecologically relevant population sub-structure and dynamics. Given that most marine species have wide ranges, spatial sampling for genetic studies is often geographically coarse, since financial and technical constraints limit the number of sampled specimens and their geographical coverage (Martinsohn et al., 2019). Apparent homogeneity over large areas in an initial screening may therefore yield the impression of panmixia and discourage further studies at finer geographic scales. While a coarse-scale sampling strategy may be adequate for uncovering the major population genetic

patterns in a species, it risks overlooking finer scale patterns that may hint at population

processes of biological relevance for local populations (Barth et al., 2019; Benestan et al.,

2021).

Many species, marine or terrestrial, live in habitats that in the context of individual movement

and dispersal may be considered two-dimensional, meaning that dispersing individuals may

reach neighbouring population(s) along several routes. Theoretical analyses found that in two

dimensional habitats, gene flow is particularly effective in counteracting genetic divergence

(Kimura & Weiss 1964; Rousset, 1997; Slatkin, 1994), implying that population genetic

structure may be weak and difficult to detect in the face of sampling noise (Waples, 1998;

Palumbi, 2004). A more powerful approach to detect mechanisms that restrict gene flow is to

consider species that occupy a linear, one-dimensional habitat where theory predicts that

gene flow is a less potent homogenizing force and where genetic structure should be more

pronounced than in a two-dimensional habitat when the total amount of gene flow is equal

(Rousset, 1997). In the marine environment species living along the coastline approximate

such a linear habitat, in contrast to the open ocean or in more complex seascapes which are

more two-dimensional. In linear habitats, limited gene flow caused by habitat discontinuities,

environmental drivers, or by geographic distance, is expected to result in more pronounced

and thereby more easily detectable genetic structures (Kanno et al., 2011).

Information on genetic structure provides important clues to the pattern and magnitude of

gene flow, but genetic data alone are insufficient to determine the demographic effects on

populations. While dispersal will leave genetic signatures due to gene flow, the exchange of

individuals among locations or local populations can also affect spatial population dynamics, as

measured by local or regional variation in abundance through time. Dispersal acts to

synchronize population dynamics across space (Bjørnstad et al. 1999), and the spatial scale of

population synchrony has been used to delineate population units in marine ecosystems

(Östman et al. 2017). Analysis of spatial population synchrony can therefore be used alongside

analyses of genetic structure to provide insights to the scales and processes affecting dispersal

and population connectivity in a more powerful approach.

In the present study we adopted a multidisciplinary approach to unravel gene flow and

population connectivity at coarse and fine geographic scales of a coastal fish species. Using a

geographically fine-scaled sampling protocol, we combine genomic data (ddRAD Seq) of the

black goby with analysis of spatial population dynamics from abundance time-series, as well as

oceanographic modelling of pelagic larval drift, and habitat analyses to identify features that

influence local- and regional-scale population connectivity and genetic structure. Specifically,

we investigate the effects of scale consideration and sampling design on genetic and

demographic inferences and the ecological drivers of the patters observed.

Material and Methods

Study Species

The black goby, *Gobius niger* Linnaeus, 1758 is widely distributed in the Eastern Atlantic and

Mediterranean Sea, from Mauritania (North Africa) to the eastern Atlantic coast northwards to

Norway and into the Baltic Sea (Froese & Pauly, 2021). The black goby is a small coastal fish

growing up to 17 cm total length and can be found mostly on sandy and muddy bottoms but

also on hard bottom among algae (Moen & Svensen, 2000). The species inhabits bottoms from

2 to 70 m depth but is most common in depths less than 30m. It can inhabit lagoons with

brackish water down to 10‰ salinity (Penthon, 2005). Sexual maturity is reached at around 2

years of age, followed by a sedentary benthic adult life (Miller, 1986). Black goby adult males

build and protect a nest where the females deposit their eggs. Spawning begins when the

water temperature reaches 12°C, and the incubation of the eggs at this temperature lasts for

about 20 days (Vaas et al., 1975). The eggs are guarded by the male until they hatch. After

hatching, the duration of the planktonic larval phase is approximately 28 days (Planes, 1998).

The species has no commercial value, but it is ecologically important in the coastal food web

complex as prey for commercially important species such as cod, haddock, and trout (e.g. Hop

et al., 1992; Knutsen et al., 2001).

Study area and sampling

In this study we focused on the Norwegian coastal system from the eastern border with

Sweden, along the southern coast of Norway, and up to Ålesund (62° N) in western Norway.

Most of the sampling sites are on the Skagerrak coast, which lies between the North Sea and

the Kattegat, bordered by Denmark, Sweden, and southern Norway (Figure 1). The Norwegian

coast is characterized by an indented landscape with numerous islands, islets and enclosed

bays, and cut by several fjords that can extend for several kilometres. The rough coastline and

the scattered landmasses create a dynamic and complex oceanography along the coast, but

with a general ocean current pattern flowing westwards along the Norwegian Skagerrak coast

(the Norwegian Coastal Current, NCC; Albretsen et al., 2012).

For more than a century, since 1919, the Institute of Marine Research has conducted an

autumn beach seine survey at a set of fixed stations along the Norwegian Skagerrak coast

(Stenseth et al., 1999; Table S1). The beach seine is 40 m long, with a stretched mesh size of

1.5 cm, and samples an area up to 700 m^2 of nearshore (< 15 m depth) habitat (Barcelo et al.,

2016; Rogers et al., 2014). In this survey black goby abundance has been counted as number of

individuals per haul yearly since 1989 at more than 130 stations distributed among 13 main

regions from Torvefjorden (TOR) in the south to the Swedish border in the east (Figure 1;

Table1; Table S1). Biogenic habitat type (e.g., eelgrass, green or brown macroalgae) and

coverage (scaled from 1-5 subjectively: 1 = bare substrate to 5 = full coverage) have been

characterized at each haul station (Table S1). Stations range from sheltered locations inside

fjords to exposed areas on the open Skagerrak coast and have been classified as "sheltered" or

"exposed" according to their degree of exposure to the open sea (see Rogers et al., 2014).

Samples of *G. niger* for genetic analyses were obtained from a total of 17 different locations

(Figure 1, Table 1 and S1) between June and October of 2018. These locations included the 13

main locations described above along the Skagerrak coast (hereafter referred to as SK) and

representing 37 hauled stations from the beach seine survey, and 4 additional locations from

Egersund (EGE) in the southwest, and along the Norwegian West coast (WC), including in

Stavanger (STA), Manger (MAN) and Ålesund (ALE). Fin clips were taken from each fish and

stored in 96% ethanol until DNA extraction.

Fish samples were collected in accordance with Norwegian regulations and with all permits

required by the Norwegian Government. The studied species (*G. niger*) is not listed in CITES

and it is listed as "Least Concern" by the International Union for Conservation of Nature (IUCN;

Carpenter et al., 2015).

Laboratory procedures

Genomic DNA was extracted using the E-Z® 96 Tissue DNA kit (Omega, USA) following

manufacturer's procedures. DNA was quantified on a Fluoroskan FL (Thermo Scientific, USA)

using the Quant-iT Broad Range dsDNA assay kit (Thermo Scientific, USA), and integrity was

checked by gel electrophoresis on a Multiskan SkyHigh spectrophotometric microplate reader

(Thermo Scientific, USA). Double-digest RADseq libraries (ddRAD, Peterson et al., 2012) were

prepared using restriction enzymes Sbf1 HF/Sph1 HF. The 95 samples (with 3 replicates) and

one blank per plate were digested at 37°C for 45 minutes, and then combined in a ligation

mixture with T4 ligase, a general P2 adapter and P1 barcoded adapter and incubated at 16°C

for 1 hour 30 minutes, 22°C for 1 hour and 30 minutes and heat inactivated at 65°C for 10

minutes. Successfully ligated samples were size selected (250-600bp) on a Blue Pippin (Sage

Science), followed by the AMPure magnetic bead clean-up (Beckman Coulter, USA) and then

amplified by PCR: 30s at 98°C, 30s at 98°C followed by 14 cycles of 10 seconds at 98°C, 30

seconds at 65°C, and 30 seconds at 72°C, followed by a final extension of 2 minutes at 72°C.

The resulting PCR was cleaned with the AMPure magnetic bead clean-up (Beckman Coulter,

USA). Library preparation success was then evaluated by running the libraries on a Bioanalyzer

(Agilent, USA), and quantified on Qubit (Thermo Scientific, USA). Libraries were diluted to

14pM, spiked with 5% PhiX and sequenced on an in-house Illumina MiSeq platform (160bp

Paired End), by using one lane per plate and the MiSeq v3 reagents kit.

Bioinformatics and statistical analyses

Sequencing reads for individual specimens were retrieved using the *process_radtags* module in

the Stacks software package (Catchen et al., 2013). De-novo assembly of reads from all

specimens, as well as variant detection and initial SNP filtering was conducted using the

ustacks, *cstacks*, *sstacks* and *populations* modules in the same software package. An average

of 549'980 PE 150bp reads were utilized per specimen in stacks formation. Minimum stack

depth (m) for SNP detection was set to 2.

Using VCFtools (Danecek et al., 2011), an initial five different SNPs datasets were generated,

each allowing different levels of missing data: 10, 30, 50, 70, and 80%, respectively. Comparing

the analyses of these datasets showed that the main genetic pattern found did not change

with different levels of missing data. Therefore, the single, final dataset was filtered to retain

SNPs with a minimum read depth of 8, a minimum allele count of 2 and a maximum of 50%

missing data for locus and individuals. Loci showing deviations of HWE at <0.001 were also

removed and the dataset was further filtered to remove loci exhibiting linkage disequilibrium

(LD at 0.5 level) with PLINK 1.9 (Purcell et al., 2007). In the end a total of 611 SNPs and 292

individuals were retained for downstream analyses.

Summary statistics, sample sizes, observed and expected heterozygosity, and F_{IS} were

calculated using the "diveRsity" (Keenan et al., 2013) R package (R Core Team, 2021). Variation

in heterozygosity across sites was investigated using a linear model in R. BayeScan (v 2.1., Foll

& Gaggiotti 2008) and PCAdapt (Luu, Bazin, & Blum, 2017; Privé et al., 2020) were used to test

if loci deviated from neutrality across localities, and between west coast and Skagerrak

samples. BayeScan was run by setting sample size to 10,000 and the thinning interval to 50 as

suggested by Foll & Gaggiotti (2008). In PCAdapt population structure was first assessed via

PCA, and outliers were thereafter detected with respect to how they relate to population

structure. The number of principal components (K=2) to identify SNPs deviating from neutrality

was then selected by Cattell's rule (Cattell, 1966). The resulting *P*-values were corrected for

false positives using false-discovery rate (FDR) approach at a 0.05 threshold in both cases

(Benjamini & Hochberg, 1995). Following a consensus approach, only those loci flagged as

outliers by both methods were considered as being under selection (Narum & Hess, 2011).

Pairwise F_{ST} (Weir & Cockerham 1984) was calculated using the Genepop R package (v1.1.7,

Rousset, 2008), and statistical significance of pairwise F_{ST} tests was assessed by G-tests for

allele frequency differences, using 10 000 dememorizations and batches, and 1000 iterations

per batch. *P*-values in multiple tests were evaluated for significance using the FDR approach.

Genetic clustering was investigated by discriminant analyses of principal components (DAPC,

Jombart et al., 2010) using the adegenet R package (Jombart & Ahmed, 2011), to provide a

graphic description of the genetic divergence among sample locations in multivariate space.

The analyses were performed for the whole dataset and for Skagerrak and Egersund

separately. The number of principal components kept was determined from the number of

PCAs needed to explain approximately 80% of variation. Optimal clustering was retrieved with

the function *find.clusters* where the best number of clusters is indicated by the lowest

Bayesian information criterion (BIC; Schwarz, 1978), inferred by the *k*-means algorithm with

10^6 simulations.

Isolation by distance (IBD) was analysed for the whole dataset and for Skagerrak only, by

testing for a correlation between pairwise genetic $F_{ST}/(1 - F_{ST})$ and geographic distances

(overwater distances) using Mantel tests in the Genepop R package, with 10 000 permutations.

Overwater distances were calculated using the marmap (Pante & Simon-Bouhet, 2013) and

fossil (Vavrek, 2011) R packages, and using bathymetric data (Amante & Eakins, 2009) hosted

by NOAA (https://www.ngdc.noaa.gov/mgg/global/relief/ETOPO1/docs/ETOPO1.pdf)

The geographic coarse-scale genetic structure was further characterized by non-linear

regression of F_{ST} against distance along the coast. The analysis was anchored to the geography

by considering only pairwise comparisons with the easternmost location (Hvaler, HVA), and

then fitting an exponential (sigmoid) function to those F_{ST} and distance measures:

$$F_{ST} = \frac{a}{1 + e^{-b*distance-c}},$$

where a, b and c are parameters estimated with the *nls* function in R.

The geographical fine-scale genetic structure of the black goby was characterized using

pairwise genetic differences between individuals (Rousset, 2000), as calculated by the

Genepop R package (estimator ê). This analysis was restricted to the more densely sampled

Skagerrak coast and performed at the beach-seine haul station level. Possible effects of ocean

currents on genetic structure were tested by classifying beach seine stations as "exposed" and

"sheltered" with respect to the coastal current, following Rogers et al. (2014), and carrying out

the analysis separately for pairs of individuals from sheltered and exposed locations (Table S1,

Figure S1). The geographic distance between individuals were calculated as distance overwater

between beach seine stations. Genetic differences (ê-estimates) for pairs of individuals from

sheltered stations were then regressed (linear regression, *lm* function in R) against geographic

distance, and similar for exposed pairs. The slopes of the regressions were tested for

significance with permutations tests. The tests randomly permuted the genetic distance (ê)

matrices repeatedly (10000 times) and regression slopes were calculated for the permuted

data. *P*-values were calculated as the fraction of permutations that resulted in a slope that was

as large or larger than the observed one.

Spatial population synchrony and decorrelation scale

Spatial population synchrony, defined as the spatial covariation in population density

fluctuations (Bjørnstad et al., 1999), was calculated by taking the pairwise correlation in

natural log-transformed catches among each pair of sampling stations, and relating pairwise

correlations to the distance between stations. A constant of 1 was added to each count to

avoid taking the logarithm of zero. Comparisons were limited to those stations sampled a

minimum of 15 years and having at least 10 years with positive (non-zero) catches (n = 132

stations).

An exponential decay model (Bjørnstad et al., 1999) was fitted to pairwise correlations by

distance, measured as distance over water between stations:

$$\rho(d) = \rho_r + \rho_0 e^{-\left(\frac{d}{v}\right)},$$

where $\rho(d)$ is the pairwise correlation at distance d, ρ_r is the asymptotic correlation, or

background regional correlation in recruitment, and $\rho_r + \rho_0$ is the estimated correlation at zero

distance, following Rogers et al. (2014). The parameter v ("decorrelation scale") describes

distance at which the pairwise correlation between time series is reduced to ~37% (e^{-1}) of that

at zero distance, relative to the background regional level of correlation. To test the hypothesis

that abundance at exposed stations would covary over longer distances than sheltered

stations due to increased connectivity, models were also fit separately to pairs of exposed

stations and pairs of sheltered stations. Confidence intervals (90%) were estimated based on

10000 bootstrap replicates, following Bjørnstad et al. (1999).

Habitat associations

Habitat associations for black goby were investigated using the Skagerrak beach seine time-

series (Table S1). Specifically, we tested for effects of habitat type and percent cover on

abundance of black goby in beach seine hauls using a negative binomial generalized mixed

effects model which included a random intercept by station. Data from all stations surveyed

for at least 15 years (n = 137) were used in this analysis, for a total of 3930 individual

observations of black goby abundance. Although recorded as a categorical variable with five

categories ("bare substrate" to "full cover"), coverage was treated in the model as a

continuous variable (ranging from 1 to 5) after preliminary model runs indicated an

approximately linear relationship between abundance and coverage. Models were fitted using

the lme4 package (Bates et al., 2015) in R.

Oceanographic modelling

A combination of high-resolution hydrodynamical modelling and particle-tracking was applied

to investigate dispersal potential of larvae from all beach seine locations, and additional sites

around the southern tip of Norway, and in the west encompassing the genetic sampling

stations between Egersund and Stavanger (see Table S1). Our circulation model system

consisted of three high-resolution fjord models run in parallel with input along the open

boundaries from the Norwegian coastal model NorKyst800 (Asplin et al., 2020). The highest

resolution models all applied 160m resolution in the horizontal, and similar experiments and

validation analysis are presented in e.g. Dalsøren et al., 2020. We applied the open-source

Regional Ocean Modeling System (ROMS) (Haidvogel et al., 2008; Shchepetkin & McWilliams,

2005) (see also http://myroms.org), which is a state-of-the-art, three-dimensional, free-

surface, hydrostatic, primitive equation ocean model that uses generalized terrain-following s-

coordinates in the vertical. The results from the three model grids using 160m resolution

covering the inner Skagerrak, the southernmost part of the Norwegian coast and the

southwestern fjords, respectively, were merged into one large grid before being used as input

in the particle-tracking model.

Our drift simulations of larvae-particles were conducted by using the open access Lagrangian

Advection and Diffusion Model (LADIM; https://github.com/bjornaa/ladim). The model solves

the Lagrangian equation of motion using a 2nd-order Runge-Kutta scheme. Due to limited

knowledge on larval swimming behaviour, we initialized the particles in 1, 2, 3 and 4m depth

and left them without any vertical movement. 250 particles were released in each depth at all

204 locations which gives 1000 particles per station. To include slightly different physical

environments, we performed three identical simulations initiating June 2nd, July 1st and August

1st, all in 2018 as a representative year, and followed every particle for 25 days. A small

horizontal random walk was also implemented to ensure variability between the particles and

to represent small-scale turbulence.

Results

Population genetic patters

The final dataset consisted of 292 individuals from 17 main localities genotyped for a total of

611 SNPs, with no deviations from HWE or LD. Genetic variability (expected heterozygosity, H_e)

ranged from 0.139 for SAN locality to 0.156 at MAN (Table 1) and displayed a significant

increasing trend from east to west (linear regression of H_e on distance over water from the

easternmost locality, Hvaler: slope = $1.285*10^{-5}$-; R^2 = 0.3731; P = 0.0055).

BayeScan identified three possible outlier loci out of the 611 SNPs analysed for the whole

dataset, while no outlier loci were identified when comparing West Coast *vs*. Skagerrak.

PCAdapt identified a total of 15 outlier loci. Following a consensus approach, two SNPs (SNP

9075_98 and 13326_52) were identified by both methods as outlier candidates under

selection. Pairwise F_{ST}, DAPC and IBD analysis were performed both with and without the two

candidate outlier loci and since no significant differences were observed in the results (Figure

S2), all loci were retained for all analyses presented below.

Cluster analyses (DAPC) comprising the whole dataset clearly showed two main clusters

separated along DF1 (Figure 2a), one grouping together the West Coast samples (ALE, MAN

and STA), and a second grouping all Skagerrak samples and Egersund. This result is supported

by the lowest BIC, which indicated that K=2 was the most likely number of clusters (Figure S3).

Using only Skagerrak and Egersund samples in a separated DAPC analyses (Figure 2b), no

further pattern stood out with samples showing no clear separation, and many locations

overlapping. This genetic pattern is also supported by F_{ST} analyses, where significant genetic

differentiation was found for overall sampling localities (F_{ST} overall =0.0181, P<0.001). Pairwise

F_{ST} values ranged from -0.0065 to 0.0657 averaged over the 611 SNP considered, with the

highest values observed for the pairwise comparisons between West Coast localities and

Skagerrak/Egersund, which were highly significant (Table 2; overall WC *vs*. SK/EGE F_{ST}= 0.0400,

P<0.001). In contrast, pairwise comparisons showed no significant differences among West

Coast localities and Stavanger nor among Skagerrak and Egersund (Table 2). Analysis of IBD

showed that for the entire Norwegian coast, genetic divergence $F_{ST}/(1-F_{ST})$ increased with

increasing distance (Figure 3a; Mantel, $P <0.001$), while no such relationship was observed

within Skagerrak samples only (Figure 3a, Mantel, $P>0.05$). This suggests that the divergence

pattern found is mostly driven by the West Coast samples and is illustrated with the sigmoid

curve in Figure 3b, evidencing a sharp increase in F_{ST} around the southern tip of Norway,

located between the EGE and STA localities.

Fine-scale genetic structure patterns

Although pairwise comparisons among Skagerrak sampling localities were non-significant

(Table 2), and low levels of genetic divergence were found among the Skagerrak sampling

localities across all loci, the overall divergence was nevertheless highly significant (overall F_{ST}

$=0.0040$, $P<0.001$), suggesting the possibility of some cryptic genetic structure within

Skagerrak. To look for fine-scale patterns of genetic divergence within Skagerrak, this dataset

was analysed separately for IBD among individuals. Although the Skagerrak samples did not

show any patterns of IBD when using the whole dataset with the individuals grouped in main

sampled localities (Figure 3a, $P> 0.05$; Table 1), when separating according to the level of

exposure for each station, two different patterns of genetic differentiation within Skagerrak

emerged. First, the overall level of genetic divergence among sheltered samples was about

three times higher (overall sheltered F_{ST} $=0.0046$) than among exposed samples (overall

exposed F_{ST} $=0.0015$). Second, genetic divergence increased significantly with distance for

sheltered stations (Figure 4; $P = 0.0018$; slope = $3.446*10^{-5}$), but not for exposed stations

(Figure 4; $P = 0.3555$; slope = $2.298*10^{-6}$).

Spatial decorrelation scale

Station-specific fluctuations of black goby abundance in the Skagerrak were positively

correlated (i.e., synchronous) at nearby stations, with synchrony decreasing with increasing

distance between stations (Figure 5). The spatial decorrelation scale (v) considering both

exposed and sheltered stations was estimated as 27.9 km (90% CI = 13.8 - 94.8 km). When only

sheltered stations were considered, the spatial decorrelation scale was 20 km (90% CI = 2.9 –

266 km), which was substantially smaller than the decorrelation scale among exposed stations

of 81 km (90% CI = 15.1 – 254 km), although confidence intervals were overlapping.

Habitat associations

The best model for black goby abundance in the Skagerrak beach seine survey included effects

of biogenic habitat type and cover. Black goby catches significantly varied by habitat type (X^2 =

44.2, df = 5, P < 0.001), with highest catches in eelgrass and mixed eelgrass/brown macroalgae

habitats, and lowest catches in habitats lacking eelgrass, but containing brown macroalgae or

green algae (Figure 6). Black goby catches were also significantly associated with habitat cover,

with higher catches at sites with higher coverage percentage (z = 8.42, P < 0.001).

Oceanographic modelling

The oceanographic modelling shows that the larvae-particles released in the surface layer

inshore and along the Skagerrak coast, from the Swedish border in the East to Egersund in the

West, are subject to three different destinies (Figure 7). (1) A portion is mixed and spread

widely by the NCC, which has a persistent westward flow out of the Skagerrak along the

Norwegian coast. The NCC in the Skagerrak is relatively strong and often has a distinct core

close to the coast. (2) Most larvae-particles entering the NCC have a stronger offshore drift and

were not able to settle along the coast at all. (3) A significant and important portion of the

larvae are retained locally and hatched inside the fjords, suggesting local settlement. The

probability for onshore settlement West of Egersund from larvae-particles released East of

Egersund is very small, since the NCC around the southern tip of Norway, transports the drift

particles offshore, towards deeper water. For larvae-particles released East of the southern tip

of Norway, the NCC acts slightly different with a broader core and occasional long-periodic

eddies interfering the continuous transport northward. The localities around Stavanger

provide the area with some larvae retention even though most of the larvae-drift from the

localities West of Egersund is offshore.

Discussion

Using a multidisciplinary approach, we combined genetic, demographic and habitat data from a long-term data series with an oceanographic particle drift model to unravel the patterns of genetic structure on different geographic scales in the black goby along the Norwegian coast. While geneflow and weak population structure is expected for marine species with high dispersal ability, we observed restricted gene flow both at a surprisingly fine (kilometres) and large/coarse (100kms) scale. Our results show a pattern of isolation by distance along the Skagerrak coast and a pronounced genetic discontinuity between populations along the Norwegian south and west coasts, likely driven by both lack of habitat and oceanography.

Our sampling design focused on the Skagerrak coast, where time series for annual abundance are available, giving the opportunity to investigate fine-scale spatial population patterns both at the genetic and demographic level. At a first glimpse, the general overall pattern showed high levels of connectivity across Skagerrak for both genetic and demographic data. In the Skagerrak, we found no genetic differentiation for pairwise comparisons among locations (Table 2), no IBD, and no clusters were perceived in the DAPC analyses. This is in accordance with the predictions of the oceanographic model where the Norwegian coastal current boosts passive dispersion of larvae promoting connectivity. However, although the overall level of genetic divergence among Skagerrak sample localities was low, it was statistically significant (SK$_{overall}$ F_{ST} =0.0040, P<0.001), suggesting possible hidden population genetic structure. The dense, fine-scale sampling design allowed us enough spatial resolution to examine samples at the individual level and compare stations that are largely sheltered from the coastal current (i.e., located inside small fjords or bays, or behind skerries) and more exposed areas. On the supposition that ocean currents transport pelagic larvae along the coast, we expect that exchange of larvae will be higher among the more exposed localities and, further, if this current-driven larval dispersal is the dominant mechanisms for gene flow in the species, that gene flow should also be higher among exposed localities than among more sheltered ones. These expectations were indeed met, as the genetic IBD analyses yielded a steeper slope for

sheltered than for exposed localities (cf. Figure 4), indicating less gene flow among sheltered

localities.

The observed spatial correlation in abundance, or population synchrony, corroborates the

genetic findings, as has been shown for other coastal species (Östman et al., 2017). First,

nearby stations only a few km apart displayed higher correlations in abundance than stations

situated farther apart on the coast (cf. Figure 5). This finding gives evidence that nearby

populations are indeed connected demographically, as expected if they also exchange genes.

Second, the sheltered stations displayed a more rapid decline in correlation with distance, i.e.

a shorter "decorrelation" scale, than did the exposed areas (20 and 81 km, respectively),

indicating that dispersal is more restricted among sheltered areas. This finding is also in

agreement with genetic data, suggesting a higher level of connectivity among exposed stations

than sheltered ones. Our particle drift model showed some level of particle larvae retention,

mostly among some of the Skagerrak most enclosed fjords (see Figure 7) supporting lower

levels of connectivity or dispersal with outside areas. Nevertheless, power at such small spatial

scales was low to test the findings at the demographic and genetic level, and more information

of larvae dynamics would be necessary before attempting to model very fine scale dispersal on

a nest scale. Still, in the light of the results obtained, it is plausible that exposed stations are

more affected by the Norwegian coastal current promoting higher rates of dispersal translating

in a higher level of population homogenization, while sheltered areas likely are less influenced

by the NCC and experience localized retention processes (i.e. Ciannelli et al., 2010; Myksvoll et

al., 2014; Hansen et al., 2021). This correspondence between genetic and demographic

patterns is relevant for the long-standing discussion about the use of genetic data for inferring

connectivity among natural populations (Lowe & Allendorf 2010; Knutsen et al., 2011), and

yields support to the notion that genetic markers provide information of ecological as well as

evolutionary relevance.

At a larger scale, when considering the entire study area, we found a strong genetic

discontinuity around the southern tip of Norway, between the two sample localities Egersund

and Stavanger (cf. Figure 1). We identified two distinct genetic clusters, one comprising the

west coast samples and the other comprising all samples from Egersund and the Skagerrak

coast, with increased levels of pairwise divergence (F_{ST} =0.0400) between West coast and

Skagerrak samples. Lower levels of genetic diversity were found within the Skagerrak samples,

suggesting restricted geneflow between these two regional groups. Genetic discontinuities, or

"genetic breaks," are characterized by rapid increase in genetic divergence over short

geographic distances (Sotka et al., 2004; Catarino et al., 2015). This particular genetic break,

around the southern tip of Norway, has been previously identified for other fish species, such

as corkwing wrasse (*Symphodus melops*; Knutsen et al., 2013; Blanco Gonzalez et al., 2016;

Mattingsdal et al., 2020) ballan wrasse (*Labrus bergylta*; Seljestad et al., 2020), and even for

some kelp species (Evankow et al., 2019). The area around this break point, while not exactly

localized for any species, is characterized by a bare stretch of sandy beaches (Jæren, about 26

km; Blanco Gonzalez et al., 2016), between Egersund and Stavanger, probably resulting in a

habitat discontinuity for some species. For wrasses, hard bottom substrate is an essential

habitat, required to build nests where females lay their eggs (Darwall et al., 1992; Villegas-Ríos

et al., 2013). Furthermore, telemetry studies showed that wrasses have small home ranges

and individual movements are likely to be restricted to less that 300m (Halvorsen et al., 2021),

suggesting that individual migration across Jæren is unlikely. For the black goby, not much is

known about individual movements, although they share similar life history traits with

wrasses: adult black gobies are mostly sedentary, and males are also territorial, building and

guarding nests where females lay their sticky eggs. Therefore, for both, black goby and

wrasses, dispersal is assumed to be mostly promoted by the pelagic larvae (Knutsen et al.,

2013; Kara & Quignard, 2019). Regarding habitat, the black goby has been described in the

literature as preferring soft bottoms and unvegetated areas (Perry et al., 2018), and with

strong habitat selection shifts based on the presence of predators (Kruschel & Schultz, 2011).

However, although black gobies are occasionally captured in bare substrates, our habitat

analysis clearly showed that the black goby is more abundant in areas with higher vegetation

coverage and prefers eelgrass and mixed eelgrass/brown macroalgae habitats (see Figure 6).

This suggests that habitat discontinuity may also be a good explanation for the lack of gene

flow across the bare stretches of sandy beaches at Jæren for the black goby population,

similarly to what has been found for wrasse species (Blanco Gonzalez et al., 2016; Seljestad et

al., 2020). Habitat discontinuities have been also shown to be a major population subdivision

in different environments and across different taxa (Riginos and Nachman, 2001; Johansson et

al., 2008; Binks et al., 2019) and therefore could be a plausible explanation. Nevertheless,

habitat discontinuity alone does not explain how a species with a long pelagic larval stage

(about 28 days) is unable to drift over a few dozens of kilometres of the stretch of sandy

beaches. Our oceanographic model suggests that particles have a high dispersal potential and

therefore larval transport should be high enough to overcome the physical barrier of lack of

habitat across Jæren. The results obtained with the oceanographic drift model offers a

plausible picture of the drivers promoting the genetic break and restricted geneflow, by

showing that particles that are released near the southern tip of Norway drift offshore towards

deep water, where recruitment of juveniles seems unlikely, restricting the particles exchange

between east and west of the southern. This pattern is driven by the Norwegian coastal

current, which flow westwards along the Skagerrak coast (Albretsen et al., 2012), but drifts

offshore at the southern tip of Norway. The oceanographic particle drift model also shows that

particles released close to Stavanger and Jæren will continue westwards and will not mix with

Skagerrak, preventing genetic connectivity and likely strengthening the barrier effect. The

model shows that few particles/larvae released from the western end of Skagerrak will reach

the Egersund area and this is likely sufficient to homogenise (probably together with some

individual migration) the local gene pool, making Egersund more genetically similar to

Skagerrak samples. The structuring effect of this oceanographic feature in species with early

life stages is also corroborated when comparing with species without such stages. A recent

study showed that for the broadnosed pipefish (*Syngnathus typhle*), which gives birth to live

young, this barrier effect is less pronounced, and that genetic divergence is likely due to

habitat effect and isolation by distance (Knutsen et al., 2022).

Our multidisciplinary approach and fine-scale sampling design allows for detecting the

importance of retention of early life history stages in shaping the genetic structure at finer

geographic scale in coastal waters. Further, oceanographic circulation also seems to play an

important role in the creation of the genetic divergence observed between West coast and

Skagerrak samples. Therefore, our results using these combined approaches, indicated a more

complex scenario of gene-flow restriction suggesting a combination of both habitat

discontinuity but also an important oceanographic component.

References

References

Albretsen J, Aure J, Sætre R, Danielssen DS (2012) Climatic variability in the Skagerrak and
coastal waters of Norway, ICES Journal of Marine Science, 69, 758-763.

Almany GR, Connolly SR, Heath DD, Hogan JD, Jones GP, McCook LJ, Mills M, Pressey RL,
Williamson DH (2009) Connectivity, biodiversity conservation, and the design of marine
reserve networks for coral reefs. Coral Reefs, 28, pages339–351.

Amante C, Eakins BW (2009) ETOPO1 1 Arc-Minute Global Relief Model: Procedures, Data
Sources and Analysis. NOAA Technical Memorandum NESDIS NGDC-24, 19 pp.

Asplin L, Albretsen J, Johnsen IA, Sandvik AD (2020) The hydrodynamic foundation for salmon
lice dispersion modeling along the Norwegian coast, Ocean Dynamics, 70, 1151–1167.

Barceló C, Ciannelli L, Olsen EM, Johannessen T, Knutsen H (2016) Eight decades of sampling
reveal a contemporary novel fish assemblage in coastal nursery habitats. Global Change
Biology, 22, 1155– 1167.

Barth JMI, Villegas-Ríos D, Freitas C, Moland E, Bastiaan S., André C, Knutsen H, Bradbury I,
Dierking JP, Christoph R, Metcalfe D, Jakobsen J, Olsen KS, Olsen EM, Jentoft S (2019)
Disentangling structural genomic and behavioral barriers in a sea of connectivity. Molecular
Ecology, 28, 1394-1411.

Bates D, Maechler M, Bolker B, Walker S (2015) Fitting Linear Mixed-Effects Models Using
lme4. Journal of Statistical Software, 67, 1-48.

Benestan L, Fietz K, Loiseau N, Guerin PE, Trofimenko E, Rühs S, Schmidt C, Rath W, Biastoch A,
Pérez-Ruzafa A, Baixauli P, Forcada A, Arcas E, Lenfant P, Mallol S, Goñi R, Velez L, Höppner M,
Kininmonth S, … Manel S (2021) Restricted dispersal in a sea of gene flow. Proceedings of the
Royal Society B: Biological Sciences, 288.

Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate: a practical and powerful
approach to multiple testing. Journal of the Royal Statistical Society B (Methodological), 57,
289–300.

Bernatchez L, Wellenreuther M, Araneda C, Ashton DT, Barth JMI, Beacham TD, et al (2017)
Harnessing the power of genomics to secure the future of seafood. Trends in Ecology &
Evolution, 32(9), 665–680.

Binks RM, Byrne M, McMahon K, Pitt G, Murray K, Evans RD (2019) Habitat discontinuities
form strong barriers to gene flow among mangrove populations, despite the capacity for long-
distance dispersal. Diversity and Distributions, 25, 298–309.

Bjørnstad ON, Ims RA, Lambin X (1999) Spatial population dynamics: Analyzing patterns and
processes of population synchrony. Trends in Ecology & Evolution, 14, 427–432.

Blanco Gonzalez E, Knutsen H, Jorde PE (2016) Habitat discontinuities separate genetically
divergent populations of a rocky shore marine fish. Plos One 11, e0163052.

Carpenter KE, Smith-Vaniz WF, de Bruyne G, de Morais L (2015) Gobius niger. The IUCN Red
List of Threatened Species 2015: e.T198570A21913403.

https://dx.doi.org/10.2305/IUCN.UK.2015-4.RLTS.T198570A21913403.en. Downloaded on 16
August 2021.

Cattell R B (1966) The Scree test for the number of factors. Multivariate Behavioral Research,
1(2), 245–276.

Catarino D, Knutsen H, Verissimo A, Olsen EM, Jorde PE, Menezes G, et al (2015) The Pillars of
Hercules as a bathymetric barrier to gene flow promoting isolation in a global deep-sea shark
(*Centroscymnus coelolepis*). Molecular Ecology, 24, 6061 - 6079.

Catchen J, Hohenlohe PA, Bassham S, Amores A, Cresko WA (2013) Stacks: an analysis tool set
for population genomics. Molecular Ecology, 22,3124–3140.

Ciannelli L, Knutsen H, Moland E, Espeland SH, Asplin L,Jelmert A, Knutsen JA, et al (2010)
Small-scale genetic structure in a marine population in relation to water circulation and egg
characteristics. Ecology, 91: 2918—2930.

Cowen RK, Paris CB, Srinivasan A (2006) Scaling of connectivity in marine populations. Science,
311, 522– 527.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G,
Marth GT, Sherry ST (2011) The Variant Call Format and VCFtools. Bioinformatics, 2715, 2156–
2158.

Dalsøren S, Albretsen J, Asplin L (2020) New validation method for hydrodynamic fjord models
applied in the Hardangerfjord, Norway. Estuarine Coastal and Shelf Science, 246, 107028.

Darwall WRT, Costello MJ, Donnelly R, Lysaght S (1992) Implication of life-history strategies for
a new wrasse fishery. Journal of Fish Biology, 41: 111–123.

Evankow A, Christie H, Hancke K, Brysting AK, Junge C, Fredriksen S, Thaulow J (2019) Genetic
heterogeneity of two bioeconomically important kelp species along the Norwegian coast.
Conservation Genetics, 20, 615–628.

Foll M, Gaggiotti O (2008) A genome-scan method to identify selected loci appropriate for both
dominant and codominant markers: A Bayesian perspective. Genetics, 180, 977–993.

Froese and Pauly, (2021) Editors. 2021. FishBase. World Wide Web electronic publication.
www.fishbase.org, version (06/2021).

Haidvogel DB, Arango H, Budgell WP, Cornuelle BD, Curchitser E, Di Lorenzo E, et al. (2008)
Ocean forecasting in terrain-following coordinates: formulation and skill assessment of the
Regional Ocean modelling system. Journal of Computational Physics, 227, 3595–3624.

Halvorsen KT, Larsen T, Browman HI, Durif C, Aasen N, Vøllestad LA, et al (2021) Movement
patterns of temperate wrasses (Labridae) within a small Marine Protected area. Journal of Fish
Biology, 99, 1513-1518.
569
570 Hansen A, Westgaard JI, Søvik G, Hanebrekke T, Nilssen EM, Jorde PE, Albretsen J, Johansen T
571 (2021) Genetic differentiation between inshore and offshore populations of northern shrimp
572 (*Pandalus borealis*). ICES Journal of Marine Science, 78(9), 3135-3146.
573

Hauser L, Carvalho GR (2008) Paradigm shifts in marine fisheries genetics: ugly hypotheses
slain by beautiful facts. Fish Fisheries, 9, 333-362.

Hop H, Gjøsæter J, Danielssen DS (1992) Seasonal feeding ecology of cod (*Gadus morhua* L.) on
the Norwegian Skagerrak coast. ICES Journal of Marine Sciences, 49, 453–461.

Johansson ML, Banks MA, Glunt KD, Hassel-Finnegan HM, Buonaccorsi VP (2008) Influence of
habitat discontinuity, geographical distance, and oceanography on fine-scale population
genetic structure of copper rockfish (Sebastes caurinus). Molecular Ecology,17, 3051–3061.

Jombart T, Ahmed I (2011) adegenet 1.3-1: New tools for the analysis of genome-wide SNP
data. Bioinformatics, 27(21), 3070–3071.

Jombart T, Devillard S, Balloux F (2010) Discriminant analysis of principal components: A new
method for the analysis of genetically structured populations. BMC Genetics, 11(1), 94.

Kanno Y, Vokoun JC, Letcher BH (2011) Fine-scale population structure and riverscape genetics
of brook trout (*Salvelinus fontinalis*) distributed continuously along headwater channel
networks. Molecular Ecology, 20(18), 3711-3729.

Kara MH, Quignard JP (2019) Fishes in lagoons and estuaries in the Mediterranean 2.
Sedentary fish. ISTE-Wiley Ed. 418 pp.

Keenan K, McGinnity P, Cross TF, Crozier WW, Prodöhl PA (2013) DiveRsity: an R package for
the estimation and exploration of population genetics parameters and their associated errors.
Methods in Ecology and Evolution, 4, 782–788.

Kelley JL, Brown AP, Therkildsen NO, Foote AD (2016) The life aquatic: advances in marine
vertebrate genomics. Nature Reviews Genetics, 17, 523–534.

Kimura M, Weiss GH (1964) The stepping stone model of population structure and the
decrease of genetic correlation with distance. Genetics, 49, 561-576.

Knutsen JA, Knutsen H, Gjøsæter J, Jonsson B (2001) Food of anadromous brown trout at sea.
Journal of Fish Biology, 59, 533– 543.

Knutsen H, Olsen EM, Jorde PE, Espeland SH, André C, Stenseth NC (2011) Are low but
statistically significant levels of genetic differentiation in marine fishes 'biologically
meaningful'? A case study of coastal Atlantic cod. Molecular Ecology, 20, 768–783.

Knutsen H, Jorde PE, Blanco Gonzalez E, Robalo J, Albretsen J, Almada V (2013) Climate Change
and Genetic Structure of Leading Edge and Rear End Populations in a Northwards Shifting
Marine Fish Species, the Corkwing Wrasse (*Symphodus melops*). Plos One, 8, e67492.

Knutsen H, Jorde PE, Blanco Gonzalez E, Eigaard OR, Pereyra RT, Sannæs H, Dahl M, André C,
Søvik G (2014) Does population genetic structure support present management regulations of
the northern shrimp (*Pandalus borealis*) in Skagerrak and the North Sea? ICES Journal of
Marine Sciences, 72, 863–871.

Knutsen H, Catarino D, Rogers L, Sodeland M, Mattingsdal M, Jahnke M, Hutchings JA,
Mellerud I, Espeland SH, Johanneson K, Roth O, Hansen MM, Jentoft S, André C, Jorde PE
(2022) Combining population genomics with demographic analyses highlights habitat

patchiness and larval dispersal as determinants of connectivity in coastal fish species.
Molecular Ecology (in press)

Kruschel C, Schultz ST (2011) Juvenile *Gobius niger* avoids seagrass in the presence and
uncertain absence of seagrass-inhabiting predators. Journal of Experimental Marine Biology
and Ecology, 409, 240–246.

Lindegren M, van Deurs M, Maureaud A, Thorson JT, Bekkevold D (2022) A spatial statistical
approach for identifying population structuring of marine fish species: European sprat as a
case study. ICES Journal of Marine Science, 79(2), 423-434.

Longmore C, Trueman CN, Neat F et al. (2014) Oceanic scale connectivity and life cycle
reconstruction in a deep-sea fish. Canadian Journal of Fisheries and Aquatic Sciences, 71,
1312–1323.

Lowe WH, Allendorf F (2010) What can genetics tell us about population connectivity?
Molecular Ecology, 19, 3038-3051.

Luu K, Bazin E, Blum MGB (2017) pcadapt: An R package to perform genome scans for selection
based on principal component analysis. Molecular Ecology Resources, 17(1), 67–77.

Martinsohn JT, Raymond P, Knott T, Glover KA, Nielsen EE, Eriksen LB, Ogden R et al. 2019.
DNA-analysis to monitor fisheries and aquaculture: too costly? Fish and Fisheries, 20, 391–401.

Mattingsdal M, Jorde PE, Knutsen H, Jentoft S, Stenseth NC, Sodeland M, Robalo JI, Hansen
MM, André C, Blanco Gonzalez E (2020) Demographic history has shaped the strongly
differentiated corkwing wrasse populations in Northern Europe. Molecular Ecology, 29, 160-
171.

Mertens LE, Treml EA, Von der Heyden S (2018) Genetic and biophysical models help define
marine conservation focus areas. Frontiers in Marine Science, 6(5), 268.

Miller PJ (1986) Gobiidae, Fishes of the North-eastern Atlantic and Mediterranean, vol. 3, Paris,
UNESCO.

Moen FE, Svensen E (2000) Dyreliv i havet – Håndbok i norsk marin fauna. 2nd edition. Kom
Forlag. Kristiansund, Norway. 576pp.

Myksvoll MS, Sandvik AD, Asplin L, Sundby S (2014) Effects of river regulations on fjord
dynamics and retention of coastal cod eggs. ICES Journal of Marine Science, 71, 943–956.

Narum SR, Hess JE (2011) Comparison of FST outlier tests for SNP loci under selection.
Molecular Ecology Resources, 11, 184–194.

Östman Ö, Olsson J, Dannewitz J, Palm S, Florin AB (2017) Inferring spatial structure from
population genetics and spatial synchrony in demography of Baltic Sea fishes: Implications for
management. Fish and Fisheries, 18(2), 324–339.

Palumbi SR (2004) Marine reserves and ocean neighborhoods: the spatial scale of marine
populations and their management. Annual Review of Environment and Resources, 29, 31-68.

Pante E, Simon-Bouhet B (2013) marmap: a package for importing, plotting and analyzing
bathymetric and topographic data in R. PLoS ONE, 8, e73051.

Penthon P (2005) Aschehougs store Fiskebok. 2nd edition. H. Aschehoug & Co, Kristiansand,
Norway, 447pp.

Perry D, Staveley TAB, Gullström M (2018) Habitat Connectivity of Fish in Temperate Shallow-
Water Seascapes. Frontiers in Marine Science, 4, 440.

Peterson BK, Weber JN, Kay EH, Fisher HS Hoekstra HE (2012) Double Digest RADseq: An
Inexpensive Method for De Novo SNP Discovery and Genotyping in Model and Non Model
Species. Plos One, 7, doi: 10.1371/journal.pone.0037135

Planes S (1998) Genetic diversity and dispersal capabilities in marine fish, in M.K. HECHT et al.
(eds), Evolutionary Biology, Plenum Press, New York.

Privé F, Luu K, Vilhjálmsson BJ, Blum MG (2020) Performing highly efficient genome scans for
local adaptation with R package pcadapt version 4. Molecular Biology and Evolution, 37, 2153–
2154.

Purcell S, Neale B, Todd-Brown K et al. (2007) PLINK: a tool set for whole-genome association
and population-based linkage analyses. American Journal of Human Genetics, 81, 559–575.

Quintela M, Kvamme C, Bekkevold D, Nash RDM, Jansson E, Sørvik AG, Taggart JB et al. (2020)
Genetic analysis redraws the management boundaries for the european sprat. Evolutionary
Applications, 13, 1906–1922.

R Core Team (2021) R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Rogers LA, Olsen EM, Knutsen H, Stenseth NC (2014) Habitat effects on population connectivity
in a coastal seascape. Marine Ecology Progress Series, 511, 153-163.

Rousset F (1997) Genetic differentiation and estimation of gene flow from F-statistics under
isolation by distance. Genetics, 145, 1219-1228.

Rousset F (2000) Genetic differentiation between individuals. Journal of Evolutionary Biology,
13, 58-62.

Rousset F (2008) genepop'007: a complete re-implementation of the genepop software for
Windows and Linux. Molecular Ecology Resources, 8, 103–106.

Schwarz G (1978) Estimating the dimension of a model. Annals of Statistics, 6, 461–464.

Seljestad GW, Quintela M, Faust E, Halvorsen KT, Besnier F, Jansson E, Dahle G, Knutsen H,
André C, Folkvord A, Glover KA (2020) "A cleaner break": Genetic divergence between
geographic groups and sympatric phenotypes revealed in ballan wrasse (*Labrus bergylta*).
Ecology and Evolution, 10, 6120-6135.

Selkoe KA, D'Aloia CC, Crandall ED, Iacchei M, Liggins L, Puritz JB, et al (2016) A decade of
seascape genetics: Contributions to basic and applied marine connectivity. Marine Ecology
Progress Series, 554, 1– 19.

Shchepetkin AF, McWilliams JC (2005) The regional oceanic modeling system (ROMS): a split-
explicit, free-surface, topography-following-coordinate oceanic model. Ocean Modelling, 9,
347–404.

Slatkin M (1994) Gene Flow and Population Structure. In: Real, L., Ed., Ecological Genetics,
Princeton University Press, Princeton, 3-17.

Stenseth NC, Bjørnstad ON, Falck W, Fromentin JM, Gjøsæter J, Gray J (1999) Dynamics of
coastal cod populations: intra- and intercohort density dependence and stochastic processes.
Proceedings of the Biological Society, 266, 1645–1654.

Sotka EE, Wares JP, Barth JA, Grosberg RK, Palumbi SR (2004) Strong genetic clines and
geographical variation in gene flow in the rocky intertidal barnacle *Balanus glandula*.
Molecular Ecology,13(8), 2143-56.

Vaas KF, Vlasblom AG, De Koeijer P (1975) Studies on the Black Goby (*Gobius niger*, Gobiidae,
Pisces) in the Veerse Meer, S.W. Netherlands. Nertherlands Journal of Sea Research, 9 (1), 56–
68.

Vavrek MJ (2011) Fossil: palaeoecological and palaeogeographical analysis tools.
Palaeontologia Electronica, 14, 1T

Villegas-Ríos D, Alós J, March D, Palmer M, Mucientes G, Saborido-Rey F (2013) Home range
and diel behavior of the ballan wrasse, *Labrus bergylta*, determined by acoustic telemetry.
Journal of Sea Research, 80, 61–71.

Waples R (1998) Separating the wheat from the chaff: patterns of genetic differentiation in
high gene flow species, Journal of Heredity, 89, 438-450.

Weir BS, Cockerham CC (1984) Estimating F-statistics for the analysis of population structure.
Evolution, 38, 1358–1370.

Acknowledgments

This study was funded by the ECOGENOME project (grant nr. grant number: 280453), through the Norwegian Research Council. Further funding was provided by MarGen II, an Intereg project under the Øresund-Kattegat-Skagerrak program, by the Ministry of Trade, Industry and Fisheries, and from University of Agder, Centre for Coastal Research. We thank the scientific team participating in the Beach seine survey. We also thank Ricardo Medeiros (IMAR, University of the Azores) for making the map and to Karen Martinez-Swatson (IMR, Norway) for helping with the laboratory methodology description. DC was funded by the University of Agder and at later stage by FCT – Foundation for Science and Technology, I.P., under the project UIDP/05634/2020 granted to Okeanos. Okeanos is funded through the FCT, I.P., under the project UIDB/05634/2020 and UIDP/05634/2020 and by the Regional Government of the Azores through the initiative to support the Research Centers of the University of the Azores and through the project M1.1.A/REEQ.CIENTÍFICO UI&D/2021/010.

Table 1. Sampling locations and summary statistics for black goby samples along the

Norwegian coast (WC, West Coast; SK, Skagerrak). Coordinates are given in decimal degrees

(the shown coordinates are an average value when more than one station was sampled). N1

refers to the total number of individuals initially genotyped, combined over sampling stations;

N2 is the number of black gobies remaining after filtering and used for downstream analyses

(292 in total); Stations referrer to the total number of stations used within a sampling location;

expected (He) and observed (Ho) heterozygosity, and inbreeding coefficient (F_{is}) averaged over

611 SNPs.

Area	Sampling locations	Site code	Latitude (N)	Longitude (E)	N1	N2	Stations	H_E	H_O	F_{IS}
WC	Ålesund	ALE	62,338	6,581	23	19	1	0.155	0.157	0.017
	Manger	MAN	60,664	5,095	21	18	2	0.156	0.156	0.020
	Stavanger	STA	59,114	5,727	24	20	1	0.152	0.149	0.047
	Egersund	EGE	58,421	6,014	23	18	1	0.148	0.144	0.052
SK	Torvefjorden	TOR	58,064	7,764	16	14	4	0.152	0.152	0.009
	Topdalsfjorden	TOP	58,144	8,050	24	21	2	0.155	0.155	0.024
	Lillesand	LIL	58,206	8,308	24	21	5	0.146	0.148	0.004
	Grimstad	GRM	58,283	8,509	17	9	2	0.145	0.150	-0.036
	Tvedestrand	TVE	58,623	9,058	24	17	3	0.145	0.142	0.042
	Risør	RIS	58,737	9,314	24	11	4	0.141	0.150	-0.038
	Kragerø	KRA	58,835	9,319	24	18	4	0.143	0.139	0.046
	Langesund	LAN	59,030	9,808	24	17	3	0.147	0.145	0.038
	Sandefjord	SAN	59,088	10,227	13	12	2	0.139	0.137	0.023
	Vrengen	VRE	59,165	10,377	25	20	3	0.153	0.146	0.056
	Håholmen	HAH	59,139	10,434	24	20	1	0.144	0.138	0.054
	Oslofjord	OSF	59,766	10,712	24	18	1	0.147	0.143	0.036
	Hvaler	HVA	59,264	10,731	25	19	3	0.147	0.148	-0.010

	ALE	MAN	STA	EGE	TOR	TOP	LIL	GRM	TVE	RIS	KRA	LAN	SAN	VRE	HAH	OSF
MAN	0.0041															
STA	0.0131	0.0116														
EGE	**0.04229***	**0.04596***	0.0234													
TOR	**0.04447***	**0.04364***	0.0267	-0.0019												
TOP	**0.03818***	**0.04046***	**0.03305***	0.0036	-0.0003											
LIL	**0.0430***	**0.05144***	**0.04781***	0.0085	0.0122	0.0051										
GRM	**0.06464***	**0.06473***	0.0361	-0.0014	0.0000	0.0115	0.0235									
TVE	**0.04023***	**0.04606***	**0.03695***	0.0011	0.0082	0.0102	0.0041	0.0079								
RIS	**0.03769**	**0.04602***	0.0377	0.0035	0.0045	0.0010	-0.0008	0.0107	-0.0040							
KRA	**0.04836***	**0.05177***	**0.03289***	0.0045	0.0049	0.0072	0.0018	0.0058	-0.0017	-0.0064						
LAN	**0.04338***	**0.04557***	0.0260	0.0032	0.0025	-0.0010	0.0091	0.0009	0.0045	0.0025	-0.0013					
SAN	**0.05817***	**0.0657***	**0.04529***	0.0027	0.0007	-0.0036	0.0113	0.0065	0.0137	0.0122	0.0080	0.0028				
VRE	**0.04393***	**0.0502***	**0.03388***	-0.0031	0.0021	-0.0025	0.0110	-0.0029	0.0084	0.0028	0.0021	0.0006	-0.0006			
HAH	**0.05965***	**0.06464***	**0.03829***	0.0018	0.0000	0.0085	0.0193	-0.0065	0.0101	0.0051	0.0069	0.0000	0.0031	0.0002		
OSF	**0.04801***	**0.05363***	**0.04173***	-0.0006	0.0035	-0.0034	0.0041	0.0081	0.0054	0.0011	0.0051	0.0005	-0.0025	-0.0015	0.0060	
HVA	**0.04554***	**0.05165***	**0.03563***	0.0038	0.0071	-0.0012	0.0089	0.0099	0.0041	0.0044	0.0031	0.0041	-0.0012	-0.0021	0.0063	0.0006

Figure 1. Map showing the main sampling locations where black goby samples were collected

for the genetic analyses. *Jæren* denotes the area of a long stretch of sandy beaches. Dashed

lines delimitate the Skagerrak region between Norway, Denmark and Sweden. Sampling

location abbreviations are given in Table 1.

Figure 2. Results from the discriminant analysis of principal components (DAPC) showing (a)

two clusters, one corresponding to West coast locations and the other to Skagerrak+ Egersund

samples; and showing (b) no distinct clustering within Skagerrak +Egersund only. The DAPC

was conducted based on 100 PCs that conserved more than 80% variation.

Figure 3. Isolation by distance (IBD) for the black goby. Effect of geographic distance (km over

water) on linearized observed genetic divergence, $F_{ST}/(1-F_{ST})$, (a) among all black goby sampling

localities (slope = 7.2*10^{-5}, P<0.001) and for Skagerrak localities only (slope= 3.8*10^{-6}, P>0.05);

(b) among all pairwise comparisons with Hvaler in bold (eastern most sample) and by fitting a

sigmoid curve to the data points including Hvaler highlighting the sharp increase in F_{ST} from the

east to the west coast of Norway.

Figure 4. Isolation by distance (IBD) among individuals for the black goby sheltered (orange)

and exposed (blue) sampling stations in Skagerrak. Sampling stations were binned into

distance classes of 1 km and size of the dots were scaled to number of individual pairs within

each distance class. Effect of linear distance on the genetic difference (ê) calculated at

individuals' level, showing a significant increase with distance for sheltered stations (P =

0.0018, slope = $3.446*10^{-5}$) and non-significate trend for exposed stations (P = 0.3555, slope =

$2.298*10^{-6}$).

Figure 5. Spatial decorrelation in abundance fluctuations of black goby at beach seine stations

along the Skagerrak coast. Results are shown for (a) all stations and (b) models fit to sheltered

(orange) and exposed (blue) stations separately. Points show pairwise correlations among

stations, and solid lines show a fitted exponential decay model along with bootstrapped 90%

confidence intervals (shaded areas). The estimated decorrelation scale (v) is shown as a

vertical dashed line.

Figure 6. Predicted effects of biogenic habitat type and percent coverage on catches of black

goby in the Skagerrak beach seine survey (with 95% prediction intervals). Predictions are

shown for intermediate coverage (10-39%) in the first panel, and for eelgrass habitat in the

second panel.

Figure 7. Maps from the oceanographic drift model showing the density of particles (colours)

after 25 days drift from the locations West (left panel) and East of Egersund (right panel). The

locations where particles were released are denoted with black circles. The colour scale

denotes probability in % relative to the total number of particles released from all locations

East and West of Egersund. The broad, zones of particle densities offshore are linked to the

open boundary of the high-resolution fjord models that inhibits further drift offshore.