**Osteocalcin binds to a GPRC6A Venus fly trap allosteric site to positively modulate**
**GPRC6A signaling**

Rupesh Agarwal*[1,2], Min Pi*[3], Ruisong Ye[3], Micholas Dean Smith[1,2], Jeremy C. Smith[1, 2], L. Dar-
ryl Quarles[3]

*Authors have contributed equally

[1]University of Tennessee/Oak Ridge National Laboratory Center for Molecular Biophysics, Oak Ridge,
Tennessee 37830; and [2]Department of Biochemistry and Cellular and Molecular Biology, University of
Tennessee, Knoxville, Tennessee 37996. [3]Departments of Medicine, University of Tennessee Health Sci-
ence Center, Memphis, Tennessee 38163.

*Running title (60 characters): Osteocalcin binds to the GPRC6A as a positive allosteric modula-*
*tor*

*Keywords: G protein-coupled receptor (GPCR), homology modeling, mutagenesis, Osteocalcin,*
*isoform-selectivity, positively allosteric modulator, extracellular domain*

## Abstract

GPRC6A is a member of the Family C G-protein coupled receptors that is activated by cations,
L-amino acids, the osteocalcin (Ocn) peptide, and testosterone. GPRC6A functions as a master
regulator of energy metabolism and sex hormone production. Based on homology to the related
receptors mGluR5 and CaSR, GPRC6A's multiple ligand specificity is likely based on an
orthosteric ligand binding site in the bilobed Venus fly trap (VFT) domain together with two posi-
tive allosteric modulator (PAM) sites, one in the VFT and the other in the 7TM domain. Here, we
show that Ocn acts as a PAM for GPRC6A by binding to a site in the VFT that is distinct from
the orthosteric site for calcium and L-amino acids. In agreement with this finding, alternatively
spliced GPRC6A isoforms 2 and 3, which lack regions of the VFT, and mutations in the predict-
ed Ocn binding site, K352E and H355P, prevent Ocn activation of GPRC6A. These observa-
tions provide a structural framework for understanding the ability of multiple distinct classes of
compounds to activate GPRC6A and set the stage to develop novel small molecules to activate
and inhibit this receptor.

## Introduction

Osteocalcin (Ocn) is a 49 residue protein in humans that is mainly produced by osteoblasts in bone. A vitamin K-dependent carboxylated form of Ocn comprises much of the non-collagenous bone protein matrix, while an undercarboxylated form (unOcn) is released into the circulation to function as a hormone. unOcn regulates energy metabolism through direct effects on metabolic functions in multiple target tissues and by indirect effects mediated by an ensemble of metaboli­cally active hormones whose release is stimulated by unOcn (1-5). Indeed, unOcn is reported to stimulate insulin secretion and β-cell proliferation in the pancreas (5), fibroblast growth factor 21 (FGF-21) release and glucose and fat metabolism in liver hepatocytes (6), interleukin 6 (IL-6) secretion and glucose utilization in skeletal muscle myocytes (7), adiponectin release and lipolytic activity in adipocytes in white fat (8), testosterone production from testicular Leydig (2,6), and glucagon-like peptide 1 (GLP-1)  secretion from gastrointestinal enterocytes (9,10).

The peripheral metabolic effects of Ocn are mediated through activation of GPRC6A, a member of the Family C G-protein coupled receptors (5,11).   GPRC6A is activated by several orthosteric ligands, including basic L-amino acids (such as L-Lys, L-Arg, and L-ornithine) and divalent cations (such as $Ca^{2+}$ and $Mg^{2+}$).  Also, positive allosteric modulators (PAMs) of GPRC6A include testosterone, NPS-568, and di-phenyl and tri-phenyl compounds (12,13). Gallate and epigallocatechin 3-gallate (ECGC), which are natural products in green tea, respec­tively activate and inhibit GPRC6A (14). The structural basis for GPRC6A interaction with multi­ple ligands is incompletely understood.

Family C GPCRs are characterized by their similarity in 7 transmembrane (7TM) domain se­quences, a necessity for dimerization to function, and in most cases, the existence of a large N-terminal ligand-binding Venus flytrap (VFT) domain (15). Recently, high resolution structural da­ta of the related mGluR5 (16) and CaSR (17-20), have identified how various ligands and drug molecules target the VFT and 7TM.  For example, mGluR5 has the orthosteric binding site for L-glutamate in the VFT, and two positive modulating allosteric (PAM) sites, one located in the VFT and the other in the 7TM (16).  The allosteric site located in the VFT was discovered by a nanobody, Nb43, that binds to Helix L and the L-M loop of mGluR5 to potentiate orthosteric ag­onist binding and receptor activation. Similarly, elucidation of the 3D structure of CaSR identified in the VFT an orthosteric $Ca^{2+}$ ion binding site that acts as a composite agonist with L-amino acids, such as L-tryptophan, to stabilize the closure of the active VFT (17). CaSR also has an

allosteric site at the dimeric interface of the VFT (proximal to residue D248) to which the peptide
etecalcitide binds and acts as a positive allosteric modulator (PAM). In CaSR, additional small
molecule drugs act as PAMs (e.g., calcimimetics), or negative allosteric modulators (NAMs)
(e.g., calcilytic) via a site in the 7TM (17-20).

Based on this new structural information, we developed a structural model for the activation of
GPRC6A by orthosteric and allosteric ligands. We investigated Ocn activation of GPRC6A
isoforms with deleted segments of the extracellular domain, mapped the peptide fragments of
Ocn necessary for activation of GPRC6A, and examined if Ocn peptides bind to the VFT and
modulate GPRC6A function.

**Results**

***Structural modeling of GPRC6A isoforms and L-Arginine orthosteric ligand binding.***

GPRC6A exists as three isoforms: isoform 1 represents the full-length receptor, and two alterna-
tively spliced receptor isoforms 2 and 3 are characterized by respective deletions of exon 3 and
exon 4 segments that encode regions of the VFT domain (21) (**Figure 1A**).  To identify regions
that differ between the isoforms, we aligned the protein sequences of isoforms 1, 2 and 3 of
GPRC6A (**Figure 1B, S1**). The TM domains of these three isoforms have 100% sequence iden-
tity, but isoforms 2 and 3 are missing residues 271-445 and 446-516 in the extracellular domain,
respectively.

To derive the structure of GPRC6A, we used AlphaFold2 (22-25) to model the full-length iso-
form 1 **(Figure 1C)**. AlphaFold2 is an AI-based technique that derives close-to-experimental ac-
curacy in protein structure prediction. These calculations confirmed that GPRC6A has 3 do-
mains: TM, CRD and VFT, similar to the related mGluR5 and CaSR Cryo-EM structures. The
residues deleted in the isoforms 2 and 3 correspond to different segments of the VFT domain. In
the case of isoform 2, a large segment of the VFT domain structure (~ 174 residues) is deleted,
whereas a relatively small region (~ 70 residues) is missing in isoform 3 (**Figure S1**).

In isoform 1, we identified an orthosteric ligand site near residues Tyr 148, Ser 149, Thr 172,
Asp 303 in the VFT based on structure superimposition with mGluR5 and CaSR (**Figure S2a**
**and S2b**). In contrast, in isoform 2, the orthosteric ligand binding site predicted based on align-
ing orthosteric bound structures of mGluR5 and CaSR structures is missing, foretelling a loss-
of-function (*vide infra*) (**Figure S2c**).  The predicted orthosteric ligand binding site is retained in
isoform 3, and hence this isoform is predicted to be responsive to orthosteric ligands (*vide infra*).

To test these predictions, GPRC6A isoforms 1, 2, or 3 were transfected into HEK-293 cells and

receptor activation was measured by cAMP accumulation in response to the orthosteric ligand

L-Arg in the presence of 1 mM calcium.  Isoform 2 resulted in loss of L-Arg stimulation of cAMP,

whereas isoform 3 retained responsiveness to L-Arg activation, similar to isoform 1 (**Figure 2A**).

***GPRC6A isoforms 2 and 3 have attenuated responses to Ocn.***

The location of the Ocn binding site in GPRC6A is still not well understood. To explore the role

of the VFT in mediating the response to Ocn, we compared the ability of Ocn to activate the

three GPRC6A isoforms (**Figures 2B and C**). In the presence of 1 mM calcium Ocn activated

GPRC6A isoform 1 in a dose dependent manner, consistent with previous reports (5). GPRC6A

isoform 2 exhibited an attenuated ERK response with a complete loss of cAMP activation in re-

sponse to Ocn treatment. In contrast, GPRC6A isoform 3 also showed attenuation of Ocn-

induced ERK stimulation but the cAMP response to Ocn persisted (**Figures 2B and C**). Taken

together, loss-of Ocn responses in isoforms 2 and 3 suggest that the VFT is necessary for Ocn

activation of GPRC6A and that the different isoforms bias ligand-induced coupling to G-protein

signaling.

***Identifying Ocn amino acids necessary for GPRC6A activation***

Ocn in humans is a 49 amino acid peptide.  Initially, we used functional assays to refine the pro-

tein-protein interface between Ocn and GPRC6A VFT. We designed Ocn fragments (**Figure**

**3A**) (11,26) and compared their ability to activate the full length GPRC6A isoform 1 (**Figures 3B**

**and C**) transfected into HEK293 cells. Ocn peptide activation was measured by assessing both

ERK (**Figure 3B**) and cAMP (**Figure 3C**) signaling. Compared to the full length Ocn (1-49), all

peptide fragments activated signaling in cell transfected with GPRC6A to some degree.  How-

ever, we found that the decapeptide fragment of Ocn 20-29 (revcelnpdc) located in the middle of

the protein and the C-terminal hexapeptide Ocn 44-49 (rfygpv) exhibited significantly higher ac-

tivity (**Figure 3B and C**).

To further investigate the bioactivity of Ocn 20-29 fragment *in vivo,* we injected the synthetic

Ocn 20-29 peptide (2 μM/kg) into wild type *Gprc6a*$^{+/+}$ and *Gprc6a*$^{-/-}$ mice by the intraperitoneal

(IP) route (27), and collected blood and serum at 24 hours. We found that Ocn 20-29 at a dose

of 2 μM/kg reduced blood glucose levels in wild-type mice by ~10% at 24 hours after

intraperitoneal administration of 2 μM/kg, whereas the vehicle (saline) had no effect on blood

glucose (Fig. 4A). Consistent with Ocn effects in stimulating IL-6 release from skeletal muscle

(7),  Ocn 20-29 at a dose of 2 μM/kg IP also resulted in a significant increase in serum IL-6 in

wild type *Gprc6a*[+/+] mice, but not in *Gprc6a* knockout (*Gprc6a*[-/-]) mice (Fig. 4C). Ocn 20-29

injection did not significantly stimulate serum insulin (Fig. 4B) or adiponectin (Fig. 4D)

levels in wild-type mice at 24 hours at a dose of 2 $\mu$M/kg.

***Protein-protein docking to generate GPRC6A-VFT: Ocn complex structure***

As noted above, there are two PAM sites in the related receptors mGluR5 and CaSR, one in the

7TM domain targeted by small molecules and another in the VFT targeted by the protein

nanobody (16) and the etecalcitide peptide (20). Specifically, the PAM function of Nb43 is medi-

ated by a nanobody binding site near a helix in the VFT domain of the mGluR5 as defined by

Cryo-EM (16). Aligning the structures of the allosteric ligand bound VFTs mGluR5 (PDB 6N4Y)

and CaSR (PDB 7E6U, 7M3G) with our model of GPRC6A and its isoforms indicated that the

dimeric interface found to bind etecalcitide and NAM in CaSR was invariant across the GPRC6A

isoforms, while the binding pocket homologous to the mGluR5 nanobody PAM site was absent

in GPRC6A's isoform 2 (**Figure S3a**). We hypothesized that, given the loss of Ocn associated

function in isoform 2, the pocket homologous to the mGluR5 binding site may serve as the bind-

ing site for Ocn in GPRC6A. Using this site as our guide, we used HADDOCK 2.4 (28) to gener-

ate a model of the GPRC6A-VFT: Ocn complex. The top 4 scoring structures within the calcu-

lated top cluster were then subjected to additional analysis and visual inspection.

Our analysis revealed that the GPRC6A-VFT forms multiple hydrogen bonds (a list is provided

in **Table 1)** with Ocn in all the models. Residues 20-29 from the highest active Ocn fragment

also form multiple hydrogen bonds with GPRC6A residues (Gln 19, Asp26, Gln381, Lys352)

(**Figure 5**) and is also in proximity (within 4 Å) to other residues (Ala385, Arg382, Asp26,

Asp349, Gln19, Gln22, Gln61, Glu356, Glu62, His351, His355, Pro20, Ser348, Ser350, Val28),

which can form other potential electrostatic interactions.  Moreover, the Ocn model shows that

there is a disulfide bond between residues Cys 23 and Cys 29 (**Figure S3b**), both of which are

present in the Ocn fragment. This is predicted to provide structural integrity to the fragment and

would potentially fold similarly to the full length Ocn peptide. In analyzing the interactions be-

tween the GPRC6A-VFT and Ocn, we observe that two GPRC6A-VFT residues, Lys 352 and

His 335, in the helix region (352-363) are common in all the models (**Figure 5, Table 1**). These

residues form hydrogen bonds with Ocn residues Asp 28, Asp 30 and Glu 31 and are near sev-

eral other residues present in 20-29 region of Ocn that was the most active fragment.

***Validation of the complex structure using single point mutagenesis***

To test the above models, we mutated Lys352 or His355 in human GPRC6A, into Glu352 (mu-
tant K352E) or Pro355 (mutant H355K), and also created double mutated Glu352/Pro355 (mu-
tant K352E/H355P) GPRC6A. Mutants and wild-type hGPRC6A cDNAs were transiently trans-
fected into HEK-293 cells and signaling was assessed in response to Ocn treatment.

We confirmed that mutants K352E, H355P, K352E/H355P, and WT mGRPC6A proteins were
equally expressed, as assessed by Western blotting using a Myc antibody, which recognized
the Myc epitope located at after signal peptides (1-18 amino acids) of the WT and mutant recep-
tors (**Figure S4 a,b**)

The full-length Ocn 1-49 peptide dose-dependently stimulated cAMP accumulation in the wild-
type receptor, but the response was significantly decreased in the K352E, H355P and
K352E/H355P mutant receptors (Figure 5A).  Cells transfected with these three mutants
hGPRC6A cDNAs also showed no response to Ocn 20-29 stimulation (**Figure 6A**). These re-
sults indicate that the binding sites identified by the computational modeling are accurate and
that the helical region (352-363) is important for Ocn binding.

The Ocn 44-49 C-terminal fragment, which was predicted to bind to the 7TM in our prior homol-
ogous model (11), also had a significantly attenuated signaling response in the single mutants.
Interestingly, Ocn 44-49 was able to partially activate the double mutant as assessed by cAMP
accumulation (**Figure 6B**).

## Discussion

Recent Cryo-EM analysis of Family C mGluR5 (16) and CaSR (17,19,20) receptors describes
the structural basis for their orthosteric ligand activation by calcium and L-amino acids and allo-
steric modulating effects of peptides and small molecules. In the current study, we developed
structural homology models of GPRC6A based on the Cryo-EM and found evidence for both
orthosteric and allosteric binding sites in its VFT, similar to mGluR5 and CaSR.

GPRC6A is activated by L-amino acids, cations, and the Ocn peptide (4). Prior work examined
the possible binding of the hexapeptide C-terminal domain of Ocn and identified a binding site in
the 7TM domain of GPRC6A for this fragment 44-48 (11). However, a comparison of recently
available Cryo-EM structures of mGluR5 and CaSR suggested that a regulatory binding site in
the VFT domains of Family C receptors may exist. These results led us to further examine the
Ocn induced activation of GPRC6A and its isoform-specific modulation. Interestingly, GPRC6A
isoforms only differ in the extracellular domain (i.e. VFT) which suggests that a second binding

site for Ocn may exist in the VFT. Taken together, the new structural models developed for
GPRC6A and complimentary experiments and peptide-protein docking calculations reported
here provide clear evidence of a binding site for full-length and/or residues 20-29 Ocn within the
GPRC6A VFT, located near the helical region (352-363).

Reports of GPRC6A activation by the peptide Ocn are inconsistent, with some studies showing
activation (2,3,5,11) and other studies finding no effect (29). Here, we confirmed Ocn activation
of GPRC6A in *in vitro* culture models.  Moreover, the lowering of glucose and stimulation of IL-6
24 hours after a single injection of Ocn 20-29 in wild-type mice but not *Gprc6a*[-/-] mice validate *in*
*vivo* effects of this peptide mediated through GPRC6A. The failure to observe expected stimula-
tion of insulin and adiponectin may be due to examining one time point, the short observation
period or the limited doses tested. Regardless, the location of the Ocn binding site in the VFT at
a site distinct from the orthosteric binding site suggests Ocn functions as a PAM rather than an
orthosteric ligand.  The fact that Ocn activation of GPRC6A requires the presence of calcium
(3,5,11) *in vitro* is consistent with Ocn function as a PAM for GPRC6A (2,11).

Our findings also suggest altered ligand specificity and downstream signaling of GPRC6A
isoforms created by alternative splicing. The loss of a segment of the VFT in isoform 2 resulted
in the loss of responses to L-Arg as well as Ocn.  In contrast, isoform 3 retains responsiveness
to orthosteric activation but exhibits biased signaling to Ocn, as evidence by loss of ERK but
preservation of cAMP signaling.  Loss of this segment may alter the conformational changes
leading to opening of binding sites in the intracellular TM 6 domain that is important for G-
protein coupling. If so, different tissue distributions or the ratio of these isoforms would be ex-
pected to change orthosteric ligand and PAM sensitivities as well as alter coupling to down-
stream signaling pathways resulting in tissue-specific regulation mechanisms.

Our modeling data shows that most contacts between Ocn and GPRC6A are driven by the
structural part of the Ocn peptide (3 helical structure).  Peptide fragments of Ocn consisting of 1-
7, 8-43, and 44-49 are released from bone (30).  We tested the GPRC6A activation potential of
multiple Ocn fragments to further identify the key regions of Ocn that interface with GPRC6A.
The hexapeptide 20-29 most strongly activated GPRC6A, consistent with its location in the cru-
cial region of the full-length Ocn used in our docking studies to the VFT helix region. Ocn 8-19
(apvpypdplepr) also activated GPRC6A (**Figure 3**). This upstream peptide is similar to the
pentadecapeptide peptide wlgapvpypdplepr that is reported to activate GPRC6A *in vitro* and to
improve fatty liver disease and insulin resistance after oral or intraperitoneal administration to a
mouse model (31).  When examining the activities associated with 44-49, it is important to recall

that previous docking and mutagenesis studies suggested that the C-terminus of Ocn binds to
the extracellular transmembrane domain of GPRC6A (11). This might explain the effects of the
rfygpv, C-terminal peptide of Ocn to activate GPRC6A in setting of mutations to disrupt the Ocn
binding domain in the VFT helix. Regardless, short peptides derived from Ocn might be devel-
oped for therapeutics to activate GPRC6A, similar to the 7 amino acid etacalcitide PAM for
CaSR.

Another interesting observation is the greater potency of the peptide 20-29 compared to Ocn 1-
49.  Hypothetically this may be due to a potential disulfide bond present in the small peptide that
creates a stable complementary conformation within the predicted binding site. Alternatively,
competitive self-interactions between the domain and the disordered N-terminus of Ocn may
inhibit long-lived binding conformations between Ocn and GPRC6A.  Instead, flexibility of the N-
terminal domain that result in N-terminal-VFT contacts may inhibit conformational changes
along the dimer interface of the VFT domains.

Although we did not examine binding of other PAMs to GPRC6A in this study, testosterone and
triphenol compounds that activate GPRC6A are predicted to bind to the 7TM domain of
GPRC6A (12,13), analogous to the  second PAM domain in the 7TM domain of mGluR5 and
CaSR (16,20).  This suggests that members of the Family C GPCRs, at least those with large
VFT domains, have a conserved structure consisting of two allosteric binding sites, one in the
VFT and the other in the 7TM domain, along with an orthosteric ligand binding domain in the
VFT.

Finally, GPR158 is another proposed Ocn sensing receptor. The Cryo-EM structure of GPR158
has recently been defined (32).  Our VFT Ocn binding region between GPRC6A and GPR158 is
not conserved and a structure superimposition of the two structures shows that the helix region
(352-363) identified for GPRC6A is missing in GPR158.

Our findings are limited by the lack of Cryo-EM structural data for GPRC6A. Cryo-EM analysis
of mGluR5 and CaSR show that each functions as homodimers, where activation leads to clo-
sure of the VFT and conformational changes the 7TMs revealing G-protein binding sites on the
cytoplasmic side necessary for signaling(17). For CaSR, which is most closely related to
GPRC6A, receptor activation results in 7TM asymmetry in the homodimer leading to only one
protomer for G-protein coupling being stabilized by PAM binding. Our biased signaling response
to Ocn in isoforms 2 and 3 suggests that there may be a structural basis for the divergent cAMP
and ERK responses to Ocn. To fully understand GPRC6A signaling we may need cryo-electron

microscopy structures of GPRC6A in inactive or active states bound to the orthosteric ligands,

PAMS and NAMs.

## Materials and methods

### System and docking details

GPRC6A and Ocn structures were modeled using AlphaFold2 (22,25,33), which is an AI system

developed by DeepMind to predict a 3D structure of a protein from its amino acid sequence.

Multiple sequence alignment (MSA) was performed using the Jackhammer method. The

'max_recycles' parameter, r which controls the maximum number of times the structure is fed

back into the neural network for refinement, was set to 3. All the other parameters were set to

default. The final models were ranked using the pLDDT score. Glycans were not added as the

details are not well known. For docking, HADDOCK 2.4 (28) was used to dock the GPRC6A-

VFP to Ocn. GPRC6A residue numbers 334,335,336,337 and mature Ocn residue numbers

21,22,23,24,25,26, and 27 were used to guide the docking. The best 4 models from the top

cluster (containing 133 models) with HADDOCK score of -81.2±1.1 and Z-score of -1.5 were

analyzed.

### Simulation details

Molecular dynamics simulation (MD) of the top ranked AlphaFold2 model was performed using

the Amber simulation engine(34) to refine the structures. The MD was performed on the protein

using Amber14 ff14SB (35) force field, with a non-bonded cutoff of 10 Å using the Particle Mesh

Ewald algorithm (36). The protein-glycan bound system was hydrated by water model TIP3P

(37) in an octahedral box of 10 Å around the protein in each direction. Initially, the protein was

held fixed with a force constant of 500 kcal mol$^{-1}$ Å$^{-2}$ while the system was energy minimized

with 500 steps of steepest descent followed by 500 steps with the conjugate gradient method. In

a second minimization step, the restraints on the protein were removed and 1000 steps of

steepest descent minimization was performed, followed by 1500 steps of conjugate gradient.

The system was heated to 300 K while holding the protein fixed with a force constant of 10 kcal

mol$^{-1}$ Å$^{-2}$ for 1000 steps. Then, the restraints were removed, and 1000 MD steps were per-

formed. The SHAKE algorithm(38) was used to constrain all bonds involving hydrogen in the

simulations. MD production runs were performed at 300 K using the NPT ensemble and a 2 fs

time step. The temperature was fixed with the Langevin dynamics thermostat (39) and the pres-

sure was fixed with the Monte Carlo barostat (40). The last snapshot from the 20ns trajectory

was used for all analyses.

**Measurement of Total and Phospho-ERK by ERK Elisa Analysis**

Ocn was purified from bovine tibial bone extracts (41,42). Decarboxylated Ocn was produced by

treating Ocn in vacuo at 110°C (42-44). The purity and decarboxylation state were confirmed by

native gel electrophoresis (41), or by blotting followed by reaction with 4-diazobenzene sulfonic

acid staining for γ-carboxyglutamic acid (42,43). The human Ocn fragments were synthesized

by BioMatik USA (Wilmington, Delaware, USA).

All culture reagents were from Invitrogen (Waltham, MA, USA). Human embryonic kidney HEK-

293 cells were obtained from American Type Culture Collection and cultured in DMEM medium

supplemented with 10% fetal bovine serum and 1% Penicillin/Streptomycin (P/S). Briefly,

HEK-293 cells transfected with/without human GPRC6A isoforms cDNA plasmids were starved

by overnight incubation in serum-free DMEM/F12 containing 0.1% bovine serum albumin (BSA)

and stimulated with various ligands at different doses. ERK activation was assessed 20 min

after treatment by using ERK1/2 (phospho-T203/Y204) ELISA Kit (Invitrogen) corrected for the

amount of total ERK using ERK1/2 (Total) ELISA Kit (Invitrogen) to measure ERK levels.

**Measurement of cAMP accumulation**

HEK-293 transfected with human GPRC6A isoform cDNA plasmids cells ($10^5$ cells/well) (45)

were cultured in triplicate in 24-well plates in DMEM supplemented with 10% fetal bovine serum

and 1% penicillin/streptomycin (100 U/mL of penicillin and 100 μg/mL of streptomycin) for 48

hours followed by 4 hours incubation in DMEM/F12 containing 0.1% BSA and 0.5 mM IBMX to

achieve quiescence. Quiescent cells were treated with vehicle control, various ligands at con-

centration as indicated for 40 minutes at 37°C. Then, the reaction was stopped, and the cells

lysed with 0.5 mL 0.1N HCl. cAMP levels were measured by using Cyclic AMP EIA kit (Cayman

Chemical) following the manufacture's protocol.

**Mouse and serum biochemistry**

8 week old wild-type (*Gprc6a$^{+/+}$*) and global knock out (*Gprc6a$^{-/-}$*) mice were injected

intraperitoneal with Ocn 20-29 (2 μM/kg body weight), or vehicle (saline; 10 μl/g body weight).

Serum was collected at 24 hours after intraperitoneal injection. Blood glucose levels were

measured by using blood glucose strips and the Accu-Check glucometer as described (7,27).

Insulin (mouse) ultrasensitive ELISA kit was obtained from ALPCO Immunoassays (Salem, NH,

USA). Mouse IL-6 ELISA Kit was purchased from Invitrogen (Waltham, MA, USA). Mouse

Adiponectin PicoKine™ Quick ELISA Kit was obtained from mybiosource.com.

The *Gprc6a*$^{-/-}$ mouse model was created by replacing exon 2 of the GPRC6A gene with the

hygromycin resistance gene, as described previously (46). Mice were maintained and used in

accordance with recommendations as described (National Research Council 1985; Guide for

the Care and Use of Laboratory Animals Department of Health and Human Services Publication

NIH 86–23, Institute on Laboratory Animal Resources, Rockville, MD) and following guidelines

established by the University of Tennessee Health Science Center Institutional Animal Care and

Use Committee. The animal study protocol was approved by the institutional review boards at

University of Tennessee Health Science Center Institutional Animal Care and Use Committee.

**Site-directed mutagenesis**

*In-vitro* mutagenesis by PCR-mediated recombination QuikChange II XL Site-Directed Muta-

genesis Kit (Agilent) was performed using the cDNA of human GPRC6A cloned in the vector

pcDNA3 (Invitrogen) as template. The primer sets as following. K352E.sen:

GTGACAGTCACGAACTCTTACATG; K352E.antisen: CATGTAAGAGTTCGAGACTGTCAC;

H355P.sen: CAAACTCTTACCTGAATATGCCATG; H355P.antisen:

CATGGCATATTCAGGTAAGAGTTTG; K352E/H355P.sen:

GTGACAGTCACGAACTCTTACGTGAATATGC; and K352E/H355P.antisen:

GCATATTCAGGTAAGAGTTCGTGACTGTCAC. All mutants were confirmed by sequencing.

**Statistics**

We evaluated differences between groups with the Student's *t* test, and for multiple groups by

two-way ANOVA, followed by *a post-hoc* Tukey's test. Significance was set at p < 0.05. All

values are expressed as means ± SEM. All computations were performed using the Statgraphic

statistical graphics system (STSC Inc., Rockville, MD, USA).

**Abbreviations**

cAMP: cyclic adenosine monophosphate; CRD: cysteine rich domain; ERK: extracellular-signal-

regulated kinase; GPRC6A: G protein-coupled receptor family C group 6 member A; NAM: neg-

ative allosteric modulator; Ocn: osteocalcin; PAM: positive allosteric modulator; VFT: venus fly-

trap; TM: transmembrane.

**Acknowledgements**

This work was supported by grants from the National Institutes of Health, NIAMS grant number
1R01AR071930 and NIDDK grant numbers 1R01DK121132 and 1R01DK120567 (LDQ)
The content is solely the responsibility of the authors and does not necessarily represent the
official views of the National Institutes of Health.

**Author Contributions**

355 Rupesh Agarwal, Micholas Dean Smith and Jeremy C. Smith were responsible for the computa-
356 tional studies and Min Pi, Ruisong Ye, and L. Darryl Quarles were responsible for the experi-
357 mental studies regarding GPRC6A.   All authors contributed to the writing of the paper.
358

**Conflict of interest**

360 The authors declare that they have no conflicts of interest with the contents of this article.

**Figures and Tables**

Figure 1. Sequence and structure of GPRC6a a) Placement of exons and its location in GPRC6A isoform 1, 2 and 3 structure; b) Sequence alignment of Venus flytrap (VFT) domain showing missing regions of isoform 2 and 3; Structure model of GPRC6A showing the three domains and the missing region of isoform 2 and 3.

364
365
366
367

Figure 2. Activities of GPRC6A isoforms on response to Ocn. A. Comparison of effects of L-Arg on GPRC6A isoforms mediated cAMP accumulation. Comparison of dose-dependent effects of Ocn on GPRC6A isoforms mediated ERK activation (B) and cAMP accumulation (C). HEK-293 cells were transfected with cDNA plasmids of GPRC6A isoform 1, 2 or 3 for 48 hours, after incubated in Dulbecco's modified Eagle's medium /F-12 containing 0.1% bovine serum albumin quiescence media for 4 hours, then exposed to L-Arg or Ocn at indicated concentrations for 15 minutes for ERK activation, or 40 minutes for cAMP accumulation details as described under "Methods". * indicates a significant difference from control and stimulation groups at p<0.05.

**Figure 3. The activities of human Ocn peptide fragments.** A, The sequence alignment of full
length mouse, bovine and human Ocn (upper panel), and synthesized peptide fragments of hu-
man Ocn (bottom panel). Red alphabets indicate different amino acids compared to human Ocn
sequence. "γ" indicates carboxylation sites in Ocn. B, Comparison of the activities of synthe-
sized fragments of human Ocn (at 20 nM) by ERK phosphorylation in HEK-293 transfected with
vector pcDNA3 (black bar) and HEK-293 cells transfected with pcDNA3-hGPRC6A cDNA (white
bar). * and  ** indicate a significant difference from control and stimulation groups at $p<0.05$ and
$p<0.01$, respectively.  C, Dose response of synthesized fragments of Ocn in HEK-293 cells
transfected with pcDNA3-hGPRC6A cDNA. The ERK phosphorylation was measured 15
minutes for Ocn fragments at concentration as indicated at indicated stimulation in HEK-293
cells transfected with pcDNA3-hGPRC6A cDNA after 4 hours quiescence.

423
**Figure 4. The effects of Ocn fragment, Ocn 20-29 on GPRC6A-mediated regulation of**
**blood glucose and serum IL-6 levels in mouse.** The biological activities of Ocn 20-29 were
assessed 24 hours after intraperitoneal injection of 2 µg/kg the Ocn 20-29 peptide (red bar) or
vehicle (saline, black bar) in 8 week-old $Gprc6a^{+/+}$ or $Gprc6a^{-/-}$ mice by measuring blood glucose
(A), serum insulin (B), IL-6 (C), and adiponectin (D). Ocn 20-29 significantly decreased the
blood glucose level, and increased serum IL-6 level compared to vehicle in wild-type mice. The-
se response were lost in $Gprc6a^{-/-}$ mice. We observed no differences in serum insulin or
adiponectin levels in Ocn 20-29 and vehicle treated mice. * and ** indicate a significant differ-
ence from vehicle and Ocn 20-29 injection groups (n≥5) at $p<0.05$ and $p<0.01$, respectively.
433

435
436

**Figure 5: Superimposition of the top 4 models of GPRC6A's VFT domain (in red) and Ocn (in blue) complex generated from HADDOCK.** The key residue interactions present in the common in helix region (352-363) of GPRC6A's VFT and residues 20-29 of Ocn.

**Figure 6. Mutagenesis of residues in predicted Ocn-binding pocket of GPRC6A.** A, Com-
parison of dose-dependent effects of Ocn on cAMP accumulation in HEK-293 cells transfected
with the cDNA plasmids of WT, K352E, H355P, or K352E/H355P hGPRC6A. * indicate a signifi-
cant difference from control and stimulation groups at $p < 0.05$ and $p < 0.01$, respectively.  B,
Comparison of Ocn small fragments, Ocn 20-29 or Ocn 44-49 at 20 nM concentration on cAMP
accumulation in HEK-293 cells transfected with the cDNA plasmids of WT, K352E, H355P, or
K352E/H355P hGPRC6A. * and  ** indicate a significant difference from control and stimulation
groups at $p < 0.05$ and $p < 0.01$, respectively.

469 **Table 1: List of hydrogen bonds between GPRC6A-VFT and Ocn in best 4 structures from**
470 **the top cluster from HADDOCK.**
471

| GPRC6A residue id | Ocn residue id | Distance (Å) | Model number |
|---|---|---|---|
| Asn389 | Asp34 | 2.79 | 1 |
| Asp26 | Tyr46 | 2.73 | 1 |
| His355 | Asp30 | 2.93 | 1 |
| Lys352 | Asp28 | 3.31 | 1 |
| Ser348 | Glu31 | 2.67 | 1 |
| Asp26 | Asn26 | 2.81 | 2 |
| Asp26 | Phe45 | 3.83 | 2 |
| Asp26 | Tyr46 | 2.73 | 2 |
| Gln381 | Glu24 | 2.8 | 2 |
| Gln381 | Pro27 | 3.06 | 2 |
| His355 | Asp30 | 3.38 | 2 |
| His355 | Glu31 | 2.8 | 2 |
| Lys352 | Asp28 | 3.12 | 2 |
| Ser348 | Glu31 | 2.71 | 2 |
| Ala385 | Asp30 | 3.47 | 3 |
| Asn389 | Asp34 | 2.81 | 3 |
| Asn389 | His35 | 4.7 | 3 |
| Asp26 | Asn26 | 2.81 | 3 |
| Gln19 | Pro18 | 3.89 | 3 |
| Gln19 | Glu21 | 3.03 | 3 |
| Gln381 | Glu24 | 2.81 | 3 |
| His355 | Asp30 | 2.91 | 3 |
| Lys352 | Asp28 | 3.18 | 3 |
| Pro20 | Pro18 | 3.48 | 3 |
| Ser348 | Glu31 | 3.6 | 3 |
| Ala385 | Asp30 | 3.57 | 4 |
| Asp26 | Phe45 | 3.38 | 4 |
| Gln19 | Pro18 | 3.86 | 4 |
| Gln381 | Glu24 | 2.8 | 4 |
| Gln381 | Asp30 | 2.87 | 4 |
| His355 | Glu31 | 2.75 | 4 |
| Lys352 | Asp28 | 3.06 | 4 |
| Pro20 | Pro18 | 3.06 | 4 |
| Ser348 | Glu31 | 2.71 | 4 |

473    **Supporting Information**

474    **Table S1. The amino acid sequence of synthesized human Ocn.**

| Name | Sequence |
|------|----------|
| Ocn 1-7 | YLYGWLG |
| Ocn 8-43 | APVPYPDPLEPRREVCELNPDCDELADHIGFQEAYR |
| Ocn 8-19 | APVPYPDPLEPR |
| Ocn 20-29 | REVCELNPDC |
| Ocn 30-39 | DELADHIGFQ |
| Ocn 44-49 | RFYGPV |

486
487

488    Figure S1: Sequence alignment of GPRC6A isoform 1, 2 and 3.

a)

Orthosteric ligand

**GPRC6A**
**CasR**
**mGlu**

b)

c)

Gln 73        Leu 411

Val 70        Phe 328

Ala 43

Glu 150        Trp 305

Thr 104        Gln 280

Residues missing
in iso 2: 271-445

490
491

Figure S2: a) Superimposed structure of AlphaFold2 minimized model of GPRC6A, CaSR with
L-Trp (PDB 7DTU) and mGluR5 with orthosteric analogue L-quisqualate (PDB 6N51) (in ribbon)
showing orthosteric ligands. b) Interaction map of L-Trp in isoform 1 c) Modeling of the
orthosteric ligand from CaSR in GPRC6A-isoform 1. Black: Trp (placed using CaSR:Trp as
template); Blue (orthosteric residues/ orthosteric binding pocket region missing in iso2); yellow:
iso1 orthosteric pocket region.
498

Figure S3: a) Superimposition of crystal structures on GPRC6A model. Yellow ribbon: GPRC6A;

Dark blue ribbon: region missing in iso2; green ribbon: part missing in iso3; Red ribbon: PAM

Nb43 from mGluR5 (PDB 6N4Y); magenta ribbon: NAM NB2D11 from CaSR (PDB 7E6U); Cy-

an sphere: etelcalcetide (calcimimetic drug) (PDB 7M3G); b) AlphaFold2 minimized model of

hOcn colored from N to C terminus (Blue to red) showing disulfide bond between Cys23 and

Cys29.

510  Figure S4. The protein expression of GPRC6A isoforms (A) and mutants (B**).**

Figure S5. GPRC6A isoform 3 loss responsive to T activation. A. Comparison of effects of $Ca^{2+}$ or T on GPRC6A isoforms mediated cAMP accumulation. B. Comparison of dose-dependent effects of T on GPRC6A isoforms mediated cAMP accumulation. HEK-293 cells were transfected with cDNA plasmids of GPRC6A isoform 1, 2 or 3 for 48 hours, after incubated in Dulbecco's modified Eagle's medium /F-12 containing 0.1% bovine serum albumin quiescence media for 4 hours, then exposed to $Ca^{2+}$ or T at indicated concentrations for 40 minutes for cAMP accumulation details as described under "Methods". * and ** indicate a significant difference from control and stimulation groups at $p<0.05$, and 0.01. C. Alignment of human GPRC6A, mouse GPRC6A and human CaSR showing predicted $Ca^{2+}$ binding sites. Red arrow shows the junction of exon 2 and exon 3. ♦ indicate expected amino acid in GPRC6A depended by CaSR.

## References

1.  Lee, N. K., Sowa, H., Hinoi, E., Ferron, M., Ahn, J. D., Confavreux, C., Dacquin, R., Mee, P. J., McKee, M. D., Jung, D. Y., Zhang, Z., Kim, J. K., Mauvais-Jarvis, F., Ducy, P., and Karsenty, G. (2007) Endocrine regulation of energy metabolism by the skeleton. *Cell* **130**, 456-469
2.  Oury, F., Sumara, G., Sumara, O., Ferron, M., Chang, H., Smith, C. E., Hermo, L., Suarez, S., Roth, B. L., Ducy, P., and Karsenty, G. (2011) Endocrine regulation of male fertility by the skeleton. *Cell* **144**, 796-809

3.    Pi, M., Nishimoto, S. K., and Darryl Quarles, L. (2021) Explaining Divergent Observations
Regarding Osteocalcin/GPRC6A Endocrine Signaling. *Endocrinology* **162**
4.    Pi, M., Nishimoto, S. K., and Quarles, L. D. (2017) GPRC6A: Jack of all metabolism (or
master of none). *Mol Metab* **6**, 185-193
5.    Pi, M., Wu, Y., and Quarles, L. D. (2011) GPRC6A mediates responses to osteocalcin in
beta-cells in vitro and pancreas in vivo. *J Bone Miner Res* **26**, 1680-1683
6.    Pi, M., Xu, F., Ye, R., Nishimoto, S. K., Williams, R. W., Lu, L., and Darryl Quarles, L. (2020)
Role of GPRC6A in Regulating Hepatic Energy Metabolism in Mice. *Sci Rep* **10**, 7216
7.    Mera, P., Laue, K., Wei, J., Berger, J. M., and Karsenty, G. (2016) Osteocalcin is necessary
and sufficient to maintain muscle mass in older mice. *Mol Metab* **5**, 1042-1047
8.    Mukai, S., Mizokami, A., Otani, T., Sano, T., Matsuda, M., Chishaki, S., Gao, J., Kawakubo-
Yasukochi, T., Tang, R., Kanematsu, T., Takeuchi, H., Jimi, E., and Hirata, M. (2021)
Adipocyte-specific GPRC6A ablation promotes diet-induced obesity by inhibiting
lipolysis. *J Biol Chem* **296**, 100274
9.    Mizokami, A., Mukai, S., Gao, J., Kawakubo-Yasukochi, T., Otani, T., Takeuchi, H., Jimi, E.,
and Hirata, M. (2020) GLP-1 signaling is required for improvement of glucose tolerance
by osteocalcin. *J Endocrinol* **244**, 285-296
10.   Mizokami, A., Yasutake, Y., Higashi, S., Kawakubo-Yasukochi, T., Chishaki, S., Takahashi,
I., Takeuchi, H., and Hirata, M. (2014) Oral administration of osteocalcin improves
glucose utilization by stimulating glucagon-like peptide-1 secretion. *Bone* **69**, 68-79
11.   Pi, M., Kapoor, K., Ye, R., Nishimoto, S. K., Smith, J. C., Baudry, J., and Quarles, L. D.
(2016) Evidence for Osteocalcin Binding and Activation of GPRC6A in beta-Cells.
*Endocrinology* **157**, 1866-1880
12.   Pi, M., Kapoor, K., Ye, R., Hwang, D. J., Miller, D. D., Smith, J. C., Baudry, J., and Quarles,
L. D. (2018) Computationally identified novel agonists for GPRC6A. *PLoS One* **13**,
e0195980
13.   Pi, M., Kapoor, K., Wu, Y., Ye, R., Senogles, S. E., Nishimoto, S. K., Hwang, D. J., Miller, D.
D., Narayanan, R., Smith, J. C., Baudry, J., and Quarles, L. D. (2015) Structural and
Functional Evidence for Testosterone Activation of GPRC6A in Peripheral Tissues. *Mol*
*Endocrinol* **29**, 1759-1773
14.   Pi, M., Kapoor, K., Ye, R., Smith, J. C., Baudry, J., and Quarles, L. D. (2018) GPCR6A Is a
Molecular Target for the Natural Products Gallate and EGCG in Green Tea. *Mol Nutr*
*Food Res* **62**, e1700770
15.   Pin, J. P., Galvez, T., and Prezeau, L. (2003) Evolution, structure, and activation
mechanism of family 3/C G-protein-coupled receptors. *Pharmacol Ther* **98**, 325-354
16.   Koehl, A., Hu, H., Feng, D., Sun, B., Zhang, Y., Robertson, M. J., Chu, M., Kobilka, T. S.,
Laeremans, T., Steyaert, J., Tarrasch, J., Dutta, S., Fonseca, R., Weis, W. I., Mathiesen, J.
M., Skiniotis, G., and Kobilka, B. K. (2019) Structural insights into the activation of
metabotropic glutamate receptors. *Nature* **566**, 79-84
17.   Chen, X., Wang, L., Cui, Q., Ding, Z., Han, L., Kou, Y., Zhang, W., Wang, H., Jia, X., Dai, M.,
Shi, Z., Li, Y., Li, X., and Geng, Y. (2021) Structural insights into the activation of human
calcium-sensing receptor. *Elife* **10**
18.   Park, J., Zuo, H., Frangaj, A., Fu, Z., Yen, L. Y., Zhang, Z., Mosyak, L., Slavkovich, V. N., Liu,
J., Ray, K. M., Cao, B., Vallese, F., Geng, Y., Chen, S., Grassucci, R., Dandey, V. P., Tan, Y.

581   Z., Eng, E., Lee, Y., Kloss, B., Liu, Z., Hendrickson, W. A., Potter, C. S., Carragher, B.,
582   Graziano, J., Conigrave, A. D., Frank, J., Clarke, O. B., and Fan, Q. R. (2021) Symmetric
583   activation and modulation of the human calcium-sensing receptor. *Proc Natl Acad Sci U*
584   *S A* **118**

585 19. Wen, T., Wang, Z., Chen, X., Ren, Y., Lu, X., Xing, Y., Lu, J., Chang, S., Zhang, X., Shen, Y.,
586   and Yang, X. (2021) Structural basis for activation and allosteric modulation of full-
587   length calcium-sensing receptor. *Sci Adv* **7**

588 20. Gao, Y., Robertson, M. J., Rahman, S. N., Seven, A. B., Zhang, C., Meyerowitz, J. G.,
589   Panova, O., Hannan, F. M., Thakker, R. V., Brauner-Osborne, H., Mathiesen, J. M., and
590   Skiniotis, G. (2021) Asymmetric activation of the calcium-sensing receptor homodimer.
591   *Nature* **595**, 455-459

592 21. Wellendorph, P., and Brauner-Osborne, H. (2004) Molecular cloning, expression, and
593   sequence analysis of GPRC6A, a novel family C G-protein-coupled receptor. *Gene* **335**,
594   37-46

595 22. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,
596   Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S.
597   A. A., Ballard, A. J., Cowie, A., Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J., Back, T.,
598   Petersen, S., Reiman, D., Clancy, E., Zielinski, M., Steinegger, M., Pacholska, M.,
599   Berghammer, T., Bodenstein, S., Silver, D., Vinyals, O., Senior, A. W., Kavukcuoglu, K.,
600   Kohli, P., and Hassabis, D. (2021) Highly accurate protein structure prediction with
601   AlphaFold. *Nature* **596**, 583-589

602 23. Jumper, J., and Hassabis, D. (2022) Protein structure predictions to atomic accuracy with
603   AlphaFold. *Nat Methods* **19**, 11-12

604 24. Varadi, M., Anyango, S., Deshpande, M., Nair, S., Natassia, C., Yordanova, G., Yuan, D.,
605   Stroe, O., Wood, G., Laydon, A., Zidek, A., Green, T., Tunyasuvunakool, K., Petersen, S.,
606   Jumper, J., Clancy, E., Green, R., Vora, A., Lutfi, M., Figurnov, M., Cowie, A., Hobbs, N.,
607   Kohli, P., Kleywegt, G., Birney, E., Hassabis, D., and Velankar, S. (2022) AlphaFold Protein
608   Structure Database: massively expanding the structural coverage of protein-sequence
609   space with high-accuracy models. *Nucleic Acids Res* **50**, D439-D444

610 25. Skolnick, J., Gao, M., Zhou, H., and Singh, S. (2021) AlphaFold 2: Why It Works and Its
611   Implications for Understanding the Relationships of Protein Sequence, Structure, and
612   Function. *J Chem Inf Model* **61**, 4827-4831

613 26. Baumgrass, R., Williamson, M. K., and Price, P. A. (1997) Identification of peptide
614   fragments generated by digestion of bovine and human osteocalcin with the lysosomal
615   proteinases cathepsin B, D, L, H, and S. *J Bone Miner Res* **12**, 447-455

616 27. Phillips, C. M., Goumidi, L., Bertrais, S., Field, M. R., Ordovas, J. M., Cupples, L. A.,
617   Defoort, C., Lovegrove, J. A., Drevon, C. A., Blaak, E. E., Gibney, M. J., Kiec-Wilk, B.,
618   Karlstrom, B., Lopez-Miranda, J., McManus, R., Hercberg, S., Lairon, D., Planells, R., and
619   Roche, H. M. (2010) Leptin receptor polymorphisms interact with polyunsaturated fatty
620   acids to augment risk of insulin resistance and metabolic syndrome in adults. *J Nutr* **140**,
621   238-244

622 28. Dominguez, C., Boelens, R., and Bonvin, A. M. (2003) HADDOCK: a protein-protein
623   docking approach based on biochemical or biophysical information. *J Am Chem Soc* **125**,
624   1731-1737

29.   Rueda, P., Harley, E., Lu, Y., Stewart, G. D., Fabb, S., Diepenhorst, N., Cremers, B.,
Rouillon, M. H., Wehrle, I., Geant, A., Lamarche, G., Leach, K., Charman, W. N.,
Christopoulos, A., Summers, R. J., Sexton, P. M., and Langmead, C. J. (2016) Murine
GPRC6A Mediates Cellular Responses to L-Amino Acids, but Not Osteocalcin Variants.
*PLoS One* **11**, e0146846
30.   Power, M. J., and Fottrell, P. F. (1991) Osteocalcin: diagnostic methods and clinical
applications. *Crit Rev Clin Lab Sci* **28**, 287-335
31.   Teng, B., Huang, C., Cheng, C. L., Udduttula, A., Yu, X. F., Liu, C., Li, J., Yao, Z. Y., Long, J.,
Miao, L. F., Zou, C., Chu, J., Zhang, J. V., and Ren, P. G. (2020) Newly identified peptide
hormone inhibits intestinal fat absorption and improves NAFLD through its receptor
GPRC6A. *J Hepatol* **73**, 383-393
32.   Jeong, E., Kim, Y., Jeong, J., and Cho, Y. (2021) Structure of the class C orphan GPCR
GPR158 in complex with RGS7-Gbeta5. *Nat Commun* **12**, 6805
33.   Varadi, M., European Molecular Biology Laboratory, E. B. I., Hinxton, UK, Anyango, S.,
European Molecular Biology Laboratory, E. B. I., Hinxton, UK, Deshpande, M., European
Molecular Biology Laboratory, E. B. I., Hinxton, UK, Nair, S., European Molecular Biology
Laboratory, E. B. I., Hinxton, UK, Natassia, C., European Molecular Biology Laboratory, E.
B. I., Hinxton, UK, Yordanova, G., European Molecular Biology Laboratory, E. B. I.,
Hinxton, UK, Yuan, D., European Molecular Biology Laboratory, E. B. I., Hinxton, UK,
Stroe, O., European Molecular Biology Laboratory, E. B. I., Hinxton, UK, Wood, G.,
European Molecular Biology Laboratory, E. B. I., Hinxton, UK, Laydon, A., DeepMind, L.,
UK, Žídek, A., DeepMind, L., UK, Green, T., DeepMind, L., UK, Tunyasuvunakool, K.,
DeepMind, L., UK, Petersen, S., DeepMind, L., UK, Jumper, J., DeepMind, L., UK, Clancy,
E., DeepMind, L., UK, Green, R., DeepMind, L., UK, Vora, A., DeepMind, L., UK, Lutfi, M.,
DeepMind, L., UK, Figurnov, M., DeepMind, L., UK, Cowie, A., DeepMind, L., UK, Hobbs,
N., DeepMind, L., UK, Kohli, P., DeepMind, L., UK, Kleywegt, G., European Molecular
Biology Laboratory, E. B. I., Hinxton, UK, Birney, E., European Molecular Biology
Laboratory, E. B. I., Hinxton, UK, Hassabis, D., DeepMind, L., UK, Velankar, S., and
European Molecular Biology Laboratory, E. B. I., Hinxton, UK. (2022) AlphaFold Protein
Structure Database: massively expanding the structural coverage of protein-sequence
space with high-accuracy models. *Nucleic Acids Research* **50**
34.   Case, D. A., Cheatham, T. E., 3rd, Darden, T., Gohlke, H., Luo, R., Merz, K. M., Jr.,
Onufriev, A., Simmerling, C., Wang, B., and Woods, R. J. (2005) The Amber biomolecular
simulation programs. *Journal of computational chemistry* **26**, 1668-1688
35.   Maier, J. A., Martinez, C., Kasavajhala, K., Wickstrom, L., Hauser, K. E., and Simmerling,
C. (2015) ff14SB: Improving the Accuracy of Protein Side Chain and Backbone
Parameters from ff99SB. *J Chem Theory Comput* **11**, 3696-3713
36.   Darden, T., York, D., and Pedersen, L. (1993) Particle Mesh Ewald - an N.Log(N) Method
for Ewald Sums in Large Systems. *J Chem Phys* **98**, 10089-10092
37.   Price, D. J., and Brooks, C. L., 3rd. (2004) A modified TIP3P water potential for simulation
with Ewald summation. *J Chem Phys* **121**, 10096-10103
38.   Miyamoto, S., and Kollman, P. A. (1992) Settle - an Analytical Version of the Shake and
Rattle Algorithm for Rigid Water Models. *Journal of computational chemistry* **13**, 952-
962

39.    Zwanzig, R. (1973) Nonlinear generalized Langevin equations. *Journal of Statistical*
*Physics* **9**, 215-220

40.    Åqvist, J., Wennerström, P., Nervall, M., Bjelic, S., and Brandsdal, B. O. (2004) Molecular
dynamics simulations of water and biomolecules with a Monte Carlo constant pressure
algorithm. *Chemical physics letters* **384**, 288-294

41.    Nishimoto, S. K., and Price, P. A. (1979) Proof that the gamma-carboxyglutamic acid-
containing bone protein is synthesized in calf bone. Comparative synthesis rate and
effect of coumadin on synthesis. *J Biol Chem* **254**, 437-441

42.    Sabek, O. M., Nishimoto, S. K., Fraga, D., Tejpal, N., Ricordi, C., and Gaber, A. O. (2015)
Osteocalcin Effect on Human beta-Cells Mass and Function. *Endocrinology* **156**, 3137-
3146

43.    Nishimoto, S. K. (1990) A colorimetric assay specific for gamma-carboxyglutamic acid-
containing proteins: its utility in protein purification procedures. *Anal Biochem* **186**, 273-
279

44.    Nishimoto, S. K., and Price, P. A. (1985) The vitamin K-dependent bone protein is
accumulated within cultured osteosarcoma cells in the presence of the vitamin K
antagonist warfarin. *J Biol Chem* **260**, 2832-2836

45.    Pi, M., Parrill, A. L., and Quarles, L. D. (2010) GPRC6A mediates the non-genomic effects
of steroids. *J Biol Chem* **285**, 39953-39964

46.    Pi, M., Chen, L., Huang, M. Z., Zhu, W., Ringhofer, B., Luo, J., Christenson, L., Li, B.,
Zhang, J., Jackson, P. D., Faber, P., Brunden, K. R., Harrington, J. J., and Quarles, L. D.
(2008) GPRC6A null mice exhibit osteopenia, feminization and metabolic syndrome.
*PLoS One* **3**, e3858
692