Utilizing multi-objective decision support tools for protected area selection

Alke Voskamp[1*], Susanne A. Fritz[1,2], Valerie Köcke[3,4], Matthias F. Biber[5], Timo Nogueira Brockmeyer[6], Bastian Bertzky[7], Matthew Forrest[1], Allie Goldstein[8], Scott Henderson[8], Thomas Hickler[1,9], Christian Hof [5], Thomas Kastner[1], Stefanie Lang[10], Peter Manning[1,11], Michael B. Mascia[8], Ian McFadden[12,13], Aidin Niamir[1], Monica Noon[8], Brian O'Donell[14], Mark Opel[14,15], Georg Schwede[16], Peyton West[17], Christof Schenck[3,4], Katrin Böhning-Gaese[1,18]

*corresponding author

Affiliations:
[1] *Senckenberg Biodiversity and Climate Research Centre, 60325 Frankfurt am Main, Germany*
[2] *Goethe University, Institut für Geowissenschaften, 60438 Frankfurt, Germany*
[3] *Frankfurt Zoological Society, 60316 Frankfurt am Main, Germany*
[4] *Frankfurt Conservation Center gGmbH, 60316 Frankfurt am Main, Germany*
[5] *Terrestrial Ecology Research Group, Technical University of Munich, 85354 Freising, Germany*
[6] *Osnabrück University, 49069 Osnabrück, Germany*
[7] *European Commission, Joint Research Centre (JRC), I-21027 Ispra, Italy*
[8] *Conservation International, Airlington, VA, USA*
[9] *Institute of Physical Geography, Goethe University, 60438 Frankfurt, Germany*
[10] *Legacy Landscapes Fund, Nature Trust Alliance, Frankfurt am Main, Germany*
[11] *Department of Biological Sciences, University of Bergen, Bergen, Norway*
[12] *Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), Zürcherstrasse 111, 8903 Birmensdorf, Switzerland &*
[13] *Department of Environmental Systems Science, ETH Zürich, 8092 Zurich, Switzerland*
[14] *Campaign for Nature, Durango, Colorado, United States*
[15] *Independent Conservation Finance Adviser, Boulder, Colorado, United States*
[16] *Campaign for Nature, Badenweiler, Germany*
[17] *Frankfurt Zoological Society – U.S., Washington DC, USA*
[18] *Department of Biological Sciences, Goethe University Frankfurt, 60438 Frankfurt, Germany*

Running title: Global priorities for conservation

Article type: Perspective

Keywords: global change, biodiversity protection, conservation funding, conservation planning, long-term protection, decision making, strategic site selection, post-2020 biodiversity targets

Summary

The establishment and maintenance of protected areas (PAs) is viewed as a key action in

delivering post-2020 biodiversity targets. PAs often need to meet multiple objectives, ranging

from biodiversity protection to ecosystem service provision and climate change mitigation,

but available land and conservation funding is limited. Therefore, optimizing resources by

selecting the most beneficial PAs is vital. Here, we advocate for a flexible and transparent

approach to selecting protected areas based on multiple objectives, and illustrate this with a

decision support tool on a global scale. The tool allows weighting and prioritization of

different conservation objectives according to user-specified preferences, as well as real-time

comparison of the selected areas that result from such different priorities. We apply the tool

across 1347 terrestrial PAs and highlight frequent trade-offs among different objectives, e.g.,

between species protection and ecosystem integrity. Outputs indicate that decision makers

frequently face trade-offs among conflicting objectives. Nevertheless, we show that

transparent decision-support tools can reveal synergies and trade-offs associated with PA

selection, thereby helping to illuminate and resolve land-use conflicts embedded in divergent

societal and political demands and values.

Introduction

Halting biodiversity loss is one of the major global challenges faced by humanity in the 21st century[1,2]. Human wellbeing, livelihoods, and economies all rely on biodiversity, and collaborative international efforts are needed to conserve it[1,3]. Protected areas (PAs) are a cornerstone of biodiversity conservation. Aichi Target 11 of the Convention on Biological Diversity called for an increase in PA coverage to 17% by 2020 for the terrestrial realm, with a focus on PAs that are of particular importance for biodiversity and ecosystem services, ecologically representative and well connected[4]; this goal has only partly been reached[5]. Further, Aichi target 11 is increasingly seen as inadequate to safeguard biodiversity[6–8]. The Kunming-Montreal Global Biodiversity Framework (GBF), which builds on the Aichi targets, has set out 23 action oriented global targets in line with an ambitious plan to implement broad action which should transform our societies' relationship with biodiversity by 2030[9]. Action Target 3 of the GBF calls for at least 30 percent of the terrestrial area to be effectively conserved by PAs or "other effective area based conservation measures"[9]. This implies not only the transformation of large land areas into new PAs over the next decade, but also stresses an urgent need for careful allocation of the long-term conservation funding necessary to effectively protect biological resources: PAs must be both sustainably funded and effectively managed, yet only about 20% of all PAs are considered to meet these criteria[10]. Meanwhile, many PAs have experienced PA downgrading, downsizing or degazettement[11] (PADDD) or are threatened by PADDD in the future[11,12].

Both the allocation of sparse conservation funding for the strengthening of current PAs and the identification of additional sites to expand PA networks frequently require the application of prioritization approaches. A wealth of methods have been developed to inform conservation efforts, which vary widely in complexity. Some approaches evaluate individual sites based on their importance for the global persistence of biodiversity, e.g. the key biodiversity area (KBA) approach,

applying different threshold-based criteria including the proportion of threatened or geographically

restricted species covered[13]. In contrast, others rely on complex algorithms to optimize conservation

networks towards specific conservation goals, e.g. by considering complementarity, connectivity, or

cost efficiency[14–16].

Priority areas for biodiversity conservation can be defined based on one or more individual

conservation objectives, to identify areas of high conservation value under each or all given

objectives. Initial approaches to identify such areas sought hotspots of various aspects of biodiversity

such as species richness or endemism[17–20]. Other approaches highlight the protection of areas that will

limit further impacts of global change on biodiversity, for example, by identifying remaining

ecologically intact ecosystems[21] or sites of high irrecoverable carbon storage[22,23]. Prioritization

approaches that focus on more than one objective often combine different conservation goals like

protecting biodiversity and maintaining ecosystem services. Here, we focus on those prioritization

approaches that allow to identify individual sites of conservation importance rather than an optimized

network of sites.

The challenge: Aligning conservation priorities

Aligning different conservation objectives has become increasingly important. For instance,

conservation strategies that address both ongoing climate warming and biodiversity loss are urgently

needed[8,24]. Still, setting priorities based on multiple goals is not always straight forward. If there are

trade-offs among conservation objectives, a very different set of sites might be optimal under each

objective, and a simple compromise among these might not select the best set for the group of

objectives as a whole. Relying on approaches tailored towards a single conservation objective, or the

identification of one key element of the GBF targets, may lead to the omission of other critical

elements of the GBF vision[25].

To date, a vast amount of literature on setting global priorities for conservation is available (see Table

S1 for an overview relevant to this study). The different approaches vary in the number of objectives

that are considered, ranging from one to multiple, and the way the included variables are weighted,

not all or with equal or uneven weights (Table 1). One of the earliest efforts to highlight global areas

of importance for biodiversity protection are the global biodiversity hotspots identified by Meyers et

al (2000)[26]. These were derived based the number of endemic species and habitat loss in the area.

With the growing volume and availability of biodiversity data, more approaches to identify areas that

are important for biodiversity protection have been introduced. Examples for individual or combined

aspects of biodiversity that have been utilized for conservation priority maps are the global species

richness patterns for terrestrial vertebrates or vascular plants as well as for various other taxonomic

groups, but also biodiversity metrics such as species endemism, phylogenetic and functional diversity,

or threat status have been used[27–31]. Similarly, increasing data availability and spatial resolution of

those data has profited approaches that focus on prioritizing conservation sites based on the intactness

of habitats and biomes or ecoregions[33]. Generally, priority maps for biodiversity protection can be

derived based on a single metric for biodiversity or based on several combined metrics, as for

example by combining the biodiversity value of an area with the level of threat, through human

impacts like habitat degradation within the area[32,34] (see Table S1 for more examples).

Several efforts have also been made to align multiple conservation objectives, such as the protection

of biodiversity, the preservation of ecosystem services and the preservation of areas important for

climate mitigation. An example (Table 1) is the comparison of the spatial alignment of terrestrial

biodiversity, carbon storage, and water quality regulation, and the identification of areas with the

highest synergies among these objectives[35–37]. However, there is also evidence for trade-offs among

conservation objectives, e.g. biodiversity hotspots do not always overlap with different ecosystem

services[38]. In summary, a wealth of spatial prioritization maps for conservation efforts has been

produced by all these different approaches, either to combine different biodiversity metrics to identify

priority areas for biodiversity conservation or to align different conservation objectives to identify

priority areas across these objectives. In fact, Cimatti et al (2012) subsequently combined 63 different

global prioritization maps to derive one spatial prioritization map and identify scientific consensus

regions among the different approaches[39]. Nevertheless, all of these selection approaches have one

aspect in common: they result in a unique solution for one or a few specific and aligned objectives

that selects a static geographic set of priorities (Table 1). Here, we advocate a more flexible approach

that can handle multiple and conflicting objectives.

The weaker the alignment is among different conservation objectives, the greater the influence of

priority setting (i.e., favoring specific conservation objectives) on the outcome of site selection

approaches. If trade-offs are prevalent, explicit values-based decision making is necessary. The

relative priority of different conservation objectives varies among different societal groups, which

differ in their demands and values[40]. Also, key local, national, and international actors – governments,

corporations, non-governmental organizations (NGOs), scientists, and funders or sponsors – are likely

to differ in their priorities[41]. Therefore, decisions as to which areas should be prioritized are often

strongly values-based, with the values underlying final compromises rarely being made entirely

explicit and transparent. Societal and political values are also likely to change over time, since the

purpose of conservation itself has been transient over time, with priorities changing to some degree

from one generation to the next[42]. All of this substantiates the need for a flexible but transparent

approach to priority-setting, where different conservation objectives can be explicitly considered and

weighed against each other, to facilitate deliberative societal and political decision making.

Table 1: A comparison of strengths and weaknesses of the approach advocated and implemented in this study vs. already existing approaches. The table

summarizes a literature review, and gives a few selected examples from this. The review focused on studies that published global prioritization maps based on

one or multiple conservation objectives and which identified individual sites of conservation importance rather than designed an optimized network of sites

(see supplement and Table S1 for details and the considered studies).

Approach	Methods (Tools)	Strength and weaknesses	Example studies	Objectives considered in the example studies
Single objective	mapping	**+** Prioritization map based on one conservation objective **-** Solution for one objective	Di Marco et al 2012[43]; Riggio et al 2020[44]	ecosystem integrity
Multiple objectives	mapping, stacked layers	**+** combined prioritization map across multiple objectives **-** static solution, all objectives equally important	Jung et al 2021[36]; Dinerstein et al 2020[8]	biodiversity, ecosystem services, climate protection
Multiple objectives + fixed weights	mapping, stacked layers, consensus score	**+** combined prioritization map across multiple objectives **+** objectives (or variables within objectives) can be weighted individually **-** static solution	Freudenberger et al 2013[45]; Girardello et al 2019[46]	biodiversity, ecosystem services, ecosystem integrity
Multiple objectives + flexible weights	mapping, stacked layers, weighted consensus score, individual ranking of sites	**+** combined prioritization map across multiple objectives **+** comparison of tradeoffs on the fly **+** flexible solution	*This study*	biodiversity, ecosystem integrity, climate protection, climatic stability, land-use stability, size

Towards a solution: flexible and transparent site selection

The allocation of conservation funding is one example where the use of a flexible and transparent prioritization approach can be advantageous since the decision process is likely to involve multiple stakeholders, each of which may have multiple objectives. Use of a decision support tool can support the identification of conservation synergies and trade-offs, facilitate deliberation and dialog among stakeholders, and enable evidence-informed, values-based collaborative decision-making. Here, we illustrate these ideas using a site selection tool that we developed for this task. We apply a transparent site selection approach that allows users to identify investment priorities among existing PAs based on various self-specified conservation objectives. In contrast to other approaches, conservation objectives in our approach are explicitly weighted by the users and the results can be immediately assessed, aiding discussions during a transparent values-based decision-making process. We implemented the approach for the terrestrial realm, exclusively using biogeographic information that is publicly available at a global scale. We aimed to identify areas with the highest potential for a range of biodiversity and climate protection goals, but excluded any information on political and economic dimensions from the site selection algorithm; although these considerations are crucial for conservation and should be evaluated equally transparently, we believe that they should be evaluated separately from biogeographic information as an additional step in the decision-making process.

We defined six different conservation objectives (Fig. 1), which represent a broad agreement on priorities for safeguarding biodiversity, climate protection (in the sense of mitigating ongoing climate change), and the present and projected future status of individual sites (identified in an initial stakeholder dialog, see also case study details below). These objectives were: 1) high current biodiversity, focusing on high biodiversity values, 2) high current ecosystem integrity, which focuses on areas that have experienced relatively few anthropogenic impacts, 3) high climate protection, which selects for sites that have large, irreplaceable carbon stocks, 4) large size, which prioritizes

bigger sites, 5) high land-use stability, which focuses on the future likelihood of land-use change in

the immediate surroundings of sites, and 6) high climatic stability, which highlights sites in which

climate change is projected to have low impacts on current biodiversity.

We collated a broad set of conservation indicators that reflect these six conservation objectives (Fig.

1). The biodiversity objective considered as indicators the total terrestrial species richness of four

vertebrate taxa (birds, mammals, amphibians and reptiles) as well as species endemism and

evolutionary diversity[47] for each taxon, to capture the amount of biodiversity as well as its

irreplaceability. The ecosystem integrity objective considered biodiversity intactness, recent land-use

change, and the human footprint within the site. The climate protection objective considered the

average amount of carbon per hectare that is stored in the vegetation and soil (up to 1 meter below

ground) of the site and its vulnerability to typical land conversion. The size objective covers the extent

of the site in km^2. The land-use stability objective considered the projected change in land-use in a

buffer zone around the site. The climatic stability objective considered the biodiversity change based

on the projected future compositional change (turnover)[48] of the four vertebrate taxa and the projected

change in tree cover within the site.

These conservation objectives and the underlying indicators were carefully selected reflecting the

demands towards the PA network based on the post-2020 GBF, as well as the current state of the

literature addressing both the biodiversity and climate crises. The biodiversity objective combines

information on the number, diversity and rarity of species across several higher taxa within the area,

to include different aspects of biodiversity[47,49–52]. Highlighting those sites that are of particular

importance for biodiversity is in line with the first part of Action Target 3 of the post-2020 GBF[9]. The

ecosystem integrity objective uses information on recent impacts on the site and the intactness of the

local ecological communities, highlighting those sites that contain ecosystems that are still largely

intact. This objective was included because remaining intact ecosystems are often not directly

addressed by conservation efforts or international policy frameworks[21,53], but provide various key

functions, such as acting as critical carbon sinks, stabilizing hydrological cycles, or providing crucial

refuge for imperiled species, intact mega-faunal assemblages, or wide-ranging or migratory

species[21,54–59]. The size objective is somewhat related to the ecosystem integrity objective, under the

assumption that larger areas have a higher potential to support populations of target species and to

maintain functioning ecosystems in the long term[60,61]. The climate protection objective is related to

Action Target 8 of the post-2020 GBF, which aims to minimize the impacts of climate change on

biodiversity.

The final two objectives were included to assess sites not only based on their current importance for

biodiversity, ecosystem functioning, and climate protection, but also based on the most major future

threats towards biodiversity, i.e. projected future climate and land-use change. The five direct drivers

of biodiversity loss with the largest impact, according to the 2019 Global Assessment Report by

IPBES, are changes in land and sea use; direct exploitation of organisms; climate change; pollution;

and invasion of alien species[1]. The climatic and land-use stability objectives provide an indication of

potential future changes within the site based on climate change responses (geographic range shifts)

of the local flora and fauna within the region and give an indication of which sites might be under

increasing pressure of land-use change in the region.

A key aspect in developing a transparent site selection approach was to make results of different

values-based objective weighting immediately accessible to a broader audience, including decision

makers. We therefore developed an open-source spatial decision support tool to facilitate the priority-

based area selection process. The tool generates a ranking of sites globally as well as for each

biogeographic realm, based on the six conservation objectives which are weighted individually by the

user. Using sliders to allocate weights to the six conservation objectives, users can design their own

conservation scenarios on the fly (examples see Fig. 1), and directly visualize the resulting ranking.

The tool allows a comparison of a far wider range of different conservation scenarios than the

examples we give here, to evaluate synergies and trade-offs among these, and select sites for a more

detailed investigation. The current version is publicly available (https://ll-evaluation-support-

tool.shinyapps.io/legacy_landscapes_dst/) and restricted to the case study dataset, objectives and

indicators presented in the paper, but the flexible approach we use can be implemented easily to other

datasets, objectives, and goals.

Fig. 1: The six conservation objectives defined to set priorities for the site selection, the indicators

considered for each objective (note that Biodiversity and Climatic stability (of biodiversity) include

indicators for four different vertebrate taxa), and examples for conservation scenarios based on these

objectives. By applying a weighting approach, user-specified objectives can be combined into

different conservation scenarios, which are therefore customized for specific conservation goals. The

High biodiversity, High ecosystem integrity and Legacy Landscapes Fund (LLF) scenarios are used in

the case study.

Illustration of the selection approach: The Legacy Landscapes Fund as a case study

The Legacy Landscapes Fund (LLF) is a recently established foundation that provides long-term funding for protected areas[62]; it is useful in this context because it uses our six conservation objectives, operates on a global level, and mostly focuses on existing sites. This allowed us to run a case study across a significant set of PAs and other sites of interest across the globe, in order to demonstrate how the newly developed decision support tool facilitates the flexible evaluation of potential priority sites for conservation and to explore the potential and limitations of this approach. We assessed synergies and trade-offs among areas according to the different objectives at a global scale, as well as within biogeographic realms. Finally, we aimed to investigate how priority setting by different societal actors affects site selection by combining the multiple conservation objectives into broader conservation scenarios that weigh each objective according to user-specified priorities.

The case study dataset for the analysis contained 1347 sites globally. These sites included formally protected areas of IUCN category I or II, listed Natural World Heritage Sites (WHS) and registered Key Biodiversity Area (KBA) (see experimental procedures and supplementary material for details on dataset and methods)[63,64]. A principal component analysis (PCA) applied to this dataset globally (Fig. 2) and at the level of biogeographic realms (Fig. 3) showed that the indicators belonging to each conservation objective tended to be closely aligned both at the global and the realm level, with the only exception being the two climatic stability indicators across the Australian realm. For example, within the biodiversity objective, species richness (SR), species endemism (SE) and evolutionary diversity (ED) were closely aligned at the global scale, as well as at the biogeographic realm level, though the alignment between SR and the other two indicators was slightly less tight in the tropical realms (Fig. 3).

Looking at the trade-offs and synergies among the objectives, we found that at the global scale the first and second PCA axes explained 31.4 and 14.2 percent of the variation in the data respectively.

These axes showed relatively clear trade-offs and synergies among the six different conservation

objectives (Fig. 3). The strongest global trade-off was found between current biodiversity and future

land-use stability (Pearson's correlation coefficient r (n=1346) = -.30, p<0.01). These two objectives

are negatively correlated, as increasing land-use pressure is often projected to occur around sites with

exceptionally high current biodiversity (e.g. deforestation of tropical forests for agriculture). The

strongest global synergies were found between current biodiversity and future climatic stability (r

(n=1346) = .41, p<0.01) and current biodiversity and high climate protection potential based on the

amount of manageable carbon stored in the site (r (n=1346) = .58, p<0.01). This suggests that sites

with exceptionally high biodiversity often coincide with areas of lower projected impacts of climate

change on vertebrate communities and tree cover and with a high potential for climate protection

through carbon storage. The identified global synergies and trade-offs between the different objectives

were only partially consistent within realms, with patterns very similar to the global analysis for the

Afrotropical realm but notably different alignments in the Palearctic and Nearctic.

Finally, to investigate how priority setting by different societal groups can affect site selection, we

compared the outcome of area selection under three different conservation scenarios. We used two

extreme and one combined scenario, to explore a broad range of values (Fig. 1). The first scenario

was a biodiversity scenario (biodiversity objective weighted by 100% and the other five objectives by

0%). The second was an ecosystem integrity scenario (ecosystem integrity 100%, all others 0%). The

third scenario was a stakeholder-driven scenario that resulted from joint discussion during an expert

workshop (LLF scenario; Fig. 1). At this two-day online workshop, which was attended by 35 experts

with a strong conservation background, we introduced the site selection approach, further developed

the indicators and objectives, and voted on the LLF scenario (see supplementary materials for more

detail). This scenario reflects the main selection criteria for potential LLF sites (high biodiversity,

ecosystem integrity and size) but considers also the other objectives weighted according to lower

priorities (biodiversity, ecosystem integrity and size weighted with 25% each, climatic stability and

land-use stability with 10% each, and climate protection with 5%).

Despite synergy between some objectives, we found that when comparing the top five sites selected

for each of the three conservation scenarios, within each biogeographic realm, there is little

congruence among these scenarios (Fig. 4). This implies that selecting sites based on their

biodiversity will in most cases result in the protection of different sites compared to a selection based

on high ecosystem integrity, or the LLF scenario. Australasia has the highest overlap of top sites for

the three different scenarios, with four sites being in the top five for both the biodiversity and the LLF

scenario. The Nearctic, Neotropic and Afrotropic realms have the least overlap among the top sites

for the investigated scenarios with only one shared site in the top five of all scenarios.

Fig. 2: Trade-offs and synergies between the conservation indicators of individual sites. Shown

are the first and second dimensions of a principal component analysis (PCA) that was performed

across 1347 sites and their variation in 13 indicator variables aggregated into six conservation

objectives (order of indicator variables in the legend aligns with Fig. 1 and 3, see these for matching

variables to objectives). The first and second PCA dimensions together explain 45.3% of the variation

in the data. Each dot represents one site. The arrows represent the indicators and the arrow length

indicates the loading of each indicator onto the PCA dimensions (i.e. their correlation with each

principal component). Opposite loadings indicate trade-offs between the variables (i.e., a site that has

a high value in one of these variables, has a low value in the other variable and vice versa). The

individual sites (points) are colored by the biogeographic realm in which they are located[65].

Fig. 3: Trade-offs and synergies between the conservation indicators of individual sites at the global and realm levels. Shown are the first two axes of the principal component analysis (PCA) for all 1347 sites included in the Legacy Landscapes case study globally and for each individual realm. These analyses reveal trade-offs between the conservation objectives, indicated by variables mapping onto opposing ends of a principal component axis. Variable colours indicate conservation objectives as in Fig. 1: biodiversity (shades of green), ecosystem integrity (shades of red and pink), climate protection (shades of blue), size (dark brown), land-use stability (light brown) and climatic stability (orange and yellow). PCA plots show the respective first two axes identified and the percentage of variation explained by each of the axes.

Fig. 4: **Spatial distribution of sites highlighting the top 5 priority sites for each of the 3 example**

conservation scenarios: prioritizing biodiversity (dark green), prioritizing ecosystem integrity (red)

and the LLF scenario (Legacy Landscapes Fund, prioritizing a combination of all objectives that

stresses high biodiversity, high ecosystem integrity and large size; blue). The top 5 sites for all three

scenarios (triangles) are shown per biogeographic realm (i.e., 30 top sites per conservation scenario in

total). The colors correspond to the three different conservation scenarios and their overlap (if a site is

in the top five for more than one objective), as shown in the Venn diagram. Only 14 of the top sites

were selected under two scenarios (light green, brown and orange) and 1 site was selected under all 3

scenarios (yellow). Grey points indicate sites included in the analysis but not selected under the top 5.

Top sites in close geographic proximity are spaced out for visualization and deviate from their exact

spatial position. Map colors indicate the different biogeographic realms.

Discussion

Our case study demonstrates that the selection of 'best' sites for nature conservation depends largely on the relative weighting of different conservation priorities and is therefore heavily influenced by decision-maker values. This is supported by the clear trade-offs among the six conservation objectives at the realm and global scale (Fig. 2, 3), as well as the limited congruence among the top sites selected under the three different conservation scenarios (Fig. 4). These results illustrate the opportunities and challenges faced by decision makers when selecting priority areas for nature conservation. Furthermore, they demonstrate the need for a global approach to nature conservation that involves multiple stakeholder groups and perspectives and a transparent decision-making process.

Here, we introduce an approach to select priority areas for biodiversity conservation at the global scale that separates 1. global biogeographic information on biodiversity, ecosystem services, etc., from 2. a value-based prioritization of different conservation objectives in the decision-making process. This allows the trade-offs between conservation objectives to be understood and acknowledged explicitly and quantitatively. It thereby enables a first transparent evaluation of sites that reflects the varying priorities among different societal or conservation actors. Furthermore, the approach allows to optimize site selection towards more than one objective, which can significantly increase the efficiency of a PA network[66]. Additionally, the transient nature of conservation goals or new drivers of biodiversity loss, such as climate change, might result in the need to adjust prioritization in the future. Both arguments highlight the advantages of a flexible site selection approach over the static selection of hotspots based on a small number of fixed objectives and indicators.

Our approach goes beyond existing studies that explore the spatial agreement of conservation objectives and present optimized solutions through aligning several objectives, by allowing the user to change the prioritization on the fly (Table 1). Instead of presenting a static conservation priority map, we present a dynamic result that ranks potential sites for protection based on user preferences. This approach puts the focus on the decision making process and allows the exploration of tradeoffs and synergies among different options. Rather than providing another method to set conservation priorities, our approach is complementary to the various approaches we found in the literature (Table

1 and S1). It could for example be used to explore the differences, synergies and tradeoffs between

any of the existing global prioritization maps, across protected areas.

Applying the tool to a specific conservation problem

For the Legacy Landscapes Fund, the three conservation objectives of size, biodiversity and

ecosystem integrity are of high priority[67]. Applying the decision support tool to the assembled dataset

revealed a trade-off between high biodiversity and high ecosystem integrity, clearly demonstrated in

the comparison between the three conservation scenarios: high biodiversity, high ecosystem integrity

and the LLF scenario, which considers multiple conservation objectives. For the actual area selection

to be financed by the LLF, the decision support tool enabled an initial screening of potential sites

globally, to evaluate the performance of individual sites under the desired conservation objectives and

to compare different weightings before proceeding with the selection of the pilot sites. Here, the

decision support tool was used in an integrative decision-making process which transparently

separated biogeographical site screening from other criteria like stakeholder consent, political

commitment, and experience of the implementing NGO (also see below).

Applying the approach beyond the case study

Our approach and the newly developed tool can be easily extended to include a broader range of

biogeographic datasets, additional conservation objectives, or additional sites into the analysis,

making the tool widely applicable to a variety of site selection tasks. Though the current set-up of the

tool already contains six objectives representing several broad conservation goals (i.e. safeguarding

biodiversity or mitigating climate change), these are still to some extent geared towards the case

study. To broaden the scope of the tool through additional objectives and opposing the focus on intact

ecosystems used in our case study, priority setting could highlight areas that harbor a high amount of

threatened biodiversity[68], e.g. by including an additional objective based on the threat status of all

occurring species (i.e. as provided in the IUCN Red List) in a site[49,69,70]. Another obvious and easy

possibility to expand the current set-up of the tool would be to allow further subsetting of the included

sites. Currently the tool allows for an initial screening of sites at the level of biogeographic realms or

at the global scale. Information such as the extent of a biogeographic realm or ecoregion that is

already protected would need to be considered separately. Adjusting the tool to rank sites not only at

the realm level but also at finer scales, as for example at the ecoregion level, would allow users to

prioritize sites in finer-scale underrepresented categories.

Action Target 8 of the post 2020 GBF also calls for a well-connected PA network[9]. Connectivity is

highly species-specific and landscape-dependent, and thus requires local and long-term studies on

individual species[71,72]. Assessments on a scale like the decision support tool shown here cannot yet

assess connectivity at that level. Still, previous efforts have estimated the connectivity of global PA

networks at a coarser scale, for example based on different levels of home range size in mammals[73] or

even by modeling the movement of large animals throughout the landscape between protected areas[74].

A first step to integrate connectivity into the decision support tool could be to use a distance matrix of

sites from surrounding existing PAs. This could give a first rough indication of how well a site is

embedded into the PA network and allow prioritization of connected sites over very isolated sites.

As currently designed, the tool is meant to allow the comparison of sites and different conservation

objectives based on biogeographic variables, which are available at a global scale. This necessitates

the use of relatively coarse-grained datasets (resolution here is mostly dependent on the biodiversity

data). The tool allows an initial screening of a large number of potential sites globally (or regionally)

and can be extremely useful in creating prioritizations of PAs based on different objectives and

indicators that can be applied flexibly. This tool, however, is only useful as a first step that allows a

range of options to be explored, as part of a much broader decision-making process. This decision-

making process should include on-site assessments of additional parameters at a higher resolution

(e.g. more detailed biological data acquired through surveys and observations) as well as non-

biological characteristics. These socio-economic factors could include, for example, the political

legitimacy of the initiative, the involvement of local communities, and the presence of a supportive

NGO. In case of pilot site selection for the LLF, these factors were considered in the next step that

followed the use of the site evaluation tool. Further, the decision support tool was designed to

facilitate value-based discussions by enabling on-the-fly comparison of sites based on different

biogeographic attributes. The tool does not facilitate the optimization of site networks (i.e. assess

different combinations of sites based on representativeness or cost efficiency).

Applying the decision support tool within the post-2020 Global Biodiversity Framework

The ambition of the Aichi Biodiversity Targets has been increasingly criticized as being too modest to

safeguard biodiversity in perpetuity[6,7]. Accordingly, the post-2020 GBF of the Convention on

Biological Diversity calls for 'at least 30 per cent of terrestrial, inland water and of coastal and marine

areas, especially areas of particular importance for biodiversity and ecosystem functions and services,

to be effectively conserved'[9]. Thus it becomes increasingly important to identify new sites for

conservation – and new ways of conserving – outside of the already delineated areas both on land and

in the oceans[8,75]. The presented decision support tool could be extended to aid these efforts, either by

adapting it to identify new sites or by expanding the case-study dataset. A first possible extension

would be the inclusion of the not yet formerly recognized Indigenous and Community Conservation

Areas (ICCAs) and of Other Effective Area-based Conservation Measures (OECMs) which are

increasingly being recognized as effective and potentially more inclusive conservation tools[76].

Going beyond global priority-setting, the post-2020 GBF aims to facilitate implementation primarily

through activities at the national level. Furthermore, unlike in the LLF case study, a vast amount of

conservation funding is not available at the global scale but rather at the national or regional level.

Our approach could be used at the national or sub-national level to help prioritize conservation

decisions through facilitating transparent value-based discussion and support implementation of the

post-2020 GBF at this scale[77]. Applying the tool at the national or regional scale would open the

possibility to add more finely resolved datasets to the conservation objectives that are not available at

the global scale (for example, species abundances or more specific land-use projections) and thus

tailor the decision support tool to specific conservation actions.

An example of a relevant adjustment that may be possible at national scales could be the adjustment

of the intended timeframe, as the decision support tool with its inclusion of future projections

(climatic and land-use stability) as well as the focus on intact ecosystems is currently geared towards

longer time horizons. Highlighting sites where there is an urgent need to act (e.g. within a couple of

years because of high conservation value in combination with high current pressure) would require

the use of very different datasets with a much higher resolution. Working at regional or national scales

would allow the inclusion of data sets on recent changes within a site that are not available or very

heterogeneous at the global scale (e.g. population trends, recent deforestation rates, or the level of

exploitation of natural resources).

In conclusion, the proposed approach facilitates a transparent initial screening of potential priority

sites that allows the trade-offs between conservation objectives to be understood and acknowledged

explicitly and quantitatively. It promotes the inclusion of multiple stakeholder positions, views and

preferences, and facilitates discourse and decision-making whilst working towards the overarching

conservation goals.

Experimental procedures

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled

by the Lead Contact, Alke Voskamp (alke.voskamp@posteo.net)

Materials availability

This study did not generate unique new materials.

Data and code availability

All codes needed to replicate the presented analysis are available from GitHub

(https://github.com/Legacy-Landscapes/LL_analysis). The decision support tool is accessible via:

https://ll-evaluation-support-tool.shinyapps.io/legacy_landscapes_dst/). All codes for the decision

support tool are available under https://github.com/Legacy-Landscapes/LL_Decision_Tool.

Conservation objectives data

The six defined conservation objectives are each based on several underlying data sets, with more detail on variable calculations and score assignations for each objective given in the supplement. The datasets behind the biodiversity objective are the global range-map polygons for all terrestrial birds, mammals, amphibians and reptiles as provided by BirdLife International, IUCN and GARD[49–51], as well as the phylogenetic supertree for all four terrestrial vertebrate taxa from Hedges *et al*. 2015[78]. From these datasets we derived species richness, species endemism (calculated as corrected range size rarity[52]) and phylogenetic endemism[47] values per site included for all four vertebrate taxa. The datasets underlying the ecosystem integrity objective are the biodiversity intactness index[79], the human footprint compiled by Venter et al 2016[80] and the recent land-use change 1992 -2018 derived from the ESA CCI Land Cover by Niamir et al 2020[81]. The climate protection objective consists of three different indicators, the amount of manageable carbon stored in the site, the amount of vulnerable carbon and the amount of irrecoverable carbon[22,23]. The size objective uses the size of each site derived in QGIS[82]. All future stability variables were derived by comparing the timespan between 1995 (average climate projections 1980 – 2009) and 2050 (average climate projections 2035 – 2064). The climatic stability objective consists of two main underlying indicators, the climatic stability of biodiversity and the projected tree cover change. The climatic stability was calculated based on modelled changes in species community compositions that resulted from projected range shifts under climate change for all four taxa[83]. The projected change in tree cover is based on the LPJ-GUESS process-based dynamic vegetation-terrestrial ecosystem model[84]. Finally, the land-use stability objective consists of projected changes in five different land-use types (rainfed crop, irrigated crop, pastures, as well as rainfed and irrigated bioenergy crops), based on the MAgPIE and REMIND-MAgPIE model[85–87] and using the assumptions of population growth and economic development as described in Frieler et al 2017[88]. These projections are based on the same climate projections as the climatic stability variables.

The six conservation objectives were developed in a discussion process among the broader

conservation community. We introduced our approach at a two-day webinar which was attended by

35 experts with a strong conservation background. These included 1) conservation scientists, 2)

international conservation NGOs, 3) the financial sector, and 4) policy sectors, in particular the

German Federal Ministry for Economic Cooperation and Development (BMZ). These experts

provided feedback on the objectives and indicators through a questionnaire (see supplementary

material). They were asked to: 1) report any missing objectives, 2) report any missing indicators that

should be included in the objectives and 3) rank the suggested objectives by their personal

preferences. To translate personal preferences into site selection, the resulting ranks for each

individual indicator were scaled from zero to one. Each objective consists of several underlying

indicators (datasets), so by taking the mean across all indicators per objective these were weighted

equally.

The case study dataset and analysis

To assess synergies and trade-offs among the conservation objectives, we used the LLF as a case

study to assemble a global dataset of sites. The LLF is a recently established foundation that provides

long-term funding of one million U.S. dollars per "legacy landscape" per year. Funding stems from

public and private sources. It aims to protect areas of outstanding biodiversity over initially 15 years –

but with a vision to ensure funding in perpetuity[67]. The LLF is based on a strategic global site-

selection approach and the strong long-term commitment of local NGOs, protected area authorities

and local communities 'on the ground'[62]. The initial requirements for sites to be considered by the

LLF are outstanding biodiversity, a minimum size of 2,000km^2 and a protection status as IUCN

protected area category I or II for at least 1,000 km^2. Based loosely on these guidelines, we assembled

a dataset and extracted site-specific values for each objective (Fig. 1) (see supplementary material for

a detailed account how the site dataset was assembled).

We then investigated global synergies and trade-offs among the final set of conservation objectives

using a principal component analysis (PCA) across sites. To further explore if synergies and trade-offs

between the objectives were different in biogeographic regions of the world, we repeated the PCA

separately for each of the six terrestrial biogeographic realms[65]. Additional analyses are described in

the supplement.

The decision support tool

To make the analysis accessible to the broader conservation community and to enable a rapid

comparison of sites based on the user-specified prioritization of the different conservation objectives,

we designed an interactive spatial decision support tool in which weightings can be modified (see

supplementary material for detailed content of the app interface). The user interface for the tool was

developed using R Shiny version 1.5.0[89].

Acknowledgements

We thank all participants of the expert workshop for valuable discussions and input to the development

of the decision support tool, and the FZS staff in the project areas who tested the tool and helped to

evaluate its use for conservation. We gratefully acknowledge the use of the Goethe-HLR HPC at the

Centre for Scientific Computing at Goethe University Frankfurt for some of the computationally heavy

aspects of this work. We also thank BirdLife International for making the KBA data available as well

as the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP) and ISIpedia - the open climate-

impacts encyclopedia for their support and data availability.

Funding statement

We thank the Temperatio Foundation for their financial support. SF was supported by the German

Research Foundation DFG (FR 3246/2-2) and the Leibniz Competition of the Leibniz Association

(P52/2017); AN was supported by European Union's Horizon 2020 research and innovation program

under grant agreement No. 689443.

Author contributions

Conceptualization: AV, SAF, VK, CS and KBG; Methodology: AV, SAF, KBG; Feedback on

Methodology: all authors, Software: AV, TNB, MFB; Writing – Original: AV, SAF, VK and KBG;

Writing – Review and Editing: all authors, Supervision: SAF and KBG.

References

1. IPBES (2019). Summary for policymakers of the global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services S. Díaz, J. Settele, E. S. Brondizio, H. T. Ngo, M. Guèze, J. Agard, A. Arneth, P. Balvanera, K. A. Brauman, S. H. M. Butchart, et al., eds. (IPBES secretariat).

2. Pereira, H.M., Leadley, P.W., Proença, V., Alkemade, R., Scharlemann, J.P.W., Fernandez-Manjarrés, J.F., Araújo, M.B., Balvanera, P., Biggs, R., Cheung, W.W.L., et al. (2010). Scenarios for Global Biodiversity in the 21st Century. Science. *330*, 1496–1501.

3. Dasgupta, P. (2021). The Economics of Biodiversity: The Dasgupta Review.

4. Convention on Biological Diversity (2010). "Strategic Plan for Biodiversity 2011 - 2020, including Aichi Biodiversity Targets." https://www.cbd.int/sp/.

5. Secretariat of the Convention on Biological Diversity (2020). Global Biodiversity Outlook 5.

6. Noss, R.F., Dobson, A.P., Baldwin, R., Beier, P., Davis, C.R., Dellasala, D.A., FRANCIS, J., Locke, H., Nowak, K., Lopez, R., et al. (2012). Bolder Thinking for Conservation. Conserv. Biol. *26*, 1–4.

7. Larsen, F.W., Turner, W.R., and Mittermeier, R.A. (2015). Will protection of 17% of land by 2020 be enough to safeguard biodiversity and critical ecosystem services? Oryx *49*, 74–79.

8. Dinerstein, E., Joshi, A.R., Vynne, C., Lee, A.T.L., Pharand-Deschênes, F., França, M., Fernando, S., Birch, T., Burkart, K., Asner, G.P., et al. (2020). A "Global Safety Net" to reverse biodiversity loss and stabilize Earth's climate. Sci. Adv. *6*.

9. Secretariat of the United Nations Convention on Biological Diversity (2022). Kunming-Montreal Global biodiversity framework Draft decision submitted by the President. https://www.cbd.int/doc/c/e6d3/cd1d/daf663719a03902a9b116c34/cop-15-l-25-en.pdf.

10. Dasgupta, P. (2021). The Economics of Biodiversity: The Dasgupta Review, Abridged Version, Chapter 19 Natures Supply and Our Demands.

11. Golden Kroner, R.E., Qin, S., Cook, C.N., Roopa, K., Pack, S.M., Bonilla, O.D., Cort-

Kansinally, K.A., Coutinho, B., Feng, M., Martínez Garcia, M.I., et al. (2019). The uncertain

future of protected lands and waters. Science. *364*, 881–886.

12.   Kuempel, C.D., Chauvenet, A.L.M., Symes, W.S., and Possingham, H.P. (2021). Predicted

protected area downsizing impedes conservation progress across terrestrial ecoregions in the

tropics and subtropics. Conserv. Sci. Pract.

13.   IUCN (2016). A global standard for the identification of Key Biodiversity Areas, Version 1.0

(IUCN).

14.   Margules, C.R., and Pressey, R.L. (2000). Systematic conservation planning. Nature *405*, 243–

253.

15.   Moilanen, A., Franco, A.M.A., Early, R.I., Fox, R., Wintle, B., and Thomas, C.D. (2005).

Prioritizing multiple-use landscapes for conservation: methods for large multi-species planning

problems. Proc. R. Soc. B Biol. Sci. *272*, 1885–1891.

16.   Moilanen, A., Wilson, K., and Possingham, H. (2009). Spatial conservation prioritization:

quantitative methods and computational tools (Oxford University Press).

17.   Jenkins, C.N., Pimm, S.L., and Joppa, L.N. (2013). Global patterns of terrestrial vertebrate

diversity and conservation. Proc. Natl. Acad. Sci. *110*, E2602–E2610.

18.   Orme, C.D.L., Davies, R.G., Burgess, M., Eigenbrod, F., Pickup, N., Olson, V.A., Webster,

A.J., Ding, T.-S., Rasmussen, P.C., Ridgely, R.S., et al. (2005). Global hotspots of species

richness are not congruent with endemism or threat. Nature *436*, 1016–1019.

19.   Satterfield, A.J., Crosby, M.J., Long, A.J., and Wege, D.C. (1998). Endemic Bird Areas of the

World: Priorities for Biodiversity Conservation (Birdlife International).

20.   Ceballos, G., and Ehrlich, P.R. (2006). Global mammal distributions, biodiversity hotspots,

and conservation. Proc. Natl. Acad. Sci. *103*, 19374–19379.

21.   Watson, J.E.M., Venter, O., Lee, J., Jones, K.R., Robinson, J.G., Possingham, H.P., and Allan,

J.R. (2018). Protect the last of the wild. Nature *563*, 27–30.

22.   Goldstein, A., Turner, W.R., Spawn, S.A., Anderson-Teixeira, K.J., Cook-Patton, S., Fargione,

J., Gibbs, H.K., Griscom, B., Hewson, J.H., Howard, J.F., et al. (2020). Protecting

irrecoverable carbon in Earth's ecosystems. Nat. Clim. Chang. *10*, 287–295.

23.   Noon, M.L., Goldstein, A., Ledezma, J.C., Roehrdanz, P.R., Cook-Patton, S.C., Spawn-Lee,

S.A., Wright, T.M., Gonzalez-Roglich, M., Hole, D.G., Rockström, J., et al. (2021). Mapping

the irrecoverable carbon in Earth's ecosystems. Nat. Sustain. *5*, 37–46.

24.   Pörtner, H.O., Scholes, R.J., Agard, J., Archer, E., Arneth, A., Bai, X., Barnes, D., Burrows,

M., Chan, L., Cheung, W.L., et al. (2021). Scientific outcome of the IPBES-IPCC co-

sponsored workshop on biodiversity and climate change.

25.   Allan, J.R., Possingham, H.P., Atkinson, S.C., Waldron, A., Di Marco, M., Butchart, S.H.M.,

Adams, V.M., Kissling, W.D., Worsdell, T., Sandbrook, C., et al. (2022). The minimum land

area requiring conservation attention to safeguard biodiversity. Science (80-.). *376*, 1094–

1101.

26.   Myers, N., Mittermeier, R.A., Mittermeier, C.G., da Fonseca, G.A.B., and Kent, J. (2000).

Biodiversity hotspots for conservation priorities. Nature *403*, 853–858.

27.   Voskamp, A., Baker, D.J., Stephens, P.A., Valdes, P.J., and Willis, S.G. (2017). Global

patterns in the divergence between phylogenetic diversity and species richness in terrestrial

birds. J. Biogeogr. *44*, 709–721.

28.   Daru, B.H., le Roux, P.C., Gopalraj, J., Park, D.S., Holt, B.G., and Greve, M. (2019). Spatial

overlaps between the global protected areas network and terrestrial hotspots of evolutionary

diversity. Glob. Ecol. Biogeogr. *28*, 757–766.

29.   Mazel, F., Guilhaumon, F., Mouquet, N., Devictor, V., Gravel, D., Renaud, J., Cianciaruso,

M.V., Loyola, R., Diniz-Filho, J.A.F., Mouillot, D., et al. (2014). Multifaceted diversity–area

relationships reveal global hotspots of mammalian species, trait and lineage diversity. Glob.

Ecol. Biogeogr. *23*, 836–847.

30.   Brum, F.T., Graham, C.H., Costa, G.C., Hedges, S.B., Penone, C., Radeloff, V.C., Rondinini,

C., Loyola, R., and Davidson, A.D. (2017). Global priorities for conservation across multiple

dimensions of mammalian diversity. Proc. Natl. Acad. Sci. *114*, 7641–7646.

31.   Kier, G., Kreft, H., Lee, T.M., Jetz, W., Ibisch, P.L., Nowicki, C., Mutke, J., and Barthlott, W.

(2009). A global assessment of endemism and species richness across island and mainland

regions. Proc. Natl. Acad. Sci. *106*, 9322–9327.

32. Mokany, K., Ferrier, S., Harwood, T.D., Ware, C., Di Marco, M., Grantham, H.S., Venter, O.,

Hoskins, A.J., and Watson, J.E.M. (2020). Reconciling global priorities for conserving

biodiversity habitat. Proc. Natl. Acad. Sci. *117*, 9906–9911.

33. Beyer, H.L., Venter, O., Grantham, H.S., and Watson, J.E.M. (2020). Substantial losses in

ecoregion intactness highlight urgency of globally coordinated action. Conserv. Lett. *13*,

e12692.

34. Gumbs, R., Gray, C.L., Böhm, M., Hoffmann, M., Grenyer, R., Jetz, W., Meiri, S., Roll, U.,

Owen, N.R., and Rosindell, J. (2020). Global priorities for conservation of reptilian

phylogenetic diversity in the face of human impacts. Nat. Commun. *11*, 2616.

35. Nature Map (2020). Nature Map Earth. See: https://naturemap.earth/ [accessed 31.01.22].

36. Jung, M., Arnell, A., de Lamo, X., García-Rangel, S., Lewis, M., Mark, J., Merow, C., Miles,

L., Ondo, I., Pironon, S., et al. (2021). Areas of global importance for conserving terrestrial

biodiversity, carbon and water. Nat. Ecol. Evol.

37. Zhu, L., Hughes, A.C., Zhao, X.-Q., Zhou, L.-J., Ma, K.-P., Shen, X.-L., Li, S., Liu, M.-Z.,

Xu, W.-B., and Watson, J.E.M. (2022). Regional scalable priorities for national biodiversity

and carbon conservation planning in Asia. Sci. Adv. *7*, eabe4261.

38. Naidoo, R., Balmford, A., Costanza, R., Fisher, B., Green, R.E., Lehner, B., Malcolm, T.R.,

and Ricketts, T.H. (2008). Global mapping of ecosystem services and conservation priorities.

Proc. Natl. Acad. Sci. *105*, 9495–9500.

39. Cimatti, M., Brooks, T.M., and Di Marco, M. (2021). Identifying science-policy consensus

regions of high biodiversity value and institutional recognition. Glob. Ecol. Conserv. *32*,

e01938.

40. Peter, S., Le Provost, G., Mehring, M., Müller, T., and Manning, P. (2021). Cultural

worldviews consistently explain bundles of ecosystem service prioritisation across rural

Germany. People Nat. *n/a*.

41. O'Connor, C., Marvier, M., and Kareiva, P. (2003). Biological vs. social, economic and

political priority-setting in conservation. Ecol. Lett. *6*, 706–711.

42. Mace, G.M. (2014). Whose conservation? Science. *345*, 1558 LP – 1560.

43.   Di Marco, M., Cardillo, M., Possingham, H.P., Wilson, K.A., Blomberg, S.P., Boitani, L., and

Rondinini, C. (2012). A novel approach for global mammal extinction risk reduction. Conserv.

Lett. *5*, 134–141.

44.   Riggio, J., Baillie, J.E.M., Brumby, S., Ellis, E., Kennedy, C.M., Oakleaf, J.R., Tait, A., Tepe,

T., Theobald, D.M., Venter, O., et al. (2020). Global human influence maps reveal clear

opportunities in conserving Earth's remaining intact terrestrial ecosystems. Glob. Chang. Biol.

26, 4344–4356.

45.   Freudenberger, L., Hobson, P., Schluck, M., Kreft, S., Vohland, K., Sommer, H., Reichle, S.,

Nowicki, C., Barthlott, W., and Ibisch, P.L. (2013). Nature conservation: priority-setting needs

a global change. Biodivers. Conserv. *22*, 1255–1281.

46.   Girardello, M., Santangeli, A., Mori, E., Chapman, A., Fattorini, S., Naidoo, R., Bertolino, S.,

and Svenning, J.-C. (2019). Global synergies and trade-offs between multiple dimensions of

biodiversity and ecosystem services. Sci. Rep. *9*, 5636.

47.   Rosauer, D.A.N., Laffan, S.W., Crisp, M.D., Donnellan, S.C., and Cook, L.G. (2009).

Phylogenetic endemism: a new approach for identifying geographical concentrations of

evolutionary history. Mol. Ecol. *18*, 4061–4072.

48.   Voskamp, A., Butchart, S.H.M., Baker, D.J., Wilsey, C.B., and Willis, S.G. (2021). Site-Based

Conservation of Terrestrial Bird Species in the Caribbean and Central and South America

Under Climate Change   . Front. Ecol. Evol.   *9*, 234.

49.   IUCN (2016). The IUCN Red List of Threatened Species. See: https://www.

iucnredlist.org/resources/spatial-data-download [accessed 15.05.17].

50.   Birdlife International and NatureServe (2015). Bird species distribution maps of the world,

Version 5.0. (Birdlife International).

51.   Roll, U., Feldman, A., Novosolov, M., Allison, A., Bauer, A.M., Bernard, R., Böhm, M.,

Castro-Herrera, F., Chirio, L., Collen, B., et al. (2017). The global distribution of tetrapods

reveals a need for targeted reptile conservation. Nat. Ecol. Evol. *1*, 1677–1682.

52.   Crisp, M.D., Laffan, S., Linder, H.P., and Monro, A. (2001). Endemism in the Australian flora.

J. Biogeogr. *28*, 183–198.

53.   Lovejoy, T.E. (2016). Conservation Biology: The Importance of Wilderness. Curr. Biol. *26*,

R1235–R1237.

54.   Betts, M.G., Wolf, C., Ripple, W.J., Phalan, B., Millers, K.A., Duarte, A., Butchart, S.H.M.,

and Levi, T. (2017). Global forest loss disproportionately erodes biodiversity in intact

landscapes. Nature *547*, 441–444.

55.   Watson, J.E.M., Evans, T., Venter, O., Williams, B., Tulloch, A., Stewart, C., Thompson, I.,

Ray, J.C., Murray, K., Salazar, A., et al. (2018). The exceptional value of intact forest

ecosystems. Nat. Ecol. Evol. *2*, 599–610.

56.   Bauer, S., and Hoye, B.J. (2014). Migratory Animals Couple Biodiversity and Ecosystem

Functioning Worldwide. Science (80-.). *344*, 1242552.

57.   Ripple, W.J., Newsome, T.M., Wolf, C., Dirzo, R., Everatt, K.T., Galetti, M., Hayward, M.W.,

Kerley, G.I.H., Levi, T., Lindsey, P.A., et al. (2022). Collapse of the world's largest

herbivores. Sci. Adv. *1*, e1400103.

58.   Klein, C.J., Wilson, K.A., Watts, M., Stein, J., Carwardine, J., Mackey, B., and Possingham,

H.P. (2009). Spatial conservation prioritization inclusive of wilderness quality: A case study of

Australia's biodiversity. Biol. Conserv. *142*, 1282–1290.

59.   Salati, E., Dall'Olio, A., Matsui, E., and Gat, J.R. (1979). Recycling of water in the Amazon

Basin: An isotopic study. Water Resour. Res. *15*, 1250–1258.

60.   Schwartz, M.W. (1999). Choosing the Appropriate Scale of Reserves for Conservation. Annu.

Rev. Ecol. Syst. *30*, 83–108.

61.   Cantú-Salazar, L., and Gaston, K.J. (2010). Very Large Protected Areas and Their

Contribution to Terrestrial Biological Conservation. Bioscience *60*, 808–818.

62.   Legacy Landscapes Fund (2021). Charter Internationaler Naturerbe Fonds - Legacy

Landscapes Fund.

63.   BirdLife International (2019). Digital boundaries of Key Biodiversity Areas from the World

Database of Key Biodiversity Areas. Developed by the KBA Partnership.

http://www.keybiodiversityareas.org/ [accessed 23.05.2019].

64.   UNEP - WCMC, and IUCN (2020). Protected Planet: The World Database on Protected Areas

(WDPA).

65. Olson, D.M., Dinerstein, E., Wikramanayake, E.D., Burgess, N.D., Powell, G.V.N.,

Underwood, E.C., D'Amico, J.A., Itoua, I., Strand, H.E., Morrison, J.C., et al. (2001).

Terrestrial ecoregions of the world: A new map of life on earth. Bioscience *51*, 933–938.

66. Sala, E., Mayorga, J., Bradley, D., Cabral, R.B., Atwood, T.B., Auber, A., Cheung, W.,

Costello, C., Ferretti, F., Friedlander, A.M., et al. (2021). Protecting the global ocean for

biodiversity, food and climate. Nature *592*, 397–402.

67. BMZ Division 102, and KfW Centre of Competence for Infrastructure Water Natural

Resources (2020). The Legacy Landscapes Fund Safeguarding outstanding biodiversity for

humanity - the next level of conservation.

68. Mittermeier, R.A., Mittermeier, C.G., Brooks, T.M., Pilgrim, J.D., Konstant, W.R., da

Fonseca, G.A.B., and Kormos, C. (2003). Wilderness and biodiversity conservation. Proc.

Natl. Acad. Sci. *100*, 10309–10313.

69. Venter, O., Fuller, R.A., Segan, D.B., Carwardine, J., Brooks, T., Butchart, S.H.M., Di Marco,

M., Iwamura, T., Joseph, L., O'Grady, D., et al. (2014). Targeting Global Protected Area

Expansion for Imperiled Biodiversity. PLOS Biol. *12*, e1001891.

70. BirdLife International (2018). IUCN Red List for birds. *2018*. http://www.birdlife.org.

71. Crooks, K.R., and Sanjayan, M. (2006). Connectivity conservation: maintaining connections

for nature K. R. Crooks and M. Sanjayan, eds. (Cambridge University Press).

72. Bélisle, M. (2005). Measuring landscape connectivity: The challenge of behavioural landscape

ecology. Ecology *86*, 1988–1995.

73. Santini, L., Saura, S., and Rondinini, C. (2016). Connectivity of the global network of

protected areas. Divers. Distrib. *22*, 199–211.

74. Brennan, A., Naidoo, R., Greenstreet, L., Mehrabi, Z., Ramankutty, N., and Kremen, C.

(2022). Functional connectivity of the world's protected areas. Science. *376*, 1101–1104.

75. Secretariat of the United Nations Convention on Biological Diversity (2021). First draft of the

Post-2020 Global Biodiversity Framework.

https://www.cbd.int/doc/c/abb5/591f/2e46096d3f0330b08ce87a45/wg2020-03-03-en.pdf.

76. Gurney, G.G., Darling, E.S., Ahmadia, G.N., Agostini, V.N., Ban, N.C., Blythe, J., Claudet, J.,

Epstein, G., Estradivari, Himes-Cornell, A., et al. (2021). Biodiversity needs every tool in the

box: use OECMs. Nature *595*, 646–649.

77. Perino, A., Pereira, H.M., Felipe-Lucia, M., Kim, H., Kühl, H.S., Marselle, M.R., Meya, J.N.,

Meyer, C., Navarro, L.M., van Klink, R., et al. (2022). Biodiversity post-2020: Closing the gap

between global targets and national-level implementation. Conserv. Lett. *15*, e12848.

78. Hedges, S.B., Marin, J., Suleski, M., Paymer, M., and Kumar, S. (2015). Tree of life reveals

clock-like speciation and diversification. Mol. Biol. Evol. *32*, 835–845.

79. Newbold, T., Hudson, L.N., Arnell, A.P., Contu, S., De Palma, A., Ferrier, S., Hill, S.L.L.,

Hoskins, A.J., Lysenko, I., Phillips, H.R.P., et al. (2016). Has land use pushed terrestrial

biodiversity beyond the planetary boundary? A global assessment. Science. *353*, 288 LP – 291.

80. Venter, O., Sanderson, E.W., Magrach, A., Allan, J.R., Beher, J., Jones, K.R., Possingham,

H.P., Laurance, W.F., Wood, P., Fekete, B.M., et al. (2016). Sixteen years of change in the

global terrestrial human footprint and implications for biodiversity conservation. Nat.

Commun. *7*, 12558.

81. Niamir, A., Salati, S., Gómez-Giménez, M., Werner, C., and Hickler, T. (2020). The fraction

of land cover classes; derived variables from ESA CCI Land Cover time-series (1992 - 2018)

(1.0) [Data set]. Zenodo.

82. QGIS Development Team (2021). QGIS Geographic Information System.

83. Hof, C., Voskamp, A., Biber, M.F., Böhning-Gaese, K., Engelhardt, E.K., Niamir, A., Willis,

S.G., and Hickler, T. (2018). Bioenergy cropland expansion may offset positive effects of

climate change mitigation for global vertebrate diversity. Proc. Natl. Acad. Sci. *115*, 13294.

84. Smith, B., Wårlind, D., Arneth, A., Hickler, T., Leadley, P., Siltberg, J., and Zaehle, S. (2014).

Implications of incorporating N cycling and N limitations on primary production in an

individual-based dynamic vegetation model. Biogeosciences *11*, 2027–2054.

85. Popp, A., Calvin, K., Fujimori, S., Havlik, P., Humpenöder, F., Stehfest, E., Bodirsky, B.L.,

Dietrich, J.P., Doelmann, J.C., Gusti, M., et al. (2017). Land-use futures in the shared socio-

economic pathways. Glob. Environ. Chang. *42*, 331–345.

86.   Popp, A., Humpenöder, F., Weindl, I., Bodirsky, B.L., Bonsch, M., Lotze-Campen, H., Müller,

783   C., Biewald, A., Rolinski, S., Stevanovic, M., et al. (2014). Land-use protection for climate

784   change mitigation. Nat. Clim. Chang. *4*, 1095–1098.

785   87.   Stevanović, M., Popp, A., Lotze-Campen, H., Dietrich, J.P., Müller, C., Bonsch, M., Schmitz,

786   C., Bodirsky, B.L., Humpenöder, F., and Weindl, I. (2016). The impact of high-end climate

787   change on agricultural welfare. Sci. Adv. *2*.

788   88.   Frieler, K., Lange, S., Piontek, F., Reyer, C.P.O., Schewe, J., Warszawski, L., Zhao, F., Chini,

789   L., Denvil, S., Emanuel, K., et al. (2017). Assessing the impacts of 1.5°C global warming -

790   simulation protocol of the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP2b).

791   Geosci. Model Dev. *10*.

792   89.   Chang, W., Cheng, J., Allaire, J., Xie, Y., and McPherson, J. (2020). shiny: Web Application

793   Framework for R. R package.

794