Fibrosis resolution involves more than collagen; new potential players of fibrosis recovery in

mouse liver

Toshifumi Sato,[1] Kimberly Z. Head,[5] Jiang Li,[1] Christine E. Dolin,[5] Daniel Wilkey,[6,7] Nolan

Skirtich,[2] Dylan D. McCreary,[2] Juliane I Beier,[1,3,4] Ryan M. McEnaney,[2] Michael L Merchant[6,7]

and Gavin E Arteel[1,4]

[1]Department of Medicine, Division of Gastroenterology, Hepatology and Nutrition, [2]Department

of Surgery, [3]Department of Environmental and Occupational Health, [4]Pittsburgh Liver Research

Center, University of Pittsburgh, Pittsburgh, PA 15213. [5]Department of Pharmacology and

Toxicology, [6]Department of Medicine, Division of Nephrology and Hypertension and [7]University

of Louisville Alcohol Research Center, University of Louisville, Louisville, KY 40292

Running title: Resolution of fibrosis: beyond collagen.

Key Words: MMP-12, Elastin, fibrosis

1   Send all correspondence to: Gavin E. Arteel, PhD, FAASLD

2   Thomas E. Starzl Biomedical Science Tower

3   West 1143

4   200 Lothrop Street

5   Pittsburgh, PA 15213

6   Phone: +1-412-648-4187

7   Email: gearteel@pitt.edu

10   Abbreviations: CCl_4, carbon tetrachloride; ECM, extracellular matrix; ELISA, enzyme-linked

11   immunosorbent assay; GO, gene ontology; LC-MS/MS, liquid Chromatography with tandem

12   mass spectrometry; MMP, matrix metalloprotease; MPM, multiphoton microscopy; rtPCR,

13   reverse-transcriptase polymerase chain reaction; SDS, sodium dodecyl sulfate.

15   Acknowledgments:

Abstract

Although most work has focused on resolution of collagen ECM, fibrosis resolution involves

changes to several ECM proteins. The purpose of the current study was two-fold: 1) to examine

the role of MMP12 and elastin; and 2) to investigate the changes in degraded proteins in plasma

(i.e., the "degradome") in a preclinical model of fibrosis resolution. Fibrosis was induced by 4

weeks carbon tetrachloride (CCl_4) exposure, and recovery was monitored for an additional 4

weeks. Some mice were treated with daily MMP12 inhibitor (MMP408) during the resolution

phase. Liver injury and fibrosis was monitored by clinical chemistry, histology and gene

expression. The release of degraded ECM peptides in the plasma was analyzed using by 1D-

LC-MS/MS, coupled with PEAKS Studio (v10) peptide identification. Hepatic fibrosis and liver

injury rapidly resolved in this mouse model. However, some collagen fibrils were still present

28d after cessation of CCl_4. Despite this persistent collagen presence, expression of canonical

markers of fibrosis were also normalized. The inhibition of MMP12 dramatically delayed fibrosis

resolution under these conditions. LC-MS/MS analysis identified that several proteins were

being degraded even at late stages of fibrosis resolution. Calpains 1/2 were identified as

potential new proteases involved in fibrosis resolution. CONCLUSION. The results of this study

indicate that remodeling of the liver during recovery from fibrosis is a complex and highly

coordinated process that extends well beyond the degradation of the collagenous scar. These

results also indicate that analysis of the plasma degradome may yield new insight into the

mechanisms of fibrosis recovery, and by extension, new "theragnostic" targets. Lastly, a novel

potential role for calpain activation in the degradation and turnover of proteins was identified.

Regardless of etiology, hepatic fibrosis is an almost universal endpoint of chronic liver injury,

when coupled with impaired/incomplete recovery from said injury (1). The clinical sequelae of

severe fibrosis/cirrhosis significantly and negatively impact patient survival and quality of life.

Given these clinical implications, the mechanisms of hepatic fibrosis initiation and progression

have been studied for decades with the goal of identifying better means to prevent and treat this

disease (2). Although much progress has been made in our understanding, the translational

value of these studies has been hampered by the fact that clinical diagnosis of liver disease is

often late-stage, where mechanism-based therapies appear to be less effective.

It has been known that for several years that hepatic fibrosis may resolve, if the underlying

causal factor is successfully removed (3). This phenomenon was consistently observed in

animal models of fibrosis, but until recently only in rare cases in humans (4). The recent

advance of highly effective disease-modifying agents (e.g., direct acting antivirals for hepatitis C

viral infection) have dramatically increased the opportunity for fibrosis resolution to occur (5).

However, not all patients for achieve fibrosis resolution when the underlying disease has been

effectively treated (6). Moreover, in contrast to the development of fibrogenesis, the

mechanisms by which fibrosis resolution is mediated are poorly understood and may not share

complete mechanistic overlap with fibrosis progression (4). To address both this clinical need

and the gap in our understanding, more investigations in this area need to be performed.

The hepatic extracellular matrix (ECM) consists of a diverse range of components that

work bi-directionally with surrounding cells to create a dynamic and responsive

microenvironment that regulates cell and tissue homeostasis. Under normal conditions, the

ECM assists in maintaining organ homeostasis and appropriate responses to injury/stress by

dynamically responding to these insults. However, failure to properly regulate these responses

can lead to qualitative and/or quantitative ECM changes that are maladaptive (7). Hepatic

fibrosis is a canonical example of ECM dyshomeostasis, leading to accumulation of fibrillary

ECM, such as collagen, which is considered the pathologic hallmark of fibrosis. Although

hepatic fibrosis is considered almost synonymous with collagen accumulation (8), the qualitative

and quantitative alterations to the hepatic ECM during fibrosis are much more diverse (9).

Indeed, hepatic fibrosis results in qualitative and quantitative changes to almost all components

of the hepatic ECM (9). This understanding has let previous work by others to investigate the

roles of non-collagen ECM components, as well as enzymes that metabolize them (e.g., matrix

metalloproteinases; MMPs) in the initiation and development of experimental and clinical fibrosis

(10, 11). In contrast, previous studies of fibrosis resolution have focused almost exclusively on

secretion and turnover of collagens and on collagen microstructure (12). The purpose of the

current study was to investigate turnover of hepatic ECM during fibrosis resolution in mice.

Experimental Procedures

Please see Supplemental Materials for additional details on Experimental Procedures.

Animals and Treatments. 8 weeks old, male C57Bl/6J mice purchased from Jackson Laboratory (Bar Harbor, ME) were housed in a pathogen-free barrier facility accredited by the Association for Assessment and Accreditation of Laboratory Animal Care, and procedures were approved by the local Institutional Animal Care and Use Committee. Animals were allowed standard laboratory chow and water *ad libitum*. Mice were administered CCl_4 (1 ml/kg i.p.; diluted 1:4 in olive oil; Sigma-Aldrich, St. Louis, MO) or vehicle 2×/wk for 4 wks; animals were sacrificed 1, 3, 7, 14 and 28 days after final CCl_4 injection (Figure 1A). Animals were administered an MMP-12 inhibitor (MMP-408; 10 mg/kg, i.g.; Merck Millipore, Burlington, MA) or vehicle (saline) daily starting after the last injection of CCl_4. At the time of sacrifice, animals were anesthetized with ketamine/xylazine (100/15 mg/kg, i.p.). Blood was collected from the vena cava just prior to sacrifice by exsanguination and citrated plasma was stored at -80°C for further analysis. Portions of liver tissue were frozen immediately in liquid nitrogen, while others were fixed in 10% neutral buffered formalin or embedded in frozen specimen medium (Tissue-Tek OCT compound, Sakura Finetek, Torrance, CA) for subsequent sectioning and mounting on microscope slides.

Biochemical assays and histology. Plasma levels of Aspartate transaminase (AST) and alanine amino transferase (ALT) were determined spectrophotometrically using standard kits (Thermo Fisher Scientific, Waltham, MA), as described previously (13). Formalin-fixed, paraffin embedded sections were cut at 5 µm and mounted on glass slides. Deparaffinized sections stained with hematoxylin and eosin and pathology was assessed in a blinded manner. ECM accumulation in liver sections was determined by staining with Sirius red/fast green and were

visualized via brightfield and polarized light (14). Bright field images of Sirius red staining was

quantitated by image analysis, as described previously (15). Specifically, a Molecular Devices

(Sunnyvale, CA) Image-1/AT image acquisition and analysis system incorporating an Axioskop

50 microscope (Carl Zeiss Inc., Thornwood, NY) was used to capture and analyze five

nonoverlapping fields per section at 100× final magnification. Image analysis was performed

using modifications of techniques described previously. Detection thresholds were set for the

red color based on an intensely labeled point and a default color threshold. The degree of

labeling in each section was determined from the area within the color range divided by the total

area.

Hepatic calpain activity. Calpain activity was measured in liver lysate, using N-succinyl-

Leucine-Leucine-Valine-Tyrosine-7-amino-4-methylcoumarin (Suc-Leu-Leu-Val-Tyr-AMC;

Supplemental Table 3) was used as the calpain substrate, as described previously (16, 17).

Forty μl of liver lysate samples (0.25 mg protein/ml in imidazole buffer containing 63 mM

imidazole, 10 mM mercaptoethanol, 1 mM EDTA, 10 mM EGTA, pH 7.3) were incubated at 37

for 1hr with 160 μl of imidazole buffer containing 50 μM Suc-Leu-Leu-Val-Tyr-AMC (dissolved in

DMSO). The formation of the cleavage product, 7-amino-4-methylcoumarin (AMC), was

measured by fluorometrically (Spectra Max M5, Molecular Devices Corpolation) at 360 nm

excitation and 430 nm emission wavelengths. Standard curves were generated with free AMC

for each experiment and results were normalized to total protein.

Determination of MMP-2 and 9 by zymography. Zymography was performed as

described previously (18). Total hepatic protein was extracted using a lysis buffer consisting of

10 mM cacodylic acid, pH 5.0, containing 150 mM NaCl, 1M ZnCl, 15 mM $CaCl_2$, 1.5 mM NaN_3, and 0.01% Triton X-100. Lysates were then diluted in 2× sample buffer (125 mM Tris HCl, pH 6.8, with 4% SDS, 20% glycerol) without boiling or reduction. The diluted samples were subjected to 0.1% gelatin–10 % SDS polyacrylamide gel electrophoresis (Novex, Carlsbad, CA). The gels were washed with 2.5 % Triton X-100 after electrophoresis and then incubated for 48 hours in the reaction buffer (50 mM Tris, 0.1 M glycine, and 0.1 M NaCl, pH 8.0). The gel was stained with Coomassie brilliant blue R-250 (0.1 % Coomassie Brilliant Blue R250, 50% methanol, and 10% acidic acid) for 1hr, destained with 45% methanol and 10% acetic acid for 30 minutes. The results were visualized and densitometric analysis was performed using an iBright imager and onboard software from Invitrogen (Waltham, MA).

Immunoblots. Liver samples were homogenized in RIPA buffer (20 mM Tris/Cl, pH 7.5, 150 mM NaCl, 1 mM EDTA, 1 mM EGTA, 1% [wt/vol] Triton X-100), containing protease and phosphatase inhibitor cocktails (Sigma). Samples were diluted in 2× sample buffer (see above) and loaded onto 10% SDS-polyacrylamide gels (Invitrogen, Waltham, MA), followed by electrophoresis and Western blotting on to PVDF membranes (Hybond P, GE Healthcare Bio Sciences, Pittsburgh, Pennsylvania). Primary antibodies against proteins of interest were compared to GAPDH as a housekeeping gene (see Supplemental Table 3). The results were visualized and densitometric analysis was performed using an iBright imager and onboard software from Invitrogen (Waltham, MA).

RNA Isolation and Real-Time RT-PCR RNA extraction and real-time RT-PCR were performed as described previously (19). RNA was extracted immediately following sacrifice from fresh liver samples using RNA Stat60 (Tel-Test, Ambion, Austin, TX) and chloroform. RNA concentrations were determined spectrophotometrically and 1µg of total RNA was reverse

transcribed using the QuantiTect Reverse Transcription Kit (Qiagen,Valencia, CA). Real-time

RT-PCR was performed using a StepOne real time PCR system (Thermo Fisher Scientific,

Grand Island, NY) using the Taqman Universal PCR Master Mix (Life Technologies, Carlsbad,

CA). Primers and probes were ordered as commercially available kits (Thermo Fisher

Scientific, Grand Island, NY; see Supplemental Table 1). The comparative C_T method was used

to determine fold differences between the target genes and an endogenous reference gene

(18S). Results were reported as copy number ($2^{-\Delta CT}$) or fold naïve control (FC; $2^{-\Delta\Delta CT}$).

Immunofluorescence. For immunofluorescent detection of elastin, sections of liver were

blocked with PBS containing 10 % goat serum for 30 min, followed by incubation overnight at

4 °C with Elastin (Abcam 21607 1:50) antibodies in blocking solution. Sections were washed

with PBS and incubated for 3 hrs with a secondary antibody conjugated to Alexa 488

(Invitrogen, Carlsbad, CA). Sections were washed with PBS and mounted with mounting

medium containing 1.5 µg/ml DAPI (VECTASHIELD w/DAPI; Vectastain, Torrance, CA).

Multiphoton microscopy imaging. Elastin and collagen content in the samples were

assessed using multiphoton microscopy (MPM), as described previously (20). MPM were

performed on frozen sections with a nonlinear optical microscope based on a confocal imaging

system (FV1000MPE, Olympus, Tokyo, Japan; purchased with grant 1S10RR025676-01

awarded to Dr Simon C. Watkins) using an optical microscope and an external tunable mode-

locked Ti:sapphire LASER (Coherent, Palo Alto, CA) set to 790-800 nm, 5.4% intensity. LASER

emission and signal detection were through a 25x, XL Plan N 1.05 N.A. water immersion

objective lens (Olympus). Detection was synchronous across three channels. Nuclei stained by

propidium iodide (Invitrogen) were detected on the RXD4 channel (663-737 nm emission filter).

Fibrillar collagen second-harmonic generation was detected on the RXD1 channel (350-450 nm emission filter). Elastin-associated fibrillar green autofluorescence was detected on the RXD2 channel (500-550 nm emission filter). A 4x line Kalman integration filter was applied to reduce stochastic noise. The laser was routed to an acousto-optic modulator (AOM) for power attenuation, then through a dichroic mirror (490 nm) and an objective to the tissue sample. Signals from second harmonic generation (SHG) were collected by the same objective lens and recorded by a photomultiplier tube (PMT).

Preparation of acid hydrolysates of liver samples and Quantitation of Desmosine Crosslinks. Liver samples were homogenized in RIPA buffer (see above) and autoclaved (3 h) twice, at 20 lb/in.2 (1.36 atm) to remove collagen and other proteins. The residue was treated with 0.1 M NaOH for 30 min at 100 °C, washed until neutral, and lyophilized. The isolated elastin was flushed with nitrogen and hydrolyzed in 6 N HCl under vacuum at 105°C for 24-36 h. The hydrolysates were evaporated to dryness and the residue was dissolved in 0.5-1.0 ml water. Appropriate aliquots were then added to 0.1 M sodium phosphate buffer, pH 7.4, containing phenol red and neutralized with 2 M NaOH. Desmosine content was determined using a commercially available ELISA kit (Cusabio, Houston, TX) and normalized to starting tissue wet weight.

Plasma peptide purification and high resolution LC-MS analysis. 100 µL PBS containing 0.1X iRT standards (Biognosis Inc, Beverly, MA) was added to 100 µL plasma. 200 µL 20% w/v TCA was mixed into each sample on ice. Samples were then be incubated for 1 hour at 4°C. The precipitate was pelleted by centrifugation at 16,000 × G for 10 min at 4°C; pellets were discarded. The centrifugation was then repeated with the supernatant samples, followed by discard of the second pellets. Samples were then desalted and concentrated using solid phase

extraction (Waters Oasis HLB µElution 30 µm plate, part no. 186001828BA) as described (21,

22). Solid phase extraction eluates were dried in a SpeedVac, and dried peptides were

resuspended in 25 µL 2% v/v acetonitrile/ 0.1% v/v formic acid. Peptide concentrations were

measured by absorbance (NanoDrop2000) at 205 nm and concentrations estimated using a

plasma tryptic digest standard curve. Peptide samples (0.2 ug) were separated using an 12cm,

360µm OD x 100µm ID fused silica tip packed with Aeris Peptide 3.6µm XB-C18 100Å material

(Phenomenex, Torrance, CA, USA). Peptides were eluted under a 90min 2%-44% acetonitrile

gradient and transferred by Nanospray Flex nanoelectrospray source (ThermoFisher) into an

Orbitrap Elite - ETD mass spectrometer (ThermoFisher) using an Nth Order Double Play

(Xcalibur v2.2, ThermoFisher) with FTMS MS1 scans (240,000 resolution) collected from 300-

2000m/z and ITMS MS2 scans collected on up to twenty peaks having a minimum signal

threshold of 5,000 counts from the MS1 scan event.

<u>Peptidomic Data Analysis</u>. RAW files were searched in Peaks Studio X (Bioinformatics

Solutions Inc., Waterloo, ON, Canada) using the Denovo, PeaksDB, PeaksPTM, and Label Free

Q algorithms and the 2/12/2019 version of the UniprotKB reviewed reference proteome

canonical and isoform Homo sapiens sequences (Proteome ID UP000005640). Search

parameters included: variable methionine, proline and lysine oxidation (+15.9949 Da), no

enzyme specified, 15 ppm precursor error for MS1 Orbitrap FTMS data, 0.5 Da error for MS2

data sets, and common PTMs in the PeaksPTM algorithm. The peptide, feature, and protein-

peptide csv files were exported from the Label Free Q result for statistical tests in Microsoft

Excel 2016 and R v3.5.0. High confidence peptide assignments (PeaksX criteria –logP scores

with p-value <0.05) were exported into comma separated values files (.csv) for upload and

analysis by the Proteasix (http://www.proteasix.org/) algorithm using the 'Peptide-centric'

prediction tool based on curated, known and observed cleavage events enabling assignment of

1 protease identities from the MEROPS database as described previously (23). Protein-protein

2 interaction network analysis of regulated proteomic data sets (q-value <0.05) was performed

3 using Search Tool for the Retrieval of Interacting Genes/Proteins, STRING v11 (24, 25), with

4 the highest confidence score (0.900).

Results

Recovery from liver injury and fibrosis. All animals survived exposure to CCl$_4$ and no

morbidity or mortality was reported during the recovery phase. As expected (26), 4 wks CCl$_4$

administration caused severe liver injury and hepatic fibrosis in mouse liver as assessed by

histology (Figure 1B) and by transaminase release (Figure 2A). Four weeks of CCl$_4$

administration increased hepatic expression of key fibrogenic genes (*Col1a1*, *Acta2*, *p4htm*,

Lox1 and *Lox2*) significantly compared with vehicle controls (Figure 2A). Hepatic fibrosis and

liver injury rapidly resolved after cessation of CCl$_4$ exposure (Figure 1B), with only thin septa of

collagen staining observable after 4 wks recovery (Figure 1B). Likewise, transaminase values

and expression of key fibrogenic genes all returned to baseline values after 4 weeks of recovery

(Figure 2A).

Effect of CCl$_4$ on expression and activity of MMPs. Even in cases where there is a net

increase in ECM in the liver (e.g., fibrosis) overall turnover is also increased (27). This turnover

is known to be mediated, at least in part, by matrix metalloproteinases (MMPs). 4 wk of CCl$_4$

administration increased the expression (Figure 3A) and activity (Figures 2B and 2C) of the

collagenases, MMP2 and MMP9. The mRNA expression of inhibitors of these MMPs (i.e.,

Timp1, *Timp2* and *Timp3*) were also induced with a similar profile (Figure 2C). As was

observed with expression of key indices of fibrogenesis (Figure 2A; see above), the increase in

expression/activity of these factors largely returned to baseline levels after 28 days of recovery.

In addition to collagen deposition, fibrosis involves quantitative and qualitative changes

to a myriad of ECM proteins and MMPs (9, 28). The expression of all hepatic MMPs was

therefore determined (Figure 3A). Analogous to findings with *Mmp2* and *Mmp9,* all hepatic

MMPs were induced by experimental fibrosis (Figure 3A). The expression pattern of the

majority of these MMPs was also similar to *Mmp2* and *Mmp9* and returned to baseline 28 days

after the cessation of CCl$_4$ (Figure 3A). The one contrast was the expression of *Mmp12*, which

was the most robustly induced MMP, both by fold change (1000-fold over control), as well as by

copy number (Figure 3A). Unlike other MMPs, expression levels of *Mmp12* were still elevated

~10-fold over baseline 28 days after cessation of CCl$_4$ (Figure 3A).

Effect of CCl$_4$ on elastin accumulation. Elastin is a key substrate for MMP12 that

increases during CCl$_4$-induced fibrosis (10). Expression and protein levels of elastin were

therefore determined (Figures 3B and C). Although the mRNA expression of elastin (*Eln*) was

only moderately (<4 f-fold) induced by 4 weeks of CCl$_4$ administration and returned to baseline

after 28 days of recovery, protein levels increased ~15-fold compared with naïve control mice

(Figures 3B and 3C). Moreover, elastin ECM was still significantly (~5 fold) elevated over naïve

control values even after 28 days of recovery. As has been observed previously (10), elastin

accumulation appeared to correlate with areas of fibrotic changes (Figure 3B). Elastin

accumulation appears to center on areas of fibrogenesis and collagen accumulation (see Figure

3B). To investigate this further, the colocalization of elastin and collagen were determined using

multi-photon microscopy (Figure 3B), as it allows observation of the elastin and collagen fiber

networks simultaneously (20). Elastin (green fluorescence) accumulation was highest on the

first day of recovery and was closely associated with collagen (blue fluorescence) signals,

consistent with the assessment of elastin immunofluorescence (Figures 3B).

MMP12 inhibition delays fibrosis resolution. Previous work with knockout mice indicated

that MMP12-mediated elastin degradation plays a protective role in the development of CCl$_4$-

induced fibrogenesis (10); however, the role of MMP12/elastin on resolution of fibrosis was

unknown. The effect of the MMP12 inhibition on resolution of fibrosis was therefore determined

(Figure 4). CCl$_4$-exposed mice were administered the MMP12 inhibitor, MMP-408, daily after

cessation of toxicant exposure (see Materials and Methods). MMP-408 did not significantly

affect recovery of transaminases after CCl$_4$ exposure (Figure 4A). MMP-408 also did not

significantly impact the expression of fibrogenic genes (e.g., *Col1A1*) during the recovery phase

(Figure 4A). On the other hand, MMP408 impacted the resolution of collagenous fibrosis, as

determined by Sirius red staining (Figures 4B and C). Image-analysis and quantification of

Sirius red staining indicated that the amount of collagen present in the liver was double 1 wk

after cessation of CCl$_4$ administration compared to the mice administered the vehicle for

MMP408 (Figure 4C). Although this effect of MMP408 was not present after 28 days of

recovery, the fibrous septa in the MMP-408 group were far less defined than in the vehicle

treated (Figure 4B and inset).

The elastin signal in the multiphoton microscopy analysis shows both immature and

mature elastin [Figure 3B (20)]. Therefore, it reflects the total amount of elastin accumulation in

the liver. In contrast, the crosslinking of mature insoluble elastin is mediated via post-

translational incorporation of the amino acid desmosine. Desmosine is unique to mature elastin

in mammalian species and can therefore be employed as a surrogate marker of mature elastin

(29). The amount of desmosine present in the liver tissues after CCl$_4$ was therefore determined

by ELISA (Figure 4C), as described in Materials and Methods. Although the total amount of

elastin detected in the liver increased by 4 wks of CCl$_4$ administration (Figures 3), the

desmosine content in the liver actually decreased to ~50% of levels in naïve mice (Figure 4C).

These data suggest that the majority of the new elastin detected during fibrosis is immature

uncrosslinked elastin. During the resolution phase desmosine content in livers from mice

administered vehicle increased, with values not significantly different that naïve control mice 28

days after cessation of CCl_4. In contrast, administration of MMP408 completely attenuated the

increase in this index of mature elastin (Figure 4C).

<u>The degradome changes dramatically during fibrosis resolution</u>. Resolution of fibrosis

requires coordinated remodeling of the liver. The net effect of fibrosis and resolution on hepatic

protein degradation has not been previously determined. The influence of CCl_4 and 4 wks

recovery on the plasma peptide "degradome" was therefore determined (Figure 5). Across the

groups, 694 peptides were identified, corresponding to 181 distinct parent proteins. The parent

proteins that contributed to the pattern of significantly changed peptides were analyzed by

StringDB, followed by K-means cluster analysis (24, 25). This database queries physical and

functional protein-protein associations that integrates both experimental and predicted

associations. The output creates both graphical and categorical enrichment patterns in the

queried dataset. Cluster analysis yielded 7 enriched clusters in the total degradome, with a

strong enrichment of proteins associated with Gene Ontology (GO) terms for "high-density

lipoprotein particle" (GO:0034464; Cluster 1), "collagen trimer" (GO:0005581; cluster 6) and

"nucleosome" (GO:0000786; Cluster 7) (Figure 5A; Supplemental Table 4).

The degradome from 1 and 28 D of recovery were compared in more detail to represent

the peak of injury (1 D) to near resolution of injury (28 D; Figures 1 and 2). A total of 255

peptides were significantly changed at least >2 fold (compared to naïve control) at these

timepoints (Figure 5B, C and D). Although there was considerable overlap between these 2

timepoints (Figure 5D), the amount of peptides altered at the 28 D timepoint was ~2.5-fold more

compared to the 1 D timepoint. A total of 144 peptides, corresponding to 82 distinct proteins,

were significantly increased at 1 D after CCl_4 and/or 28 D after CCl_4 (Figure 5C, Supplemental

Table 5). All peptides that significantly increased at 28 D after recovery tended to be at

relatively low levels 3 and 7 days after cessation of CCl_4, then increasing at the 14 D timepoint

(Figure 5B).

Many proteases cleave substrates with high specificity with limited cleavage promiscuity.

Thus, the fragment sequence of degraded proteins can be informative on proteases that may

have generated this pattern. Proteasix is an open-source peptide-centric tool to predict in silico

the proteases involved in generating the observed peptide features (30). Figure 6A is a

frequency distribution figure comparing the predicted proteases that generated the significantly

increased peptides at the 1 and 28 D timepoint (vs. naïve control). The top predicted proteins,

calpains 1 and 2 were shared between the timepoints under these conditions (Figure 6A). The

expression of these splice variants (*Capn1* and *Capn2*) was determined via real-time rtPCR

(Figure 6B). The copy number of Capn2 was approximately an order of magnitude higher than

Capn1 in liver (3.1×10^{-5} vs. 4.3×10^{-6}, respectively) under basal conditions. The expression of

both isoform variants was increased by CCl_4 administration but returned to baseline by 28 D of

recovery (Figure 6B). In contrast, predicted enzyme activity of CAPN, both by cleavage of α-

fodrin and by enzymatic analysis (Figure 6B), indicated that CAPN activity was elevated at the

28 D timepoint. Several of the predicted CAPN1/2 substrates for these enzymes (Figure 6C)

were analogous to the key clusters identified by StringDB analysis for the degradome as a

whole (Figure 5A and Supplemental Table I), such as fibrinogens (cluster 1), collagens (cluster

6) and histones (cluster 7) and their corresponding GO terms.

Discussion

Direct-Acting Antiviral therapy has for all intents and purposes cured the hepatitis C virus (HCV) epidemic in immunocompetent adults. However, one of the most important issues is whether patients with significant HCV-induced fibrosis/cirrhosis will recover from this injury once a sustained virologic response is achieved (31). For example, improvement of liver stiffness in HCV cirrhosis after successful antiviral therapy is reportedly 30-65%, depending on the study referenced (32-36). Although promising at first glance, these data indicate that around half of the HCV-fibrosis/cirrhosis patients with successful viral clearance do not improve in their liver disease. Furthermore, another alarming concern is that the commonly used non-invasive transient elastographic analysis of liver stiffness tends to overpredict fibrosis/cirrhosis regression in HCV, compared to biopsy-based histologic assessment (31, 37). Indeed, despite achievement of viral clearance, histological assessment of patient livers often continues to show lingering inflammation, hepatocyte and ductular reaction and fibrosis (31, 37). It is therefore an unmet clinical need to accurately predict determining factors of fibrosis resolution. Although accumulation of collagen ECM is the *de facto* definition of fibrosis, the changes to the hepatic ECM that happen during both processes are much diverse than those that happen to collagen (9). Likewise, although it is understood that hepatic fibrosis resolution involves more than simple clearance of collagen ECM, the process and its mediators are incompletely understood, at best (38).

Previous studies have indicated that MMP12, most likely via mediating elastin ECM accumulation, is key for the development of hepatic fibrosis (10). Specifically, genetic or pharmacologic inhibition of the elastase, MMP12, exacerbated the development of experimental fibrosis. Furthermore, prevention of elastin crosslinking protected against experimental fibrosis (39). More recently, Chen et al. (40) demonstrated that elastin fibers degrade slowly during

fibrosis resolution, a finding recapitulated in the current study (Figure 3). The authors of that

study suggested that cross-linked elastin stabilizes the fibrotic ECM and may thereby protect it

from degradation (40). However, whether or not MMP12 activity mediates resolution of fibrotic

ECM was not clear. Here, it was demonstrated that the MMP12 inhibitor, MMP408, delayed

resolution of Sirius red-positive areas of the liver (Figure 4), which supports the hypothesis that

elastin degradation is key to overall fibrosis resolution. Although elastin protein levels increased

(Figure 3), the amount of mature crosslinked elastin (as determined by desmosine content;

Figure 4C) actually was decreased by CCl_4 exposure. Desmosine content did not return to

basal levels until after 28 D of recovery. These results suggest that the increase in elastin

content in the liver during fibrosis represents less mature (crosslinked) protein oligomers (Figure

7). This finding is similar in models of photoaging in which damaged skin accumulates

disorganized elastin fibrils (41). Interestingly, inhibition of MMP12 with MMP408 prevented the

recovery of desmosine content during fibrosis resolution, suggesting that MMP12 is critical not

only for the degradation of accumulated elastin, but also the reorganization to mature elastin

(Figure 7).

Hypothesis-driven approaches are key to elucidating mechanism. However, the nature

of the approach by definition requires prior knowledge. Discovery-based 'omic techniques

coupled with informatic analyses can yield new information and insight into disease progression

and regression. The peptidome is defined as the population of low molecular weight biologically

derived peptides (0.5-3 kDa), within the cells and biologic fluids. These compartments contain

peptides critical for normal organismal function. The study of the peptidome (i.e., 'peptidomics')

is a discipline related to proteomics, but with significant methodological and analytical

differences (42, 43). For example, as the original structure of the peptides is of interest,

samples are not trypsin digested prior to LC-MS/MS analysis, as is the case for bottom-up

proteomic approaches. This difference originally limited peptide identification, as the available

databases were based on trypsin-digested peptide fragments (44). However, the development

of peptidome-specific identification and analysis tools has addressed this concern (44, 45). In

addition to synthesized peptides, the peptidome also contains fragments of proteins degraded

by normal and/or abnormal processes (i.e., 'degradome'). These degradomic profiles are

conditionally unique to the metabolic/pathologic status of the organism. The latter subset of the

peptidome has generated key interest in some areas of human health as possible (surrogate)

biomarkers for disease. Here, LC-MS/MS analysis of the degradome profile in plasma during

recovery from experimental fibrosis was analyzed to investigate the general pattern of protein

turnover/remodeling during recovery, as well as to identify potential new key players.

When the degradome was analyzed, an interesting pattern resolved. Specifically,

although livers were almost histologically-normal after 4 weeks of recovery from CCl_4-induced

experimental fibrosis (Figure 1), there was a relative increase in the turnover of proteins

observed in the plasma compartment (Figure 5), as determined by analysis of the degradome.

This finding is in-line with previous observations by this group that hepatic remodeling in

response to injury is much more diverse than can be observed histologically (13). In addition to

the appearance of degraded ECM proteins canonically associated with fibrosis (e.g., collagens),

several other degraded ECM proteins were present, especially after 28 days of recovery. A

limitation of degradome analysis is that the original source of the parent protein cannot be

completely verified. However, the CCl_4 model is relatively specific for liver injury and the

majority of the significantly changed peptides were hepatic by Brenda Tissue Ontology analysis

(BTO:0000759; $p=2.2\times10^{-14}$).

One of the most robust families of degraded proteins that were enriched 28 days after

cessation of CCl_4 were associated with the coagulation cascade, especially the fibrin(ogens)

(Figure 5 and Supplemental Tables 4 and 5). Previous studies have investigated the potential

role of fibrin(ogens) in the development of hepatic injury and fibrosis, with somewhat conflicting

results (46-48). However, few if any studies have investigated the role of fibrin(ogen) ECM in

resolution of hepatic fibrosis. In contrast, the turnover of fibrin(ogen) ECM has been heavily

studied in other diseases of remodeling and recovery. For example, fibrin(ogen) ECM plays a

key role in sub-cutaneous wound-healing (49), and is the rationale for the delivery fibrin-

enriched products to promote wound healing (50). However, the clearance of the fibrin-

containing provisional matrix is also a key step in late stages of wound-healing (49). Since

hepatic fibrosis and subcutaneous wound-healing qualitatively and quantitatively share many

similarities, it is possible this increase in fibrin(ogen) fragments during the late stage of recovery

from hepatic fibrosis represents a similar clearance of fibrin ECM from the provisional matrix.

Future studies should specifically investigate this possibility, especially given the use of

anticoagulants in fibrotic liver disease and cirrhosis (51).

As mentioned in the Introduction, collagenous ECM accumulation has been the major

focus in the field of hepatic fibrosis (2). Likewise, proteases involved in resolution of

collagenous ECM (e.g., MMP2 and MMP9) have also received a sizeable amount of attention

(38). Once the technical limitations to identifying peptide fragments of degraded proteins

informatically was overcome (42, 43), the fragment sequence and putative cleavage site may

yield new information on proteases involved. The predicted protease that generated each

peptide was therefore determined by in silico analysis of the cleavage site using the online

open-source tool, Proteasix (www.proteasix.org) (30). Although several proteases with

established roles in fibrosis and/or resolution were identified (e.g., MMP2), several novel players

were predicted by the algorithm.

The most common predicted protease both 1 and 28 d after cessation of CCl_4 was

calpain1 (Capn1) (Figure 6). It should be noted that Capn1 and its isozyme Capn2 cannot be

delineated by substrate or activity assays, and expression of both followed a similar pattern in

liver after recovery from fibrosis. This finding was validated by investigating levels of the

intrinsic substrate for Capn1/2 (α-fodrin) and via an enzymatic assay (Figure 6B). The calpain

family are cysteine proteases that cleave a myriad of protein substrates. Although they are

generally localized intracellularly, they can be excreted when proteolytically activated (52).

Calpain activation has been identified as a key driver of inflammation in other diseases, such as

atherosclerosis and cardiac disease (52, 53). Moreover, calpain activation has been identified

to be involved in diseases of remodeling (i.e., fibrosis) via mechanisms that remain unclear (54).

Interestingly, the predicted substrates for calpain activation not only included key ECM proteins

[e.g., collagens and fibrin(ogens)], but also nuclear proteins, such as histones (Figure 6C),

which is in-line with previous findings (55). Few studies have identified calpain activation to play

a role in hepatic injury. These data suggest that the role of calpains in liver disease and fibrosis

may be underappreciated at this time.

Taken together, the results of this study indicate that remodeling of the liver during

recovery from fibrosis is a complex and highly coordinated process that extends well beyond the

degradation of the collagenous scar. A key role of MMP12, and by extension elastin, was

demonstrated, which is in-line with previous studies. Moreover, it was shown that even during

late stages of recovery (i.e., 28 D after CCl_4 cessation), there is significant enhanced turnover of

proteins. These results also indicate that analysis of the plasma degradome may yield new

insight into the mechanisms of fibrosis recovery, and by extension, new "theragnostic" targets.

Lastly, a novel potential role for calpain activation in the degradation and turnover of proteins

was identified.

Reference List

1.   Seki E, Brenner DA. Recent advancement of molecular mechanisms of liver fibrosis. J
Hepatobiliary Pancreat Sci 2015;22:512-518.
2.   Arteel GE, Naba A. The liver matrisome - looking beyond collagens. JHEP Rep 2020;2:100115.
3.   Iredale JP. Hepatic stellate cell behavior during resolution of liver injury. Semin Liver Dis
2001;21:427-436.
4.   Campana L, Iredale JP. Regression of Liver Fibrosis. Semin Liver Dis 2017;37:1-10.
5.   Shahid I, AlMalki WH, Hassan S, Hafeez MH. Real-world challenges for hepatitis C virus
medications: a critical overview. Crit Rev Microbiol 2018;44:143-160.
6.   Ramachandran P, Iredale JP, Fallowfield JA. Resolution of liver fibrosis: basic mechanisms and
clinical relevance. Semin Liver Dis 2015;35:119-131.
7.   Bonnans C, Chou J, Werb Z. Remodelling the extracellular matrix in development and disease.
Nat Rev Mol Cell Biol 2014;15:786-801.
8.   Friedman SL. Stellate cell activation in alcoholic fibrosis--an overview. Alcohol Clin Exp Res
1999;23:904-910.
9.   Gressner OA, Weiskirchen R, Gressner AM. Evolving concepts of liver fibrogenesis provide new
diagnostic and therapeutic options. Comp Hepatol 2007;6:7.
10.   Pellicoro A, Aucott RL, Ramachandran P, Robson AJ, Fallowfield JA, Snowdon VK, Hartland SN, et
al. Elastin accumulation is regulated at the level of degradation by macrophage metalloelastase (MMP-
12) during experimental liver fibrosis. Hepatology 2012;55:1965-1975.
11.   Ehling J, Bartneck M, Fech V, Butzbach B, Cesati R, Botnar R, Lammers T, et al. Elastin-based
molecular MRI of liver fibrosis. Hepatology 2013;58:1517-1518.
12.   Kisseleva T, Brenner D. Molecular and cellular mechanisms of liver fibrosis and its regression.
Nat Rev Gastroenterol Hepatol 2021;18:151-166.
13.   Massey VL, Dolin CE, Poole LG, Hudson SV, Siow DL, Brock GN, Merchant ML, et al. The hepatic
"matrisome" responds dynamically to injury: Characterization of transitional changes to the extracellular
matrix in mice. Hepatology 2017;65:969-982.
14.   Rittié L. Method for Picrosirius Red-Polarization Detection of Collagen Fibers in Tissue Sections.
Methods Mol Biol 2017;1627:395-407.
15.   Arteel GE, Iimuro Y, Yin M, Raleigh JA, Thurman RG. Chronic enteral ethanol treatment causes
hypoxia in rat liver tissue in vivo. Hepatology 1997;25:920-926.
16.   Kohli V, Madden JF, Bentley RC, Clavien PA. Calpain mediates ischemic injury of the liver through
modulation of apoptosis and necrosis. Gastroenterology 1999;116:168-178.
17.   Edelstein CL. Calpain activity in rat renal proximal tubules. An in vitro assay. Methods Mol Biol
2000;144:233-238.
18.   Bergheim I, Guo L, Davis MA, Duveau I, Arteel GE. Critical role of plasminogen activator
inhibitor-1 in cholestatic liver injury and fibrosis. J Pharmacol Exp Ther 2006;316:592-600.
19.   Isaacson RH, Beier JI, Khoo NK, Freeman BA, Freyberg Z, Arteel GE. Olanzapine-induced liver
injury in mice: aggravation by high-fat diet and protection with sulforaphane. J Nutr Biochem
2020;81:108399.
20.   Koch RG, Tsamis A, D'Amore A, Wagner WR, Watkins SC, Gleason TG, Vorp DA. A custom image-
based analysis tool for quantifying elastin and collagen micro-architecture in the wall of the human
aorta from multi-photon microscopy. J Biomech 2014;47:935-943.
21.   Keshishian H, Addona T, Burgess M, Kuhn E, Carr SA. Quantitative, multiplexed assays for low
abundance proteins in plasma by targeted mass spectrometry and stable isotope dilution. Mol Cell
Proteomics 2007;6:2212-2229.

22. Carr SA, Abbatiello SE, Ackermann BL, Borchers C, Domon B, Deutsch EW, Grant RP, et al. Targeted peptide measurements in biology and medicine: best practices for mass spectrometry-based assay development using a fit-for-purpose approach. Mol Cell Proteomics 2014;13:907-917.

23. Merchant ML, Barati MT, Caster DJ, Hata JL, Hobeika L, Coventry S, Brier ME, et al. Proteomic Analysis Identifies Distinct Glomerular Extracellular Matrix in Collapsing Focal Segmental Glomerulosclerosis. J Am Soc Nephrol 2020;31:1883-1904.

24. Szklarczyk D, Gable AL, Lyon D, Junge A, Wyder S, Huerta-Cepas J, Simonovic M, et al. STRING v11: protein-protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. Nucleic Acids Res 2019;47:D607-D613.

25. Szklarczyk D, Gable AL, Nastou KC, Lyon D, Kirsch R, Pyysalo S, Doncheva NT, et al. The STRING database in 2021: customizable protein-protein networks, and functional characterization of user-uploaded gene/measurement sets. Nucleic Acids Res 2021;49:D605-d612.

26. von Montfort C, Beier JI, Kaiser JP, Guo L, Joshi-Barve S, Pritchard MT, States JC, et al. PAI-1 plays a protective role in CCl4-induced hepatic fibrosis in mice: role of hepatocyte division. Am J Physiol Gastrointest Liver Physiol 2010;298:G657-666.

27. Roderfeld M. Matrix metalloproteinase functions in hepatic injury and fibrosis. Matrix Biol 2018;68-69:452-462.

28. Hemmann S, Graf J, Roderfeld M, Roeb E. Expression of MMPs and TIMPs in liver fibrosis - a systematic review with special emphasis on anti-fibrotic strategies. J Hepatol 2007;46:955-975.

29. Sage EH, Gray WR. Evolution of elastin structure. Adv Exp Med Biol 1977;79:291-312.

30. Klein J, Eales J, Zurbig P, Vlahou A, Mischak H, Stevens R. Proteasix: a tool for automated and large-scale prediction of proteases involved in naturally occurring peptide generation. Proteomics 2013;13:1077-1082.

31. Grgurevic I, Bozin T, Madir A. Hepatitis C is now curable, but what happens with cirrhosis and portal hypertension afterwards? Clin Exp Hepatol 2017;3:181-186.

32. D'Ambrosio R, Aghemo A, Fraquelli M, Rumi MG, Donato MF, Paradis V, Bedossa P, et al. The diagnostic accuracy of Fibroscan for cirrhosis is influenced by liver morphometry in HCV patients with a sustained virological response. J Hepatol 2013;59:251-256.

33. Bachofner JA, Valli PV, Kroger A, Bergamin I, Kunzler P, Baserga A, Braun D, et al. Direct antiviral agent treatment of chronic hepatitis C results in rapid regression of transient elastography and fibrosis markers fibrosis-4 score and aspartate aminotransferase-platelet ratio index. Liver Int 2017;37:369-376.

34. Chan J, Gogela N, Zheng H, Lammert S, Ajayi T, Fricker Z, Kim AY, et al. Direct-Acting Antiviral Therapy for Chronic HCV Infection Results in Liver Stiffness Regression Over 12 Months Post-treatment. Dig Dis Sci 2018;63:486-492.

35. Dolmazashvili E, Abutidze A, Chkhartishvili N, Karchava M, Sharvadze L, Tsertsvadze T. Regression of liver fibrosis over a 24-week period after completing direct-acting antiviral therapy in patients with chronic hepatitis C receiving care within the national hepatitis C elimination program in Georgia: results of hepatology clinic HEPA experience. Eur J Gastroenterol Hepatol 2017;29:1223-1230.

36. Persico M, Rosato V, Aglitti A, Precone D, Corrado M, De Luna A, Morisco F, et al. Sustained virological response by direct antiviral agents in HCV leads to an early and significant improvement of liver fibrosis. Antivir Ther 2017.

37. Vispo E, Barreiro P, Del Valle J, Maida I, de Ledinghen V, Quereda C, Moreno A, et al. Overestimation of liver fibrosis staging using transient elastography in patients with chronic hepatitis C and significant liver inflammation. Antivir Ther 2009;14:187-193.

38. Friedman SL, Pinzani M. Hepatic Fibrosis 2022: Unmet Needs and a Blueprint for the Future. Hepatology 2021.

39.   Zhao W, Yang A, Chen W, Wang P, Liu T, Cong M, Xu A, et al. Inhibition of lysyl oxidase-like 1
(LOXL1) expression arrests liver fibrosis progression in cirrhosis by reducing elastin crosslinking. Biochim
Biophys Acta Mol Basis Dis 2018;1864:1129-1137.
40.   Chen W, Yan X, Xu A, Sun Y, Wang B, Huang T, Wang H, et al. Dynamics of elastin in liver fibrosis:
Accumulates late during progression and degrades slowly in regression. J Cell Physiol 2019;234:22613-
22622.
41.   Dhital B, Durlik P, Rathod P, Gul ENF, Wang Z, Sun C, Chang EJ, et al. Ultraviolet radiation
reduces desmosine cross-links in elastin. Biochem Biophys Rep 2017;10:172-177.
42.   Schrader M, Schulz-Knappe P, Fricker LD. Historical perspective of peptidomics. EuPA Open
Proteomics 2014;3:171-182.
43.   Schrader M, Schulz-Knappe P. Peptidomics technologies for human body fluids. Trends
Biotechnol 2001;19:S55-60.
44.   Menschaert G, Vandekerckhove TT, Baggerman G, Schoofs L, Luyten W, Van Criekinge W.
Peptidomics coming of age: a review of contributions from a bioinformatics angle. J Proteome Res
2010;9:2051-2061.
45.   Greening DW, Kapp EA, Simpson RJ. The Peptidome Comes of Age: Mass Spectrometry-Based
Characterization of the Circulating Cancer Peptidome. Enzymes 2017;42:27-64.
46.   Lee KC, Hsu WF, Hsieh YC, Chan CC, Yang YY, Huang YH, Hou MC, et al. Dabigatran Reduces Liver
Fibrosis in Thioacetamide-Injured Rats. Dig Dis Sci 2019;64:102-112.
47.   Poole LG, Pant A, Baker KS, Kopec AK, Cline-Fedewa HM, Iismaa SE, Flick MJ, et al. Chronic liver
injury drives non-traditional intrahepatic fibrin(ogen) crosslinking via tissue transglutaminase. J Thromb
Haemost 2019;17:113-125.
48.   Beier JI, Luyendyk JP, Guo L, von Montfort C, Staunton DE, Arteel GE. Fibrin accumulation plays a
critical role in the sensitization to lipopolysaccharide-induced liver injury caused by ethanol in mice.
Hepatology 2009;49:1545-1553.
49.   Kearney KJ, Ariëns RAS, Macrae FL. The Role of Fibrin(ogen) in Wound Healing and Infection
Control. Semin Thromb Hemost 2021.
50.   Yu P, Zhong W. Hemostatic materials in wound care. Burns Trauma 2021;9:tkab019.
51.   Naymagon L, Tremblay D, Zubizarreta N, Moshier E, Mascarenhas J, Schiano T. Safety, Efficacy,
and Long-Term Outcomes of Anticoagulation in Cirrhotic Portal Vein Thrombosis. Dig Dis Sci
2021;66:3619-3629.
52.   Zhang M, Wang G, Peng T. Calpain-Mediated Mitochondrial Damage: An Emerging Mechanism
Contributing to Cardiac Disease. Cells 2021;10.
53.   Miyazaki T, Miyazaki A. Dysregulation of Calpain Proteolytic Systems Underlies Degenerative
Vascular Disorders. J Atheroscler Thromb 2018;25:1-15.
54.   Letavernier E, Zafrani L, Perez J, Letavernier B, Haymann JP, Baud L. The role of calpains in
myocardial remodelling and heart failure. Cardiovasc Res 2012;96:38-45.
55.   Arnandis T, Ferrer-Vicens I, Torres L, García C, Garcia-Trevijano ER, Zaragoza R, Viña JR.
Differential functions of calpain 1 during epithelial cell death and adipocyte differentiation in mammary
gland involution. Biochem J 2014;459:355-368.

Figure 1. Recovery from liver injury and fibrosis.

Panel A: Mice were administered CCl_4 for 4 weeks to induce hepatic fibrosis as described in

Materials and Methods. Animals were sacrificed 1, 3, 7, 14 and 28 days after cessation of CCl4

to track fibrosis resolution. Panel B: Representative photos of livers and photomicrographs

(100×) of stained liver sections are shown.   Hematoxylin and eosin (H&E) shows general

histology, while Sirius red brightfield (BF) and polarized light (PL) show fibrosis.

Figure 2. Effect of CCl_4 and recovery on indices of liver injury and fibrosis.

Experimental groups and approaches are as described in Experimental Procedures. Panel A

shows plasma transaminase (AST and ALT; IU/L) levels and hepatic mRNA expression of key

indices of fibrosis (*Col1a1*, *Acta2*, *Lox1*, *Lox2* and *P4htm*; Log2FC) are shown. Panel B shows

representative zymographs demonstrating gelatin lysis by MMP-2 and -9. Panel C shows

densitometric analysis of zymography (left; FC) and hepatic expression of key tissue inhibitors

of metalloproteinases (*Timp1*, *Timp2* and *Timp3*; right; Log2FC). Quantitative data are reported

as means ± SEM (n=4-6). [a]P <0.05 compared with absence of CCl_4 by ANOVA using

Bonferroni's post hoc test.

Figure 3. Hepatic expression of MMPs and elastin after fibrosis and recovery.

Experimental groups and approaches are as described in Experimental Procedures. Panel A

shows hepatic mRNA expression of key indices of fibrosis of all hepatic matrix

metalloproteinases (Mmps) both as Log2FC (left) and copy number (right). Panel B shows

elastin protein in the liver by Western blot (upper), immunofluorescence (100×; middle) and

multiphoton microscopy (MPM, 100×, lower). MPM images depict elastin (green), collagen

(blue) and stained nuclei (red). Panel C summarized densitometric quantitation of elastin by

Western blot (FC). Quantitative data are reported as means ± SEM (n=4-6). [a]P <0.05

compared with absence of CCl_4 by ANOVA using Bonferroni's post hoc test.

Figure 4. The effect of MMP-408 on recovery from liver injury and fibrosis.

Experimental groups and approaches are as described in Experimental Procedures. The effect

of MMP-408 on liver transaminase levels (Figure 3A, left) and fibrosis genes (Figure 3A ,middle

and right) during recovery were determined. The effect of MMP-408 on resolution of fibrosis

was assessed by Sirius Red staining (100×) (Figure 3B). Intensity of Sirius red staining was

assessed by image analysis (Figure 3C, left). Quantitative data are reported as means ±SEM,

(n = 8-10). [a]P <0.05 compared with absence of CCl_4; [b]P <0.05 compared with vehicle group by

ANOVA using Bonferroni's post hoc test.

Figure 5. Effect of fibrosis and recovery on the plasma degradome.

Experimental groups and approaches are as described in Experimental Procedures. Plasma

peptides were measured by LC-MS/MS, as described in experimental procedures and

compared naive control mice (Log2FC). Panel A depicts StringDB clustering analysis of

peptides significantly increased 28 days after cessation of CCl_4. Cluster groups are

summarized in Supplemental Table 4. Panel B shows the pattern of changes in peptides over

the course of the recovery period that were increased at 28 days. Panel C compares the

change in peptides after 28 days of recovery, versus those a 1 day. Panel D is a Venn diagram

comparing the peptides increased at 28 days of recovery to those increased at 1 day of

recovery.

Figure 6. Expression and activity of predicted protease from ProteaSix.

Experimental groups and approaches are as described in Experimental Procedures. ProteaSix

was used to predict proteases that formed these peptides (Figure 6A). Panel B (upper) depicts

western blot analysis of α-fodrin. The 240 kDa band represents intact α-fodrin, while the 150

and 120 kDa bands represent calpain-cleaved fragments. Densitometric analysis is also shown

(lower, right). The expression (lower, left) of *Capn1* and *Capn1* were determined by real-time

rtPCR. Panel 6C summarizes the type of parent proteins identified a potential substrates for

Capn1/2 cleavage by ProteaSix, organized by subcellular localization. The relative size of the

pie charts (1 D vs 28 D) is reflective of the amount of peptides identified at each timepoint.

Quantitative data are reported as means ±SE, n=4-6. [a]P <0.05 compared with absence of CCl_4

by ANOVA using Bonferroni's post hoc test.

Figure 7. Proposed role of MMP-12 and elastin during recovery from liver fibrosis

Hepatic activity of MMPs was robustly induced; but MMP-12 returned to baseline after recovery

from fibrosis. Elastic fibers, which are substrates of MMP-12, are also critical in fibrotic ECM

formation, and appeared after CCl_4 administration. Fibrosis resolved and histology was nearly

normal after 28 D recovery, but elastin ECM still remained elevated. Mature tropoelastin chains

are connected with desmosine cross-links. Newly synthesized immature elastin lacks this post-

translational modification. As recovery progresses, total elastin decreases, but desmosine

content increases; interestingly, blocking MMP-12 activity prevented both of these phenomena.

3   Figure 1-Sato et al.

Days of Recovery after CCl$_4$

                          Figure 2-Sato et al.

2   Figure 3-Sato et al.

3   Figure 4-Sato et al.

3   Figure 5-Sato et al.

Figure 7-Sato et al.