Fibrosis resolution in the mouse liver: role of Mmp12 and potential role of Calpain 1/2

Toshifumi Sato,[1] Kimberly Z. Head,[7] Jiang Li,[1] Christine E. Dolin,[7] Daniel Wilkey,[8,9] Nolan

Skirtich,[3] Dylan D. McCreary,[3] Sylvia Liu[4,5] Juliane I Beier,[1,5,6] Ryan M. McEnaney,[3] Michael L

Merchant[8,9] and Gavin E Arteel[1,5]

[1]Department of Medicine, Division of Gastroenterology, Hepatology and Nutrition, [2]Department

of Pharmacology and Chemical Biology, [3]Department of Surgery, [4]Department of Pathology,

[5]Pittsburgh Liver Research Center,[6] Department of Environmental and Occupational Health,

University of Pittsburgh, Pittsburgh, PA 15213. [7]Department of Pharmacology and Toxicology,

[8]Department of Medicine, Division of Nephrology and Hypertension and [9]University of Louisville

Alcohol Research Center, University of Louisville, Louisville, KY 40292

Running title: Resolution of fibrosis: beyond collagen.

Key Words: MMP-12, Elastin, fibrosis

1   Send all correspondence to: Gavin E. Arteel, PhD, FAASLD

2   Thomas E. Starzl Biomedical Science Tower

3   West 1143

4   200 Lothrop Street

5   Pittsburgh, PA 15213

6   Phone: +1-412-648-4187

7   Email: gearteel@pitt.edu

10   Abbreviations: CCl_4, carbon tetrachloride; ECM, extracellular matrix; ELISA, enzyme-linked

11   immunosorbent assay; GO, gene ontology; LC-MS/MS, liquid Chromatography with tandem

12   mass spectrometry; MMP, matrix metalloprotease; MPM, multiphoton microscopy; rtPCR,

13   reverse-transcriptase polymerase chain reaction; SDS, sodium dodecyl sulfate.

15   Acknowledgments:

Abstract

Although most work has focused on resolution of collagen ECM, fibrosis resolution involves

changes to several ECM proteins. The purpose of the current study was two-fold: 1) to examine

the role of MMP12 and elastin; and 2) to investigate the changes in degraded proteins in plasma

(i.e., the "degradome") in a preclinical model of fibrosis resolution. Fibrosis was induced by 4

weeks carbon tetrachloride (CCl_4) exposure, and recovery was monitored for an additional 4

weeks. Some mice were treated with daily MMP12 inhibitor (MMP408) during the resolution

phase. Liver injury and fibrosis was monitored by clinical chemistry, histology and gene

expression. The release of degraded ECM peptides in the plasma was analyzed using by 1D-

LC-MS/MS, coupled with PEAKS Studio (v10) peptide identification. Hepatic fibrosis and liver

injury rapidly resolved in this mouse model. However, some collagen fibrils were still present

28d after cessation of CCl_4. Despite this persistent collagen presence, expression of canonical

markers of fibrosis were also normalized. The inhibition of MMP12 dramatically delayed fibrosis

resolution under these conditions. LC-MS/MS analysis identified that several proteins were

being degraded even at late stages of fibrosis resolution. Calpains 1/2 were identified as

potential new proteases involved in fibrosis resolution. CONCLUSION. The results of this study

indicate that remodeling of the liver during recovery from fibrosis is a complex and highly

coordinated process that extends well beyond the degradation of the collagenous scar. These

results also indicate that analysis of the plasma degradome may yield new insight into the

mechanisms of fibrosis recovery, and by extension, new "theragnostic" targets. Lastly, a novel

potential role for calpain activation in the degradation and turnover of proteins was identified.

Regardless of etiology, hepatic fibrosis is an almost universal endpoint of chronic liver injury,

when coupled with impaired/incomplete recovery from said injury (1). The clinical sequelae of

severe fibrosis/cirrhosis significantly and negatively impact patient survival and quality of life.

Given these clinical implications, the mechanisms of hepatic fibrosis initiation and progression

have been studied for decades with the goal of identifying better means to prevent and treat this

disease (2). Although much progress has been made in our understanding, the translational

value of these studies has been hampered by the fact that clinical diagnosis of liver disease is

often late-stage, where mechanism-based therapies appear to be less effective.

It has been known that for several years that hepatic fibrosis may resolve, if the underlying

causal factor is successfully removed (3). This phenomenon was consistently observed in

animal models of fibrosis, but until recently only in rare cases in humans (4). The recent

advance of highly effective disease-modifying agents (e.g., direct acting antivirals for hepatitis C

viral infection) have dramatically increased the opportunity for fibrosis resolution to occur (5).

However, not all patients for achieve fibrosis resolution when the underlying disease has been

effectively treated (6). Moreover, in contrast to the development of fibrogenesis, the

mechanisms by which fibrosis resolution is mediated are poorly understood and may not share

complete mechanistic overlap with fibrosis progression (4). To address both this clinical need

and the gap in our understanding, more investigations in this area need to be performed.

The hepatic extracellular matrix (ECM) consists of a diverse range of components that

work bi-directionally with surrounding cells to create a dynamic and responsive

microenvironment that regulates cell and tissue homeostasis. Under normal conditions, the

ECM assists in maintaining organ homeostasis and appropriate responses to injury/stress by

dynamically responding to these insults. However, failure to properly regulate these responses

can lead to qualitative and/or quantitative ECM changes that are maladaptive (7). Hepatic

fibrosis is a canonical example of ECM dyshomeostasis, leading to accumulation of fibrillary

ECM, such as collagen, which is considered the pathologic hallmark of fibrosis. Although

hepatic fibrosis is considered almost synonymous with collagen accumulation (8), the qualitative

and quantitative alterations to the hepatic ECM during fibrosis are much more diverse (9).

Indeed, hepatic fibrosis results in qualitative and quantitative changes to almost all components

of the hepatic ECM (9). This understanding has let previous work by others to investigate the

roles of non-collagen ECM components, as well as enzymes that metabolize them (e.g., matrix

metalloproteinases; MMPs) in the initiation and development of experimental and clinical fibrosis

(10, 11). In contrast, previous studies of fibrosis resolution have focused almost exclusively on

secretion and turnover of collagens and on collagen microstructure (12). The purpose of the

current study was to investigate turnover of hepatic ECM during fibrosis resolution in mice.

Experimental Procedures

Please see Supplemental Materials for additional details on Experimental Procedures.

Animals and Treatments. 8 weeks old, male C57Bl/6J mice purchased from Jackson Laboratory (Bar Harbor, ME) were housed in a pathogen-free barrier facility accredited by the Association for Assessment and Accreditation of Laboratory Animal Care, and procedures were approved by the local Institutional Animal Care and Use Committee. Animals were allowed standard laboratory chow and water *ad libitum*. Mice were administered CCl_4 (1 ml/kg i.p.; diluted 1:4 in olive oil; Sigma-Aldrich, St. Louis, MO) or vehicle 2×/wk for 4 wks; animals were sacrificed 1, 3, 7, 14 and 28 days after final CCl_4 injection (Figure 1A). Animals were administered an MMP-12 inhibitor (MMP-408; 10 mg/kg, i.g.; Merck Millipore, Burlington, MA) or vehicle (saline) daily starting after the last injection of CCl_4. At the time of sacrifice, animals were anesthetized with ketamine/xylazine (100/15 mg/kg, i.p.). Blood was collected from the vena cava just prior to sacrifice by exsanguination and citrated plasma was stored at -80°C for further analysis. Portions of liver tissue were frozen immediately in liquid nitrogen, while others were fixed in 10% neutral buffered formalin or embedded in frozen specimen medium (Tissue-Tek OCT compound, Sakura Finetek, Torrance, CA) for subsequent sectioning and mounting on microscope slides.

Biochemical assays and histology. Plasma levels of Aspartate transaminase (AST) and alanine amino transferase (ALT) were determined spectrophotometrically using standard kits (Thermo Fisher Scientific, Waltham, MA), as described previously (13). Formalin-fixed, paraffin embedded sections were cut at 5 μm and mounted on glass slides. Deparaffinized sections stained with hematoxylin and eosin and pathology was assessed in a blinded manner. ECM accumulation in liver sections was determined by staining with Sirius red/fast green and were

visualized via brightfield and polarized light (14). Bright field images of Sirius red staining was

quantitated by image analysis, as described previously (13). Specifically, a Molecular Devices

(Sunnyvale, CA) Image-1/AT image acquisition and analysis system incorporating an Axioskop

50 microscope (Carl Zeiss Inc., Thornwood, NY) was used to capture and analyze five

nonoverlapping fields per section at 100× final magnification. Image analysis was performed

using modifications of techniques described previously. Detection thresholds were set for the

red color based on an intensely labeled point and a default color threshold. The degree of

labeling in each section was determined from the area within the color range divided by the total

area.

Hepatic calpain activity. Calpain activity was measured in liver lysate, using N-succinyl-

Leucine-Leucine-Valine-Tyrosine-7-amino-4-methylcoumarin (Suc-Leu-Leu-Val-Tyr-AMC;

Supplemental Table 3) was used as the calpain substrate, as described previously (15). Forty µl

of liver lysate samples (0.25 mg protein/ml in imidazole buffer containing 63 mM imidazole, 10

mM mercaptoethanol, 1 mM EDTA, 10 mM EGTA, pH 7.3) were incubated at 37 for 1hr with

160 µl of imidazole buffer containing 50 µM Suc-Leu-Leu-Val-Tyr-AMC (dissolved in DMSO).

The formation of the cleavage product, 7-amino-4-methylcoumarin (AMC), was measured by

fluorometrically (Spectra Max M5, Molecular Devices Corporation) at 360 nm excitation and 430

nm emission wavelengths. Standard curves were generated with free AMC for each experiment

and results were normalized to total protein.

Determination of MMP-2 and 9 by zymography. Zymography was performed as

described previously (16). Total hepatic protein was extracted using a lysis buffer consisting of

10 mM cacodylic acid, pH 5.0, containing 150 mM NaCl, 1M ZnCl, 15 mM $CaCl_2$, 1.5 mM NaN_3,

and 0.01% Triton X-100. Lysates were then diluted in 2× sample buffer (125 mM Tris HCl, pH

6.8, with 4% SDS, 20% glycerol) without boiling or reduction. The diluted samples were

subjected to 0.1% gelatin–10 % SDS polyacrylamide gel electrophoresis (Novex, Carlsbad, CA).

The gels were washed with 2.5 % Triton X-100 after electrophoresis and then incubated for 48

hours in the reaction buffer (50 mM Tris, 0.1 M glycine, and 0.1 M NaCl, pH 8.0). The gel was

stained with Coomassie brilliant blue R-250 (0.1 % Coomassie Brilliant Blue R250, 50%

methanol, and 10% acidic acid) for 1hr, destained with 45% methanol and 10% acetic acid for

30 minutes. The results were visualized and densitometric analysis was performed using an

iBright imager and onboard software from Invitrogen and normalized to total protein (Waltham,

MA).

Immunoblots. Liver samples were homogenized in RIPA buffer (20 mM Tris/Cl, pH 7.5,

150 mM NaCl, 1 mM EDTA, 1 mM EGTA, 1% [wt/vol] Triton X-100), containing protease and

phosphatase inhibitor cocktails (Sigma). Samples were diluted in 2× sample buffer (see above)

and loaded onto 10% SDS-polyacrylamide gels (Invitrogen, Waltham, MA), followed by

electrophoresis and Western blotting on to PVDF membranes (Hybond P, GE Healthcare Bio

Sciences, Pittsburgh, Pennsylvania). Primary antibodies against proteins of interest were

compared to GAPDH as a housekeeping gene (see Supplemental Table 3). The results were

visualized and densitometric analysis was performed using an iBright imager and onboard

software from Invitrogen (Waltham, MA). Protein levels were normalized either to an internal

housekeeping gene (GADPH) or, as in the case for α-fodrin, the ratio of cleaved to total α-fodrin

protein signal.

<u>RNA Isolation and Real-Time RT-PCR</u> RNA extraction and real-time RT-PCR were

performed as described previously (16). RNA was extracted immediately following sacrifice

from fresh liver samples using RNA Stat60 (Tel-Test, Ambion, Austin, TX) and chloroform. RNA

concentrations were determined spectrophotometrically and 1µg of total RNA was reverse

transcribed using the QuantiTect Reverse Transcription Kit (Qiagen,Valencia, CA). Real-time

RT-PCR was performed using a StepOne real time PCR system (Thermo Fisher Scientific,

Grand Island, NY) using the Taqman Universal PCR Master Mix (Life Technologies, Carlsbad,

CA). Primers and probes were ordered as commercially available kits (Thermo Fisher Scientific,

Grand Island, NY; see Supplemental Table 1). The comparative C_T method was used to

determine fold differences between the target genes and an endogenous reference gene (18S).

Results were reported as copy number ($2^{-\Delta CT}$) or fold naïve control (FC; $2^{-\Delta\Delta CT}$).

<u>MMP12 activity</u>. MMP-12 assay kit was purchased from AnaSpec (Fremont, CA).

Frozen liver tissue (50-75 mg) was homogenized in the provided assay buffer containing 0.1%

(v/v) Triton-X 100, and subsequently centrifuged 10,000×G for 15 min. MMP-12 activity was

measured by FRET ($490_{ex}/520_{em}$) in the resultant supernatant according to the kit's instructions

and calibrated to commercially-available purified MMP-12 (AnnSpec) and normalized to protein

levels in the original homogenate.

<u>Immunofluorescence.</u> For immunofluorescent detection of elastin, sections of liver were

blocked with PBS containing 10 % goat serum for 30 min, followed by incubation overnight at

4 °C with Elastin (Abcam 21607 1:50) antibodies in blocking solution. Sections were washed

with PBS and incubated for 3 hrs with a secondary antibody conjugated to Alexa 488 (Invitrogen,

Carlsbad, CA). Sections were washed with PBS and mounted with mounting medium containing

1.5 µg/ml DAPI (VECTASHIELD w/DAPI; Vectastain, Torrance, CA). Specificity of staining was validated by comparison to negative controls in the absence of primary antibody.

Multiphoton microscopy imaging. Elastin and collagen content in the samples were assessed using multiphoton microscopy (MPM), as described previously (17). MPM were performed on frozen sections with a nonlinear optical microscope based on a confocal imaging system (FV1000MPE, Olympus, Tokyo, Japan; purchased with grant 1S10RR025676-01 awarded to Dr Simon C. Watkins) using an optical microscope and an external tunable mode-locked Ti:sapphire LASER (Coherent, Palo Alto, CA) set to 790-800 nm, 5.4% intensity. LASER emission and signal detection were through a 25x, XL Plan N 1.05 N.A. water immersion objective lens (Olympus). Detection was synchronous across three channels. Nuclei stained by propidium iodide (Invitrogen) were detected on the RXD4 channel (663-737 nm emission filter). Fibrillar collagen second-harmonic generation was detected on the RXD1 channel (350-450 nm emission filter). Elastin-associated fibrillar green autofluorescence was detected on the RXD2 channel (500-550 nm emission filter). A 4x line Kalman integration filter was applied to reduce stochastic noise. The laser was routed to an acousto-optic modulator (AOM) for power attenuation, then through a dichroic mirror (490 nm) and an objective to the tissue sample. Signals from second harmonic generation (SHG) were collected by the same objective lens and recorded by a photomultiplier tube (PMT).

Preparation of acid hydrolysates of liver samples and Quantitation of Desmosine Crosslinks. Liver samples were homogenized in RIPA buffer (see above) and autoclaved (3 h) twice, at 20 lb/in.2 (1.36 atm) to remove collagen and other proteins. The residue was treated with 0.1 M NaOH for 30 min at 100 °C, washed until neutral, and lyophilized. The isolated

elastin was flushed with nitrogen and hydrolyzed in 6 N HCl under vacuum at 105°C for 24-36 h.

The hydrolysates were evaporated to dryness and the residue was dissolved in 0.5-1.0 ml water.

Appropriate aliquots were then added to 0.1 M sodium phosphate buffer, pH 7.4, containing

phenol red and neutralized with 2 M NaOH. Desmosine content was determined using a

commercially available ELISA kit (Cusabio, Houston, TX) and normalized to starting tissue wet

weight.

<u>Plasma peptide purification and high resolution LC-MS analysis</u>. 100 µL PBS containing

0.1X iRT standards (Biognosis Inc, Beverly, MA) was added to 100 µL plasma. 200 µL 20% w/v

TCA was mixed into each sample on ice. Samples were then be incubated for 1 hour at 4°C.

The precipitate was pelleted by centrifugation at 16,000 × G for 10 min at 4°C; pellets were

discarded. The centrifugation was then repeated with the supernatant samples, followed by

discard of the second pellets. Samples were then desalted and concentrated using solid phase

extraction (Waters Oasis HLB µElution 30 µm plate, part no. 186001828BA) as described (18).

Solid phase extraction eluates were dried in a SpeedVac, and dried peptides were resuspended

in 25 µL 2% v/v acetonitrile/ 0.1% v/v formic acid. Peptide concentrations were measured by

absorbance (NanoDrop2000) at 205 nm and concentrations estimated using a plasma tryptic

digest standard curve. Peptide samples (0.2 ug) were separated using an 12cm, 360µm OD x

100µm ID fused silica tip packed with Aeris Peptide 3.6µm XB-C18 100Å material

(Phenomenex, Torrance, CA, USA). Peptides were eluted under a 90min 2%-44% acetonitrile

gradient and transferred by Nanospray Flex nanoelectrospray source (ThermoFisher) into an

Orbitrap Elite - ETD mass spectrometer (ThermoFisher) using an Nth Order Double Play

(Xcalibur v2.2, ThermoFisher) with FTMS MS1 scans (240,000 resolution) collected from 300-

2000m/z and ITMS MS2 scans collected on up to twenty peaks having a minimum signal

threshold of 5,000 counts from the MS1 scan event.

Peptidomic Data Analysis. RAW files were searched in Peaks Studio X (Bioinformatics

Solutions Inc., Waterloo, ON, Canada) using the Denovo, PeaksDB, PeaksPTM, and Label Free

Q algorithms and the 2/12/2019 version of the UniprotKB reviewed reference proteome

canonical and isoform Homo sapiens sequences (Proteome ID UP000005640). Search

parameters included: variable methionine, proline and lysine oxidation (+15.9949 Da), no

enzyme specified, 15 ppm precursor error for MS1 Orbitrap FTMS data, 0.5 Da error for MS2

data sets, and common PTMs in the PeaksPTM algorithm. The peptide, feature, and protein-

peptide csv files were exported from the Label Free Q result for statistical tests in Microsoft

Excel 2016 and R v3.5.0. High confidence peptide assignments (PeaksX criteria –logP scores

with p-value <0.05) were exported into comma separated values files (.csv) for upload and

analysis by the Proteasix (http://www.proteasix.org/) algorithm using the 'Peptide-centric'

prediction tool based on curated, known and observed cleavage events enabling assignment of

protease identities from the MEROPS database as described previously (19) The positive

predictive value (PPV) cut-off for the prediction algorithm output was set to 80%. Protein-protein

interaction network analysis of regulated proteomic data sets (q-value <0.05) was performed

using Search Tool for the Retrieval of Interacting Genes/Proteins, STRING v11 (20), with the

highest confidence score (0.900).

Data Sharing. Acquired LCMS data (.RAW). Uniprot mouse sequence database annotated

with Biognosys iRT standards, metadata for sample key and PeaksX search results exported as

an excel file (.xlsx) files have been deposited in MassIVE

(massive.ucsd.edu/ProteoSAFe/static/massive.jsp) data repository (MassIVE ID: 000089311)

with the Center for Computational Mass Spectrometry at the University of California, San Diego.

Public data mining. The expression of Capn1 and Capn2 in hepatic from fibrosis was

explored, using data from public study GSE135251 [GSE135251-1, GSE135251-2] (21, 22),

which includes 46, 48, 54, 54 and 14 patients at fibrosis stage from 0 to 4 [by METAVIR score

(23)], respectively. Gene expression profile and clinical information was downloaded from GEO

database. The gene count data was normalized by R package DESeq2 normTransform function

(24). Expression for *CAPN1* and *CAPN2* as a function of fibrosis stages were visualized by

violin plot.

Results

Recovery from liver injury and fibrosis. All animals survived exposure to CCl_4 and no

morbidity or mortality was reported during the recovery phase. As expected (25), 4 wks CCl_4

administration caused severe liver injury and hepatic fibrosis in mouse liver as assessed by

histology (Figure 1B) and by transaminase release (Figure 2A). Four weeks of CCl_4

administration increased hepatic expression of key fibrogenic genes (*Col1a1*, *Acta2*, *P4htm*, *Lox*

and *Loxl2*) significantly compared with vehicle controls (Figure 2A). Hepatic fibrosis and liver

injury rapidly resolved after cessation of CCl_4 exposure (Figure 1B), with only thin septa of

collagen staining observable after 4 wks recovery (Figure 1B). Likewise, transaminase values

and expression of key fibrogenic genes all returned to baseline values after 4 weeks of recovery

(Figure 2A).

Effect of CCl_4 on expression and activity of MMPs. Even in cases where there is a net

increase in ECM in the liver (e.g., fibrosis) overall turnover is also increased (26). This turnover

is known to be mediated, at least in part, by matrix metalloproteinases (MMPs). 4 wk of CCl_4

administration increased the expression (Figure 2A) and activity (Figures 2B and 2C) of the

collagenases, MMP2 and MMP9. The mRNA expression of inhibitors of these MMPs (i.e.,

Timp1, *Timp2* and *Timp3*) were also induced with a similar profile (Figure 2C). As was

observed with expression of key indices of fibrogenesis (Figure 2A; see above), the increase in

expression/activity of these factors largely returned to baseline levels after 28 days of recovery.

In addition to collagen deposition, fibrosis involves quantitative and qualitative changes

to a myriad of ECM proteins and MMPs (9, 27). The expression of all hepatic MMPs was

therefore determined (Figure 3A). Analogous to findings with *Mmp2* and *Mmp9*, all hepatic

MMPs were induced by experimental fibrosis (Figure 3A). The expression pattern of the

majority of these MMPs was also similar to *Mmp2* and *Mmp9* and returned to baseline 28 days

after the cessation of CCl_4 (Figure 3A). The one contrast was the expression of *Mmp12*, which

was the most robustly induced MMP, both by fold change (1000-fold over control), as well as by

copy number (Figure 3A). Unlike other MMPs, expression and protein levels of *Mmp12* were

still elevated ~10-fold over baseline 28 days after cessation of CCl_4 (Figure 3A).

Effect of CCl_4 on elastin accumulation. Elastin is a key substrate for MMP12 that

increases during CCl_4-induced fibrosis (10). Expression and protein levels of elastin were

therefore determined (Figures 3B and C). Although the mRNA expression of elastin (*Eln*) was

only moderately (<4 f-fold) induced by 4 weeks of CCl_4 administration and returned to baseline

after 28 days of recovery, protein levels increased ~15-fold compared with naïve control mice

(Figures 3B and 3C). Moreover, elastin ECM was still significantly (~5 fold) elevated over naïve

control values even after 28 days of recovery. As has been observed in previous studies (10),

elastin accumulation appeared to correlate with areas of fibrotic changes (Figure 3B). Elastin

accumulation appears to center on areas of fibrogenesis and collagen accumulation (see Figure

3B). To investigate this further, the colocalization of elastin and collagen were determined using

multi-photon microscopy (Figure 3B), as it allows observation of the elastin and collagen fiber

networks simultaneously (17). Elastin (green fluorescence) accumulation was highest on the

first day of recovery and was closely associated with collagen (blue fluorescence) signals,

consistent with the assessment of elastin immunofluorescence (Figures 3B).

MMP12 inhibition delays fibrosis resolution. Previous work with knockout mice indicated

that MMP12-mediated elastin degradation plays a protective role in the development of CCl_4-

induced fibrogenesis (10); however, the role of MMP12/elastin on resolution of fibrosis was

unknown. The effect of the MMP12 inhibition on resolution of fibrosis was therefore determined

(Figure 4). CCl_4-exposed mice were administered the MMP12 inhibitor, MMP-408, daily after

cessation of toxicant exposure (see Materials and Methods). MMP-408 did not significantly

affect recovery of transaminases after CCl_4 exposure (Figure 4A). MMP-408 also did not

significantly impact the expression of fibrogenic genes (e.g., *Col1A1*) during the recovery phase

(Figure 4A). On the other hand, MMP408 impacted the resolution of collagenous fibrosis, as

determined by Sirius red staining (Figures 4B and C). Image-analysis and quantification of

Sirius red staining indicated that the amount of collagen present in the liver was double 1 wk

after cessation of CCl_4 administration compared to the mice administered the vehicle for

MMP408 (Figure 4C). Although this effect of MMP408 was not present after 28 days of

recovery, the fibrous septa in the MMP-408 group were far less defined than in the vehicle

treated (Figure 4B and inset).

The elastin signal in the multiphoton microscopy analysis shows both immature and

mature elastin [Figure 3B (17)]. Therefore, it reflects the total amount of elastin accumulation in

the liver. In contrast, the crosslinking of mature insoluble elastin is mediated via post-

translational incorporation of the amino acid desmosine. Desmosine is unique to mature elastin

in mammalian species and can therefore be employed as a surrogate marker of mature elastin

(28). The amount of desmosine present in the liver tissues after CCl_4 was therefore determined

by ELISA (Figure 4C), as described in Materials and Methods. Although the total amount of

elastin detected in the liver increased by 4 wks of CCl_4 administration (Figures 3), the

desmosine content in the liver actually decreased to ~50% of levels in naïve mice (Figure 4C).

These data suggest that the majority of the new elastin detected during fibrosis is immature

uncrosslinked elastin. During the resolution phase desmosine content in livers from mice

administered vehicle increased, with values not significantly different that naïve control mice 28

days after cessation of CCl_4. In contrast, administration of MMP408 completely attenuated the

increase in this index of mature elastin (Figure 4C).

The degradome changes dramatically during fibrosis resolution. Resolution of fibrosis

requires coordinated remodeling of the liver. The net effect of fibrosis and resolution on hepatic

protein degradation has not been previously determined. The influence of CCl_4 and 4 wks

recovery on the plasma peptide "degradome" was therefore determined (Figure 5). Across the

groups, 694 peptides were identified, corresponding to 181 distinct parent proteins. The parent

proteins that contributed to the pattern of significantly changed peptides were analyzed by

StringDB, followed by K-means cluster analysis [Figure 6 (20)]. This database queries physical

and functional protein-protein associations that integrates both experimental and predicted

associations. The output creates both graphical and categorical enrichment patterns in the

queried dataset. Cluster analysis yielded 7 enriched clusters in the total degradome, with a

strong enrichment of proteins associated with Gene Ontology (GO) terms for "high-density

lipoprotein particle" (GO:0034464; Cluster 1), "collagen trimer" (GO:0005581; cluster 6) and

"nucleosome" (GO:0000786; Cluster 7) (Figure 6; Table 1).

The degradome from 1 and 28 D of recovery were compared in more detail to represent

the peak of injury (1 D) to near resolution of injury (28 D; Figures 1 and 2). A total of 255

peptides were significantly changed at least >2 fold (compared to naïve control) at these

timepoints (Figure 5A, B and C). Although there was considerable overlap between these 2

timepoints (Figure 5C), the amount of peptides altered at the 28 D timepoint was ~2.5-fold more

compared to the 1 D timepoint. A total of 144 peptides, corresponding to 82 distinct proteins,

were significantly increased at 1 D after CCl_4 and/or 28 D after CCl_4 (Figure 5A and B,

Supplemental Table 4). All peptides that significantly increased at 28 D after recovery tended to

be at relatively low levels 3 and 7 days after cessation of CCl_4, then increasing at the 14 D

timepoint (Figure 5A).

Many proteases cleave substrates with high specificity with limited cleavage promiscuity.

Thus, the fragment sequence of degraded proteins can be informative on proteases that may

have generated this pattern. Proteasix is an open-source peptide-centric tool to predict in silico

the proteases involved in generating the observed peptide features (29). Figure 7A is a

frequency distribution figure comparing the predicted proteases that generated the significantly

increased peptides at the 1 and 28 D timepoint (vs. naïve control). The top predicted proteins,

calpains 1 and 2 were shared between the timepoints under these conditions (Figure 7A). The

expression of these splice variants (*Capn1* and *Capn2*) was determined via real-time rtPCR

(Figure 7B). The copy number of *Capn2* was approximately an order of magnitude higher than

Capn1 in liver (3.1×10^{-5} vs. 4.3×10^{-6}, respectively) under basal conditions. The expression of

both isoform variants was increased by CCl_4 administration but returned to baseline by 28 D of

recovery (Figure 7B). In contrast, predicted enzyme activity of CAPN, both by cleavage of α-

fodrin and by enzymatic analysis (Figure 7B), indicated that CAPN activity was elevated at the

28 D timepoint. Several of the predicted CAPN1/2 substrates for these enzymes (Figure 7A)

were analogous to the key clusters identified by StringDB analysis for the degradome as a

whole (Figure 6 and Table I), such as fibrinogens (cluster 1), collagens (cluster 6) and histones

(cluster 7) and their corresponding GO terms.

Capn2 is induced in human NASH fibrosis. These data indirectly suggest that calpains

may be novel mediators of hepatic fibrosis and resolution. However, the relevance of these

findings to human fibrotic liver disease is unclear. A recent study showed that the expression of

calpain proteases was induced in human livers with fibrosis caused by HBV (30). Here, the

expression of CAPN1 and CAPN2 was determined in hepatic NASH fibrosis (see Experimental

Procedures; Figure 7C). The expression of CAPN1 was slightly increased during early (F2)

fibrosis; in contrast, the expression of CAPN2 was increased in a step-wise fashion at every

fibrosis stage (Figure 7C).

Discussion

Direct-Acting Antiviral therapy has for all intents and purposes cured the hepatitis C virus (HCV) epidemic in immunocompetent adults. However, one of the most important issues is whether patients with significant HCV-induced fibrosis/cirrhosis will recover from this injury once a sustained virologic response is achieved (31). For example, improvement of liver stiffness in HCV cirrhosis after successful antiviral therapy is reportedly 30-65%, depending on the study referenced (32-36). Although promising at first glance, these data indicate that around half of the HCV-fibrosis/cirrhosis patients with successful viral clearance do not improve in their liver disease. Furthermore, another alarming concern is that the commonly used non-invasive transient elastographic analysis of liver stiffness tends to overpredict fibrosis/cirrhosis regression in HCV, compared to biopsy-based histologic assessment (31, 37). Indeed, despite achievement of viral clearance, histological assessment of patient livers often continues to show lingering inflammation, hepatocyte and ductular reaction and fibrosis (31, 37). It is therefore an unmet clinical need to accurately predict determining factors of fibrosis resolution. Although accumulation of collagen ECM is the *de facto* definition of fibrosis, the changes to the hepatic ECM that happen during both processes are much diverse than those that happen to collagen (9). Moreover, recent studies have suggested that livers that recover from fibrosis do not completely revert to the naive state and are more sensitive to reinduction of fibrosis (12). In toto, although it is understood that hepatic fibrosis resolution involves more than simple clearance of collagen ECM, the process and its mediators are incompletely understood, at best (38).

Previous studies have indicated that MMP12, most likely via mediating elastin ECM accumulation, is key for the development of hepatic fibrosis (10). As in other solid organs, MMP-12 is almost exclusively expressed at high levels in a subset of macrophages in the liver. In previous work Ramachandran et al. (39) identified a subset of Ly6C^{low} macrophages that

were enriched in the resolving livers; these macrophages expressed high levels of MMP-12.

Previous studies have shown that genetic or pharmacologic inhibition of MMP12 exacerbates

the development of experimental fibrosis. Furthermore, prevention of elastin crosslinking

protected against experimental fibrosis (40). More recently, Chen et al. (41) demonstrated that

elastin fibers degrade slowly during fibrosis resolution, a finding recapitulated in the current

study (Figure 3). The authors of that study suggested that cross-linked elastin stabilizes the

fibrotic ECM and may thereby protect it from degradation (41). However, whether MMP12

activity mediates resolution of fibrotic ECM was not clear. Here, it was demonstrated that the

MMP12 inhibitor, MMP408, delayed resolution of Sirius red-positive areas of the liver (Figure 4),

which supports the hypothesis that elastin degradation is key to overall fibrosis resolution.

Although elastin protein levels increased (Figure 3), the amount of mature crosslinked elastin

(as determined by desmosine content; Figure 4C) actually was decreased by CCl_4 exposure.

Desmosine content did not return to basal levels until after 28 D of recovery. These results

suggest that the increase in elastin content in the liver during fibrosis represents less mature

(crosslinked) protein oligomers (Figure 8). This finding is similar in models of photoaging in

which damaged skin accumulates disorganized elastin fibrils (42). Interestingly, inhibition of

MMP12 with MMP408 prevented the recovery of desmosine content during fibrosis resolution,

suggesting that MMP12 is critical not only for the degradation of accumulated elastin, but also

the reorganization to mature elastin (Figure 8). These findings should be explored in future

studies.

Although fibrosis resolution was delayed by MMP12 inhibition, there were no detectable

differences in total sirius red staining between vehicle and MMP12 inhibitor 28 days after the

cessation of CCl_4 intoxication (Figure 4C). However, the pattern of staining in the MMP12

inhibitor group was more diffuse and less well-defined compared to vehicle controls at this

timepoint (Figure 4B). These data do not necessarily indicate that MMP12 is equivocal in

fibrosis resolution. First, as shown by studying the degradome (see Figures 5-6), there is still

significant difference in protein turnover 28 days after cessation of CCl_4, a time at which the liver

is almost histologically normal (Figure 1). Moreover, spontaneous resolution of fibrosis in

rodents is much more robust than in humans and by extension, humans are likely more

sensitive to delays in fibrosis resolution (12). Although not yet studied in human liver disease,

previous work has shown that remodeling in other organ systems (e.g., atherosclerosis) are

sensitive to variants in MMP12 activity; for example, a gain-of-function promoter polymorphism

in MMP12 is associated with protection against coronary artery disease in diabetes (43).

Hypothesis-driven approaches are key to elucidating mechanism. However, the nature

of the approach by definition requires prior knowledge. Discovery-based 'omic techniques

coupled with informatic analyses can yield new information and insight into disease progression

and regression. The peptidome is defined as the population of low molecular weight biologically

derived peptides (0.5-3 kDa), within the cells and biologic fluids. These compartments contain

peptides critical for normal organismal function. The study of the peptidome (i.e., 'peptidomics')

is a discipline related to proteomics, but with significant methodological and analytical

differences (44). For example, as the original structure of the peptides is of interest, samples

are not trypsin digested prior to LC-MS/MS analysis, as is the case for bottom-up proteomic

approaches. This difference originally limited peptide identification, as the available databases

were based on trypsin-digested peptide fragments (45). However, the development of

peptidome-specific identification and analysis tools has addressed this concern (46). In addition

to synthesized peptides, the peptidome also contains fragments of proteins degraded by normal

and/or abnormal processes (i.e., 'degradome'). These degradomic profiles are conditionally

unique to the metabolic/pathologic status of the organism and reflect not only quantitative

changes (i.e., more/less parent protein to degrade), but also qualitative changes (i.e., change in

the pattern of protease digestion of parent proteins). The latter subset of the peptidome has

generated key interest in some areas of human health as possible (surrogate) biomarkers for

disease. Here, LC-MS/MS analysis of the degradome profile in plasma during recovery from

experimental fibrosis was analyzed to investigate the general pattern of protein

turnover/remodeling during recovery, as well as to identify potential new key players.

When the degradome was analyzed, an interesting pattern resolved. Specifically,

although livers were almost histologically-normal after 4 weeks of recovery from CCl_4-induced

experimental fibrosis (Figure 1), there was a relative increase in the turnover of proteins

observed in the plasma compartment (Figures 5-6), as determined by analysis of the

degradome. This finding is in-line with previous observations by this group that hepatic

remodeling in response to injury is much more diverse than can be observed histologically (13).

In addition to the appearance of degraded ECM proteins canonically associated with fibrosis

(e.g., collagens), several other degraded ECM proteins were present, especially after 28 days of

recovery. A limitation of degradome analysis is that the original source of the parent protein

cannot be completely verified. However, the CCl_4 model is relatively specific for liver injury and

the majority of the significantly changed peptides were hepatic by Brenda Tissue Ontology

analysis (BTO:0000759; $p=2.2\times10^{-14}$).

One of the most robust families of degraded proteins that were enriched 28 days after

cessation of CCl_4 were associated with the coagulation cascade, especially the fibrin(ogens)

(Figure 6, Table 1 and Supplemental Table 4). Previous studies have investigated the potential

role of fibrin(ogens) in the development of hepatic injury and fibrosis, with somewhat conflicting

results (47, 48). However, few if any studies have investigated the role of fibrin(ogen) ECM in

resolution of hepatic fibrosis. In contrast, the turnover of fibrin(ogen) ECM has been heavily

studied in other diseases of remodeling and recovery. For example, fibrin(ogen) ECM plays a

key role in sub-cutaneous wound-healing (49), and is the rationale for the delivery fibrin-

enriched products to promote wound healing (50). However, the clearance of the fibrin-

containing provisional matrix is also a key step in late stages of wound-healing (49). Since

hepatic fibrosis and subcutaneous wound-healing qualitatively and quantitatively share many

similarities, it is possible this increase in fibrin(ogen) fragments during the late stage of recovery

from hepatic fibrosis represents a similar clearance of fibrin ECM from the provisional matrix.

Future studies should specifically investigate this possibility, especially given the use of

anticoagulants in fibrotic liver disease and cirrhosis.

As mentioned in the Introduction, collagenous ECM accumulation has been the major

focus in the field of hepatic fibrosis (2). Likewise, proteases involved in resolution of

collagenous ECM (e.g., MMP2 and MMP9) have also received a sizeable amount of attention

(38). Once the technical limitations to identifying peptide fragments of degraded proteins

informatically was overcome (44), the fragment sequence and putative cleavage site may yield

new information on proteases involved. The predicted protease that generated each peptide

was therefore determined by in silico analysis of the cleavage site using the online open-source

tool, Proteasix (www.proteasix.org) (29). Although several proteases with established roles in

fibrosis and/or resolution were identified (e.g., MMP2), several novel players were predicted by

the algorithm.

The most common predicted protease both 1 and 28 d after cessation of CCl_4 was

calpain1 (Capn1) (Figure 7). It should be noted that Capn1 and its isozyme Capn2 cannot be

delineated by substrate or activity assays, and expression of both followed a similar pattern in

liver after recovery from fibrosis. This finding was validated by investigating levels of the

intrinsic substrate for Capn1/2 (α-fodrin) and via an enzymatic assay (Figure 7B). The calpain

family are cysteine proteases that cleave a myriad of protein substrates. Although they are

generally localized intracellularly, they can be excreted when proteolytically activated (51).

Calpain activation has been identified as a key driver of inflammation in other diseases, such as

atherosclerosis and cardiac disease (51, 52). Moreover, calpain activation has been identified

to be involved in diseases of remodeling (i.e., fibrosis) via mechanisms that remain unclear (53).

Interestingly, the predicted substrates for calpain activation not only included key ECM proteins

[e.g., collagens and fibrin(ogens)], but also nuclear proteins, such as histones (Figure 7C),

which is in-line with previous findings in other organs (54) and the liver (30). Few studies have

identified calpain activation to play a role in hepatic injury. However, a recent study indicated

that CAPN2 is induced in human livers in HBV-induced fibrosis (30), and it was demonstrated

here that CAPN2 expression correlated with the severity of fibrosis in NASH (Figure 7C). The

mechanisms by which CAPN2 may mediate these effects are unclear, but work in other organ

systems suggest a role in not only ECM metabolism, but also in organelle damage [e.g., (51)].

These data suggest that the role of calpains in liver disease and fibrosis may be

underappreciated at this time.

Taken together, the results of this study indicate that remodeling of the liver during

recovery from fibrosis is a complex and highly coordinated process that extends well beyond the

degradation of the collagenous scar. A key role of MMP12, and by extension elastin, was

demonstrated, which is in-line with previous studies. Moreover, it was shown that even during

late stages of recovery (i.e., 28 D after CCl_4 cessation), there is significant enhanced turnover of

proteins. These results also indicate that analysis of the plasma degradome may yield new

insight into the mechanisms of fibrosis recovery, and by extension, new "theragnostic" targets.

Lastly, a novel potential role for calpain activation in the degradation and turnover of proteins

was identified.

Reference List

1. Seki E, Brenner DA. Recent advancement of molecular mechanisms of liver fibrosis. J Hepatobiliary Pancreat Sci 2015;22:512-518.

2. Arteel GE, Naba A. The liver matrisome - looking beyond collagens. JHEP Rep 2020;2:100115.

3. Iredale JP. Hepatic stellate cell behavior during resolution of liver injury. Semin Liver Dis 2001;21:427-436.

4. Campana L, Iredale JP. Regression of Liver Fibrosis. Semin Liver Dis 2017;37:1-10.

5. Shahid I, AlMalki WH, Hassan S, Hafeez MH. Real-world challenges for hepatitis C virus medications: a critical overview. Crit Rev Microbiol 2018;44:143-160.

6. Ramachandran P, Iredale JP, Fallowfield JA. Resolution of liver fibrosis: basic mechanisms and clinical relevance. Semin Liver Dis 2015;35:119-131.

7. Bonnans C, Chou J, Werb Z. Remodelling the extracellular matrix in development and disease. Nat Rev Mol Cell Biol 2014;15:786-801.

8. Friedman SL. Stellate cell activation in alcoholic fibrosis--an overview. Alcohol Clin Exp Res 1999;23:904-910.

9. Gressner OA, Weiskirchen R, Gressner AM. Evolving concepts of liver fibrogenesis provide new diagnostic and therapeutic options. Comp Hepatol 2007;6:7.

10. Pellicoro A, Aucott RL, Ramachandran P, Robson AJ, Fallowfield JA, Snowdon VK, Hartland SN, et al. Elastin accumulation is regulated at the level of degradation by macrophage metalloelastase (MMP-12) during experimental liver fibrosis. Hepatology 2012;55:1965-1975.

11. Ehling J, Bartneck M, Fech V, Butzbach B, Cesati R, Botnar R, Lammers T, et al. Elastin-based molecular MRI of liver fibrosis. Hepatology 2013;58:1517-1518.

12. Kisseleva T, Brenner D. Molecular and cellular mechanisms of liver fibrosis and its regression. Nat Rev Gastroenterol Hepatol 2021;18:151-166.

13. Massey VL, Dolin CE, Poole LG, Hudson SV, Siow DL, Brock GN, Merchant ML, et al. The hepatic "matrisome" responds dynamically to injury: Characterization of transitional changes to the extracellular matrix in mice. Hepatology 2017;65:969-982.

14. Rittié L. Method for Picrosirius Red-Polarization Detection of Collagen Fibers in Tissue Sections. Methods Mol Biol 2017;1627:395-407.

15. Kohli V, Madden JF, Bentley RC, Clavien PA. Calpain mediates ischemic injury of the liver through modulation of apoptosis and necrosis. Gastroenterology 1999;116:168-178.

16. Bergheim I, Guo L, Davis MA, Duveau I, Arteel GE. Critical role of plasminogen activator inhibitor-1 in cholestatic liver injury and fibrosis. J Pharmacol Exp Ther 2006;316:592-600.

17. Koch RG, Tsamis A, D'Amore A, Wagner WR, Watkins SC, Gleason TG, Vorp DA. A custom image-based analysis tool for quantifying elastin and collagen micro-architecture in the wall of the human aorta from multi-photon microscopy. J Biomech 2014;47:935-943.

18. Keshishian H, Addona T, Burgess M, Kuhn E, Carr SA. Quantitative, multiplexed assays for low abundance proteins in plasma by targeted mass spectrometry and stable isotope dilution. Mol Cell Proteomics 2007;6:2212-2229.

19. Merchant ML, Barati MT, Caster DJ, Hata JL, Hobeika L, Coventry S, Brier ME, et al. Proteomic Analysis Identifies Distinct Glomerular Extracellular Matrix in Collapsing Focal Segmental Glomerulosclerosis. J Am Soc Nephrol 2020;31:1883-1904.

20. Szklarczyk D, Gable AL, Nastou KC, Lyon D, Kirsch R, Pyysalo S, Doncheva NT, et al. The STRING database in 2021: customizable protein-protein networks, and functional characterization of user-uploaded gene/measurement sets. Nucleic Acids Res 2021;49:D605-d612.

21. Govaere O, Cockell S, Tiniakos D, Queen R, Younes R, Vacca M, Alexander L, et al. Transcriptomic profiling across the nonalcoholic fatty liver disease spectrum reveals gene signatures for steatohepatitis and fibrosis. Sci Transl Med 2020;12.

22. Pfister D, Núñez NG, Pinyol R, Govaere O, Pinter M, Szydlowska M, Gupta R, et al. NASH limits anti-tumour surveillance in immunotherapy-treated HCC. Nature 2021;592:450-456.

23. Bedossa P, Poynard T. An algorithm for the grading of activity in chronic hepatitis C. The METAVIR Cooperative Study Group. Hepatology 1996;24:289-293.

24. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol 2014;15:550.

25. von Montfort C, Beier JI, Kaiser JP, Guo L, Joshi-Barve S, Pritchard MT, States JC, et al. PAI-1 plays a protective role in CCl4-induced hepatic fibrosis in mice: role of hepatocyte division. Am J Physiol Gastrointest Liver Physiol 2010;298:G657-666.

26. Roderfeld M. Matrix metalloproteinase functions in hepatic injury and fibrosis. Matrix Biol 2018;68-69:452-462.

27. Hemmann S, Graf J, Roderfeld M, Roeb E. Expression of MMPs and TIMPs in liver fibrosis - a systematic review with special emphasis on anti-fibrotic strategies. J Hepatol 2007;46:955-975.

28. Sage EH, Gray WR. Evolution of elastin structure. Adv Exp Med Biol 1977;79:291-312.

29. Klein J, Eales J, Zurbig P, Vlahou A, Mischak H, Stevens R. Proteasix: a tool for automated and large-scale prediction of proteases involved in naturally occurring peptide generation. Proteomics 2013;13:1077-1082.

30. Feng R, Du W, Lui P, Zhang J, Liu Y. CAPN2 acts as an indicator of hepatitis B virus to induce hepatic fibrosis. J Cell Biochem 2020;121:2428-2436.

31. Grgurevic I, Bozin T, Madir A. Hepatitis C is now curable, but what happens with cirrhosis and portal hypertension afterwards? Clin Exp Hepatol 2017;3:181-186.

32. D'Ambrosio R, Aghemo A, Fraquelli M, Rumi MG, Donato MF, Paradis V, Bedossa P, et al. The diagnostic accuracy of Fibroscan for cirrhosis is influenced by liver morphometry in HCV patients with a sustained virological response. J Hepatol 2013;59:251-256.

33. Bachofner JA, Valli PV, Kroger A, Bergamin I, Kunzler P, Baserga A, Braun D, et al. Direct antiviral agent treatment of chronic hepatitis C results in rapid regression of transient elastography and fibrosis markers fibrosis-4 score and aspartate aminotransferase-platelet ratio index. Liver Int 2017;37:369-376.

34. Chan J, Gogela N, Zheng H, Lammert S, Ajayi T, Fricker Z, Kim AY, et al. Direct-Acting Antiviral Therapy for Chronic HCV Infection Results in Liver Stiffness Regression Over 12 Months Post-treatment. Dig Dis Sci 2018;63:486-492.

35. Dolmazashvili E, Abutidze A, Chkhartishvili N, Karchava M, Sharvadze L, Tsertsvadze T. Regression of liver fibrosis over a 24-week period after completing direct-acting antiviral therapy in patients with chronic hepatitis C receiving care within the national hepatitis C elimination program in Georgia: results of hepatology clinic HEPA experience. Eur J Gastroenterol Hepatol 2017;29:1223-1230.

36. Persico M, Rosato V, Aglitti A, Precone D, Corrado M, De Luna A, Morisco F, et al. Sustained virological response by direct antiviral agents in HCV leads to an early and significant improvement of liver fibrosis. Antivir Ther 2017.

37. Vispo E, Barreiro P, Del Valle J, Maida I, de Ledinghen V, Quereda C, Moreno A, et al. Overestimation of liver fibrosis staging using transient elastography in patients with chronic hepatitis C and significant liver inflammation. Antivir Ther 2009;14:187-193.

38. Friedman SL, Pinzani M. Hepatic Fibrosis 2022: Unmet Needs and a Blueprint for the Future. Hepatology 2021.

39. Ramachandran P, Pellicoro A, Vernon MA, Boulter L, Aucott RL, Ali A, Hartland SN, et al. Differential Ly-6C expression identifies the recruited macrophage phenotype, which orchestrates the regression of murine liver fibrosis. Proc Natl Acad Sci U S A 2012;109:E3186-3195.

40. Zhao W, Yang A, Chen W, Wang P, Liu T, Cong M, Xu A, et al. Inhibition of lysyl oxidase-like 1 (LOXL1) expression arrests liver fibrosis progression in cirrhosis by reducing elastin crosslinking. Biochim Biophys Acta Mol Basis Dis 2018;1864:1129-1137.

41. Chen W, Yan X, Xu A, Sun Y, Wang B, Huang T, Wang H, et al. Dynamics of elastin in liver fibrosis: Accumulates late during progression and degrades slowly in regression. J Cell Physiol 2019;234:22613-22622.

42. Dhital B, Durlik P, Rathod P, Gul ENF, Wang Z, Sun C, Chang EJ, et al. Ultraviolet radiation reduces desmosine cross-links in elastin. Biochem Biophys Rep 2017;10:172-177.

43. Jormsjö S, Ye S, Moritz J, Walter DH, Dimmeler S, Zeiher AM, Henney A, et al. Allele-specific regulation of matrix metalloproteinase-12 gene activity is associated with coronary artery luminal dimensions in diabetic patients with manifest coronary artery disease. Circ Res 2000;86:998-1003.

44. Schrader M, Schulz-Knappe P, Fricker LD. Historical perspective of peptidomics. EuPA Open Proteomics 2014;3:171-182.

45. Menschaert G, Vandekerckhove TT, Baggerman G, Schoofs L, Luyten W, Van Criekinge W. Peptidomics coming of age: a review of contributions from a bioinformatics angle. J Proteome Res 2010;9:2051-2061.

46. Greening DW, Kapp EA, Simpson RJ. The Peptidome Comes of Age: Mass Spectrometry-Based Characterization of the Circulating Cancer Peptidome. Enzymes 2017;42:27-64.

47. Lee KC, Hsu WF, Hsieh YC, Chan CC, Yang YY, Huang YH, Hou MC, et al. Dabigatran Reduces Liver Fibrosis in Thioacetamide-Injured Rats. Dig Dis Sci 2019;64:102-112.

48. Beier JI, Luyendyk JP, Guo L, von Montfort C, Staunton DE, Arteel GE. Fibrin accumulation plays a critical role in the sensitization to lipopolysaccharide-induced liver injury caused by ethanol in mice. Hepatology 2009;49:1545-1553.

49. Kearney KJ, Ariëns RAS, Macrae FL. The Role of Fibrin(ogen) in Wound Healing and Infection Control. Semin Thromb Hemost 2021.

50. Yu P, Zhong W. Hemostatic materials in wound care. Burns Trauma 2021;9:tkab019.

51. Zhang M, Wang G, Peng T. Calpain-Mediated Mitochondrial Damage: An Emerging
Mechanism Contributing to Cardiac Disease. Cells 2021;10.

52. Miyazaki T, Miyazaki A. Dysregulation of Calpain Proteolytic Systems Underlies
Degenerative Vascular Disorders. J Atheroscler Thromb 2018;25:1-15.

53. Letavernier E, Zafrani L, Perez J, Letavernier B, Haymann JP, Baud L. The role of calpains
in myocardial remodelling and heart failure. Cardiovasc Res 2012;96:38-45.

54. Arnandis T, Ferrer-Vicens I, Torres L, García C, Garcia-Trevijano ER, Zaragoza R, Viña
JR. Differential functions of calpain 1 during epithelial cell death and adipocyte
differentiation in mammary gland involution. Biochem J 2014;459:355-368.

55. Naba A, Clauser KR, Ding H, Whittaker CA, Carr SA, Hynes RO. The extracellular matrix:
Tools and insights for the "omics" era. Matrix Biol 2016;49:10-24.

Table 1-Clusters identified by StringDB to be enriched in the degradome

Cluster Number	Enrichment (-log10PV)	Top GO term (-log10PV)	Genes in Cluster
1	15/37 (>16)	high-density lipoprotein particle GO:0034364 (9.0)	Aamdc, Apoa2, Apoa4, Apoc1, Apoc3, Apoe, Arpp19, Enthd2, Eps8l3, F7, Fga, Fgb, Hnrnpdl, Hpx, Inadl, Pah, Pard3, Saa2, Serpina1d, Tat, Tmbim6, Wipf1
2	12/37 (>16)	nucleosomal DNA binding GO:0031492 (5.7)	Aco1, Bach2, Clrn1, Cyp2c29, Cyp2c37, Cyp2c54, Ddx3y, Emilin1, Gck, Gsk3a, Gstp1, Gstp2, Gsx2, Hist1h1a, Hist1h1b, Hist1h1d, Hist1h1e, Hmgb1, Hmgb2, Hmgn2, Impdh2, Maf, Mug1, Pdcd7, Postn, Pygl, Samm50, Scn4a, Sipa1l2, Skida1, Slc1a5, Tuba3a, Tuba8, Tubal3, Wdr65, Wnk2, Zfp36l3
3	10/37 (4.8)	NuA4 histone acetyltransferase complex GO:0035267 (1.9)	Actb, Adra1d, Alyref, Arhgef33, Arid1b, Cbx2, Cbx4, Col8a1, Cryz, Cstf2t, Dctn2, Decr2, Ehd2, Eln, Eno1, Hist1h1c, Hnrnpk, Hrnr, Morf4l1, Morf4l2, Mug2, Nadk2, Nbea, Psmb1, Rbpr2, Ryr1, Sdr42e1, Serpina3k, Setd1a, Sspo, Tmem218, Tmem8b, Tmsb4x, Uox, Zcchc11, Zeb1, Zfp687
4	13/15 (11.6)	acetyl-CoA C-myristoyltransferase activity GO:0050633 (4.3)	Acaa1a, Acaa1b, Cat, Cct4, Ech1, Eef1a1, Eif2s2, Notch4, Rpl27, Rpl28, Rps14, Rps8, Scp2, Serbp1, Timm23
5	11/11 (>16)	structural constituent of cytoskeleton GO:0005200 (8.8)	Acaa1a, C3, C4b, Hspe1, Tagln2, Ttn, Tuba1a, Tuba1b, Tuba1c, Tuba4a, Vim, Wdr1
6	13/42 (>16)	collagen trimer GO:0005581 (15.6)	Adh1, Adipor2, Atp8a1, Bglap3 Bhmt2, C77080, Ccdc85a, Cebpa, Col11a1, Col13a1, Col1a1, Col1a2, Col4a1, Col4a3, Col4a4, Col5a1, Col5a2, Col6a1, Col6a4, Col7a1, Dab2, Gdf7, Gfi1, Hba-a2, Hdgfrp3, Hist1h2bl, Ilk, Inhbe, Itih4, Mast2, Mat1a, Mki67, Nckap5l, Neurl4, Npnt, Pcnxl3, Prpsap2, Rcor3, Sntb1, Tmem41a, Ufd1l, Ybx3
7	16/21 (>16)	nucleosome GO:0000786 (15.9)	Carkd, Dhx9, H2afv, H2afz, H3f3a, Hist1h2ba, Hist1h2bb, Hist1h2bc, Hist1h2bh, Hist1h2bm, Hist1h2bp, Hist2h2aa2, Hist2h2bb, Hist3h2ba, Hpd, Polg2, Saa1, Set, Trp53bp1, Vav2, Ybx1

1   Enriched clusters were identified using StringDB (20), as described in Experimental Procedures. The top identified gene ontology
2   (GO) term for each cluster are also shown. Red parent proteins are defined as "Matrisome" by the matrisome project
3   (www.matrisomeproject.mit.edu) (55). See also Figure 6.

	Figure Legends

Figure 1. Recovery from liver injury and fibrosis.

Panel A: Mice were administered CCl_4 for 4 weeks to induce hepatic fibrosis as described in

Materials and Methods. Animals were sacrificed 1, 3, 7, 14 and 28 days after cessation of CCl4

to track fibrosis resolution. Panel B: Representative photos of livers and photomicrographs

(100×) of stained liver sections are shown.   Hematoxylin and eosin (H&E) shows general

histology, while Sirius red brightfield (BF) and polarized light (PL) show fibrosis.

Figure 2. Effect of CCl_4 and recovery on indices of liver injury and fibrosis.

Experimental groups and approaches are as described in Experimental Procedures. Panel A

shows plasma transaminase (AST and ALT; IU/L) levels and hepatic mRNA expression of key

indices of fibrosis (*Col1a1*, *Acta2*, *Lox1*, *Lox2* and *P4htm*; Log2FC) are shown. Panel B shows

representative zymographs demonstrating gelatin lysis by MMP-2 and -9. Panel C shows

densitometric analysis of zymography (left; FC) and hepatic expression of key tissue inhibitors

of metalloproteinases (*Timp1*, *Timp2* and *Timp3*; right; Log2FC). Quantitative data are reported

as means ± SEM (n=4-6). [a]P <0.05 compared with absence of CCl_4 by ANOVA using

Bonferroni's post hoc test.

Figure 3. Hepatic expression of MMPs and elastin after fibrosis and recovery.

Experimental groups and approaches are as described in Experimental Procedures. Panel A

shows hepatic mRNA expression of key indices of fibrosis of all hepatic matrix

metalloproteinases (Mmps) both as Log2FC (left) and copy number (right). Panel B shows

elastin protein in the liver by Western blot (upper), immunofluorescence (100×; middle) and

multiphoton microscopy (MPM, 100×, lower). Elastin immunofluorescence is depicted as intese

green staining against the background of hepatic tissue autofluorescence (light green). MPM

images depict elastin (green), collagen (blue) and stained nuclei (red). Panel C summarized

densitometric quantitation of elastin by Western blot (FC). Quantitative data are reported as

means ± SEM (n=4-6). [a]P <0.05 compared with absence of CCl_4 by ANOVA using Bonferroni's

post hoc test.

Figure 4. The effect of MMP-408 on recovery from liver injury and fibrosis.

Experimental groups and approaches are as described in Experimental Procedures. The effect

of MMP-408 on liver transaminase levels (Figure 3A, left) and fibrosis genes (Figure 3A ,middle

and right) during recovery were determined. The effect of MMP-408 on resolution of fibrosis

was assessed by Sirius Red staining (100×) (Figure 3B). Intensity of Sirius red staining was

assessed by image analysis (Figure 3C, left). Quantitative data are reported as means ±SEM,

(n = 8-10). [a]P <0.05 compared with absence of CCl_4; [b]P <0.05 compared with vehicle group by

ANOVA using Bonferroni's post hoc test.

Figure 5. Effect of fibrosis and recovery on the plasma degradome.

Experimental groups and approaches are as described in Experimental Procedures. Plasma

peptides were measured by LC-MS/MS, as described in experimental procedures and

compared naive control mice (Log2FC). Panel A shows the pattern of changes in peptides over

the course of the recovery period that were increased at 28 days. Panel B compares the

change in peptides after 28 days of recovery, versus those a 1 day. Panel C is a Venn diagram

comparing the peptides increased at 28 days of recovery to those increased at 1 day of

recovery. Box and whisker plots depict the range, median and interquartile range of the

changes of the degraded proteins over the period of recovery (Panel A).

Figure 6.

StringDB clustering analysis of peptides significantly increased 28 days after cessation of CCl_4.

Red circled parent proteins are defined as "Matrisome" by the matrisome project

(www.matrisomeproject.mit.edu) (55). Cluster groups are summarized in Table 1.

Figure 7. Expression and activity of predicted protease from ProteaSix.

Experimental groups and approaches are as described in Experimental Procedures. ProteaSix

was used to predict proteases that formed these peptides (Figure 6A). Panel B (upper) depicts

western blot analysis of α-fodrin. The 240 kDa band represents intact α-fodrin, while the 150

and 120 kDa bands represent calpain-cleaved fragments. Densitometric analysis is also shown

(lower, right). The expression (lower, left) of *Capn1* and *Capn1* were determined by real-time

rtPCR. Panel 6C summarizes the type of parent proteins identified a potential substrates for

Capn1/2 cleavage by ProteaSix, organized by subcellular localization. The relative size of the

pie charts (1 D vs 28 D) is reflective of the amount of peptides identified at each timepoint.

Quantitative data are reported as means ±SE, n=4-6. [a]P <0.05 compared with absence of CCl_4

by ANOVA using Bonferroni's post hoc test.

Figure 8. Proposed role of MMP-12 and elastin during recovery from liver fibrosis

Hepatic activity of MMPs was robustly induced; but MMP-12 returned to baseline after recovery

from fibrosis. Elastic fibers, which are substrates of MMP-12, are also critical in fibrotic ECM

formation, and appeared after CCl_4 administration. Fibrosis resolved and histology was nearly

normal after 28 D recovery, but elastin ECM still remained elevated. Mature tropoelastin chains

are connected with desmosine cross-links. Newly synthesized immature elastin lacks this post-

1   translational modification. As recovery progresses, total elastin decreases, but desmosine

2   content increases; interestingly, blocking MMP-12 activity prevented both of these phenomena.

Days of Recovery after CCl$_4$

Figure 2-Sato et al.

A

B

| | control | 1 day | 3 days | 7 days | 28 days |

Elastin

GAPDH

IF

MPM

C

Figure 3-Sato et al.

A

B

1 Day 7 Days 28 Days

Vehicle

MMP408

C

10   Figure 5-Sato et al.

10   Figure 6-Sato et al.

Figure 7-Sato et al.

11   Figure 8-Sato et al.