Engineering *Agrobacterium tumefaciens* adhesion to target cells

Xavier Pierrat, Alix Pham, Jeremy P. H. Wong, Zainebe Al-Mayyah and Alexandre Persat*

Institute of Bioengineering and Global Health Institute, School of Life Sciences, Ecole

Polytechnique Fédérale de Lausanne, CH-1015 Lausanne, Switzerland

*Corresponding author: alexandre.persat@epfl.ch

Abstract

Agrobacterium tumefaciens is a plant pathogen commonly repurposed for genetic
modification of crops. Despite its versatility, it remains inefficient at transferring DNA to many
hosts, including to animal cells. Like many pathogens, physical contact between
A. tumefaciens and host cells promotes infection efficacy. Thus, improving the strength and
specificity of *A. tumefaciens* to target cells has the potential for enhancing DNA transfer for
biotechnological and therapeutic purposes. Here we demonstrate a methodology for
engineering genetically-encoded exogeneous adhesins at the surface of *A. tumefaciens*. We
identified an autotransporter gene we named Aat, that is predicted to show canonical β-barrel
and passenger domains. We engineered the β-barrel scaffold and linker (Aat_β) to display
synthetic adhesins susceptible to rewire *A. tumefaciens* to alternative host targets. As a proof
of concept, we leveraged the versatility of a VHH domain to rewire *A. tumefaciens* adhesion
to yeast and mammalian hosts displaying a GFP target receptor. Finally, to demonstrate how
synthetic *A. tumefaciens* adhesion can improve transfer to host cells, we showed improved
protein translocation into HeLa cells using a sensitive split luciferase reporter system.
Engineering *A. tumefaciens* adhesion has therefore a strong potential in generating complex
heterogeneous cellular assemblies and in improving DNA transfer efficiency against non-
natural hosts.

Introduction

A. tumefaciens is a soil bacterium able to transfer DNA fragments tens of kilobases-long to

plant host cells. In its natural environment, *A. tumefaciens* transfers a pathogenic DNA

fragment called transferred-DNA (T-DNA). *A. tumefaciens* has been repurposed into a potent

gene delivery tool for a broad range of plants, yeasts and fungi[1–3]. Using disarmed strains

lacking pathogenic T-DNA, plant engineers introduce new genes in target organisms such as

maize, rice and wheat[4–6]. Genetic modifications include random integration or targeted

editing using zinc-finger nucleases or CRISPR/Cas9 [4].

A. tumefaciens uses the type IV secretion (T4SS) system to inject DNA into host cells.

The T4SS functions upon host surface contact, so that attachment to target cells promotes T-

DNA transfer[7–9]. For synthetic application, forcing cell-cell interaction, for instance by

wounding the plant or by using syringe- or vacuum-driven agroinfiltration, is usually

recommended[10]. The addition of extracellular cellulose increases bacterial adhesion to

improve T-DNA transfer efficiency to recalcitrant plant cells[11]. Altogether, these studies

suggest that increased bacterial adhesion favors T-DNA transfer. However, our

understanding of how *A. tumefaciens* adheres to host plant surfaces remains incomplete[8,9].

Despite this knowledge gap, engineering adhesion has the potential to improve gene delivery

to plants or alternative targets, for example by broadening the host range of *A. tumefaciens.*

In Gram-negative bacteria, outer-membrane proteins called autotransporters help display

functional proteins moieties at the cell surface, including adhesins[12]. Autotransporters are

promising candidate scaffolds for surface display of exogenous proteins[13]. They belong to the

type V secretion system (T5SS) family, and are composed of a β-barrel domain fused to a

passenger domain with an α-helical linker[14]. The β-barrel anchors the autotransporter to the

outer membrane. The passenger domain translocates by traversing the β-barrel, thereby

exposing itself at the cell surface and conferring function. In contrast to other secretion

systems, autotransporters are expressed as a single protein. This allows for substitution of

the passenger domain to other peptides as a way to switch autotransporter function. For

example, the *Pseudomonas aeruginosa* EstA autotransporter has been used to display a

variety of lipases, with applications in whole-cell biocatalysis and screening of enzyme

libraries[14].

Adhesion plays a crucial function in host microbe interactions, which can be repurposed

for synthetic applications. Fusing an alternative receptor to an autotransporter is a

convenient way to engineer bacterial adhesion to a specific host target. For example, fusions

of non-natural receptors to the *E. coli* intimin autotransporter efficiently rewires adhesion to a

wide range of host cells. Specifically, antigen-binding domain from camelid heavy-chain

antibodies (variable heavy chain of heavy-chain antibodies, VHH) fused to intimin provide

adhesion properties to target ligands[15]. Utilizing VHH has many advantages as they are

single-chained, short (less than 130 amino acids-long) and have a robust structure[16].

Moreover, they are more efficiently displayed at the surface of bacteria than human single-
chain fragment variables (scFv)[17]. Finally, screening for VHH forms with affinity to new
targets is streamlined, for example via phage display starting from naïve DNA libraries[18].
Using the intimin-VHH display system, Fraile *et al.* targeted *P. putida* to abiotic surface
coated with the target antigen[19]. Salema *et al.* screened libraries against cancer biomarkers
by selecting bacteria binding to live mammalian cells[20]. Glass *et al.* created an adhesin
toolbox that enables the assembly of complex *E. coli* communities[21]. Finally, Ting *et al.*
performed targeted bacterial killing using programmed adhesion that directs toxin injection[22].
Thus, synthetic adhesion can confer modularity and specificity in assembling bacterial
consortia or attaching to alternative hosts[23,24].

Here, we aimed at engineering *A. tumefaciens* adhesion to improve protein delivery to
alternative targets, with the ultimate goal to program DNA transfer into alternative hosts.
While using an intimin scaffold is intuitive, it is not a practical solution. Intimin has a topology
belonging to the reverse autotransporter family, also called type Ve secretion system[25–27].
A. tumefaciens C58 does not possess any annotated reverse autotransporter, and
consequently likely does not express the right variants of the associated chaperones that
enable proper insertion in the outer membrane and/or translocation of the passenger
domain[25]. Therefore, we explored alternative autotransporters in order to engineer synthetic
adhesins at the surface of *A. tumefaciens*.

Here, we engineered a modular synthetic adhesin display system for *A. tumefaciens*. To
achieve this, we investigated the putative autotransporter Atu5364 which we renamed Aat
(as <u>A</u>grobacterium <u>a</u>uto<u>t</u>ransporter) as candidate scaffold for adhesin display. We
successfully repurposed Aat to display multiple functional receptors targeting a variety of
ligands, including VHH, lectins and arginine-glycine-aspartic acid (RGD) peptide. We then
demonstrated that displaying a nanobody-based VHH anti-GFP by Aat fusion strongly
improved *A. tumefaciens* attachment to alternative hosts including yeast and mammalian
cells displaying GFP. Finally, and as a proof of concept, we provide preliminary evidence that
synthetic display of VHH improves transfer of type IV secretion system proteins to
mammalian host cells.

Results

Identification of an A. tumefaciens autotransporter scaffold

The genome of *A. tumefaciens* is annotated with two uncharacterized autotransporters of the

type Va secretion system (T5aSS), namely *atu5354* and *atu5364*, both located on the cryptic

megaplasmid pAtC58. We only found a signal peptide encoded within the sequence of

atu5364, but not in *atu5354,* which we therefore disregarded as a potential candidate

scaffold (Figure S1). We then modeled <u>A</u>grobacterium <u>a</u>uto<u>t</u>ransporter Atu5364 (Aat) using

RoseTTaFold and obtained an estimate of its three dimensional conformation (Figure

1A)[28,29]. The predicted structure shows a C-terminal scaffold (β-barrel and α-helix) holding a

passenger formed of repeated parallel β-strand repeats organized in a β-solenoid, a

prevalent structure in T5aSS autotransporters[30]. Aat harbors a proline-rich linker between the

scaffold and the folded region of the passenger domain. A potential function for this linker is

to remain unfolded and extend to help the passenger domain reach distant targets, for

example in attachment[31]. We did not identify a clear homolog for the passenger domain, but

our analysis points toward a function in adhesion (Table S1)[32–37]. Altogether, the predicted

scaffold, signal peptide and long linker make Aat a promising candidate for the

autotransporter-based display of non-natural passenger domains.

Protein display in A. tumefaciens

We then moved on to use Aat as a scaffold for protein display in *A. tumefaciens*. We first

fused an HA-tag between the endogenous passenger domain and the linker domain. We

expressed the fusion under a cumic acid-inducible promoter to decouple expression from

virulence[38]. After staining with FITC-conjugated anti-HA antibodies, these cells showed low

expression with patchy localization at the cell surface (Figure S2B). We noticed that the

native passenger of Aat contains a disulfide bond located at the N-terminal end of the native

passenger domain, which may prevent translocation (Figure 1A)[39]. We generated a cysteine-

free mutant version of the passenger domain with the HA-tag fusion. Staining and

microscopy showed that expression levels of the cysteine-free mutant were much stronger

than wild-type (WT) and that the localization was uniform along the bacterium surface (Figure

S2C).

To demonstrate the capacity of Aat in displaying alternative functional passenger

domains, we fused a VHH domain targeting GFP to the Aat β-barrel scaffold with its proline-

rich linker (which we call Aat$_\beta$). We initially screened a variety of fusion strategies, which

consisted in either replacing parts of the passenger domain (amino acids 35-160, 35-512 or

161-512) or in introducing VHH in front, within or after the folded domain of the passenger (at

C35, D161 or A513) (Supplementary figure S3A). The constructs were expressed from

binary vectors under the control of the pVirE promoter (Supplementary table 2)[40]. Incubating

bacteria expressing these constructs with GFP shows that the scaffold is properly folded and

inserted in the outer membrane. However, these constructs caused frequent cell death

(Figure S3B, C). We hypothesize that VHH is folded but accumulates in the periplasm,

causing toxicity.

We noticed that VHH contains a disulfide bond that again could prevent passenger

domain translocation[39]. Mutating cysteines from VHH anti-GFP has little effect on affinity[41].

We therefore cloned a disulfide-free VHH by site-directed mutagenesis of C24A and C98V.

We then displayed the disulfide-free VHH anti-GFP (VHH$_{GFP}$) by fusion to Aat$_\beta$

(VHH$_{GFP}$_Aat$_\beta$) in *A. tumefaciens* and compared it to the native VHH. We expressed this

construct in a strain constitutively expressing cytoplasmic mScarlet as viability reporter. After

expressing and staining VHH$_{GFP}$_Aat$_\beta$ with GFP, we observed enhanced fluorescent signal at

the bacterium surface compared to the native, disulfide-containing version, while not

affecting bacterial viability (Figure 1B, Figure S3C, D). Of note, the distribution of the GFP

signal seemed more concentrated at the narrower side of the bacteria, which might be due to

intrinsic, yet unknown, properties of the Aat$_\beta$ scaffold. In summary, Aat$_\beta$ efficiently displays

functional VHH$_{GFP}$ at the surface of live bacteria.

Figure 1: Protein display in *A. tumefaciens* by fusion to Aat$_\beta$. (A) Structure of Aat modeled by
RoseTTaFold. Aat is a T5aSS autotransporter with a C-terminal β-barrel inserted at the outer
membrane. A passenger domain consists of β-strands capped by a disulfide bond (C35-C48, S-S,
framed in red). Aat$_\beta$ is composed of the β-barrel, α-helix and proline-rich linker (p-rich). SP = signal
peptide. (B) Fusion of cysteine-free VHH anti-GFP (VHH$_{GFP}$) to Aat$_\beta$ is displayed at the surface of
A. tumefaciens. (i) Diagram of the construct placed under a VirE promoter (pVirE). (ii) Illustration of
A. tumefaciens expressing VHH$_{GFP}$_Aat$_\beta$ (red) binding to GFP (green stars). (iii) *A. tumefaciens*
expressing VHH$_{GFP}$_Aat$_\beta$ binds diffusible GFP at its surface. Cytosolic mScarlet highlights the
surface GFP signal. (C) Display of arginine-glycine-aspartate (RGD), *P. aeruginosa* LecA, LecB
and disulfide-free *E. coli* FimH at the surface of *A. tumefaciens*. The constructs are placed under a
pVirE or a synthetic cumic acid-inducible promoter (pCuO) with an HA-tag allowing for
visualization. Bacteria were stained with anti-HA antibody conjugated with fluorescein (FITC). (D)
Negative control: overexpression of Aat wild-type under a VirE promoter, followed by staining with
GFP. Bacteria have aberrant shapes and are not stained with GFP. Bars, 2 µm.

To illustrate the modularity of the display scaffold, we sought to display alternative

receptors. We turned to the arginine-glycine-aspartate (RGD) motif which targets integrin

ligands, and two disulfide-free lectins from *Pseudomonas aeruginosa*, namely LecA and

LecB which respectively bind to galactose and fucose[42–45]. We fused each of the receptors to

Aat_β. In each, we also inserted an HA-tag between the adhesin and Aat_β for characterization

of display. Upon induction and staining with a green fluorescent anti-HA antibody, all

constructs showed fluorescence at their periphery. All maintained WT levels of a

constitutively-expressed cytosolic mScarlet demonstrating cell viability (Figures 1C, S4A-E).

To further demonstrate the process of engineering adhesins for Aat_β-based display, we

displayed the FimH fimbrial tip of uropathogenic *E. coli*. FimH harbors a disulfide bond

between C3 and C44. Mutations of one of FimH cysteines to a serine does not affect affinity

to mannose at no- and low shear rates[46]. Hence, we generated a FimH(C3S) mutant that is

deficient in forming disulfide bonds and compared the display efficiency with the WT. Like for

VHH display, the removal of the disulfide bond greatly improved display (Figures 1C, right

panel and S4F and G). Altogether, this shows that Aat_β is highly modular for passenger

display and likely can display a wide variety of short peptides, tags, and other disulfide-free

β-stranded proteins. For clarity and demonstration purposes, we subsequently focused on

the display of VHH_{GFP} adhesin as it enables quantitative control at the adhesin level, but also

at the host ligand level.

To improve the genetic stability of the constructs, we generated markerless genomic

insertion of $VHH_{GFP}\_Aat_\beta$ and compared it to plasmid-based expression. More specifically,

we integrated $VHH_{GFP}\_Aat_\beta$ at the *virE2* locus on the Tumor-inducing (Ti) megaplasmid

(Supplementary figure S5). We next induced bacteria with increasing concentrations of

inducer (acetosyringone titration) in both of the plasmid-borne and chromosomally-integrated

version. We stained the bacteria with recombinant GFP to quantify the display efficiency of

$VHH_{GFP}\_Aat_\beta$ (Figure 2). Both Ti plasmid-integrated and binary vector-based displayed

VHH_{GFP} at high and very high levels, a small difference probably likely due to plasmid copy

number. In conclusion, both binary vector-based and Ti plasmid-integrated versions are

compatible with VHH_{GFP} display.

Figure 2: pVirE is a potent inducible promoter for VHH$_{GFP}$_Aat$_β$ on binary vector and Tumor-inducing megaplasmid. (A) Acetosyringone induction of VHH$_{GFP}$_Aat$_β$ from a VirE promoter on a binary vector. *A. tumefaciens* transformed with pVirE – VHH$_{GFP}$_Aat$_β$ (pXP213, table S2) was induced with increasing concentrations of acetosyringone and stained with recombinant eGFP. GFP fluorescence intensity scale is identical between images. (B) Mean bacterial GFP intensity as a function of inducer concentration. (C) VHH$_{GFP}$_Aat$_β$ can also be expressed from a stable insertion in the disarmed tumor-inducing plasmid (*A. tumefaciens VHH$_{GFP}$_Aat$_β$::ΔVirE2*, see table S3). Bars, 1 μm.

Attachment of engineered A. tumefaciens to yeast

Synthetic adhesin display opens the possibility of engineering bacteria with emerging

functions, such as the assembly of complex multicellular structures, the destruction of cancer

cells, or can be applied in investigations of adhesion to host cells[21,47]. With these applications

in mind, we explored the potential of *A. tumefaciens* with synthetic Aat$_β$ display in providing

attachment to non-plant hosts. Forcing bacteria-host contact improves Agrobacterium-

mediated transformation of yeast cells[1,48]. We therefore first focused on testing whether the

synthetic VHH$_{GFP}$_Aat$_β$ construct promotes attachment to yeast cells. By fusing GFP to cell-

wall-anchoring proteins, we constructed a GFP-displaying *S. cerevisiae* in the strain eby100,

commonly used in yeast display libraries (Figure 3A)[49]. Then, we separately induced GFP-

display in yeast and expression of VHH$_{GFP}$_Aat$_β$ in *A. tumefaciens*. We mixed the two strains

and imaged the consortium by confocal microscopy. Confocal sections showed that bacteria

displaying VHH$_{GFP}$ strongly bound to the yeast cell wall, to the extent of imprinting their shape

into the cell walls (Figure 3B). We then compared the binding to GFP-displaying yeast in the

absence of VHH$_{GFP}$_Aat$_\beta$ and observed a decrease in the number of bacteria bound per

yeast cell (Figure 3C). In order to unambiguously attribute binding to the specificity of VHH$_{GFP}$

to GFP, we compared this to GFP-negative yeast and *A. tumefaciens* displaying a cysteine-

free VHH anti-mCherry (VHH$_{mCherry}$). The number of bacteria attached per yeast cell only

increased when the VHH$_{GFP}$_Aat$_\beta$ construct was expressed and when yeast displayed GFP

(Figure 3D). This shows that our scaffold can rewire *A. tumefaciens* to the yeast cells,

opening the possibility of generating complex interkingdom assemblies. The low binding of

A. tumefaciens displaying VHH$_{mCherry}$ additionally demonstrates the specificity of synthetic

adhesion.

Figure 3: Synthetic adhesion of *A. tumefaciens* VHH$_{GFP}$_Aat$_\beta$ to GFP-displaying yeast cells.
(A) *A. tumefaciens mScarlet* VHH$_{GFP}$_Aat$_\beta$ binds to GFP displayed at the surface of *S. cerevisiae.*
(B) Representative confocal images of *A. tumefaciens mScarlet* VHH$_{GFP}$_Aat$_\beta$ attaching to GFP-displaying yeast cell. In the lower panels, bacterial shape imprints are visible. (C) 3D visualization of *A. tumefaciens mScarlet* (orange) binding to *S. cerevisiae* displaying GFP (green) in the absence or presence of VHH$_{GFP}$ display. (D) Number of *A. tumefaciens mScarlet* cells attached per yeast cell [no VHH: empty vector pFGL815, αGFP: pVirE - VHH$_{GFP}$_Aat$_\beta$, αmCh: pVirE - cysteine-free VHH anti-mCherry (VHH$_{mCherry}$_Aat$_\beta$), see table S2). Bars represent the mean of biological triplicates. Statistical tests: two-way ANOVA and Sidak *post-hoc* test (****, P<10^{-4}).

To demonstrate the capability of the synthetic Aat$_\beta$-based display for therapeutic purposes,

we tested how engineered *A. tumefaciens* could specifically bind human cells. To achieve

this, we employed HeLa cells displaying GFP at their surface by fusion to a CD80

transmembrane domain[47]. We incubated HeLa GFP-display with bare or displaying VHH$_{GFP}$

A. tumefaciens. After washing of unbound bacteria, we acquired confocal images to quantify

the average number of bacteria per HeLa cell. We observed an increase in the number of

bound bacteria when HeLa displayed GFP and *A. tumefaciens* displayed VHH$_{GFP}$

simultaneously. However, there was no increase in binding when only the receptor or its

ligand were expressed (Figure 4Ai-iii and B). By preincubating *A. tumefaciens* with GFP, we

were able to prevent binding to host cells, demonstrating specificity and the potential for

external inhibition of adhesion (Figure 4iv and B). In addition, *A. tumefaciens* displaying

VHH$_{mCherry}$ could not attach to HeLa displaying GFP, further demonstrating specificity (Figure

4Av and B). These results held across cells lines, as *A. tumefaciens* attached in a similar

manner to GFP-displaying HEK293T cells (Figure S6).

Figure 4: Synthetic adhesins promote specific *A. tumefaciens* binding to HeLa cells. (A) Maximum intensity projections of *A. tumefaciens mScarlet* binding to HeLa cells. (i) no VHH$_{GFP}$_Aat$_\beta$ and GFP-display, (ii) VHH$_{GFP}$_Aat$_\beta$ and no GFP-display, (iii) VHH$_{GFP}$_Aat$_\beta$ and GFP display, (iv) VHH$_{GFP}$_Aat$_\beta$ and GFP-display prevented by soluble GFP and (v) VHH$_{mCherry}$_Aat$_\beta$ and GFP display. All images are overlays of GFP (green) and bacteria (orange) except (ii) where we use brightfield to visualize non-fluorescent HeLa. (B) Average number of bacteria per HeLa cells. HeLa were uninduced (GFP -) or induced for GFP display (GFP +), *A. tumefaciens mScarlet* was transformed with an empty vector (VHH -), pVirE – VHH$_{GFP}$_Aat$_\beta$ (αGFP) or pVirE – VHH$_{mCherry}$_Aat$_\beta$ (αmCh). Dark green set of bars show conditions where soluble eGFP was added to saturate VHH$_{GFP}$ receptors. Each symbol is a biological replicate, bars represent the means of four biological replicates. Statistical tests: two-way ANOVA and Sidak post hoc test (****, P<10^{-4}).

Monitoring VirE2 transfer using split NanoLuc

We next aimed at targeted delivery into mammalian cells using synthetic adhesion. Helper

proteins VirE2 and VirD2 increase T-DNA transfer efficiency in experiments involving artificial

T-DNA transfection[50]. Following binding to the target cell and T4SS assembly, VirD4 actively

recruits helpers proteins such as VirD2 and VirE2 to the T4SS machinery for injection into

the target cell's cytosol. Consequently, the cytosol of the target cell is the first compartment

through which the T-DNA and helper proteins navigate.

To sensitively monitor protein transfer, we employed a split luciferase (NanoLuc)

approach where the translocated protein is fused to one fragment (HiBit) and the host cell

expresses a complementary fragment (LgBit). Successful injection into the host leads to

NanoLuc complementation and hence to an increase in luminescence signal from the host[51−]

[53]. VirE2 is the most abundant Vir protein and is often fused to the short fragment of split

fluorescent proteins by insertion into an internal permissive site[54,55]. We therefore generated

an *A. tumefaciens* strain expressing a HiBit::VirE2 internal fusion under the VirE promoter

(Figure 5A). As reporter cell line, we further engineered the GFP-displaying HeLa cell line to

constitutively express the LgBit fragment. We controlled LgBit complementation by

transiently transfecting cells with a plasmid driving HiBit::VirE2 expression. We measured a

strong increase in luminescence signal, suggesting that HiBit internally fused to VirE2

efficiently complements LgBit (Figure S7).

We next challenged HeLa GFP display expressing LgBit with *A. tumefaciens* expressing

$VHH_{GFP}\_Aat_\beta$ and HiBit::VirE2. We first modulated either inducer concentration or multiplicity

of infection. Luciferase signal intensity increased with induction levels at constant multiplicity

of infection (Figure 5B), and with multiplicity of infection at constant induction (Figure 5C).

This implies that increased in bacterial virulence or increased in number of bacteria both

increase VirE2 transfer to HeLa cells.

Figure 5: VHH_GFP_Aat_β improves VirE2 transfer to HeLa cells. (A) Overview of the split luciferase (NanoLuc) system: NanoLuc is split in two fragments, LgBit and HiBit. *A. tumefaciens VHH_GFP_Aat_β::ΔVirE2* expresses an internal fusion of HiBit into VirE2. The target host cell expresses LgBit. Upon successful VirE2 transfer, NanoLuc activity should be restored. (B) HiBit::VirE2 expressing bacteria were induced with increasing concentrations of acetosyringone and added to GFP-displaying HeLa cells expressing LgBit. (C) VirE2 transfer as a function of bacterial load, controlled by volume of culture added to GFP-displaying HeLa cells expressing LgBit. 100 µM inducer concentration. (B and C) Error bars represent the standard deviation of biological triplicates. (D) VHH_GFP_Aat_β promotes VirE2 transfer in a GFP-display dependent manner. Bacteria were induced with 100 µM acetosyringone and added to LgBit-expressing reporter HeLa cells with or without GFP ligand at their surface. Each bar represents the means of biological triplicates. Statistical tests: one-way ANOVA followed by Dunnett's post hoc test (**** $P<10^{-4}$, ** $P<0.01$, * $P<0.05$).

We then investigated the specificity of VirE2 transfer to the synthetic binding. Based on

the titrations (Figure 5B, C), we infected HeLa cells with 22 μL of bacterial culture induced

with 100 μM acetosyringone. We first verified that the signal was specific to HiBit::VirE2

translocation by measuring luminescence of a native VirE2 in the same system. We

measured only background luminescence levels, order of magnitude smaller than for the

HiBit::VirE2 fusion (Figure 5D). In addition, infecting HeLa with or without GFP display with

an *A. tumefaciens* strain lacking VHH adhesin display but expressing HiBit::VirE2 abolished

luminescence. These experiments demonstrate that VHH display is required for the increase

in VirE2 transfer. We however found one discrepancy: luminescence was still high—while

statistically significantly lower than when both adhesins are present—when VirE2 is

transferred to HeLa without expression of GFP receptors (Figure 5D). To explain this

discrepancy, we checked for the possibility of leaky GFP expression. We inhibited binding by

pre-incubating *A. tumefaciens* VHH with recombinant GFP or VHH to block adhesion from

bacterial or host side respectively. We included a control by adding recombinant mCherry,

which should not interfere with VHH in binding GFP on host cells. Luminescence decreased

about 2-fold compared to non-induced GFP-display, still partly explaining the GFP-unspecific

signal increase (Figure S8A).

We attribute the remaining signal to VirE2 transfer independent of adhesion through an

unknown mechanism. We therefore investigated the impact of the Aat_β scaffold on transfer.

We displayed either a disulfide-free VHH anti-mCherry, or an empty passenger domain, and

repeated the experiment. In the absence of specific adhesion to GFP, we observed a similar

VirE2 transfer as in the absence of GFP-display (Figure S8B, second to last columns). This

suggests that Aat_β has an intrinsic contribution to VirE2 transfer. We next verified if this

intrinsic contribution of Aat_β occurs before contact (e.g. lysis) or upon contact. By comparing

the signal of reporter cells onto which we applied either overnight supernatant, resuspended

bacteria, we observed that the signal increase was due to VirE2 translocation from bacteria

but not VirE2 secreted in the supernatant (Figure S8C). However, lysate of *A. tumefaciens*

expressing $VHH_{GFP}\_Aat_\beta$ and HiBit::VirE2 showed a strong increase in luminescence when

incubated with HeLa GFP (Figure S8C), which suggests that mammalian cells actively

uptake VirE2, by a mechanism that remains to be elucidated. One unifying hypothesis about

the non-specific fraction of VirE2 transfer would be that overexpression of Aat_β slightly

increases unspecific binding to target cells (Figure 4B), reaching a sufficient threshold for

VirE2 transfer, disregard of the specificity of the VHH (Figures 5D and S8B). In conclusion,

we demonstrated that synthetic adhesion promotes VirE2 transfer to host cells, a process

which is in part stimulated by Aat_β in an unspecific, ligand-independent manner.

Discussion

Here we investigated *A. tumefaciens* Aat autotransporter as a candidate scaffold for the

display of synthetic passenger domains, aiming at designing a custom, target-specific

adhesion system. We repurposed it by fusing the scaffold to lectins, RGD and a VHH. VHH

are extremely versatile and can be engineered towards almost any biomarker of interest[56].

As a proof of concept, we rewired *A. tumefaciens* using a cysteine-free VHH anti-GFP to

target soluble and cell-displayed GFP.

In animal cells, only low *Agrobacterium*-mediated T-DNA transfection efficiencies have

been reported[57,58]. In recalcitrant plants, several reports demonstrate a positive correlation

between adhesion and T-DNA delivery efficiency[11,59]. Hence, one possible explanation of the

limited progress in *Agrobacterium*-mediated delivery in mammalian cells might be the low

affinity of bacteria to such non-natural hosts. We showed that our novel Aat_β scaffold has a

strong potential in engineering *A. tumefaciens* adhesion to non-natural hosts, including

mammalian cells. Thus, our design opens the possibility to further examine the possibility of

A. tumefaciens-mediated DNA transfer to non-plant hosts. In addition, it provides synthetic

biologists with a novel tool to program adhesion of complex multispecies microbial consortia,

or the display of enzymes at the surface of other alphaproteobacteria that could be used as

additives features in biotechnology or bioremediation[14].

We next developed a highly sensitive split luciferase assay that enabled us to monitor

VirE2 delivery into mammalian cells. The VirE2 transfer in a cell-cell contact-dependent but

partly GFP-independent manner has yet to be resolved. Bacteria displaying VHH might

release VirE2 upon contact, potentially by VHH display-facilitated lysis. Like in plants, VirE2

could then trigger clathrin-mediated endocytosis[60]. Our results however provide a strong

basis for further engineering adhesion to improve *A. tumefaciens* DNA transfer to non-natural

host.

Improved *A. tumefaciens*-mediated protein delivery is of importance, as scientists

repurposed the T4SS of the bacterium for protein delivery and gene editing, as an alternative

to using T-DNA. As an example, Vergunst *et al.* fused VirE2 to the Cre recombinase and

leveraged minute amount of proteins transferred to the target cells. They monitored Cre-

mediated recombination in plant cells by conferring resistance to cells undergoing

recombination[61]. More recently, Cas9 fusions to the VirF peptide responsible for translocation

allowed Schmitz *et al.* to target both yeast and plant reporter cells expressing gRNA[62].

Consequently, future experiments could consist in trying such fusions in combination with the

adhesin display system to optimize helper proteins delivery at high throughput.

A. tumefaciens is an attractive candidate for gene delivery to human cells. The availability

of safe, efficient and precisely targeting gene delivery vectors is currently one of the main

limiting factors for gene editing therapies *in vivo*. Adeno-associated viruses (AAVs) are the

current gold standard for gene delivery. Due to their limited packaging capacity, AAVs

however remain a bottleneck for the development of therapies based on CRISPR/Cas9[63,64].

On the other hand, wild-type *A. tumefaciens* injects T-DNA of 25 kb, a size that can easily

accommodate several times the CRISPR machinery and repair templates. We thus

anticipate that, granted important engineering efforts, *A. tumefaciens* will constitute a

powerful tool as an alternative therapeutic DNA delivery vector into human cells.

Material and methods

Chemicals are purchased from Sigma, unless otherwise stated.

Cloning

In silico cloning was performed using Benchling software and the cloning strategy of individual plasmids is described in Supplementary table 2.

Phusion polymerase (Thermo) was used for PCR using primers from Microsynth (Switzerland) and restriction enzymes (NEB) for digestion. Plasmid, genomic and gel-purified DNA was extracted using commercially available kits. We performed Gibson assembly using NEB HiFi DNA assembly kit, or classical ligation using the T4 Ligase (Thermo). We noted an improvement of the Gibson assembly efficiency when gel-purifying DNA using the Monarch kit (NEB) compared to other kits. Constructs were transformed by heat shock in XL10Gold (Agilent) prepared in 100 mM calcium chloride and 15% glycerol. Bacteria were plated on Luria broth (LB) plates containing the respective antibiotics and plasmids were screened by Sanger sequencing (Microsynth).

Composition of home-made media and agar plates:

- 20x AB salts (per 200mL): 4 g NH4Cl, 1.2 g $MgSO_4 \cdot 7H2O$, 0.6 g KCl, 0.04 g $CaCl_2$, 10 mg $FeSO_4 \cdot 7H_2O$. Sterile-filtered.

- 20x AB buffer (per 200mL): 12 g K_2HPO_4, 4 g NaH_2PO_4, pH to 7.0 using either KOH or H_3PO_4, as required, before autoclaving.

- Induction medium (IM): 1x AB salts, 0.5% glucose, 2 mM phosphate buffer pH 5.6, 50 mM 2-(4-morpholino)-ethane sulfonic acid (MES)

- Agrobacterium minimal medium: 1x AB salts, 1x AB buffer, 0.5% sucrose, antibiotics.

- ATGN plates: 1x AB salts, 1x AB buffer, 1% glucose, 1.5% noble agar, antibiotics

- ATSN plates: 1x AB salts, 1x AB buffer, 5% sucrose, 1.5% noble agar.

- SDCAA: 18.2% Sorbitol, 2% Glucose, 0.67% Yeast Nitrogen Base, 0.5% Casamino Acids, 0.54% Disodium Phosphate, 0.86% Monosodium Phosphate (Add 1.5% Agar for plates.

- SGCAA: 18.2% Sorbitol, 0.8% Glucose, 8% galactose, 0.67% Yeast Nitrogen Base, 0.5% Casamino Acids, 0.54% Disodium Phosphate, 0.86% Monosodium Phosphate

Bacterial culture and induction

E. coli were cultured at 37°C in LB containing either 100 µg/mL ampicillin (Huberlab), 50

µg/mL kanamycin, 25 µg/mL chloramphenicol, 50 µg/mL spectinomycin (Chemie

Brunschwig).

In this study, we used the disarmed strain *A. tumefaciens* C58C1 pMP90 (GV3101, see

Supplementary table 3, genome accession number: GCA_000092025.1[65]). *A. tumefaciens*

were cultured at 28-30°C in LB containing 60 µg/mL gentamycin (Biochemica) or, when

required, either 50 µg/mL kanamycin, 50 µg/mL spectinomycin or 100 µg/mL carbenicillin.

Unless otherwise stated, *A. tumefaciens* was inoculated at an optical density of 0.1 for 8h in

LB containing antibiotics and early stationary cells were induced overnight by addition of one

volume of induction medium IM and 100 µM acetosyringone (AS) for virulence induction

using the VirE promoter. For cumic acid-inducible constructs, early stationary cells were

induced overnight in LB by addition of cumic acid at 10 µM.

Bacterial strain engineering

For replicative plasmids, 1 mL of exponential culture of bacteria was washed 3 times in bi-

distilled water, concentrated 20 times and electroporated with 100 ng of plasmid in 1 mm

cuvette, rescued for 60 min in Super Optimal broth with Catabolite repression (SOC) medium

and plated on the corresponding antibiotics plates. Electro-competent bacteria were snap-

frozen in 15% glycerol solution.

For markerless genetic engineering of *A. tumefaciens*, we followed Morton and Fuqua,

2012[66], using *E. coli* S17-1 for conjugation and pNPTS138 suicide vector (see supplementary

Tables 2 and 3), with the following modifications: we added rifampicin (Axon Lab) at 25

µg/mL during selection and counterselection steps (ATGN and ATSN plates) to better kill

donor *E. coli*. As kanamycin is inhibited by phosphate-buffered media, we increased the

concentration to 1200 µg/mL during selection. LB plates containing rifampicin at 25 µg/mL

and kanamycin at 300 µg/mL were sometimes more efficient than the aforementioned ATGN

plates. Mutants were screened by colony-PCR using primers flanking the knockin or

knockout sites and validated by Sanger sequencing.

Bacterial staining, titration and quantification

Bacteria displaying VHH were washed with PBS and stained with recombinant GFP at 100

µg/mL for 10 minutes prior to two PBS washes. Bacteria harboring an HA tag were washed

with PBS and stained with anti-HA antibody conjugated with FITC (Abcam ab1208) at 10

µg/mL for 75 minutes in the dark on ice, washed once with PBS. For mScarlet-negative cells,

viability was checked by concomitant addition of 10 µg/mL propidium iodide (PI) during

staining. Wide field fluorescent pictures of bacteria on a coverslip under 1% agarose PBS

pads were taken at 100x and 1.5x lens magnification.

Bioinformatics and modeling

Protein sequences were submitted to the deep learning structure prediction online server

RoseTTaFold, provided by the Baker lab: robetta.bakerlab.org [28].

Protein sequences were submitted to the online deep neural network software SignalP-5.0

for signal peptide prediction (services.healthtech.dtu.dk/service.php?SignalP-5.0)[29].

Average amino acid usage was extracted from kazusa.or.jp/codon/cgi-

bin/showcodon.cgi?species=260551.

Engineering of S. cerevisiae

For GFP display, we fused eGFP to Aga2p under the control of a galactose-inducible

promoter pGal1. Aga2p form disulfide bonds with Aga1p anchored in the cell wall, resulting in

eGFP being anchored in the cell wall.

S. cerevisiae eby100 was retransformed using 1 µg of pGal1 – Aga2p_eGFP following the

protocol of the EZ yeast transformation kit II (Zymo). 100 µL of cells were selected on

SDCAA plates, which do not contain tryptophan. Colonies were directly selected by induction

and visualization with fluorescent microscopy.

A. tumefaciens binding to S. cerevisiae

Yeast GFP display induced overnight in SGCAA were washed twice in PBS to remove GFP

in suspension and concentrated 10 times. Induced *A. tumefaciens* were washed once in PBS

and added to concentrated yeast at a 1 to 1 volume ratio for 60-90 min. Five µL of the cell

mixture were transferred to and sandwiched between two coverslips, the yeast cells were left

to settle down and imaging was performed at 100x with a confocal microscope and 0.3 µm

step. We used NIS Elements (Nikon) for three-dimensional rendering of z-stack pictures and

manual cell counting. Per biological replicate, a total of 20 to 50 yeast cells were considered

on four different fields of view. For each field of view, the number of bound A. tumefaciens

was divided by the number of yeast cell considered. The mean of these values was

accounted for one biological replicate.

Mammalian cell culture and engineering

HEK293T and HeLa cells were cultured in high-glucose Dulbecco's Modified Eagle Medium

(DMEM, Thermofisher) supplemented with 10% FBS (Life Technologies) at 37°C and 5%

CO_2. A GFP-displaying monoclonal HEK cell line was generated by transfecting PvuI-

linearized pXP145 and FACS sorting after 7 days of culture.

HeLa cells stably expressing the doxycycline-inducible GFP display were further engineered

to stably express LgBit by transduction (pXP499), using lentivector as described previously[47].

After removal of the lentiviruses, a polyclonal cell line was obtained by selection with

puromycin (Labforce) at 2 µg/mL for one week.

Mammalian cell transient transfection

Mammalian cells were transfected with Lipofectamine 3000 (Life Technologies) overnight

with 100 ng of purified plasmid per well of 96-well plates, following the manufacturer's

instructions.

A. tumefaciens binding to mammalian cells

Mammalian cells were washed with medium twice prior to the addition of 10 µL of bacteria

induced with 100 µM AS per well of 96-well plates. Bacteria were homogenized by pipetting

and left to adhere for 5 hours at 30°C and 5% CO2. For the prevention of binding using

soluble GFP, bacteria were incubated with 100 µg/mL recombinant eGFP for 5 minutes prior

to the addition to mammalian cells. Consequently, recombinant eGFP was also present

during coculture at a concentration of 10 µg/mL. After coculture, wells were washed 5 times

with mammalian culture medium and imaging was performed in the region of the well the

closest to the dispensing of the medium. Biological replicates were performed on different

days and one biological replicate is as follows: confocal microscope Z-stacks were acquired

over 12 µm and 2 µm steps in three to five representative fields of view. On each field of

view, we estimated the number of HeLa cells and counted bacteria on maximum intensity

projection using ImageJ and Trackmate[67].

Split luciferase assays

A polyclonal cell line of HeLa cells stably expressing LgBit and doxycycline-inducible GFP

display was used. Cells were seeded in 96-well plates (Costar 3603). 6 h later, they were

induced when required with doxycycline at 1 µg/mL for overnight expression of GFP-display.

Cells were washed twice with medium to remove shed GFP from the supernatant and 22 µL

of induced *A. tumefaciens* (LB-IM AS 100 µM) were added to 100 µL of culture medium for 2

h. To reduce unspecific binding and VirE2 transfer, wells were washed 3 times with DMEM

with 20% LB-IM and 100 µM AS, and infection was prolonged for 3 more hours before

readout (Fig. 5D and S8A-B), else infection was performed for 5 hours.

Wells were washed with OptiMEM (Thermo) twice and 20 µL of a 1:19 mixture of
substrate:buffer from the Nano-Glo Live Cell Assay (Promega) were added to 80 µL of
OptiMEM per well. After 3 min incubation with the reagent, luminescence activity was
acquired using a multiwell plate reader (Tecan Spark) with 5 s integration time per well.
Background from wells with only reagent and OptiMEM was subtracted from the values.

Biological replicates were performed on different days and one biological replicate consists in
three technical replicates (separate wells).

Microscopy

For widefield visualizations of bacteria, we used a Nikon TiE epifluorescence microscope
equipped with a Hamamatsu ORCA Flash 4 camera and an oil immersion 100x Plan APO
N.A. 1.45 objective.

For bacterial adhesion to yeast and mammalian cells, we used a Nikon Eclipse Ti2-E
inverted microscope coupled with a Yokogawa CSU W2 confocal spinning disk unit and
equipped with a Prime 95B sCMOS camera (Photometrics). We used either a 40x air
objective with N.A. of 1.15 to acquire z-stacks for mammalian cells or a 100x oil immersion
objective with N.A. of 1.45 for yeast cells.

Production of recombinant proteins

6x-His tagged eGFP on a pET28a vector (see Supplementary table 2) was retransformed
into BL21 strain. We induced production with 1 mM IPTG (Fisher bioreagents) at room
temperature overnight. We pelleted and lysed bacteria by sonication in lysis buffer (Tris
100mM, NaCl 0.5M, glycerol 5%) and eGFP was purified using fast flow His-affinity columns
(GE Healthcare) and eluted with 0.5 M imidazole. We exchanged buffer to PBS using 30kDa
ultracentrigation spin columns (Merck) and adjusted the concentration to 1 mg/mL. Aliquots
were snap frozen for further use.

Supporting information

Supplementary figure 6: Analysis of the signal peptides of the two putative autotransporters
of *A. tumefaciens*

Supplementary figure 7: Removal of the disulfide bonds of the native passenger domain in
Atu5364 promotes even surface distribution of the passenger domain.

Supplementary figure 8: VHH display strategy using Atu5364 as scaffold.

Acknowledgments

The authors are grateful to Dr. Ingmar Riedel-Kruse (The University of Arizona) and Dr. Luis

Angel Fernández (Centro Nacional de Biotecnologia) for the nanobody anti-GFP construct, to

Dr. Bruno Correia, Stéphane Rosset and Dr. Leo Scheller (Ecole Polytechnique Fédérale de

Lausanne) for the production and purification of recombinant proteins, *S. cerevisiae* eby100

strain, NanoLuc and split NanoLuc constructs, to Dr. Csaba Koncz (Max Planck Institute for

plant breeding research) for *A. tumefaciens* GV3101.

Funding

521   We are grateful for the funding provided by the Gebert Rüf Foundation, project number GRS-

522   057/16 and the Novartis FreeNovation 2020 program.

Author contributions

525   X.P. and A. Persat conceived the work. X.P. designed and performed the experiments. A.

526   Pham and J.W. performed experiments for the revision of the manuscript. Z.A-M. provided

527   support for cell culture, DNA purification and colony PCR. The EPFL Flow Cytometry Core

528   Facility performed single cell sorting. X.P. and A. Pham analyzed the data. X.P. and A.

529   Persat wrote the manuscript. All authors gave input and approved the manuscript.

Conflicts of interest

532   None.

533   References

1.   Hooykaas PJJ, Heusden GPH Van, Niu X, et al. Agrobacterium -Mediated
Transformation of Yeast and Fungi. doi:10.1007/82

2.   Gelvin SB. Traversing the Cell: Agrobacterium T-DNA's Journey to the Host Genome.
Front Plant Sci. 2012;3(March):1-11. doi:10.3389/fpls.2012.00052

3.   Lacroix B, Citovsky V. Pathways of DNA transfer to plants from Agrobacterium
tumefaciens and related bacterial species. *Annu Rev Phytopathol*. 2019;57:231-251.
doi:10.1146/annurev-phyto-082718-100101

4.   Sardesai N, Subramanyam S. Agrobacterium: A Genome-Editing Tool-Delivery
System. *Curr Top Microbiol Immunol*. 2018;418:463-488. doi:10.1007/82_2018_101

5.   Hwang H-H, Gelvin SB, Lai E-M. *Editorial: "Agrobacterium Biology and Its Application*
to Transgenic Plant Production." Vol 6.; 2015. doi:10.3389/fpls.2015.00265

6.   Hwang H-H, Yu M, Lai E-M. Agrobacterium -Mediated Plant Transformation: Biology
and Applications . *Arab B*. 2017;15:e0186. doi:10.1199/tab.0186

7.   Krenek P, Samajova O, Luptovciak I, Doskocilova A, Komis G, Samaj J. Transient
plant transformation mediated by Agrobacterium tumefaciens: Principles, methods and
applications. *Biotechnol Adv*. 2015;33(6):1024-1042.
doi:10.1016/j.biotechadv.2015.03.012

8.   Thompson MA, Onyeziri MC, Fuqua C. Function and Regulation of Agrobacterium
tumefaciens Cell Surface Structures that Promote Attachment. *Curr Top Microbiol*
Immunol. 2018. doi:10.1007/82

9.   Matthysse AG. Attachment of Agrobacterium to plant surfaces. *Front Plant Sci*.
2014;5(June):1-8. doi:10.3389/fpls.2014.00252

10.   Leuzinger K, Dent M, Hurtado J, et al. Efficient agroinfiltration of plants for high-level
transient expression of recombinant proteins. *J Vis Exp*. 2013;(77):1-9.
doi:10.3791/50521

11.   Gürel F, Uçarlı C, Tufan F, Kalaskar DM. Enhancing T-DNA Transfer Efficiency in
Barley (Hordeum vulgare L.) Cells Using Extracellular Cellulose and Lectin. *Appl*
Biochem Biotechnol. 2015;176(4):1203-1216. doi:10.1007/s12010-015-1640-0

12.   Van Ulsen P, Zinner KM, Jong WSP, Luirink J. On display: Autotransporter secretion
and application. *FEMS Microbiol Lett*. 2018;365(18):1-10. doi:10.1093/femsle/fny165

13.   Tozakidis IEP, Sichwart S, Jose J. Going beyond E. coli: autotransporter based
surface display on alternative host organisms. *N Biotechnol*. 2015;32(6):644-650.
doi:10.1016/j.nbt.2014.12.008

14.   Nicolay T, Vanderleyden J, Spaepen S. Autotransporter-based cell surface display in
Gram-negative bacteria. *Crit Rev Microbiol*. 2015;41(1):109-123.
doi:10.3109/1040841X.2013.804032

15.   Salema V, Fernández LÁ. Escherichia coli surface display for the selection of
nanobodies. *Microb Biotechnol*. 2017;10(6):1468-1484. doi:10.1111/1751-7915.12819

16.   González C, Chames P, Kerfelec B, Baty D, Robert P, Limozin L. Nanobody-CD16
Catch Bond Reveals NK Cell Mechanosensitivity. *Biophys J*. 2019;116(8):1516-1526.
doi:10.1016/j.bpj.2019.03.012

17.   Veiga E, De Lorenzo V, Fernández LA. Structural tolerance of bacterial
autotransporters for folded passenger protein domains. *Mol Microbiol*.
2004;52(4):1069-1080. doi:10.1111/j.1365-2958.2004.04014.x

18.   Olichon A, De Marco A. Preparation of a naïve library of camelid single domain
antibodies. *Methods Mol Biol*. 2012;911:65-78. doi:10.1007/978-1-61779-968-6_5

19.   Fraile S, Briones M, Revenga-Parra M, De Lorenzo V, Lorenzo E, Martínez-García E.

Engineering Tropism of Pseudomonas putida toward Target Surfaces through Ectopic
Display of Recombinant Nanobodies. *ACS Synth Biol.* 2021;10(8):2049-2059.
doi:10.1021/acssynbio.1c00227

20.   Salema V, Mañas C, Cerdán L, et al. High affinity nanobodies against human
epidermal growth factor receptor selected on cells by E. coli display. *MAbs.*
2016;8(7):1286-1301. doi:10.1080/19420862.2016.1216742

21.   Glass DS, Riedel-Kruse IH. A Synthetic Bacterial Cell-Cell Adhesion Toolbox for
Programming Multicellular Morphologies and Patterns. *Cell.* 2018;174(3):649-658.e16.
doi:10.1016/j.cell.2018.06.041

22.   Ting S-Y, Martınez-Garcia E, Huang S, Peterson SB, De Lorenzo V, Mougous JD.
Short Article Targeted Depletion of Bacteria from Mixed Populations by Programmable
Adhesion with Antagonistic Competitor Cells Short Article Targeted Depletion of
Bacteria from Mixed Populations by Programmable Adhesion with Antagonistic
Competitor Cells. *Cell Host Microbe.* 2020:1-9. doi:10.1016/j.chom.2020.05.006

23.   Chien T, Doshi A, Danino T. Advances in bacterial cancer therapies using synthetic
biology. *Curr Opin Syst Biol.* 2017;5:1-8. doi:10.1016/j.coisb.2017.05.009

24.   Piñero-Lambea C, Bodelón G, Fernández-Periáñez R, Cuesta AM, Álvarez-Vallina L,
Fernández LÁ. Programming controlled adhesion of E. coli to target surfaces, cells,
and tumors with synthetic adhesins. *ACS Synth Biol.* 2015;4(4):463-473.
doi:10.1021/sb500252a

25.   Grijpstra J, Arenas J, Rutten L, Tommassen J. Autotransporter secretion: Varying on a
theme. *Res Microbiol.* 2013;164(6):562-582. doi:10.1016/j.resmic.2013.03.010

26.   Meuskens I, Saragliadis A, Leo JC, Linke D. Type V secretion systems: An overview
of passenger domain functions. *Front Microbiol.* 2019;10(MAY):1-19.
doi:10.3389/fmicb.2019.01163

27.   Fairman JW, Dautin N, Wojtowicz D, et al. Crystal structures of the outer membrane
domain of intimin and invasin from enterohemorrhagic E. coli and enteropathogenic Y.
pseudotuberculosis. *Structure.* 2012;20(7):1233-1243. doi:10.1016/j.str.2012.04.011

28.   Baek M, DiMaio F, Anishchenko I, et al. Accurate prediction of protein structures and
interactions using a three-track neural network. *Science.* 2021;373(6557):871-876.
doi:10.1126/SCIENCE.ABJ8754

29.   Almagro Armenteros JJ, Tsirigos KD, Sønderby CK, et al. SignalP 5.0 improves signal
peptide predictions using deep neural networks. *Nat Biotechnol 2019 374.*
2019;37(4):420-423. doi:10.1038/s41587-019-0036-z

30.   Rojas-lopez M, Zorgani MA, Kelley LA, Bailly X, Rosini R, Desvaux M. Identification of
the Autochaperone Domain in the Type Va Secretion System (T5aSS): Prevalent
Feature of Autotransporters with a β -Helical Passenger. 2018;8(January):1-10.
doi:10.3389/fmicb.2017.02607

31.   Imai K, Mitaku S. Mechanisms of secondary structure breakers in soluble proteins.
Biophysics (Oxf). 2005;1:55-65. doi:10.2142/BIOPHYSICS.1.55

32.   Brotcke Zumsteg A, Goosmann C, Brinkmann V, Morona R, Zychlinsky A. IcsA is a
Shigella flexneri adhesin regulated by the type III secretion system and required for
pathogenesis. *Cell Host Microbe.* 2014;15(4):435-445.
doi:10.1016/J.CHOM.2014.03.001

33.   Leininger E, Roberts M, Kenimer JG, et al. Pertactin , an Arg-Gly-Asp-containing
Bordetella pertussis surface protein that promotes adherence of mammalian cells.
Proc Natl Acad Sci. 1991;88(2):345-349. doi:10.1073/PNAS.88.2.345

34.   Garnham CP, Campbell RL, Davies PL. Anchored clathrate waters bind antifreeze
proteins to ice. *Proc Natl Acad Sci U S A.* 2011;108(18):7363-7367.
doi:10.1073/PNAS.1100429108/

35. Herrmann J, Jabbarpour F, Bargar PG, et al. Environmental Calcium Controls
Alternate Physical States of the Caulobacter Surface Layer. *Biophys J*.
2017;112(9):1841-1851. doi:10.1016/J.BPJ.2017.04.003

36. Lau JHY, Nomellini JF, Smit J. Analysis of high-level S-layer protein secretion in
Caulobacter crescentus. *Can J Microbiol*. 2010;56(6):501-514. doi:10.1139/W10-036

37. Heras B, Totsika M, Peters KM, et al. The antigen 43 structure reveals a molecular
Velcrolike mechanism of autotransporter-mediated bacterial clumping. *Proc Natl Acad*
Sci U S A. 2014;111(1):457-462. doi:10.1073/PNAS.1311592111

38. Denkovskiene E, Paskevicius S, Werner S, Gleba Y, Razanskiene A. Inducible
Expression of Agrobacterium Virulence Gene VirE2 for Stringent Regulation of T-DNA
Transfer in Plant Transient Expression Systems. 2015;28(11):1247-1255.

39. Leyton DL, Sevastsyanovich YR, Browning DF, et al. Size and conformation limits to
secretion of disulfide-bonded loops in autotransporter proteins. *J Biol Chem*.
2011;286(49):42283-42291. doi:10.1074/jbc.M111.306118

40. Hirooka T, Rogowsky PM, Kado CI. Characterization of the virE locus of
Agrobacterium tumefaciens plasmid pTiC58. *J Bacteriol*. 1987;169(4):1529-1536.
doi:10.1128/jb.169.4.1529-1536.1987

41. Saerens D, Conrath K, Govaert J, Muyldermans S. Disulfide Bond Introduction for
General Stabilization of Immunoglobulin Heavy-Chain Variable Domains. *J Mol Biol*.
2008;377(2):478-488. doi:10.1016/j.jmb.2008.01.022

42. Park S, Zheng JH, Nguyen VH, et al. RGD Peptide Cell-Surface Display Enhances the
Targeting and Therapeutic Efficacy of Attenuated Salmonella -mediated Cancer
Therapy. *Theranostics*. 2016;6(10). doi:10.7150/thno.16135

43. Diggle SP, Stacey RE, Dodd C, Cámara M, Williams P, Winzer K. The galactophilic
lectin, LecA, contributes to biofilm development in Pseudomonas aeruginosa. *Environ*
Microbiol. 2006;8(6):1095-1104. doi:10.1111/j.1462-2920.2006.001001.x

44. Tielker D, Hacker S, Loris R, et al. Pseudomonas aeruginosa lectin LecB is located in
the outer membrane and is involved in biofilm formation. *Microbiology*.
2005;151(5):1313-1323. doi:10.1099/mic.0.27701-0

45. Chemani C, Imberty A, De Bentzmann S, et al. Role of LecA and LecB lectins in
Pseudomonas aeruginosa-induced lung injury and effect of carbohydrate ligands.
Infect Immun. 2009;77(5):2065-2075. doi:10.1128/IAI.01204-08

46. Nilsson LM, Yakovenko O, Tchesnokova V, et al. The cysteine bond in the Escherichia
coli FimH adhesin is critical for adhesion under flow conditions. *Mol Microbiol*.
2007;65(5):1158-1169. doi:10.1111/j.1365-2958.2007.05858.x

47. Pierrat X, Wong J, Al-Mayyah Z, Persat A. The Mammalian Membrane
Microenvironment Regulates the Sequential Attachment of Bacteria to Host Cells.
MBio. 2021;12(4).

48. Piers KL, Heath JD, Liang X, Stephens KM, Nester EW. Agrobacterium tumefaciens-
mediated transformation of yeast. *Proc Natl Acad Sci U S A*. 1996;93(4):1613-1618.
doi:10.1073/pnas.93.4.1613

49. Gera N, Hussain M, Rao BM. Protein selection using yeast surface display. *Methods*.
2013;60(1):15-26. doi:10.1016/j.ymeth.2012.03.014

50. Pelczar P, Kalck V, Gomez D, Hohn B. Agrobacterium proteins VirD2 and VirE2
mediate precise integration of synthetic T-DNA complexes in mammalian cells. *EMBO*
Rep. 2004;5(6):632-637. doi:10.1038/sj.embor.7400165

51. Dixon AS, Schwinn MK, Hall MP, et al. NanoLuc Complementation Reporter
Optimized for Accurate Measurement of Protein Interactions in Cells. *ACS Chem Biol*.
2016;11(2):400-408. doi:10.1021/acschembio.5b00753

52.   Westerhausen S, Nowak M, Torres-Vargas CE, et al. A NanoLuc luciferase-based
assay enabling the real-time analysis of protein secretion and injection by bacterial
type III secretion systems. *Mol Microbiol*. 2020;113(6):1240-1254.
doi:10.1111/mmi.14490

53.   Lettl C, Haas R, Fischer W. Kinetics of CagA type IV secretion by Helicobacter pylori
and the requirement for substrate unfolding. *Mol Microbiol*. 2021;116(3):794-807.
doi:10.1111/mmi.14772

54.   Zhou X, Christie PJ. Mutagenesis of the Agrobacterium VirE2 Single-Stranded DNA-
Binding Protein Identifies Regions Required for Self-Association and Interaction with
VirE1 and a Permissive Site for Hybrid Protein Construction. 1999;181(14):4342-4352.

55.   Li X, Yang Q, Peng L, et al. Agrobacterium-delivered VirE2 interacts with host
nucleoporin CG1 to facilitate the nuclear import of VirE2-coated T complex. *Proc Natl*
Acad Sci U S A. 2020;117(42):26389-26397. doi:10.1073/pnas.2009645117

56.   Muyldermans S. Applications of Nanobodies. *Annu Rev Anim Biosci*. 2021;9:401-421.
doi:10.1146/annurev-animal-021419-083831

57.   Kunik T, Tzfira T, Kapulnik Y, Gafni Y, Dingwall C, Citovsky V. Genetic transformation
of HeLa cells by Agrobacterium. *Proc Natl Acad Sci*. 2001;98(4):1871-1876.
doi:10.1073/pnas.98.4.1871

58.   Petrunia I V., Frolova OY, Komarova T V., Kiselev SL, Citovsky V, Dorokhov YL.
Agrobacterium tumefaciens-induced bacteraemia does not lead to reporter gene
expression in mouse organs. *PLoS One*. 2008;3(6):2-6.
doi:10.1371/journal.pone.0002352

59.   Gaspar YM, Nam J, Schultz CJ, et al. Characterization of the Arabidopsis Lysine-Rich
That Results in a Decreased Efficiency of Agrobacterium Transformation 1.
2004;135(August):2162-2171. doi:10.1104/pp.104.045542.2162

60.   Li X, Pan SQ. Agrobacterium delivers VirE2 protein into host cells via clathrin-
mediated endocytosis. *Sci Adv*. 2017;3(3):1-12. doi:10.1126/sciadv.1601528

61.   Vergunst AC, Schrammeijer B, Den Dulk-Ras A, De Vlaam CMT, Regensburg-Tuink
TJG, Hooykaas PJJ. VirB/D4-dependent protein translocation from Agrobacterium into
plant cells. *Science*. 2000;290(5493):979-982. doi:10.1126/science.290.5493.979

62.   Schmitz DJ, Ali Z, Wang C, et al. CRISPR/Cas9 Mutagenesis by Translocation of
Cas9 Protein Into Plant Cells via the Agrobacterium Type IV Secretion System. *Front*
Genome Ed. 2020;2(July):1-11. doi:10.3389/fgeed.2020.00006

63.   Davidsson M, Wang G, Aldrin-Kirk P, et al. A systematic capsid evolution approach
performed in vivo for the design of AAV vectors with tailored properties and tropism.
Proc Natl Acad Sci U S A. 2019;116(52):27053-27062. doi:10.1073/pnas.1910061116

64.   Schmit PF, Pacouret S, Zinn E, et al. Cross-Packaging and Capsid Mosaic Formation
in Multiplexed AAV Libraries. *Mol Ther - Methods Clin Dev*. 2020;17(June):107-121.
doi:10.1016/j.omtm.2019.11.014

65.   Wood DW, Setubal JC, Kaul R, et al. The genome of the natural genetic engineer
Agrobacterium tumefaciens C58. *Science*. 2001;294(5550):2317-2323.
doi:10.1126/science.1066804

66.   Morton ER, Fuqua C. Unit 3D.2 Genetic manipulation of Agrobacterium. *Curr Protoc*
Microbiol. 2012:1-17. doi:10.1002/9780471729259.mc03d02s25

67.   Tinevez JY, Perry N, Schindelin J, et al. TrackMate: An open and extensible platform
for single-particle tracking. *Methods*. 2017;115:80-90.
doi:10.1016/j.ymeth.2016.09.016

727