1 **Title**
2 **Substrate topology and cognate immunity modulate antibacterial toxicity of Tae cell wall**
3 **hydrolases**
4

5 **Authors**
6 Atanas Radkov[1], Anne Sapiro[1], Sebastian Flores[2], Corey Henderson[3], Hayden Saunders[1], Rachel
7 Kim[4], Steven Massa[5], Samuel Thompson[6], Chase Mateusiak[7], Jacob Biboy[8], Ziyi Zhao[1], Lea M
8 Starita[9,10], William Hatleberg, Waldemar Vollmer[8], Alistair B Russell[11], Jean-Pierre Simorre[12], Spencer
9 Anthony-Cahill[13], Peter Brzovic[14], Beth Hayes[1], Seemay Chou[1]*
10
1 Department of Biochemistry & Biophysics, University of California – San Francisco, San Francisco,
CA
2 University of Miami, Miami, FL
3 InBios International, Seattle, WA
4 Pacific Northwest University of Health Sciences, Yakima, WA
5 Department of Biology, Stanford University, Stanford, CA
6 Department of Bioengineering, Stanford University, Stanford, CA
7 Computer Science Department, University of Washington in St Louis, St. Louis, MO
8 Center for Bacterial Cell Biology, Newcastle University, Newcastle upon Tyne, United Kingdom
9 Department of Genome Sciences, University of Washington, Seattle, WA
10 Brotman Baty Institute for Precision Medicine, Seattle, WA
11 Division of Biological Sciences, University of California, San Diego, La Jolla, CA
12 Université Grenoble Alpes, CNRS, CEA, IBS, Grenoble, France
13 Chemistry Department, Western Washington University, Bellingham, WA
14 Department of Biochemistry, University of Washington, Seattle, WA
26
*Corresponding author: seemay.chou@ucsf.edu

Abstract
Members of the bacterial T6SS amidase effector (Tae) superfamily of toxins are delivered between
competing bacteria to degrade cell wall peptidoglycan. Although Taes share a common substrate, they
exhibit distinct antimicrobial potency across different competitor species. To investigate the molecular
basis governing these differences, we quantitatively defined the functional determinants of Tae1 from
Pseudomonas aeruginosa PAO1 using a combination of nuclear magnetic resonance (NMR) and a
high-throughput *in vivo* genetic approach called deep mutational scanning (DMS). As expected,
combined analyses confirmed the role of critical residues near the Tae1 catalytic center. Unexpectedly,
DMS revealed substantial contributions to enzymatic activity from a much larger, ring-like functional hot
spot extending around the entire circumference of the enzyme. Comparative DMS across distinct
growth conditions highlighted how functional contribution of different surfaces is highly context-
dependent, varying alongside composition of targeted cell walls. These observations suggest that Tae1
engages with the intact cell wall network through a more distributed three-dimensional interaction
interface than previously appreciated, providing an explanation for observed differences in antimicrobial
potency across divergent Gram-negative competitors. Further binding studies of several Tae1 variants
with their cognate immunity protein demonstrate that requirements to maintain protection from Tae
activity may be a significant constraint on the mutational landscape of *tae1* toxicity in the wild. In total,
our work reveals that Tae diversification has likely been shaped by multiple independent pressures to
maintain interactions with binding partners that vary across bacterial species and conditions.

48
49
50
51

Introduction

Bacteria live in dense polymicrobial communities where competition for nutrients and space impacts survival[1]. To combat neighboring rivals, some Gram-negative bacteria use a contact-dependent toxin delivery system called the Type 6 secretion system (T6SS)[2,3] that injects a cocktail of effectors from the donor cell directly into nearby recipient cells[4–6]. A bacterium's T6S arsenals include several types of protein effectors with distinct activities that each compromise essential cellular pathways or components, such as recipient peptidoglycan (PG) or membranes. Although many T6S bacteria deploy similar toxins, interbacterial outcomes can vary considerably depending on the bacterial species engaged in T6S-mediated competition[7,8]. The molecular basis of this organism-level specificity of interbacterial T6SSs remains poorly understood.

A potential source of specificity could be the T6S-delivered toxins themselves[9]. T6S toxins can belong to large superfamilies comprised of diverse homologs. One such example is the T6S amidase effector (*tae*) superfamily[7]. The *tae*s encode hydrolase enzymes that induce recipient cell lysis by breaking down the major structural component of bacterial cell envelopes, PG[5]. Tae toxins are known to be structurally and biochemically diverse[10]. Given that targeted PG substrates can also vary in a species-dependent manner[11], such diversity could potentially impact toxin antibacterial specificity. Tae toxins divide into four phylogenetically distinct subfamilies (Tae1-4)[7] that hydrolyze different amide bonds within PG, and toxins within the same subfamily can have unique PG compositional preferences.[10]

These observations raise the possibility that T6SS organismal specificity may be linked, in part, to the biochemical diversity of delivered toxins. Such a model would predict that physiological features of recipient microbes may have driven divergent evolution of T6S effector diversity. To address this hypothesis, a quantitative, molecular understanding of T6S effector specificity is needed. However, comprehensive functional analyses are challenging for toxins such as the Taes, given that they act on cellular substrates with complex, three-dimensional structures *in vivo*. In cells, PG is a cross-linked macromolecule that forms a contiguous, mesh-like network that is structurally heterogeneous, insoluble, and dynamically regulated[12,13]. These features are not well captured through *in vitro* functional studies that necessarily focus on soluble PG fragments.

To probe Tae–cell wall interactions in a physiologically relevant context, we developed a high throughput genetic approach known as deep mutational scanning (DMS)[16] that surveys the entire Tae enzyme for functional determinants of toxicity in live *Escherichia coli* cells. We focused on Tae1 from the common soil bacterium *Pseudomonas aeruginosa*, which is known to interact with a wide range of microbes, including *E. coli*. We found that an extended surface across the entire Tae1 enzyme is important for mediating *in vivo* degradation of cell walls. Furthermore, we present evidence that Tae1 functional determinants can be quite distinct depending on the compositional context of PG. Finally, we discovered that the evolutionary potential for Tae1 potency against a broad range of cell wall types may be constrained by orthogonal functional requirements, such as Tae1 interactions with its cognate immunity protein during T6S-mediated competition.

RESULTS

Deep mutational scanning identifies determinants of Tae1 function *in vivo*

To understand how sequence diversity within the Tae protein superfamily impacts interbacterial toxicity, we must first probe how each amino acid residue contributes to Tae function. Biochemical assays of Tae protein binding to and degradation of peptidoglycan (PG) use isolated, purified sacculi[10], which differ from the turgor-stretched PG layer in the periplasm of living cells. Thus, these assays likely identify only a subset of determinants that govern how Tae engages with PG *in vivo*. To define comprehensively and quantitatively all the amino acid residues in Tae that are important for its function *in vivo*, we developed a DMS experiment in live bacterial cells that allows us to examine the contributions of amino acid substitutions at every Tae residue in a physiologically relevant context.[14,15]

We adapted a DMS approach[16,17] to study the Tae1 protein from *P. aeruginosa*, which is injected into *E. coli* and leads to cell lysis (**Figure 1A**). We based our screen on a lysis assay that measures the toxic effects of Tae1 on *E. coli* when it is ectopically expressed in the periplasm, which contains the PG[10]. Periplasmic expression of wild-type Tae1 (Tae1[WT]), but not catalytically inactive Cys30Ala Tae (Tae1[C30A]), results in rapid cell lysis upon induction due to osmotic stress resulting cell wall degradation by the toxin. Cells expressing *tae1[WT]* are depleted from the population faster than cells expressing inactive Tae1 (**Figure 1B**). Mutations that increase or decrease the rate of Tae1-induced lysis can be identified by sequencing and analyzing plasmid copy numbers in the population over time.

We generated a barcoded *tae1* plasmid library for massively parallel screening in *E. coli* (**Figure 1C**). Mutations were introduced by error-prone PCR, and library diversity was characterized by sequencing. We focused our analysis on copies of *tae1* that had a single mutation. In total, there were 902 different Tae1 variants, which encompassed at least one amino acid substitution at every position in the 154-residue protein. To ensure Tae1 delivery to the bacterial cell wall, we

Figure 1. Deep mutational scanning identifies amino acid residues important for Tae1 function *in vivo*. (A) Illustration of the function of the *P. aeruginosa* Tae1 toxin in lysis of *E. coli*. The donor *P. aeruginosa* cell injects Tae1 into the periplasm of the recipient *E. coli* cell. Tae1 cleaves cell wall PG at the cleavage site, resulting in cell lysis. (B) Illustration of lysis assays expressing Tae1 in the periplasm of *E. coli*. When wild-type Tae1 is expressed, cells are depleted over time (gray). When a loss-of-function mutation such as Tae1[C30A] is expressed, cells grow normally (red). When a gain-of-function mutation is expressed, cells are depleted more quickly over time (blue). T1 = time 1, T2 = time 2. (C) Schematic of deep mutational scanning screen for functional determinants of Tae1. A library of Tae1 mutants was transformed into *E. coli* and periplasmic expression of the mutant toxins was induced. The number of mutant cells remaining at T2 compared to the number at T1 yield a fitness score for each variant. Increased fitness indicates loss-of-function mutations in Tae1, which decrease cell lysis, whereas decreased fitness indicates gain-of-function mutations, which increase lysis. (D) DMS-derived relative fitness scores for all substitutions in the library at the Cys30 and His91 positions. Data shown are means +/- standard deviation, n = 3. DMS experiment in M9 media was repeated 3 independent times, each time with at least 3 technical replicates.

expressed the variants as fusion proteins with N-terminal periplasm-targeting *pelB* signal sequences.
We transformed *E. coli* (BW25113) grown in M9 minimal media with our *tae1* library and sequenced
plasmid barcodes, both before and two hours after library induction (**Figure 1C**). Tae1-dependent lytic
effects were quantified by using the DMS analysis pipeline Enrich2[17], which calculates output-to-input
ratios for individual variants, yielding fitness scores that are normalized to wild type (**Supplemental**
Table 1). In our assay, the fitness score reflects a difference in the antibacterial capacity of the Tae1
variant *in vivo* when compared to the wild type: relative enrichment of a variant indicates reduced Tae1-
dependent cell death or loss-of-function (LOF) and yields a positive fitness score, whereas relative
depletion of a variant indicates increased cell death or gain-of-function (GOF) and yields a negative
fitness score.
156
To validate our approach, we examined the canonical catalytic dyad residues of Tae1 to assess
whether our screen accurately identified known functional determinants. Any substitutions of these
catalytic residues, Cys30 and His91, should inhibit enzyme activity and lead to LOF. As anticipated, all
non-synonymous amino acid changes at these sites resulted in increased fitness scores, indicating
their crucial roles in Tae1 toxicity (**Figure 1C and D**) and confirming that the DMS screen was indeed
selecting *in vivo* functional determinants of Tae PG-degrading toxicity.
163
164 **DMS identifies functional hotspots extending around the surface of Tae1**
165
To obtain a global view of the amino acid residues in Tae1 that affect fitness in our screen, we
generated an averaged sequence-function map of variant fitness scores (**Figure 2A**). The majority of all
unique Tae1 protein variants (54%) were LOF variants with positive fitness scores, whereas only 6%
were GOF variants. To determine the functional role of each residue of Tae1, we collapsed and
averaged scores across all substitutions at each Tae1 site (**Figure 2A**) and superposed these onto the
three-dimensional structure of Tae1 (**Supplemental Figure 1**). For some Tae1 residues, position
averages obscured how the degree or direction of functional change depended greatly on the specific
amino acid change introduced. Therefore, we used *k*-means clustering, a statistical method for
analyzing DMS data[18], to classify Tae1 residues in an unbiased way into one of four categories:
residues where mutations generally decreased cell lysis (LOFclus), those where mutations generally
increased cell lysis (GOFclus), those where mutations had variable effects on lysis (mixed), and those
where mutations had little to no effect (tolerant). The vast majority of residues were categorized as

Figure 2. DMS identifies functional hotspots extending around the surface of Tae1. (A) Heatmap representation of fitness scores relative to wild-type Tae1 for all mutations in DMS. Each amino acid position is represented on the x-axis, and each possible amino acid substitution is represented on the y-axis, * = stop codon. Each box represents the average fitness score of three replicate experiments. The average fitness score of all substitutions at each residue is depicted across the bottom row. Catalytic residues Cys30 and His91 are indicated with asterisks. Variants in grey were not present in our library. (B) Results of *k*-means clustering of DMS data. Based on fitness effects, each residue of Tae1 was labelled loss-of-function (LOFclus), gain-of-function (GOFclus), mixed, or tolerant. (C) Ribbon diagram of Tae1 structure. LOFclus residues are shown in red, GOFclus residues are shown in blue, and all other residues are in gray. The catalytic residues of Tae1 (Cys30 and His91) are indicated. Mutations in the enzyme core lead to loss-of-function phenotypes. (D) Functional residues on the surface of Tae1. The catalytic surface of the enzyme is shown on the left and the opposite surface, after 180-degree rotation, is on the right. Color code as in part C. LOFclus mutations are located in large areas of the surface on both sides of the enzyme, and most GOFclus mutations are also found on the surface. See also supplemental figure 1.

either LOF[clus] (56%) or mixed (36%), and
fewer were categorized as GOF[clus] (5%) or
tolerant (3%) (**Figure 2B**). We mapped
these classifications onto the Tae1 structure
and found, consistent with other mutational
studies, a large majority of buried residues
(78%) including the catalytic residues, were
categorized as LOF[clus] (**Figure 2C**).
Residues on the enzyme surface, by
contrast, were a mix of categories including most of the GOF[clus] residues.
188
We anticipated that only surface residues within or near to the catalytic cleft would show loss-of-
function phenotypes. Instead, we found that many surface-exposed residues more distant from the
catalytic cleft were also categorized as LOF[clus] (**Figure 2D**). Indeed, a zone of LOF[clus] residues forms a
ring around the entire enzyme, suggesting a more distributed set of surfaces may play key roles in cell
lysis *in vivo* than was predicted by *in vitro* studies[10]. Many of the surface LOF[clus] residues that our
screen indicates are crucial for cell lysis *in vivo* are poorly conserved across the *tae1* superfamily[10]. We

Figure S1. Average fitness score across each residue. (A) Ribbon diagram and (B) surface diagrams of fitness scores when averaging scores for all substitutions at each residue. Red residues indicate loss-of-function on average and blue residues indicate gain-of-function on average. Cys30 and His91 of the active site are indicated.

195 posit that while these mutations are LOF
196 when applied to *E. coli* sacculi, divergent
197 bacterial physiologies *in situ* may instead
198 drive diversification of this extended
199 surface.
200
201 **Tae1 surfaces distal from catalytic**
202 **core mediate binding and hydrolysis**
203 **of cell wall network**
204
Our DMS results suggested surfaces
outside the Tae1 catalytic core are
functionally important. We observed that
several of these distal surface LOF[clus]
sites encoded serine and glycine
residues, which have been previously
implicated in sugar-binding[19,20]. Thus, it is
possible that the distal LOF[clus] surfaces
are important for interacting with the
glycan chains of PG. Alternatively, these
distal surfaces could mediate Tae1
interactions with surrounding, untargeted
peptide stems within the PG network. To
validate that these residues play a role in
Tae1 function, we chose nine LOF[clus]
variants distributed across the Tae1
surface (**Figure 3A**) and manually
assayed their individual effects through
E. coli lysis assays (**Figure 3B and**
Supplemental Figure 2). We focused
mainly on serine and glycine residues
and introduced the most conservative
substitutions represented in the screen
(Alanines or Serines, respectively).

Figure 3. Tae1 surfaces distal from catalytic core mediate binding and hydrolysis of cell wall network. (A) Nine distal residues on Tae1 that had loss of function phenotypes. The catalytic core is highlighted in yellow. (B) Lysis assays measuring relative lysis of *E. coli* expressing wild-type Tae1 (WT), negative controls (vector and C30A) and loss-of-function single amino acid substitutions (G42S, V34S, A52P, S144A, S149A, L21S, G48S, G56S). Data shown are means +/- standard deviation, n = 3, * p < 0.05. Lysis experiment was repeated 3 independent times, each time with at least 3 technical replicates. Western blots below show the beta subunit of RNA polymerase (RNAP β) and Tae1 proteins in each assay. All variants and controls showed reduced lysis when compared to wild-type Tae1. (C) Schematic of the multimeric glycan chains used as the substrate in NMR experiments. (D) CSPs calculated from NMR spectra of Tae1 with or without multimeric PG chains mapped onto the Tae1 surface structure. Numerous residues on both sides of the Tae1 display CSPs consistent with direct substrate interactions. (E) 1H-15N Best-Trosy NMR spectra from Tae1 alone (black) and Tae1 incubated with multimeric peptidoglycan chains (red) for six residues from A that show CSPs. Residues distal from the Tae1 active site show shifts indicative of interactions with PG. Numerous residues display CSPs consistent with direct substrate interactions. (F) HPLC traces showing degradation of peptidoglycan in the presence of wild type (black, WT) or mutant (red) Tae1 *in vitro*. Arrows indicate peaks containing the peptide stem that can be cleaved by Tae1. Mutant Tae1 (S144A and G48S) cleaves peptidoglycan less well than wild-type Tae1 (29.1 +/- 3.2% and 36.4 +/- 4.2% less PG., respectively) See also supplemental figure 2, 3, 4.

229 Consistent with DMS results, all nine variants showed
230 significantly less lytic activity than Tae1[WT] (**Figure 3B**).
231
LOF phenotypes could stem from a variety of
mechanisms, including reduced Tae1 protein
expression, disruption of protein folding or stability, or
direct inhibition of Tae1 PG binding or hydrolysis. To
rule out expression and possibly stability-dependent
effects, we performed Western blot analyses to
measure protein levels of Tae1 variants expressed in
E. coli (**Figure 3B**). We found that all variants were
expressed; five substitutions did not exhibit major
visible decreases in protein levels (G42S, S144A,
S147A, G48S, G56S). Our results led us to

Figure S2. Lysis assay growth curves. Cell growth as measured by OD_{600} for wild type Tae1 (WT), negative controls (Empty, C30A), and the nine distal residues on Tae1 in Figure 3B. Expressing Tae1 proteins containing the nine distal residue substitutions decreases cell lysis, suggesting these are loss-of-function residues.

243 hypothesize that these five surface substitutions may play direct roles in Tae1–PG interactions.
244
We hypothesized that an extended surface of Tae1 may be required for substrate binding and
hydrolysis *in vivo*. Given that Tae1 likely engages with larger and even sometimes insoluble fragments
of PG within the intact cell wall, we conducted a series of *in vitro* biochemical assays of LOF[clus] variants
using either intact PG sacculi or fragments that encompass more than the minimal peptide region. To
address whether distal surfaces of Tae1 bind PG, we experimentally mapped Tae1 binding sites by
two-dimensional (2D) nuclear magnetic resonance (NMR) using [15]N-labeled, catalytically inactive
Tae1[C30A] (Tae1[N15]) in complex with larger, soluble fragments of PG. Backbone [13]C, [15]N, and [1]HN NMR
resonances of Tae1 (>89% complete) were assigned using conventional heteronuclear techniques[21].
To prepare larger PG fragments for binding experiments, we digested *E. coli* cell wall sacculi using
another Tae superfamily member (Tae3 from *Salmonella* Typhi)[7], which hydrolyzes peptide crosslinks
resulting in long multimeric PG fragments that include both Tae cleavage sites and intact glycan chains
(**Figure 3C**) with an average length range of 10Å to over 300Å (1 to over 30 dimeric units)[22].
257
To map enzyme residues that bind cell wall substrate, we compared NMR spectra of Tae1[N15] alone or
in the presence of digested multimeric PG (**Supplemental Figure 3**). Observed spectral differences
allowed us to identify and quantitate chemical shift
perturbations (CSPs) across the protein, which serve as
indicators of residues involved in direct substrate
interactions[19,20]. Mapping these shifts onto the Tae1
structure, we observed CSPs for residues located within
the catalytic core as well as for a variety of both buried
and exposed residues outside of the core (**Figure 3D**).
While CSPs for buried residues could be due to
interaction-dependent enzyme conformational or
dynamical changes[23], contiguous clusters of CSPs
localized to the protein surface are suggestive of Tae1–
PG binding. Such putative binding regions extended all
the way to surfaces on the protein face opposite of the
catalytic core. In fact, we identified shifts in six of the
validated LOF[CLUS] residues: Leu21, Gly42, Gly48,
Ala52, Gly56, and Ser144 (**Figure 3E**), suggesting that
these residues are necessary for Tae1-induced lysis
because they bind PG.
278

Figure S3. [1]H-[15]N Best-Trosy NMR spectra of Tae1 with and without multimeric PG. NMR spectra from Tae1 alone (black) and Tae1 incubated with multimeric peptidoglycan chains (red). Numerous residues display CSPs consistent with direct substrate interactions.

To determine if these functional distal residues that bind PG are also required for hydrolysis, we
focused our studies on two highly expressed variants, Tae1^{S144A} and Tae1^{G48S}, for which we expressed
recombinant forms and purified. Size exclusion chromatography analysis of these variants showed
similar elution volumes and apparent hydrodynamic radii compared to Tae1^{WT}. We then conducted
degradative assays with PG sacculi purified from *E. coli*. Tae enzymes were incubated with PG sacculi,
and the resulting fragments were quantitatively assessed using high-performance liquid
chromatography (HPLC, **Figure 3F**). Tae1^{S144A} and Tae1^{G48S} cleaved 29.1 +/- 3.2% and 36.4 +/- 4.2%
less PG, respectively, than observed for the wild-type enzyme. Despite being relatively far from the
active site, these amino acid substitutions decreased Tae1 activity. Together, these data reveal that
numerous sites distributed around the Tae1 surface physically interact with PG, and that these distal
interaction sites are important for cell wall-degrading toxicity *in vivo*. Moreover, these results suggest
our DMS-based approach can faithfully uncover novel and unexpected functional determinants *in vivo*,
providing important clues to evolutionary pressures that may be acting on *tae1*.
292
293 **The functional landscape of Tae1 is sensitive to PG differences**
294
295 Our results point to a model in which Tae1 might interact with more structures of PG outside of the
296 catalytic site, and these interactions are important for *in vivo* toxicity. This model predicts that Tae
297 enzymes are sensitive to differences in cell wall architecture, and broad changes to PG composition

Figure 4. The functional landscape of Tae1 is sensitive to cell wall differences. (A) Cartoon illustration of cells grown in media containing D-Met showing altered PG architecture. D-Met becomes incorporated into the peptide stems and there are overall fewer crosslinks in the macromolecule. (B) Schematic of the deep mutational scanning experiments performed in media containing D-Met. DMS was completed as in Figure 1 with media containing D-Met. (C) Number of residues classified as each of the four cluster types in D-Met DMS screen. The functional classification of residues is dramatically changed from the original screen. (D) DMS-derived loss- and gain-of-function residues mapped onto the surface structure of Tae1 from the initial screen (left) and the D-Met screen (right). Compared with regular media, there were fewer loss-of-function and more gain-of-function residues in the presence of D-Met on both the catalytic face and back of the enzyme. (E) Heatmap representing average fitness scores for the top 50 GOF individual amino acid substitutions in D-Met and M9 media. n = 3. DMS in D-Met media was repeated 3 independent times, each time with at least 3 technical replicates. The majority of the GOF substitutions in D-Met are condition specific. (F) Lysis assays with wild-type Tae1, empty vector and catalytically inactive Tae1^{C30A} as negative controls, and 3 gain-of-function Tae1 variants (A116T, D134N, S147I). Data are means +/- standard deviation, n = 3, * p < 0.05. Lysis experiment was repeated 3 independent times, each time with at least 3 technical replicates. Western blots below measure Tae1 protein expression with RNAP as a control. All three gain-of-functions from DMS show increased lysis phenotype in individual lysis assays. (G) Lysis assays with the same Tae1 variants as in F but in M9 media. Western blots below measure Tae1 protein expression with RNAP as a control. The three D-Met GOF substitutions are condition specific. See also supplemental figure 5.

would alter Tae1–PG interactions. We hypothesized that such alterations could be captured by
comparative DMS analysis across different screening conditions. To address this hypothesis, we
conducted a second DMS screen in *E. coli* grown in the presence of D-methionine (D-Met), a condition
known for inducing major featural changes in cell wall composition and organization (**Figure 4A**).
Specifically, cell walls of *E. coli* grown in the presence of D-Met (*E. coli*[DMet]) have generally reduced PG
density and fewer modified PG crosslinks relative to cell walls of *E. coli* grown in the absence of D-
Met[24,25]. We verified these trends by HPLC analysis, and we confirmed that Tae1 maintained its ability
to cleave this distinct cell wall type *in vitro* (**Supplementary Figure 4**).

We performed the same suite of DMS analyses
for the second screen as for the first (**Figure 4B**)
and found that the calculated fitness landscape
shifted dramatically. Approximately 50% fewer
residues were categorized as LOF[clus] (38/86),
while the number of mixed residues increased
(94/56) as did the number of GOF[clus] residues
(18/8). The number of tolerant residues did not
change (4 in both) (**Figure 4C**). Overall, 58% of
residues were classified differently than in our
initial screen in M9 media. We mapped the
LOF[clus] and GOF[clus] residues onto the Tae1
structure (**Figure 4D**). We also saw that the
previous ring-like band of LOF[clus] determinants

Figure S4. D-met alters density and type of crosslinks. (A) HPLC traces comparing peptidoglycan from cells grown in either M9 and D-met media. (B) HPLC traces showing Tae1 still degrades peptide crosslinks containing D-met. (C) Summary of PG differences between M9 and D-met media. (D) There are no major morphological changes in the presence of D-met.

emanating from the catalytic groove was greatly diminished in *E. coli*[DMet]. Given that our biochemical
experiments suggest these sites are important for extended PG-binding, we hypothesize that such
contact interfaces may be less functionally critical in cellular conditions where the cell wall matrix is
more loosely packed around the enzyme. Alternatively, these changes could be indirectly due to other
metabolic or physical changes to the cell under D-Met growth conditions. Overall, our comparative
screening approach revealed that compositional changes to PG dramatically alter the functional
landscape of Tae1, suggesting these variants are sensitive to differences in cell wall architecture.

The *E. coli*[DMet] DMS screen also yielded a larger and distinct set of top-performing GOF[clus] variants with
amino acid substitutions that localized to both the catalytic groove and opposing faces of Tae1. When
we compared fitness scores for the GOF[clus] variants from the D-Met screen to the initial M9 screen, we
found that only a small subset of substitutions showed similar effects across both conditions (**Figure
4E**). Condition-specific phenotypes suggests that Tae1 toxicity *in vivo* is highly context-dependent and
optimizable for specific PG forms. To test this hypothesis, we conducted lysis assays in D-Met and M9
media for three GOF variants uniquely identified in the D-Met screen (**Figure 4F and G**). All three
substitutions increased cell lysis in *E. coli* grown in D-Met but not M9 minimal media. For two of the
variants, Asp134Asn and Ser147Ile, the increase in lysis may be due to higher expression levels based
on Western blot analysis (**Figure 4F and G**). Together, these results suggest Tae enzymes could
possibly be engineered to target specific bacteria with certain cell wall structures. Given that the
structure and composition of PG varies across bacterial species, our results also potentially point to a
selective pressure underlying natural sequence-based diversity in the Tae1 superfamily. Namely, cell
wall features associated with particular rival species a given bacterium must antagonize may drive
evolved substrate specificity differences in interbacterial competition toxins such as the Tae enzymes.

A hyperactive variant of Tae1 is naturally encoded by other *tae* homologs

A number of context-independent Tae1 variants that significantly out-performed wild-type enzyme
across both DMS conditions were also identified through our comparative approach. We were

particularly struck by one variant, Tae1^{C110S}, which is modified at a partially buried residue sitting
directly behind the catalytic Cys30 site (**Figure 5A**). The Cys110Ser substitution was the top-scoring
GOF variant in the D-Met screen and ranked 11th amongst GOF variants in the M9 screen. Using the *E.*
coli lysis assay, we first confirmed that Tae1^{C110S} indeed lysed cells at a faster rate than Tae1^{WT} in cells
grown in both D-Met and M9 minimal media (**Figure 5B**). To rule out indirect effects on toxicity caused
by increased protein levels, we assessed Tae1^{C110S} expression by Western blot analysis and did not
observe any increase relative to
Tae1^{WT} (**Figure 5B**).
357
The observation that Tae1 activity
could generally be tuned for greater
toxicity led us to ask why *tae1* from *P.*
aeruginosa does not naturally encode
Ser110. We considered several
explanations, none of which are
mutually exclusive. First, it is possible
that structural rearrangements
introduced by the Cys110Ser
substitution may disrupt Tae1
interactions with critical molecular
partners present in natural T6S-
mediated interbacterial interactions but
not in our DMS set-up. During T6S
competition, Tae1 from *P. aeruginosa*
must be delivered to neighboring cells
by a complex T6S secretion apparatus
that we bypassed with an engineered
secretion signal in our screen.
Furthermore, Tae1 injected into nearby
kin cells is deactivated through binding
by a cognate immunity protein, T6S
amidase immunity protein 1 (Tai1),
preventing lysis. Tai1 is specifically
encoded by *P. aeruginosa* and not *E.*
coli. Second, it is possible that we have
not sampled Tae1 fitness across
contexts that accurately represent
bacterial rivals or competition
conditions most relevant for *P.*
aeruginosa in the wild.

Figure 5. A hyperactive variant of Tae1 is naturally encoded by other *tae* homologs. (A) Ribbon structure diagram at the Tae1 catalytic site. Catalytic dyad residues Cys30 and His91 are indicated along with GOF Cys110. (B) Lysis assays with wild-type Tae1, empty vector and catalytically inactive Tae1^{C30A} as negative controls, and gain-of-function variant Tae1^{C110S} in D-Met and M9 media. Data are means +/- standard deviation, n = 3, * p < 0.05. Lysis experiment was repeated 3 independent times, each time with at least.3 technical replicates. Western blots below measure Tae1 protein expression with RNAP as a control. The C110S GOF substitution does is not sensitive to the D-Met induced changes in cell wall architecture. (C) Tae1 family phylogenetic tree. Species with Ser at position 110 are indicted in blue, while those with Cys are in black and those with Trp are in purple. Some but not all Tae1 proteins have evolved to include a Ser at position 110.

389
We therefore investigated whether Ser110 naturally occurs in any Tae1 representatives outside of *P.*
aeruginosa by searching for family members that encode a serine at residue positions comparable to
Cys110, as predicted by structure-based alignments. Surprisingly, we found that approximately half of
tae1-carrying bacterial species encoded serine residues at the Cys110 position, while some encoded
tryptophan (**Figure 5C**). To examine how these three variant classes are taxonomically distributed, we
generated a phylogenetic tree of all associated bacterial species and observed correlated divisions
across taxonomic clades and the residues at the Cys110 position, with the exception of one
Pseudomonas species (**Figure 5C**). Considering this observation, we hypothesized that *tae1* evolution
may have been locally constrained due to the requirement for Tai1-dependent immunity to prevent self-
antagonism.

Hyperactive variant Tae1^{C110S} evades binding and inhibition by cognate immunity

We reasoned that if Tae1 proteins are too structurally divergent, the toxin may evade binding by the cognate Tai1 partner in *P. aeruginosa* (Tai1^{PA}) and consequently antagonize kin cells or other closely related species. To assess whether the Tae1^{C110S} variant affects Tae1–Tai1 interactions, we first assessed whether Tae1^{C110S} adopted any major structural changes by solving its three-dimensional crystal structure (PDB ID: 7TVH) (**Table 2**). We did not observe any major structural rearrangements that would immediately implicate distinct modes of Tai1^{PA}-binding (RMSD=0.746, **Figure 6A**). However, we did observe some key side chain alterations near the Tae1^{C110S} active site, which is the enzyme region most intimately engaged with Tai1^{PA} upon binding (**Supplemental Figure 5**). Most notably, we found that His91 protruded away from the active site of Tae1^{C110S}, which would sterically clash with a loop of Tai1^{PA} that typically inserts into the catalytic cleft (**Figure 6B**). Our structural observations are consistent with the hypothesis that the Cys110 position is important for Tai1 recognition; therefore, serine at this position in *P. aeruginosa* may be disfavored to preserve immunity to the toxin. We

Figure 6. Hyperactive variant Tae1^{C110S} evades binding and inhibition by cognate immunity. (A) Tae1^{WT} ribbon structure diagram (silver) and Tae1^{C110S} ribbon structure diagram (blue) with the His91 side chain shown. C110S mutation does not cause major structural changes, but alters side chains in the active site. (B) Ribbon diagrams of the active sites of Tae1^{WT} (left) and Tae1C110S (right) in complex with Tai1 (pink). Side chains of Cys30, His91, and Cys110 are indicated in gray, and Ser110 in blue. His91 clashes with Tai1 in the active site. (C) Isothermal titration calorimetry determination of binding constants between wild-type Tae1 and Tai1 (left) and Tae1^{C110S} and Tai1 (right). Tae1^{C110S} binds poorly to Tai1. (D) Schematic depicting Tae1 and Tai1 interactions in wild-type *P. aeruginosa* and Tae1^{C110S} *P. aeruginosa*. When wild-type Tae1 is injected into a neighboring cell of the same species, Tai1 binds to it, conferring immunity. Tai1 cannot bind Tae1^{C110S}, thus the mutant Tae1 can cause cell lysis. (E) Fitness assays of *P. aeruginosa* expressing wild-type Tae1 or Tae1^{C110S} from the endogenous locus. *P. aeruginosa* lacking a Type 6 Secretion System (-T6SS) or expressing catalytically inactive Tae1 (C30A) were used as controls. Data are means +/- standard deviation, n = 3, * p < 0.05. Competition experiment was repeated 3 independent times, each time with at least.3 technical replicates. Cells expressing Tae1^{C110S} have decreased fitness when compared with wild-type Tae1. See also supplementary figures 6, 7.

generated a tree for the Tai1 protein family, similar to our tree for the Tae1 protein family. Consistent with this possibility, the tree for Tai1 has a similar branching pattern that is coincident with the residue variation at this position (**Supplemental Figure 6**).

We tested our prediction through a series of quantitative *in vitro* Tae1–Tai1 binding experiments. We measured Tai1-binding affinity (Kd) for recombinant Tae1^{WT} and Tae1^{C110S} proteins by isothermal titration calorimetry and found that Tae1^{WT} bound Tai1 with ~36-fold greater affinity than Tae1^{C110S} (9.5 nM Kd and 349.6 nM Kd, respectively) (**Figure 6C**). Toxin–immunity complexes are typically associated with binding affinities in the nM range; thus,

Figure S5. Tae1 – Tai1 interactions. (A) Ribbon diagram of Tae1 (silver) and Tai1 (pink). (B) Tae1 surface diagram, with red residues representing contacts with Tai1. The contacts are based on PDB ID 4EQA.

reduction may preclude the ability of Tai1 to effectively bind its toxin partner *in vivo*. To test whether
Tae1[C110S] evades Tai1-dependent immunity *in vivo*, we constructed a strain of *P. aeruginosa* harboring
this mutation (*PA[C110S]*) with a wild-type strain *(PA[WT])*. We reasoned that if Tae1[C110S] escapes Tai1-
binding and therefore immune protection, we would observe a decrease in *PA[C110S]* fitness compared to
a wild-type strain (*PA[WT]*) specifically when grown on solid media which induces T6S (**Figure 6D**).

Indeed, for *PA[C110S]* we observed substantially
reduced growth compared to *PA[WT]* on solid media
(**Figure 6E**) but not in liquid media, where T6S is
not induced (**Supplemental Figure 7**). This
phenotype strongly suggests that Tae1[C110S] lyses
kin cells at a faster rate than Tae1[WT] can bind and
inhibit killing, leading to a fitness cost for this
strain.

Together, our *in vitro* and *in vivo* results suggest
that the fitness cost of evading Tai1-binding likely
limits the proliferation of Tae1[C110S] in the wild. This
interaction presents an evolutionary bottleneck
that impedes certain toxin variants, such as
Tae1[C110S], from broadly taking hold across all
species. More generally, our DMS screening
approach allowed us to systematically probe
function across different *in vivo* contexts,
identifying two distinct but important selective
pressures – substrate-specificity and toxin–
immunity interactions – that may, in combination,
shape diversity across the Tae1 toxin superfamily.

Figure S6. Tai1 family phylogenetic tree. The residue at position 109 is either a Val (in Tai1s corresponding to Tae1_S110 proteins) or a Ser (in Tai1s corresponding to Tae1 non-S110 proteins). The tree contains the same species found in the Tae1 tree, except for *Mitsuaria* (unclassified) and *Pseudomonas* (multiple). Coloring of the species also matches that in the Tae1 tree. The sequences in the tree were identified using a published heuristic approach that specifically searches for T6S effector toxins. ML tree was inferred using PhyML v20160115 ran with model JTT and parameters: --alpha e -o tlr --bootstrap -2 -f m --nclasses 4 --pinv e. Branch supports are the Chi2-based parametric values returned by the approximate likelihood ratio test. Major clades had 90 or higher support values. The species marked with an asterisk have a Ser at position 110 in the corresponding Tae1 protein (for *Pseudomonas* sp. TC-11) or have an Ala at position 109 for the Tai1 protein (for *Singulisphaera acidiphila*).

DISCUSSION

The Tae superfamily of interbacterial toxins is phylogenetically, structurally, and functionally diverse[7,10].
Here, we show that sequence-based differences in *tae* homologs shape Tae proteins and their complex
substrate, the PG that constitutes the bacterial cell wall. Our studies revealed that Tae1 engages PG
through a circumferential interface that is more extensive than most proteases or other enzymes. This
unexpectedly large interaction surface may be key for dictating species-specific toxicity of Tae proteins
and consequently serve as a major driver of Tae superfamily diversification. This surface could also
mediate processivity across PG. Taes have many other binding partners as well. These toxins interact
with the T6SS apparatus for secretion, which likely limits effector size to a single, minimal catalytic
unit[4,26]. Therefore, unlike other larger multi-domain
housekeeping PG hydrolases[27], Tae toxins rely on a
relatively restricted set of surfaces to mediate many
distinct, functional interactions. Our unbiased, high
throughput DMS analyses revealed that determinants for
Tae1 cell wall-binding and immunity-binding partially
overlap. Our findings suggest that accommodation of
multiple, distinct functions likely constrains evolution of Tae
enzymes in nature despite intrinsic potential for enhanced
toxicity. Extensive Tae functional surfaces may also point
to the existence of additional unknown interaction partners
within bacteria. This principle may prove true for other T6S

Figure S7. Competition assay in liquid culture. Growth in *P. aeruginosa* expressing wild-type Tae1 or Tae1[C110S] from the endogenous locus in liquid culture. Data are means +/- standard deviation, n = 3. Growth experiment was repeated 3 independent times, each time with at least 3 technical replicates.

501 toxins and proteins; comprehensive mapping of functional determinants by DMS may power further
502 functional discovery.
503
Complementing traditional biophysical approaches with DMS allowed us to probe molecular
interactions in physiological states that are difficult to capture *in vitro*. Our current understanding of
enzyme interactions with the bacterial cell wall draws mostly from biochemical analyses conducted in
solution, posing a fundamental challenge for proteins with three-dimensional highly-crosslinked
macromolecular substrates. Considering the architectural complexity of the bacterial cell wall, a large
ring-like surface looping around Tae1 seems critical for toxicity *in vivo*. This observation could be due to
at least two possible modes of binding that we posit may be unique to enzymes with macromolecular
substrates. First, we theorize that a more extensive, non-linear substrate binding domain may enable
an enzyme to specifically recognize a particular conformational state. Three-dimensional configurations
of the cell wall may be an important feature for a toxin like Tae1 to detect, given that this could affect
protein localization to specific cell wall features that have distinct shapes. It has been previously
observed that a critical hole size in the cell wall is necessary for bacterial lysis[28]; thus localized action of
PG enzymes may provide an important advantage for Tae toxins. Another potential advantage of
binding a specific substrate configuration is that this would enable a Tae enzyme to distinguish between
different species or incorporation states of PG fragments. Indeed, certain endogenous enzymes, such
as FtsN interact exclusively with complex PG chains of a certain length[29].
520
Another possibility is that the large functional surface comprises several different PG-binding domains
that streamline substrate recognition or underlie enzyme processivity. This has been suggested for a
cell wall digesting enzyme, Lysostaphin from *Staphylococcus*, which exhibits two large PG-binding
surfaces on opposite sides of the enzyme[23]. Multi-step binding would support the "smart autolysins"
model postulated by Arthur Koch[30]. This model hints at the ability of hydrolase enzymes to recognize
"stretched" areas of PG. The resulting advantage for hydrolytic cell wall enzymes is that this type of
recognition decreases substrate search dimensionality from random 3-D diffusion to facilitated diffusion
by searching along the PG sacculus one plane at a time[31]. This may be particularly important for rapid,
effective recognition of substrates within a large cage-like 3D target like the cell wall, which is key for
potent, fast-acting toxins. This mechanism is also commonly adopted by DNA-binding proteins, which
dock nonspecifically to DNA before searching for specific cut sites[32,33]. Multi-step binding may also lead
to processive activity along PG by ensuring that Tae enzymes remain tethered to the cell wall substrate
after each hydrolysis event, which would allow for "stepwise chewing." This would be yet another
mechanism to increase the odds of localized, lytic activity.
535
In total, our combined use of biophysical and high throughput genetic approaches uncovered novel
insights into Tae toxin activity inside bacterial cells, providing a window into how small, single-domain
enzymes carry out several, distinct functions *in vivo*. Our leveraging of a multi-pronged strategy was
critical for overcoming technical obstacles inherent to proteins that interact with complex, insoluble
molecules, providing access to a class of macromolecular interactions that are not well understood. The
biophysical and kinetic parameters of proteins with three-dimensional highly-crosslinked
macromolecular substrates [34,35] are dissimilar from those well-established for small, soluble substrates,
and future adoption of combinatorial approaches such as ours may improve our understanding of
general binding principles across these important enzymes that are ubiquitous across life. Additionally,
the ability to quickly generate rich quantitative heatmaps for enzymes across different experimental
conditions enables more sophisticated computational analyses. Predictive representation learning of
unbiased, functional datasets across multiple biological conditions could help us anticipate *a priori* the
functional consequences of sequence-based phylogenetic diversity. A deeper understanding of
functional design principles would also enable more effective leveraging of evolutionary innovations to
engineer new molecules tailored to physiological contexts with important, useful applications.
551

ACKNOWLEDGEMENTS

We would like to thank Dr Doug Fowler (University of Washington) and Dr Eric Chow (University of California – San Francisco) for their help with setting up the deep mutational scanning pipeline. We are also grateful to Dr David Mavor (University of Massachusetts Chan Medical School), Dr Robert Newberry (University of Texas – Austin), Dr Martin Kampmann (University of California – San Francisco), and Dr James Fraser (University of California – San Francisco) for their assistance with sequencing data analysis. We say thanks to Dr Kari Herrington and Dr SoYeon Kim (University of California – San Francisco) for their contribution with microscopy data collection and analysis. We also thank Dr George Meigs and Dr James Holton (Lawrence Berkeley National Lab), and Dr Michael Thompson (University of California – Merced), for their help with x-ray crystallography data collection and analysis. We are grateful to Dr Mark Kelly for his help with setting up NMR data collection (University of California – San Francisco). We also thank to Dr Parnian Lak (University of California – San Francisco) for her help with setting up the ITC experiments. Lastly, we greatly appreciate the input regarding the experiments and the manuscript from members of the Chou lab at the University of California – San Francisco (Kristine Trotta, Calla Martyn, Fauna Yarza, Elizabeth A. Martinez-Bond, Patrick Grimes, Dr Domokos Lauko). Anne Sapiro received a Life Sciences Research Foundation fellowship from the SVCF-Wave Fund. Financial support from the IR-RMN-THC FR3050 CNRS for conducting the research on the 950 MHz NMR spectrometer is gratefully acknowledged. We would also like to thank the Chan Zuckerberg Biohub as well as the Sanghvi-Agarwal Innovation Award for their financial support.

COMPETING INTERESTS

The authors declare no competing interests.

METHODS

Docking analysis

Modeling was conducted with SwissDock using the default parameters[36]. The Tae1 monomer with PDB ID: 4EOB was used here[10]. The crosslinked PG dimer was constructed with MarvinSketch, based on ChEBI:61388. This viable pose was chosen based on the proximity of the active site nucleophile Cys30 to the scissile bond within the PG substrate.

Preparation of long chains of PG

PG isolation and HPLC analysis were conducted according to an established protocol[37] with the following modifications. Briefly, *E. coli* BW25113 cells were grown in 200ml LB media to mid-log phase. The cells were harvested by centrifugation at 5000 x g for 15min, after which they were suspended in 300μl of fresh LB media and added to 600μl 6% SDS solution in a 2ml Eppendorf tube. The cells were boiled for 3hrs in a heat block and then maintained in the SDS solution at 4°C until needed. The boiled sacculi were washed 5 times with MQ water to remove the SDS. Each wash consisted of 15min centrifugation at 21000 x g followed by decanting of the supernatant and resuspending in 1ml of MQ water by vortexing. After the final wash, the sacculi were suspended in Pronase E digestion buffer (10mM Tris-HCl pH 7.2 and 0.06% NaCl) and were treated for 2hrs at 60°C with activated Pronase E (100μg/ml final concentration, pre-treated for 2hrs at 60°C in the digestion buffer; supplier: VWR). The sacculi were incubated at 100°C for 10min to inactivate the Pronase E and were washed 3 times with 1ml of MQ water, with vortexing to resuspend. The sacculi were treated with purified Tae3 enzyme (10μM) overnight at 37°C in 20mM HEPES pH7.0 and 100mM NaCl buffer. The PG fragments were finally dialyzed using 500Da molecular weight cutoff tubing (Biotech CE Tubing, Spectra/Por) and dried on a Vacufuge concentrator to remove all liquid. They were stored at -80°C and were redissolved

602   before each NMR experiment with minimal volume to yield the desired enzyme to substrate
603   concentrations.
604
NMR experiments
NMR experiments for backbone assignments were performed at 25°C on a Bruker 950 MHz AVANCE
(Institut de Biologie Structurale in Grenoble) or Varian INOVA spectrometers (600 and 800 MHz
instruments at Pacific Northwest National Laboratories). NMR samples for determination of Tae1
resonance assignments consisted of 0.3 to 0.6 mM [^{13}C,^{15}N]-Tae1 in citrate buffer pH 5.5 with 150 mM
NaCl, 100 μM EDTA containing 5% D$_2$O. Assignment of backbone resonances was accomplished by
analysis of standard triple-resonance experiments (^{1}H,^{15}N)-HSQC, HNCA, HNCOCA, HNCACB,
CBCA(CO)NH, and HNCO spectra[21]). Assignments of Tae1 resonances in 20mM HEPES or NaP$_i$,150
mM NaCl, 100 μM EDTA at pH 7.0 were determined by collecting a series of [^{1}H,^{15}N]-HSQC spectra
titrating the sample from pH 5.5 to 7.0. These assignments were confirmed by collecting an HNCO
spectrum under the final conditions at pH 7.0. NMR data were processed and analyzed using
NMRPipe[38] and NMRView[39,40] (for the residue assignments, see **Supplemental Figure 8**). See
Supplemental Table 2 for the peak values and **Supplemental Table 3** for CSP ($\Delta\delta$) values.
618

Figure S8. (^{1}H,^{15}N)-TROSY spectrum of 150mM Tae1 in 20mM HEPES, 100mM NaCl, pH 7.0 showing resonance assignments for backbone amides. B) Expanded view of a smaller region of the same spectrum.

619
All titration experiments were conducted at the Institut de Biologie Structurale in Grenoble or at the
UCSF NMR facility in Genentech Hall (San Francisco). PG ligand was dialyzed against water using a
float dialysis membrane device with a 100-500 Da cut-off (Spectrum Laboratories, Inc) and lyophilized
before preparation of a concentrated stock solution in NMR buffer. Before aliquoting the samples or
before lyophilization, the concentration of the stock solution was estimated by liquid-state NMR using
the Eretic pulse sequence[41] and a reference 1 mM sucrose sample in 90%:10% H$_2$O:D$_2$O. Interaction
studies with ^{15}N-labelled Tae1 and PG substrates were monitored by superimposition of ^{15}N-BEST-
TROSY spectra at 25°C for different substrate-to-protein ratio[42]. A 50μM ^{15}N labeled Tae1_C30A was
titrated with 500 and 1000 μM of purified muropeptide. CcpNmr2.4 was used to monitor protein CSP for
every assigned amide resonance by superimposition of the ^{15}N-BEST-TROSY spectra. CSPs ($\Delta\delta$)
were calculated on a per-residue basis for the highest substrate-to-protein ratio.
631
E. coli lysis assays
The pBAD vector was used to clone Tae1 variants of interest[43]. The Tae1 wild-type was initially
amplified from a previously published pET22::Tae1 construct[10], together with the pelB signal sequence.
All variants were constructed based on this pelB_pBAD::Tae1 template. See **Table 1** for all the primers

used to clone Tae1 and the different variants. SOE-PCR was used to introduce the individual
mutations, followed by restriction enzyme digestion and ligation. pBAD::Tae1 constructs were always
transformed freshly into *E. coli* BW25113 chemically competent cells and a LB culture with antibiotic
was immediately started after the transformation outgrowth period. The culture was diluted 1 to 100 on
the following day into LB media in the presence of antibiotic selection. After reaching OD_{600} of 0.2,
protein expression was induced with 0.25% arabinose and cell lysis was monitored overnight on a
Biotek plate reader.
643
DMS library construction and assay
pET22b vector was chosen for library construction. This vector contains the pelB signal sequence that
would allow the expression of Tae1 variants in the cell periplasm. The preparation of variants was
achieved through an error-prone PCR mutagenesis kit (GeneMorph II Random Mutagenesis Kit –
Agilent Technologies), following the manufacturer's instructions in order to insure high proportion of
single-mutant variants. The variants were cloned via restriction enzyme cloning. In order to track the
behavior of variants, each one was barcoded with a unique DNA sequence. The barcodes were
introduced immediately after the Tae1 coding sequence via Gibson assembly cloning (In-Fusion HD
Cloning Kit – Takara Bio). Our Next-Generation sequencing strategy comprised two stages. First, the
entire library was sequenced to determine the barcode corresponding to each variant. This was
accomplished using 300bp forward and reverse reads on the MiSeq Illumina platform (MiSeq Reagent
Kit v3). Using this information, a dictionary was prepared to demonstrate the relationships between
variants and barcodes. The dictionary was prepared by following previously published scripts[44] with
some modification (see the GitHub directory "ChouLab dictionary" for the steps and scripts). For each
of the subsequent DMS experiments, 25bp reads on a NextSeq Illumina platform (conducted at Chan
Zuckerberg BioHub) were used to sequence only the barcode region. For data analysis and DMS
scores calculations, we used Enrich2 software[17]. The barcode dictionary was used to convert barcodes
to variants. Please see the "Choulab DMS data analysis" GitHub directory for detailed information on
data analysis in Enrich2. For the DMS experiments in this study, the DMS library was grown for 3hrs at
37°C in M9 minimal medium (or M9 supplemented with 20mM D-Methionine) before distributing across
a 96-well plate. A set of 24 wells was combined for each experimental time point in order to obtain
enough DNA for library amplification. Libraries were amplified for 9 cycles using Phusion high-fidelity
polymerase to minimize any masking of differences in abundance. The libraries were gel purified,
quantified using Qubit, and their concentrations were adjusted according to Illumina instructions before
sequencing. The 0hrs time point (DMS library before any growth) was chosen as the input of the 96-
well plate incubation. After growth to OD_{600} 0.2, protein expression was induced with 1mM IPTG. Cells
were collected 2hrs after induction that would serve as the output time point. All of the scripts used in
the paper are available on GitHub at https://github.com/AtanasDRadkov/ChouLab_DMS. The data files
are available on NCBI, Sequence Read Archive – PRJNA803461.
673
Clustering analysis
K-means clustering analysis was conducted using previously published scripts[18] to group Tae1
positions into categories based on their fitness scores. We calculated spatial clusters by 1) sorting the
fitness score vectors for each position 2) trimming the vectors to match vector lengths (discarding no-
data values) 3) calculating a difference by subtracting the two sorted and trimmed vectors, and 4)
calculating the distance – mean of the absolute value of the vector difference. The distance between a
candidate position and a cluster of positions is calculated as the average of the distance between the
candidate position and the three closest non-self positions in the cluster. Clustering was performed over
10 rounds following the initial seeded round, and convergence was confirmed by observing that five
repetitions gave identical clusters. The results yielded five clusters, including a loss-of-function
phenotype cluster which was the focus of the study. The same analysis was performed for both DMS
experiments. See the GitHub directory "ChouLab clustering" for the steps and scripts. All of the scripts
used in the paper are available on GitHub at https://github.com/AtanasDRadkov/ChouLab_DMS.

687
HPLC PG analysis
PG isolation and HPLC analysis were conducted as described[37] with the following modifications. Briefly,
E. coli BW25113 cells were grown in 200ml LB (or LB plus 20mM D-Met) to mid-log phase. The cells
were harvested by centrifugation at 5000 x g for 15min, after which they were suspended in 300μl fresh
LB and added to 600μl of 6% SDS solution in a 2ml Eppendorf tube. The cells were boiled for 3hrs in a
heat block and then maintained in the SDS solution at 4°C until needed. Before any treatments, the
boiled sacculi were washed 5 times with MQ water to remove the SDS. Each wash consisted of 15min
centrifugation at 21000 x g followed by decanting of the supernatant and resuspending in 1ml MQ water
by vortexing. After the final wash, the sacculi were suspended in Pronase E digestion buffer (10mM
Tris-HCl pH 7.2 and 0.06% NaCl) and were treated for 2hrs at 60°C with Pronase E (100μg/ml final
concentration, pre-treated for 2hrs at 60C in the digestion buffer; supplier: VWR). The sacculi were
incubated at 100°C for 10min to inactivate the Pronase E. Enzymatic treatments with Tae1 variant
enzymes were conducted at this point using 20mM HEPES pH7.0, 100mM NaCl buffer. Tae1 enzyme
was added to 1μM final concentration and the reactions were incubated at room temperature for 10min,
followed by 10min at 100°C to inactivate the enzyme. The sacculi were washed 3 times as described
above. Finally the sacculi were solubilized using 25μg/ml final concentration of mutanolysin (Sigma-
Aldrich) in 10mM Tris-HCl pH 7.2 and 0.06% NaCl after overnight digestion in a 37°C shaker. To
prepare PG fragments for HPLC analysis, the mutanolysin treatment was incubated at 100°C for 10min,
followed by centrifugation to remove any debris, alkalinization with 0.5M borate buffer and reduction
with sodium borohydride for 1hr at room temperature. The reduced PG fragments were acidified to pH
3-4 with 50% o-phosphoric acid and filtered (PVDF syringe filters, 0.22μm, 0.4mm - Tisch Scientific)
before the HPLC injection. The HPLC instrument, method, and buffers were the same as described
previously[45].
711
Protein purification
All Tae proteins, and Tai1 protein, used in this study were purified according to an established
protocol[10] using pET29b+ vector and 3hrs induction in a 37°C shaker with 1mM IPTG. Buffers for metal
affinity chromatography were 20mM HEPES pH7, 100mM NaCl, 20mM imidazole and 20mM HEPES
pH7, 100mM NaCl, 400mM imidazole. Buffer for size exclusion chromatography (SEC) was 20mM
HEPES pH7, 100mM NaCl. The SEC was performed using an isocratic gradient on an AKTA pure
instrument with a HiPrep 16/60 Sephacryl S-200 HR column. For isotopically labeled Tae1_C30A, the
same plasmid constructs were used, but the expression was conducted in M9 minimal medium
prepared with ^{15}N-ammonium chloride (5g per liter media) (M9 minimal medium prepared according to
Cold Spring Harbor Protocols). The cells were grown and induced as described above. The same
purification procedure was used as described above.
723
Tae1 phylogenetic tree
Initial candidates of the Tae1 family as identified in Russell et al.[7] were aligned and used as a seed in a
PSI-BLAST search. Significant results were downloaded and aligned using MAFFT. Candidates
containing the critical Cys-His dyad were kept, and other candidates discarded. Final candidates were
re-aligned. Default parameters were used for all steps. Multisequence alignment (ClustalW algorithm)
and phylogenetic trees (PhyML analysis) were prepared on GenomeNet (at www.genome.jp).
730
P. aeruginosa strain construction
P. aeruginosa strains used in this study were derived from the sequenced strain PAO1[46]. *E. coli* strain
SM10 was used for conjugal transfer of pEXg2 plasmids for homologous recombination as described
previously[5]. All deletions were in-frame and were generated by exchange with deletion alleles
constructed by SOE PCR, followed by restriction enzyme cloning into pEXg2 vector. See **Table 1** for
the primers used to delete genes *pppA* and *retS*, and to introduce the C110S mutation into *tae1*.
737

Contact-dependent fitness assays

Fitness assays were performed with overnight *P. aeruginosa* cultures. The assays were performed on solid media (LB without added NaCl containing 3% agar) using nitrocellulose membranes (Amersham Biosciences Hybond-N+, 0.45um) to insure close contact between the cells[5]. The OD_{600} was adjusted to 1.0 for all strains. 10μl were spotted onto a membrane to initiate the contact-dependent fitness assays. The assays were incubated for 6 hrs at 37°C after which the spots were removed with a razor blade, placed in a tube with glass beads, shaken on a bead beater (BeadBug microtube homogenizer; Z763705, Sigma) for 20sec at 4000rpm, and the suspended cells were distributed across a 96-well plate (3 x 200μl wells for each competition tube) for OD_{600} measurements on a Biotek plate reader.

Isothermal titration calorimetry

Isothermal titration calorimetry experiments were performed at 298°K using a Microcal ITC200 (GE Healthcare, Northampton, MA). Stock solutions of Tae1 and Tai1 were dialyzed overnight at 277°K in the same buffer (50mM NaCl, 20mM Tris 7.0). The cell was filled with 200μL of Tai1 protein at a concentration of 20μM to which 150μM of Tae1 protein was titrated. For 25 titrations of 1.5μL of Tae1 into Tai1 the heat of binding was recorded. Data were extracted and fitted using the single-site binding model following manufacturer's instructions.

Tae1^{C110S} crystal structure

Tae1^{C110S} protein crystals were generated by hanging drop vapor diffusion at 25°C from a 1:1 mixture of 10 mg/ml protein (in 20mM HEPES pH7, 100mM NaCl, 5% glycerol) with 0.2 M sodium thiocyanate, 0.1 M HEPES pH 7.5, 20% PEG3350 for 24hrs (same conditions that were used for initial structure of Tae1[10]). Crystals were directly used for diffraction data collection at the Lawrence Berkeley National Laboratory Advanced Light Source Beamline 8.3.1[47]. Coot[48] and Maximum-Likelihood[49] refinement with PHENIX were used for iterative building and refinement. Structure was validated by MOLProbity[50] and visualized with Chimera[51]. Coordinates and structure factors were deposited in the Protein Data Bank (PDB ID: 7TVH).

Tables

791 **Table1.** Primers used in this study

Name	Sequence
pET22_Tae1_C110S_F1	CCGATGTGCTGGTCCGGCAGCATCGCCGGC
pET22_Tae1_C110S_F2	GGTGGTGGTGCTCGAG
pET22_Tae1_C110S_R1	GCCGGCGATGCTGCCGGACCAGCACATCGG
pET22_Tae1_C110S_R2	TGGATATCGGAATTAATTCGGATC
pET29_Tae1_C110S_F1	CCGATGTGCTGGTCCGGCAGCATCGCCGGC
pET29_Tae1_C110S_F2	GGTGGTGGTGCTCGAG
pET29_Tae1_C110S_R1	GCCGGCGATGCTGCCGGACCAGCACATCGG
pET29_Tae1_C110S_R2	GGGAATTCCATATGGACAGTCTCGATCAATGC
pBAD_Tae1_F	TAGTACAGAGAATTCACCATGAAATACCTGCTGCCGACCGCTGCTGC
pBAD_Tae1_R	TCAATCAGTATCTAGATTACTCGAGACTGGCCCTGGGCAGGCTG
pEXG2_pppA_F1	TGTTAAGCTAAAGCTTGTCGGTCGCTATTTCCCGCTGAC
pEXG2_pppA_F2	GCATACCCGGTTAAGGACAACTGATGGTGAACTCGAGCCGCAAGCATGCTGAA
pEXG2_pppA_R1	TCAATCAGTATCTAGAGCCGGTGAGGATCTCGTAGAGCAC
pEXG2_pppA_R2	CCGGAAGTTCTATGTCCATGTCCGTCTCAGAATTCAGCATGCTTGCGGCTCGAGTT
pEXG2_retS_F1	TCAATCAGTATCTAGACGGAAACCCAGCCGATCATGG
pEXG2_retS_F2	AACTCGAGCCGCAAGCATGCTGAAGGCGAAGTCCCTTCGAAGGTG
pEXG2_retS_R1	TGTTAAGCTAAAGCTTCCAGCATCTTCAGGTAGACGC
pEXG2_retS_R2	TTCAGCATGCTTGCGGCTCGAGTTGCCTACTGGGTCGGCGAACGC
pEXG2_Tae1C110S_F1	TCAAGTACTAGAGCTCGTGGTGCACCGCGAGGACATCTC
pEXG2_Tae1C110S_F2	CTGCCACTCCAGCACATCGGGTAC
pEXG2_Tae1C110S_R1	TATCAGAAACCTGCAGGTAGGCCAGCGACTGCATGCCGTAG
pEXG2_Tae1C110S_R2	GATGTGCTGGAGTGGCAGCATCGCC
pBAD_Tae1L21S_F	AGCTGGGACAAGAGCTACAGCGCCGGCACCCCG
pBAD_Tae1L21S_R	CTTGTTCGGGGTGCCGGCGCTGTAGCTCTTGTC
pBAD_Tae1G48S_F	GGCGTACCGATGCCCCGCAGCAACGCCAACGCC
pBAD_Tae1G48S_R	GACCATGGCGTTGGCGTTGCTGCGGGGCATCGG
pBAD_Tae1G56S_F	GCCAACGCCATGGTCGACAGCCTGGAGCAGAGC
pBAD_Tae1G56S_R	GGTCCAGCTCTGCTCCAGGCTGTCGACCATGGC
pBAD_Tae1G42S_F	TCGGTGGCCGCCGAGCTGAGCGTACCGATGCCC
pBAD_Tae1G42S_R	GCCGCGGGGCATCGGTACGCTCAGCTCGGCGGC
pBAD_Tae1V43S_F	GTGGCCGCCGAGCTGGGCAGCCCGATGCCCCGC
pBAD_Tae1V43S_R	GTTGCCGCGGGGCATCGGGCTGCCCAGCTCGGC
pBAD_Tae1A52P_F	CCCCGCGGCAACGCCAACCCGATGGTCGACGGC
pBAD_Tae1A52P_R	CTCCAGGCCGTCGACCATCGGGTTGGCGTTGCC
pBAD_Tae1S144A_F	CTCAACTACTACGTCTACGCCCTGGCCAGTTGC
pBAD_Tae1S144A_R	CAGGCTGCAACTGGCCAGGGCGTAGACGTAGTA
pBAD_Tae1S147A_F	TACGTCTACTCCCTGGCCGCCTGCAGCCTGCCC
pBAD_Tae1S147A_R	GGCCCTGGGCAGGCTGCAGGCGGCCAGGGAGTA
pBAD_Tae1S149A_F	TACTCCCTGGCCAGTTGCGCCCTGCCCAGGGCC
pBAD_Tae1S149A_R	GAGACTGGCCCTGGGCAGGGCGCAACTGGCCAG
pBAD_Tae1A116T_F	TGCGGCAGCATCGCCGGCACTGTCGGCCAGAGC
pBAD_Tae1A116T_R	GCCCTGGCTCTGGCCGACAGTGCCGGCGATGCT
pBAD_Tae1D134N_F	CAGGTGTGGAATCGCACCAACCGCGACCGCCTC
pBAD_Tae1D134N_R	GTAGTTGAGGCGGTCGCGGTTGGTGCGATTCCA
pBAD_Tae1S147I_F	TACGTCTACTCCCTGGCCATTTGCAGCCTGCCC
pBAD_Tae1S147I_R	GGCCCTGGGCAGGCTGCAAATGGCCAGGGAGTA
DMS_Lib_Amplify_F	AATGATACGGCGACCACCGAGATCTACACCTTTCGGGCTTTGTTAGCAG
DMS_Lib_Amplify_R1	CAAGCAGAAGACGGCATACGAGATGGAACGGAGTGACTGGAGTTCAGACGTGTGC TCTTCCGATCTTCGATAGTAGCCTATGAAGCATGCTTACTA

DMS_Lib_Amplify_R2	CAAGCAGAAGACGGCATACGAGATCGGCACATGTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTTCGATAGTAGCCTATGAAGCATGCTTACTA
DMS_Lib_Amplify_R3	CAAGCAGAAGACGGCATACGAGATTACCCTTGGTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTTCGATAGTAGCCTATGAAGCATGCTTACTA
DMS_Lib_Amplify_R4	CAAGCAGAAGACGGCATACGAGATTAGGTGTCGTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTTCGATAGTAGCCTATGAAGCATGCTTACTA
DMS_Lib_Amplify_R5	CAAGCAGAAGACGGCATACGAGATGAATCACTGTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTTCGATAGTAGCCTATGAAGCATGCTTACTA
DMS_Lib_Amplify_R6	CAAGCAGAAGACGGCATACGAGATCATAGAGTGTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTTCGATAGTAGCCTATGAAGCATGCTTACTA
DMS_custom_seq_F	GGTGGTGCTCGACGATAGCACGTCTCTGACACCGAATTC
DMS_custom_seq_R	GCCCAGCCGGCGATGGCCATGGATATCGGAATTAATTC

829 **Table2.** X-ray data collection and refinement statistics for Tae1[C110S] (PDB ID: 7TVH)

	Crystal 1	831
Data collection		
Space group	P 1 21 1	
Cell dimensions		
a, b, c (Å)	39.121 108.512 82.402	
Resolution (Å)	65.56 - 1.71 (1.77 - 1.71)	
R_{sym} or R_{merge}	0.03896 (0.9019)	
I / I	10.48 (0.80)	
Completeness (%)	98.45 (87.02)	
Redundancy	2.00 (2.00)	
$CC_{1/2}$	0.999 (0.469)	
Refinement		
Resolution (Å)	65.56 - 1.71	
No. reflections	72921 (6436)	
R_{work} / R_{free}	0.2113 / 0.2430	
No. atoms	4673	
Protein	4277	
Water	396	
B-factors		
Protein	38.21	
Water	43.00	
R.m.s. deviations		
Bond lengths (Å)	0.006	
Bond angles	0.77	
Ramachandran favored (%)	98.09	
Ramachandran allowed (%)	1.91	
Ramachandran outliers (%)	0	
Rotamer outliers	0	

832 *Values in parentheses are for highest-resolution shell.

References:

1. Peterson, S. B., Bertolli, S. K. & Mougous, J. D. The central role of interbacterial antagonism in bacterial life. *Current Biology* **30**, R1203–R1214 (2020).

2. Jani, A. J. & Cotter, P. A. Type VI Secretion: Not just for pathogenesis anymore. *Cell Host & Microbe* **8**, 2–6 (2010).

3. Schwarz, S., Hood, R. D. & Mougous, J. D. What is type VI secretion doing in all those bugs? *Trends in Microbiology* **18**, 531–537 (2010).

4. Russell, A. B., Peterson, S. B. & Mougous, J. D. Type VI secretion system effectors: poisons with a purpose. *Nat Rev Microbiol* **12**, 137–148 (2014).

5. Russell, A. B. *et al.* Type VI secretion delivers bacteriolytic effectors to target cells. *Nature* **475**, 343–347 (2011).

6. Hood, R. D. *et al.* A Type VI Secretion System of *Pseudomonas aeruginosa* targets a toxin to bacteria. *Cell Host & Microbe* **7**, 25–37 (2010).

7. Russell, A. B. *et al.* A widespread bacterial Type VI Secretion effector superfamily identified using a heuristic approach. *Cell Host & Microbe* **11**, 538–549 (2012).

8. Russell, A. B. *et al.* A Type VI Secretion-related pathway in Bacteroidetes mediates interbacterial antagonism. *Cell Host & Microbe* **16**, 227–236 (2014).

9. LaCourse, K. D. Conditional toxicity and synergy drive diversity among antibacterial effectors. *Nature Microbiology* **3**, 9 (2018).

10. Chou, S. *et al.* Structure of a peptidoglycan amidase effector targeted to Gram-negative bacteria by the Type VI Secretion System. *Cell Reports* **1**, 656–664 (2012).

11. Vollmer, W., Blanot, D. & De Pedro, M. A. Peptidoglycan structure and architecture. *FEMS Microbiol Rev* **32**, 149–167 (2008).

12. Vollmer, W. & Seligman, S. J. Architecture of peptidoglycan: more data and more models. *Trends in Microbiology* **18**, 59–66 (2010).

13. Furchtgott, L., Wingreen, N. S. & Huang, K. C. Mechanisms for maintaining cell shape in rod-

shaped Gram-negative bacteria: Rod-shape maintenance in Gram-negative bacteria. *Molecular*

Microbiology **81**, 340–353 (2011).

14. Elazar, A. *et al.* Mutational scanning reveals the determinants of protein insertion and association

energetics in the plasma membrane. *eLife* **5**, e12125 (2016).

15. Barad, B. A. *et al.* Differences in the chitinolytic activity of mammalian chitinases on soluble and

insoluble substrates. *Protein Science* **29**, 952–963 (2020).

16. Fowler, D. M. & Fields, S. Deep mutational scanning: a new style of protein science. *Nat Methods*

11, 801–807 (2014).

17. Rubin, A. F. *et al.* A statistical framework for analyzing deep mutational scanning data. *Genome*

Biol **18**, 150 (2017).

18. Thompson, S., Zhang, Y., Ingle, C., Reynolds, K. A. & Kortemme, T. Altered expression of a quality

control protease in *E. coli* reshapes the in vivo mutational landscape of a model enzyme. *eLife* **9**,

e53476 (2020).

19. Schanda, P. *et al.* Atomic model of a cell-wall cross-linking enzyme in complex with an intact

bacterial peptidoglycan. *J. Am. Chem. Soc.* **136**, 17852–17860 (2014).

20. Maya-Martinez, R. *et al.* Recognition of peptidoglycan fragments by the transpeptidase Pbp4 from

Staphylococcus aureus. Front. Microbiol. **9**, 3223 (2019).

21. Sattler, M. Heteronuclear multidimensional NMR experiments for the structure determination of

proteins in solution employing pulsed field gradients. *Progress in Nuclear Magnetic Resonance*

Spectroscopy **34**, 93–158 (1999).

22. Harz, H., Burgdorf, K. & Höltje, J.-V. Isolation and separation of the glycan strands from murein of

Escherichia coli by reversed-phase high-performance liquid chromatography. *Analytical*

Biochemistry **190**, 120–128 (1990).

23. Gonzalez-Delgado, L. S. *et al.* Two-site recognition of *Staphylococcus aureus* peptidoglycan by

lysostaphin SH3b. *Nat Chem Biol* **16**, 24–30 (2020).

24. Caparrós, M., Pisabarro, A. G. & de Pedro, M. A. Effect of D-amino acids on structure and

synthesis of peptidoglycan in *Escherichia coli*. *J Bacteriol* **174**, 5549–5559 (1992).

25. Lam, H. *et al.* D-amino acids govern stationary phase cell wall remodeling in bacteria. *Science* **325**,

1552–1555 (2009).

26. Basler, M., Pilhofer, M., Henderson, G. P., Jensen, G. J. & Mekalanos, J. J. Type VI Secretion

requires a dynamic contractile phage tail-like structure. *Nature* **483**, 182–186 (2012).

27. Vollmer, W., Joris, B., Charlier, P. & Foster, S. Bacterial peptidoglycan (murein) hydrolases. *FEMS*

Microbiol Rev **32**, 259–286 (2008).

28. Daly, K. E., Huang, K. C., Wingreen, N. S. & Mukhopadhyay, R. Mechanics of membrane bulging

during cell-wall disruption in Gram-negative bacteria. *Phys. Rev. E* **83**, 041922 (2011).

29. Ursinus, A. *et al.* Murein (peptidoglycan) binding property of the essential cell division protein FtsN

from *Escherichia coli*. *J Bacteriol* **186**, 6728–6737 (2004).

30. Koch, A. L. Additional arguments for the key role of "smart" autolysins in the enlargement of the wall

of Gram-negative bacteria. *Research in Microbiology* **141**, 529–541 (1990).

31. Kari, J., Andersen, M., Borch, K. & Westh, P. An inverse Michaelis–Menten approach for interfacial

enzyme kinetics. *ACS Catal.* **7**, 4904–4914 (2017).

32. Terry, B. J., Jack, W. E. & Modrich, P. Facilitated diffusion during catalysis by EcoRI endonuclease.

Nonspecific interactions in EcoRI catalysis. *Journal of Biological Chemistry* **260**, 13130–13137

(1985).

33. Stracy, M. *et al.* Transient non-specific DNA binding dominates the target search of bacterial DNA-

binding proteins. *Molecular Cell* **81**, 1499-1514.e6 (2021).

34. Broendum, S. S. *et al.* High avidity drives the interaction between the streptococcal C1 phage

endolysin, PlyC, with the cell surface carbohydrates of Group A *Streptococcus*. *Mol Microbiol* **116**,

397–415 (2021).

35. McLaren, A. D. & Packer, L. Some aspects of enzyme reactions in heterogeneous systems. in

Advances in enzymology and related areas of molecular biology (ed. Nord, F. F.) vol. 33 245–308

(Interscience Publishers, 1970).

36. Grosdidier, A., Zoete, V. & Michielin, O. SwissDock, a protein-small molecule docking web service

based on EADock DSS. *Nucleic Acids Research* **39**, W270–W277 (2011).

37. Stankeviciute, G. *et al.* Differential modes of crosslinking establish spatially distinct regions of

peptidoglycan in *Caulobacter crescentus*. *Mol Microbiol* **111**, 995–1008 (2019).

38. Delaglio, F. *et al.* NMRPipe: A multidimensional spectral processing system based on UNIX pipes.

J Biomol NMR **6**, (1995).

39. Johnson, B. A. & Blevins, R. A. NMR View: A computer program for the visualization and analysis

of NMR data. *J Biomol NMR* **4**, 603–614 (1994).

40. Downing, A. K. *Protein NMR Techniques*. (Humana Press, 2004).

41. Frank, O., Kreissl, J. K., Daschner, A. & Hofmann, T. Accurate determination of reference materials

and natural isolates by means of quantitative ^{1}H NMR spectroscopy. *J. Agric. Food Chem.* **62**,

2506–2515 (2014).

42. Favier, A. & Brutscher, B. Recovering lost magnetization: polarization enhancement in biomolecular

NMR. *J Biomol NMR* **49**, 9–15 (2011).

43. Guzman, L. M., Belin, D., Carson, M. J. & Beckwith, J. Tight regulation, modulation, and high-level

expression by vectors containing the arabinose P_{BAD} promoter. *J Bacteriol* **177**, 4121–4130 (1995).

44. Mavor, D. *et al.* Determination of ubiquitin fitness landscapes under different chemical stresses in a

classroom setting. *eLife* **5**, e15802 (2016).

45. Hayes, B. M. *et al.* Ticks resist skin commensals with immune factor of bacterial origin. *Cell* **183**,

1562-1571.e12 (2020).

46. Stover, C. K. *et al.* Complete genome sequence of *Pseudomonas aeruginosa* PAO1, an

opportunistic pathogen. *Nature* **406**, 6 (2000).

47. MacDowell, A. A. *et al.* Suite of three protein crystallography beamlines with single superconducting

bend magnet as the source. *J Synchrotron Rad* **11**, 447–455 (2004).

48. Emsley, P. & Cowtan, K. *Coot*: model-building tools for molecular graphics. *Acta Crystallogr D Biol*

Crystallogr **60**, 2126–2132 (2004).

49. Adams, P. D. *et al. PHENIX*: a comprehensive Python-based system for macromolecular structure

solution. *Acta Crystallogr D Biol Crystallogr* **66**, 213–221 (2010).

50. Davis, I. W., Murray, L. W., Richardson, J. S. & Richardson, D. C. MOLPROBITY: structure

validation and all-atom contact analysis for nucleic acids and their complexes. *Nucleic Acids*

Research **32**, W615–W619 (2004).

51. Pettersen, E. F. *et al.* UCSF Chimera - A visualization system for exploratory research and

analysis. *J. Comput. Chem.* **25**, 1605–1612 (2004).

963