Title: Experimental infection of mink with SARS-COV-2 Omicron (BA.1) variant leads to

symptomatic disease with lung pathology and transmission

Authors: Jenni Virtanen[1], Kirsi Aaltonen[1], Kristel Kegler, Vinaya Venkat, Thanakorn Niamsap,

Lauri Kareinen, Rasmus Malmgren, Olga Kivelä, Nina Atanasova, Pamela Österlund, Teemu

Smura, Antti Sukura, Tomas Strandin, Lara Dutra, Olli Vapalahti, Heli Nordgren, Ravi Kant, Tarja

Sironen

[1]These authors contributed equally to this article

Author affiliations: University of Helsinki, Helsinki, Finland (J. Virtanen, K. Aaltonen, K. Kegler,

V. Venkat, T. Niamsap, L. Kareinen, R. Malmgren, O. Kivelä, N. Atanasova, T. Smura, A. Sukura,

T. Strandin, L. Dutra, O. Vapalahti, H. Nordgren, R. Kant, T. Sironen); Finnish Meteorological

Institute, Helsinki (N. Atanasova); Finnish Institute for Health and Welfare, Helsinki (P. Österlund);

Helsinki University Hospital, Helsinki (O. Vapalahti)

Abstract

We report an experimental infection of American mink with SARS-CoV-2 Omicron variant and

show that minks remain virus RNA positive for days, develop clinical signs and histopathological

changes, and transmit the virus to uninfected recipients warranting further studies and preparedness.

Text

SARS-CoV-2 has been detected in farmed and feral American mink in multiple countries with

evidence of extensive environmental contamination and human-to-mink and mink-to-human

transmission *(1-5)*. This has led to strict measures in mink farms and mink farming countries to

prevent the spread of the disease. In late 2021, a new SARS-CoV-2 variant (Omicron),

characterized by possibly milder symptoms and more efficient human-to-human transmission, was

detected, but its infectivity and spread in American mink in unknown *(6, 7)*.

We tested the response of the American mink to the Omicron variant by infecting three male mink

intranasally with 4×10^5 PFU of the virus (see Appendix for methods). Infected minks were

followed for seven days, sampled daily for saliva, and subjected to histopathological evaluation of

upper and lower respiratory tracts on the last day of follow-up.

All experimentally infected mink showed mild to moderate signs of illness including lethargy,

anorexia, diarrhea, nasal and lacrimal discharge, and sneezing. Consistent with earlier experiments

with other variants *(8, 9)*, saliva samples tested PCR positive 1-day post-infection (dpi) and

remained that way throughout the follow-up (Table and Appendix). Even though some of the

clinical signs may be due to other factors such as stress from the change of environment,

consistency of symptoms to studies with other variants combined with PCR results, demonstrate

that the Omicron variant also causes a symptomatic infection in mink.

To study if mink can transmit the virus, two minks were used as uninfected recipients in separate

cages located 10-20 cm away from the cages of the infected mink and followed for ten days. Both

recipients developed similar symptoms to the experimentally infected mink and were consistently

PCR positive from day three onwards (Table), indicating mink-to-mink transmission. Even though

there is currently no evidence of mink-to-human transmission of the Omicron variant, it seems

likely based on our results and the information from other variants.

 Table: PCR results from saliva of three experimentally infected mink and two uninfected recipient mink

Mink Id	1 dpi	2 dpi	3 dpi	4 dpi	5 dpi	6 dpi	7 dpi	8 dpi	9 dpi	10 dpi
Infected mink saliva										
451	+	+	+	+	+	+	+			
453	+	+	+	+	+	+	+			
455	+	+	+	+	+	+	+			
Recipient mink saliva										
452	(+)	-	+	+	+	+	+	+	+	(+)
454	-	-	(+)	+	+	+	+	+	+	+

dpi, days post infection of experimentally infected mink; +, signal detected with both primers of Luna SARS-CoV-2

RT-qPCR Multiplex Assay Kit; (+), signal detected only with one out of two primers; -, no signal with either of the

primers

Gross findings in the nasal cavity and lungs were subtle in both experimentally infected and

recipient mink consisting of hyperemia of respiratory mucosa with small amounts of viscous

exudate and non-collapsed, dark-red, and wet pulmonary lobes. All mink showed consistent

histopathological changes in the upper and lower respiratory tracts. Multifocal degeneration and

loss of respiratory epithelium with variable mucosal and submucosal neutrophilic infiltration was

observed in the nose. The lumen contained sloughed epithelial cells, mucinous material, and

degenerated neutrophils (Figure A, C). Viral nucleoprotein was widely distributed beyond intact

cells, within sloughed cells, and mucosal respiratory epithelium (Figure B, D). The olfactory

epithelium was inconsistent and mildly affected with only focal viral antigen detection. Unlike in

some experimental infections reported in rodents, clear pathology was observed in the lungs (*10*). In

two inoculated and both recipient mink, pulmonary lesions (Figure E, G) were associated with viral

antigen expression (Figure F, H) and characterized by multifocal to coalescing alveolar damage

with degeneration and/or necrosis of alveolar septa, infrequent hyalin membrane formation, and

variable proliferation of type II pneumocytes (Figure I, J). Alveolar spaces contained macrophages, sloughed cells, edema, and hemorrhage. Bronchiolar epithelial degeneration and hyperplasia were variably present (Figure K), and the lumen filled with few sloughed cells and neutrophils. Bronchi were lined by hyperplastic epithelium with increased numbers of goblet cells. Other consistent findings were vasculitis (Figure L), perivasculitis, perivascular, and peri-bronchial edema. One inoculated mink had markedly thickened alveolar septa by mononuclear cells, marked proliferation of type II pneumocytes, intra-alveolar macrophages, few syncytial cells, bronchi and bronchiolar epithelial cell hyperplasia, vasculitis, and perivasculitis. Viral antigen could not be detected in this mink. Strikingly, all evaluated mink lacked viral antigen in the epithelium of bronchi and bronchioles.

The Omicron variant is different from other variants due to its more efficient spread, primarily attributable to immune escape and likely milder symptoms in humans (*6, 7*). This makes it more difficult to prevent virus introduction into the mink farms through asymptomatic humans, creating a more significant risk for the formation of virus reservoirs among farmed or feral mink. This study shows that mink can be infected by Omicron and importantly, efficiently transmit the virus to other mink. Despite the reports of lower virulence of Omicron, mink develop clinical disease and nasal and pulmonary microscopic lesions closely resemble infection with previously reported variants in mink and humans. A better understanding of the clinical symptoms helps detect the virus among mink earlier. Further studies are needed to determine the risk of transmission to humans, emergence of mink-specific mutations, the pathogenesis of pulmonary involvement, and prepare for this easily transmitted variant among farmed and feral mink.

Figure. Histopathological changes and SARS-CoV-2 expression in the upper and lower respiratory tracts in

experimentally infected mink with Omicron variant at 7 dpi and recipient mink after 10 days of follow up. A)

Respiratory segment of the nose from an intranasally infected mink showing luminal accumulation of exudate (*) and

degeneration of mucosal epithelium (arrow, bar 500µm. B) Viral antigen is widely detected within nasal lumen and

respiratory epithelium (bar 500µm). C) Respiratory epithelium from a recipient mink depicting marked degeneration

and loss (arrow), intraluminal accumulation of sloughed cells and neutrophils (*), and D) intraepithelial viral expression

(C, D bar 50 µm). E-H) Lungs from intranasally infected (E, F) and recipient mink (G-H) showing alveolar damage

with intralesional presence of viral nucleoprotein (E-G bar 200µm, H bar 50 µm). I, J) Marked degeneration and

necrosis of alveolar septa and focal hyalin membrane (arrow, I), and prominent proliferation of type II pneumocytes (J)

in an intranasally infected mink (I, J bar 25µm). K, L) Recipient mink showing bronchiolar epithelial degeneration and

hyperplasia (K), and vasculitis (L) with complete destruction of blood vessel wall and mononuclear cell infiltration (K

bar 50µm, L bar 100µm). Haematoxylin and esosin (HE) stain and immunohistochemistry, haematoxylin counterstain.

Acknowledgements: We thank animal Jari Elemo and Mari Elemo and other animal care takers for

handling the animals and assessing their health and Esa Pohjalainen, Sanna Mäki, Tiina Sihvonen,

Johanna Rintamäki, Hanna Valtonen, Marika Skön, Larissa Laine, and Elina Väisänen for technical

assistance. We thank Kati Kuipers, Anne Kujanpää, Laura Vähälä, and the Finnish Centre for

Laboratory Animal Pathology (FCLAP) for expert technical help as well as Johanna Korpela and

Jussi Peura from Finnish Fur Breeders Association and Jan Segervall and Maarit Mohaibes from the

Kannus Research Farm Luova Ltd. for providing the animals. We also thank E3 Excellence in

Pandemic Response and Enterprise Solutions co-innovation project and all its parties.

This study was supported by the Academy of Finland (grant number 336490, 339510), VEO -

European Union's Horizon 2020 (grant number 874735), Business Finland E3 (4917/31/2021),

Finnish Institute for Health, and Welfare and the Jane and Aatos Erkko Foundation.

References

1.   Aguilo-Gisbert J, Padilla-Blanco M, Lizana V, Maiques E, Munoz-Baquero M, Chillida-Martinez
E, et al. First Description of SARS-CoV-2 Infection in Two Feral American Mink (Neovison vison) Caught in
the Wild. Animals (Basel). 2021 May 16;11(5).
2.   Oreshkova N, Molenaar RJ, Vreman S, Harders F, Oude Munnink BB, Hakze-van der Honing
RW, et al. SARS-CoV-2 infection in farmed minks, the Netherlands, April and May 2020. Euro Surveill. 2020
Jun;25(23).
3.   Rabalski L, Kosinski M, Mazur-Panasiuk N, Szewczyk B, Bienkowska-Szewczyk K, Kant R, et al.
Zoonotic spill-over of SARS-CoV-2: mink-adapted virus in humans. Clin Microbiol Infect. 2021 Dec 15.
4.   Rabalski L, Kosinski M, Smura T, Aaltonen K, Kant R, Sironen T, et al. Severe Acute
Respiratory Syndrome Coronavirus 2 in Farmed Mink (Neovison vison), Poland. Emerg Infect Dis. 2021
Sep;27(9):2333-9.
5.   Hammer AS, Quaade ML, Rasmussen TB, Fonager J, Rasmussen M, Mundbjerg K, et al. SARS-
CoV-2 Transmission between Mink (Neovison vison) and Humans, Denmark. Emerg Infect Dis. 2021
Feb;27(2):547-51.
6.   Viana R, Moyo S, Amoako DG, Tegally H, Scheepers C, Althaus CL, et al. Rapid epidemic
expansion of the SARS-CoV-2 Omicron variant in southern Africa. Nature. 2022 Jan 7.
7.   Wolter N, Jassat W, Walaza S, Welch R, Moultrie H, Groome M, et al. Early assessment of the
clinical severity of the SARS-CoV-2 omicron variant in South Africa: a data linkage study. Lancet. 2022 Jan
29;399(10323):437-46.
8.   D. Adney, J. Lovaglio, J. Schulz, C. Yinda, V. Avanzato, E. Haddock, et al. Severe acute
respiratory disease in American mink (Neovison vison) experimentally infected with SARS-CoV-2. bioRxiv
preprint 2022.
9.   Song Z, Bao L, Deng W, Liu J, Ren E, Lv Q, et al. Integrated histopathological, lipidomic, and
metabolomic profiles reveal mink is a useful animal model to mimic the pathogenicity of severe COVID-19
patients. Signal Transduction and Targeted Therapy. 2022;7(29).
10.   Halfmann PJ, Iida S, Iwatsuki-Horimoto K, Maemura T, Kiso M, Scheaffer SM, et al. SARS-CoV-
2 Omicron virus causes attenuated disease in mice and hamsters. Nature. 2022 Jan 21.

Address for Correspondence: Jenni Virtanen, Department of Veterinary Biosciences, Faculty of

Veterinary Medicine, P.O. Box 66, and Department of Virology, Faculty of Medicine, P.O. Box 21,

00014 University of Helsinki, Helsinki, Finland; tel: +358407768832, email:

jenni.me.virtanen@helsinki.fi

149