Detecting clonemates in multicellular clonal species based on

"Shared Heterozygosity (SH)" and technical replicates

Lei Yu[1*], John J. Stachowicz[2,3], Katie DuBois[2,3], Thorsten B. H. Reusch[1]

[1]Marine Evolutionary Ecology, GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel, Germany

[2]Center for Population Biology, University of California, Davis, CA, USA

[3]Department of Evolution and Ecology, University of California, Davis, CA, USA

[*]Corresponding author, lyu@geomar.de

Abstract

Clonal reproduction, the formation of new individuals that are genetically nearly identical to

the parent via mitosis in the absence of genetic recombination, is a very common reproductive

mode across plants, fungi, and animals. Current genetic marker based methods fail to detect

clonal structure when all collected samples belong to one single clone, which seems to be

more common than previously thought. Here we propose a new similarity index, "Shared

Heterozygosity (SH)" based on the number of genetic markers (typically SNPs, single-

nucleotide polymorphisms) that are identically heterozygous among two or more genomes

(i.e., N_{SH}). Ideally N_{SH} should be on the order of approximately $>=3{,}000$, which can be easily

achieved nowadays via Restriction-site Associated DNA (RAD) sequencing or whole-genome

resequencing. One should be careful when N_{SH} is small (roughly $<1{,}000$). We analyze two

large seagrass clones (*Posidonia australis*, *Zostera marina*) along with two *Z. marina* clones

of known age (17-yrs), and show that SH can potentially extend the detection of clonemates

to any pair of samples with the aid of technical replicates. Another potential application of SH

is to detect possible parent-descendant pairs under selfing, because the heterozygous loci in

the selfing-produced descendants represent a subset of those in the parent. Our proposed

workflow takes advantage of the availability of the larger number of genetic markers in the

genomic era, and fills a gap in detecting clonemates in the growing number of cases where

many or all samples at a location belong to one single clone.

Key words: shared heterozygosity, technical replicates, clonal reproduction, clonemate,

partial clonality

Introduction

Clonal reproduction refers to the formation of new individuals that are genetically nearly identical to the parent via mitosis in the absence of genetic recombination (De Meeûs et al., 2007). This reproductive mode is widespread in multicellular organisms, and has many different forms, such as vegetative multiplication (animals and plants), parthenogenesis (animals), apomixis (plants), and gynogeneis or pseudogamy (De Meeûs et al., 2007; Orive & Krueger-Hadfield, 2021). In multicellular clonal species, a module or ramet (Harper & White, 1974) is the basic modular unit with an ability to live independently, while a genet (synonymous with clone) is the product of a single zygote from gamete fusion (Noble et al., 1979), and comprises all ramets that emerge via mitosis (i.e., clonemates) during its lifetime. The majority of multicellular clonal species can conduct both clonal and sexual reproduction (Schön et al., 2009), which is called partial clonality (Stoeckel et al., 2021a; Stoeckel et al., 2021b). Partially clonal taxa thus have two levels of biological organization, an asexual population of ramets belonging to a single clone/genet, which is nested into the population of clones/genets, the "classical", sexually reproducing population (Reusch et al., 2021).

Ramets belonging to the same clone vs. different clones are mostly phenotypically indistinguishable and can also rarely be physically tracked once they obtain autonomy and break off the parental module. Hence, genetic markers such as microsatellites or AFLP have been widely used for detecting clonemates (Arnaud-Haond et al., 2007). A ramet can be genotyped with multiple independent markers, and the genotypes at all these loci collectively form a multilocus genotype (MLG). In case of co-dominant markers such as microsatellites, identical or nearly identical MLGs are then interpreted as being members of the same clone or genet. However, identical or nearly identical MLGs are not definitive evidence for clonal

reproduction (Halkett et al., 2005), if for example there is low (or no) allelic diversity at

particular loci. In order to calculate the probability of finding identical MLGs resulting from

distinct zygotes, the population allelic frequencies are required, and the population is assumed

to be under Hardy-Weinberg equilibrium (Parks & Werth, 1993). Evidently, estimates of such

population-level frequencies are inaccurate in populations dominated by a few clones

(Arnaud-Haond et al., 2007), and meaningless if a single site is dominated by one clone

(Smith et al., 1992; Reusch et al., 1999; Arnaud-Haond et al., 2012; Japaud et al., 2015;

Edgeloe et al., 2022), thereby rendering distinction of clonemates from non-clonemates

difficult.

Another difficulty arises because members of the same clone may have slightly different

MLGs (Douhovnikoff & Dodd, 2003; Yu et al., 2020; Edgeloe et al., 2022) owing to somatic

genetic polymorphisms or technical errors. To overcome this issue and also to work with

dominant markers that do not allow for the application of MLGs, an alternative suite of

approaches is the analysis of pairwise genetic distances/similarities. The genomes of the

ramets from the same clone are identical by descent (IBD), except at those sites that have

experienced somatic mutations, as the genomes are inherited from the same asexual ancestor

via mitosis without recombination. Thus, ramets from the same clone will show much lower

levels of genetic distances or much higher levels of genetic similarities than those from

different clones. This will lead to a bimodal pattern in the frequency distribution of pairwise

genetic distances/similarities. Then, a threshold can be defined to separate the intra-clone vs.

inter-clone ramet pairs (Douhovnikoff & Dodd, 2003; Bailleul et al., 2016; Meirmans, 2020).

Such a threshold can be defined arbitrarily by the user, or by simulating a pseudo-observed

generation of pure sexual reproduction from the genotyped population (Bailleul et al., 2016).

Obviously, one problem is that the collected samples have to belong to more than one

clones/genets. Otherwise, it would be impossible to identify the intra-clone sample pairs.

Here, we propose a novel genetic similarity index, Shared Heterozygosity (SH), which

corresponds to the number of heterozygous sites shared by all individuals (ramets) divided by

the number of heterozygous sites observed in the individual with the highest count.

SH is an extension to Jaccard similarity index, with some modifications. SH involves a

selection of marker loci, taking into account only the ones where at least one sample is

heterozygous. Homozygous sites are usually not informative, as even distantly related

individuals are identically homozygous at most of the genomic sites (The 1000 Genomes

Project Consortium, 2015; The 1001 Genomes Consortium, 2016). On the contrary,

heterozygous sites are more informative for detecting clonemates, because they show the

potential for segregation in sexual reproduction, while remaining identically heterozygous in

clonal reproduction.

The genetic differences between clonemates are determined by both technical errors and

somatic genetic polymorphisms. By analyzing some of the largest detected clones ever (Yu et

al., 2020; Edgeloe et al., 2022), we provided a conservative estimate as to how large the

within-clone divergence in terms of non-shared heterozygosity may get. Even for these very

large (and possibly old) clones, the level of non-shared heterozygosity caused by somatic

genetic polymorphisms was low. Therefore, after having quantified the influence of technical

errors with the aid of technical replicates (i.e., multiple pieces of tissues from the same sample

are used for all the processes from DNA extraction to genotype calling), it may be possible to

identify the intra-clone sample pairs without inter-clone information, which will extend the

clonemate detection to any pair of samples.

Our workflow is most suitable for large number of SNPs obtained from whole-genome

resequencing, restriction-site associated DNA (RAD) sequencing, or SNP arrays. The

application of SH requires a sufficient number of genetic markers where the two target

samples are identically heterozygous (i.e., N_{SH}, ideally on the order of >= 3,000). Since the

number of markers used does not always reflect the level of N_{SH}, one should pay attention to

N_{SH} and be careful when the level of N_{SH} is low (roughly <1,000). The goal of this study is to

develop the workflow of detecting clonemates based on shared heterozygosity and technical

replicates (Fig. 1). We apply our workflow to two of the largest clones in marine systems

detected so far, seagrass clones in Finland (*Zostera marina*, covering >300 m x 200 m) and

Australia (*Posidonia australis*, spanning >180 km), respectively (Yu et al., 2020; Edgeloe et

al., 2022), which likely represent the far end of accumulating somatically generated variation.

We also apply the workflow to two 17-year-old *Z. marina* clones cultured in outdoor flow-

through mesocosms, representing the near end of accumulating somatically generated

variation.

Materials and Methods

Shared Heterozygosity (SH) indexes

For n samples in multicellular clonal species (X_1, …, X_n), $N_{SH}(X_1, …, X_n)$ denotes the

number of the genetic markers where n samples are identically heterozygous; $N_{Het}(X_n)$

denotes the number of the heterozygous loci in X_n.

$SH(X_n) = N_{SH}(X_1, …, X_n)/N_{Het}(X_n)$

$SH(X_1, …, X_n) = min(SH(X_1), …, SH(X_n))$.

Detecting clonemates based on a published dataset on seagrass _Posidonia australis_

Edgeloe et al. (2022) reported a polyploid _P. australis_ clone spanning at least 180 km. The

SNP dataset (all.vcf) containing 18,021 SNPs for 133 individuals (including 4 technical

replicates) from 10 populations was used here. According to Edgeloe et al. (2022), population

GU was diploid, while the other 9 populations were polyploid and belonged to the same

clone. SH indexes were calculated for each pair of samples based on the SNP dataset, and

clonemate detection was conducted following the workflow illustrated by Fig. 1.

Distribution of SH indexes within two _Z. marina_ clones that are 17 years old

Two _Z. marina_ clones were selected from the eight clones cultured in adjacent 300-l outdoor

flow-through mesocosms at Bodega Marine Laboratory (BML) under ambient light and

temperature conditions (Hughes et al., 2009). Each mesocosm was initiated with a single

terminal shoot collected from one of eight different locations in Bodega Harbor, California, in

July 2004. The genetic distinctiveness of each of the original shoots was verified before

planting. Six shoots (= ramets) were collected from each clone for genomic analysis in 2021,

after the clones had grown in mesocosm for 17 years. The two clones were referred to as

"GREEN" and "RED", respectively.

Genomic DNA was extracted using NucleoSpin plant II kit from Macherey-Nagel following

the instructions. The plant tissue (160-200 mg fresh weight) was ground with liquid nitrogen.

Library construction and Illumina sequencing (PE150, paired end 150 bp) were conducted by

BGI (Beijing Genomics Institute, Hong Kong). The quality of the raw Illumina reads was

assessed by FastQC V0.11.5 (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/).

BBDuk (last modified November 7, 2019, https://jgi.doe.gov/data-and-tools/bbtools/bb-tools-

user-guide/bbduk-guide/) was used to remove adapter contamination and conduct a basic

quality filtering: (1) reads with more than one N were discarded (maxns=1); (2) reads shorter

than 50 bp after trimming were discarded (minlength=50); (3) reads with average quality

below 10 after trimming were discarded (maq=10). FastQC V0.11.5 was used for second

round of quality check for the filtered reads.

The quality-filtered Illumina reads were mapped against the chromosome-level *Z. marina*

reference genome V3.1 (Ma et al., 2021) using BWA MEM V0.7.17 (Li & Durbin, 2009).

The alignments were converted to BAM format and then sorted using Samtools V1.11 (Li et

al., 2009). The MarkDuplicates module in GATK V4.1.1.0 (Van der Auwera & O'Connor,

2020) was used to identify and tag duplicate reads in the BAM files. Sequencing depth and

percentage of covered genome region for each ramet were calculated using Samtools V1.11

and Bedtools V1.11 (Quinlan & Hall, 2010), respectively. HaplotypeCaller (GATK V4.1.1.0)

was used to generate GVCF format file for each ramet, and all the GVCF files were combined

by CombineGVCFs (GATK V4.1.1.0). GenotypeGVCFs (GATK V4.1.1.0) was used to call

genetic variants.

BCFtools V1.11 (Li, 2011) was used to remove SNPs within 20 base pairs of an indel or other

variant type. Then we extracted only bi-allelic SNPs located in the six main chromosomes

(2,969,820 SNPs). INFO annotations were extracted using VariantsToTable (GATK

V4.1.1.0). SNPs meeting one or more than one of the following criteria were marked by

VariantFiltration (GATK V4.1.1.0): MQ < 40.0; FS > 60.0; QD < 13.0; MQRandSum > 2.5

or MQRandSum < -2.5; ReadPosRandSum < -2.5; ReadPosRandSum > 2.5; SOR > 3.0; DP >

1192.45 (2 * average DP). Those SNPs were excluded by SelectVariants (GATK V4.1.1.0). A

total of 1,105,628 SNPs were retained. VCFtools V0.1.15 (Danecek et al., 2011) was used to

convert individual genotypes to missing data when GQ < 30 or DP < 20. Individual

homozygous reference calls with one or more than one reads supporting the variant allele, and

individual homozygous variant calls with one or more than one reads supporting the reference

allele, were also converted to missing data using a custom Python script

(https://github.com/leiyu37/Detecting-clonemates.git). Finally, we kept only genomic sites

showing polymorphism (604,066 SNPs).

SH indexes were calculated by a custom Phython script

(https://github.com/leiyu37/Detecting-clonemates.git) for each pair of samples. We only

focused on the nucleotide positions in the reference genome where both samples had available

genotype calls (not missing data).

Detecting clonemates based on a published dataset on seagrass *Zostera marina*

Yu et al. (2020) conducted whole-genome resequencing of 24 ramets of a seagrass *Z. marina*

clone located in Finland. The clone was estimated to be between 750 and 1,500 years old.

Here the clone was referred to as "the Finnish clone". All the 24 ramets were sequenced by

Illumina Hiseq, and three of them were also sequenced independently by Illumina Novaseq.

The Hiseq and Novaseq data for the three ramets provided three pairs of technical replicates.

The Novaseq data reached a coverage of ~1000x. A subset was selected from the Novaseq

data (~80x) to achieve a similar coverage to the Hiseq data. Then we re-analyzed the Hiseq

and Novaseq (~80x) data together using the same method above for the two 17-year-old

clones. All the SNP calling and filtering processes were exactly same with those for the 17-

year-old clones, except the threshold for DP during hard filtering (DP > 2945.57). SH indexes

were calculated for each pair of samples.

To understand the influence of $N_{SH}(X_1, X_2)$ on $SH(X_1, X_2)$, we randomly selected 1/10, 1/100

and 1/1000 from the 38,831 SNPs for the 24 ramets available in Yu et al. (2020), based on

which $N_{SH}(X_1, X_2)$ and $SH(X_1, X_2)$ were calculated again for each pair of samples. Each

selection was repeated 100 times, and $N_{SH}(X_1, X_2)$ and $SH(X_1, X_2)$ from the 100 repeats were

pooled together to analyze.

Results

We applied our workflow to two of the largest clones in marine systems detected so far,

seagrass clones in Finland (*Zostera marina*, covering >300 m x 200 m) and Australia

(*Posidonia australis*, spanning >180 km), respectively (Yu et al., 2020; Edgeloe et al., 2022).

In the larger clone of *P. australis* in Shark Bay (Western Australia), SNPs were obtained by

RAD (Edgeloe et al., 2022). Based on the SNP dataset encompassing 18,021 SNPs for 133

individuals (including 4 technical replicates) from 10 populations, the pairwise N_{SH} ranged

from 1,047 to 15,428. The minimum pairwise SH between technical replicates was 0.9809.

The pairwise SH between clonemates were smaller than 0.9809, due to somatic genetic

variation. They were distributed within a narrow range bordered by the technical replicates

(Fig. 2a). Including technical replicates made it possible to identify intra-clone sample pairs

without inter-clone information (Fig. 2a). Our analyses based on SH indexes and technical

replicates confirmed that all populations except population GU belonged to the same clone

(Fig. 2b).

In the relatively smaller clone of *Z. marina* in Finland, whole-genome resequencing was

conducted for 24 ramets (Yu et al., 2020). After re-analyzing the data (Hiseq data for 24

ramets, and Novaseq data for 3 of the ramets), we obtained 41,885 SNPs. The Hiseq and

Novaseq data for the three ramets provided three pairs of technical replicates. Based on the

dataset, the pairwise N_{SH} ranged from 28,619 to 32,210 (Fig. 3). The minimum pairwise SH

between technical replicates was 0.9998 (Fig. 3a). The pairwise SH between clonemates were

distributed within a narrow range from 0.9548 to 0.9993, bordered by the technical replicates

(Fig. 3b). Our analyses based on SH indexes and technical replicates confirmed that all 24

ramets belonged to the same clone. By calculating SH indexes based on the the subsets of the

SNP dataset, we found that smaller N_{SH} led to larger variation of pairwise SH (Fig. 4). The

consistently high SH when N_{SH} is at least on the order of >=3,000 suggests this as a

reasonable minimum target to apply this method. When N_{SH} <1,000, variability in SH

increased considerably, which would reduce confidence in distinguishing clonemates from

non-clonemates.

We also calibrated our approach using 2 relatively young *Z. marina* clones of known age (17-

year-old clones). Unfortunately, technical replicates were also not available. The whole-

genome resequencing data of the 12 cultured *Z. marina* ramets covered >93% of the reference

genome, and reached >66x coverage (Table 1). The pairwise N_{SH} ranged from 108,585 to

288,606 (Fig. 5a). Although technical replicates were not available, the pairwise SH between

technical replicates should be distributed within the range from 0.9980 to 1 (Fig. 5b). Within

the clone "GREEN", the minimum $SH(X_1, X_2)$ was 0.9975 (Fig. 5a). As for the clone "RED",

we detected one "contaminant" ramet (i.e., R10), obviously not belonging to that clone. The

five ramet pairs composed of R10 and one of the other five "RED" ramets showed SH(R10,

X_2) ranging from 0.4103 to 0.4114 (Fig. 5a), indicating that R10 does not belong to this clone.

Apart from R10, the minimum SH(X_1, X_2) for the ramet pairs among the other five "RED"

ramets was 0.9974 (Fig. 5a). The inter-clone SH(X_1, X_2) ranged from 0.4790 to 0.4803. Since

the pairwise SH between clonemates were distributed within a narrow range from 0.9974 to

0.9980 (Fig. 5b), they could be easily identified.

Discussion

Both sexual and clonal reproduction occur in almost all clonal taxa, but the length and extent

of the clonal phase varies widely within and among species (for seagrasses, e.g. Reusch &

Boström, 2011). Examples accumulate that some populations can propagate clonally to such

an extent that the whole population is dominated by one or a few clones, such as fungi (Smith

et al., 1992), seagrass species *Zostera marina* (Reusch et al., 1999; Yu et al., 2020), *Posidonia*

oceanica (Arnaud-Haond et al., 2012) and *P. australis* (Edgeloe et al., 2022), and corals

(Japaud et al., 2015). The situation of having a collection of samples belonging to one single

clone (Yu et al., 2020) is thus more widespread than previously thought. The currently

available methods cannot work in this extreme case (Arnaud-Haond et al., 2007). We have

proposed a workflow based on the concept of "Shared Heterozygosity (SH)", meaning that

two or more than two genomes are identically heterozygous at some genetic markers. The

pairwise SH between clonemates are determined by both technical errors and somatic genetic

variation. However, even for the two very old seagrass clones, the level of pairwise

differences caused by somatic genetic variation is relatively low (Fig. 2&3). We have shown

that intra-clone sample pairs can be easily identified with the aid of the technical replicates

even without inter-clone information (Fig. 3). Our workflow can be applied to any pair of

ramets when technical replicates are available, even if there is only one genet occurring

within a "population"-level sampling. To the best of our knowledge, this is the only method

that can work in cases where all collected ramets belong to one single clone.

When applying the workflow to the two 17-year-old *Z. marina* clones, we found one

"contaminant" ramet (i.e., R10) that clearly did not belong to the clone "RED". One possible

reason is that R10 was produced by sexual event between members of the clone "RED" (i.e.,

selfing). In the first 15 years of growth, this would be the only way for contamination to

occur, as the clones were cultured in isolated containers. Under selfing, each of the

heterozygous loci in the parent can either remain identically heterozygous or change to a

homozygous state in R10. As a consequence, the heterozygous loci in R10 should be a subset

of those in the parent, and thus SH(R10) for the five ramet pairs composed of R10 and one of

the other five "RED" ramets should be close to 1. However, that is not the case, as SH(R10)

for the five ramet pairs ranges from 0.4320 to 0.4331. Hence, R10 is not likely to be derived

from selfing. In late 2019 or early 2020, these clones were transferred and planted in separate

containers within in a single larger tank (~4 m diameter). All clones were in the same tank for

at most one year. It is possible that either (a) outcrossing occurred and seeds set and

germinated or (b) shoots dislodged and re-attached or were able to grow between pots.

SH can potentially be used to detect possible parent-descendant pairs under selfing, because

this reproductive mode leads to loss of heterozygosity. Under selfing, a subset of the

heterozygous loci in the parent will change to a homozygous state in the descendants. If not

considering the ramet-specific heterozygous loci derived from mutations and genotyping

errors, the heterozygous loci in the descendant should be a subset of those in the parent,

which can potentially be used to detect possible parent-descendant pairs under selfing.

The application of SH requires a sufficient number of genetic markers where the two target

samples are identically heterozygous (i.e., N_{SH}). We were able to correctly detect clonemates

based on the datasets with minimum N_{SH} of 1,047. With the development of sequencing

technology, it is now easy to obtain thousands of genetic markers using whole-genome

resequencing, restriction-site associated DNA (RAD) sequencing, or SNP arrays. However,

the level of N_{SH} can still be low based on genome-wide maker sets, when the target species

shows very low levels of genome heterozygosity. One should pay attention to N_{SH} and use

caution in the interpretation of results when N_{SH} is low. Our workflow takes advantage of the

availability of the larger number of genetic markers in the genomic era. It can potentially

extend the detection of clonemates to any pair of samples against the baseline of technical

replicates and fills a gap in detecting clonemates in extreme cases where all sampled

genotypes at a site belong to one single clone.

Acknowledgements

This study was supported by a 4-year PhD scholarship from the China Scholarship Council

(CSC) to L.Y. (No. 201704910807), and by a Human Frontiers of Science (HFSP) grant to

T.B.H.R (RGP0042_2020). We thank D. Gill for lab assistance. We thank Dr. Y. Li for the

comments on the early version of the manuscript.

References

318   Arnaud-Haond, S., Duarte, C. M., Diaz-Almela, E., Marbà, N., Sintes, T. & Serrão, E. A.
319   (2012). Implications of extreme life span in clonal organisms: millenary clones in

meadows of the threatened seagrass *Posidonia oceanica. PLoS ONE, 7*(2), e30454. doi:10.1371/journal.pone.0030454

Arnaud-Haond, S., Duarte, C. M., Alberto, F. & Serrao, E. A. (2007). Standardizing methods to address clonality in population studies. *Mol. Ecol., 16*(24), 5115-5139. doi:10.1111/j.1365-294X.2007.03535.x

Bailleul, D., Stoeckel, S. & Arnaud-Haond, S. (2016). RClone: a package to identify MultiLocus Clonal Lineages and handle clonal data sets in r. *Methods Ecol. Evol., 7*(8), 966-970. doi:10.1111/2041-210X.12550

Buss, L. W. (1983). Evolution, development, and the units of selection. *Proc. Natl. Acad. Sci. U.S.A, 80*(5), 1387-1391. doi:10.1073/pnas.80.5.1387

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., . . . Genomes Project Analysis Group. (2011). The variant call format and VCFtools. *Bioinformatics, 27*(15), 2156-2158. doi:10.1093/bioinformatics/btr330

De Meeûs, T., Prugnolle, F. & Agnew, P. (2007). Asexual reproduction: genetics and evolutionary aspects. *Cell. Mol. Life Sci., 64*(11), 1355-1372. doi:10.1007/s00018-007-6515-2

Douhovnikoff, V. & Dodd, R. S. (2003). Intra-clonal variation and a similarity threshold for identification of clones: application to Salix exigua using AFLP molecular markers. *Theor. Appl. Genet., 106*(7), 1307-1315. doi:10.1007/s00122-003-1200-9

Edgeloe, J. M., Severn-Ellis, A. A., Bayer, P. E., Mehravi, S., Breed, M. F., Krauss, S. L., . . . Sinclair, E. A. (2022). Extensive polyploid clonality was a successful strategy for seagrass to expand into a newly submerged environment. *Proc. Royal Soc. B, 289*(1976), 20220538. doi:10.1098/rspb.2022.0538

Halkett, F., Simon, J. & Balloux, F. (2005). Tackling the population genetics of clonal and partially clonal organisms. *Trends Ecol. Evol. , 20*(4), 194-201. doi:10.1016/j.tree.2005.01.001

Harper, J. L. & White, J. (1974). The demography of plants. *Annu. Rev. Ecol. Syst., 5*(1), 419-463. doi:10.1146/annurev.es.05.110174.002223

Honnay, O. & Bossuyt, B. (2005). Prolonged clonal growth: escape route or route to extinction? *Oikos, 108*(2), 427-432. doi:10.1111/j.0030-1299.2005.13569.x

Hughes, R. A., Stachowicz, J. J. & Williams, S. L. (2009). Morphological and physiological variation among seagrass (*Zostera marina*) genotypes. *Oecologia, 159*(4), 725-733. doi:10.1007/s00442-008-1251-3

Japaud, A., Bouchon, C., Manceau, J. & Fauvelot, C. (2015). High clonality in *Acropora palmata* and *Acropora cervicornis* populations of Guadeloupe, French Lesser Antilles. *Mar. Freshw. Res., 66*(9), 847-851. doi:10.1071/MF14181

Li, H. (2011). A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. *Bioinformatics, 27*(21), 2987-2993. doi:10.1093/bioinformatics/btr509

Li, H. & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics, 25*(14), 1754-1760. doi:10.1093/bioinformatics/btp324

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., . . . Genome Project Data Processing, S. (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics, 25*(16), 2078-2079. doi:10.1093/bioinformatics/btp352

Ma, X., Olsen, J. L., Reusch, T. B. H., Procaccini, G., Kudrna, D., Williams, M., . . . Van de Peer, Y. (2021). Improved chromosome-level genome assembly and annotation of the seagrass, *Zostera marina* (eelgrass). *F1000research, 10.* doi:10.12688/f1000research.38156.1

368   Meirmans, P. G. (2020). genodive version 3.0: Easy-to-use software for the analysis of
369   genetic data of diploids and polyploids. *Mol. Ecol. Resour., 20*(4), 1126-1131.
370   doi:10.1111/1755-0998.13145
371   Noble, J. C., Bell, A. D. & Harper, J. L. (1979). The population biology of plants with clonal
372   growth: I. The morphology and structural demography of *Carex arenaria*. *J. Ecol., 67*,
373   983-1008. doi:10.2307/2259224
374   Orive, M. E. & Krueger-Hadfield, S. A. (2021). Sex and Asex: A clonal lexicon. *J. Hered.,*
375   *112*(1), 1-8. doi:10.1093/jhered/esaa058
376   Parks, J. C. & Werth, C. R. (1993). A study of spatial features of clones in a population of
377   bracken fern, *Pteridium aquilinum* (Dennstaedtiaceae). *Am. J. Bot., 80*(5), 537-544.
378   doi:10.1002/j.1537-2197.1993.tb13837.x
379   Quinlan, A. R. & Hall, I. M. (2010). BEDTools: a flexible suite of utilities for comparing
380   genomic features. *Bioinformatics, 26*(6), 841-842. doi:10.1093/bioinformatics/btq033
381   Reusch, T. B. H., Boström, C., Stam, W. T. & Olsen, J. L. (1999). An ancient eelgrass clone
382   in the Baltic. *Mar. Ecol. Prog. Ser., 183*, 301-304. doi:10.3354/meps183301
383   Reusch, T. B. H. & Boström, C. (2011). Widespread genetic mosaicism in the marine
384   angiosperm *Zostera marina* is correlated with clonal reproduction. *Evol. Ecol., 25*(4),
385   899-913. doi:10.1007/s10682-010-9436-8
386   Reusch, T. B. H., Baums, I. B. & Werner, B. (2021). Evolution via somatic genetic variation
387   in modular species. *Trends Ecol. Evol., 36*(12), 1083-1092.
388   doi:10.1016/j.tree.2021.08.011
389   Schön, I., Martens, K. & van Dijk, P. (2009). *Lost sex: The Evolutionary Biology of*
390   *Parthenogenesis*. Dordrecht: Springer Dordrecht.
391   Smith, M. L., Bruhn, J. N. & Anderson, J. B. (1992). The fungus *Armillaria bulbosa* is among
392   the largest and oldest living organisms. *Nature, 356*(6368), 428-431.
393   doi:10.1038/356428a0
394   Stoeckel, S., Arnaud-Haond, S. & Krueger-Hadfield, S. A. (2021a). The combined effect of
395   haplodiplonty and partial clonality on genotypic and genetic diversity in a finite
396   mutating population. *J. Hered., 112*(1), 78-91. doi:10.1093/jhered/esaa062
397   Stoeckel, S., Porro, B. & Arnaud-Haond, S. (2021b). The discernible and hidden effects of
398   clonality on the genotypic and genetic states of populations: improving our estimation
399   of clonal rates. *Mol. Ecol. Resour., 21*(4), 1068-1084. doi:10.1111/1755-0998.13316
400   The 1000 Genomes Project Consortium. (2015). A global reference for human genetic
401   variation. *Nature, 526*(7571), 68-74. doi:10.1038/nature15393
402   The 1001 Genomes Consortium. (2016). 1,135 genomes reveal the global pattern of
403   polymorphism in *Arabidopsis thaliana*. *Cell, 166*(2), 481-491.
404   doi:10.1016/j.cell.2016.05.063
405   Van der Auwera, G. A. & O'Connor, B. D. (2020). *Genomics in the cloud: using Docker,*
406   *GATK, and WDL in Terra*: O'Reilly Media.
407   Yu, L., Boström, C., Franzenburg, S., Bayer, T., Dagan, T. & Reusch, T. B. H. (2020).
408   Somatic genetic drift and multilevel selection in a clonal seagrass. *Nat. Ecol. Evol.,*
409   *4*(7), 952-962. doi:10.1038/s41559-020-1196-4
410

Data Accessibility

DNA sequence data are available in the NCBI short read archive, BioProject no.

PRJNA806459, SRA accession no. SRR18000159- SRR18000170. Custom-made computer

code is available at Github https://github.com/leiyu37/Detecting-clonemates.

Authors' contributions

L.Y. and T.B.H.R. conceived and designed the study, J.J.S and K.D. cultured and provided

the samples of the two 17-year-old seagrass *Zostera marina* clones, L.Y. extracted DNA and

analyzed the sequencing data, L.Y. performed the bioinformatic analyses, L.Y. and T.B.H.R.

wrote the paper, with input from all authors.

Samping permit

The original sampling of *Zostera marina* clones was conducted under the permit no. SC-4874.

Competing interests

The authors declare that they have no competing interests.

428 **Table 1: Coverage of the whole-genome resequencing data for the two cultured *Zostera***
429 ***marina* clones.**

Sample	Average depth for the covered region	Genome region covered (%)
G01	68.64	93.33
G02	80.42	93.66
G06	79.05	93.72
G07	69.94	93.36
G10	77.49	93.52
G16	73.31	93.26
R02	81.55	93.86
R03	85.13	94.06
R05	68.14	93.57
R08	66.44	93.51
R09	71.73	93.59
R10	67.97	93.69

1. **DNA extraction** 2. **Genotyping**	Generating technical replicates (i.e., multiple pieces of tissues of the same sample are used independently for all the processes from DNA extraction to genotyping).
3. Calculate Pairwise SH indexes	For each pair of samples, focus on only the loci meeting the following criteria: 1. Both samples have available genotype calls (i.e., not missing data). 2. At least one sample is heterozygous. Calculate N_{SH}, $SH(X_1)$, $SH(X_2)$, and $SH(X_1, X_2)$.
4. Histogram of pairwise SH

 |
| **5. Point plot based on N_{SH} and $SH(X_1, X_2)$ of all sample pairs** |

 |

Figure 1: Workflow for detecting clonemates in multicellular clonal species based on SH indexes and technical replicates.

Figure 2: Detecting clonemates using SH indexes and technical replicates based on a published dataset on

seagrass *Posidonia australis*. a, Histogram of pairwise SH. The dataset contains four pairs of technical

replicates. The orange vertical line indicates the minimum pairwise SH between technical replicates. The

pairwise SH between clonemates are expected to be smaller than the minimum pairwise SH between technical

replicates, due to somatic genetic variation. They form a distribution to the left of the orange vertical line. **b,**

Point plot based on all sample pairs. Each data point represents one pair of samples. The sample pairs are

categorized into four different types: 1, TR: a pair of technical replicates for the same sample; 2, non-GU:

neither samples are from population GU; 3, GU: both samples are from population GU; 4, GU_non-GU: one

sample is from population GU, and the other sample is from the other 9 populations.

438
439
448
449

Figure 3: Detecting clonemates using SH indexes and technical replicates based on a published dataset on seagrass *Zostera marina*. a, Histogram of pairwise SH. The dataset contains 24 ramets of the same clone. Three of them were sequenced independently on both Illumina Hiseq and Novaseq platform, providing three pairs of technical replicates. The orange vertical line indicates the minimum pairwise SH between technical replicates. The pairwise SH between clonemates are expected to be smaller than the minimum pairwise SH between technical replicates, due to somatic genetic variation. They form a distribution to the left of the orange vertical line. **b, Point plot based on all sample pairs.** Each data point represents one pair of samples. The sample pairs are categorized into two different types: 1, TR: a pair of technical replicates for the same sample; 2, Clone_FI: the samples pairs that do not belong to TR. The blue line indicates the minimum pairwise SH within Clone_FI.

Figure 4: Distribution of pairwise SH indexes within the Finnish *Zostera marina* clone based on subsets (1/10, 1/100, and 1/1000) of SNPs. The variation of $SH(X_1, X_2)$ increases when N_{SH} decreases.

Figure 5: Distribution of pairwise SH indexes within the two 17-year-old *Zostera marina* clones. a, SH indexes for all sample pairs. Each data point represents one sample pair, which is either from the same clone ("intra-GREEN" or "intra-RED") or from different clones ("inter-clone"). One ramet from clone "RED" (i.e., R10) obviously belongs to another clone, leading to much lower levels of pairwise SH with ramets of clone "RED". b, SH indexes for sample pairs of the same clone. The orange and the blue dotted lines indicate the minimum and maximum values for $SH(X_1, X_2)$, respectively.