1　Identification of Mebendazole as an ethylene signaling activator reveals a role of ethylene
2　signaling in the regulation of lateral root angles
3

4　Wenrong He1, Hai An Truong 1, Ling Zhang 1, Min Cao 1, Neal Arakawa 3, Yao Xiao 4,
5　Kaizhen Zhong1, Yingnan Hou 5, 6, Wolfgang Busch1,2,7

1. Plant Molecular and Cellular Biology Laboratory, Salk Institute for Biological Studies, La Jolla, California, USA
2. Integrative Biology Laboratory, Salk Institute for Biological Studies, La Jolla, California, USA
3. Environmental and Complex Analysis Laboratory (ECAL), Department of Chemistry and Biochemistry, University of California, San Diego, La Jolla, USA
4. Department of Integrative Structural and Computational Biology, The Scripps Research Institute, La Jolla, California, USA
5. Department of Microbiology and Plant Pathology, Center for Plant Cell Biology, University of California, Riverside, California, USA
6. School of Agriculture and Biology, Shanghai Jiao Tong University, Shanghai, China
7. Lead contact

Correspondence: wbusch@salk.edu

In brief: He *et al.* identify a small molecule that regulates lateral root angle. They show that the compound increases lateral root angle by inhibiting CTR1 kinase activity, which in turn activates ethylene signaling. Therefore, they uncover that the ethylene pathway is involved in lateral root angle regulation.

Highlights:
- MBZ increases lateral root angle
- MBZ regulates lateral root angle by activating ethylene signaling
- MBZ inhibits CTR1 kinase activity
- The ethylene pathway is involved in lateral root angle regulation

SUMMARY

The lateral root angle or gravitropic set-point angle (GSA) is an important trait for root system architecture (RSA) that determines the radial expansion of the root system. The GSA therefore plays a crucial role for the ability of plants to access nutrients and water in the soil. Despite its importance, only few regulatory pathways and mechanisms that determine GSA are known, and these mostly relate to auxin and cytokinin pathways. Here, we report the identification of a small molecule, Mebendazole (MBZ) that modulates GSA in *Arabidopsis thaliana* roots and acts via the activation of ethylene signaling. We uncover that MBZ directly acts on the serine/threonine protein kinase CTR1, which is a negative regulator of ethylene signaling. Our study not only reveals that the ethylene signaling pathway is essential for GSA regulation, but it also identifies a small molecular modulator of RSA and the first small molecule that acts downstream of ethylene receptors and that directly activates ethylene signaling.

KEYWORDS

Root System Architecture, Lateral Root Angle, Root Growth, Ethylene Signaling, Chemical genetics, Mebendazole

INTRODUCTION

The root systems of plants are vital for plant survival and productivity. They anchor plants and they absorb water and nutrients from the soil. The distribution of roots in the soil, root system architecture (RSA), is of enormous relevance. For instance, deeper rooting is associated with higher carbon permanence (Poirier et al., 2018), a critical trait for soil carbon sequestration and the ability of plants to survive terminal droughts (Uga et al., 2013). Besides, to enhance the capacity for root systems to efficiently uptake nitrogen and water, deeper roots are required, yet for improving the uptake of phosphate and other nutrients, shallower soil layers have to be accessed by roots (Lynch, 2019). A key determinant of RSA is the lateral root angle or gravitropic set-point angle (GSA). Previous studies have shown that auxin plays a crucial role in regulating GSA (Rosquete et al., 2013a; Roychoudhry et al., 2013; Xiao and Zhang, 2020). Application of exogenous IAA or synthetic auxin leads to lateral root growth toward the vector of gravity thereby

decreasing GSA (Rosquete et al., 2013a; Roychoudhry et al., 2013). Another phytohormone, cytokinin, was also found to be involved in the regulation of GSA. Application of cytokinin showed an anti-gravitropic effect and thereby increased GSA (Waidmann et al., 2019b). Finally, members of the IGT gene family, including DRO1/LAZY4 have been shown to play a major role in lateral root angles via auxin associated mechanisms (Ge and Chen, 2016; Guseman et al., 2017; Waite and Dardick, 2021; Yoshihara and Spalding, 2017).

Ethylene is an ancient plant hormone. The ethylene signaling pathway is highly conserved and was present already in the aquatic progenitors of land plants (Ju et al., 2015). Ethylene is produced from the conversion of S-adenosyl-L-methionine (SAM) to 1-aminocy-clopropane-1-carboxylic acid (ACC) and then to ethylene, which is catalyzed by enzymes including ACC synthase (ACS) and ACC oxidase (ACO) (Pattyn et al., 2021; S F Yang and Hoffman, 1984). When ethylene binds to its receptors (ETRs) (Chang et al., 1993; Hua et al., 1998; Sakai et al., 1998), it inactivates a serine/threonine protein kinase, CONSTITUTIVE TRIPLE RESPONSE1(CTR1) (Kieber et al., 1993). In the absence of ethylene, CTR1 blocks ethylene downstream responses by phosphorylating the C-terminal end of ETHYLENE-INSENSITIVE2 (EIN2), an ER–associated membrane protein that works as a positive regulator of ethylene signaling (Alonso et al., 1999; Ju et al., 2012; Wen et al., 2012). This leads to the degradation of EIN2-CEND by the Ub/26S proteasome (Ju et al., 2012; Qiao et al., 2012a; Wen et al., 2012). When CTR1 is inactivated by ethylene binding to ETRs, EIN2-CEND is stabilized and moves into the nucleus to transduce the ethylene signal to downstream transcription factors, such as EIN3, EIN3 LIKE1 (EIL1) (An et al., 2010; Chao et al., 1997) and ETHYLENE RESPONSE FACTORs (ERFs) (Muller and Munne-Bosch, 2015). These transcription factors activate the downstream ethylene response. The ethylene pathway was found to play an important role in numerous growth and developmental processes in the root, such as the inhibition of root elongation (Kieber et al., 1993; Le et al., 2001; Vaseva et al., 2018), induction of root hair growth (Feng et al., 2017), and inhibition of the initiation of lateral roots (LRs) (Negi et al., 2008).

To identify potential regulators of GSA, we performed a chemical genetics screen and
identified a small molecule, Mebendazole (MBZ), which dramatically increased GSA.
Genetic analysis revealed that MBZ increased GSA through rapidly activating the
ethylene signaling pathway. Both MBZ and ACC treatment induced ethylene signaling,
which led to a modulation of lateral root angle. Further molecular and biochemistry studies
showed that MBZ activates ethylene signaling by specifically targeting CTR1 and thereby
blocking its kinase activity. Our study not only identified MBZ as an efficient small
molecular activator of ethylene signaling, but also uncovered that ethylene signaling plays
an important role in regulating GSA.
105
Results
A small compound profoundly affects the lateral root gravitropic set-point angle
(GSA) in Arabidopsis
To identify small molecules that affect GSA, we screened chemicals for their ability to
change GSA of young seedlings from a small molecular library with 2000 diverse generic
chemicals (SP 2000)(He et al., 2011; Sun et al., 2017; Zhu et al., 2019). We found that
treatment with Mebendazole (MBZ) (Figure S1A) led to a distinct increase of lateral root
angles that caused lateral roots to grow in a more horizontal direction (Figure 1A). The
lateral root GSA of untreated two-week-old Arabidopsis seedlings was approx. 46°, and
the GSA of lateral roots of seedlings grown on 1 µM MBZ plates was approx. 70° (Figure
1A, B). While the lateral root angle was the most striking phenotype, at a much earlier
developmental stage (one week after germination), seedlings on 1 µM MBZ displayed a
decreased root length (in average reduced by 37%) and an increased root hair initiation
and elongation (Figure 1C, D, Figure S1B, C). To test how quickly MBZ would affect
lateral root growth, we grew seedlings for 14 days on ½ MS medium with DMSO, then
transferred them to ½ MS medium supplemented with 1 µM MBZ and acquired images
every 10 minutes. All of the 5 measured lateral root tips started to grow at a more
horizontal (less steep) angle within 3 hours of MBZ treatment (Figure 1E upper panel, F),
which coincided with the first timepoint at which root angles were significantly different
between control and MBZ (Welch Two Sample t-test; p-value <0.05). To test whether the
action of MBZ was reversible, we germinated and grew seedlings on ½ MS for 5 days,

and then transferred them to medium containing 1 µM MBZ for 14 days, thereby inducing

lateral roots of MBZ treated seedlings to grow more horizontally. We note that we used

this later timepoint (compared to the data presented in the upper panel) to be able to

consider enough lateral roots that were growing at the more horizontal angle. We then

transferred the seedlings back to control conditions and acquired images every 10

minutes. Already after one hour root angles were significantly different between control

(MBZ) and treatment (removal from MBZ) conditions (Welch Two Sample t-test; p-value

<0.05), and within 4 hours all 5 lateral root tips we measured started to grow at a distinctly

steeper angle upon removal from MBZ (Figure 1E bottom panel, F). These data showed

that the treatment of Arabidopsis roots with MBZ changes lateral root angles within 3

hours and that this effect is fully reversible. In summary, we found that MBZ changes

lateral root angles and important RSA traits associated with the development of a shallow

root system.

Application of MBZ induces the ethylene pathway

To identify the pathway that MBZ acted upon, we utilized a genome-scale expression

profiling approach. For this, we performed an mRNA-seq experiment using whole roots

of 14-day-old Arabidopsis grown on ½ MS media treated with 10 µM MBZ for 4 hours.

The analysis revealed that 677 genes were induced compared to mock treatment

(FDR<0.05; log2 fold change > 1) and 658 genes were repressed (FDR<0.05; log2 fold

change < -1). A gene ontology (GO) analysis revealed that among the genes that were

upregulated by MBZ, genes involved in ethylene responses were strongly

overrepresented in this early response. In particular, the GO process "negative regulation

of ethylene-activated signaling pathway" (GO:0010105) was the most enriched process

(~21 fold enrichment, P-value: $6.07*10^{-4}$) upon MBZ treatment (Figure 2A), and the 4th

most enriched GO process was "ethylene-activated signaling pathway" (GO:0009873)

(~11 fold enrichment, P-value:$7.45*10^{-5}$) (Figure 2A). Other GO categories were also

significantly enriched, most notably responses to light. However, given that ethylene

signaling was the most enriched GO term, we first looked at the responses of genes

involved in the ethylene hormone pathway. A major portion of these genes were

differentially regulated by MBZ compared to mock treatment (Figure 2A, B). We therefore

hypothesized that MBZ elicited the ethylene pathway. To test this hypothesis, we first

compared the set of genes that had responded to MBZ with a published transcriptome

dataset that had been generated by exposing roots to 4 hours of ethylene treatment

(Chang et al., 2013). Of the 1335 differentially expressed genes (DEGs) upon the 4 hours

of MBZ treatment, 306 were overlapping with 1397 DEGs upon 4 hours of ethylene

treatment (P-value for the overlap using a hypergeometric test: $p < 4.657*10^{-143}$) (Figure

S3A). Notably, more than 90% of these common DEGs retained the directionality of their

expression change (i.e. genes upregulated by MBZ were also upregulated by ethylene

treatment and vice versa) (Figure S3B, C). While this overlap was highly significant and

supported the hypothesis that MBZ elicited ethylene responses, there was a high number

of differentially expressed genes that didn't overlap. We reasoned that this might be due

to differences in developmental and growth conditions of our MBZ-treated plants and of

those plants used for the published ethylene treatment dataset (Chang et al., 2013). For

instance, we had used 14-day-old seedlings grown under long day conditions before we

conducted the treatment and the published data had utilized 3-day-old, dark-grown

seedlings for eliciting the classical ethylene triple response phenotype. To compare

ethylene responses to MBZ responses under the same growth conditions and plant age,

we performed another mRNA-seq experiment using roots of 14-day-old Arabidopsis

grown on ½ MS media treated with 10 µM MBZ or 10 µM ACC (the ethylene precursor)

for 3 hours. We then again compared the overlap of the responsive genes. Out of 826

DEGs upon the MBZ treatment, 524 were overlapping with 741 DEGs upon 3 hours of

ACC treatment, accounting for 63 and 71% of DEGs upon MBZ and ACC treatment

respectively (Figure 2C). 77% upregulated DEGs of the MBZ treatment overlapped with

65% upregulated DEGs of the ACC treatment (Figure 2D). These fractions were 54% and

75% for downregulated DEGs (Figure 2E). Taken together, our data strongly suggested

that application of MBZ induces the ethylene pathway but in addition might also elicit non-

ethylene related effects.

MBZ induces ethylene responses

To investigate how MBZ treatment effects the ethylene response, we compared plant

responses to MBZ and the ethylene precursor ACC. The most characteristic response to

ethylene treatment is the triple response of etiolated seedlings (Guzman and Ecker, 1990)

(Figure 3A to C), which includes shortened and thickened roots and hypocotyls, and an

exaggerated apical hook. Consistent with our hypothesis that MBZ activates ethylene

responses, 3-day-old etiolated seedlings treated with 10 µM MBZ showed a prominent

triple response phenotype (Figure 3A to C). Using different doses of MBZ, we found that

2.5 µM of MBZ was sufficient to induce the triple response phenotype in a dark

environment (Figure S3A to F), we therefore started to use 2.5 µM of MBZ to observe

triple responses. Consistent with MBZ mimicking ethylene treatment, MBZ induced triple

response was abolished in the ethylene insensitive mutants *ein2-5* and *ein3 eil1* (Figure

S3D to F). Ethylene treatment inhibits root elongation mainly by repressing cell elongation

rather than reducing the length of meristem (De Cnodder et al., 2005; Vaseva et al., 2018).

We therefore measured the length of root cells of 4-day-old etiolated seedlings treated

with 2.5 µM MBZ and 10 µM ACC, and found that similar to the ACC treatment, the mature

cell size was reduced by MBZ treatment (Figure 3D to F). Altogether, our data strongly

suggested that MBZ elicits its effects via the ethylene pathway.

As we wanted to identify the mechanism by which MBZ changes the steepness of lateral

root angles and given our data that MBZ acts on the ethylene pathway, we conducted an

RNA-seq experiment on lateral root tips that had been treated either with DMSO or MBZ

for 3 hours. As sample collection was very challenging and took an hour per replicate, the

sampled lateral roots had been treated for a time between 3-4 hours. The most enriched

gene ontology categories were "regulation of transcription", "response to auxin", "ethylene

activated signaling pathway" and "cytokinin-signaling pathway" (Figure 2F). Given that

the time window for the RNA-seq coincided with the change of lateral root angle, it was

not surprising to detect auxin and cytokinin related genes, as these two hormonal

pathways had been shown to be involved in determining the GSA of lateral roots

(Rosquete et al., 2013a; Roychoudhry et al., 2013; Waidmann et al., 2019b). However,

given the enrichment of genes assigned to these pathways we wanted to check whether

these pathways are activated before or after ethylene responses were elicited by MBZ.

To do this, we obtained the *DR5-GFP V2* auxin response reporter line (Liao et al., 2015a),

the *35S-EIN3-GFP* ethylene reporter line in the *ein3 eil1* background (He et al., 2011),

and the *TCN-GFP* cytokinin reporter line (Zurcher et al., 2013). We then conducted time-

lapse spinning disk confocal microscopy on lateral root tips from 14-day-old seedlings
that were either treated with a DMSO control solution, MBZ, or the respective positive
control (the auxin IAA, the cytokinin Zeatin, or the ethylene precursor ACC). All positive
control treatments swiftly induced the expression of the reporter genes in lateral roots
(Figure 2G-L; Suppl. Movies 1, 3, 4, 6, 7, 9). Consistent with our hypothesis that MBZ
induced the ethylene pathway, MBZ treatment drastically induced the ethylene reporter
within the first 20 minutes after treatment, even before the ethylene precursor ACC
induced the reporter (Figure 2G, J; Suppl. Movies 1-3). MBZ treatment did not lead to an
increased signal of the auxin response reporter, while the DMSO control led to a slight
decrease of the reporter signal after 40 minutes, and the IAA treatment led to a strong
increase throughout the whole duration of the experiment (Figure 2H, K; Suppl. Movies
4-6). There was no induction of the cytokinin reporter by MBZ in the timeframe we
observed the roots (Figure 2I, L; Suppl. Movies 7-9). Overall, these data strongly
suggested that MBZ strongly and swiftly induces the ethylene pathway in lateral root tips.
234

MBZ activates ethylene signaling
Next, we set out to locate the target of MBZ in the ethylene pathway. First, we tested
whether MBZ induces ethylene biosynthesis, or acts on the perception of ethylene, or on
the ethylene signaling pathway. To test whether ethylene biosynthesis is affected, we
measured ethylene production by GC-MS in 3-day-old etiolated Col-0 seedlings that were
treated with 10 µM MBZ or ACC. Ethylene production was dramatically induced by 10 µM
ACC, but not detected in DMSO or 10 µM MBZ-treated-seedlings (Figure 4A). The results
suggested that MBZ acts downstream of ethylene biosynthesis. To further test this
hypothesis, we utilized gain-of-function mutants in the ethylene receptor *ETR1*. Root
growth and GSA in strong *ETR1* alleles *etr1-1 and etr1-3,* were still strongly altered upon
MBZ treatment (Figure 4B to E). Moreover, 4-day-old etiolated seedlings of *etr1-3* treated
with 2.5 µM MBZ showed the typical triple response phenotype, demonstrating that
ethylene responses are triggered by MBZ even without induced ethylene production or
the perception of ethylene by ETR1 (Figure S3A to C). In contrast to the response to MBZ,
etr1-3 plants were completely insensitive to the treatment with 10 µM of ACC (Figure S3A

to C). Collectively, these results show that MBZ induces ethylene responses by targeting
factors downstream of ethylene biosynthesis and perception.

To uncover which components of ethylene signaling are targeted by MBZ, we studied
MBZ responses in key mutants of the ethylene signaling pathway. First, we checked the
root inhibition phenotype and GSA in *ctr1-1* mutant treated with MBZ and compared these
responses to the Col-0 wild type. *CTR1* is the first negative regulator of ethylene signaling
acting downstream of ethylene receptors, and *EIN2* and *EIN3/EIL1* are positive regulators
that act downstream of *CTR1* to trigger ethylene signaling (Figure 5A). Roots of Col-0
were sensitive to 1.2 µM MBZ treatment and showed short roots and increased root hair
development (Figure 5B, C). However, *ctr1-1* mutants in which ethylene signaling is
constitutively active showed short roots and abundant root hairs without any treatment at
an early stage. Due to this pronounced early-seedling phenotype, we could not
confidently assess changes in response to MBZ. However, we noticed that root growth
remained insensitive to MBZ (Figure 5B, C). We further measured GSA at a later stage
of development, which we had found to be a major effect of MBZ treatment. 3-week-old
ctr1-1 plants were insensitive to MBZ with regards to GSA (Figure 5D, E). Interestingly,
ctr1-1 without MBZ treatment showed an increased lateral root angle phenotype, which
mimics MBZ treatment, suggesting that CTR1 might be the target of MBZ in ethylene
signaling. Consistent with this hypothesis, root growth and GSA of *ein2-5* and *ein3 eil1*
was clearly not affected by MBZ treatment (Figure 5B, C, F, G) and in fact, *ein2-5* mutant
plants, which are insensitive to ethylene, displayed steeper lateral root angles under
control conditions compared to the wildtype (Figure 5F, G). We further tested triple
responses in *ctr1-1*, *ein2-5* and *ein3 eil1* mutants. Consistent with the observation of root
inhibition and GSA in *ctr1-1*, these plants showed a constitutive triple response regardless
of MBZ treatment (Figure S3D to F). *ein2-5* and *ein3 eil1* plants were also insensitive to
MBZ treatment (Figure S3D to F). Taken together our data showed that MBZ activates
ethylene signaling most likely by interfering with CTR1 in the ethylene signaling pathway.
277

MBZ is a potent inhibitor of CTR1 kinase activity
To pinpoint whether CTR1 is the target of MBZ, we capitalized on what is known about
the activity of MBZ in mammalian systems. There, MBZ has been reported to be an

inhibitor of the serine/threonine kinase *homo sapiens* (*hs*) MAPK14 (Ariey-Bonnet et al.,

2020). Because CTR1 is also a serine/threonine protein kinase (Kieber et al., 1993), we

hypothesized that CTR1 is the direct target of MBZ in plants. Since it was shown that

there are four critical amino acid residues in the catalytic site of *hs*MAPK14 that are

relevant for MBZ binding (Ariey-Bonnet et al., 2020), we first tested *in silico* whether these

residues were conserved between *hs*MAPK14 and CTR1. For this, we aligned the amino

acid sequence of the Arabidopsis CTR1 and *hs*MAPK14 kinase domains (Figure 6A,

Figure S4A, B). We found three (Lys 578, Thr 625, and Asp 694) of the four critical amino

acid residues were conserved between the CTR1 kinase domain and that of *hs*MAPK14

(Figure 6A). The 4th critical residue is Met 109 in *hs*MAPK14, which corresponds to Leu

628 in CTR1. Because both Met and Leu are non-polar amino acids with hydrophobic

side chains and similar structure, it appeared possible that the replacement of Met by Leu

in the CTR1 kinase domain does not significantly change the pocket structure for MBZ

binding. We therefore hypothesized that the kinase domains of CTR1 and MAPK14 might

have similar structures and function with regards to binding MBZ (Figure S4B).

To test this hypothesis, we first performed an *in silico* docking analysis using the published

crystal structures of CTR1 (Mayerhofer et al., 2012)(PDB: 3PPZ) and MBZ (CHEMBL685).

We found that MBZ is predicted to bind into the pocket of the CTR1 kinase domain (Figure

6B) formed by the aforementioned key amino acids (Figure 6A, B, Figure S4B). Overall,

these analyses indicated that CTR1 could be the target of MBZ in Arabidopsis.

To further test this model, we expressed the recombinant kinase domains (KD) of CTR1

and MAPK4 (a MAPK protein in Arabidopsis with a highly similar amino acid sequence to

*hs*MAPK14) (Table S1, Figure S4C, D). We then confirmed the kinase inhibitory activity

of MBZ through an *in vitro* phosphorylation assay via anti-Thiophosphate-Ester antibodies

(Allen et al., 2007). Using the phosphorylation state of MBP (substrate) as the readout of

the kinase activity, we found that MBZ inhibited the kinase activities of both CTR1-KD

and MPK4 in a concentration-dependent manner, however, the inhibition for CTR1-KD

was much more pronounced (Figure 6C, D). In particular, 10 µM MBZ strongly inhibited

the kinase activity of CTR1-KD, while 100 µM MBZ totally blocked its activity. This assay

demonstrated that MBZ can directly inhibit the kinase activity of CTR1-KD. Together with

our genetic data that *ctr1-1* mutant mimicked the effect of MBZ on GSA and the triple

response phenotypes (Figure 5D, E, Figure S3D-F), and our molecular docking data
(Figure 6B), our data strongly suggest that CTR1 is a direct target of MBZ in Arabidopsis.
To further test the hypothesis that MBZ acts upon CTR1, we utilized a readout directly
downstream of CTR1. Previous studies had shown that functional CTR1 phosphorylates
the carboxyl end of EIN2, leading to its degradation (Ju et al., 2012; Qiao et al., 2012a;
Wen et al., 2012). When ethylene inactivates CTR1, EIN2-CEND is stabilized and moves
into the nucleus to activate downstream ethylene responses. We therefore tested if MBZ
treatment mimicked ethylene for inducing EIN2 accumulation in the nucleus using 3-day-
old etiolated seedlings of two independent transgenic *EIN2-GFP* reporter lines in which
the GFP was fused to EIN2's C-terminus (*pEIN2::EIN2-GFP* in the *ein2-5* background (Fu
et al., 2021)). As expected, treatment with ACC quickly increased the nuclear
accumulation of EIN2-GFP in root cells (Figure 7A, B). Consistent with our hypothesis
that MBZ acted on CTR1, MBZ also significantly increased accumulation of EIN2-GFP in
the nucleus (Figure 7A, B). Taken together, our data support a model in which MBZ is a
small molecular inhibitor of the CTR1 kinase activity and thereby triggers the ethylene
signaling pathway by permitting C-terminal of EIN2 translocation into the nucleus, which
in turn triggers ethylene downstream responses (Figure 7C, D).
329
DISCUSSION
Our chemical genetic screen led to the identification of MBZ as a potent inhibitor of the
kinase activity of CTR1, which is a negative modulator of the ethylene signaling pathway
(Figure 7C). MBZ therefore is an activator of ethylene signaling, and to our knowledge
the first small molecule that acts downstream of ethylene receptors and that directly
activates ethylene signaling. In previous studies, ethylene gas and ACC have been used
to activate the ethylene signaling pathway. However, there were some disadvantages to
use these approaches. In the case of treatment with ethylene gas, it is difficult to maintain
a continuous treatment over longer periods of time due to the technical challenges for
constantly maintaining the required ethylene concentrations. Because its controlled
application is technically less challenging, ACC has been used more widely to study the
role of ethylene. However, ACC was recently found to be a signaling molecule which can
act independently of the ethylene pathway (Mou et al., 2020). Moreover, the necessity for

enzymatic ACC conversion to ethylene in plants might be limiting in different tissues or
conditions. While MBZ is not a perfect ethylene substitute and might have unspecific
effects beyond activating ethylene signaling, as a direct activator of the ethylene signaling
pathway, it promises to be a useful tool for the field. Such tools have been lacking so far
despite several attempts that have been made to identify small molecular regulators of
ethylene signaling. Neither approaches targeting ethylene biogenesis and perception, nor
approaches targeting downstream auxin biosynthesis had yielded in molecules acting on
ethylene signaling (He et al., 2011; Li et al., 2017; Sun et al., 2017; Zhu et al., 2019).

Because CTR1 contains the critical MBZ binding residues that had been elucidated in the
human MBZ target *hs*MPK14, and because of the profound genetic evidence for CTR1,
it is somewhat unlikely that other components in the ethylene pathway such as the E3
ligases ETP1/2 and EBF1/2, which negatively regulate EIN2 and EIN3 respectively (Guo
and Ecker, 2003; Qiao et al., 2012b), are targets of MBZ. Another signaling pathway that
impinges on ethylene signaling is the glucose-activated TOR signaling pathway, which
had recently been shown to regulate EIN2 activity through phosphorylation (Fu et al.,
2021). However, the TOR kinase does not contain any of the 4 key residues known for
MBZ binding. Moreover, unlike MBZ, none of the inhibitors of TOR, including Rap, Torin2,
and AMA, did induce the ethylene induced triple response phenotype (Fu et al., 2021). In
addition, *ein2-5* and *ein3 eil1* mutants displayed a shorter hypocotyl when the Glu-TOR-
EIN2 pathway was inhibited (Fu et al Nature 2021, Figure 1b), but these mutants did not
respond to MBZ treatment (Figure S3D, E). Altogether, it is most likely that the observed
response of MBZ on the ethylene signaling is exerted via its binding to CTR1.

While most of the effects of MBZ are related to its activation of ethylene signaling, our
RNA profiling indicated additional effects of MBZ. Consistently, treatment with high
concentration of MBZ or long-term treatment with it led to strongly inhibited plant growth
and development, including a disrupted root apical meristem and morphological changes
of leaves. These phenotypes have not been observed upon ethylene treatment (Figure
S5A, B), raising the possibility that MBZ may have other targets in plants. MAPK4 has the
highest amino acid sequence similarity to the human MBZ target *hs*MAPK14 (Table S1).

Compared to CTR1, the effect of MBZ on MAPK4 kinase activity was much less
pronounced (Figure 6C), consistent with the lack of conservation of key amino acids
relevant for MBZ binding (Figure S4C). Although MBZ couldn't inhibit the kinase activity
of MAPK4 as strongly as that of CTR1-KD, a decrease of the weak MAPK4
autophosphorylation band might indicate that MBZ treatment can disrupt MAPK4
autophosphorylation (Figure 6C). This raises the possibility that MBZ might alter
autophosphorylation of MAPK4 and probably other CTR1-like kinases, based on the
sequence conservation of the four critical residues (Figure 6A, Figure S4B). This might
explain the additional effects of MBZ. Additionally, there are a number of CTR1 paralogs
in Arabidopsis. This includes the kinases encoded by AT4G24480, AT1G08720,
At1g18160, AT1G67920, and AT5G11850. These kinases contain all four critical amino
acids, raising the possibility that MBZ may also bind to these paralogs. Another
explanation for the observed effects of MBZ that go beyond observed ethylene effects,
might be that continuous activation of ethylene signaling via CTR1 might have more
complicated or compounded effects than are elicited by external ethylene or ACC
treatment (Figure S5C). Support for this comes from the *ctr1-1* mutant, which is dwarfed
and shows aberration in leaf shape and thereby displays much stronger phenotypes than
elicited with ACC or ethylene treatment (Kieber et al., 1993) (Figure 5B, D, Figure S5C).
While MBZ's predominant effect on ethylene signaling via CTR1 is strongly supported by
our study, further studies will be needed to elucidate the details of MBZ specificity.

One of the most intriguing effects of MBZ on the plant is its potent ability to change GSA.
This suggests that ethylene signaling is involved in the regulation of the gravitropic
setpoint angle in lateral roots. While important regulatory roles for GSA are known for
auxin and cytokinin, and there can be complex interplays between auxin, cytokinin and
ethylene, the induction of the ethylene signaling pathway clearly dominated responses in
lateral roots after they were treated with MBZ (Figure 2). This activation of the ethylene
responses occurred before angles of lateral roots were changing and was maintained
throughout the timescale at which GSA was changed by the MBZ treatment, which
occurred after 1-4 hours of MBZ treatment (Figure 1E, F, Figure 2, Figure S2I). In our
study, we found that MBZ didn't induce ethylene biosynthesis (Figure 4A). ACC treatment

had a weak but distinct effect on root angle and displayed concentration-dependent effects when applying a large range of concentrations (Figure S5C). Strong direct evidence for the impact of ethylene signaling in regulating lateral root set point angle is provided by the RSA phenotype of the *ctr1-1* mutant. In *ctr1-1*, ethylene signaling is constitutively activated and leads to an increased lateral root angle (Figure 5D, E). Conversely, we observed that the ethylene insensitive *ein2-5* mutant displays steeper lateral roots (Figure 5F, G). Overall, this exposes a yet unknown role of ethylene signaling in the regulation of lateral root set point angle. One reason why the role of ethylene in the regulation of lateral root angles hasn't been recognized so far might be that it reduces lateral root formation by itself (Negi et al., 2008) making the study of lateral root angles more difficult. Our study strongly suggests that it will be worthwhile to explore the role of ethylene and its interaction with auxin and cytokinin in root gravitropism and RSA. Moreover, MBZ might also be used as a versatile tool for exploring the molecular mechanisms of root gravitropism and lateral root set point angles since as a small molecule, and due to its reversible mode of action, it can be easily applied to elicit controlled changes of lateral root angles. It might also be useful to reveal the mechanisms underlying GSA regulation by phytohormones and known GSA signals, such as the *LAZY* family genes (Furutani and Morita, 2021). Finally, since ethylene signaling is a widely conserved process in land plants, exploring the ability of MBZ or genetic perturbation of the ethylene pathway to change RSA in crop species might be highly worthwhile for RSA engineering.

Acknowledgements

The research was supported by start-up funds from the Salk Institute for Biological Studies and funds from the Harnessing Plants Initiative to WB, a Salk Women & Science Special Award and a Pioneer Fund Postdoctoral Scholar Award to WH. We are grateful to Dr. Glenn R. Hicks and Dr. Natasha Raikhel for sharing the SP2000 small molecule library. We thank Dr. Cris Argueso, Dr. Dolf Weijers, and Dr. Liwen Fu for sharing seeds. We also thank Dr. Joseph R. Ecker and Dr. Joanne Chory for critical discussions. We thank the Next Generation Sequencing Core of Salk institute for NGS Library preparation

435 & sequencing. We also thank Matthieu Platre, Nicole Gibbs, Anna Malolepszy, Björn
436 Willige and Travis Lee for fruitful discussions and suggestions.
437

Author contributions
Wenrong He and Wolfgang Busch conceived the project and the experiments. Wenrong
He performed the SP2000 library screen and analysis with help from Yingnan Hou.
Wenrong He performed the phenotyping, qPCR, alignment, docking, protein purification,
kinase activity assay. Wenrong He conducted the microscopy and mRNA-seq on primary
roots with help from Kaizhen Zhong. Hai An Truong performed the time lapse reporter
analysis using spinning disk microscopy, as well as the whole root system and lateral root
RNA-seq experiments. Ling Zhang conducted the mRNA-seq analysis and the statistical
analysis. Wenrong He conducted heatmap analysis of mRNA-seq. Wenrong He and Ling
Zhang conducted Gene ontology analysis. Min Cao provided the protocol and
suggestions for kinase activity assay. Neal Arakawa performed the GC test. Yao Xiao
performed the FPLC. Wenrong He and Wolfgang Busch wrote the manuscript with input
from all authors. Wolfgang Busch provided materials, funding, supervision, guidance and
advice on the project, the experiments, and the analysis of the results.
452

453 **Declaration of Interests**
454 W.B. is a co-founder of Cquesta, a company that works on crop root growth and carbon
455 sequestration.
456

458 **FIGURE LEGENDS**
459

Figure1. MBZ treatment perturbs root system architecture (RSA).
(A) 14-day-old Arabidopsis seedlings of grown on DMSO and MBZ (1 µM) plates.
(B) Quantification of GSA in (A). 15 LRs per seedling on DMSO plates, 12 LRs per
seedling on MBZ plates, and 6 seedlings for each condition were used. Unpaired, two-
tailed Student's t-tests was used for statistical analysis. **** p < 0.0001.
(C) 7-day-old seedlings on DMSO and MBZ (1 µM) plates.

(D) Quantification of primary root length (cm) in (B). 15-20 seedlings for each treatment
were used. **** p < 0.0001.
(E) 14-day-old (upper panel) Arabidopsis seedlings on DMSO plates were transferred to
MBZ (1 µM) plates, and 19-day-old (bottom panel) seedlings on MBZ (1 µM) plates were
transferred to DMSO plates, followed by continuous scanning for 24 h.
(F) Changes of lateral root angle after plate transfer from DMSO to MBZ (red and yellow
lines) or MBZ to DMSO (blue, purple lines) compared to timepoint 0.
(Scale bar: A, C, G, 1 cm; E, 5 mm.). Boxplots: Whiskers: min/max values; hinges: 25th
to 75th percentile; mid-line: median.
475
Figure 2. MBZ treatment regulates the ethylene pathway.
(A) Gene ontology (GO) analysis upregulated genes (abs(log2) fold change > 1) from
DEGs upon MBZ treatment in roots of 14-day-old Arabidopsis seedlings.
(B) Heatmap of DEGs involved in the ethylene pathway, which were detected to be
differentially expressed upon MBZ treatment in the RNA-seq analysis. The left 4 columns
of DMSO are data from 4 replicates of samples from DMSO (Mock) plates. The right 4
columns of MBZ are data from 4 replicates of samples from MBZ treatment plates. The
heatmap was made online by heatmapper (http://heatmapper.ca/expression/), The
cluster method is complete linkage. Color bar indicates the relative expression level.
(C-E) Venn diagrams showing differentially expressed genes in 3 h MBZ treatment and 3
h ACC treatment. All genes (FDR < 0.05; abs(log2) fold change > 1) (C), upregulated
genes (FDR < 0.05; log2 fold change > 1) (D), downregulated genes (log2 fold change <
-1) (E).
(F) Dot plots showing the most enriched GO terms with KS p-value < 0.05 from DEGs
upon MBZ treatment in lateral root tips of 14-day-old seedlings.
(G-I**)** Representative images of lateral root tips of two-week-old plants in response to
negative control, positive control and MBZ treatment using ethylene (*EIN3-GFP*)(G),
auxin (*DR5v2-GFP*)(H), and cytokinin (*TCSn-GFP*)(I) reporter lines at 0 and 60 minutes
of treatment.
(J) Relative fluorescent intensity of *EIN3-GFP* lateral root tips treated over a course of 4
h with DMSO, MBZ, and ACC.

(K) Relative fluorescent intensity of *DR5v2-GFP* lateral root tips treated over a course of 4 h with DMSO, MBZ, and IAA.

(L) Relative fluorescent intensity of *TCSn-GFP* lateral root tips treated over a course of 4 h with DMSO, MBZ, and tZ. 6 lateral roots for each treatment were used. Unpaired, two-tailed Student's t-tests were used to compare the data of MBZ or the corresponding positive controls to DMSO at each timepoint. *: p<0.05; **: p<0.01; ***: p<0.001.

Figure 3. MBZ induces ethylene responses and mimics ACC treatment.

(A) 4-day-old etiolated seedlings grown on MS, 10 µM ACC, DMSO, 10 µM MBZ plates. 20 Col-0 seedlings for each treatment were observed and 2 representative seedlings for each treatment are shown in the panel.

(B-C) Quantification of hypocotyl length (B) and primary root length (C) of seedlings in (A). 20 Col-0 seedlings for each treatment were used. **** p < 0.0001.

(D) Images of median longitudinal optical sections of primary root meristem and mature zone stained with propidium iodide (PI) of 4-day-old etiolated Col-0 on DMSO, 2.5 µM MBZ, and 10 µM ACC plates, 6-10 seedlings were observed for each treatment.

(E-F) Meristem length (E) and mature cell size (F) of seedlings in (D). 6-10 seedlings for each treatment were used. Different letters label significant different values (p<0.0001). One-way ANOVA and *post hoc* Tukey testing were used for statistical analysis in (E) and (F). (Scale bar: **A**, 5 mm; **D**, 100 µm.). Boxplots: Whiskers: min/max values; hinges: 25^{th} to 75^{th} percentile; mid-line: median.

Figure 4. MBZ acts downstream of ethylene biosynthesis.

(A) Quantification of ethylene production by GC-MS of 4-day-old etiolated seedlings of Col-0 in vials with DMSO, 10 µM MBZ, and 10 µM ACC medium. 100 seedlings for every vial, 4 biological replicates for each treatment, ND: not detectable.

(B) 6-day-old seedlings of Col-0, *etr1-1*, and *etr1-3* grown on DMSO and 1.2 µM MBZ plates. 20 seedlings were observed.

(C) Quantification of root length in (A). 20 seedlings were countified.

(D) 16-day-old seedlings of Col-0 and *etr1-1* grown on DMSO and 1.2 µM MBZ plates. 20 seedlings were observed.

(E) Quantification of GSA in (C). 20 seedlings were countified.

One-way ANOVA and *post hoc* Tukey testing were used for statistical analysis in b and

d. Different letters label significant different values (p<0.0001). (Scale bar: **B**, **D**, 1 cm.)

Boxplots: Whiskers: min/max values; hinges: 25th to 75th percentile; mid-line: median.

Figure 5. MBZ targets the ethylene signaling pathway.

(A) Diagram of ethylene signaling pathway.

(B) 7-day-old seedlings of Col-0, *ctr1-1*, and *ein2-5* grown on DMSO and 1.2 µM MBZ

plates.

(C) Quantification of primary root length of seedlings in (B) 15-20 seedlings were

quantified. Similar results were obtained from three biological replicates of the

experiment.

(D) 19-day-old seedlings of Col-0 and *ctr1-1* grown on DMSO and 1.2 µM MBZ plates.

(E) Quantification of GSA of seedlings in (D). 10 seedlings were quantified. Similar results

were obtained from three biological replicates of the experiment.

(F) 16-day-old seedlings of Col-0 and *ein2-5*, *ein3 eil1* grown on DMSO and 1.2 µM MBZ

plates.

(G) Quantification of GSA of seedlings in (F). *dm* stands for *ein3 eil1* double mutants.

One-way ANOVA and *post hoc* Tukey testing were used for statistical analysis in **C**, **E**,

and **G**. Different letters label significant different values (p<0.0001). (Scale bar: **B**, **D**, **F**,

1 cm.). Boxplots: Whiskers: min/max values; hinges: 25th to 75th percentile; mid-line:

median.

Figure 6. MBZ inhibits CTR1 kinase activity.

(A) Sequence alignment of *hs*MAPK14 and CTR1-KD in Arabidopsis around 4 core amino

acids (Highlighted by blue and pink frames) for MBZ binding.

(B) Molecular modeling of the interaction between MBZ (coppery) and CTR1-KD (silvery).

In the left panel, the key residues that contribute to the binding with CTR1-KD were

highlighted. The right panel showed the surface (gray) of the catalytic pocket of CTR1-

KD.

(C) Western blot using Recombinant Anti-Thiophosphate ester antibody under different

concentration of MBZ treatment. Left panel: CTR1-KD; Right panel: MAPK4. MBP is the

substrate in both panels, MBZ concentration is 0, 10, and 100 μM as labeled in different

reactions, [ATPγS]CTR1-KD: ATPγS binding CTR1-KD, [ATPγS]MAPK4: ATPγS binding

MAPK4, MBP that obtained ATPγS from CTR1-KD or MAPK4, which is labeled as

[ATPγS]MBP. Note: ATPγS: Adenosine 5'-(3-thiotriphosphate) tetralithium salt (Sigma,

Cat# A1388)

(D) Coomassie blue staining of the gel shown in (C). CTR1: ~100 ng, MAPK4: ~1 μg.

Myelin Basic Protein (MBP, a common kinase substrate, Sigma Cat. M1891): 2 μg,

ATPγS: 1 mM. MBZ: 0, 10, or 100 μM. Arrows label the position of these bands on the

gel.

Figure 7. MBZ activates ethylene signaling.

(A) Different patterns of nuclear accumulation of *EIN2-GFP* in root cells. 3-day-old

etiolated seedlings treated with Mock, ACC (100 μM), or MBZ (10 μM) for 3 hours. Arrows

indicate nuclei. (Scale bar: 20 μm).

(B) Quantification of Relative GFP fluorescent intensity between the nucleus and cytosol

of *EIN2-GFP* in **A**. Boxplots: Whiskers: min/max values; hinges: 25^{th} to 75^{th} percentile;

mid-line: median. One-way ANOVA and *post hoc* Tukey testing were used for statistical

analysis.

(C,D) Model for MBZ activation of ethylene signaling. (C) Ethylene suppresses CTR1

activity to promote EIN2C translocation and activates ethylene signaling. (D) MBZ inhibits

CTR1 activity by binding to its kinase domain and promotes EIN2C translocation to

activate the ethylene signaling.

Note: EIN2C: C-terminal of EIN2; ATP: Adenosine triphosphate; ADP: adenosine

diphosphate; C2H4: ethylene; P: Phosphate group. The CTR1 with orange color means

active CTR1 proteins and the CTR1 with gray color means inactive CTR1.

Figure S1. MBZ treatment affects root growth and development.

(A) Chemical structure of MBZ.

(B-C) Root hair phenotype of 7-day-old seedlings on DMSO and 1 µM MBZ plates
recorded by a scanner (B) or a microscope (C). 20 seedlings were observed on left panel
and 10 seedlings were observed on right panel. (Scale bar: **B**, 13 mm; **C**, 1000µm)
591
Figure S2. Analysis of differentially expressed genes (DEGs) upon MBZ treatment.
(A-C) Venn diagrams showing differentially expressed genes in 4 h MBZ treatment in
roots of 14-day-old green seedlings and published 4 h ethylene treatment in 3-day-old
etiolated seedlings. All genes (FDR<0.05; abs(log2) fold change > 1) (A), upregulated
genes (FDR<0.05; log2 fold change > 1) (B), downregulated genes (FDR<0.05; log2 fold
change < -1) (C).
(D) Gene ontology (GO) analysis of upregulated genes (FDR<0.05; log2 fold change > 1)
from DEGs upon MBZ treatment.
600
Figure S3. Etiolated seedling phenotypes in mutants of the ethylene pathway.
(A) 4-day-old etiolated seedlings of Col-0 and *etr1-3* grown on DMSO, 2.5 µM MBZ, and
10 µM ACC plates.
(B-C) Quantification of hypocotyl length (B) and root length (C) of seedlings in (A). 20
seedlings were quantified.
(D) 4-day-old etiolated seedlings of Col-0, *ctr1-1, ein2-5*, and *ein3 eil1* grown on DMSO,
2.5 µM MBZ, and 10 µM ACC plates.
(E-F) Quantification of hypocotyl length (E) and root length (F) of seedlings in **D**. 20
seedlings were quantified. Statistical analysis in **B, C** and **E, F** were done using unpaired,
two-tailed Student's t-tests. P-values are indicated in the figure. (Scale bar: **A, D**, 1 cm).
Whiskers: min/max values; hinges: 25^{th} to 75^{th} percentile; mid-line: median.
612
Figure S4. Alignment of CTR1-KD in Arabidopsis and MAPK14 in human.
(A) The structure alignment of *hs*MAPK14 (coppery) and CTR1-KD (celeste).
(B) The protein binding sites in *hs*MAPK14 (coppery) and CTR1-KD (periwinkle) with
highlighted key residues.
(C) Sequence alignment of kinase domains. Key residues in MAPK14_human and CTR1-
KD in Arabidopsis are indicated by red boxes.

(D) Phylogenetic analysis of kinases aligned in (C).

Figure S5. High concentrations or long-term treatment of MBZ induce phenotypes

that go beyond ethylene induced effects.

(A) Confocal microscopy images of a root meristem of 4-day-old etiolated WT seedling

grown on 10 µM MBZ plate.

(B) 38-day-old Col-0 plants grown on DMSO and 1.2 µM MBZ plates.

(C)14-day-old seedlings of Col-0 grown on increasing concentrations MBZ and ACC (0.1,

0.2, 0.4, 1, 2, 4, 10 µM) 20 seedlings were observed. Similar results were obtained from

three biological replicates of the experiment. (Scale bar: **A**, 100 µm, **B**, **C**, 1 cm)

Table S1. The list of blast hits in Arabidopsis using the MAPK14_human sequence

as BLASTP.

 KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Recombinant Anti-Thiophosphate ester antibody	Abcam	Cat# Ab92570
Chemicals, Peptides, and Recombinant Proteins		
SP 2000 small molecule library	(He et al., 2011)	Gift from UC Riverside
Mebendazole	ACROS	Cat# 101720050
DMSO	Alfa Aesar	Cat# 36480
ACC	Sigma	Cat# A3903
IAA	Alfa Aesar	Cat# A10556
Maxima™ H Minus cDNA Synthesis Master Mix, with dsDNase	Thermo Scientific	Cat# M1682
Luna® Universal qPCR Master Mix	NEB	Cat# M3003L
MBP	Sigma	Cat# M1891
ATP γ S	Sigma	Cat# A1388
PNBM	CAYMANCHEMICAL	Cat# 21456-1
Experimental Models: Organisms/Strains		
Arabidopsis: *etr1-1*	ABRC	CS237
Arabidopsis: *etr1-3*	ABRC	CS3070
Arabidopsis: *ctr1-1*	(Kieber et al., 1993)	N/A
Arabidopsis: *ein2-5*	(Alonso et al., 1999)	N/A
Arabidopsis: *ein3-1eil1-1*	(Alonso et al., 2003)	N/A
Arabidopsis: *ein3 eil1 35S: EIN3-GFP*	(He et al., 2011)	N/A
Arabidopsis: *UBQ10p:TUB6-GFP*	(Nakamura et al., 2004)	N/A
Arabidopsis: *35S:DR5v2:3xGFP-NLS*	(Liao et al., 2015b)	Weijers lab
Arabidopsis: *TCSn-GFP*	(Zürcher et al. 2013)	Argueso lab
Arabidopsis: *EIN2-GFP in ein2-5*	(Fu et al., 2021)	Fu Lab
Oligonucleotides		
ERF1Q-PCRF: ATTCTTTCTCATCCTCTTCTTCT	This paper	N/A
ERF1Q-PCRR: CGAATCTCTTATCTCCGCCG	This paper	N/A
AtACTIN2F: TTGACTACGAGCAGGAGATGG	This paper	N/A
AtACTIN2R: ACAAACGAGGGCTGGAACAAG	This paper	N/A
UBQ10F: CGTCAAGGCCAAGATCCAAG	This paper	N/A
UBQ10F: CAGCAAGAGTTCTGCCATCC	This paper	N/A
EcoR1-CTR1-atg-540-F: GGGGAATTCATGGATGGTGATGATATGGACATCCCG	This paper	N/A
sal-CTR1-821-R: GGGGTCGACTTACAAATCCGAGCGGTTGGGCGGA	This paper	N/A
EcoR1-MPK4-F: GGGGAATTCATGTCGGCGGAGAGTTGTTTC	This paper	N/A

sal-MPK4-R: GGGGTCGACTCACACTGAGTCTTGAGGATTGAAC	This paper	N/A
Recombinant DNA		
pET28a-CTR1-KD	This paper	N/A
pET28a-MAPK4	This paper	N/A
Commercial Kits		
Spectrum™ Plant Total RNA Kit	Sigma	Cat# STRN250
Software and Algorithms		
Fiji	(Schindelin et al., 2012)	(http://fiji.sc/Fiji).
GraphPad Prism	GraphPad	Version 9.3.1
Heatmapper	(Babicki et al., 2016)	(http://heatmapper.ca/)
AutoDock Vina	(Trott and Olson, 2010)	(https://vina.scripps.edu/)
Chimera	(Pettersen et al., 2004)	(https://www.cgl.ucsf.edu/chimera/)
Other		

CONTACT FOR REAGENT AND RESOURCE SHARING

636 Further information and requests for resources and reagents should be directed to and
637 will be fulfilled by the Lead Contact, Wolfgang Busch (wbusch@salk.edu).

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Arabidopsis thaliana

Arabidopsis mutants and transgenic lines used in this study are in the Col-0 background
if not mentioned otherwise. *etr1-1* (CS237) and *etr1-3* (CS3070) were ordered from ABRC,
ctr1-1 (Kieber et al., 1993) *ein2-5* (Alonso et al., 1999), *ein3-1 eil1-1*(Alonso et al., 2003),
ein3 eil1 35S: EIN3-GFP (He et al., 2011), and *UBQ10p:TUB6-GFP* (Nakamura et al.,
2004) were lab stock. *35S:DR5v2:3xGFP-NLS* (Liao et al., 2015b) was gift from Weijers
lab (Col-Ultrecht background). *TCSn-GFP* (Zurcher et al., 2013) was a gift from Argueso

lab. *EIN2-GFP* (Fu et al., 2021) in *ein2-5* was a gift from Fu lab. Plants were sown on half

Murashige and Skoog (MS) medium with 1% (wt/vol) sugar, 0.1 % (wt/vol) MES (pH 5.75),

containing 1% (wt/vol) agar supplemented with different chemicals as described in the

article. Following a 3-day stratification in the cold room (4~10°C), for green seedling assay,

plants were grown on vertical plates in a walk-in growth chamber with long day conditions

(16/8 h) at 21°C, 50 µM light intensity, 60% humidity. During nighttime, temperature was

decreased to 15°C. For etiolated seedling assay, the plates were placed in light for 4 h

and then germinated in the dark for 3 day in the growth chamber at 22 °C.

Plants were grown on vertical plates in a walk-in growth chamber with long day conditions

(16/8 h) at 21°C, 50 µM light intensity, 60% humidity. During nighttime, temperature was

decreased to 15°C.

Accession Numbers

Sequence data from this article can be found in the Arabidopsis Genome Initiative or

GenBank/EMBL databases or Uniprot databases under the following accession numbers:

ERF1 (AT3G23240), *ETR1* (AT1G66340), *CTR1* (AT5g03730), *EIN2* (AT5G03280), *EIN3*

(AT3g20770), *EIL1* (AT2G27050), *MAPK4* (AT4G01370), and *hsMAPK14* (HGNC:6876).

Accession numbers of protein and chemicals structures used in the molecular modeling

in Figure 6 are as follows: CTR1 (PDB: 3PPZ), MAPK14 (PDB: 6SFO) and MBZ

(CHEMBL685).

METHOD DETAILS

Small Molecule Library and Screen Information

The SP2000 small molecule library was received from the University of California,

Riverside. The library was first screened at a concentration of 50 ~ 100 µM in 200 µL of

liquid MS media using wild type Arabidopsis. After being grown in the liquid media with

small molecules for 7 days, seedlings were transferred to vertical plates to grow for

another 2 days. The root phenotypes were acquired by CCD flatbed scanners. After

candidates were picked, a second round of screen was performed by growing Col-0

seedlings on vertical plates with gradient concentration of candidate small molecules,

from 1 µM to 100 µM, for 14 days to confirm the phenotype and optimal concentration.

Cloning and constructs

To produce recombinant CTR1 kinase domain (CTR1-KD) and MAPK4 proteins in *E. coli*,
the coding sequences of 540-821 aa of CTR1and full length MAPK4 were PCR amplified
from cDNA of Arabidopsis Col-0, using gene-specific primers (listed in KEY
RESOURCES TABLE). The PCR products were inserted into pET28a using restriction
digestion and ligation with EcoR 1 and Sal 1 (NEB) sites.
682

683 **Chemical Solutions**
684 The stock solutions for preparing media for the plant experiments were Mebendazole (10
685 mM) in DMSO, ACC (10 mM) in dH2O, IAA (1 mM) in ethanol.
686

687 **Root phenotyping**
688 Images of root phenotypes on plates were acquired with CCD flatbed scanners (EPSON).
689 Root lengths and lateral root angles were measured using Fiji (http://fiji.sc/Fiji).
690

691 **Lateral root angle measurement**
692 The lateral root angle of individual LRs was measured according to (Rosquete et al.,
693 2013b) using Fiji. LRs after stage III from each seedling were measured according to
694 (Rosquete et al., 2013b; Waidmann et al., 2019a) .
695

Time lapse of lateral root angle phenotyping
14-day-old seedlings on DMSO plate or 19 d old seedling on mebendazole plate were
transferred to DMSO or 1 µM mebendazole plates as described (Figure 1E). Root
phenotype images were acquired by CCD vertical scanners (EPSON) every 10 mins over
a period of 24 hours using the tool described(Ogura et al., 2022). Root length and lateral
root angles were measured using Fiji (http://fiji.sc/Fiji).
702

mRNA-seq
A. Tissue preparation
Roots or lateral root tips of 14-day-old Col-0 seedlings with different treatment were
collected for the mRNA-seq analysis. In detail:
a. Roots

Sterilized seeds of Col-0 on ½ MS media were planted on a mesh in one row (~more than

30 seeds/plate). 5 plates were used for one treatment condition for every replicate, and 4

replicates were used for every treatment. After 14 days of vertical growth in the chamber,

liquid media of ½ MS with DMSO (0.1 %, v/v), 10 μM MBZ, and 10 μM ACC (to be

consistent, we also added the same concentration of DMSO (0.1 %, v/v) in the ACC

treatment plates) were prepared. 10 mL liquid treatment medium was used on each plate.

The mesh with seedlings from ½ MS plates was transferred the respective liquid

treatment plates and kept in growth chamber. After 4 hours for the mRNA-seq of MBZ

treatment and 3 hours for the mRNA-seq of MBZ and ACC treatment, roots were collected

using scalpel and tweezers. Excess liquid on the roots was removed using a paper towel.

Root samples were flash-frozen in liquid nitrogen and then stored at -80 freezer before

RNA extraction.

For RNA extraction, root tissues were ground in liquid nitrogen and RNA was extracted

using Quick-RNA Kits (Zymo research). RNA libraries were prepared for sequencing

using standard Illumina protocols at the Salk Next Generation Sequencing Core (NGS).

Note: We used liquid media for DMSO, MBZ, and ACC treatment in the mRNA-seq of

roots to detect transient gene regulation as these chemicals in liquid media can be

sufficiently and equally absorbed into lateral roots and reduce time for sample collection.

b. Lateral root tips

Sterilized seeds were grown for 14 days on ½ MS medium. The seedlings were then

transferred to freshly prepared DMSO or 1μM MBZ plates for 4 hours on 15 plates (5

plants/plate) for each treatment. 4 biological replications were performed. A stereo

microscope (Leica) was used to dissect the meristem area of lateral root tips in order to

collect the lateral root tips. The cut lateral root tips were quickly placed into 1.5

microcentrifuge tubes that floated on liquid nitrogen. This process was repeated until

approximately 400 root tips were collected per sample.

The NucleoSpin RNA extraction kit (Macherey-Nagel) was used to extract total RNA. The

concentration and integrity of extracted RNA were determined using Qubit and

TapeStation 2200 before being sent to Salk Next Generation Sequencing Core (NGS) for

library preparation and sequencing.

B. Data analysis

a. Read alignment

The TAIR10 genome file and annotation file were obtained from the Arabidopsis Information resource web site (http://www.arabidopsis.org)(Berardini et al., 2015). The An aligner, called the Splice Transcripts Alignments to Reference (STAR) version 2.7.0a (Dobin et al., 2013), was used to align short reads in the FASTQ files.

b. Differential expression analysis

Differential expression of genes between treatments and control was determined using the R package, *edgeR* (version 3.36.0)(Robinson et al., 2010). The *CPM* (counts per million) function from *edgeR* was used to normalize the counts and differentially expressed genes (DEGs) were identified using *glmLRT* function from edgeR. A false discovery rate (FDR < 0.05) and |log2FC| > 1 were used as the criterial values for identification of up-regulated and down-regulated DEGs.

GO analysis

Gene Ontology analysis of up-regulated DEGs (FDR < 0.05) and |log2FC| > 1 by MBZ treatment in roots of 14-day-old Arabidopsis seedlings was performed with PANTHER Overrepresentation Test (Released 20230705). The Annotation Version used GO Ontology database (DOI: 10.5281/zenodo.7942786 (Released 20230510)). Enriched genes (Bonferroni-corrected for P < 0.05) for biological processes were listed (Figure 2A). We used R package topGO (version 2.42.0)(Alexa et al., 2006) with the weight01 algorithm and the mappings from the "org.At.tair.db" R package to perform functional enrichment analysis of DEGs by MBZ treatment in roots and lateral root tips of 14-day-old Arabidopsis seedlings. We carried out KS test for statistical comparison. Then we used "ggplot2" package in R to plot 30 most enriched gene ontology terms (GO) which were ordered by Kolmogorov-Smirnov p-values for biological processes (Figure 2F, S2D).

Measurement of ethylene production.

100 seeds of Col-0 were planted and sealed in a screw-top 20 mL vial containing 10 mL ½ MS medium supplemented with DMSO (control), 10 µM mebendazole, and 10 µM ACC treatments (4 replicates). These vials were incubated at 22 °C for 3 days under dark

environment. 500-1000 μL sample of the headspace gas was sampled with an
autosampler (TriPlus RSH, Thermo Scientific) and injected into a gas chromatograph
(Trace 1310 GC, Thermo Scientific) that was equipped with a HP-PLOTQ column (30 mm,
320 μm, 20 μm) and a mass spectrometer (TSQ8000 Evo MS, Thermo Scientific),
scanning from 25-27.5 m/z. Separations were carried out at 35°C using He as the carrier
gas. The area of the ethylene peak (RT: 4 min) was integrated using Thermo Xcalibur
Qual Browser. A calibration curve was generated by varying the injection volume (100
μL, 250 μL, 500 μL, and 1000 μL) of a 10 ppm ethylene standard (in nitrogen), and sample
results are expressed as concentrations calculated from linear regression of calibration
samples.
780
qRT-PCR
For measurement of marker genes of plant hormone pathways, 20 7d old seedlings of
Col-0 grown on ½ MS media were treated with DMSO or 10μM MBZ, dH2O or 10 μM
ACC for 2h, or 3, 6, 12h as described in the manuscript. Then whole roots were collected.
RNA extraction was performed using the Spectrum™ Plant Total RNA Kit (Sigma),
followed by cDNA synthesis using Maxima™ H Minus cDNA Synthesis Master Mix, with
dsDNase (Thermo Scientific). Quantitative PCR was performed with Luna® Universal
qPCR Master Mix (NEB). All procedures were carried out according to manufacturer
instructions. qRT-PCR primer information were listed in the KEY RESOURCES TABLE.
790
Microscopy
A Zeiss 710 confocal microscope with a 40x objective was used to detect propidium iodide
(PI) fluorescence in Figure 3F. A Keyence BZ-X810 microscope with a 20x objective was
used to detect bright field images and GFP fluorescence in Figure 3B, Figure S1B right
panel. A Leica TCS SP5 confocal microscope with 40x objective was used to detect GFP
fluorescence in Figure 7A.
797
Accumulation of EIN2-GFP in root cells
3-day-old etiolated seedlings of two independent transgenic *EIN2-GFP* reporter lines in
which the GFP was fused to EIN2's C-terminus (*pEIN2::EIN2-GFP* in the *ein2-5*

background (Fu et al., 2021)) were moved into different wells of 6-well plate with 5 mL
Mock (dH2O+ 5 μL DMSO), ACC (100 μM + 5 μL DMSO), or MBZ (10 μM). The plate
was then covered with tin foil to keep the dark environment for 3 hour. Then samples
were directly mounted on glass slides in their treatment liquid and observed under a Leica
SP5 microscope (Leica).

Time lapse microscopy of reporter genes in lateral root tips
To observe the response of ethylene (*EIN3-GFP*), auxin (*DR5v2-GFP*), and cytokinin
(*TCSn-GFP*) reporter lines to MBZ, 14-day-old seedlings grown on ½ MS media were
transferred to a Chambered Coverglass (Thermo Scientific) that was filled with treatment
medium. For chamber preparation, 6 mL of ½ MS medium (1 % agar) supplemented with
the different treatments (DMSO, 10 μM MBZ, 1 μM IAA, 50 μM ACC, and 10 μM tZ) was
filled to each chamber. After 20 minutes of medium solidification, the seedlings were
placed on the media surface and the lateral roots were adjusted to positions suitable for
imaging.
For imaging, a Zeiss CSU Spinning Disk Confocal Microscope equipped with laser 488
mm and a Yokogawa spinning disk scan head with EM-CCD camera were used to set up
the image acquisition every 20 minutes for a total of 4 hours. Three Z-stack positions were
automatically captured at each time point under 10X objective and Fiji was used to
quantify the average intensity of GFP fluorescence. For proper comparison at different
treatment conditions of each reporter lines, we normalized the average fluorescence data
as the percentages (or relative fluorescence intensity as text label). In detail, the first time
point of each treatment is set as 100% and the relative fluorescence intensity from the
second time point is calculated as the fold-change in percentages comparing the first time
point.

Quantification of relative GFP fluorescent intensity between the nucleus and
cytosol
The quantification of the microscopy localization data of *EIN2-GFP* in Figure 7B was done
according to (Fu et al., 2021) using LAS X (Leica Application Suite v. X3.1.1.15751) and
GraphPad Prism V9.5.1. In brief, the signal intensity of nucleus (Ni) and whole cell (Wi)

was quantified by LAS X (Mean value of nucleus * ROI Area of nucleus, mean value of
whole cell * ROI Area of whole cell), and then the signal intensity of cytosol (Ci) was
obtained by Wi minus Ni, and the area of cytosol (Sc) was obtained by the area of whole
cell (ROI Area of whole cell) minus the area of nucleus (ROI Area of nucleus). The mean
value of the signal intensity of cytosol (Mean Ci) equals Ci/Sc, and the relative GFP
fluorescent intensity between the nucleus and cytosol equals Mean value of nucleus/
Mean Ci.
839

840 **Computational Docking and Molecular Modeling**
841 The AutoDock Vina and Chimera software were used to dock mebendazole
842 (CHEMBL685) into crystal structure of CTR1 (PDB: 3PPZ) and MAPK14 (PDB: 6SFO).
843 Figures for those structures were visualized and generated by Chimera and Pymol.
844

Protein purification
The CTR1 Kinase domain (540 aa-821 aa) (CTR1-KD) and MAPK4 were cloned into a
pET28a. Proteins were expressed in Rosetta™ 2(DE3)pLysS Competent Cells. Cultures
were incubated at 37°C (OD 600:0.7 to 0.9) and induced by adding 0.4 mM
isopropylthioβ-D-galactoside (IPTG) at 20°C for 22 h, 200 rpm. Cells were harvested and
resuspended in lysis buffer (50 mM pH 7.5 Tris-HCl, 1mM EDTA, 100 mM NaCl, 20%
glycerol, EDTA-free protease inhibitor cocktail, 10 mM imidazole). followed by sonication
and centrifugation (13,000 × g × 20 min). The His-tagged proteins were first purified using
QIAGEN Ni-NTA Agarose. The elute from Ni-NTA was collected and further purified by
FPLC using superdex 200 Increase 10/300 column (Mayerhofer et al., 2011). The
fractions corresponding to single peak from FPLC were combined, and concentrated to a
desired volume. Protein purity was verified by Coomassie blue gel staining. Protein
concentration was measured by their absorbance at 280 nm.
858

In vitro **kinase assay with non-radioactive ATP-γ-S**
The *in vitro* kinase assay followed the previous published method (Allen et al., 2007). 100
ng purified recombinant CTR1-KD or 1 μg MPK4 was mixed with 2 μg Myelin Basic
Protein (MBP, a common kinase substrate, Sigma Cat. M1891) and 1 mM ATPγS in

kinase reaction buffer (10 mM HEPES pH 7.4, 150 mM NaCl, 10 mM MgCl2, 1xRoche
Complete Protease Inhibitor mixture) and different concentration of MBZ at room
temperature and vortexed for 1 hour. The reactions were terminated by 20 mM EDTA,
and the proteins were alkylated by 1.5 μL of 50 mM PNBM (CAYMAN CHEMICAL CO,
Cat. 21456-1) at room temperature and vortexed for 2 hours. This step was terminated
by denaturing in NuPAGE™ LDS Sample Buffer and 1x NuPAGE™ Sample Reducing
Agent (Invitrogen™). Samples were subjected to SDS-PAGE. The phosphorylation status
of MBP, CTR1-KD and MPK4 were analyzed by western blot using Recombinant Anti-
Thiophosphate ester antibody (Abcam, Cat. Ab92570, 1:5000).

Statistical analysis

A two-sided Student's t-test was used for comparisons of two conditions. The comparison
of meristem length, mature cell length, root length, GSA between multiple treatments or
genotypes was performed by One-way ANOVA and *post hoc* Tukey testing. Detailed
information was prepared in figure legends. Statistical significance of overlap between
DEGs from MBZ treatment and ethylene treatment was assessed by hypergeometric
distribution test.

Data availability

Data that support the findings of this study have been deposited in GEO. The GEO
number is GSE201648 .

References
Alexa, A., Rahnenfuhrer, J., and Lengauer, T. (2006). Improved scoring of functional
groups from gene expression data by decorrelating GO graph structure. Bioinformatics
22, 1600-1607.
Allen, J.J., Li, M., Brinkworth, C.S., Paulson, J.L., Wang, D., Hubner, A., Chou, W.H.,
Davis, R.J., Burlingame, A.L., Messing, R.O., et al. (2007). A semisynthetic epitope for
kinase substrates. Nat Methods 4, 511-516.
Alonso, J.M., Hirayama, T., Roman, G., Nourizadeh, S., and Ecker, J.R. (1999). EIN2, a
bifunctional transducer of ethylene and stress responses in Arabidopsis. Science 284,
2148-2152.
Alonso, J.M., Stepanova, A.N., Solano, R., Wisman, E., Ferrari, S., Ausubel, F.M., and
Ecker, J.R. (2003). Five components of the ethylene-response pathway identified in a
screen for weak ethylene-insensitive mutants in Arabidopsis. Proc Natl Acad Sci U S A
100, 2992-2997.

An, F., Zhao, Q., Ji, Y., Li, W., Jiang, Z., Yu, X., Zhang, C., Han, Y., He, W., Liu, Y., et
al. (2010). Ethylene-induced stabilization of ETHYLENE INSENSITIVE3 and EIN3-
LIKE1 is mediated by proteasomal degradation of EIN3 binding F-box 1 and 2 that
requires EIN2 in Arabidopsis. Plant Cell 22, 2384-2401.
Ariey-Bonnet, J., Carrasco, K., Le Grand, M., Hoffer, L., Betzi, S., Feracci, M., Tsvetkov,
P., Devred, F., Collette, Y., Morelli, X., et al. (2020). In silico molecular target prediction
unveils mebendazole as a potent MAPK14 inhibitor. Mol Oncol 14, 3083-3099.
Babicki, S., Arndt, D., Marcu, A., Liang, Y.J., Grant, J.R., Maciejewski, A., and Wishart,
D.S. (2016). Heatmapper: web-enabled heat mapping for all. Nucleic Acids Res 44,
W147-W153.
Berardini, T.Z., Reiser, L., Li, D.H., Mezheritsky, Y., Muller, R., Strait, E., and Huala, E.
(2015). The arabidopsis information resource: Making and mining the "gold standard"
annotated reference plant genome. Genesis 53, 474-485.
Chang, C., Kwok, S.F., Bleecker, A.B., and Meyerowitz, E.M. (1993). Arabidopsis
ethylene-response gene ETR1: similarity of product to two-component regulators.
Science 262, 539-544.
Chang, K.N., Zhong, S., Weirauch, M.T., Hon, G., Pelizzola, M., Li, H., Huang, S.S.C.,
Schmitz, R.J., Urich, M.A., Kuo, D., et al. (2013). Temporal transcriptional response to
ethylene gas drives growth hormone cross-regulation in Arabidopsis. Elife 2.
Chao, Q., Rothenberg, M., Solano, R., Roman, G., Terzaghi, W., and Ecker, J.R.
(1997). Activation of the ethylene gas response pathway in Arabidopsis by the nuclear
protein ETHYLENE-INSENSITIVE3 and related proteins. Cell 89, 1133-1144.
De Cnodder, T., Vissenberg, K., Van Der Straeten, D., and Verbelen, J.P. (2005).
Regulation of cell length in the Arabidopsis thaliana root by the ethylene precursor 1-
aminocyclopropane- 1-carboxylic acid: a matter of apoplastic reactions. The New
phytologist 168, 541-550.
Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner.
Bioinformatics 29, 15-21.
Feng, Y., Xu, P., Li, B., Li, P., Wen, X., An, F., Gong, Y., Xin, Y., Zhu, Z., Wang, Y., et
al. (2017). Ethylene promotes root hair growth through coordinated EIN3/EIL1 and
RHD6/RSL1 activity in Arabidopsis. Proc Natl Acad Sci U S A 114, 13834-13839.
Fu, L.W., Liu, Y.L., Qin, G.C., Wu, P., Zi, H.L., Xu, Z.T., Zhao, X.D., Wang, Y., Li, Y.X.,
Yang, S.H., et al. (2021). The TOR-EIN2 axis mediates nuclear signalling to modulate
plant growth. Nature 591, 288-292.
Furutani, M., and Morita, M.T. (2021). LAZY1-LIKE-mediated gravity signaling pathway
in root gravitropic set-point angle control. Plant Physiology 187, 1087-1095.
Ge, L.F., and Chen, R.J. (2016). Negative gravitropism in plant roots. Nature plants 2.
Guo, H., and Ecker, J.R. (2003). Plant responses to ethylene gas are mediated by
SCF(EBF1/EBF2)-dependent proteolysis of EIN3 transcription factor. Cell 115, 667-677.
Guseman, J.M., Webb, K., Srinivasan, C., and Dardick, C. (2017). DRO1 influences root
system architecture in Arabidopsis and Prunus species. Plant J 89, 1093-1105.
Guzman, P., and Ecker, J.R. (1990). Exploiting the triple response of Arabidopsis to
identify ethylene-related mutants. Plant Cell 2, 513-523.
He, W.R., Brumos, J., Li, H.J., Ji, Y.S., Ke, M., Gong, X.Q., Zeng, Q.L., Li, W.Y., Zhang,
X.Y., An, F.Y., et al. (2011). A Small-Molecule Screen Identifies L-Kynurenine as a

Competitive Inhibitor of TAA1/TAR Activity in Ethylene-Directed Auxin Biosynthesis and
Root Growth in Arabidopsis. Plant Cell 23, 3944-3960.
Hua, J., Sakai, H., Nourizadeh, S., Chen, Q.G., Bleecker, A.B., Ecker, J.R., and
Meyerowitz, E.M. (1998). EIN4 and ERS2 are members of the putative ethylene
receptor gene family in Arabidopsis. Plant Cell 10, 1321-1332.
Ju, C., Van de Poel, B., Cooper, E.D., Thierer, J.H., Gibbons, T.R., Delwiche, C.F., and
Chang, C. (2015). Conservation of ethylene as a plant hormone over 450 million years
of evolution. Nature plants 1, 14004.
Ju, C., Yoon, G.M., Shemansky, J.M., Lin, D.Y., Ying, Z.I., Chang, J., Garrett, W.M.,
Kessenbrock, M., Groth, G., Tucker, M.L., et al. (2012). CTR1 phosphorylates the
central regulator EIN2 to control ethylene hormone signaling from the ER membrane to
the nucleus in Arabidopsis. Proc Natl Acad Sci U S A 109, 19486-19491.
Kieber, J.J., Rothenberg, M., Roman, G., Feldmann, K.A., and Ecker, J.R. (1993).
CTR1, a negative regulator of the ethylene response pathway in Arabidopsis, encodes a
member of the raf family of protein kinases. Cell 72, 427-441.
Le, J., Vandenbussche, F., Van Der Straeten, D., and Verbelen, J.P. (2001). In the early
response of Arabidopsis roots to ethylene, cell elongation is up- and down-regulated
and uncoupled from differentiation. Plant Physiol 125, 519-522.
Li, W., Lacey, R.F., Ye, Y., Lu, J., Yeh, K.C., Xiao, Y., Li, L., Wen, C.K., Binder, B.M.,
and Zhao, Y. (2017). Triplin, a small molecule, reveals copper ion transport in ethylene
signaling from ATX1 to RAN1. PLoS Genet 13, e1006703.
Liao, C.Y., Smet, W., Brunoud, G., Yoshida, S., Vernoux, T., and Weijers, D. (2015a).
Reporters for sensitive and quantitative measurement of auxin response. Nat Methods
12, 207-210, 202 p following 210.
Liao, C.Y., Smet, W., Brunoud, G., Yoshida, S., Vernoux, T., and Weijers, D. (2015b).
Reporters for sensitive and quantitative measurement of auxin response (vol 12, pg
207, 2015). Nat Methods 12, 1098-1098.
Liao, C.Y., Smet, W., Brunoud, G., Yoshida, S., Vernoux, T., and Weijers, D. (2015).
Reporters for sensitive and quantitative measurement of auxin response. Nat Methods
12, 207-210, 202 p following 210.
Lynch, J.P. (2019). Root phenotypes for improved nutrient capture: an underexploited
opportunity for global agriculture. New Phytol 223, 548-564.
Mayerhofer, H., Mueller-Dieckmann, C., and Mueller-Dieckmann, J. (2011). Cloning,
expression, purification and preliminary X-ray analysis of the protein kinase domain of
constitutive triple response 1 (CTR1) from Arabidopsis thaliana. Acta Crystallogr Sect F
Struct Biol Cryst Commun 67, 117-120.
Mayerhofer, H., Panneerselvam, S., and Mueller-Dieckmann, J. (2012). Protein kinase
domain of CTR1 from Arabidopsis thaliana promotes ethylene receptor cross talk. J Mol
Biol 415, 768-779.
Mou, W., Kao, Y.T., Michard, E., Simon, A.A., Li, D., Wudick, M.M., Lizzio, M.A., Feijo,
J.A., and Chang, C. (2020). Ethylene-independent signaling by the ethylene precursor
ACC in Arabidopsis ovular pollen tube attraction. Nat Commun 11, 4082.
Muller, M., and Munne-Bosch, S. (2015). Ethylene Response Factors: A Key Regulatory
Hub in Hormone and Stress Signaling. Plant Physiol 169, 32-41.

Nakamura, M., Naoi, K., Shoji, T., and Hashimoto, T. (2004). Low concentrations of
propyzamide and oryzalin alter microtubule dynamics in Arabidopsis epidermal cells.
Plant Cell Physiol 45, 1330-1334.
Negi, S., Ivanchenko, M.G., and Muday, G.K. (2008). Ethylene regulates lateral root
formation and auxin transport in Arabidopsis thaliana. Plant J 55, 175-187.
Ogura, T., Goeschl, C., and Busch, W. (2022). A Multiplexed, Time-Resolved Assay of
Root Gravitropic Bending on Agar Plates. Methods Mol Biol 2368, 61-70.
Pattyn, J., Vaughan-Hirsch, J., and Van de Poel, B. (2021). The regulation of ethylene
biosynthesis: a complex multilevel control circuitry. New Phytol 229, 770-782.
Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng,
E.C., and Ferrin, T.E. (2004). UCSF Chimera--a visualization system for exploratory
research and analysis. J Comput Chem 25, 1605-1612.
Poirier, V., Roumet, C., and Munson, A.D. (2018). The root of the matter: Linking root
traits and soil organic matter stabilization processes. Soil Biology and Biochemistry 120,
246-259.
Qiao, H., Shen, Z., Huang, S.S., Schmitz, R.J., Urich, M.A., Briggs, S.P., and Ecker,
J.R. (2012a). Processing and subcellular trafficking of ER-tethered EIN2 control
response to ethylene gas. Science 338, 390-393.
Qiao, H., Shen, Z.X., Huang, S.S.C., Schmitz, R.J., Urich, M.A., Briggs, S.P., and
Ecker, J.R. (2012b). Processing and Subcellular Trafficking of ER-Tethered EIN2
Control Response to Ethylene Gas. Science 338, 390-393.
Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR: a Bioconductor
package for differential expression analysis of digital gene expression data.
Bioinformatics 26, 139-140.
Rosquete, M.R., von Wangenheim, D., Marhavy, P., Barbez, E., Stelzer, E.H., Benkova,
E., Maizel, A., and Kleine-Vehn, J. (2013a). An auxin transport mechanism restricts
positive orthogravitropism in lateral roots. Current biology : CB 23, 817-822.
Rosquete, M.R., von Wangenheim, D., Marhavy, P., Barbez, E., Stelzer, E.H.K.,
Benkova, E., Maizel, A., and Kleine-Vehn, J. (2013b). An Auxin Transport Mechanism
Restricts Positive Orthogravitropism in Lateral Roots. Curr Biol 23, 817-822.
Rosquete, M.R., von Wangenheim, D., Marhavy, P., Barbez, E., Stelzer, E.H., Benkova,
E., Maizel, A., and Kleine-Vehn, J. (2013). An auxin transport mechanism restricts
positive orthogravitropism in lateral roots. Current biology : CB 23, 817-822.
Roychoudhry, S., Del Bianco, M., Kieffer, M., and Kepinski, S. (2013). Auxin controls
gravitropic setpoint angle in higher plant lateral branches. Current biology : CB 23,
1497-1504.
S F Yang, a., and Hoffman, N.E. (1984). Ethylene Biosynthesis and its Regulation in
Higher Plants. Annual Review of Plant Physiology 35, 155-189.
Sakai, H., Hua, J., Chen, Q.G., Chang, C., Medrano, L.J., Bleecker, A.B., and
Meyerowitz, E.M. (1998). ETR2 is an ETR1-like gene involved in ethylene signaling in
Arabidopsis. Proc Natl Acad Sci U S A 95, 5812-5817.
Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source
platform for biological-image analysis. Nat Methods 9, 676-682.

Sun, X.Z., Li, Y.X., He, W.R., Ji, C.G., Xia, P.X., Wang, Y.C., Du, S., Li, H.J., Raikhel, N., Xiao, J.Y., et al. (2017). Pyrazinamide and derivatives block ethylene biosynthesis by inhibiting ACC oxidase. Nat Commun 8.

Trott, O., and Olson, A.J. (2010). AutoDock Vina: improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading. J Comput Chem 31, 455-461.

Uga, Y., Sugimoto, K., Ogawa, S., Rane, J., Ishitani, M., Hara, N., Kitomi, Y., Inukai, Y., Ono, K., Kanno, N., et al. (2013). Control of root system architecture by DEEPER ROOTING 1 increases rice yield under drought conditions. Nature genetics 45, 1097-1102.

Vaseva, II, Qudeimat, E., Potuschak, T., Du, Y., Genschik, P., Vandenbussche, F., and Van Der Straeten, D. (2018). The plant hormone ethylene restricts Arabidopsis growth via the epidermis. Proc Natl Acad Sci U S A 115, E4130-E4139.

Waidmann, S., Rosquete, M.R., Scholler, M., Sarkel, E., Lindner, H., LaRue, T., Petrik, I., Dunser, K., Martopawiro, S., Sasidharan, R., et al. (2019a). Cytokinin functions as an asymmetric and anti-gravitropic signal in lateral roots. Nat Commun 10.

Waidmann, S., Ruiz Rosquete, M., Scholler, M., Sarkel, E., Lindner, H., LaRue, T., Petrik, I., Dunser, K., Martopawiro, S., Sasidharan, R., et al. (2019b). Cytokinin functions as an asymmetric and anti-gravitropic signal in lateral roots. Nat Commun 10, 3540.

Waidmann, S., Ruiz Rosquete, M., Scholler, M., Sarkel, E., Lindner, H., LaRue, T., Petrik, I., Dunser, K., Martopawiro, S., Sasidharan, R., et al. (2019). Cytokinin functions as an asymmetric and anti-gravitropic signal in lateral roots. Nat Commun 10, 3540.

Waite, J.M., and Dardick, C. (2021). The roles of the IGT gene family in plant architecture: past, present, and future. Curr Opin Plant Biol 59.

Wen, X., Zhang, C., Ji, Y., Zhao, Q., He, W., An, F., Jiang, L., and Guo, H. (2012). Activation of ethylene signaling is mediated by nuclear translocation of the cleaved EIN2 carboxyl terminus. Cell Res 22, 1613-1616.

Xiao, G., and Zhang, Y. (2020). Adaptive Growth: Shaping Auxin-Mediated Root System Architecture. Trends in plant science 25, 121-123.

Yoshihara, T., and Spalding, E.P. (2017). LAZY Genes Mediate the Effects of Gravity on Auxin Gradients and Plant Architecture. Plant Physiology 175, 959-969.

Zhu, Y., Li, H.J., Su, Q., Wen, J., Wang, Y.F., Song, W., Xie, Y.P., He, W.R., Yang, Z., Jiang, K., et al. (2019). A phenotype-directed chemical screen identifies ponalrestat as an inhibitor of the plant flavin monooxygenase YUCCA in auxin biosynthesis. J Biol Chem 294, 19923-19933.

Zurcher, E., Tavor-Deslex, D., Lituiev, D., Enkerli, K., Tarr, P.T., and Muller, B. (2013). A robust and sensitive synthetic sensor to monitor the transcriptional output of the cytokinin signaling network in planta. Plant Physiol 161, 1066-1075.

Figure 1. MBZ treatment perturbs root system architecture (RSA).

A

GO biological process

negative regulation of ethylene-activated signaling pathway (GO:0010105) — P = 6.21 X 10⁻³
response to far red light (GO:0010218) — P = 1.73 X 10⁻⁷
response to nitrate (GO:0010167) — P = 1.17 X 10⁻²
response to red light (GO:0010114) — P = 2.10 X 10⁻⁷
ethylene-activated signaling pathway (GO:0009873) — P = 2.42 X 10⁻³
response to UV-B (GO:0010224) — P = 1.16 X 10⁻²
response to blue light (GO:0009637) — P = 1.52 X 10⁻²
response to hypoxia (GO:0001666) — P = 2.39 X 10⁻⁷
cellular response to extracellular stimulus (GO:0031668) — P = 8.54 X 10⁻³
response to jasmonic acid (GO:0009753) — P = 2.86 X 10⁻³
response to oxidative stress (GO:0006979) — P = 4.65 X 10⁻³
response to wounding (GO:0009611) — P = 1.29 X 10⁻³
secondary metabolic process (GO:0019748) — P = 1.87 X 10⁻²
response to organic cyclic compound (GO:0014070) — P = 9.93 X 10⁻³
defense response to fungus (GO:0050832) — P = 1.92 X 10⁻²
defense response to bacterium (GO:0042742) — P = 3.79 X 10⁻³
regulation of DNA-templated transcription (GO:0006355) — P = 1.48 X 10⁻⁶
response to abscisic acid (GO:0009737) — P = 1.28 X 10⁻²
macromolecule biosynthetic process (GO:0009059) — P = 4.35 X 10⁻²
RNA metabolic process (GO:0016070) — P = 1.61 X 10⁻⁴
gene expression (GO:0010467) — P = 1.93 X 10⁻⁵
protein localization (GO:0008104) — P = 3.99 X 10⁻²
intracellular transport (GO:0046907) — P = 3.79 X 10⁻²
amide metabolic process (GO:0043603) — P = 4.07 X 10⁻²

Fold Enrichment   0   5   10   15   20

Figure 2. MBZ treatment regulates the ethylene pathway.

Figure 3. MBZ induces ethylene response and mimics ACC treatment

Figure 4. MBZ acts downstream of ethylene biosynthesis

Figure 5. MBZ targets the ethylene signaling pathway

Figure 6. MBZ inhibits CTR1 kinase activity

Figure 7. MBZ activates ethylene signaling

A

Mebendazole (MBZ)

B

DMSO   1 µM MBZ

C

DMSO   1 µM MBZ

Figure S1. MBZ treatment affects root growth and development

Figure S2. The analysis of differentially expressed genes (DEGs) by MBZ treatment.

Figure S3. Etiolated seedling phenotypes in mutants of the ethylene pathway.

Figure S4. Alignment of *At*CTR1-KD and *hs*MAPK14

Figure S5. High concentrations or long-term treatment of MBZ induce phenotypes that go beyond ethylene induced effects

Table S1. The list of blast hits in Arabidopsis using the MAPK14_human sequence as BLASTP

JVD9H54701R-Alignment-Descriptions

Description	Scientific Name	Max Score	Total Score	Query Cover	E value	Per. ident	Acc. Len	Accession
MAP kinase 4 [Arabidopsis thaliana]	Arabidopsis thaliana	346	346	93%	3E-117	51.18	376	NP_192046.1
mitogen-activated protein kinase 4 [Arabidopsis thaliana]	Arabidopsis thaliana	346	346	93%	3E-117	51.18	376	ABR46146.1
mitogen-activated protein kinase 4 [Arabidopsis thaliana]	Arabidopsis thaliana	345	345	93%	8E-117	50.88	376	ABR46148.1
MAP kinase [Arabidopsis thaliana]	Arabidopsis thaliana	345	345	93%	9E-117	50.88	376	BAA04867.1
mitogen-activated protein kinase 4 [Arabidopsis thaliana]	Arabidopsis thaliana	344	344	93%	1E-116	50.88	376	ABR46162.1
unnamed protein product [Arabidopsis thaliana]	Arabidopsis thaliana	340	340	95%	8E-115	47.71	376	VYS62299.1
MPK5 [Arabidopsis thaliana]	Arabidopsis thaliana	339	339	95%	9E-115	47.71	376	OAO98154.1
MAP kinase 11 [Arabidopsis thaliana]	Arabidopsis thaliana	337	337	93%	6E-114	49.71	369	NP_001117210.1
unnamed protein product [Arabidopsis thaliana]	Arabidopsis thaliana	337	337	93%	6E-114	49.71	369	CAA0154293.1
MAP kinase 5 [Arabidopsis thaliana]	Arabidopsis thaliana	337	337	95%	8E-114	47.43	376	NP_567378.4
unnamed protein product [Arabidopsis thaliana]	Arabidopsis thaliana	336	336	95%	2E-113	46.84	370	CAD5324945.1
MAP kinase [Arabidopsis thaliana]	Arabidopsis thaliana	335	335	95%	2E-113	46.55	370	BAA04866.1
mitogen-activated protein kinase 3 [Arabidopsis thaliana]	Arabidopsis thaliana	335	335	95%	2E-113	46.55	370	NP_190150.1
MAP kinase [Arabidopsis thaliana]	Arabidopsis thaliana	330	330	95%	3E-111	46.57	376	BAA04868.1
mitogen-activated protein kinase 12 [Arabidopsis thaliana]	Arabidopsis thaliana	329	329	92%	9E-111	49.26	372	NP_182131.2
mitogen-activated protein kinase 12 [Arabidopsis thaliana]	Arabidopsis thaliana	330	330	92%	2E-110	49.26	406	NP_001325006.1
Mitogen-activated protein kinase 6 [Arabidopsis thaliana]	Arabidopsis thaliana	323	323	95%	2E-108	46.84	364	6DTL_A
Crystal structure of Arabidopsis thaliana MPK6 [Arabidopsis thaliana]	Arabidopsis thaliana	323	323	95%	3E-108	46.84	370	5CI6_A
unnamed protein product [Arabidopsis thaliana]	Arabidopsis thaliana	323	323	92%	3E-108	46.31	393	VYS60924.1
MAP kinase 6 [Arabidopsis thaliana]	Arabidopsis thaliana	323	323	95%	4E-108	46.84	395	NP_181907.1
MAP kinase 10 [Arabidopsis thaliana]	Arabidopsis thaliana	323	323	92%	4E-108	46.31	393	NP_191538.1
unnamed protein product [Arabidopsis thaliana]	Arabidopsis thaliana	323	323	92%	5E-108	46.31	393	CAA0387508.1
MPK10 [Arabidopsis thaliana]	Arabidopsis thaliana	323	323	92%	6E-108	46.31	393	OAP05435.1
unnamed protein product [Arabidopsis thaliana]	Arabidopsis thaliana	321	321	92%	2E-107	46.02	393	CAD5326291.1
MAP kinase [Arabidopsis thaliana]	Arabidopsis thaliana	318	318	95%	2E-106	46.29	373	CAB51417.1
Protein kinase superfamily protein [Arabidopsis thaliana]	Arabidopsis thaliana	317	317	92%	4E-106	47.79	363	NP_001030990.1
unnamed protein product [Arabidopsis thaliana]	Arabidopsis thaliana	316	316	92%	8E-106	47.79	363	CAD5312051.1
mitogen-activated protein kinase homolog 2 [Arabidopsis thaliana]	Arabidopsis thaliana	316	316	85%	2E-105	49.68	376	NP_564746.1
ATMPK1 [Arabidopsis thaliana]	Arabidopsis thaliana	315	315	93%	2E-105	46.33	370	BAA03535.1
unnamed protein product [Arabidopsis thaliana]	Arabidopsis thaliana	315	315	93%	3E-105	46.33	370	CAA0186271.1
AT1G10210 [Arabidopsis thaliana]	Arabidopsis thaliana	315	315	93%	3E-105	46.33	370	BAH19911.1
mitogen-activated protein kinase 1 [Arabidopsis thaliana]	Arabidopsis thaliana	315	315	93%	3E-105	46.33	370	NP_001031017.1
mitogen-activated protein kinase 14 [Arabidopsis thaliana]	Arabidopsis thaliana	315	315	95%	3E-105	45.04	361	NP_195363.1
ATMPK2 [Arabidopsis thaliana]	Arabidopsis thaliana	315	315	85%	4E-105	49.35	376	BAA03536.1
unnamed protein product [Arabidopsis thaliana]	Arabidopsis thaliana	313	313	95%	1E-104	45.04	361	CAA0397686.1
MPK14 [Arabidopsis thaliana]	Arabidopsis thaliana	311	311	95%	8E-104	44.76	361	OAO98488.1
MAP kinase 5-CT [Arabidopsis thaliana]	Arabidopsis thaliana	310	310	81%	9E-104	50	337	ADZ76491.1
MAP kinase 7 [Arabidopsis thaliana]	Arabidopsis thaliana	310	310	85%	2E-103	48.71	368	NP_179409.1
MPK7 [Arabidopsis thaliana]	Arabidopsis thaliana	310	310	85%	2E-103	48.71	368	OAP11028.1
MAP Kinase [Arabidopsis thaliana]	Arabidopsis thaliana	309	309	86%	4E-103	50.48	354	AAB61033.1
unnamed protein product [Arabidopsis thaliana]	Arabidopsis thaliana	310	310	91%	4E-103	48.49	389	CAD5321516.1
unnamed protein product [Arabidopsis thaliana]	Arabidopsis thaliana	309	309	85%	5E-103	48.71	368	CAA0364931.1
MAP kinase 11 [Arabidopsis thaliana]	Arabidopsis thaliana	307	307	78%	2E-102	52.08	346	NP_001322112.1
MAP kinase [Arabidopsis thaliana]	Arabidopsis thaliana	306	306	85%	1E-101	48.06	368	BAA04870.1
unnamed protein product [Arabidopsis thaliana]	Arabidopsis thaliana	310	310	86%	3E-99	48.12	716	CAD5330118.1
mitogen-activated protein kinase 12 [Arabidopsis thaliana]	Arabidopsis thaliana	297	297	80%	5E-99	51.03	324	NP_001189755.1
mitogen-activated protein kinase 16 [Arabidopsis thaliana]	Arabidopsis thaliana	287	287	91%	7E-92	42.77	567	NP_197402.1
Unknown protein [Arabidopsis thaliana]	Arabidopsis thaliana	287	287	91%	7E-92	42.77	567	AAL32607.1
unnamed protein product [Arabidopsis thaliana]	Arabidopsis thaliana	287	287	91%	9E-92	42.77	569	CAD5332107.1
MAP kinase [Arabidopsis thaliana]	Arabidopsis thaliana	276	276	75%	6E-90	50	360	BAD 95376.10
putative MAP kinase [Arabidopsis thaliana]	Arabidopsis thaliana	271	271	90%	4E-87	42.6	431	AAK62464.1
MPK15 [Arabidopsis thaliana]	Arabidopsis thaliana	274	274	92%	1E-86	41.86	575	OAP17567.1
MAP kinase 19 [Arabidopsis thaliana]	Arabidopsis thaliana	268	268	91%	3E-86	42.18	425	NP_001326968.1
MAP kinase 15 [Arabidopsis thaliana]	Arabidopsis thaliana	272	272	92%	4E-86	41.86	576	NP_565070.2
MPK8 [Arabidopsis thaliana]	Arabidopsis thaliana	269	269	92%	1E-84	42.15	588	OAP18797.1