Retrotransposon instability dominates the acquired mutation landscape of mouse induced pluripotent stem cells

Patricia Gerdes[1,10], Sue Mei Lim[2,3,4,10], Adam D. Ewing[1,10], Michael R. Larcombe[2,3,4], Dorothy Chan[1], Francisco J. Sanchez-Luque[1,5], Lucinda Walker[1], Alexander L. Carleton[1], Cini James[1], Anja S. Knaupp[2,3,4], Patricia E. Carreira[1], Christian M. Nefzger[2,3,4], Ryan Lister[6,7], Sandra R. Richardson[1,*], Jose M. Polo[2,3,4,9,*], and Geoffrey J. Faulkner[1,8,*]

[1]Mater Research Institute - University of Queensland, TRI Building, Woolloongabba QLD 4102, Australia.

[2]Department of Anatomy & Developmental Biology, Monash University, Melbourne VIC 3800, Australia.

[3]Development and Stem Cells Program, Monash Biomedicine Discovery Institute, Melbourne VIC 3800, Australia.

[4]Australian Regenerative Medicine Institute, Monash University, Melbourne VIC 3800, Australia.

[5]GENYO. Pfizer-University of Granada-Andalusian Government Centre for Genomics and Oncological Research, PTS Granada 18016, Spain.

[6]Australian Research Council Centre of Excellence in Plant Energy Biology, School of Molecular Sciences, The University of Western Australia, Perth WA 6009, Australia.

[7]Harry Perkins Institute of Medical Research, Perth WA 6009, Australia.

[8]Queensland Brain Institute, University of Queensland, Brisbane QLD 4072, Australia.

[9]Adelaide Centre for Epigenetics and The South Australian Immunogenomics Cancer Institute, Faculty of Health and Medical Sciences, The University of Adelaide, Adelaide SA 5005, Australia.

[10]These authors contributed equally.

[*]Corresponding authors: sandra.richardson@mater.uq.edu.au (S.R.R.); jose.polo@monash.edu (J.M.P.); faulknergj@gmail.com (G.J.F.)

Abstract

Induced pluripotent stem cells (iPSCs) can in principle differentiate into any cell of the body, and have revolutionized biomedical research and regenerative medicine. Unlike their human counterparts, mouse iPSCs (miPSCs) are reported to silence transposable elements (TEs) and prevent TE-mediated mutagenesis. Here we applied short-read or Oxford Nanopore Technologies (ONT) long-read genome sequencing to 38 bulk miPSC lines reprogrammed from 10 parental cell types, and 18 single-cell miPSC clones. While single nucleotide variants and structural variants restricted to miPSCs were rare, we found 83 *de novo* TE insertions, including examples intronic to *Brca1* and *Dmd*. LINE-1 (L1) retrotransposons were profoundly hypomethylated in miPSCs, beyond other TEs and the genome overall, and harbored alternative protein-coding gene promoters. Treatment with the L1 inhibitor lamivudine did not hinder reprogramming and efficiently blocked endogenous retrotransposition, as detected by ONT sequencing. These experiments reveal the complete spectrum and potential significance of mutations acquired by miPSCs.

Introduction

Induced pluripotent stem cells (iPSCs) resemble embryonic stem cells (ESCs) in their near unlimited capacity for self-renewal and differentiation potential[1]. These properties have driven widespread uptake of iPSCs in clinical and research applications[2–4]. Despite their immense therapeutic promise, the reprogramming process required to generate iPSCs can produce genomic and epigenomic aberrations[4–8]. These abnormalities could undermine the functional equivalence of iPSCs and ESCs, or alter the phenotype of iPSC-derived differentiated cells, and hence necessitate genetic and functional screening of iPSCs prior to their use in the clinic[9]. Fortunately, whole genome sequencing (WGS) analyses of single nucleotide variants (SNVs), copy number variants, and structural variants (SVs) restricted to human and mouse iPSC lines have found relatively few conclusive reprogramming-associated mutations[10–12]. Instead, most mutations acquired by iPSCs appear to occur before and after reprogramming[10,11,13], implying they are not caused by molecular processes intrinsic to iPSC generation. Transposable elements (TEs) may present an important exception to this rule, where the attainment of a pluripotent state via reprogramming leaves iPSCs vulnerable to TE-mediated mutagenesis.

The retrotransposon long interspersed element 1 (LINE-1, or L1) is active in nearly all mammals[14]. L1 autonomously mobilizes via a copy-and-paste process called target-primed reverse transcription (TPRT), which involves reverse transcription of L1 mRNA *in cis*, and is characterized by the generation of target site duplications (TSDs) upon L1 integration[15–20]. The C57BL/6 mouse reference genome contains ~3,000 potentially mobile L1 copies belonging to three subfamilies (T_F, G_F and A) defined by their monomeric 5′ promoter sequences, in addition to several active endogenous retrovirus (ERV) and short interspersed element (SINE) families[21–23]. By contrast, only ~100 mobile L1s from the transcribed subset Ta (-Ta)[24] subfamily are present in each individual human genome, with the vast majority of retrotransposition potential concentrated in fewer than 10 of these elements[25,26]. Perhaps owing to the disparate count of mobile TEs in each species, the rate of L1 mobilization in the mouse germline is estimated to be at least an order of magnitude higher than that of humans[27–30].

TE mobility is regulated by DNA methylation and histone modifications, as well as various post-transcriptional and post-translational mechanisms[31–41]. Reprogramming somatic cells to generate human iPSCs (hiPSCs) and mouse iPSCs (miPSCs) leads to epigenome-wide remodeling, including broad de-repression of L1 promoters[7,42–47]. L1 mRNA abundance increases strongly during reprogramming, and remains approximately 10-fold higher in cultured miPSCs than in parental mouse embryonic fibroblasts (MEFs)[46]. As a corollary, the early mouse embryo is a major niche for new heritable L1 retrotransposition events[28]. Mouse ESCs cultured in standard media containing serum, or naïve ground state media incorporating two small-molecule kinase inhibitors (2i), permit genome-wide L1 hypomethylation, express endogenous L1 proteins and support engineered L1 mobilization[37,41,48,49]. Engineered and endogenous L1 retrotransposition are supported by hiPSCs and human ESCs[45,50–52]. These observations collectively suggest L1 hypomethylation may be an inherent aspect of pluripotency accentuated by the molecular roadmap to an induced pluripotent state. Consequently, miPSCs are likely to harbor *de novo* retrotransposition events. However, a prior WGS analysis of 3 miPSC lines, employing paired-end 42mer reads and ~11× genome-wide sequencing depth, found no *de novo* TE insertions, and concluded that endogenous retrotransposition did not occur during miPSC production[12]. The apparent lack of TE mobility in this context remains an unresolved and yet potentially important source of miPSC mutagenesis[4]. Here, we analyze a diverse panel of miPSC genomes with short-

or long-read sequencing and, as reported for hiPSCs and other pluripotent human cells[45,50–52], we

detect numerous *de novo* TE insertions acquired by miPSCs.

Results

Mutational spectra of bulk miPSC populations generated from diverse cell lineages

To survey genomic variation among miPSC lines generated from a broad range of parental cell

types, we bred triple transgenic C57BL/6×129S4Sv/Jae animals carrying a GFP reporter knocked

into the *Oct4* locus (*Oct4*-GFP), a transcriptional activator (m2rtTA) under the control of the

ubiquitously expressed Rosa26 locus (R26-m2rtTA), and a doxycycline-inducible polycistronic

reprogramming cassette (Col1a1-tetO-OKSM)[53]. From each of three animals (labeled A67, A82

and A172), we used fluorescence activated cell sorting (FACS) and a range of surface markers to

isolate nine isogenic primary cell populations, including three representing each germ layer (**Fig.**

1a). Bulk cultures were then treated with doxycycline to induce reprogramming, followed by

FACS to purify *Oct4*-GFP⁺ miPSCs. Twenty-six miPSC lines were expanded and cultured in

standard media containing serum (**Supplementary Table 1** and **Extended Data Fig. 1**). Illumina

paired-end 150mer read WGS (~41× average genome-wide depth) was then applied to each miPSC

line at passage 4 (p4), as well as to 3 MEF genotypic controls (**Supplementary Table 1**).

Concordant SNVs detected by GATK HaplotypeCaller and freebayes[54,55] were filtered to

remove known mouse strain germline variants[56], yielding 3,603 SNVs private to a single miPSC

line (average ~140 per line) and absent from the corresponding MEF samples (**Supplementary**

Table 2). Of these, 27 in total were non-synonymous exonic mutations. We then called concordant

SVs using Delly and GRIDSS[57,58], finding 34 private SVs (~1 per line). These included a 210kbp

deletion of the *de novo* methyltransferase *Dnmt3a* in miPSCs derived from the hematopoietic stem

cells of animal A172 (**Supplementary Table 2**). Considering private SNVs and SVs together, we

observed no significant (p<0.05, one-way ANOVA with Tukey's multiple comparison test)

difference in miPSC variant counts associated with parental cell type or germ layer, and SNV and

SV rates resembled those found previously for fibroblast-derived miPSCs[10,12]. Choice of primary

cell type, at least among the diverse panel assembled here, may therefore not significantly impact

the frequency of SNVs and SVs later found in miPSC lines.

Fig. 1: *De novo* **L1 insertions in germ layer specific bulk expanded miPSC lines. a,** Experimental design of bulk miPSC generation using a Col1a1-tetO-OKSM mouse model containing a doxycycline-inducible reprogramming cassette. Tissues were isolated and sorted by FACS to obtain 9 primary cell types (named and numbered 1-9) from each of 3 mice (A67, A82, A172). After reprogramming with doxycycline (+DOX), *Oct4*-GFP positive miPSCs were sorted and expanded in cell culture. DNA was extracted from miPSCs, sequenced via WGS and mRC-seq, and analyzed for *de novo* TE insertions with TEBreak. The histogram displayed underneath indicates the mean reprogramming efficiency (*Oct4*-GFP+) ± SD for each primary cell type, with individual replicate values marked by black dots. Red numerals indicate the count of reprogramming-associated L1 insertions found in the miPSCs obtained from each primary cell type. Note: astrocyte-derived miPSCs were not produced for animal A172. **b,** A full-length (6.8 monomers) intergenic *de novo* L1 T_F insertion. Promoter monomers are shown as triangles within the L1 5′UTR. PolyA (A_n) tract length is indicated immediately 3′ of the L1. Target site duplications (TSDs) are depicted as gray arrows flanking the L1. PCR validation primers are shown as red arrows. A PCR validation agarose gel containing the full-length PCR product (red arrow) only in the fibroblast-derived miPSC line where the L1 was detected by genomic analysis is shown. miPSC line numbers are provided in panel (a). Note: given the variable, and in some cases very low, reprogramming efficiencies shown in panel (a), and the objective to obtain a full set of 9 miPSC lines from each animal, we entirely used each sorted primary cell population for reprogramming, relying on PCR amplification in a single miPSC line to validate reprogramming-associated retrotransposition events. DNA from other animals included in the study are however shown at right as additional controls. **c,** As for panel (b), except for an L1 T_F with 5.8 promoter monomers. Note: the very faint, smaller-sized gel band observed in most of the samples was an off-target product. **d,** As for panel (b), except for an L1 T_F with 5.3 promoter monomers, and using an empty/filled PCR design where both primers are outside of the L1 insertion, generating "filled" L1 (red arrow) and "empty" wild-type (blue arrow) products. **e,** As for panel (b), except showing a 5′ truncated and inverted/deleted L1 T_F insertion and using an empty/filled PCR validation design, as per panel (d). **f,** Locus-specific methylation analysis schematic representation for 3 full-length *de novo* L1 insertions (panels b-d). After bisulfite conversion, the 5′ monomeric sequences of each L1 were PCR amplified using primer pairs (red arrows) specific to that locus. Amplicons were then pooled and sequenced as 2×300mer Illumina reads. Orange strokes indicate CpG dinucleotides covered by the assay. **g,** Methylation of the 3 L1 promoter sequences shown in panel (f), in the miPSC line where each *de novo* L1 insertion was identified. Each cartoon panel corresponds to an amplicon and displays 50 non-identical sequences (black circle, methylated CpG; white circle, unmethylated CpG; ×, mutated CpG) extracted at random from a much larger pool of available Illumina reads. The percentage of methylated CpG is indicated in the lower right corner of each cartoon in red. **h,** *top*: Rationale of a cultured cell retrotransposition assay[19,67]. A mouse L1 driven by its native promoter alone, or with the addition of a CMV promoter (CMVp), is tagged with an antisense orientated neomycin (NEO) reporter cassette interrupted by an intron. Cells harboring this construct become neomycin (G418) resistant upon retrotransposition. *bottom*: Retrotransposition assays conducted in HeLa cells. Constructs included: L1SM[68], a highly mobile synthetic L1 (positive control); L1SMmut2, L1SM with endonuclease and reverse transcriptase active site mutations (negative control); TG_F21, a mobile L1 G_F element[21]; $L1_{spa}$, a mobile L1 T_F element[22]; miPSC_1_L1 (panel b); miPSC_4_L1 (panel d). Data were normalized to $L1_{spa}$ and are shown as mean ± SD of three independent biological replicates, each of which comprised three technical replicates. Representative well pictures are shown below each construct. Note: L1SM retrotransposed very efficiently, leading to cell colony crowding in wells, and a likely underestimate of retrotransposition. Unless otherwise stated, L1 constructs incorporated a CMVp element.

Bulk miPSC populations harbor *de novo* L1 insertions

As *de novo* TE insertions can be overlooked by generalized SV calling algorithms[59], we used TEBreak[60] to identify non-reference TE insertions. Known non-reference genome TE insertions[56], and those found in MEF genotypic controls or multiple miPSC lines, were filtered, leaving 4 putative *de novo* L1 T_F insertions (**Fig. 1b-d**, **Table 1**, **Extended Data Fig. 2**, **Supplementary Table 3**). To achieve even greater coverage of potentially active TEs, we performed mouse retrotransposon capture sequencing (mRC-seq), which uses sequence capture probes to enrich Illumina libraries for the 5′ and 3′ genomic junctions of mobile TEs, including T_F, G_F and A subfamily L1s, B1 and B2 SINEs, and IAP and ETn ERVs (**Supplementary Table 1**)[28,61]. The combination of WGS and mRC-seq identified an additional 4 putative *de novo* L1 G_F and T_F insertions (**Extended Data Fig. 3**, **Table 1** and **Supplementary Table 3**).

We PCR amplified and fully characterized each putative L1 insertion sequence. Six events were full-length, retaining 2-7 monomers at their 5′ end, and could only be PCR amplified in the miPSC line where they were detected by genomic analysis (**Fig. 1b-d**, **Extended Data Fig. 2** and **Extended Data Fig. 3**). An additional L1 (labeled miPSC_6_L1) was very heavily 5′ truncated and confirmed by PCR to be private to one miPSC line (**Extended Data Fig. 3**). The final example (miPSC_2_L1) was heavily 5′ truncated and inverted[62] and could be PCR amplified in 7/9 miPSC lines representing all 3 germ layers of animal A67 (**Fig. 1e** and **Extended Data Fig. 2**). miPSC_2_L1 most likely represented a mosaic insertion that arose early in the embryonic development of animal A67, as found previously[28,30,63,64]. Each insertion carried TSDs of 13-19nt, a long and pure 3′ polyA tract, and integrated at a degenerate L1 endonuclease recognition motif (5′-TTTT/AA-3′) (**Table 1**). These hallmarks were consistent with *bona fide* TPRT-mediated L1 retrotransposition events[15,16,19,65,66]. Including the mosaic miPSC_2_L1 insertion, 10/26 miPSC lines harbored at least one PCR validated *de novo* L1 insertion. Not counting miPSC_2_L1, miPSCs from all 3 animals and 4/9 primary cell types, representing each germ layer, presented at least one *de novo* L1 insertion (**Fig. 1a** and **Supplementary Table 3**). Of these insertions, 4/7 were detected in astrocyte-derived miPSCs and 6/7 in miPSCs obtained from primary cells in the bottom 50% of reprogramming efficiencies (**Fig. 1a**), though neither of these proportions were statistically significant (binomial test). Notably, down-sampling to 11× depth WGS, as per[12], indicated an expected 95% probability of finding none of the validated *de novo* insertions (**Extended Data Fig.**

4a).

Comprehensive capillary sequencing of the 3 full-length insertions (miPSC_1_L1,

miPSC_3_L1 and miPSC_4_L1) revealed that each had intact ORFs (**Fig. 1b-d**). To assess the

potential for further mobilization of these newly retrotransposed elements, we first used

multiplexed L1 locus-specific bisulfite sequencing[34,61] to measure CpG methylation of their most

5′ promoter monomers (**Fig. 1f**). All 3 full-length insertions were fully unmethylated in a subset

of miPSCs, and their methylation decreased with distance from the L1 5′ end (**Fig. 1g**). Next, we

cloned and tested miPSC_1_L1 and miPSC_4_L1 in a cultured cell retrotransposition assay[19,67],

using the natural elements L1$_{spa}$ (T$_F$ subfamily)[22] and TG$_F$21 (G$_F$ subfamily)[21] as positive controls,

as well as the highly mobile synthetic L1 T$_F$ element L1SM[68]. As evidence of their *in vitro*

mobility, miPSC_1_L1 and miPSC_4_L1 retrotransposed efficiently in HeLa cells, when

expressed from their native promoter alone or with the addition of a cytomegalovirus promoter

(**Fig. 1h**). Thus, endogenous L1 retrotransposition in miPSCs is driven by highly mobile donor

L1s that can produce incompletely methylated, retrotransposition-competent offspring L1s.

Single-cell miPSC clones reveal extensive L1-mediated endogenous retrotransposition

Despite *de novo* L1 insertions being present in 10/26 miPSC lines, we were concerned that

additional mutations could be obscured by the heterogeneous mixture of cellular clones contained

in bulk reprogrammed miPSCs. We therefore reprogrammed MEFs from one of our

C57BL/6×129S4Sv/Jae animals (labeled I222e2), isolated individual miPSCs via FACS, and

expanded 18 clones cultivated in serum until p3 (**Fig. 2a**). To assess the impact of standard and

naïve culture conditions, respectively, upon L1 activity, each clone was divided and then further

expanded in serum or 2i until p6 (**Fig. 2a**). We then applied ~41× average genome-wide depth

Illumina WGS and mRC-seq to miPSC single-cell clones 1-9, and mRC-seq only to clones 10-18,

with each clone analyzed after culture in serum or 2i media (**Fig. 2a**, **Extended Data Fig. 1** and

Supplementary Table 1). Deep WGS was performed on the parental I222e2 MEF population,

attaining cumulative 117× genome-wide depth, in addition to mRC-seq (**Supplementary Table**

1). Using the WGS data, we again called concordant SNVs and SVs private to one miPSC clone,

while excluding known germline variants and those found in the parental MEFs. We found, on

average, ~100 and ~1 private SNVs and SVs per miPSC clone, respectively, almost all of which

Fig. 2: Frequent *de novo* TE insertions in MEF-derived clonally expanded miPSC lines. a, Experimental design to generate single-cell miPSC clonal lines. Bulk MEFs from a Col1a1-tetO-OKSM mouse (animal I222e2) were purified and reprogrammed by addition of doxycycline. Individual *Oct4*-GFP positive miPSCs were then isolated via FACS, expanded in serum for 3 passages, and then cultured in serum- or 2i-containing miPSC media for 3 additional passages. DNA was then extracted and analyzed by WGS and mRC-seq for 9 single-cell clones (for both serum and 2i conditions), with 9 further clones analyzed only with mRC-seq. **b,** A full-length *de novo* B2 inserted and orientated in antisense to intron 15 of *Brca1*. PolyA tract length is indicated immediately 3′ of the B2. TSDs are depicted as gray arrows flanking the B2. PCR validation (gel pictures shown) involved an empty/filled PCR design where both primers (red arrows) are outside of the B2, generating "filled" B2 (red arrow) and "empty" wild-type (blue arrow) products. The B2 was amplified only in either the serum or 2i conditions for the single-cell clone (number 7) where the B2 was detected by genomic analysis, and not in the matched parental MEFs, the C57BL/6 strain, or a single-cell clone (number 16) selected at random. **c,** A full-length (3 monomers) L1 A subfamily element inserted *de novo* antisense to intron 7 of *Gpc5*. Sequence characteristics and PCR validation results are shown as in panel (b). Promoter monomers are shown as triangles within the L1 5′UTR. **d,** As in panel (b), except showing an unusual intergenic B1 insertion flanked by both 5′ and 3′ polyA tracts. **e,** A full-length L1 G_F inserted *de novo* antisense to intron 60 of *Dmd*. PCR validation involved a 5′ genomic primer and a 3′ junction primer (red arrows). As indicated by a gray box with black stripes, the number of monomers is unknown but was >3. **f,** A heavily 5′ truncated, intergenic *de novo* L1 T_F insertion validated by empty/filled PCR, as per panel (b). Sequence features are annotated as per panel (b), with the addition of a 34nt 3′ transduction matching a donor L1 T_F located on Chromosome 9. PCR using primers (purple arrows) designed to amplify the entire donor L1 indicated it was polymorphic in our colony. Capillary sequencing indicated the donor L1 retained a promoter of 10 monomers and had intact ORFs. **g,** Locus-specific bisulfite sequencing analysis of the donor L1 promoter identified in panel (f), in MEFs, single-cell miPSC clones, and miPSC lines derived from primary cells. *top*: Assay design and primer locations. CpGs located in the first 3 monomers of the donor L1 were assessed. Orange and gray strokes indicate CpGs covered and not covered, respectively, by sequencing the amplicon with 2×300mer Illumina reads. *middle*: Mean percentages of donor L1 CpG methylation for 50 non-identical sequences selected at random from each sample. A two-tailed t test (*p<0.05) was used to compare serum and 2i culture conditions for single-cell miPSC clones 1-4. *bottom*: Percentages of fully unmethylated (mCpG=0, filled bars) and heavily unmethylated (0<mCpG<5, white bars) reads using the same sequencing data as displayed in the above histogram. **h,** Percentages of fully unmethylated (mCpG=0) reads corresponding to the donor L1 promoter identified in panel (f), for miPSCs cultured in serum or 2i conditions. Data represent mean methylation ± SD observed for single-cell miPSC clones 1-4. Significance testing was via two-tailed t test (***p<0.0001). **i,** As for panel (h), except using an assay targeting the L1 T_F subfamily monomer.

were detected in both the serum and 2i conditions for each clone (**Supplementary Table 2**). These

frequencies resembled those found by genomic analysis of bulk miPSCs, underlining that

heterogeneous and homogeneous fibroblast-derived miPSC populations are relatively free of

genomic abnormalities[10,12]. This experiment also indicated choice of serum or 2i media did not

impact the frequency of SNVs or SVs present in miPSCs.

By contrast, TEBreak revealed 35 putative *de novo* TE insertions absent from the parental

MEFs, all of which were found in both serum and 2i culture conditions for at least one miPSC

clone. Of these, 27 were detected by both WGS and mRC-seq, 6 by mRC-seq only, and 2 by WGS

only (**Supplementary Table 3**). Note that the 6 events found only by mRC-seq were detected in

the 9 miPSC clones (numbered 10-18) not analyzed with WGS. The 2 insertions found by WGS

alone were both moderately 5′ truncated and carried a 3′ transduction, two features that reduced

their probability of detection by mRC-seq, where enrichment probes target the 5′ and 3′ ends of

L1 consensus sequences[28,61]. We were able to PCR amplify 32 insertions in full and capillary

sequence at least their 5′ and 3′ junctions (**Fig. 2b-f**, **Extended Data Fig. 3** and **Supplementary**

Table 3). Two other putative TE insertions could only be amplified at their 5′ genome junction;

one of these (miPSC_29_L1) however also had strong 3′ WGS and mRC-seq support. We therefore

considered 33 TE insertions as validated *de novo* events (**Table 1**). Thirty-one of these were PCR

validated as private to only one miPSC clone, whereas the remaining two events were found in

either 2 clones (miPSC_23_B2) or 4 clones (miPSC_43_L1) (**Extended Data Fig. 3**). These last

two insertions were therefore present in subclones of the parental MEF population.

The 33 *de novo* insertions included 20, 3 and 1 T_F, G_F and A L1 subfamily members,

respectively, as well as 2 B1 and 7 B2 elements (**Supplementary Table 3**). All insertions

generated TSDs and a 3′ polyA tract, and integrated at a degenerate L1 endonuclease motif (**Table**

1). 14/24 L1 insertions retained at least one promoter monomer and were therefore considered full-

length (**Table 1**). Of the remaining 10 L1s, 3 were 5′ inverted. One unusual B1 insertion,

miPSC_18_B1, was flanked by 5′ and 3′ polyA tracts as well as TSDs (**Fig. 2d**), likely arising via

a variant of TPRT[69]. While no TE insertions were found in protein-coding exons, 14 were intronic,

including a B2 antisense to the tumor suppressor gene *Brca1* (**Fig. 2b**) and an L1 G_F antisense to

the dystrophin gene *Dmd* (**Fig. 2e**). 16/18 miPSC clones (88.9%) harbored at least one *de novo*

TE insertion, including all clones analyzed with both WGS and mRC-seq (**Supplementary Table**

1). Clone 2 contained the most (6) insertions. No *de novo* ERV insertions were found.

Among 277 high confidence heterozygous non-reference TE insertions (**Supplementary**
Table 4) found in the parental MEF population, 97.0% were detected on average in each miPSC
clone surveyed with WGS and mRC-seq. Down-sampling followed by seeking at least one WGS
or mRC-seq read in support of these non-reference insertions suggested our approach would
distinguish approximately 50%, 95% and 99% of *de novo* TE insertions from pre-existing
subclonal TE insertions present in 1%, 5% and 10% of cells, respectively (**Extended Data Fig.**
4b), whereas mRC-seq alone would achieve respective sensitivities of approximately 22%, 69%
and 89%. Consistently, only 2/33 PCR validated TE insertions in the miPSC clones were subclonal
in the parental MEFs (**Table 1**, **Extended Data Fig. 3**). An additional down-sampling analysis
indicated *de novo* TE insertions were likely to be detected at a lower average WGS depth in the
single-cell miPSC clones than insertions found in the bulk miPSC experiments (**Extended Data**
Fig. 4a), in agreement with the greater homogeneity of the clonal miPSC cultivars. Deep
sequencing of miPSCs and parental MEFs therefore enabled reliable detection and distinction of
TE insertions arising before and during reprogramming.

A polymorphic retrotransposition-competent L1 eludes methylation
Six *de novo* L1 insertions found in the miPSC clones carried 3′ transductions, flanking sequences
generated when PolII bypasses the native L1 polyA signal in favor of a downstream alternative[70–
74] (**Table 1**). Of these transductions, 5 were either too short to reliably map to the genome, or
mapped to multiple locations (**Supplementary Table 3**). The remaining 34bp transduction
accompanied a 5′ truncated L1 T_F insertion on Chromosome 12 (miPSC_10_L1) (**Fig. 2f**). While
the transduction aligned uniquely to Chromosome 9, a donor L1 was not present adjacent to this
reference genome location. However, PCR amplification revealed an L1 T_F immediately upstream
of the transduced sequence (**Fig. 2f**). This donor L1 was polymorphic in our
C57BL/6×129S4Sv/Jae animals and retained a 5′ promoter comprising an unusually high number
of monomers (10). Capillary sequencing confirmed the donor L1 possessed intact ORFs. L1 locus-
specific bisulfite sequencing revealed that few (24.1%) of the CpG dinucleotides in the first two
monomers of the donor L1 promoter were methylated in MEFs (**Fig. 2g** and **Extended Data Fig.**
5), as opposed to 7.3% in a subset of single-cell miPSC clones cultured in serum, and 1.3% for the

same miPSC clones when cultured in 2i conditions (**Fig. 2g**). This difference in CpG methylation between culture conditions was significant (p<0.05, two-tailed t test). The donor L1 promoter was fully unmethylated in nearly all miPSCs cultured in 2i (**Fig. 2g** and **Extended Data Fig. 5**). Indeed, significantly more (p<0.0001, two-tailed t test) fully unmethylated sequences were found for the donor L1 promoter in 2i conditions than in serum, possibly as a consequence of global naïve state hypomethylation (**Fig. 2h**). Among the bulk reprogrammed miPSCs obtained from animals A67 and A172, which carried the donor L1 (**Fig. 2f**), only 9.1% of CpG dinucleotides were methylated in the donor L1 promoter, and fully unmethylated sequences were identified in all miPSC lines (**Fig. 2g** and **Extended Data Fig. 5**). By contrast, in MEFs, 83.6% of CpG dinucleotides in L1 T_F promoter monomers genome-wide were methylated, compared to 45.2% among the A67 and A172 miPSC lines (**Extended Data Fig. 6**), and resembled L1 T_F promoter methylation reported elsewhere for differentiated primary cells[48,61]. L1 T_F subfamily monomers were also significantly (p<0.001, two-tailed t test) less methylated in 2i (34.3%) miPSC conditions than serum (53.5%), leading to an increase in fully unmethylated monomers (**Fig. 2i** and **Extended Data Fig. 6**). These bisulfite sequencing analyses highlighted genome-wide and persistent relaxation of L1 T_F methylation in miPSCs, leaving mobile L1 promoters completely unmethylated.

Reprogramming is unaffected by L1 reverse transcriptase inhibition

Lamivudine (3TC) is a potent nucleoside reverse transcriptase inhibitor known to limit engineered L1 retrotransposition without impacting telomerase or ERV mobility[75,76]. In previous retrotransposition assays conducted in cultured HeLa cells, 3TC was tested at a maximum concentration of 25µM against the codon-optimized L1SM element, reducing its mobility by ~50%[75]. By performing titration experiments to optimize the use of 3TC during miPSC generation, we determined that 3TC concentrations of up to 100µM did not reduce viability of cultured MEFs or miPSCs (**Extended Data Fig. 7a**) or MEF reprogramming efficiency (**Fig. 3a** and **Extended Data Fig. 7b-e**). Using a wild-type L1 T_F carrying an mCherry retrotransposition indicator cassette (L1-mCherry), we found 100µM 3TC reduced mouse L1 retrotransposition by ~95% in cultured HeLa cells (**Fig. 3b**). Thus, 3TC could be used, without apparent drawbacks, to limit L1-mediated mutagenesis arising during reprogramming and miPSC cultivation.

Fig. 3: Long-read detection of retrotransposition in HeLa cells and MEF-derived bulk miPSC lines. a, Bulk MEFs were reprogrammed by the addition of doxycycline. *Oct4*-GFP positive miPSCs were then sorted and expanded in serum. Six miPSC lines were reprogrammed and cultured in media containing 100µM lamivudine (3TC), and six miPSC lines generated without lamivudine (CTRL). DNA was extracted from MEFs and miPSCs and then ONT sequenced. **b,** *top left*: retrotransposition indicator plasmid L1-mCherry consists of the pCEP4 backbone (CMV promoter, black; SV40 polyadenylation signal, open lollipop; hygromycin resistance gene, white) containing a wild-type L1 T_F element (5′UTR, light purple; ORFs, dark purple). An mCherry reporter gene equipped with an EF1α promoter and HSVtk polyadenylation signal (black lollipop) is inserted into the L1 3′UTR antisense to the L1. The mCherry sequence is interrupted by an intron in sense orientation relative to the L1, ensuring mCherry expression only upon retrotransposition. *bottom left*: retrotransposition assay timeline. Cells were split (S), transfected (T), and cultured in hygromycin-containing medium with and without 100µM 3TC. Retrotransposition efficiency was assessed by flow cytometry 8 days post-transfection (R). *top right*: fluorescence microscopy images showing representative wells at 8 days post-transfection with L1 T_F (left), reverse transcriptase mutant (RT⁻) L1 T_F (middle), and L1 T_F treated with 100µM 3TC (right). *bottom right*: Retrotransposition efficiency assessed by flow cytometry, relative to L1 T_F. Histogram depicts the mean and standard deviation of three independent biological replicates (black dots) consisting of three technical replicates each. **c,** Target site duplication (TSD) size distributions for L1-mCherry retrotransposition events recovered from HeLa cells via TLDR analysis of ONT sequencing, divided into integrants detected by >1 (top) or 1 (bottom) spanning read. Inset sequence logos[120] display the observed integration site motif, as preferred by the L1 endonuclease (EN). **d,** Integrative genomics viewer (IGV)[121] inspection of 3 example *de novo* TE insertions. Cartoons (top) show a full-length (>6 monomers) L1 T_F insertion, a heavily 5′ truncated L1 T_F insertion, and a SINE B2 insertion, each flanked by TSDs (gray triangles) and a 3′ polyA tract, and integrated at the indicated L1 EN motif. IGV snapshots (bottom) show ONT read alignments from the miPSC line carrying each TE insertion (purple box). **e,** A 5′ truncated intergenic *de novo* L1 T_F insertion. PolyA (A_n) tract length is indicated immediately 3′ of the L1. PCR validation primers are shown as red arrows. Underneath is shown a PCR validation agarose gel containing the 5′ junction PCR product (red arrow) only in the miPSC line where the L1 was detected by ONT sequencing. Note: DNA was obtained from two parental MEF aliquots, one corresponding to miPSC CTRL/3TC lines 1-3 and one to miPSC CTRL/3TC lines 4-6. **f,** TSD size distributions for *de novo* L1 (top) and SINE (bottom) insertions detected in miPSC lines via ONT sequencing, with inset integration site sequence logo, as per panel (c). **g,** Putative reprogramming-associated L1-mediated insertion counts detected by ONT sequencing in CTRL and 3TC-treated miPSCs. Replicate (*n*=6 per group) data points are marked by black dots and represented as the mean ± SD, *p<0.02, two-tailed t test.

Nanopore genomic analysis of TE insertions in bulk miPSCs

A single DNA sequencing read, if of sufficient length and quality, can completely resolve a *de novo* TE insertion present in a heterogeneous cell population, as well its genomic flanks and accompanying TPRT hallmarks[77]. Long-read sequencing, as for example developed by Oxford Nanopore Technologies (ONT), is well suited to this application, and can locate TE insertions within repetitive genomic regions refractory to short-read methods[35,77–79]. As a proof-of-principle before analyzing miPSCs, we applied PCR-free ONT sequencing to 5 HeLa cell lines expanded from single colonies (~5× genome-wide depth per colony) harboring mouse L1-mCherry retrotransposition events (**Supplementary Table 1**). Using the TLDR long-read TE insertion detection pipeline[35], we identified 41 L1-mCherry insertions spanned by at least one ONT read (**Supplementary Table 3**). L1-mCherry insertions overwhelmingly bore TPRT hallmarks regardless of whether they were detected by one, or more than one, ONT read (**Fig. 3c**), showing that single spanning ONT reads can reliably recover *bona fide* retrotransposition events.

Next, to survey endogenous retrotransposition in miPSCs, we ONT sequenced (~20× average genome-wide depth) 6 bulk miPSC lines reprogrammed in the presence of 100µM 3TC, 6 control miPSC lines not treated with 3TC, and matched parental MEFs (**Fig. 3a** and **Supplementary Table 1**). Using these data, TLDR identified 3,975 non-reference TE insertions carried by the parental MEFs (**Supplementary Table 4**). Of these, 3,429 (86.3%) corresponded to previously known insertions[56] and 99.6% were found in each miPSC line, on average (**Supplementary Table 4**). To gauge the general tractability of PCR validation applied to this dataset, we used a panel of 4 heterozygous non-reference TE insertions (**Supplementary Table 3**) and confirmed these all PCR amplified in the MEFs and miPSCs (**Extended Data Fig. 8a**).

An additional 43 TE insertions were each detected in only one miPSC line and not the parental MEFs, or in the earlier Illumina sequencing, and were supported by at least one spanning ONT read (**Fig. 3d** and **Supplementary Table 3**). Performing PCR validation of these insertions, a step mainly intended to exclude them being polymorphisms carried by the parental or feeder MEF populations, we could amplify one (miPSC_50_B1) in the parental MEFs (**Extended Data Fig. 8b**). The remaining 42 putative *de novo* events comprised 16, 4 and 2 L1 T_F, G_F, and A insertions, respectively, as well as 5 and 13 SINE B1 and B2 insertions, respectively, and 2 ERV insertions. One L1 T_F insertion (miPSC_69_L1) PCR amplified in multiple miPSC lines

(**Extended Data Fig. 8c**) and one L1 T$_F$ insertion (miPSC_87_L1) amplified only in the miPSC

line where it was detected by ONT sequencing (**Fig. 3e**). While the remaining insertions could not

be PCR amplified in any sample, all 40 of the putative *de novo* L1 and SINE insertions carried

clear TPRT hallmarks (**Fig. 3f** and **Supplementary Table 3**) and on this basis we considered them

bona fide retrotransposition events. As well, both ERVs corresponded to the mobile intracisternal

A-particle (IAP) family, presented a typical proviral structure of two long terminal repeats flanking

an internal coding sequence, and generated TSDs of the expected size (6bp)[80,81] (**Supplementary**

Table 3). It was not clear whether the difference in (two) *de novo* ERV insertions being found here

by ONT sequencing, and none by the earlier Illumina sequencing of different miPSC lines, was

due to chance or unknown technical reasons. Significantly fewer (p<0.02, two-tailed t test) putative

reprogramming-associated L1-mediated insertions were found on average in the 3TC-treated

miPSCs (~1.3 per line) than in the control miPSCs (~5.2 per line) (**Fig. 3g**), consistent with L1

inhibition by 3TC (**Fig. 3b**). Overall, detection of endogenous retrotransposition events in bulk

miPSCs by ONT sequencing yielded results orthogonal and complementary to our short-read

genomic analyses.

Genome-wide DNA demethylation during reprogramming focused on young L1 loci

A major feature of reprogramming mouse fibroblasts to a pluripotent state is globally reduced

DNA methylation[43,44,47,48]. Although bisulfite sequencing can estimate the overall methylation of

TE families, at specific genomic loci it can typically only resolve CpGs close to the termini of full-

length L1s not located in highly repetitive regions. To generate a comprehensive genome-wide

view of DNA methylation changes during reprogramming, and complement our bisulfite

sequencing data, we analyzed the ONT data obtained from MEFs and miPSCs (control and 3TC-

treated) using Methylartist[35,82]. While methylation in control miPSCs was reduced genome-wide

when compared to MEFs, on protein-coding gene promoters and amongst all of the TE families

considered, the very youngest L1 subfamilies (T$_{FI}$ and T$_{FII}$) displayed by far the greatest median

methylation change (-54.0%) (**Fig. 4a**, **Extended Data Fig. 9a,b** and **Supplementary Table 5**).

90.3% of full-length L1 T$_{FI}$ and T$_{FII}$ copies were significantly (p<0.01, Fisher's exact test with

Bonferroni correction, and ΔmC >25%) less methylated in control miPSCs than MEFs

(**Supplementary Table 5**), with this demethylation most pronounced in the monomeric L1 5′UTR

Fig. 4: Genome-wide methylation analyses via ONT sequencing. a, CpG methylation ascertained by ONT sequencing of MEFs (brown), control miPSCs (CTRL, green), and miPSCs derived in the presence of lamivudine (3TC, pink). Results are shown for the whole genome (10kbp windows), the proximal promoters (-1000,+500) of protein-coding genes[119], the 5′UTR of T_F, G_F, and A-type L1s >6kbp, B1 and B2 SINEs, and MERVL MT2 and IAP long terminal repeats. Note: replicates from control and 3TC-treated miPSCs are displayed in aggregate. **b,** Composite L1 T_F methylation profiles. Each graph displays 100 profiles. A schematic of the T_F consensus is provided at top. Average values are indicated by more thickly colored lines. **c,** Methylation profile of the *Fsd1l* locus obtained by ONT sequencing. The first panel shows an L1 T_F orientated in sense to intron 6 of *Fsd1l*, as well as an expressed sequence tag (EST) obtained from a mouse ESC sample and supporting a transcript initiated in the T_F 5′UTR and spliced into a downstream *Fsd1l* exon. The second panel displays ONT read alignments, with unmethylated CpGs colored in brown (MEFs), green (control miPSCs) and pink (3TC-treated miPSCs), methylated CpGs colored black, and CpGs not confidently called, i.e. abs(log-likelihood ratio) > 2.5, omitted. The displayed miPSC data were downsampled to the approximate depth of the MEF data with Picard DownsampleSam. The third panel indicates the relationship between CpG positions in genome space and CpG space, including those corresponding to the T_F 5′UTR (shaded light blue). The fourth panel indicates the fraction of methylated CpGs. Note: this L1 T_F is polymorphic amongst inbred mouse strains[56].

(**Fig. 4b**). By contrast, we observed no significant local or global differences in L1 T_F or protein-

coding gene promoter methylation between control and 3TC-treated miPSCs (**Fig. 4** and

Supplementary Table 5). Thirty-six L1s initiated transcription of a spliced mRNA from their

5′UTR, as defined by GenBank expressed sequence tags, including alternative promoters for

protein-coding genes expressed in pluripotent cells, such as *Fsd1l* (**Fig. 4c** and **Supplementary**

Table 5). We also identified full-length L1s demethylated in both MEFs and miPSCs (**Extended**

Data Fig. 9c), in line with prior human data suggesting certain L1 loci evade DNA methylation in

differentiated cells[34,83]. In sum, ONT analysis showed global reprogramming-associated

demethylation was most accentuated for the youngest L1s, where retrotransposition potential is

concentrated, creating opportunities for L1-driven mobilization and protein-coding gene

alternative promoters.

Discussion

This study demonstrates miPSCs incompletely silence mobile TE families and routinely harbor *de*

novo TE insertions. While some TE insertions occur in parental cells and are inherited by miPSCs,

our data suggest the majority arise during reprogramming or very early upon reaching

pluripotency. In support of this view, firstly, we observed profound hypomethylation of young L1

promoters in miPSCs and not parental cells. As shown elsewhere, L1 mRNA abundance is low in

fibroblasts and increases greatly upon reprogramming[45,46,52], while engineered L1 reporter genes

retrotranspose >10-fold more frequently in hiPSCs and human ESCs than in fibroblasts[51,52].

Secondly, 38/41 *de novo* TE insertions detected by Illumina sequencing PCR validated in only one

miPSC line each. These and the 42 putative *de novo* TE insertions identified by ONT sequencing

were absent from all other samples in the study, as assayed by PCR and deep WGS. Finally, given

the similar mutation calling thresholds of Illumina sequencing applied to heterogeneous (bulk) and

homogenous (single-cell clone) miPSC populations, we note that private SNVs (~100 per line) and

SVs (~1 per line) were detected at similar frequencies in either experiment, whereas far more *de*

novo retrotransposition events were found in the latter dataset. One explanation for this result is

that a relatively small number of clones dominate bulk reprogramming experiments[84] and most

SNVs and SVs predate reprogramming[10,11,13], while retrotransposons mainly mobilize during

reprogramming. This position is consistent with a prior WGS analysis that, alongside thousands

of SNVs, identified no somatic L1 insertions among 10 human fibroblast clones generated from

single cells[85]. It is nonetheless possible that additional somatic variants would have been annotated

if primary single-cell clones, as analyzed elsewhere[86], were reprogrammed. However, the

introduction of multiple bottlenecks followed by clonal expansion prolongs cell culture and could

thereby exaggerate mutation frequencies. The parental MEFs used here cannot in any case be

clonally expanded from single cells, and reprogram extremely inefficiently after more than three

passages in culture[87,88], and for these reasons we did not prepare single-cell MEF clones prior to

reprogramming.

Previous experiments employing hiPSCs and mouse and human ESCs showed L1 de-

repression and mobilization were likely to take place in pluripotent cells[34,41,42,45,46,48,50–52,89].

Notably, 31/56 (55.4%) *de novo* L1 insertions found here in miPSCs were full-length, a similar

percentage to that observed previously in hiPSCs (57.1%)[45]. New full-length L1 insertions have

potential for further retrotransposition and were largely unmethylated in miPSCs. Their CpG

dinucleotides presented a "sloping shore" of methylation, as found elsewhere for newly

retrotransposed CpG islands[34,35,48,90], where methylation decreases from the L1 5′ genome junction

and forms a trough before sharply increasing over the L1 ORFs. Only 3/56 (5.4%) L1 insertions

corresponded to the A subfamily, while the remainder were T_F and G_F elements, consistent with

relative activity levels revealed by sequencing extended mouse pedigrees and mouse tumors[28,61].

De novo SINE B1 and B2 insertions, mediated *in trans* by the L1 protein machinery[91], were also

detected in miPSCs, in line with L1-mediated *Alu* SINE insertions arising in hiPSCs and human

ESCs[45,89,92]. Discovery of *de novo* TE insertions in low-passage miPSCs derived from multiple

parental cell types suggests endogenous retrotransposition may be an intrinsic risk of the

epigenome remodeling required for the acquisition of pluripotency[7,43–45,47].

Exonic retrotransposon insertions are clear in their potential to cause disease[93]. However, the

L1 endonuclease does not favor exons[65,66], which are depleted for its AT-rich recognition motif

and make up only ~2% of the genome[94], and none of the 81 *de novo* L1-mediated insertions

reported here in miPSCs were exonic. By comparison, L1 integration within introns occurs much

more frequently. We observed 30/81 (37.0%) L1-mediated insertions in introns, in line with the

proportion of the genome occupied by these regions[65,66]. Whilst intronic events are less likely to

be pathogenic than exonic insertions, they can perturb gene expression, for example through

reduced transcriptional elongation[95] or, as shown here, the provision of new promoter elements[96].

Retrotransposition into the introns of protein-coding genes, as observed here for *Brca1* and *Dmd*,

could therefore undermine miPSC models of human disease. Such mutations necessitate screening

of miPSC lines[4]. Fortunately, strategies to minimize TE-mediated mutagenesis, including via the

use of 3TC or another L1 reverse transcriptase inhibitor[75], appear achievable without affecting

DNA methylation or reprogramming efficiency. While the frequency of *de novo* endogenous L1-

mediated insertions found in miPSCs was significantly reduced by 3TC, a handful of events, likely

arising prior to reprogramming, were nonetheless observed in 3TC-treated miPSCs. Reverse

transcriptase inhibitors thus may be incorporated into future miPSC derivation protocols to

attenuate *de novo* retrotransposition, in addition to genomic screening of miPSC lines.

Methods

Ethics statement

All animal experimentation was performed under the auspices and approval of the Monash

University Animal Research Platform Animal Ethics Committee (Approval Numbers MARP-

2011-172-Polo, MARP-2011-171-BC-Polo, MARP-2017-151-BC-Polo, and ERM# 21634).

Adult *Oct4*GFP-OKSM-M2rtTA mouse tissue somatic cell isolation and reprogramming

Induced pluripotent stem cells were generated from adult and embryonic *Oct4*GFP-OKSM-

M2rtTA doxycycline inducible reprogrammable mice[53]. These animals are heterozygous for an

Oct4-GFP reporter and an OKSM cassette targeted to the *Collagen1α1* locus, and homozygous for

the ROSA26-M2rtTA allele from the ubiquitous *ROSA26* locus. The polycistronic cassette is under

the control of a tetracycline-dependent promoter *(tetOP)*. Hence, upon the addition of doxycycline,

M2rtTA binds to the *tetOP*, thereby inducing OKSM expression. *Oct4*GFP-OKSM-M2rtTA mice

were housed at the Monash University Animal Research Platform animal facility.

Bone marrow extraction and FACS purification of granulocytes and hematopoietic stem

(LSK) cells were performed as previously described[97]. In brief, harvested bone marrow cells were

labeled using a two-step sequential antibody labeling procedure using the following primary

conjugated antibodies: 1:200 dilution of Anti-Mouse CD5 FITC antibody (BD Biosciences, Cat#:

553020), 1:100 dilution of Anti-Mouse B220 FITC antibody (BD Biosciences, Cat#: 557669),

1:200 dilution of Anti-Mouse TER-119 FITC antibody (BD Biosciences, Cat#: 557915), 1:400 dilution of Anti-Mouse Sca-1 PB antibody (Biolegend, Cat#: E13-161.7), 1:200 dilution of Anti-Mouse cKit APC antibody (BD Biosciences, Cat#: 553356), 1:200 dilution of Anti-Mouse SSEA1 Biotinylated antibody (Thermo Fisher Scientific, Cat#: 13-8813-80), 1:200 dilution of Anti-Mouse Gr-1 APC-Cy7 antibody (Biolegend, Cat#: 108423) and 1:1000 dilution of Anti-Mouse Mac1 PE antibody (Biolegend, Cat#: 101207). This was followed by the secondary labeling step with 1:200 dilution of Streptavidin PE-Cy7 antibody (BD Biosciences, Cat#: 557598). Cells were isolated and sorted using an Influx Cell Sorter Instrument (BD Biosciences) with a 100µm nozzle. Samples were resuspended in phosphate buffered saline (PBS) supplemented with 2% fetal bovine serum (Thermo Fisher Scientific, Cat#: SH30071.03FBS, Hyclone). FACS sorting for these and the cell types below were performed with 2µg/mL Propidium Iodide (PI) (Sigma Aldrich, Cat#: P4864) in order to exclude non-viable cells. Granulocytes were isolated using the following cell surface marker profile: $CD5^-/B220^-/Ter119^-/Sca1^-/cKit^-/SSEA1^-/Gr1^+/Mac1^+$, whilst LSK cells were isolated from bone marrow using the following cell surface marker profile: $CD5^-/B220^-/Ter119^-/Sca1^+/cKit^+/SSEA1^-/Gr1^-/Mac1^-$.

Fibroblasts were isolated from both ear lobes from each mouse. Tissue pieces were resuspended in 0.25% Trypsin-EDTA (Thermo Fisher Scientific, Cat#: 25200-072) solution, and after 5min incubation at room temperature, were mechanically minced using two surgical blades for a further 2min. iPSC medium was used to inactivate trypsin, and dissociated pieces were transferred to a 15mL centrifuge tube (Corning). Tissue pieces were then transferred to a gelatin coated T-75 flask (Corning) and cells were left to grow for a further 7 days. $CD45^-/CD31^-/Thy1.2^{hi+}$ fibroblasts were fractionated by FACs using the following antibodies: a 1:100 dilution of Anti-Mouse CD31 antibody conjugated to FITC (Thermo Fisher Scientific, Cat#: 11-0311-81), a 1:100 dilution of Anti-Mouse CD45 antibody conjugated to FITC (Thermo Fisher Scientific, Cat#: 11-0451-810) and a 1:400 dilution of Anti-Mouse Thy-1.2 antibody conjugated to APC (Thermo Fisher Scientific, Cat#: 17-0902-81).

Liver epithelial cells were isolated according to an adaptation of a previously described method[98]. Briefly, 3mg/mL Collagenase Type 1 (Sigma-Aldrich, Cat#: C1639) solution was prepared in sterile PBS. Whole liver was transferred into a sterile 6cm petri dish and finely minced using fine dissecting scissors. Minced liver pieces were transferred to 15mL tube with preheated

Collagenase Type 1 (Sigma, Cat#: C1639). Tubes were left to agitate on a Thermomix (Eppendorf)

at 750rpm, 37°C for 15min. Following digestion, the tube was removed and the cellular suspension

was triturated with an 18G needle, until tissue chunks were mostly dissociated. Sample tubes were

then left to agitate for an additional 15min, until liver fragments were completely digested. The

sample suspension was again triturated, with a 21G needle, to generate a single cell suspension,

and then processed through a 40µm cell strainer into a clean 50mL centrifuge tube (Corning). After

rinsing in 2% FCS/PBS (wash buffer) and centrifuging for 5min at 1380rpm for 4°C, the

supernatant was removed and cells were resuspended in wash buffer and centrifuged once again.

Cells were counted and 5×10^6 cells were resuspended for sorting. Cells were labeled with primary

antibodies using a 1:100 dilution of Anti-mouse CD31 antibody conjugated to FITC (Thermo

Fisher Scientific, Cat#: 11-0311-81), followed by a 1:100 dilution of Anti-mouse CD45 antibody

conjugated to FITC (Thermo Fisher Scientific, Cat#: 11-0451-81) and 1:100 dilution of Anti-

mouse EpCAM antibody conjugated to eFluor450 (Thermo Fisher Scientific, Cat#: 48-5791-82).

Liver epithelial cells were isolated using the following cell surface marker profile: CD45$^-$/CD31$^-$

/EpCAM^{+hi}.

Thymus tissue was processed for thymic epithelial cell isolation as previously described[99].

Cells were labeled with the following antibodies: 1:400 dilution of Anti-mouse CD45 antibody

conjugated to APC-Cy7 (BD Biosciences, Cat#: 557659), 1:200 dilution of Anti-mouse TER-119

antibody conjugated to APC-Cy7 (BD Biosciences, Cat#: 560509), 1:6000 dilution of Anti-mouse

MHC Class II antibody conjugated to PB (Biolegend, Cat#: 107620) and 1:1000 dilution of Anti-

mouse EpCAM antibody conjugated to APC (Biolegend, Cat#: 118214). Thymic epithelial cells

were sorted according to the following cell surface marker profile: CD45$^-$/Ter119$^-$/MHC Class

II$^+$/EpCAM$^+$.

Intestinal stem cells were purified as previously described[100]. Cells were labeled with a

1:200 dilution of Anti-mouse CD45 antibody conjugated to BV510 (BD Biosciences, Cat#:

563891), 1:200 dilution of Anti-mouse CD31 antibody conjugated to BV510 (BD Biosciences,

Cat#: 563089), a 1:100 dilution of Anti-mouse CD24 antibody conjugated to Pe-Cy7 (Thermo

Fisher Scientific, Cat#: 25-0242-82), a 1:100 dilution of Anti-mouse EpCAM antibody conjugated

to eFluor450 (Thermo Fisher Scientific, Cat#: 48-5791-82), and 1:100 Anti-EphrinB2

unconjugated antibody (BD Biosciences, Cat#: 743763). In the secondary labeling step, a 1:200

dilution of Anti-mouse Alexa Fluor 555 polyclonal antibody (Thermo Fisher Scientific, Cat#: A-
31570) was used to detect the EphrinB2 antibody. Intestinal stem cells were fractionated according
to the following cell surface marker profile: CD45$^-$/CD31$^-$/CD24$^+$/EpCAM$^+$/Ephrin$^+$.

To obtain astrocytes, brain tissue was processed using a MACS Neural Tissue Dissociation
Kit (T) (Miltenyi Biotec, Cat#: 130-093-231) and manually dissected according to manufacturer's
instructions. Cells were then collected and incubated with antibodies directed against Glast1
(Allophycocyanin-conjugated, ACSA-1, 1:10 dilution) (Miltenyi Biotec, Cat#: 130-098-803),
1:100 dilution of Anti-mouse CD133 antibody conjugated to PE (Thermo Fisher Scientific, Cat#:
12-1331-80), 1:200 dilution of Anti-mouse CD45 antibody conjugated to PE-Cy7 (BD
Biosciences, Cat#: 552848) and 1:200 dilution of Anti-mouse CD31 antibody conjugated to PE-
Cy7 (Thermo Fisher Scientific, Cat#: 25-0311-82). Astrocytes were sorted and purified according
to the following cell surface marker profile: CD45$^-$/CD31$^-$/CD133$^-$/GLAST1$^+$.

Keratinocytes and bulge stem cells were isolated from epidermis as previously
described[101]. Cells were collected and incubated with antibodies against Anti-Mouse Integrin alpha
6 antibody (GoH3) conjugated to PE (1:600) (Abcam, Cat#: ab95703), a 1:200 CD104 antibody
conjugated to FITC (Biolegend, Cat#: 346-11A) and a 1:100 dilution of Anti-mouse CD34
biotinylated antibody (Thermo Fisher Scientific, Cat#: 13-0341-85) for 20min at 4°C. For
secondary antibody labeling, cells were incubated with 1:200 APC-Streptavidin antibody
(Biolegend, Cat#: 405207) to detect CD34 biotinylated antibody for 20min at 4°C. They were
then washed and resuspended in PI (2µg/mL) 1% BSA/PBS (Sigma-Aldrich, Cat#: A8412) and
passed through a 40µm cell strainer (BD Falcon) to produce single cell suspensions. Cells with the
surface marker profile of CD104$^+$/CD34$^+$/α6-integrin$^+$ were defined as bulge stem cells, and those
marked as α6-integrin$^-$/CD34$^+$ were defined as keratinocytes.

Reprogramming of the above 9 primary cell types was performed as follows: cells were
seeded into gelatinized tissue culture treated 6-well plates (Corning Costar, Cat#: CLS3506) and
cultured at 37°C and 5% CO$_2$ in iPSC media containing KnockOut DMEM (Thermo Fisher
Scientific, Cat#: 10829-018), 15% Fetal Bovine Serum (FBS) (Thermo Fisher Scientific, Cat#:
SH30071.03), GlutaMAX Supplement (Thermo Fisher Scientific, Cat#: 35050061), Penicillin-
Streptomycin (Thermo Fisher Scientific, Cat#: 15070063), MEM Non-Essential Amino Acids
Solution (Thermo Fisher Scientific, Cat#: 11140050), 2-Mercaptoethanol (Thermo Fisher

Scientific, Cat#: 21985023) and 1000U/mL Leukemia Inhibitory Factor (LIF) (Merck Millipore,

Cat#: ESG1107), supplemented with 2µg/mL of doxycycline (dox) (Sigma-Aldrich, Cat#: 33429-

100MG-R). iPSC medium supplemented with dox was replaced every alternate day after the first

3 days of reprogramming and withdrawn 4 days after the presence of iPSC-like colonies had

formed, with typical dome-shaped iPSC morphology. Cells were then cultured to confluency on a

layer of irradiated MEFs prior to further FACs purification and enrichment for Oct-GFP$^+$ cells.

Purified *Oct4*-GFP iPSCs were then bulk expanded in 175cm^2 cell culture flasks (Corning, Cat#:

CLS430825) and then frozen at a density of 1×10^6 cells/vial.

Mouse embryonic fibroblast isolation and reprogramming

Reprogrammable mouse embryonic fibroblast (MEF) cultures were derived as described

previously[102] from a E13.5dpc *Oct4*GFP-OKSM-M2rtTA embryo (animal I222e2) and cultivated

at 37°C, 5% O$_2$, 5% CO$_2$ in MEF medium containing DMEM High Glucose (Thermo Fisher

Scientific, Cat# 11960-044) with 10% FBS (Thermo Fisher Scientific, Cat#: SH30071.03), 1mM

Sodium Pyruvate (Thermo Fisher Scientific, Cat#: 11360-070), GlutaMAX Supplement (Thermo

Fisher Scientific, Cat#: 35050061), Penicillin-Streptomycin (Thermo Fisher Scientific, Cat#:

15070063), MEM Non-Essential Amino Acids Solution (Thermo Fisher Scientific, Cat#:

11140050) and 2-Mercaptoethanol (Thermo Fisher Scientific, Cat#: 21985023). MEFs were

reprogrammed by being placed in iPSC medium supplemented with 2µg/mL dox (Sigma-Aldrich,

Cat#: 33429-100MG-R) and cultured on irradiated MEFs at 37°C, 5% CO$_2$. iPSC colonies were

discerned according to GFP expression in the absence of dox. In addition to bulk iPSC cultures

(see below), single *Oct4*-GFP$^+$ cells were deposited via FACS individually into 96-well pre-

gelatinized tissue culture plates (Falcon, Cat#: 353072). Eighteen single-cell clones were bulk

expanded on 6-well pre-gelatinized tissue culture plates (Falcon, Cat#: 353046) and maintained in

serum or 2i conditions (see below).

Oct4-GFP$^+$ iPSC flow cytometry

For flow cytometry, cells were harvested by dissociating in 0.25% Trypsin EDTA (Life

Technologies) to yield a single cell suspension, and then resuspended in FACS wash (Phosphate

Buffered Saline with 2% Fetal Calf Serum) containing PI. Live cells were gated on the basis of

forward scatter, side scatter and PI exclusion. Flow cytometric gates were set using control iPSCs

that did not have endogenous GFP expression. Tubes were sorted according to GFP expression

using an Influx Cell Sorter Instrument (Becton Dickinson). Data collected were analyzed and

presented using FlowJo software. Sorted GFP$^+$ cells were then plated down on T-25 flasks

(Corning) and expanded onto T-150 flasks (Corning), before being frozen down at a density of

1×10^6 cells/vial.

Serum/LIF and 2i/LIF serum-free iPSC culture

Mouse iPSCs were maintained on irradiated primary MEFs, as previously described[8,103]. Briefly,

iPSCs were cultured on 0.2% Porcine Gelatin (Sigma-Aldrich, Cat#: G1890-500G) coated tissue

culture plates and flasks (Corning) on a feeder layer of irradiated MEFs (2×10^4 cells/cm^2). iPSC

medium was changed daily and cells were cultured at 37°C and 5% CO$_2$. Passaging was performed

when iPSCs reached 70% confluency. Alternatively, iPSCs were cultured on irradiated MEFs in

serum-free media containing knockout serum replacement (KOSR) and 2i/LIF[104]. Here, cells were

cultured in DMEM (Thermo Fisher Scientific, Cat#: 11960-044), 1000U/mL LIF (Merck

Millipore, Cat#: ESG1107), 0.1mM 2-Mercaptoethanol (Thermo Fisher Scientific, Cat#:

21985023), 1mM GlutaMAX Supplement (Thermo Fisher Scientific, Cat#: 35050061), 1%

Sodium Pyruvate (Thermo Fisher Scientific, Cat#: 11360-070), 0.1mM MEM Non-Essential

Amino Acids Solution (Thermo Fisher Scientific, Cat#: 11140050), 1% Penicillin-Streptomycin

(Thermo Fisher Scientific, Cat#: 15070063), with medium supplemented with 15% KOSR

(Thermo Fisher Scientific, Cat#: 10828-028), 1µm Mek1/2 Inhibitor (PD0325901) (Tocris, Cat#:

4192) and 3µm GSK3a/b inhibitor (CHIR99021) (Tocris, Cat#: 4423). Prior to genomic DNA

extraction, iPSCs depleted from irradiated feeders were dissociated with 0.5% Trypsin EDTA

(Thermo Fisher Scientific, Cat#: 25200-072). The irradiated MEFs were feeder depleted with

10mL of iPSC media for 45min in non-gelatinized T-25 flasks (Corning, Cat#: CLS3056). The

resultant iPSCs were collected as a supernatant in suspension medium.

Lamivudine titration experiments

iPSCs were cultured with primary irradiated MEFs, as above, for 9 days in concentrations of

lamivudine (3TC, Sigma-Aldrich, Cat#: L1295-10MG) ranging from 0 to 200µM and cell survival

calculated as a % of the starting population. Reprogrammable OKSM, rtTA3 MEFs were isolated

from embryonic day 13.5 embryos from *Oct4*-GFP;ROSA-rtTA-out;OKSM-72 mice as previously

described[44,105]. Doxycycline inducible reprogrammable MEFs were grown in media containing

2µg/mL dox (Sigma Aldrich Cat#: 33429-100MG-R) and 0-200µM 3TC for 15 days, with the

percentage cell survival calculated at days 3, 7, 10 and 15. Once 100µM was identified as the

optimal concentration of 3TC to assess its impact on L1 retrotransposition, 30,000

reprogrammable MEFs at passage 2 were seeded onto gelatinized 6-well plates and reprogrammed

in dox for 12 days, then cultured for an additional 4 days without dox. *Oct4*-GFP⁺ iPSCs were then

purified via flow cytometric sorting and expanded on irradiated MEFs for an additional 11 days,

then feeder depleted prior to DNA extraction for ONT sequencing. Reprogramming and iPSC

media contained serum, and either 100µM 3TC or no 3TC.

Illumina sequencing and genomic analysis

Genomic DNA was harvested from MEFs and iPSCs using a DNeasy Blood and Tissue Kit

(Qiagen, Cat#: 60594). DNA was quantified by a Qubit dsDNA HS Assay Kit (Life Technologies,

Cat#: Q32851) on a Qubit Fluorometer 3.0 (Life Technologies). For WGS, libraries were

generated using an Illumina TruSeq DNA PCR-free kit (Illumina, Cat#: 20015962) and sequenced

separately on an Illumina HiSeq X Ten platform (Macrogen, Korea).

For mRC-seq, libraries were prepared as follows: 1µg genomic DNA was sheared using a

Covaris M220 Focused Ultrasonicator in a 130µL microTUBE AFA fiber snap-cap vial (Covaris,

Cat#: 520045). The following parameters were used to gain 500bp insert libraries: 50W, duty

factor 20%, 200 cycles per burst, duration 55s. Size selection to remove fragments <300 bp was

performed using Agencourt AMPure XP beads (Beckman Coulter, Cat#: A63881) with a 1:0.6

DNA:beads ratio. Libraries were then generated by TruSeq Nano DNA LT kit (Illumina, Cat#:

20015964) using TruSeq DNA Single Indexes (Illumina, Cat#: 20015960 and 20015961) and run

on a 2% agarose gel (Bioline, Cat#: BIO-41025) pre-stained with SYBR Safe Nucleic Acid Gel

Stain (Invitrogen, Cat#: S33102). For ~500bp insert size libraries the target gel fragment size was

600-650bp, which was excised under a Safe Imager 2.0 Blue-Light Transilluminator (Invitrogen).

DNA was purified using a MinElute Gel Extraction Kit (Qiagen, Cat#: 28606) according to the

manufacturer's instructions. DNA was eluted in 25µL molecular grade water. Enrichment of DNA

fragments was performed as described for Illumina TruSeq Nano DNA LT Kit (Illumina, Cat#:
20015964). Sample clean up was performed with Agencourt AMPure XP beads (Beckman
Coulter, Cat#: A63881) using a 1:1.1 ratio of DNA to beads. Amplified libraries were eluted in
30μL molecular grade water and quantified using a Bioanalyzer DNA 1000 chip (Agilent
Technologies, Cat#: 5067-1504).

mRC-seq hybridization was performed as previously described[28]. Hybridization reactions
were washed using SeqCap Hybridization and Wash Kit (Roche, Cat#: 05634261001) and DNA
eluted in 50μL molecular grade water. Two post-hybridization LM-PCR reactions per sample were
performed using 20μL Enhanced PCR Mix, 5μL PCR Primer Cocktail from the Illumina TruSeq
Nano DNA LT Kit (Illumina, Cat#: 20015964) and 25μL sample. PCR was performed with the
following cycling conditions: 95°C for 3min, 8 cycles of 98°C for 20s, 60°C for 15s, and 72°C for
30s, followed by 72°C for 5min. The two PCR reactions for each sample were pooled and cleaned
up using the QIAquick PCR Purification Kit (Qiagen) and samples eluted in 15μL Elution Buffer
(Qiagen, Cat#: 28706). Quantity and fragment size were determined using a Bioanalyzer DNA
1000 chip (Agilent Technologies, Cat#: 5067-1504). Libraries were pooled and sequenced on an
Illumina HiSeq X Ten platform (Macrogen, Korea).

Reads were aligned to the mm10 reference genome using bwa-mem[106] version 0.7.12 with
parameters -M -Y. Duplicate reads were marked via Picard MarkDuplicates version 1.128. Indel
Realignment was carried out via GATK IndelRealigner (3.7). SNVs were called by GATK
HaplotypeCaller 3.7[54] to generate GVCFs and GenotypeGVCFs to obtain cohort-level calls. SNVs
were also called using freebayes[55] filtered to remove known mouse strain germline variants[56]. SVs
were called using Delly2 and GRIDSS 2.0.0[57,58], using calls with concordant non-filtered precise
breakends. SNVs and SVs, including potential private variants, were genotyped using the full
catalog of variants obtained from all samples, including MEFs. Variant impact prediction and
annotation was carried out using SnpEff version 4.3T[107]. WGS and mRC-seq aligned BAMs were
processed to identify non-reference TE insertions using TEBreak[60]
(https://github.com/adamewing/tebreak) as previously described[61].

TE insertion PCR validation experiments

Reads supporting putative *de novo* TE insertions were manually examined using Serial Cloner

(http://serialbasics.free.fr/Serial_Cloner.html), the UCSC Genome Browser BLAT tool[108] and the

Repbase CENSOR tool[109]. PCR primers were designed with Primer3[110] against TE insertion

sequences and their 5′ and 3′ genomic flanks (**Supplementary Table 3**). Empty/filled PCRs

(combining 5′ and 3′ flanking primers) and full-length PCRs (using junction-spanning primers)

were performed using an Expand Long Range dNTPack (Roche, Cat#: 4829034001). Reaction

mixes contained 5μL 5× Expand Long Range Buffer with 12.5mM MgCl$_2$, 1.25μL dNTP Mix

(dATP, dCTP, dGTP, dTTP at 10mM each), 1.25μL DMSO (100%), 1μL primer mix (25μM of

each primer), 0.35μL Expand Long Range Enzyme Mix (5U/μL), 4-10ng genomic DNA template,

and molecular grade water up to a total volume of 25μL. PCR was performed with the following

cycling conditions: 92˚C for 3min, 10 cycles of 92˚C for 30s, 56-60˚C for 30s, and 68˚C for 7min

30s 25 cycles of 92˚C for 30s, 56-60˚C for 30s, and 68˚C for 7min + 20s cycle elongation for each

successive cycle, followed by 68˚C for 10min. TE-genome junction validation PCRs were

performed using MyTaq HS DNA Polymerase (Bioline, Cat#: BIO-2111). Reaction mixes

contained 5μL 5× MyTaq Reaction Buffer, 0.5μL primer mix (25μM of each primer), 0.2μL

MyTaq HS DNA Polymerase, 2-4ng genomic DNA template, and molecular grade water up to a

total volume of 25μL. PCRs were performed using the following conditions: 95˚C for 2min, 35

cycles of 95˚C for 15s, 55/57˚C for 15s, and 72˚C for 10s, followed by 72˚C for 10min. PCR

products were run on 0.8-2% agarose gels (Bioline, Cat#: BIO-41025), depending on fragment

size, pre-stained with SYBR Safe Nucleic Acid Gel Stain (Invitrogen, Cat#: S33102). A Typhoon

FLA 9000 (GE Healthcare Life Sciences) was used for gel imaging. Gel fragments were excised

under a Safe Imager 2.0 Blue-Light Transilluminator (Invitrogen). DNA purification was

performed using the QIAquick Gel Extraction Kit (Qiagen, Cat#: 28706) or MinElute Gel

Extraction Kit (Qiagen, Cat#: 28606) according to the manufacturer's instructions. PCR fragments

were either sequenced directly or cloned using the pGEM-T Easy Vector System (Promega, Cat#:

A1360) and Sanger sequenced to resolve insertion characteristics, as shown in **Supplementary**

Table 3.

L1-mCherry retrotransposition assays

The L1-mCherry construct was derived from the construct pTN201, a pCEP4-based vector

containing the native mouse element L1$_{spa}$[22]. The L1$_{spa}$ coding sequence was modified by site-

directed mutagenesis to include two nonsynonymous nucleotide substitutions, rendering the

ORF1p amino acid sequence identical to that of the L1 T$_F$ subfamily consensus sequence[111,112].

The 3′UTR was interrupted by a reporter cassette based on previously described L1

retrotransposition indicator plasmids[19,113]. This reporter cassette consisted of the mCherry coding

sequence in antisense orientation to the L1 and was equipped with an EF1α promoter and HSVtk

polyadenylation signal. The mCherry ORF was interrupted by a β-globin intron oriented in sense

to the L1. The mCherry cassette was cloned using G-block double-stranded DNA fragments

synthesized by Integrated DNA Technologies (IDT) and PCR products generated using Q5 DNA

polymerase (New England Biolabs, Cat#: M0492). The mCherry coding sequence was synthesized

with silent mutations ablating potential splice donor and splice acceptor sites that could interfere

with intended splicing of the intron. In the L1-mCherry construct, the final 157bp of the L1$_{spa}$

3′UTR, which included a conserved poly-purine tract, was situated downstream of the mCherry

cassette and immediately upstream of the pCEP4 SV40 polyadenylation signal[112]. The L1-

mCherry_RT- mutant contained a missense mutation in the reverse transcriptase domain of ORF2

(D709Y)[22]. Plasmids were prepared using a Qiagen Plasmid Plus Midi Kit and a QIAvac vacuum

manifold (Qiagen, Cat#: 12145).

HeLa-JVM cells[19] were cultured at 37°C and 5% CO_2 in HeLa complete medium (DMEM,

Life Technologies, Cat#: 11960044) supplemented with 10% FBS (Life Technologies, Cat#:

10099141), 1% Glutamax (Life Technologies, Cat#: 35050061) and 1% penicillin-streptomycin

(Life Technologies, Cat#: 15140122). Cells were passaged at 70-80% confluency using 0.25%

Trypsin-EDTA (Life Technologies, Cat#: 25200072). Cultured cell retrotransposition assays were

then performed as described previously[67,113], except retrotransposition was detected by mCherry

fluorescence instead of EGFP fluorescence. Briefly, 1×10^5 HeLa-JVM cells were seeded per well

of a 6-well plate. Eighteen hours later, cells were transfected with 1µg L1-mCherry or L1-

mCherry_RT- plasmid per well using 3µL FuGENE HD transfection reagent (Promega, Cat#:

E2311) and 97µL Opti-MEM (Life Technologies, Cat#: 31985047) per well according to the

manufacturer's protocol. Twenty-four hours post-transfection, medium was replaced with either

HeLa complete medium with 200µg/mL Hygromycin (Life Technologies, Cat#: 10687010), or

HeLa complete medium with 200µg/mL Hygromycin and 100µM Lamivudine (Sigma-Aldrich,

Cat#: L1295-10MG). Medium was replaced every other day, and at 8 days post-transfection cells

were collected by trypsinization, resuspended in sterile PBS, and analyzed on a CytoFLEX flow

cytometer (Beckman Coulter) to determine the percentage of mCherry positive cells. Three

biological replicate assays were performed, each consisting of 3 assayed wells per condition

(technical replicates).

L1-mneoI retrotransposition assays

To prepare reporter constructs, miPSC_1_L1 and miPSC_4_L1 were amplified from genomic

DNA using an Expand Long Range dNTPack (Roche, Cat#: 4829034001). Reaction mixes

contained 5μL 5× Expand Long Range Buffer with 12.5mM MgCl$_2$, 1.25μL dNTP Mix (dATP,

dCTP, dGTP, dTTP at 10mM each), 1.25μL DMSO (100%), 1μL primer mix (50μM of each

primer), 0.35μL Expand Long Range Enzyme Mix (5U/μL), 10ng genomic DNA template and

molecular grade water, up to a total volume of 25μL. PCRs were performed with the following

cycling conditions: 92˚C for 3min, 10 cycles of 92˚C for 30s, 58˚C for 30s, and 68˚C for 7min 30s;

25 cycles of 92˚C for 30sec, 58˚C for 30s, and 68˚C for 7min plus 20s elongation for each

successive cycle, followed by 68˚C for 10min. Primers introduced a *NotI* restriction site at the L1

5′ end (miPSC_1_L1_F, 5′-tttgcggccgcagaaagggaataatcgaggtg-3′; miPSC_1_L1_R, 5′-

gctaagcttgagaataagtgaagga-3′; miPSC_4_L1_F, 5′-agggcggccgcaggattaagaacccaatcaccag-3′;

miPSC_4_L1_R, 5′-aaaatgcctgttgtgccaat-3′). Reactions were purified using agarose gel

electrophoresis. Target fragments were excised and purified using either traditional phenol-

chloroform extraction or QIAquick and MinElute Gel Extraction Kits (Qiagen, Cat#: 28706 and

28604). Each L1 was then cloned into pGEMT Easy Vector (Promega, Cat#: A1360). Ligations

were incubated overnight at 4˚C. Ligation reactions were transformed using One Shot TOP10

chemically competent *E. coli* (Invitrogen, Cat#: C404010). Blue/white screening was performed

using LB/ampicillin/IPTG/X-Gal plates. At least 3 positive colonies per L1 were chosen for

Miniprep culture and plasmid DNA was isolated using a QIAprep Spin Miniprep Kit (Qiagen,

Cat#: 27106). At least three clones per element were capillary sequenced and compared to identify

PCR-induced mutations. Full-length L1s were then reconstructed by combining PCR-mutation

free fragments from different clones using restriction enzymes (New England Biolabs) recognizing

the L1 sequence. Reactions were purified using agarose gel electrophoresis and target fragments

were excised and purified using QIAquick and MinElute Gel Extraction Kits (Qiagen, Cat#: 28706
and 28604).

pTN201 was used to generate L1 reporter constructs. pTN201 is composed of a pCEP4
backbone (Life Technologies) containing $L1_{spa}$, a retrotransposition-competent L1 T_F[22] and a
downstream mneoI retrotransposition reporter cassette[114]. The mneoI cassette is driven by an SV40
promoter and holds the neomycin resistance gene, which is interrupted by an intron and is
positioned antisense to L1spa. In this assay, neomycin (or its analog, Geneticin/G418) resistance
only occurs via transcription, splicing and integration of the L1 and mneoI cassette into genomic
DNA[19,67]. To measure miPSC_1_L1 and miPSC_4_L1 retrotransposition efficiency, $L1_{spa}$ was
removed from the pCEP4 backbone by digesting with *NotI* and *PacI*. The pCEP4 backbone was
dephosphorylated using Calf Intestinal Alkaline Phosphatase (CIP) (New England Biolabs, Cat#:
M0290). The backbone and fragments of either miPSC_1_L1 or miPSC_4_L1 were combined in
a single ligation reaction using T4 DNA Ligase (New England Biolabs, Cat#: M0202) and
incubated overnight at 16˚C. Ligations were transformed using One Shot TOP10 chemically
competent *E. coli* (Invitrogen, Cat#: C404010) and plasmid DNA of positive clones was obtained
using QIAprep Spin Miniprep Kit (Qiagen, Cat#: 27106). Clones were verified as mutation-free
by capillary sequencing. Plasmid DNA for retrotransposition assays was obtained using a Plasmid
Maxi Kit (Qiagen, Cat#: 12163). Each construct was built with and without a cytomegalovirus
promoter (CMVp) preceding the L1. In addition, the following controls, each based on a pCEP4
backbone containing the mneoI cassette, were employed: TGF21, a retrotransposition-competent
L1 G_F[21]; L1SM, a synthetic codon optimized mouse L1[68]; L1SMmut2, L1SM immobilized by
reverse transcriptase and endonuclease domains mutations[68].

Retrotransposition assays were performed as previously described[67], with minor
modifications. HeLa-JVM cells were grown in HeLa complete medium (DMEM, Life
Technologies, Cat#: 11960044) supplemented with 10% FBS (Life Technologies, Cat#:
10099141), 1% Glutamax (Life Technologies, Cat#: 35050061) and 1% penicillin-streptomycin
(Life Technologies, Cat#: 15140122), and then seeded at a density of 4×10^4 cells/well in 6-well
tissue culture plates. 14-16h after plating, cells were transfected with L1 reporter constructs using
4μL FuGENE HD transfection reagent (Promega, Cat#: E2311) 96μL Opti-MEM (Life
Technologies, Cat#: 31985047) and 1μg plasmid DNA per well. Transfection efficiencies were

determined in parallel by preparing transfection mixes containing 4μL FuGENE HD transfection

reagent (Promega, Cat#: E2311), 96μL Opti-MEM (Life Technologies, Cat#: 31985047), 0.5μg

L1 expression plasmid and 0.5μg pCEP4-eGFP. The transfection mixture was added to each well

containing 2mL DMEM-complete medium. Plates were incubated at 37°C and 5% CO_2, medium

replaced 24h post-transfection, and transfection efficiency determined 72h post-transfection.

pCEP4-eGFP transfected wells were trypsinized and cells were collected from each well and

centrifuged at 2000g for 5min. Cell pellets were resuspended in 300-500μL 1× PBS. The number

of eGFP-positive cells was determined using a CytoFLEX flow cytometer (Beckman Coulter). The

percentage of eGFP-positive cells was used to normalize the G418-resistant colony counts for each

L1 reporter construct[67]. G418 (400μg/mL) (Thermo Fisher Scientific, Cat#: 10131035) selection

was started 3 days post-transfection and performed for 12 days. G418 foci were washed with 1×

PBS and fixed using 2% Formaldehyde/0.2% Glutaraldehyde in 1× PBS (Sigma-Aldrich) fixing

solution at room temperature for 30min. Staining was done using 0.1% Crystal Violet solution

(Sigma-Aldrich) at room temperature for 10min. Foci were counted in each well to quantify

retrotransposition.

L1 bisulfite sequencing experiments

Bisulfite conversion was performed with 200ng input genomic DNA from miPSC lines and MEFs

using a EZ DNA Methylation-Lightning Kit (Zymo Research, Cat#: D5030), following the

manufacturer's instructions. DNA was eluted in 10μL Elution Buffer. The internal sequences of

L1 T_F monomers were amplified genome-wide with the following primers: BS_TfIII_mono_F, 5′-

GGAAATTAGTTTGAATAGGTTAGAGGGTG;   BS_TfIII_mono_R,   5′-

TCCTAAATTCCAAAAAATCCTAAAACCAAA. The following locus-specific primers were

used to target the 5′ promoter region of the following elements of interest: BS_miPSC_1_L1_F,

5′-TGATTTATTTTTGATTGAATTTATTTTTAT;   BS_miPSC_1_L1_R/donor_L1_R,

5′-CTATTCAAACTAATTTCCTAAATTCTACTA;   BS_miPSC_3_L1_F,   5′-

TAGTTGGGGTTGTATGATGTAAGTT;   BS_miPSC_3_L1_R,   5′-

TCCCAAAAACTATCTAATTCTCTAAC;   BS_miPSC_4_L1_F,   5′-

TTTATATTGAAGGTTTGGATGATTTTATAT;   BS_miPSC_4_L1_R,   5′-

TCCAATTCTCTAATACACCCTCTAAC;   BS_donor_L1_F,   5′-

TTAAAGAAGTTAGTGATTTTTTAGAATTTT. PCRs were performed using MyTaq HS DNA

Polymerase (Bioline, Cat#: BIO-21111). Reaction mixes contained 5μL 5× MyTaq Reaction

Buffer, 0.5μL primer mix (25μM of each primer), 0.2μL MyTaq HS DNA Polymerase, DMSO at

a final concentration of 0.1%, 2μL bisulfite converted DNA template, and molecular grade water

up to a total volume of 25μL. PCR cycling parameters were as follows: 95˚C for 2min, 40 cycles

of 95˚C for 30s, 54˚C for 30s, and 72˚C for 30s, followed by 72˚C for 5min. PCR products were

run on a 2% agarose gel, excised and purified using a MinElute Gel Extraction Kit (Qiagen, Cat#:

28604) according to the manufacturer's instructions. Illumina libraries were constructed using a

NEBNext Ultra[TM] II DNA Library Prep Kit (New England Biolabs, Cat#: E7645). Libraries were

quantified using a Bioanalyzer DNA 1000 chip (Agilent Technologies, Cat#: 5067-1504).

Barcoded libraries were pooled in equimolar amounts and sequenced as 2x300mer reads on an

Illumina MiSeq platform using a MiSeq Reagent Kit v3 (Illumina, Cat#: MS-102-3003). 50% PhiX

Control v3 (Illumina, Cat#: FC-110-3001) was used as a spike-in. Sequencing data were analyzed

as described previously[34]. To summarize, for the L1 T_F genome-wide analysis, paired-end reads

were considered separately and those with the L1 T_F bisulfite PCR primers at their termini were

retained and aligned to the mock converted T_F monomer target amplicon sequence with blastn.

Reads where non-CpG cytosine bisulfite conversion was <95%, or ≥5% of CpG dinucleotides were

mutated, or ≥5% of adenine and guanine nucleotides were mutated, were removed. 50 reads per

sample, excluding identical bisulfite sequences, were randomly selected and analyzed using

QUMA[115] with default parameters, with strict CpG recognition. Specific L1 loci were analyzed in

a similar fashion, except paired-end reads were assembled into contigs, as described elsewhere[34],

prior to blastn alignment to the mock converted L1 locus target amplicon.

Nanopore sequencing analyses

Genomic DNA was extracted from 6 miPSC lines reprogrammed in the presence of 100μM 3TC,

6 miPSC lines generated without 3TC, and the parental MEFs, with a Nanobind CBB Big DNA

Kit (Circulomics, Cat#: NB-900-001-01) according to the manufacturer's instructions. DNA

libraries were prepared at the Kinghorn Centre for Clinical Genomics (KCCG, Australia) using

3μg input DNA, without shearing, and an SQK-LSK110 ligation sequencing kit. Libraries were

each sequenced separately on a PromethION (Oxford Nanopore Technologies) flow cell (FLO-

PRO002, R9.4.1 chemistry) (**Supplementary Table 1**). Bases were called with guppy 5.0.13

(Oxford Nanopore Technologies).

Non-reference TE insertions were detected with TLDR[35]. Briefly, this involved aligning

ONT reads to the mm10 reference genome using minimap2[116] version 2.20 (index parameter: -x

map-ont; alignment parameters: -ax map-ont -L -t 32) and SAMtools[117] version 1.12. BAM files

were then processed as a group with TLDR[35] version 1.2.2 (parameters -e teref.mouse.fa -p 128 -

m 1 -r mm10.fa -n nonref.collection.mm10.chr.bed.gz --keep_pickles). The files teref.mouse.fa,

composed of TE family consensus sequences, and nonref.collection.mm10.chr.bed.gz, a collection

of known non-reference retrotransposon insertions, are available from

github.com/adamewing/tldr/. The TLDR output table was further processed to remove calls not

passing relevant TLDR filters, where family = "NA" or remappable = "FALSE" or UnmapCover

< 0.5 or LengthIns < 100 or EndTE-StartTE < 100 or strand = "None" or SpanReads < 1. 3′

truncated TE insertions, and B1 or B2 insertions 5′ truncated by more than 2bp, were removed.

Events detected in only one miPSC line and not matching a known non-reference insertion were

designated as putative *de novo* insertions (**Supplementary Table 3**).

Reference TE methylation was assessed for parental MEFs and miPSC lines aggregated by

condition (reprogrammed with or without 3TC), using Methylartist version 1.2.4[82]. Briefly, CpG

methylation calls were generated from ONT reads using nanopolish version 0.13.2[118]. Using

Methylartist commands db-nanopolish, segmeth and segplot with default parameters, methylation

statistics were generated for the genome divided into 10kbp bins, protein-coding gene promoters

defined the Eukaryotic Promoter Database (-1000bp,+500bp)[119], and reference TEs defined by

RepeatMasker coordinates (http://www.repeatmasker.org/). TE families displayed in **Fig. 4a**

included T_F, G_F, and A-type L1s >6kbp, B1 and B2 SINEs, and MERVL MT2 and IAP elements

represented by their long terminal repeats. Methylation values were calculated for L1 5′UTRs only,

excluding the L1 body. Methylation profiles for individual loci were generated using the

Methylartist command locus. L1 T_F methylation profiles shown in **Fig. 4b** were generated for

elements >7kbp with the Methylartist command composite. To identify individual differentially

methylated TEs (**Supplementary Table 5**), we required elements to have at least 4 reads and 20

methylation calls in each of the MEF, aggregated control miPSC and aggregated 3TC-treated

miPSC datasets. Statistical comparisons were performed based on methylated and unmethylated

CpG call counts, using Fisher's exact test with Bonferroni correction for multiple testing.

L1-mCherry retrotransposition events were recovered via ONT sequencing in a similar

fashion to the analysis of mouse samples, except genomic DNA obtained from 5 HeLa cell lines

expanded over 3-5 passages from single L1-mCherry insertion-harboring colonies was barcoded

using a SQK-MLK111.96-XL kit, pooled in equimolar amounts, and sequenced on a single

PromethION flow cell. Reads were aligned to the hg38 human reference genome build. L1-

mCherry insertions were then identified using TLDR, as for the endogenous mouse TE analysis

described above.

Data availability

All Oxford Nanopore Technologies and Illumina sequencing data generated by this study were

deposited in the European Nucleotide Archive (ENA) under project PRJEB20569.

Code availability

TEBreak, TLDR and Methylartist, and instructions for their use, are available at

https://github.com/adamewing/tebreak,   https://github.com/adamewing/TLDR   and

https://github.com/adamewing/methylartist, respectively.

References

1.   Takahashi, K. & Yamanaka, S. Induction of pluripotent stem cells from mouse embryonic

and adult fibroblast cultures by defined factors. *Cell* **126**, 663–676 (2006).

2.   Mandai, M. *et al.* Autologous Induced Stem-Cell–Derived Retinal Cells for Macular

Degeneration. *N. Engl. J. Med.* **376**, 1038–1046 (2017).

3.   Takahashi, K. & Yamanaka, S. A decade of transcription factor-mediated reprogramming to

pluripotency. *Nat. Rev. Mol. Cell Biol.* **17**, 183–193 (2016).

4.   Tapia, N. & Schöler, H. R. Molecular Obstacles to Clinical Translation of iPSCs. *Cell Stem

Cell* **19**, 298–309 (2016).

5.   Gore, A. *et al.* Somatic coding mutations in human induced pluripotent stem cells. *Nature*

471, 63–67 (2011).

6.   Hussein, S. M. *et al.* Copy number variation and selection during reprogramming to

pluripotency. *Nature* **471**, 58–62 (2011).

7.   Lister, R. *et al.* Hotspots of aberrant epigenomic reprogramming in human induced

pluripotent stem cells. *Nature* **471**, 68–73 (2011).

8.   Polo, J. M. *et al.* Cell type of origin influences the molecular and functional properties of

mouse induced pluripotent stem cells. *Nat. Biotechnol.* **28**, 848–855 (2010).

9.   Laurent, L. C. *et al.* Dynamic changes in the copy number of pluripotency and cell

proliferation genes in human ESCs and iPSCs during reprogramming and time in culture.

Cell Stem Cell **8**, 106–118 (2011).

10.   Young, M. A. *et al.* Background mutations in parental cells account for most of the genetic

heterogeneity of induced pluripotent stem cells. *Cell Stem Cell* **10**, 570–582 (2012).

11.   Cheng, L. *et al.* Low incidence of DNA sequence variation in human induced pluripotent

stem cells generated by nonintegrating plasmid expression. *Cell Stem Cell* **10**, 337–344

(2012).

12.   Quinlan, A. R. *et al.* Genome sequencing of mouse induced pluripotent stem cells reveals

retroelement stability and infrequent DNA rearrangement during reprogramming. *Cell Stem*

Cell **9**, 366–373 (2011).

13.   Abyzov, A. *et al.* Somatic copy number mosaicism in human skin revealed by induced

pluripotent stem cells. *Nature* **492**, 438–442 (2012).

14.   Kazazian, H. H., Jr & Moran, J. V. Mobile DNA in Health and Disease. *N. Engl. J. Med.*

377, 361–370 (2017).

15. Doucet, A. J., Wilusz, J. E., Miyoshi, T., Liu, Y. & Moran, J. V. A 3' Poly(A) Tract Is

Required for LINE-1 Retrotransposition. *Mol. Cell* **60**, 728–741 (2015).

16. Jurka, J. Sequence patterns indicate an enzymatic involvement in integration of mammalian

retroposons. *Proc. Natl. Acad. Sci. U. S. A.* **94**, 1872–1877 (1997).

17. Luan, D. D., Korman, M. H., Jakubczak, J. L. & Eickbush, T. H. Reverse transcription of

R2Bm RNA is primed by a nick at the chromosomal target site: a mechanism for non-LTR

retrotransposition. *Cell* **72**, 595–605 (1993).

18. Monot, C. *et al.* The specificity and flexibility of l1 reverse transcription priming at

imperfect T-tracts. *PLoS Genet.* **9**, e1003499 (2013).

19. Moran, J. V. *et al.* High frequency retrotransposition in cultured mammalian cells. *Cell* **87**,

917–927 (1996).

20. Wei, W. *et al.* Human L1 retrotransposition: cis preference versus trans complementation.

Mol. Cell. Biol. **21**, 1429–1439 (2001).

21. Goodier, J. L., Ostertag, E. M., Du, K. & Kazazian, H. H., Jr. A novel active L1

retrotransposon subfamily in the mouse. *Genome Res.* **11**, 1677–1685 (2001).

22. Naas, T. P. *et al.* An actively retrotransposing, novel subfamily of mouse L1 elements.

EMBO J. **17**, 590–597 (1998).

23. Nellåker, C. *et al.* The genomic landscape shaped by selection on transposable elements

across 18 mouse strains. *Genome Biol.* **13**, R45 (2012).

24. Skowronski, J., Fanning, T. G. & Singer, M. F. Unit-length line-1 transcripts in human

teratocarcinoma cells. *Mol. Cell. Biol.* **8**, 1385–1397 (1988).

25. Beck, C. R. *et al.* LINE-1 retrotransposition activity in human genomes. *Cell* **141**, 1159–

1170 (2010).

26. Brouha, B. *et al.* Hot L1s account for the bulk of retrotransposition in the human
population. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 5280–5285 (2003).

27. Ewing, A. D. & Kazazian, H. H., Jr. High-throughput sequencing reveals extensive
variation in human-specific L1 content in individual human genomes. *Genome Res.* **20**,
1262–1270 (2010).

28. Richardson, S. R. *et al.* Heritable L1 retrotransposition in the mouse primordial germline
and early embryo. *Genome Res.* **27**, 1395–1405 (2017).

29. Stewart, C. *et al.* A comprehensive map of mobile element insertion polymorphisms in
humans. *PLoS Genet.* **7**, e1002236 (2011).

30. Feusier, J. *et al.* Pedigree-based estimation of human mobile element retrotransposition
rates. *Genome Res.* **29**, 1567–1577 (2019).

31. de la Rica, L. *et al.* TET-dependent regulation of retrotransposable elements in mouse
embryonic stem cells. *Genome Biol.* **17**, 234 (2016).

32. Warkocki, Z. *et al.* Uridylation by TUT4/7 Restricts Retrotransposition of Human LINE-1s.
Cell **174**, 1537–1548.e29 (2018).

33. Deniz, Ö., Frost, J. M. & Branco, M. R. Regulation of transposable elements by DNA
modifications. *Nat. Rev. Genet.* **20**, 417–431 (2019).

34. Sanchez-Luque, F. J. *et al.* LINE-1 Evasion of Epigenetic Repression in Humans. *Mol. Cell*
75, 590–604 (2019).

35. Ewing, A. D. *et al.* Nanopore Sequencing Enables Comprehensive Transposable Element
Epigenomic Profiling. *Mol. Cell* **80**, 915–928.e5 (2020).

36. Bourc'his, D. & Bestor, T. H. Meiotic catastrophe and retrotransposon reactivation in male
germ cells lacking Dnmt3L. *Nature* **431**, 96–99 (2004).

37.   Walter, M., Teissandier, A., Pérez-Palacios, R. & Bourc'his, D. An epigenetic switch

ensures transposon repression upon dynamic loss of DNA methylation in embryonic stem

cells. *Elife* **5**, e11418 (2016).

38.   Greenberg, M. V. C. & Bourc'his, D. The diverse roles of DNA methylation in mammalian

development and disease. *Nat. Rev. Mol. Cell Biol.* **20**, 590–607 (2019).

39.   Goodier, J. L. Restricting retrotransposons: a review. *Mob. DNA* **7**, 16 (2016).

40.   Chelmicki, T. *et al.* m6A RNA methylation regulates the fate of endogenous retroviruses.

Nature **591**, 312–316 (2021).

41.   MacLennan, M. *et al.* Mobilization of LINE-1 retrotransposons is restricted by Tex19.1 in

mouse embryonic stem cells. *Elife* **6**, e26152 (2017).

42.   Salvador-Palomeque, C. *et al.* Dynamic Methylation of an L1 Transduction Family during

Reprogramming and Neurodifferentiation. *Mol. Cell. Biol.* **39**, e00499–18 (2019).

43.   Lee, D.-S. *et al.* An epigenomic roadmap to induced pluripotency reveals DNA methylation

as a reprogramming modulator. *Nat. Commun.* **5**, 5619 (2014).

44.   Knaupp, A. S. *et al.* Transient and Permanent Reconfiguration of Chromatin and

Transcription Factor Occupancy Drive Reprogramming. *Cell Stem Cell* **21**, 834–845.e6

(2017).

45.   Klawitter, S. *et al.* Reprogramming triggers endogenous L1 and Alu retrotransposition in

human induced pluripotent stem cells. *Nat. Commun.* **7**, e10286 (2016).

46.   Friedli, M. *et al.* Loss of transcriptional control over endogenous retroelements during

reprogramming to pluripotency. *Genome Res.* **24**, 1251–1259 (2014).

47.   Maherali, N. *et al.* Directly reprogrammed fibroblasts show global epigenetic remodeling

and widespread tissue contribution. *Cell Stem Cell* **1**, 55–70 (2007).

48.   Gerdes, P. *et al.* Mouse L1s fade with age: a methylation-enforced mechanism for

attenuation of L1 retrotransposition potential. *bioRxiv* (2022)

doi:10.1101/2022.08.06.500894.

49.   Habibi, E. *et al.* Whole-genome bisulfite sequencing of two distinct interconvertible DNA

methylomes of mouse embryonic stem cells. *Cell Stem Cell* **13**, 360–369 (2013).

50.   Garcia-Perez, J. L. *et al.* LINE-1 retrotransposition in human embryonic stem cells. *Hum.*

Mol. Genet. **16**, 1569–1577 (2007).

51.   Macia, A. *et al.* Engineered LINE-1 retrotransposition in nondividing human neurons.

Genome Res. **27**, 335–348 (2017).

52.   Wissing, S. *et al.* Reprogramming somatic cells into iPS cells activates LINE-1

retroelement mobility. *Hum. Mol. Genet.* **21**, 208–218 (2012).

53.   Stadtfeld, M., Maherali, N., Borkent, M. & Hochedlinger, K. A reprogrammable mouse

strain from gene-targeted embryonic stem cells. *Nat. Methods* **7**, 53–55 (2010).

54.   McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing

next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).

55.   Garrison, E. & Marth, G. Haplotype-based variant detection from short-read sequencing.

arXiv [q-bio.GN] arXiv:1207.3907 (2012).

56.   Keane, T. M. *et al.* Mouse genomic variation and its effect on phenotypes and gene

regulation. *Nature* **477**, 289–294 (2011).

57.   Rausch, T. *et al.* DELLY: structural variant discovery by integrated paired-end and split-

read analysis. *Bioinformatics* **28**, i333–i339 (2012).

58.   Cameron, D. L. *et al.* GRIDSS: sensitive and specific genomic rearrangement detection

using positional de Bruijn graph assembly. *Genome Res.* **27**, 2050–2060 (2017).

59. Ewing, A. D. Transposable element detection from whole genome sequence data. *Mob.*

DNA **6**, 24 (2015).

60. Carreira, P. E. *et al.* Evidence for L1-associated DNA rearrangements and negligible L1

retrotransposition in glioblastoma multiforme. *Mob. DNA* **7**, 21 (2016).

61. Schauer, S. N. *et al.* L1 retrotransposition is a common feature of mammalian

hepatocarcinogenesis. *Genome Res.* **28**, 639–653 (2018).

62. Ostertag, E. M. & Kazazian, H. H., Jr. Twin priming: a proposed mechanism for the

creation of inversions in L1 retrotransposition. *Genome Res.* **11**, 2059–2065 (2001).

63. Kano, H. *et al.* L1 retrotransposition occurs mainly in embryogenesis and creates somatic

mosaicism. *Genes Dev.* **23**, 1303–1312 (2009).

64. van den Hurk, J. A. J. M. *et al.* L1 retrotransposition can occur early in human embryonic

development. *Hum. Mol. Genet.* **16**, 1587–1592 (2007).

65. Sultana, T. *et al.* The Landscape of L1 Retrotransposons in the Human Genome Is Shaped

by Pre-insertion Sequence Biases and Post-insertion Selection. *Mol. Cell* **74**, 555–570

(2019).

66. Flasch, D. A. *et al.* Genome-wide de novo L1 Retrotransposition Connects Endonuclease

Activity with Replication. *Cell* **177**, 837–851 (2019).

67. Kopera, H. C. *et al.* LINE-1 cultured cell retrotransposition assay. *Methods Mol. Biol.* **1400**,

139–156 (2016).

68. Han, J. S. & Boeke, J. D. A highly active synthetic mammalian retrotransposon. *Nature*

429, 314–318 (2004).

69. Meyer, T. J., Srikanta, D., Conlin, E. M. & Batzer, M. A. Heads or tails: L1 insertion-

associated 5' homopolymeric sequences. *Mob. DNA* **1**, 7 (2010).

70. Miki, Y. *et al.* Disruption of the APC gene by a retrotransposal insertion of L1 sequence in

a colon cancer. *Cancer Res.* **52**, 643–645 (1992).

71. Goodier, J. L., Ostertag, E. M. & Kazazian, H. H., Jr. Transduction of 3'-flanking sequences

is common in L1 retrotransposition. *Hum. Mol. Genet.* **9**, 653–657 (2000).

72. Pickeral, O. K., Makałowski, W., Boguski, M. S. & Boeke, J. D. Frequent human genomic

DNA transduction driven by LINE-1 retrotransposition. *Genome Res.* **10**, 411–415 (2000).

73. Moran, J. V., DeBerardinis, R. J. & Kazazian, H. H., Jr. Exon shuffling by L1

retrotransposition. *Science* **283**, 1530–1534 (1999).

74. Holmes, S. E., Dombroski, B. A., Krebs, C. M., Boehm, C. D. & Kazazian, H. H., Jr. A new

retrotransposable human L1 element from the LRE2 locus on chromosome 1q produces a

chimaeric insertion. *Nat. Genet.* **7**, 143–148 (1994).

75. Banuelos-Sanchez, G. *et al.* Synthesis and Characterization of Specific Reverse

Transcriptase Inhibitors for Mammalian LINE-1 Retrotransposons. *Cell Chem Biol* **26**,

1095–1109.e14 (2019).

76. Dai, L., Huang, Q. & Boeke, J. D. Effect of reverse transcriptase inhibitors on LINE-1 and

Ty1 reverse transcriptase activities and on LINE-1 retrotransposition. *BMC Biochem.* **12**, 18

(2011).

77. Smits, N. *et al.* No evidence of human genome integration of SARS-CoV-2 found by long-

read DNA sequencing. *Cell Rep.* **36**, 109530 (2021).

78. Lanciano, S. & Cristofari, G. Measuring and interpreting transposable element expression.

Nat. Rev. Genet. **21**, 721–736 (2020).

79. Zhou, W. *et al.* Identification and characterization of occult human-specific LINE-1

insertions using long-read sequencing technology. *Nucleic Acids Res.* **48**, 1146–1163

(2020).

80. Wolf, G. *et al.* KRAB-zinc finger protein gene expansion in response to active

retrotransposons in the murine lineage. *Elife* **9**, e56337 (2020).

81. Heidmann, O. & Heidmann, T. Retrotransposition of a mouse IAP sequence tagged with an

indicator gene. *Cell* **64**, 159–170 (1991).

82. Cheetham, S. W., Kindlova, M. & Ewing, A. D. Methylartist: Tools for Visualising

Modified Bases from Nanopore Sequence Data. *Bioinformatics* **38**, 3109–3112 (2022).

83. Scott, E. C. *et al.* A hot L1 retrotransposon evades somatic repression and initiates human

colorectal cancer. *Genome Res.* **26**, 745–755 (2016).

84. Shakiba, N. *et al.* Cell competition during reprogramming gives rise to dominant clones.

Science **364**, eaan0925 (2019).

85. Saini, N. *et al.* The Impact of Environmental and Endogenous Damage on Somatic

Mutation Load in Human Skin Fibroblasts. *PLoS Genet.* **12**, e1006385 (2016).

86. Kuijk, E. *et al.* The mutational impact of culturing human pluripotent and adult stem cells.

Nat. Commun. **11**, 2493 (2020).

87. Nefzger, C. M., Alaei, S., Knaupp, A. S., Holmes, M. L. & Polo, J. M. Cell surface marker

mediated purification of iPS cell intermediates from a reprogrammable mouse model. *J. Vis.*

Exp. e51728 (2014).

88. Utikal, J. *et al.* Immortalization eliminates a roadblock during cellular reprogramming into

iPS cells. *Nature* **460**, 1145–1148 (2009).

89. Macia, A. *et al.* Epigenetic control of retrotransposon expression in human embryonic stem

cells. *Mol. Cell. Biol.* **31**, 300–316 (2011).

90. Grandi, F. C. *et al.* Retrotransposition creates sloping shores: a graded influence of

hypomethylated CpG islands on flanking CpG sites. *Genome Res.* **25**, 1135–1146 (2015).

91. Dewannieux, M. & Heidmann, T. L1-mediated retrotransposition of murine B1 and B2

SINEs recapitulated in cultured cells. *J. Mol. Biol.* **349**, 241–247 (2005).

92. Dewannieux, M., Esnault, C. & Heidmann, T. LINE-mediated retrotransposition of marked

Alu sequences. *Nat. Genet.* **35**, 41–48 (2003).

93. Kazazian, H. H., Jr *et al.* Haemophilia A resulting from de novo insertion of L1 sequences

represents a novel mechanism for mutation in man. *Nature* **332**, 164–166 (1988).

94. International Human Genome Sequencing Consortium. Initial sequencing and analysis of

the human genome. *Nature* **409**, 860–921 (2001).

95. Han, J. S., Szak, S. T. & Boeke, J. D. Transcriptional disruption by the L1 retrotransposon

and implications for mammalian transcriptomes. *Nature* **429**, 268–274 (2004).

96. Faulkner, G. J. *et al.* The regulated retrotransposon transcriptome of mammalian cells. *Nat.*

Genet. **41**, 563–571 (2009).

97. Firas, J., Liu, X., Nefzger, C. M. & Polo, J. M. GM-CSF and MEF-conditioned media

support feeder-free reprogramming of mouse granulocytes to iPS cells. *Differentiation* **87**,

193–199 (2014).

98. Bertoncello, I. & McQualter, J. Isolation and clonal assay of adult lung epithelial

stem/progenitor cells. *Current Protocols in Stem Cell Biology* **16**, (2011).

99. Seach, N., Wong, K., Hammett, M., Boyd, R. L. & Chidgey, A. P. Purified enzymes

improve isolation and characterization of the adult thymic epithelium. *J. Immunol. Methods*

385, 23–34 (2012).

100. Merlos-Suárez, A. *et al.* The intestinal stem cell signature identifies colorectal cancer stem

cells and predicts disease relapse. *Cell Stem Cell* **8**, 511–524 (2011).

101. Jensen, K. B., Driskell, R. R. & Watt, F. M. Assaying proliferation and differentiation

capacity of stem cells using disaggregated adult mouse epidermis. *Nat. Protoc.* **5**, 898–911

(2010).

102. Alaei, S. *et al.* An improved reprogrammable mouse model harbouring the reverse

tetracycline-controlled transcriptional transactivator 3. *Stem Cell Res.* **17**, 49–53 (2016).

103. Lim, S. M. *et al.* Temporal restriction of pancreatic branching competence during

embryogenesis is mirrored in differentiating embryonic stem cells. *Stem Cells Dev.* **21**,

1662–1674 (2012).

104. Liu, K. *et al.* KSR-based medium improves the generation of high-quality mouse iPS cells.

PLoS One **9**, e105309 (2014).

105. Knaupp, A. S. *et al.* TINC—A Method to Dissect Regulatory Complexes at Single-Locus

Resolution—Reveals an Extensive Protein Complex at the Nanog Promoter. *Stem cell*

reports **15**, 1246–1259 (2020).

106. Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.

arXiv [q-bio.GN] arXiv:1303.3997 (2013).

107. Cingolani, P. *et al.* A program for annotating and predicting the effects of single nucleotide

polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118;

iso-2; iso-3. *Fly* **6**, 80–92 (2012).

108. Kent, W. J. *et al.* The human genome browser at UCSC. *Genome Res.* **12**, 996–1006 (2002).

109. Bao, W., Kojima, K. K. & Kohany, O. Repbase Update, a database of repetitive elements in

eukaryotic genomes. *Mob. DNA* **6**, 11 (2015).

110. Untergasser, A. *et al.* Primer3--new capabilities and interfaces. *Nucleic Acids Res.* **40**, e115

(2012).

111. Martin, S. L. *et al.* A single amino acid substitution in ORF1 dramatically decreases L1

retrotransposition and provides insight into nucleic acid chaperone activity. *Nucleic Acids*

Res. **36**, 5845–5854 (2008).

112. Richardson, S. R. *et al.* Revisiting the impact of synthetic ORF sequences on engineered

LINE-1 retrotransposition. *bioRxiv* (2022) doi:10.1101/2022.08.29.505632.

113. Ostertag, E. M., Prak, E. T., DeBerardinis, R. J., Moran, J. V. & Kazazian, H. H., Jr.

Determination of L1 retrotransposition kinetics in cultured cells. *Nucleic Acids Res.* **28**,

1418–1423 (2000).

114. Freeman, J. D., Goodchild, N. L. & Mager, D. L. A modified indicator gene for selection of

retrotransposition events in mammalian cells. *Biotechniques* **17**, 46, 48–9, 52 (1994).

115. Kumaki, Y., Oda, M. & Okano, M. QUMA: quantification tool for methylation analysis.

Nucleic Acids Res. **36**, W170–5 (2008).

116. Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–

3100 (2018).

117. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–

2079 (2009).

118. Simpson, J. T. *et al.* Detecting DNA cytosine methylation using nanopore sequencing. *Nat.*

Methods **14**, 407–410 (2017).

119. Dreos, R., Ambrosini, G., Groux, R., Cavin Périer, R. & Bucher, P. The eukaryotic

promoter database in its 30th year: focus on non-vertebrate organisms. *Nucleic Acids Res.*

45, D51–D55 (2017).

120. Crooks, G. E., Hon, G., Chandonia, J.-M. & Brenner, S. E. WebLogo: a sequence logo

generator. *Genome Res.* **14**, 1188–1190 (2004).

121. Robinson, J. T. *et al.* Integrative genomics viewer. *Nat. Biotechnol.* **29**, 24–26 (2011).

Acknowledgements

The authors thank Jef D. Boeke and John V. Moran for sharing L1SM plasmids and the HeLa-

JVM cell line, respectively. This study was supported by an Australian Government Research

Training Program (RTP) Scholarship and a Mater Research Frank Clair Scholarship awarded to

P.G., ARC Discovery Early Career Researcher Award (DE150101117), Discovery Project

(DP170101198) and Australian Department of Health Medical Frontiers Future Fund (MRFF)

(MRF1175457) grants awarded to A.D.E., NHMRC Project Grant (GNT1051117), ARC Future

Fellowship (FT180100674) and Sylvia and Charles Viertel Senior Medical Research Fellowship

funds awarded to J.M.P., NHMRC Investigator Grants (GNT1178460 to R.L., GNT1173476 to

S.R.R., GNT1173711 to G.J.F.), and CSL Centenary Fellowship and NHMRC Project Grant

(GNT1106206, GNT1125645, GNT1126393, GNT1138795) funding awarded to G.J.F.. A.D.E.,

S.R.R. and G.J.F. additionally acknowledge support from the Mater Foundation.

1103

Author contributions

1104

1105   P.G., S.M.L., M.R.L., D.C., F.J.S-L., L.W., A.L.C., C.J., A.S.K., P.E.C., C.M.N. and S.R.R.

1106   performed experiments. R.L., J.M.P. and G.J.F. provided resources. A.D.E. and G.J.F. performed

1107   bioinformatic analyses. P.G., S.M.L., S.R.R. and G.J.F. generated figures. P.G., S.R.R., and G.J.F.

1108   conceived the study. P.G., S.M.L., S.R.R., J.M.P, and G.J.F. designed and supervised experiments.

1109   G.J.F. wrote the manuscript and directed the study. All authors commented on the manuscript.

1110

Competing interests

1111

1112   The authors declare no competing interests.

 Supplementary Table Legends

 Supplementary Table 1: WGS and mRC-seq library statistics.

 Supplementary Table 2: Genomic variants private to individual bulk or single-cell miPSC
 lines, and their predicated impacts. Note: Variants (SNVs, INDELs, SVs) were called using
 Illumina WGS data only.

 Supplementary Table 3: TE insertion validation details. Four tabs provide information for *de*
 novo insertions detected via (i) Illumina sequencing of bulk miPSCs reprogrammed from various
 tissues, (ii) Illumina sequencing of single-cell miPSC clones derived from MEFs, (iii) ONT
 sequencing of bulk miPSCs derived from MEFs and (iv) ONT sequencing to detect L1-mCherry
 retrotransposition events in cultured HeLa cells. Illumina data were analyzed with TEBreak, while
 ONT data were analyzed with TLDR. In tabs (i), (ii) and (iii), Column 1 lists a universally unique
 identifier (UUID) for each insertion. Columns 2-6 show the insertion name used in other figures
 and tables, the chromosomal location including genomic 5′ and 3′ end positions and whether the
 insertion is intergenic or intragenic. For tabs (i) and (ii), columns 7-14 provide the 5′ and 3′ end
 orientations and whether these provisionally indicate a 5′ inversion, the TE family, the 5′ and 3′
 end nucleotide positions of each insertion relative to the TE consensus sequence, and the
 WGS/mRC-seq read counts for the 5′ and 3′ ends. Column 15 displays the TSD sequence and
 columns 16 and 17 respectively contain the 5′ and 3′ junction consensus sequences constructed
 from WGS/mRC-seq reads. Columns 18 and 19 contain the name of the samples where the TE
 insertion was detected at its 5′ and 3′ end, respectively, and the number of reads present for each
 sample. For tab (iii), columns 7-17 provide the orientation of the insertion, the TE family and
 subfamily, the 5′ and 3′ end nucleotide positions of each insertion relative to the TE consensus
 sequence, the length of the insertion, whether a 5′ inversion is present, the total number of ONT
 reads supporting the insertion, the total number of spanning reads supporting the insertion, the
 name of the samples where the TE insertion was detected and the number of reads present for each
 sample, and whether the TE insertion has previously been observed in genomic analysis of mouse
 strains. Columns 18-20 display the TSD sequence, the consensus sequencing constructed from

ONT reads, and any associated TLDR flags. Columns 21-23 indicate TSD sequences and other

TPRT hallmarks associated with TE insertions obtained by manual inspection of consensus

sequences. For tab (iv), Column 1 lists a universally unique identifier (UUID) for each insertion.

Columns 2-4 show the chromosomal location including genomic 5′ and 3′ end positions. Columns

5-13 provide the orientation of the insertion, the TE family (all L1-mCherry), the 5′ and 3′ end

nucleotide positions of each insertion relative to the spliced L1-mCherry sequence, the length of

the insertion, whether a 5′ inversion is present, the total number of ONT reads supporting the

insertion, the total number of spanning reads supporting the insertion, and the name of the HeLa

colony carrying the TE insertion along with the corresponding number of reads. Columns 14-16

display the TSD sequence, the consensus sequencing constructed from ONT reads, and any

associated TLDR flags. Columns 17-19 indicate TSD sequences and other TPRT hallmarks

obtained by manual inspection of consensus sequences. Finally, in tabs (i), (ii) and (iii), the last 10

columns comprise the validation PCR type, validation PCR primers and capillary sequencing reads

for each insertion, and any relevant notes.

1157

1158   **Supplementary Table 4: Non-reference TE insertions detected in MEFs.** Two tabs provide

1159   information for (i) 277 gold standard heterozygous insertions detected via Illumina sequencing

1160   and (ii) insertions detected via ONT sequencing. Each tab is arranged approximately as per

1161   **Supplementary Table 3**, except without PCR validation information. All listed insertions were

1162   detected in the parental MEFs.

1163

Supplementary Table 5: Reference genome TE methylation values. Tab (i) contains

information for TEs and tab (ii) contains information for gene promoters, as defined by the

Eukaryotic Promoter Database[119]. Columns are as follows. seg_chrom, seg_start, seg_end:

position of the reference TE or gene promoter. seg_name: TE subfamily or gene promoter name.

seg_strand: + or - orientation relative to the genome assembly. *_meth_calls: number of

methylated CpG calls for each sample. *_unmeth_calls: number of unmethylated CpG calls for

each sample. *_no_calls: number of ambiguous CpG calls for each sample (log likelihood ratio

between -2.5 and 2.5). *_methfrac: fraction of non-ambiguous calls indicating methylation.

*_readcount: number of reads aligned to the element. methfrac_difference: difference between the

indicated sample methylation percentages. uncorrected_pvalue: Fisher's exact test p-value from

comparison of methylation/demethylation counts between samples. fdr_corrected_pvalue: the

value in "uncorrected_pvalue" Bonferroni corrected for multiple testing. significant: whether the

corrected p-value is less than 0.01 and the absolute methylation difference (ΔmC) is >25%. In tab

(i), L1_TSS_support indicates whether the 5′ end of a spliced EST or mRNA obtained from the

UCSC Genome Browser coincides with the listed L1. Note: full-length L1 methylation values were

calculated for the given 5′UTR coordinates.

Table 1: *De novo* TE insertions detected and PCR validated in miPSC lines.

Insertion #	Subfamily	Location	Monomers	EN motif	TSD (bp)	PolyA (bp)	Origin	Detected by
miPSC_1_L1	T_F	1q	6.8	TCTT/AG	16	~125	Reprogramming	WGS + mRC-seq
miPSC_2_L1	T_F	10q	0	TTCT/GT	14	>100	Mosaic	WGS + mRC-seq
miPSC_3_L1	T_F	13q	5.8	ATTC/AA	15	~50	Reprogramming	WGS + mRC-seq
miPSC_4_L1	T_F	3q	5.3	TCTT/AA	13	~54	Reprogramming	WGS + mRC-seq
miPSC_5_L1	G_F	19q	2	TTAT/AT	14	~50	Reprogramming	WGS + mRC-seq
miPSC_6_L1	T_F	7q	0	TTTA/AA	17	~51	Reprogramming	WGS + mRC-seq
miPSC_7_L1	G_F	Xq	5	TCTT/AT	16	>80	Reprogramming	WGS + mRC-seq
miPSC_8_L1	T_F	19q	3.7	TTTC/AA	19	~24	Reprogramming	WGS + mRC-seq
miPSC_9_B2	B2	11q	NA	TCTT/AC	16	>60	Reprogramming	WGS + mRC-seq
miPSC_10_L1	T_F	12q	0	TTTT/GT	6	~36*	Reprogramming	WGS
miPSC_11_B2	B2	13q	NA	TTTT/GA	14	>73	Reprogramming	WGS + mRC-seq
miPSC_12_L1	T_F	13q	0	TCTT/AG	17	~97	Reprogramming	WGS + mRC-seq
miPSC_13_L1	A	14q	3	TTTC/AT	13	~46	Reprogramming	WGS + mRC-seq
miPSC_14_B2	B2	15q	NA	TTTT/AC	16	>66	Reprogramming	WGS + mRC-seq
miPSC_15_L1	G_F	2q	0	TTTC/AA	17	~28*	Reprogramming	WGS
miPSC_16_L1	T_F	2q	>3	TTTT/AA	16	>100	Reprogramming	WGS + mRC-seq
miPSC_17_L1	T_F	3q	>3	ACTT/AA	14	~45	Reprogramming	WGS + mRC-seq
miPSC_18_B1	B1	3q	NA	TTTT/AA	15	~30	Reprogramming	WGS + mRC-seq
miPSC_19_L1	T_F	3q	>3	GTTT/AT	15	>80	Reprogramming	WGS + mRC-seq
miPSC_21_L1	T_F	4q	0	TTTT/CA	17	>150	Reprogramming	WGS + mRC-seq
miPSC_22_B2	B2	6q	NA	TCTT/GA	15	~52	Reprogramming	WGS + mRC-seq
miPSC_23_B2	B2	9q	NA	TTTT/AT	16	~50	Mosaic	WGS + mRC-seq
miPSC_24_B2	B2	Xq	NA	TTTT/AA	15	>100	Reprogramming	WGS + mRC-seq
miPSC_26_L1	T_F	1q	>3	TCTT/AT	22	~58	Reprogramming	WGS + mRC-seq
miPSC_27_B2	B2	11q	NA	TTTC/AA	14	>60	Reprogramming	WGS + mRC-seq
miPSC_28_L1	T_F	13q	3.6	TCCT/AA	15	~93*	Reprogramming	mRC-seq
miPSC_29_L1	T_F	15q	0	TCTT/AA	16	>80	Reprogramming	WGS + mRC-seq
miPSC_30_L1	T_F	6q	>3	TCTT/AT	16	~72	Reprogramming	mRC-seq
miPSC_31_L1	T_F	7q	>3	TTTG/AC	15	~43	Reprogramming	mRC-seq
miPSC_32_L1	T_F	Xq	2	TCTT/AT	13	~37	Reprogramming	WGS + mRC-seq
miPSC_33_L1	G_F	Xq	>3	TTTT/AA	15	~47	Reprogramming	mRC-seq
miPSC_34_L1	T_F	8q	0	TCTT/AA	6	~36*	Reprogramming	WGS + mRC-seq
miPSC_35_L1	T_F	1q	0	TTTA/AA	15	~38	Reprogramming	WGS + mRC-seq
miPSC_36_L1	G_F	8q	0	ATGT/GA	6	~42	Reprogramming	WGS + mRC-seq
miPSC_37_L1	T_F	1q	1.2	TTTT/GT	14	~20	Reprogramming	WGS + mRC-seq
miPSC_38_L1	T_F	10q	0	TTCT/AA	15	~55	Reprogramming	WGS + mRC-seq
miPSC_39_L1	T_F	10q	0	TTTT/AA	8	>140*	Reprogramming	WGS + mRC-seq
miPSC_40_L1	T_F	11q	>3	TTTT/GA	14	>120	Reprogramming	WGS + mRC-seq
miPSC_41_L1	T_F	12q	2.6	TCTT/GC	16	~49	Reprogramming	mRC-seq
miPSC_42_B1	B1	14q	NA	TTCT/AA	15	>50	Reprogramming	mRC-seq
miPSC_43_L1	T_F	16q	>3	ATTT/AA	14	~42*	Mosaic	WGS + mRC-seq
miPSC_69_L1	T_F	13q	2	CTTT/AT	16	~61	Mosaic	WGS (ONT)
miPSC_87_L1	T_F	17q	0	TTTT/GT	16	~21	Reprogramming	WGS (ONT)

Insertions marked with an asterisk carry a 3′ transduction. miPSC_1_L1 - miPSC_8_L1 were detected in adult primary cell-derived bulk miPSCs, miPSC_9_B2 - miPSC_43_L1 in MEF-derived single-cell miPSC clones, and miPSC_69_L1 and miPSC_87_L1 in MEF-derived bulk miPSCs. Unless stated otherwise, insertions were detected by Illumina sequencing.

Extended Data Fig. 1: miPSC line genotypic relationships. a, Clustering of miPSC lines derived from 9 primary cell types isolated from 3 animals (A67, A82, A172), and 3 MEF genotypic controls. For each pairwise comparison, the Jaccard index (J) was calculated as the ratio of the union and intersection of SNP/INDEL variants called from WGS data and shared by the sample pair. Known SNPs/INDELs were removed and filtered as described in the Methods. J=1 (light color on key) indicates an identical variant profile between a sample pair, whereas J=0 (dark color on key) indicates no variants in common. Hierarchical clustering was performed using average linkage and a Euclidean distance metric via the seaborn clustermap function. **b,** As for panel (a), except for 9 single-cell clones derived from animal I222e2 MEFs and cultured in serum or 2i conditions.

Extended Data Fig. 2: Sequence characteristics of *de novo* L1 insertions detected in bulk tissue-derived miPSCs. For each of four insertions, the following information is provided: the chromosomal location; a filled site illustration indicating target site duplication (TSD) sequences in red, the number of promoter monomers (black triangles) if applicable, and 3′ polyA tract length (A_n/T_n); an empty site illustration depicting TSD sequence and first strand endonuclease (EN) cleavage motif (underlined); summary characteristics (L1 subfamily, number of monomers, TSD length and EN motif).

Chr7

miPSC_31_L1$_{TF}$

A$_{43}$ ORF2 ORF1

Clone 18

mw Serum 2i MEFs Clone 16 (-) C57 (-) H$_2$O

7000

ChrX

miPSC_32_L1$_{TF}$

A$_{37}$ ORF2 ORF1

Clone 2

mw Serum 2i MEFs Clone 16 (-) C57 (-) H$_2$O

7000

Chr8

miPSC_34_L1$_{TF}$

A$_9$ 104 nt A$_{36}$

Clone 4

mw Serum 2i MEFs Clone 16 (-) C57 (-) H$_2$O

1000
500
200

Chr6

$_{15}$A 104 nt A$_{11}$ ORF2 ORF1

retrotransposition

Chr18

$_9$A ORF2 ORF1

?

?

Chr1

Pard3b

miPSC_35_L1$_{TF}$

A$_{38}$

Clone 4

mw Serum 2i MEFs Clone 16 (-) C57 (-) H$_2$O

1500

Chr8

Tbc1d9

miPSC_36_L1$_{GF}$

A$_{42}$

Clone 4

mw Serum 2i MEFs Clone 16 (-) C57 (-) H$_2$O

500
300

Chr1

miPSC_37_L1$_{TF}$

ORF1 ORF2 A$_{20}$

Clone 2

mw Serum 2i MEFs Clone 16 (-) C57 (-) H$_2$O

7000

700
500

Chr10

miPSC_38_L1$_{TF}$

A$_{55}$

Clone 7

mw Serum 2i MEFs Clone 16 (-) C57 (-) H$_2$O

1500

Chr10

miPSC_39_L1$_{TF}$

A$_{22}$ 11 nt A$_{>140}$

Clone 6

mw Serum 2i MEFs Clone 16 (-) C57 (-) H$_2$O

1000

Chr11

miPSC_40_L1$_{TF}$

ORF1 ORF2 A$_{>120}$

Clone 2

mw Serum 2i MEFs Clone 16 (-) C57 (-) H$_2$O

7000

Chr12

Gm31508

miPSC_41_L1$_{TF}$

A$_{49}$ ORF2 ORF1

Serum (Clone #)

mw 1 2 3 4 5 6 7 8 9 10 11 12 mw

7000

Serum (Clone #) 2i (Clone #) MEFs C57 (-) H$_2$O

mw 13 14 15 16 17 18 13 14 mw

7000

Extended Data Fig. 3: Additional *de novo* TE insertion validation and characterization. *De novo* TE insertions found in 26 bulk miPSC lines generated from primary cells or 18 single-cell miPSC clones derived from MEFs, each subjected to Illumina sequencing. For each insertion, the chromosomal location and orientation are shown. L1 and SINE B1 and B2 insertions are represented by white rectangles. L1 5′UTR promoter monomers, if present, are indicated by triangles or, if the number of monomers is unknown, a gray box with black stripes. Poly(A) tracts and their length are indicated (A_n), and target site duplications (TSDs) are depicted as gray arrows. 3′ transductions are shown as orange lines. PCR validation primers are shown as red arrows. PCR products in agarose gels used to confirm TE insertions are indicated by red arrows. Empty site (wild-type) amplicons are indicated by blue arrows, where applicable.

Extended Data Fig. 4: TE detection sensitivities at simulated sequencing depths. a, To assess whether PCR validated *de novo* TE insertions would have been initially overlooked by lower coverage WGS, we down sampled our ~41× average depth WGS in percentile increments. In order to be called as present, *de novo* insertions found in the bulk (*top*) and single-cell (*bottom*) miPSC experiments required ≥1 WGS read at each of their 5′ and 3′ junctions, and ≥10 WGS reads in total. **b,** To estimate the likelihood of a mosaic TE insertion being overlooked in the parental animal I222e2 MEF population, and called as *de novo* in one of the associated clonal miPSC lines, we defined a set of 277 heterozygous germline TE insertions found in I222e2 and that were detected by ≥25 WGS or mRC-seq reads at each of their 5′ and 3′ junctions. We then simulated the probability of at least one read being found for an insertion when the reads assigned to that insertion were assigned probabilities to achieve random sampling depths ranging from 0.01% to 100% of the parental MEF bulk sequencing data. Note: at each depth in panels (a) and (b), simulations were repeated 10,000 times.

Extended Data Fig. 5. Donor L1 hypomethylation in MEFs and miPSCs. a, *top left*: Locus-specific methylation analysis design for a donor L1 found to generate insertion miPSC_10_L1 in a MEF-derived single-cell miPSC clone (Clone 1). CpGs located in the first 3 monomers of the donor L1 were assessed. Orange and gray strokes indicate CpGs covered and not covered, respectively, by sequencing the amplicon with 2×300mer Illumina reads. *bottom right*: Methylation of the donor L1 promoter sequence in four single-cell miPSC clones, including Clone 1, cultured in either serum or 2i conditions, and the parental MEF population. Each cartoon panel corresponds to an amplicon and displays 50 non-identical randomly selected sequences (black circle, methylated CpG; white circle, unmethylated CpG; ×, mutated CpG). The percentage of methylated CpG is indicated underneath each cartoon. **b,** Donor L1 methylation data as per (a) except for bulk miPSC lines derived from two animals (A67 and A172) carrying the polymorphic donor L1.

Extended Data Fig. 6. L1 T_F subfamily promoter monomer methylation. a, L1 T_F monomer CpG methylation in MEFs, single-cell miPSC clones, and bulk miPSCs derived from primary cells. *top*: Assay design and primer locations with respect to L1 T_F monomer structure. Orange strokes indicate CpGs covered by the assay. *bottom*: Histogram data represent the mean percentage methylation of 50 non-identical bisulfite converted sequences selected at random from each sample. A two-tailed t test (p<0.001) was used to compare serum and 2i culture conditions for single-cell miPSC clones 1-4. **b,** L1 T_F methylation in four single-cell miPSC clones and parental MEFs. Each cartoon panel corresponds to an amplicon and displays 50 non-identical randomly selected sequences (black circle, methylated CpG; white circle, unmethylated CpG; ×, mutated CpG). Methylated CpG percentage is indicated underneath each cartoon. **c,** As per (b) except for bulk miPSCs derived from animals A67 and A172. Note: this assay surveys CpG methylation for T_F monomers genome-wide without retaining their position within individual L1 loci.

Extended Data Fig. 7: Lamivudine does not impact reprogramming efficiency or miPSC survival. a, Cultured MEF and miPSC viability as a function of lamivudine (3TC) concentration. miPSCs were tested for 3 and 9 days in culture with 3TC, and MEFs tested for 9 days. **b,** Cell viability during MEF reprogramming to miPSCs in the presence of varying concentrations of 3TC, as a function of days since reprogramming was induced by the addition of doxycycline. **c,** Reprogrammed MEFs stained for alkaline phosphatase (AP) and *Oct4*-GFP pluripotency markers 15 days after the addition of doxycycline, in media containing 100μM 3TC or no 3TC. Results are shown as microscopy images of stained cultured cells (left) and as histograms (right) representing the mean ± SD observed AP positive colony counts, *Oct4*-GFP positive colony counts, and reprogramming efficiency (percentage *Oct4*-GFP positive cells at day 15 divided by the number of cells seeded at day 0). Differences were not significant (two-tailed t test). **d,** Representative microscopy images of day 15 miPSC colonies obtained when MEFs were reprogrammed in the presence of 100μM 3TC or no 3TC. **e,** Representative immunostaining of Nanog (red), an additional pluripotency marker, in day 15 miPSCs.

Extended Data Fig. 8: Supporting data for ONT sequencing analysis of miPSCs. a, Non-reference polymorphic TE insertions found by ONT sequencing, used as positive controls for PCR validation experimental designs. **b,** A putative *de novo* B1 insertion detected in one miPSC line by ONT sequencing and annotated as a false positive based on PCR amplification in the parental MEF template DNA. **c,** A putative *de novo* L1 T_F insertion detected in one miPSC line by ONT sequencing and amplified in multiple miPSC lines by PCR. Here, DNA was obtained from two parental MEF aliquots, one corresponding to miPSC CTRL/3TC lines 1-3 and one to miPSC CTRL/3TC lines 4-6. Note: in each panel, the insertion chromosomal location and orientation are shown. L1 and SINE B1 and B2 insertions are represented by white rectangles. L1 5′UTR promoter monomers are indicated by triangles. PolyA tracts and their length are indicated (A_n), and target site duplications (TSDs) are depicted as gray arrows. 3′ transductions are shown as orange lines. PCR validation primers are shown as red arrows. PCR products in agarose gels used to confirm TE insertions are indicated by red arrows. Empty site (wild-type) amplicons are indicated by blue arrows, where applicable.

Extended Data Fig. 9: Examples of protein-coding gene and TE methylation, as surveyed by ONT sequencing. Methylation profiles are shown for **a,** the *Hsf5* gene promoter **b,** an IAP LTR intronic to *Ylpm1*, and **c,** an intergenic L1 T$_F$. For each example, the panels are arranged as per **Fig. 4c.** ONT data are shown for MEFs (brown) and for control (green) and 3TC-treated (pink) miPSC datasets. miPSC replicates were aggregated for each condition and down-sampled to approximate MEF dataset read depth.