Prey Size Spectra and Predator to Prey Size Ratios of Southern Ocean Salps

Authors: Christian K. Fender[1,*], Moira Décima[2], Andres Gutiérrez-Rodríguez[3], Karen E. Selph[4],

Natalia Yingling[1], Michael R. Stukel[1,5]

[1]Department of Earth, Ocean & Atmospheric Science, Florida State University, Tallahassee, FL,

USA

[2]Integrative Oceanography Division, Scripps Institution of Oceanography, La Jolla, CA, USA

[3]National Institute of Water and Atmospheric Research, Wellington, New Zealand

[4]Department of Oceanography, University of Hawaii, Honolulu, Hawaii

[5]Center for Ocean-Atmospheric Prediction Studies, Florida State University, Tallahassee, FL,

USA

* Correspondance to: Christian K. Fender (ckf18b@fsu.edu)

Abstract

Salp grazing is important in shaping planktonic food-web structure. However, little is

known about the size ranges of their prey in the field or how grazing impacts size structure. This

study investigated the feeding habits of both the solitary and aggregate life stages of 7 different

species of salps, representing a variety of sizes across subtropical and subantarctic waters east of

New Zealand. Scanning electron microscopy was used to examine the gut contents of 58 salps,

which were then compared to water column plankton communities characterized via

epifluorescence microscopy, FlowCam, and flow cytometry. While most of the gut contents

resembled ambient waters, substantial differences were found amongst some co-occurring

species, such as increased retention of submicron bacteria by *Thalia democratica*. We found that

even for those salps capable of feeding on bacteria efficiently, nanoplankton and small

microplankton still made up the majority of gut biomass. Larger microplankton were rarer in the

guts than in the water column, suggesting an upper size-threshold in addition to the lower size-

threshold that has been the focus of most previous work. Salp carbon-weighted predator to prey

size ratios were variable, with the majority falling between 1,000:1 and 10,000:1 depending

largely on the size of the salp. Taken together our results indicate that despite being able to feed

on submicron particles, picoplankton make up at most 26.4% (mean = 6.4%) of salp gut carbon

and are relatively unimportant to the energetics of most salps in this region compared to

nanoplankton such as small dinoflagellates and diatoms.

1. Introduction

Salps are a group of large (~1 – 30 cm as adults), suspension-feeding gelatinous zooplankton that filter water through a fine mucous mesh at hourly rates of more than 1000 times the organism's biovolume (Madin et al. 2006). With population doubling times as fast as 8 hours due to their dual staged life history composed of both solitary and aggregate forms (Alldredge and Madin 1982; Deibel and Lowen 2012), salps can form swarms of up to 1000 individuals m^{-3} covering up to 9065 km^2 in response to favorable conditions (Anderson 1998; Berner 1967). Salps can have clearance rates >100,000 mL ind^{-1} day^{-1} (Madin and Kremer 1995; Perissinnotto and Pakhamov 1998; Sutherland et al. 2010), equivalent to the clearance rate of 450 copepods (Harbison and Gilmer 1976), allowing these salp blooms to clear water columns of prey so quickly that they can prematurely end spring diatom blooms before surface nutrients are depleted (Bathmann 1988). Despite a growing appreciation for the impact salps may have on ecosystem structure and biogeochemical cycles, there are still gaps in our understanding of their feeding ecology such as the distribution of sizes over which they feed. The ratio of a predator's size to that of its prey has long been linked to the total length of an ecosystem's food web, where large predator:prey size ratios (hereafter PPSR) result in fewer trophic levels and more efficient transport of primary production to large taxa rather than remineralization (Lindeman 1942; Sheldon et al. 1977; Sherr and Sherr 1988). While it is less frequently recognized, the range of prey sizes a predator can feed upon has similar impacts on ecosystem structure. More generalist predators display higher standard deviations in prey size (hereafter SD$_{PPSR}$) resulting in ecosystems with lower connectance, smaller phytoplankton, and less diversity in phytoplankton size classes (Fuchs and Franks 2010). In short, the PPSR and SD$_{PPSR}$ of the dominant predators in an ecosystem play a strong role in determining its trophic structure, and only a handful of studies have quantitatively investigated the size of salps' prey in the field (Dadon-Pilosof et al. 2019; Madin 1974; Madin and Purcell 1992; Vargas and Madin 2004).

It has long been accepted that the relationship between predator and prey size has a strong impact on the ecology of an ecosystem (Sheldon et al. 1977, Jennings et al. 2002) and that PPSR broadly varies between zooplankton of different feeding modes (Hansen et al. 1994). For instance, while most zooplankton tend to feed with PPSRs between 10:1 and 100:1 (Hansen et al. 1994, Fuchs & Franks 2010), raptorial protists often feed with PPSRs of closer to 3:1 and both pallium-feeding dinoflagellates and some siphonophores can feed at a ~1:1 PPSR (Purcell et al.

1981, Fenchel 1987, Sherr et al. 1991, Naustvoll 2000, Sherr & Sherr 2002). At the other end of

the spectrum, for filter-feeding plankton it is common to observe anywhere from a 5:1 to 100:1

ratio between an organism's size and that of its prey (Hansen et al. 1994; Conley et al. 2018).

These variations are the product of physiological differences between raptorial and filter feeders

that lead to significant differences in the portion of available prey biomass that is utilized (Figure

1). For example, raptorial crustaceans can be expected to exhibit a preferred or optimal prey size

whereas the size distribution of prey for filter feeders is more broadly determined by what is

available in the water column. Filter feeders also tend to have higher feeding efficiencies for

small particles compared to raptorial feeders of similar size, differentiating the niche space such

that filter feeders are often in competition with organisms far smaller than themselves (Stukel et

al. 2021). Depending on the size of the filter feeder, this is manifested in substantially higher

PPSRs. High feeding efficiencies for a broader range of prey sizes likewise result in a higher

SD_{PPSR}.

Figure 1. Feeding efficiencies for two theoretical raptorial feeders with an optimal prey size (a)

and for two theoretical filter feeders whose feeding efficiency as a function of prey size is

determined by the mesh size of their filters and the size of their oral opening. (b) Blue and purple

lines represent organisms with higher SD_{PPSR} than the red and green organisms. Hypothetical

prey biomass as a function of size is shown in black in (c) and (d) along with feeding rates as a

function of size, which is equal to prey biomass times feeding efficiency for hypothetical

raptorial (c) and filter (d) feeders. Vertical dashed lines show the carbon-weighted mean prey

size (i.e., the prey size for which half of prey biomass consumed was greater than that size and

half was less than that size), while prey sizes within one standard deviation of the mean prey size

are shown in the shaded colors. PPSR for a given salp was computed as the salp's length divided

by its carbon-weighted mean prey size.

As an extreme example of the efficiency of filter feeders, pelagic tunicates (salps,

doliolids, pyrosomes, and appendicularians) often feed with PPSRs >1,000:1 (Madin 1974,

Crocker et al. 1991, Madin and Purcell 1992, Vargas and Madin 2004, Katechakis et al. 2004).

They can feed directly on micron-sized picoplankton, but also consume ~1-mm organisms

including copepod nauplii, radiolarians, foraminifera (Madin 1974), and ostracods (Decima et al.

2019). Consequently, salps have the potential to simultaneously act as competitors to and

predators of a wide variety of phagotrophic protists (Stukel et al. 2021), the group responsible for

the majority of phytoplankton grazing (Calbet and Landry 2004). Furthermore, as salps

themselves are part of the diet of at least 202 marine species, salps may not act as a net carbon

sink as classically believed but instead increase trophic transfer and vertical export efficiencies

(Michaels and Silver, 1988; Henschke et al. 2016). Their presence could thus support

economically important species like mackerel (Nishimura 1958), bluefin tuna (Cardona et al.

2012), anchovies (Mianzan et al. 2001), and other demersal fishes (Horn et al. 2011; Forman et

al. 2016). However, the physiological ability to feed at PPSRs up to 10,000:1 does not

necessarily mean that this is the range over which most salps feed. As non-selective filter

feeders, it is likely that this value is largely dependent on the phytoplankton community present.

Differing retention efficiencies for small particles have also been reported for a variety of salp

species based on removal studies in deck-board incubators (Kremer and Madin 1992; Vargas and

Madin 2004; Stukel et al. 2021), though these techniques often struggle to accurately resolve

very small or very large prey. The size of the salp itself likely plays some role as well, as there is

evidence that small salps may retain smaller particles more efficiently than larger individuals

even of the same species (Harbison and McAlister 1979; Kremer and Madin 1992; Stukel et al.

2021), and that the size of the oral opening will limit consumption of larger-sized phytoplankton

cells and/or diatom chains (Figure 1b). Whether feeding differs intrinsically between species or

life stages or if it is purely a function of size is also unknown.

Although traditional microscopic analyses of salp stomach contents have long supported

their non-selective nature (Silver 1975; Vargas and Madin 2004; Tanimura et al. 2008), these

methods are limited in terms of image resolution such that only large, hard-bodied plankton can

be easily identified. Similarly, modern tracer and genetic analyses have also been applied to gut

contents to determine relative proportions of different types of prey (von Harbou et al. 2011;

Metfies et al. 2014; Pauli et al. 2021), but these methods do not allow for determining prey size.

A handful of studies have also attempted to directly assess salp gut contents via scanning

electron microscopy (SEM), which can resolve particles from the submicron to millimeter size

range. Unfortunately, preparatory methods vary, and the focus has mostly been on larger, more

easily identifiable prey items like diatoms and thecate dinoflagellates (Madin and Purcell 1992;

von Harbou et al. 2011; Ahmad-Ishak 2017). This, along with observations of the mesh spacing

in their mucous filters, have led to the commonly accepted view that most salps predominantly

feed on particles >1-2 µm in diameter due to the inefficiency with which smaller particles are

retained (Harbison and McAlister 1979; Kremer and Madin 1992; Madin and Purcell 1992).

Some studies leveraging numerical simulations as well as removal experiments using beads of

known size have recently challenged this, suggesting that direct interception of submicron

particles by the mucous fibers may be a more important source of nutrition than originally

believed (Sutherland et al. 2010). Even if retained less efficiently than larger particles, submicron

phytoplankton like cyanobacteria are present at far higher concentrations than larger taxa such

that feeding on them may still satisfy a large portion of a salp's carbon quota (Sutherland et al.

2010; Dadon-Pilosof et al. 2019).

In this study, we sought to compare the size composition of prey items in salp stomachs

to that of the water column and determine if the resulting size spectra are a function of salp

species, size class, or life stage, then quantify how these demographics impact the PPSR, SD_{PPSR},

and retention efficiencies of salps. We utilized gut content SEM analysis of 58 individuals

comprised of both life stages of 7 different salp species ranging from 10-155 mm to quantify

their diets and compare these diets to water column plankton populations characterized by flow

cytometry, epifluorescence microscopy, and FlowCam imaging. To our knowledge, this is the

most diverse collection of salp diets sampled to date in a single study.

2. Materials and Methods

2.1 Field collection

Samples were collected over Aotearoa New Zealand's Chatham Rise during October 21 - November 21 2018 as part of the Salp Particle Export and Ocean Production (SalpPOOP) study, which was designed to investigate how salps affect the ecology and biogeochemistry of the region (Décima et al., in review). The Chatham Rise is notable because it sits within the Subtropical Front of the southwest Pacific, which defines the boundary between warm, salty and low-nitrate subtropical waters and cold, fresh, nitrate-rich, and iron-poor subantarctic waters (Heath 1985; Sokolov & Rintoul, 2009; Zentara and Kamykowski, 1981). This creates a dynamic frontal zone with high mixing that supports high phytoplankton biomass and productivity (Currie and Hunter, 1998) as well as, anecdotally, periodic summertime salp blooms. The cruise consisted of 5 quasi-Lagrangian experiments (hereafter "cycles") lasting 4-8 days each, with only 4 cycles (Cycle 1-4) showing significant salp presence (Figure 2). Cycle locations were chosen based on net tows through regions with high potential for salp presence according to previous observations on habitat distributions of salp-eating demersal fish (Forman et al. 2016; Horn et al. 2011).

At the beginning of each cycle, we deployed a surface-tethered drifting array to track the chosen water parcel (Landry et al. 2009). Daily dawn and noon deployments of a 24-bottle CTD-Niskin rosette were made to collect water used in bottle-incubation experiments as well as water column profiling. To characterize the prey community, we analyzed 4 types of samples collected from Niskin bottles: (1) 250 mL subsamples from the base of the mixed layer and the deep chlorophyll max (DCM) were concentrated by gravity filtration to 10 mL over a 2 µm 47 mm filter, and 2 mL of this concentrate was imaged using a FlowCam's 10X objective lens to quantify the larger (>3 µm) phytoplankton (Sieracki et al. 1998); (2) an Accuri C6 Plus flow cytometer was used at sea to determine the abundance of *Synechococcus* and phototrophic eukaryotes collected at 6 depths spanning the euphotic zone with cell size estimated using the forward light scatter of polystyrene beads of known size (Stukel et al. 2021); (3) additional preserved samples were collected from the same casts (same 6 depths) and analyzed on a Beckman Coulter CytoFLEX S flow cytometer to enumerate heterotrophic bacteria and *Prochlorococcus* (Selph 2021); (4) preserved and stained microscopy samples were collected

from 6 depths per cast and later analyzed on an epifluorescence microscope where cell biomass

and abundance were calculated using ImageJ (Taylor et al. 2015). Each of these four methods

allowed for the enumeration of overlapping parts of the phytoplankton size spectra, namely 0.4-

50 µm for flow cytometry, 2-200 µm for epifluorescence microscopy, and >4-300 µm for

FlowCam. In total, 21 CTD casts were sampled over the course of the 4 cycles from which salps

were collected.

To determine salp biomass and demographic structure, we conducted twice daily oblique

Bongo tows down to 200 m. Ring net surface tows with a 30 L non-filtering cod-end were also

conducted daily to collect additional salps. Once onboard, salps from ring net and Bongo tows

were identified to the species level, sorted by life stage (i.e., solitary or aggregate), measured,

and sexed (Lüskow et al. 2020). Triplicate representative samples for each salp species from a

total of 10 of these casts were preserved in 5% formalin < 30 minutes after collection the first

time each species was encountered. While this means there was potential for prey digestion

between the time the salp was caught and when preservation occurred, processing time was

always substantially less than the estimated gut pigment turnover time of 2.8 hours for even the

smallest salp caught (von Harbou et al. 2011). Whenever possible these preserved samples

included a distribution of size classes and both solitary and aggregate life stages.

Figure 2. a) TAN1810 cruise study area (blue square) located east of Aotearoa New Zealand with

color denoting bottom depth. b) Magnification of study region with color denoting sea surface

temperature. Yellow diamonds represent CTD deployments to determine water column

properties and red circles represent the locations of salp collections for SEM via Bongo or ring

net tow.

2.2 Scanning electron microscopy

In total, 58 salps representing the species *Salpa thompsoni*, *Pegea confoederata*, *Thalia*

democratica, *Soestia zonaria, Thetys vagina*, *Salpa fusiformis*, and *Ihlea magalhanica,* including

both solitary and aggregate stages of the first four and a distribution of sizes for the first two,

were collected. Once ashore, SEM samples were prepared from each preserved organism by

excising guts under a HEPA-filter equipped laminar flow exhaust hood using acid-cleaned

plastic dissection equipment to minimize contamination. Guts were then placed in either 15 or 50

mL plastic Falcon tubes with a small volume of brine, lacerated, and then vortex mixed for two

minutes to release gut contents into solution while minimizing damage to the more fragile

phytoplankton (Jung et al. 2010; von Harbou et al. 2011; Ahmad-Ishak 2017). An aliquot of this

solution was then filtered onto a 0.2 μm Nuclepore filter, followed by six rinses of decreasing

salinity in 5 ppt increments for a minimum of 5 minutes each with the final MilliQ water rinse

performed twice. This was immediately followed by a dehydration series of increasing ratios of

Ethanol:MilliQ to purge water from the sample, with the final 100% anhydrous ethanol step

again performed twice. Finally, a substitution series of increasing ratios of the chemical drying

agent hexamethyldisilazane (HMDS):anhydrous ethanol was conducted with each step lasting a

minimum of 10 minutes, with the final HMDS step being allowed to air dry. We chose this

chemical drying agent over more traditional critical point drying both to minimize changes in

cell size as well as maintain material on the filter (Jung et al. 2010). Each step was conducted

under either a light vacuum or gravity filtration depending on material concentration to minimize

loss between treatments. Similar procedures have proven successful in the preparation of both

delicate dinoflagellates (Botes et al. 2002; Jung et al. 2010) and bacteria (Koon et al. 2019).

The dried filter was then affixed to an aluminum SEM stub using carbon conductive

adhesive tabs and further grounded with a thin piece of carbon conductive tape touching the edge

of the filter and the bottom of the stub. Samples were then sputter coated with 10 nm iridium and

visualized using an FEI Nova 400 NanoSEM set to an accelerating voltage of 10 kV. Twenty

random regions of each filter were imaged at 3 different magnifications: ~500x, ~2,500x, and

~12,000x to target microplankton (20-200 µm), nanoplankton (2-20 µm), and picoplankton (<2

µm), respectively. Since sufficient structural detail could not be observed to definitively identify

very small particles, spherical particles within the size range of ~0.4-1.5 µm and resembling

control images from lab cultures of *Prochlorococcus sp.* and *Synechococcus sp.* are instead

referred to as bacteria-like particles. We also note that while formalin preservation can lead to

cell shrinkage (Choi and Stoecker 1989; Zinabu and Bott 2000), we assume that the shrinkage

would be roughly proportional amongst all cells and therefore does not impact our relative

contributions of different groups to the nutrition of the salps. Furthermore, we assume that this

shrinkage will have an effect on the dimensions of most phytoplankton taxa that is within our

margin of error with the exception of ciliates, for which we apply a carbon conversion

specifically for formalin fixed cells, and nanoflagellates. Studies have shown ~40% decreases in

cell volume for cultured flagellates due to formaldehyde fixation (Choi and Stoecker 1989;

Zinabu and Bott 2000), which translates to a 10-20% decrease in ESD. This roughly corresponds

with the 20% decrease in total length of salps preserved in 5% formalin noted by Madin et al.

(2006) which will have a small impact on the absolute lengths of both predator and prey reported

here (although this small difference is within our margin of error) but is unlikely to affect the

PPSR.

2.3 Image processing

Particles in SEM and epifluorescence microscopy images were manually outlined using

ImageJ (v. 1.52a or 1.53c) to extract the maximum feret length (L), minimum feret length, and

area (A). These measurements were used to estimate equivalent spherical diameter (ESD),

biovolume (BV), and carbon biomass assuming a prolate spheroid. To avoid overestimating the

size of irregularly shaped particles, we calculated width (W) for a prolate spheroid of measured

A and L such that:

256   1) $W = \frac{4\pi}{3} \times \frac{A}{L}$

257   2) $ESD = \sqrt[3]{L \times W \times \frac{W}{2}}$

258
$$3) \quad BV = \frac{4\pi}{3} \times \frac{L}{2} \times \frac{W}{2} \times \frac{W}{4}$$

Note that because we estimated the three-dimensional size of particles using a two-

dimensional image, we determined the height of each particle from its width. To account for the

~50% flattening of height caused by filtration (Taylor et al. 2011), we furthermore assumed

height to be equivalent to half of the width. The biomass of formalin-preserved ciliates was

estimated as 0.14 pg C μm^{-3} (Putt and Stoecker 1989) while rhizarians were 0.001 pg C μm^{-3}

(Stukel et al. 2018). The biomass of diatoms was estimated allometrically as $0.288*BV^{0.811}$ while

other protists and unidentified particles were estimated using $0.216*BV^{0.939}$ (Menden-Deuer and

Lessard 2000). Because we could not differentiate between types of prokaryotes in the SEM

images, we calculated a single average biomass conversion for all bacteria-like particles in the

salp guts using published allometric relationships weighted by the ratio of each of the key

bacterial groups to each other in the water column from flow cytometry data for a given cycle

(see Supplementary Methods).

FlowCam image analyses were conducted using FlowCam's dedicated classification

software VisualSpreadsheet (v. 4.18.5). First, duplicate images resulting from parabolic flow

within the flow cell were manually removed. The particles within the remaining images were

then classified based on the quality with which VisualSpreadsheet detected their outlines. For

particles where the outline appeared to provide good estimates of L and W, size was calculated

for a prolate spheroid as above using Equations 1-3 with the exception that W was left as the

directly measured minimum feret length and no correction for flattening was applied. While this

is likely less accurate than utilizing particle area as was done for the SEM images, we frequently

noted that particle outlines possessed large holes not representative of overall shape such that

area was severely underestimated. For particles where the quality of automatic outline detection

was poor, alternative methods used to determine size can be found in the Supplementary

Materials. Biomass was calculated as described above for the salp gut contents with the

exception of a non-formalin preserved conversion for ciliates of 0.19 pg C μm^{-3} (Putt and

Stoecker 1989). For epifluorescence microscopy, biomass was similarly calculated using either

the allometric equation from Menden-Deuer and Lessard (2000) for diatoms or for other protists.

Carbon content for cells detected via flow cytometry were also calculated based on biovolume

using the relationship for protists from Menden-Deuer and Lessard (2000), with the exception of

Prochlorococcus (36 fg cell^{-1}; Buitenhuis et al. 2012) and heterotrophic bacteria (11 fg cell^{-1};

Garrison et al. 2000) for which we used fixed carbon conversions.

2.4 Size spectra and feeding distribution parameter calculations

For each cycle, the prey items in the salp guts were separated into size bins and their

abundance was averaged for each salp species. This abundance was then divided by the size bin

width to calculate normalized abundance size spectra (NASS). To avoid biasing our size spectra

low or overcounting more fragile species, we chose to exclude all particles in the salp guts which

were clearly broken (less than ¾ their assumed true length). This had negligible impacts on the

initial counts of small dinoflagellates and centric diatoms but did result in 82% of the initial

pennate diatom and >90% of large dinoflagellate counts being disqualified. However, sensitivity

analysis on the inclusion of these particles revealed little impact on any of our results and did not

yield substantially different size spectra since these were relatively rare organisms. NASS were

similarly produced for each of the methods used to assay the prey field (FlowCam, flow

cytometry, and epifluorescence microscopy samples) at each depth and vertically integrated over

the euphotic zone. We also present a single composite water column spectrum representing the

geometric mean of each method over the size ranges where it was relevant. Carbon biomass

estimates for both salp and water column measurements were then used in the same way to

compute normalized biomass size spectra (NBSS). 95% confidence intervals were calculated

using Monte Carlo random resampling (see Supplementary Methods for details).

The carbon-weighted mean prey size of each salp was calculated by sorting all particles

in a gut from smallest to largest and finding the prey size at which 50% of the total biomass in a

salp gut was achieved (Supplementary Figure 2). The total length of a given salp was then

divided by this value to estimate the carbon-weighted predator:prey size ratio (PPSR). We also

calculated the SD_{PPSR} by finding the particle size at which ±1 standard deviation, or 15.9% and

84.1%, of a given salps total gut carbon content was met and then dividing the difference of the

log transformed values by 2. The results are thus feeding distribution values that are more

representative of the energetic value of the salps' diets.

The degree to which each salp can efficiently retain picoplanktonic prey is determined by

the mesh spacing of their feeding filter, which varies between species (Bone et al 2003). We

estimated retention efficiency (hereafter RE) by dividing the prey NASS consumed by each salp

species by that occurring in the water column for each size class and dividing the result by the

average clearance rate for the 8-32 µm particles as this is the size range expected to be retained

with 100% efficiency. Here we chose to average over 8-32 µm particles as ESDs slightly smaller

than this will sometimes possess aspect ratios allowing them to pass through the feeding mesh

and our results show that particles larger than this were frequently absent from salp guts.

3. Results

3.1 Salp vs. ambient phytoplankton community comparisons

Cycle 1, which occurred closest to the coast of the 4 cycles and was most strongly

influenced by the Southland Current, a coastal expression of the Subtropical Front (Sutton 2003),

had the largest contribution of microplankton to the water column plankton community (66.5%

of total vertically integrated plankton carbon, Figure 3). It was also the only cycle where

phytoplankton larger than 120 µm were detected. Nanoplankton and picoplankton contributed

20.6% and 12.9% of the carbon, respectively. Qualitative analysis of FlowCam images from

Cycle 1 suggests the microplankton were mostly centric and pennate diatoms. Flow cytometry

data showed that *Synechococcus*, picoeukaryotes, and heterotrophic bacteria were present in all 4

cycles, while *Prochlorococcus* sp. was only present during Cycle 2. Cycle 2 was farther from the

coast and supported a smaller proportion of microplankton (29.6% of total carbon) characterized

by diatom assemblages similar to Cycle 1 that may have been advected off the coast. The

contribution of nano- and picoplankton, however, was approximately twice that of Cycle 1 at

38.1% and 32.3%, respectively. Cycle 4, which was conducted approximately 1° north of Cycle

2 in waters 3.1°C warmer (Figure 2), had very similar size contributions albeit with nearly all of

the microplankton portion made up by large dinoflagellates rather than diatoms. Cycle 3 had

only 21.2% of its total carbon in the microplankton size range, with no >60 µm cells observed.

Instead, 67.9% was comprised of nanoplankton, twice that of Cycles 2 and 4 and four times that

of Cycle 1. The contribution of picoplankton was comparable to Cycle 1 at 11.0%.

Figure 3. Contribution of particles of various size classes to the total biomass present in the water

column of each cycle (outlined by black boxes) compared to that of the guts of each of the 58

salps sampled (as indicated by their number on the x-axis), with salps grouped by species as

indicated by their initials along the top of each section. Within species salps are ordered from

smallest to largest. See Supplementary Table 1 for more information regarding the individually

numbered salps. WC = water column, SF = *Salpa fusiformis*, ST = *Salpa thompsoni*, TV =

Thetys vagina, SZ = *Soestia zonaria*, PC = *Pegea confoederata*, IM = *Ihlea magalhanica*, TD =

Thalia democratica.

Microplankton within salp guts were broadly representative of the taxa present in the

water column with some notable exceptions elaborated on in the Discussion. When considering

the contribution of various size classes to the gut content of all 58 salps, picoplankton made up

between 0.6% and 26.4% of total biomass (Figure 3) with a mean and standard deviation of

6.4±5.0%. This is substantially lower than the cycle averaged mean contribution in the water

column of 21.9±7.4%. The most abundant objects observed in the guts of all salps regardless of

size or species were unidentifiable, smooth, spherical particles (hereafter referred to as "white

spheres") lacking any other notable morphological features and generally ~2-7 µm in diameter

but ranging from 0.4-22.5 µm (Supplementary Figure 3). Given the broad range in size, it is

likely that this category comprises several unrelated taxonomic groups that had been partially

digested, potentially including prymnesiophytes, pelagophytes, prasinophytes, cysts (diatoms or

dinoflagellates), and/or other nanoflagellate taxa. These, along with large numbers of similarly

sized dinoflagellates, contributed to high concentrations of nanoplankton within the gut contents

(range = 36.8% - 99.2% of total gut biomass, with a mean of 81.9±13.5%). This is overall

substantially higher than what was observed in the water column as nanoplankton made up only

35.3±20.1% of the biomass on average. Microplankton were generally much rarer in the salp

guts (range = 0% - 60.3% of biomass, mean = 11.7±14.2%) in contrast to the available prey in

the water column (water column microplankton = 42.8±24.3% of biomass).

We also compared the average size spectra of prey for each salp species to that of the

water columns of each of the four cycles from which samples were obtained (Figure 4). Since the

volume filtered could not be determined for the salps caught, the units for the normalized

abundance size spectra (NASS) of the prey and normalized biomass size spectra (NBSS) differ

between salp and ambient measurements such that only the shape of the spectra can be

compared. This shape compares the relative importance of small to large particles with steeper

(i.e., more negative) spectra indicating more small particles relative to large ones and vice versa.

In general, most of the salp ingested-prey spectra match up well to both the water column NASS

and NBSS as would be expected for non-selective filter feeders. However, every salp species

shows the same sharp decline in the abundance of consumed submicron particles, which supports

the assumption of inefficient capture of particles <1 µm. There are several other notable

departures from the ambient spectra as well. For example, in *Thalia democratica* of Cycle 3 the

NBSS spectra is much flatter than that of the flow cytometry. While this trend is less apparent for

T. democratica observed in Cycle 4, it could suggest an increased importance of small particles

for this species relative to the others. In contrast, the NASS and NBSS spectra for *Ihlea*

magalhanica is far more peaked at 2-4 µm compared to ambient; rather than suggesting

difficulty in retaining small particles, this shape is likely due to the far higher number of white

spheres relative to any smaller or larger particles observed in their guts. *Salpa thompsoni* and

Soestia zonaria guts of Cycle 1 also display a surprising lack of particles >50 µm despite their

presence in the water, as shown by FlowCam and epifluorescence microscopy. Similarly, the

maximum particle size was only 24.7 µm in the *Thetys vagina* guts from Cycle 1. All other salp

species regardless of cycle also display this trend, with the exception of *Pegea confoederata* for

which the contribution of large particles is still far lower than predicted by the ambient spectra of

Cycle 2.

Figure 4. Average normalized abundance (left) and normalized biomass (right) size spectra of

salp prey (NASS and NBSS, respectively) as a function of equivalent spherical diameter (ESD)

for Cycles 1-4. Warm-colored lines denote salp gut contents while cool-colored lines represent

water column measurements from samples collected from the CTD rosette. The thicker, dark

blue, dashed line (labeled composite) represents the geometric mean of the water column

measurements. Shaded regions represent 95% confidence intervals for each spectrum. Note the

difference in units for y-axes and that some water column measurements are not visible due to

overlap with the composite spectrum.

3.2 Feeding distribution parameters

Most salp species obtained the majority of their carbon from particles less than 13 µm

(Figure 5; Supplementary Table 1), likely due to the predominance of nanoplankton present in all

four cycles coupled with the salps' apparent difficulty in capturing large particles. Notably, the

smallest salp species (*Thalia democratica*) had the second lowest average prey size of 6.8 µm

ranging from 4.4 to 9.3 µm (Figure 6) which may be due to better adaptation for feeding on

small particles or simply because the majority of this species was collected from Cycle 3 which

had relatively few microplankton (Figure 3). Species from Cycles 1 and 2 (where there were

more large particles in the water) such as *Salpa thompsoni* and *Soestia zonaria* typically fed on

slightly larger prey, ranging from 2.7 to 33.9 µm with a mean of 11.9 µm for *S. thompsoni* and

7.4 to 11.4 µm with a mean of 9.6 µm for *S. zonaria*. Within species, the size of the salp did not

seem to substantially affect the mean prey size. For example, the mean prey size of *Pegea*

confoederata was 9.1 µm ranging from 3.5 to 12.4 µm for individuals varying in size by almost

an order of magnitude. Even between species, no trend with size was observed as the mean prey

size of the largest salp *Thetys vagina* is quite comparable at 7.2 µm with a range of 6.0 to 8.4 µm

to that of the smallest species *T. democratica*. Likewise, there does not seem to be any trend with

salp life stage as the mean prey size was similar for all species for which we encountered both

stages: *S. zonaria* (aggregate=9.1 µm; solitary=10.0 µm), *P. confoederata* (aggregate=9.2 um;

solitary=8.9 µm), *S. thompsoni* (aggregate=12.5 µm; solitary=9.9 µm), *T. democratica*

(aggregate=6.3 µm; solitary=7.7 µm). *Ihlea magalhanica* had the smallest mean prey size (4.4

µm ranging from 3.8 to 4.7 µm) because their guts almost exclusively contained small white

spheres and bacteria-like particles. The largest degree of variability in mean prey size was seen

in *S. thompsoni*, which had several individuals where more than 50% of their gut biomass was

contained in one or two very large particles, such as polycystine radiolarians, substantially

inflating the mean prey size. *S. thompsoni* was also the most abundant salp species and one of the

taxa for which we collected the most samples (n=14) such that the probability of catching a *S.*

thompsoni with a large, rare prey particle in its gut was higher.

Figure 5. Prey size (left) and predator:prey size ratio (right) at which 50% of gut content biomass
is achieved as a function of salp size for each of the 58 salps collected. Color denotes salp
species while shape denotes the cycle in which the sample was collected. Filled symbols denote
aggregates and empty symbols denote solitaries.

Figure 6. Box and whisker plot of the distribution of carbon-weighted mean prey sizes averaged
by salp species. SF = *Salpa fusiformis*, ST = *Salpa thompsoni*, TV = *Thetys vagina*, SZ = *Soestia
zonaria*, PC = *Pegea confoederata*, IM = *Ihlea magalhanica*, TD = *Thalia democratica*.

Most salps had predator-prey size ratios (PPSR) between 1,000:1 and 10,000:1, with only
the two largest *S. thompsoni* individuals and the two *T. vagina* feeding at or above 10,000:1. *T.
vagina* had the highest PPSR of 21,548:1, followed by *P. confoederata* (5,275:1), *S. zonaria*
(4,529:1), *Salpa fusiformis* (4,490:1), *I. magalhanica* (3,845:1), *S. thompsoni* (3,403:1), and *T.
democratica* (1,846:1). When averaging by life stage as well, solitary stage salps always had a
higher PPSR than aggregates of the same species although this may be due to the fact that the
solitary stage individuals tended to be larger. There was a clear trend of increasing PPSR with
increasing salp size because of the larger variance in salp sizes compared to the relatively similar
mean prey sizes. However, a great deal of variation in PPSR was seen amongst individuals of the
same species and size. The PPSRs of smaller *S. thompsoni* range by almost an order of
magnitude, while *P. confoederata* and *T. democratica* possessed many individuals of
approximately the same size yet also show PPSRs that vary by a factor of 2-3 in a given size
class, reflecting similar variability in mean prey size. Also of note, PPSR was largely
independent of which cycle the salp was collected from; 35-40 mm *S. fusiformis* and *S. zonaria*

collected during Cycle 4, which was much farther from the coast and displayed relatively higher

concentrations of dinoflagellates were comparable to *S. zonaria* and *P. confoederatea* of the same

size from the coastal, diatom rich Cycle 1. We caution that this does not imply that the

differences in prey community composition across cycles have no bearing on PPSR, but rather

that the greater degree of variability seen in salp size is the primary determinant.

4. Discussion

4.1 Feeding distribution parameters

The relative abundance of predators with different PPSR has the potential to substantially

alter relationships between trophic level and size, with commensurate impacts on energy transfer

to larger taxa of commercial importance and carbon export into the deep ocean (Barnes et al.

2010; Michaels & Silver 1988). In addition to a predator's mean PPSR, the range of sizes upon

which a predator can feed has the potential to structure food webs. Fuchs and Franks (2010)

concluded that a dominance of specialized predators which feed on a narrow range of prey sizes

(low SD_{PPSR}) or are large relative to their prey (high PPSR), such as copepods, would lead to an

ecosystem state with lower connectivity between trophic levels, fewer omnivores, more top

predators, and greater transfer efficiency to higher trophic levels due to the reduced number of

trophic interactions. Conversely, high SD_{PPSR} and/or low PPSR predators like dinoflagellates,

raptorial ctenophores, or ciliates led to an ecosystem state with the opposite characteristics.

We carefully quantified the mean PPSR for 7 salp species and the range of sizes that

comprised the majority of their carbon-weighted prey, finding Southern Ocean PPSRs that were

typically higher than the summary of estimates for salps reported in a meta-analysis by Fuchs

and Franks (2010) from other regions. We saw typical values of 1,000:1 to 10,000:1, with all but

3 individual salps >1,000:1 and 74% of our dataset above the maximum reported range of Fuchs

and Franks (2010) at 2,236:1 (Figure 7). As salps are non-selective feeders, some of this

difference may be due to the different prey communities present in the regions from which Fuchs

and Franks (2010) acquired their data, which include the Mid-Atlantic Bight (Vargas and Madin

2004), Subarctic Pacific (Madin and Purcell 1992), Florida Current, and Gulf of California

(Madin 1974). Their data also include several species not observed in our study, such as

Cyclosalpa affinis, C. pinnata, C. bakeri, and *Weelia cylindrica,* which are quite physiologically

distinct from the species investigated here and may also exhibit different feeding characteristics

(Harbison and McAlister 1979; Bone et al. 2003). The individuals in Fuchs and Franks (2010)

dataset are also on the small end of our size range, with 3 out of their 5 datasets containing only

salps <20 mm and none containing salps >100 mm (whereas our *T. vagina* were as large as 163

mm). This also likely plays a role in the larger values we observed, because we found little

difference in prey size between large and small salps. Considering our Southern Ocean data

along with the data from other regions described above, salp PPSRs range over two orders of

magnitude, which is comparable to the difference between pallium-feeding dinoflagellates and

suspension-feeding copepods. The SD_{PPSR} for our study ranged from 0.14 to 0.53, while Fuchs

and Franks (2010) estimated similar values ranging from 0.22 to 0.50. Thus, the range of prey

sizes salps feed on can be broader than most other planktonic predators or as narrow as the

protists they compete with, again largely varying with the specific conditions of a given study

region. Attempts to characterize salp predator:prey interactions in models using a single set of

generalized parameterizations would fail to capture this regional variability and may lead to

significant error when applied to large, heterogeneous areas such as the Southern Ocean. Even

for global modelling attempts, it is clear that ascribing a single PPSR of 10,000:1 will lead to

misleading results. Models should endeavor to use appropriate PPSRs depending upon the prey

field and salp population in the region studied.

Figure 7. Predator:prey size ratios and standard deviation of prey size for a variety of planktonic predators from Fuchs and Frank (2010; black circles) with bars representing the range. Squares represent individual salp data from this study, with color representing salp species. Filled symbols denote aggregates and empty symbols denote solitaries. Note that values from this study are carbon-weighted whereas those of Fuchs and Franks (2010) are not.

While our estimates of SD_{PPSR} are fairly comparable to those of Fuchs and Franks (2010), ~1/4 of our dataset is below their minimum value of 0.22. This is probably due in part to the dominance of ~10 µm prey particles that were ubiquitous across nearly all salp's guts sampled in our study (Figure 5). While many of these were clearly small dinoflagellates or centric diatoms, the most common particulate items found in salp guts were ~2-7 µm smooth white spheres (Supplementary Fig. 3H). These spheres made up 25% of all identified particles and were present even in samples where little to no other recognizable taxa were found. Unknown particles matching this description were also described by both Madin and Purcell (1992) and Ahmad-Ishak et al. (personal communication) and can be seen in the SEM images taken by Caron et al.

(1989) although they were not discussed. Due to their size and generally spherical shape, we

assume that the majority of these particles were nanophytoplankton such as prasinophytes,

prymnesiophytes, or pelagophytes for which characteristic features such as flagella had been

digested or broken off as these groups had a high abundance and contribution to phytoplankton

community biomass (Décima et al. in review). It is, however, likely that this morphological

categorization includes particles of various origins. For example, some displayed significant

silicon signals under electron diffusion spectroscopy and therefore may have instead been the

resting stage cysts of diatoms. Alternatively, some may have been debris or tissue associated

with the preparation of the salp itself rather than ingested prey particles. How these particles are

treated has a strong impact on how some of our results should be interpreted. For example, the

NBSS of *I. magalhanica* showed a strong peak at 2-8 µm due to the exceptionally high numbers

of white spheres observed in the guts of all 3 individuals imaged (Figure 4). When these are

treated as nanoplankton, as we believe is most likely, our data suggests feeding on particles of

this size class is even more important for this species of salp relative to the others investigated

here. If white spheres were not treated as particles and instead excluded from our analysis, the

resulting spectra for *I. magalhanica* would be much the same as that of *T. democratica* from the

same cycle and would instead suggest their feeding dynamics are similar (Supplementary Figure

4). Both salp spectra, however, would then be significantly lacking in particles of this size

relative to the ambient spectra. This is also true of the other salp spectra across all cycles and

species, with even more noticeable declines in nanoplankton for the salps of Cycles 2 and 4. We

interpret this as evidence that the majority of these white spheres indeed originated in the water

column but caution that future work, especially if seeking to automate image processing, should

take special care with particles in this size class.

4.2 Retention efficiency

For most salps, we find 100% retention for 1-16 µm particles (Figure 8), but much lower

retention for submicron particles, which agrees with many previous studies (Harbison and

Gilmer 1976; Kremer and Madin 1982; Caron et al. 1989; Sutherland et al. 2010; Nishikawa and

Tsuda 2021; Stukel et al. 2021). With respect to the larger prey size classes, however, the most

striking result of our calculations is the rapid decline in retention of 16-64 µm particles for many

salps. While our methods were optimized for finding the lower threshold of retention, making it

difficult to pinpoint a single particle size for the drop-off, it's clear that this resulted in surprisingly low contributions of microplankton to the total salp gut biomass compared to what was found in the water column.

Figure 8. Average retention efficiency as a function of prey equivalent spherical diameter (ESD) for each salp species, organized by cycle, assuming filtration rate is equivalent to the clearance rate on 8-32 µm cells. Shaded areas represent 95% confidence intervals.

Typically, all particles above the width of a salp's mucous feeding mesh are considered to be retained with 100% efficiency. However, previous studies used only a selection of particles of up to ~10 µm in size, so no data existed for larger particles and their relative absence from salp diets may have been easily missed. Alternatively, several factors could have led to an underestimate of RE for these larger size classes in the present study. For example, it is possible that larger, fragile organisms like chains of pennate diatoms were separated into solitary cells and/or further fragmented as the salp's filter was passed through the esophagus and into the

stomach or even as a result of our preparations in the lab. Indeed, deck-board prey disappearance

experiments with *S. thompsoni* during our cruise, which would not suffer this bias, did not show

a consistent decrease in RE for particles up to 30 µm ESD (Stukel et al. 2021). However, the

incubation experiments of Vargas and Madin (2004) did find 30-40% lower retention efficiencies

for 60 µm diatoms in Mediterranean *Salpa cylindrica* and *Cyclosalpa affinis*. They proposed the

existence of a size threshold for large particles at which the likelihood of bolus formation and

subsequent expulsion of the feeding filter (Harbison and Gilmer 1976; Harbison et al. 1986;

Vargas and Madin 2004) further decreases their already low relative abundance in the guts. To

this end we noted a complete absence across all 58 salps of large, spiny centric diatoms like

Chaetoceros sp. (both solitary cells and chains) as well as those which appeared to be either

Proboscia sp. or *Rhizosolenia* sp. which dominated the >60 µm size classes in the FlowCam.

Similar discrepancies were observed by Ahmad-Ishak et al. (2017) who noted that while both

Chaetoceros and *Proboscia* made up large biovolumes in their study region according to light

microscopy, only *Proboscia alata* was found in the guts of *S. fusiformis* and *T. democratica* via

SEM. Our understanding of the conditions under which boluses form is still too poor to allow for

more than speculation here.

While only a handful of studies have attempted to quantify the mesh spacing of the

mucous feeding filters in salps (Silver and Bruland 1981; Bone et al. 1991; Bone et al. 2000;

Sutherland et al. 2010), the mesh of *T. democratica* has been observed to be quite fine at 0.9 µm

x 0.9 µm (Bone et al. 2003). This is one explanation for their markedly higher retention

efficiencies for submicron particles compared to other species (Figure 8). Indeed, there are

reports of *T. democratica* feeding at varying levels of efficiency on bacteria (Mullen 1983;

Vargas and Madin 2004). Higher retention of small particles could also be related to the size of

the individual as several studies have reported higher retention efficiencies for small particles in

smaller salps (Harbison and McCalister 1979; Kremer and Madin, 1992, Stukel et al. 2021)

potentially due to isometric scaling of the mesh width to organism size (Sutherland et al. 2010).

Indeed, *Salpa fusiformis* REs for submicron particles are as high as those for *T. democratica* and

stand in stark contrast to the closely related *Salpa thompsoni* which struggled to retain anything

smaller than 1 µm. We postulate that this is due to the lower mean size of *S. fusiformis* (34.7

mm) captured compared to that of our *S. thompsoni* (50.3 mm). When instead averaging *S.*

thompsoni retention across different size classes, we again find smaller individuals display

higher REs for submicron particles. These observations match reasonably well to those predicted

by the equation for *S. thompsoni* size specific clearance rate given in Stukel et al. (2021), which

describes mesh diameter as an allometrically-scaling function of salp size (Figure 9). Note that

these results do not contradict our earlier observations of mean prey size being relatively

independent of salp size. As mean prey size is a function of both retention efficiency and the

prey field available, the impact of the differences in RE for small particles we find here on mean

prey size, and thus PPSR, will be proportional to that size classes contribution to the ambient

carbon pool. Submicron particles never comprised >31.5% (and usually substantially less) of the

total carbon in the water column, so it's not surprising that the mean prey size and PPSR of any

given salp was not strongly impacted by its ability to retain these particles.

Figure 9. Retention Efficiencies (RE) for different sizes of *S. thompsoni* assuming a filtration rate

equal to the clearance rate for 8-32 um particles. Solid lines represent REs averaged by salp size

class, dashed lines represent REs calculated using the size resolved clearance rate equation from

Stukel et al. (2021) for *S. thompsoni*, and shaded intervals represent 95% confidence intervals.

4.3 Ecological implications

Variation in how differently sized particles are consumed by salps is of great ecological

importance as it determines how salp feeding structures the plankton community around them,

but the contribution of different size classes to the total biomass ingested by the salp is also

relevant from the perspective of salp energetics. As our results have shown, the latter is not

exclusively predicated on the former and, as in all filter feeders, also depends on the prey size

structure available to feed on. For example, even if inefficient submicron feeding results in larger

numbers of bacteria being removed from the water than traditionally believed (Sutherland et al.

2010; Dadon-Pilosof et al. 2019) and smaller salps do indeed display more efficient retention for

submicron particles (Harbison and McCalister 1979; Kremer and Madin, 1992, Stukel et al.

2021), the lower total carbon content per cell attributable to small particles compared to those

that are larger and rarer but more efficiently retained and carbon-rich means these submicron

particles may still be relatively inconsequential to a salp's diet. Indeed, we found large numbers

of bacteria-like submicron particles in the salp's guts, however <1 µm particles made up only

3.2±3.8% of the average salps gut content. Consequently, the contributions by picoplankton to

salp gut content were lower than their contribution to the water column in 93% of the individuals

sampled (Figure 3). Most salps instead showed carbon-weighted mean prey sizes of 5-10 µm

(Figure 5), which is in agreement with Stukel et al.'s (2021) results for *S. thompsoni* that found

carbon-weighted median prey sizes of 8-9 µm across all cycles.

Likewise, a variety of reports suggest that small diatoms and dinoflagellates make up the

bulk of the diet for many different species of salps (Vargas and Madin 2004; Tanimura et al.

2008; Ahmad-Ishak et al. 2017), regardless of sampling location or season (von Harbou et al.

2011), implying this ~10 µm size class may represent a sweet spot between numerical abundance

in the water column and carbon content. While diatoms and thecate dinoflagellates contain hard

frustules or theca that are often resistant to digestion and may therefore be expected to contribute

disproportionately to gut contents, molecular and fatty acid composition analyses of *S. thompsoni*

and *I. racovitzai* in the Southern Ocean also found higher abundances of dinoflagellates and

diatoms in the guts (von Harbou et al. 2011; Metfies et al. 2014; Pauli et al. 2021). This contrasts

with the findings of Sutherland et al. (2010), who reported submicron particles alone could make

up more than 100% of the carbon requirement for *P. confoederata*. However, this conclusion

depended on their assumption that only the outer 0.1 µm of ingested cells were digested. When

instead assuming complete digestion of all particles, as we do here, Sutherland et al. (2010) also

found the majority of ingested carbon came from 1-10 µm particles such as nanoflagellates and

small diatoms. The true degree of digestion will vary based on many characteristics of a given

particle and the salp's gut turnover time. Until such time as when we can better predict the

nutritional values for different plankton types, our results suggest that the true mean prey size of

salps in the Southern Ocean is close to 10 µm.

5. Conclusion

   Overall, our results indicate that within the 1-16 µm prey size range, salp diets for the seven species investigated here largely reflect the size composition of plankton in the surrounding water regardless of species, size, or life stage. Feeding on submicron particles, however, appears to be dependent on both salp species and/or size. *T. democratica*, perhaps due to its smaller average size, was able to retain even submicron particles efficiently, whereas most of the larger species such as *T. vagina*, *S. thompsoni*, and *P. confoederata* showed reduced retention efficiencies below 1 µm. Size resolved retention efficiencies for *S. thompsoni* also showed this trend. Retention for particles >1 µm was generally high, with a decrease in efficiency for >16 µm particles across all salp species studied, possibly as a result of difficulty ingesting larger particles. Coupled to their prevalence in the water column, this caused nanoplankton to comprise the majority of the carbon in salp guts across all samples. This led to predator:prey size ratios ranging from 1,124:1 for the smallest *T. democratica* to 23,929:1 for the largest *T. vagina* with most falling between 1,000:1 and 10,000:1. Rather than being due to systematic differences in filtration physiology, however, our results indicate this order of magnitude variability is primarily due to the large range of sizes over which salps can occur as well as the spectra of the ambient prey field. In other words, while the ability of salps to feed on small particles is variable and ecologically important, this variation is less important with respect to salp energetics. Future work to investigate prey size spectra across a broader range of salp sizes and species is necessary to disentangle the potentially confounding impacts of these factors.

7. Acknowledgements

We would like to thank the captain and crew of the R/V Tangaroa for help in the deployment of equipment as well as our many collaborators in the SalpPOOP project including Scott Nodder, Sadie Mills, Florian Lüskow, Morgan Meyers, Sarah Searson, Lana Young, Siobhan O'Connor, Karl Safi, Adriana Lopes dos Santos, and Fenella Deans.

8. Funding

This study was made possible by funding from the Ministry for Business, Innovation and Employment (MBIE) of New Zealand, NIWA Coast and Oceans Food Webs (COES) and Ocean Flows (COOF), and the Royal Society of New Zealand Marsden Fast-track award to M. Décima, and by U.S. National Science Foundation awards OCE- 1756610 and 1756465 to M.R.S. and K.E.S.

9. Data Availability

All data is accessible on the Biological and Chemical Oceanography Data Management Office website (https://www. bco-dmo.org/project/754878).

10. Conflict of Interests

The authors declared no conflict of interests.

11. Ethical Approval

All applicable international, national, and/or institutional guidelines for sampling and experimental use of organisms for the study have been followed and all necessary approvals have been obtained.

12. Author Contributions

CKF and MRS wrote the manuscript and conducted the data analysis. KS conducted FlowCam and flow cytometry sampling. NY conducted epifluorescence microscopy sampling and image analysis. MD and CKF were responsible for net deployments and sample collection. CKF was responsible for SEM and FlowCam image analysis. All authors contributed to editing the manuscript.

7. References

Alldredge AL, Madin LP (1982) Pelagic tunicates: unique herbivores in the marine plankton. Bioscience 32(8): 655-663

Anderson V (1998) Salp and pyrosomid blooms and their importance in biogeochemical cycles. The biology of pelagic tunicates. 125-137

Barnes C, Maxwell D, Reuman DC, Jennings S (2010) Global patterns in predator–prey size relationships reveal size dependency of trophic transfer efficiency. Ecology 91(1): 222-232. http://dx.doi.org/10.1890/08-2061.1

Bathmann UV (1988) Mass occurrence of Salpa fusiformis in the spring of 1984 off Ireland : implications for sedimentation processes. Mar Biol 97:127-135

Berner LD (1967) Distributional atlas of Thaliacea in the California Current region. Calcofi Atlas No. 8, California Marine Research Committee

Bone Q (1998) The biology of pelagic tunicates.

Bone Q, Braconnot JC, Ryan KP (1991) On the pharyngeal feeding filter of the salp Pegea confoederata (Tunicata: Thaliacea). Acta Zoologica 72(1):55-60

Bone Q, Carre C, Ryan KP (2000) The endostyle and the feeding filter in salps (Tunicata). Journal of the Marine Biological Association of the United Kingdom 80(3):523-534

Bone Q, Carre C, Chang P (2003) Tunicate feeding filters. Journal of the Marine Biological Association of the United Kingdom 83(5):907-919

Botes L, Price B, Waldron M, Pitcher GC (2002) A simple and rapid scanning electron microscope preparative technique for delicate "Gymnodinioid" dinoflagellates. Microscopy research and technique 59(2):128-130

Buitenhuis ET, Li WK, Vaulot D, Lomas MW, Landry MR, Partensky F, Karl DM, Ulloa O, Campbell L, Jacquet S, Lantoine F (2012) Picophytoplankton biomass distribution in the global ocean. Earth System Science Data 4(1):37-46

Calbet A, Landry MR (2004) Phytoplankton growth, microzooplankton grazing, and carbon cycling in marine systems. Limnology and Oceanography 49(1):51-57

Cardona L, Álvarez de Quevedo I, Borrell A, Aguilar A (2012) Massive consumption of

gelatinous plankton by Mediterranean apex predators. PloS one 7(3):e31329

Conley KR, Lombard F, Sutherland KR (2018) Mammoth grazers on the ocean's minuteness: a

review of selective feeding using mucous meshes. Proceedings of the Royal Society B:

Biological Sciences 285(1878)

Crocker KM, Alldredge AL, Steinberg DK (1991) Feeding rates of the doliolid, Dolioletta

gegenbauri, on diatoms and bacteria. Journal of plankton research 13(1):77-82

Currie KI, Hunter KA (1998) Surface Water Carbon Dioxide in the Waters Associated with the

Subtropical Convergence, East of New Zealand. Deep Sea Research Part I: Oceanographic

Research Papers 45(10):1765–77. https://doi.org/https://doi.org/10.1016/S0967-0637(98)00041-

7

Dadon☐Pilosof A, Lombard F, Genin A, Sutherland KR, Yahel G (2019) Prey taxonomy rather

than size determines salp diets. Limnology and Oceanography 64(5):1996-2010

Décima M, Stukel MR, López☐López L, Landry MR (2019) The unique ecological role of

pyrosomes in the Eastern Tropical Pacific. Limnology and Oceanography 64(2): 728-743

Deibel D, Lowen B (2012) A review of the life cycles and life-history adaptations of pelagic

tunicates to environmental conditions. ICES Journal of Marine Science 69(3):358-369

Fenchel T (1987) Ecology of Protozoa: The biology of free-living phagotropic protists. Springer-

Verlag

Forman JS, Horn PL, Stevens DW (2016) Diets of deepwater oreos (Oreosomatidae) and orange

roughy Hoplostethus atlanticus. J Fish Biol 88:2275-2302

Fuchs HL, Franks PJ (2010) Plankton community properties determined by nutrients and size-

selective feeding. Marine Ecology Progress Series 413:1-15.

Garrison DL, Gowing MM, Hughes MP, Campbell L, Caron DA, Dennett MR, et al. (2000)

Microbial food web structure in the Arabian Sea: a US JGOFS study. Deep Sea Research Part II:

Topical Studies in Oceanography 47(7-8):1387-1422

Hansen B, Bjornsen PK, Hansen PJ (1994) The size ratio between planktonic predators and their

prey. Limnology and Oceanography 39(2): 395-403. https://doi.org/10.4319/lo.1994.39.2.0395

Harbison GR, Gilmer RW (1976) The feeding rates of the pelagic tunicate Pegea confederata and

two other salps 1. Limnology and Oceanography 21(4):517-528

Harbison GR, McAlister VL (1979) The filter□feeding rates and particle retention efficiencies of

three species of Cyclosalpa (Tunicata, Thaliacea) 1. Limnology and Oceanography 24(5):875-

892

Harbison GR, McAlister VL, Gilmer RW (1986) The response of the salp, Pegea confoederata,

to high levels of particulate material: Starvation in the midst of plenty 1. Limnology and

Oceanography 31(2):371-382

Heath, RA (1985) A review of the physical oceanography of the seas around New Zealand—

1982. New Zealand journal of marine and freshwater research 19:79-124.

Henschke N, Everett JD, Richardson AJ, Suthers IM (2016) Rethinking the role of salps in the

ocean. Trends in ecology & evolution 31(9):720-733

Horn PL, Burrell T, Connell A, Dunn MR (2011) A comparison of the diets of silver (Seriolella

punctata) and white (Seriolella caerulea) warehou. Mar Biol Res 7:576-591

Ishak NHA, Clementson LA, Eriksen RS, van den Enden RL, Williams GD, Swadling KM

(2017) Gut contents and isotopic profiles of Salpa fusiformis and Thalia democratica. Marine

Biology 164(6):1-10

Jennings S, Warr KJ, Mackinson S (2002) Use of size-based production and stable isotope

analyses to predict trophic transfer efficiencies and predator-prey body mass ratios in food

webs. Marine Ecology Progress Series 240:11-20

Jung SW, Joo HM, Park JS, Lee JH (2010) Development of a rapid and effective method for

preparing delicate dinoflagellates for scanning electron microscopy. Journal of applied

phycology 22(3):313-317

Katechakis A, Stibor H (2004) Feeding selectivities of the marine cladocerans Penilia avirostris,

Podon intermedius and Evadne nordmanni. Marine Biology 145(3):529-539

Koon MA, Ali KA, Speaker RM, McGrath JP, Linton EW, Steinhilb ML (2019) Preparation of

Prokaryotic and Eukaryotic Organisms Using Chemical Drying for Morphological Analysis in

Scanning Electron Microscopy (SEM). Journal of Visualized Experiments 143:e58761

Kremer P, Madin LP (1992) Particle retention efficiency of salps. Journal of Plankton

Research 14(7):1009-1015

Landry MR, Ohman MD, Goericke R, Stukel MR, Tsyrklevich K (2009) Lagrangian studies of

phytoplankton growth and grazing relationships in a coastal upwelling ecosystem off Southern

California. Progress in Oceanography 83(1-4):208-216

Lindeman RL (1942) The trophic-dynamic aspect of ecology. Ecology, 23(4):399-417

Lüskow F, Pakhomov EA, Stukel MR, Décima M (2020) Biology of Salpa thompsoni at the

Chatham Rise, New Zealand: demography, growth, and diel vertical migration. Marine

Biology 167(12):1-18

Madin LP (1974) Field observations on the feeding behavior of salps (Tunicata:

Thaliacea). Marine Biology 25(2):143-147

Madin, LP (1982) Production, composition and sedimentation of salp fecal pellets in oceanic

waters. Marine Biology 67(1):39-45

Madin, LP, Purcell JE (1992) Feeding, metabolism, and growth of Cyclosalpa bakeri in the

subarctic Pacific. Limnology and Oceanography 37(6):1236-1251

Madin, LP, Kremer P (1995) Determination of the filter-feeding rates of salps (Tunicata,

Thaliacea). ICES Journal of Marine Science 52(3-4):583-595

Madin, LP et al. (2006) Periodic swarms of the salp Salpa aspera in the Slope Water off the NE

United States: Biovolume, vertical migration, grazing, and vertical flux. Deep Sea Res. Part I

53:804-819

Menden-Deuer S, Lessard EJ (2000) Carbon to volume relationships for dinoflagellates, diatoms,

and other protist plankton. Limnology and oceanography 45(3):569-579

Metfies K, Nicolaus A, Harbou LV, Bathmann U, Peeken I (2014) Molecular analyses of gut
contents: elucidating the feeding of co-occurring salps in the Lazarev Sea from a different
perspective. Antarctic Science, 26(5):545-553

Mianzan H et al. (2001) Feeding on survival-food: gelatinous plankton as a source of food for
anchovies. Hydrobiologia 451:45-53

Michaels AF, Silver MW (1988) Primary production, sinking fluxes and the microbial food web.
Deep-Sea Research 35(4):473-490

Naustvoll LJ (2000) Prey size spectra and food preferences in thecate heterotrophic
dinoflagellates. Phycologia 39(3):187-198

Nishimura S (1958) Quelques remarques sur l'ingestion du salpe, Salpa fusiformis Curvier, chez
les maquereaux peches dans la mer du Japan orientale. Annu Rep Jap Sea Reg Fis Res Lab
4:105–112

Nodder SD, Chiswell SM, Northcote LC (2016) Annual cycles of deep☐ocean biogeochemical
export fluxes in subtropical and subantarctic waters, southwest Pacific Ocean. Journal of
Geophysical Research: Oceans 121(4):2405-2424

Pauli N, Metfies K, Pakhomov EA, Neuhaus S, Graeve M, Wenta P, Flintrop CM, Badewien
TH, Iversen MH, Meyer B (2021) Selective Feeding in Southern Ocean Key Grazers—Diet
Composition of Krill and Salps. Communications Biology 4(1):1061.
https://doi.org/10.1038/s42003-021-02581-5.

Purcell JE (1981) Dietary composition and diel feeding patterns of epipelagic siphonophores.
Mar Biol 65:83–90

Putt M, Stoecker DK (1989) An experimentally determined carbon: volume ratio for marine
"oligotrichous" ciliates from estuarine and coastal waters. Limnology and
Oceanography 34(6):1097-1103

Selph KE (2021) Enumeration of marine microbial organisms by flow cytometry using near☐UV
excitation of Hoechst 34580☐stained DNA. Limnology and Oceanography:
Methods 19(10):692-701

Sieracki CK, Sieracki ME, Yentsch CS (1998) An imaging-in-flow system for automated
analysis of marine microplankton. Marine Ecology Progress Series 168:285-296

Sheldon RW, Sutcliffe Jr WH, Paranjape MA (1977) Structure of pelagic food chain and
relationship between plankton and fish production. Journal of the Fisheries Board of
Canada 34(12):2344-2353

Sherr E, Sherr B (1988) Role of microbes in pelagic food webs: a revised concept. Limnology
and oceanography 33(5):1225-1227

Sherr EB, Sherr BF (2002) Significance of predation by protists in aquatic microbial food webs.
Antonie Van Leeuwenhoek International Journal of General and Molecular Microbiology 81(1-
4):293-308

Sherr EB, Sherr BF, McDaniel J (1991) Clearance Rates of Less-Than-6 Mu-M Fluorescently
Labeled Algae (Fla) by Estuarine Protozoa - Potential Grazing Impact of Flagellates and Ciliates.
Marine Ecology-Progress Series 69(1-2):81-92

Silver MW (1975) The habitat of Salpa fusiformis in the California Current as defined by
indicator assemblages 1. Limnology and Oceanography 20(2):230-237

Silver MW, Bruland KW (1981) Differential feeding and fecal pellet composition of salps and
pteropods, and the possible origin of the deep-sea flora and olive green 'cells'. Marine Biology
62:263-273

Stukel MR, Biard T, Krause J, Ohman MD (2018) Large Phaeodaria in the twilight zone: Their
role in the carbon cycle. Limnology and Oceanography 63(6):2579-2594

Stukel MR, Décima M, Selph KE, Gutiérrez☐Rodríguez A (2021) Size☐specific grazing and
competitive interactions between large salps and protistan grazers. Limnology and
Oceanography, 66(6):2521-2534

Sokolov S, Rintoul SR (2009) Circumpolar structure and distribution of the Antarctic
Circumpolar Current fronts: 1. Mean circumpolar paths. Journal of Geophysical Research-
Oceans 114(C11018). https://doi.org/10.1029/2008JC005108

Sutherland KR, Madin LP (2010) A comparison of filtration rates among pelagic tunicates using

kinematic measurements. Marine biology 157(4):755-764

Sutton PJ (2003) The Southland Current: a subantarctic current. New Zealand Journal of Marine

and Freshwater Research 37(3):645-652

Tanimura A, Kawaguchi S, Oka N, Nishikawa J, Toczko S, Takahashi KT et al. (2008)

Abundance and grazing impacts of krill, salps and copepods along the 140^ oE meridian in the

Southern Ocean during summer. Antarctic Science 20(4):365

Taylor AG, Landry MR, Selph KE, Yang EJ (2011) Biomass, size structure and depth

distributions of the microbial community in the eastern equatorial Pacific. Deep Sea Research

Part II: Topical Studies in Oceanography 58(3-4), 342-357

Taylor AG, Landry MR, Selph KE, Wokuluk JJ (2015) Temporal and spatial patterns of

microbial community biomass and composition in the Southern California Current

Ecosystem. Deep Sea Research Part II: Topical Studies in Oceanography 112:117-128

Vargas CA, Madin LP (2004) Zooplankton feeding ecology: clearance and ingestion rates of the

salps Thalia democratica, Cyclosalpa affinis and Salpa cylindrica on naturally occurring particles

in the Mid-Atlantic Bight. Journal of plankton research 26(7):827-833

von Harbou L, Dubischar CD, Pakhomov EA, Hunt BP, Hagen W, Bathmann UV (2011) Salps

in the Lazarev Sea, Southern Ocean: I. Feeding dynamics. Marine Biology 158(9):2009-2026

Zentara SJ, Kamykowski D (1981) Geographic variations in the relationship between silicic acid

and nitrate in the South Pacific Ocean. Deep Sea Res. Part A 28(5):455–465. doi:10.1016/0198-

0149(81)90137-0.