The *Mycobacterium tuberculosis* protein *O*-phosphorylation landscape

Authors:
Andrew Frando[1,2], Vishant Boradia[1], Marina Gritsenko[3], Michael-Claude Beltejar[1], Le Day[3],
David R. Sherman[4], Shuyi Ma[1,2,5], Jon M. Jacobs[3], Christoph Grundner[1,2,6,7*]

Affiliations:
[1]Center for Global Infectious Disease Research, Seattle Children's Research Institute, Seattle, WA.
[2]Department of Global Health, University of Washington, Seattle, WA.
[3]Pacific Northwest National Laboratory, Richland, WA.
[4]Department of Microbiology, University of Washington, Seattle, WA
[5]Department of Chemical Engineering, University of Washington, Seattle, WA
[6]Department of Pediatrics, University of Washington, Seattle, WA.
[7]Lead contact
*Correspondence

SUMMARY

Bacterial phosphosignaling has long been synonymous with the histidine kinases of the two
component systems, but many bacteria, including *Mycobacterium tuberculosis* (*Mtb*), also code
for Ser/Thr protein kinases (STPKs). STPKs are the main phosphosignaling enzymes in
eukaryotes, but the full extent of phosphorylation on protein Ser/Thr and Tyr (*O*-
phosphorylation) in bacteria remains unclear. Here, we explored the global signaling capacity of
the STPKs in *Mtb*. We generated STPK loss- and gain-of-function strains and measured the
resulting *O*-phosphorylation and transcriptional changes. This deep phosphoproteome shows
that *O*-phosphorylation in *Mtb* is an underexplored protein modification that affects >70% of the
proteome. The substrate-STPK interactions show an extensive interface with the transcriptional
machinery, resulting in regulation of gene expression of ~30% of *Mtb* genes. *Mtb O*-
phosphorylation gives rise to an expansive, distributed, and cooperative network of a complexity
that has previously only been associated with eukaryotic phosphosignaling networks.

INTRODUCTION

Bacteria signal through two branches of phosphosignaling with different chemistries of protein
modification (1,2). The His kinases of two component systems (TCSs) have long been
considered the mainstay of bacterial phosphosignaling, but phosphorylation on protein Ser/Thr
and Tyr (*O*-phosphorylation) is emerging as another important bacterial phosphosignaling
mechanism catalyzed by the Ser/Thr protein kinases (STPKs). The bacterial STPKs are similar
to the primary phosphosignaling enzymes in eukaryotes (3-5) and have been termed
"eukaryotic-like". The pervasive presence of bacterial STPKs throughout the bacterial kingdom
(6) as well as in archaea (7) and the presence of archetypical STPK sequences in bacteria,
however, all suggest a common origin at the root of the tree of life. *Mycobacterium tuberculosis*
(*Mtb*) encodes 11 STPKs, a relatively large number in relation to its genome size, and nearly as
many as TCSs (8,9). All but two STPKs are transmembrane receptors that are likely detecting
changes in the environment (8). In this way, the *Mtb* STPKs are primed to be a major conduit for
turning extracellular signals into bacterial adaptations.

Many studies have defined physiological substrates and functions for several of the *Mtb* STPKs,
in particular the essential PknA and PknB, which broadly affect transcription and metabolism
(10,11) and have central roles in coordinating replication and cell division (12). These and other
studies point to a more dominant role of *O*-phosphorylation in *Mtb* and perhaps other bacteria
than is presently recognized. Phosphoproteomic studies provided the first insights into the full
extent of *Mtb O*-phosphorylation (13,14). Despite this emerging picture of *Mtb O*-
phosphorylation as an important arm of bacterial and especially *Mtb* cell signaling, the full scope
of *O*-phosphorylation is not known, and most STPK cellular functions, signaling pathway
organization, regulatory mechanisms, and overall signaling network architecture remain to be
characterized.

The similarity between eukaryotic and bacterial STPKs has made the much better understood
eukaryotic STPKs a template for understanding STPK signaling in prokaryotes (15,16). In
eukaryotes, STPKs engage in often extensive crosstalk that can modulate pathway output, and
unlike bacterial TCSs that typically only phosphorylate their specific response regulator, can
form branched signaling networks with multiple substrates (17). Bacteria have between none
and hundreds of STPKs. For example, *S. cellulosum* codes for 317 STPKs (18), a number close
to that found in many eukaryotic organisms that perhaps also implies a similar network
organization. However, comprehensive analysis of hundreds of STPKs is challenging, and a
systems-wide analysis of a larger bacterial kinome is currently lacking. *Mtb*, on the other hand,
expresses eleven STPKs, a number that is small enough to be experimentally tractable yet
large enough to potentially give rise to a network of *O*-phosphorylation signaling. A more
complex *O*-phosphorylation network structure in *Mtb* was previously suggested by a study of *in*
vitro kinase-kinase interactions (19) and initial phosphoproteomic studies using single STPK
deletion or chemical inactivation approaches (10,11), but the presence and architecture of a
higher-order bacterial STPK *O*-phosphorylation system remains undetermined. Importantly,
STPKs are one of the major classes of drug targets in humans, with >60 drugs in the clinic (20).
A deeper understanding of *Mtb* STPK function could help translate these successes to this
human pathogen that is fueling an enduring global pandemic for which few effective
countermeasures are available.
80
To understand the full ensemble of *Mtb* STPKs and the signaling network they establish, we
generated a panel of loss (LoF) and gain-of-function (GoF) STPK mutant strains and
comprehensively characterized the STPK's phosphoproteomic and transcriptional effects. We
show that >70% of *Mtb* proteins are *O*-phosphorylated and draw a detailed map of the
substrate-kinase relationships of the STPKs, providing >3,700 high confidence substrates of
individual STPKs. Remarkably, ~30% of *Mtb* gene expression was altered by STPK
perturbation. This large impact of STPK signaling on transcription could be explained by the
pervasive interactions between STPKs, TCSs, and transcription factors (TFs), 54% of which
were differentially phosphorylated by STPKs. Together, these data provide the deepest bacterial
O-phosphoproteome to date, provide thousands of *Mtb* STPK-substrate relationships, identify a
large interface between *O*-phosphorylation and transcription in *Mtb*, predict and validate
functional STPK-TF interactions, and together establish *O*-phosphorylation as a dominant
signaling mechanism with a complexity that is comparable to eukaryotic *O*-phosphorylation
systems.
95
96
97 **RESULTS**
98
An STPK gain- and loss-of-function mutant panel

We sought to comprehensively characterize the *Mtb* phosphoproteome and test the effects of
individual STPKs by measuring changes in phosphorylation and gene expression in STPK
mutant strains. Nine of the *Mtb* STPKs have an extracellular sensor domain that reaches into
the periplasmic space (21). We anticipated that several of these have low basal activity in broth
culture and are only fully activated by signals that are likely absent *in vitro*. The loss of such an
inactive enzyme would not produce measurable effects. STPKs are typically activated by
binding to ligands that promote oligomerization, and this mode of activation appears to be
generally conserved in *Mtb* (15) and may be replicated by STPK induction. Thus, we employed
a complementary perturbation approach that combines STPK loss-of-function (LoF; knockdown
or knockout) and gain-of-function (GoF; induction) to reveal substrates for all STPKs
irrespective of their basal activation state.

All mutant strains were generated in the *Mtb* H37Rv background. GoF strains were engineered
by inserting a plasmid expressing the complete STPK coding region under control of a
tetracycline-inducible promoter. LoF strains were engineered by replacing all STPK genes
except for the *in vitro* essential *pknA* and *pknB* by a hygromycin cassette using recombineering
(22). Expression of PknB was reduced by CRISPRi (23), with the degree of knockdown titrated
so as not to affect cell growth *in vitro*. PknA was not amenable to our complementary strategy,
as the PknA kinase activity caused toxicity that precluded expression above normal levels in
Mtb. Also, a study of the phosphoproteomic consequences of *pknA* deletion was recently
presented (11), prompting us to exclude PknA from our study. We grew all cultures to early
stationary phase, as slowed replication under nutrient limitation arguably mimics restrictive *in*
vivo growth conditions more faithfully than exponential growth, and because the largest degree
of *O*-phosphorylation in *E. coli* was reported in stationary phase (24). We confirmed STPK
knockout and induction by quantitative, targeted mass spectrometry (MS) using Selected
Reaction Monitoring. We observed a range of STPK abundances in WT, with several STPKs
present at low concentrations (Fig. 1A). All STPKs were absent or reduced to below the level of
detection in the LoF strains, confirming knockout or knockdown, and all GoF mutants showed
robust induction (Fig. 1A). In total, we generated 20 STPK mutant strains, a complete LoF/GoF
panel except for PknA. To rule out altered viability of the mutant strains that may lead to a non-
specific stress or death responses, we tested the viability of mutants in early stationary phase
by regrowth after dilution in fresh medium. All mutant strains were viable and grew with wild type
(WT) kinetics, except for PknF, which grew slightly slower than WT (Fig. S1).

A deep *Mtb* phosphoproteome
To define a comprehensive *Mtb* *O*-phosphophoproteome, we analyzed WT and mutant strains
by MS-based phosphoproteomics. In pilot studies, we observed that label-free MS identified
more total phosphosites, but tandem mass tag (TMT) labeling provided more reliable
quantitation, and both methods identified unique phosphosites. To maximize the number of total
sites in combination with precise quantitation, we measured all samples by both methods.
Together, we analyzed total protein abundance and phosphorylation in three biological
replicates for each of the 20 mutant strains and WT by label-free and TMT, >150
phosphoproteomes and >10 million MS^2 spectra in total. We only considered phosphosites that
had a false discovery rate of <0.01% at the peptide level, a PEP score of <0.05, a MaxQuant
localization score >0.99 (label-free), or an Ascore of >19 (TMT), corresponding to 99%
confidence in phosphosite localization assignment. An overview of the MS workflow is given in
Fig. S2.

We detected 3,232 of the ~4,200 predicted *Mtb* proteins, providing high coverage when
compared to previous *Mtb* proteogenomic studies (25) and in light that only ~80% of *Mtb* genes
are expressed at a time (26). We detected 14,452 unique *O*-phosphorylation sites on 2,617 *Mtb*

proteins that met our statistical criteria, or 80% of detected *Mtb* proteins. These data show pervasive *O*-phosphorylation comparable to the highest level of phosphorylation detected in eukaryotes, where between 50-70% of proteins are typically found to be phosphorylated (27-29). Our study identified >5 times more phosphosites than all previously published *Mtb* phosphoproteome studies combined (Fig. 1B, Table S1) and represents the largest bacterial phosphoproteome to date. We identified 7,533 pThr, 5,816 pSer, and 1,103 pTyr sites with precise phosphosite assignment. The label-free and TMT approach each added a large number of unique phosphosites, 4,208 and 6,125, respectively (Fig. 1C). The distribution of localization scores for sites detected by label-free and TMT data showed that most sites could be assigned with high confidence, with ~50% of assignments reaching >99% confidence and over 60% reaching 90% confidence (Fig. 1D). The relative distribution of pSer, pThr, and pTyr detected in our study was 40%, 52%, and 8%, a distinct phosphorylation profile compared to either eukaryotic or the *E. coli* phosphoproteome, where pSer is the most common modification (Fig. 1E). The share of pTyr was higher than that previously detected in *Mtb* (13), and >10-fold higher than in humans (29), indicating a more prominent role of pTyr in *Mtb* (Fig. 1E). Most phosphoproteins (>500) had a single phosphosite, while 413 proteins had two, and 1,330 proteins had between 3 and 10 sites. The number of proteins with more than ten sites was small, but one group of proteins had a remarkably large number of up to 84 phosphorylation sites (Fig. 1F). These highly phosphorylated proteins included PknK, ClpC1, Pks13, and FhaA and could suggest a function of these sites outside of canonical phospho-switches, perhaps based on the physicochemical properties of a large accumulation of negative charges in a cellular subcompartment. We next determined the number of phosphoproteins that are essential as defined by *in vitro* transposon mutagenesis (30). Essential gene products were ~1.5-fold enriched in the phosphoproteome overall. However, a strong correlation between essentiality and the number of phosphorylation sites was apparent. For example, >36% of proteins with more than nine phosphosites were essential, and ~88% of those with 31-40 sites were essential (p<0.0001), compared to ~20% essential genes in the genome overall (Fig 1G).

The *Mtb* kinase-substrate relationships
We next sought to create a directory of individual STPKs and their cellular substrates. We defined substrates of a given STPK as either hypophosphorylated in that kinase's LoF or hyperphosphorylated in the GoF mutant strain, or both. Proteins with phosphorylation changes in the opposite direction were considered indirect substrates. To this end, we compared the phosphorylation pattern of each site in every STPK mutant to that of WT and calculated their abundance changes. Due to higher quality quantitation data, we used TMT data for this analysis. We defined a significant change as one detected in all three replicates, changed by at least 2-fold compared to WT, and with a p-value of <0.005. We also included on/off changes with no detected peptides in one set of replicates, although we could not generate p-values for these sites. We identified 3,503 significant changes in unique phosphosites across all mutants, or 24% of all phosphosites (Fig. 2). The overall shifts in the phosphoproteome in response to kinase perturbation were largely as expected, with GoF leading predominantly to increases in phosphorylation and LoF to decreases. Together, we detected 296 and 3,454 unique changes in phosphosites as a result of LoF and GoF, respectively (Fig. 2). The number of substrates varied largely by STPK from 15 to 968 (Fig. 2), suggesting widely varying signaling capacity of the different STPKs.

Consistent with our premise, many STPKs had low basal activity in the WT strain in broth culture. PknD, F, I, K, and L showed little substrate hypophosphorylation (<30 phosphosites) when deleted (Fig. 2, 3A). However, these silent kinases could be effectively activated to phosphorylate substrates in the GoF strains (Fig. 2, 3A), allowing for the definition of a distinct substrate set for each STPK (Table S2). We next tested if STPK induction in the GoF strains

may lead to off-target phosphorylation. We found that the average magnitude and number of
phosphorylation changes did not correlate with the abundance of the respective kinase (Fig.
S3A). For example, the PknG GoF strain, which showed the highest degree of STPK induction,
generated only 41 phosphorylation changes. These data are consistent with GoF leading to
predominantly physiologic phosphorylation events. With a total number of >14,000 unique
phosphosites, each of the STPKs would on average be expected to phosphorylate ~1,400
substrates. Only the PknK and PknL GoF strains phosphorylated as many substrates, indicating
that induction in our strains may still not fully capture the full activity for most STPKs.
210
Differential phosphorylation can result from changes in phosphoprotein abundance and/or from
changes in occupancy of phosphosites. To understand the relationship between protein
abundance and phosphosite occupancy, we determined the total abundance of proteins in all
strains by MS and correlated abundance with phosphorylation (Fig. 3B). There was no
significant correlation between the two (r=0.09) indicating that changes in phosphorylation
primarily occur on the level of site occupancy. Kinases interact with each other more than with
other protein families and can profoundly affect each other's activity by phosphorylation (17,19).
A corollary of these interactions is that phosphorylation events detected upon kinase
perturbation may be indirect, which is difficult to discriminate from direct effects (31). To identify
indirect phosphorylation events, we examined the number of hyperphosphorylated sites in the
LoF mutants and of hypophosphorylated sites in the GoF mutants as these cannot be explained
by a direct STPK-substrate interaction (Fig. 3C). While all STPKs showed some of these indirect
effects, their number was ~8% of all altered phosphosites (Fig. 3C). All indirect sites are
indicated in Table S2. Additional indirect substrates might also be among the substrates
hypophosphorylated in the LoF and hyperphosphorylated in the GoF mutant, but these are
indistinguishable from direct substrates in our data.
227
To identify potential substrate sequence preferences of the STPKs, we compiled the substrates
that showed significant changes in LoF and/or GoF mutants and identified the amino acids most
commonly found adjacent to the phosphorylation site using the pLogo tool (32) (Fig. S3B). The
Mtb STPKs' preferences were different from the often basophilic and proline-directed motifs in
eukaryotes and were generally less well-defined (33,34). Motif preferences were largely shared
among the pSer and pThr substrates, but not the pTyr substrates. Motifs associated with pSer
and pThr sites showed preferences in the -1 (D), +1 (G/D/E), +2 (P/E) position relative to the
phosphosite (Fig S3B). Charged residues (D, E, K) were generally favored in the -5 to +5
positions, and hydrophobic residues were disfavored in most positions. The preferred pTyr motif
was less clearly defined, with moderate preference for G/Q/D in -1 and hydrophobic residues (V,
L, I) in the +1 positions. To further parse individual STPKs' motifs, we mapped preferred sites of
the STPKs individually. Motifs of many STPKs predictably reflected the overall preference, and
motifs were generally weak, suggesting other substrate selectivity determinants than the
primary sequence of the phosphosite (data not shown).
242
Several STPKs are dual specificity kinases
Protein Tyr phosphorylation was long thought to be absent in *Mtb* but has now conclusively
been shown (13). Given that canonical bacterial Tyr kinases are absent in *Mtb*, the identity of
the Tyr kinases remained an open question. Based on *in vitro* biochemical activity, an atypical
kinase and dual specificity kinases among the STPKs have previously been suggested as
candidates (13,35). Our data further expanded the number of pTyr sites in *Mtb* and provided the
opportunity to test the hypothesis of dual specificity of the STPKs in a cellular context. While all
STPKs showed a preference for phosphorylating Ser/Thr over Tyr residues, many also
phosphorylated Tyr residues. The pTyr sites affected by STPK permutation account for 231
pTyr sites, or 24% of all pTyr sites detected in our study. In particular PknK and PknL LoF and

GoF mutants showed a marked decrease and increase of pTyr, respectively (Fig. 3D). These
data suggest that some *Mtb* STPKs have evolved dual activity similar to the dual specificity
kinases in many eukaryotes to perform a protein modification that is typically executed by
specialized bacterial Tyr kinases.
257
The STPKs constitute a distributed network
We next analyzed the STPK-substrate relationships globally to detect larger features of the
network by visualizing all STPK-substrate interactions. While 469 substrate proteins were only
phosphorylated by a single STPK, 935 were phosphorylated by more than one STPK (Fig. 3E).
Of the proteins phosphorylated by more than one STPK, most were phosphorylated by two
STPKs (292), and one, Rv0020c, was phosphorylated by all ten STPKs. The high redundancy
of this phosphorylation network may explain the relative paucity of *in vitro* phenotypes of KO
strains of most of the STPKs despite their large number of substrates and is reminiscent of
distributed eukaryotic phosphosignaling networks (36). We next inspected the STPK substrates
showing significant changes upon kinase perturbation for enrichment in several protein families
and GO terms. Among the most significantly enriched were components of the proteasome and
Clp protease systems (Fig. 3F), and RNA Polymerase (Fig. 3G), almost all of which were
phosphorylated by multiple STPKs suggesting coordinated regulation by several STPK inputs of
these central cellular functions.
272
STPKs have global transcriptional effects
We identified 158 phosphorylated TFs (77% of *Mtb* TFs (37)) and 112 TFs (54% of TFs) with
significant phosphorylation changes associated with STPK perturbation, suggesting potentially
widespread regulation of TF activity by *O*-phosphorylation (Fig. 4A). In a simple transcriptional
relay, an STPK and gene expression can be directly linked by the phosphorylation of a TF,
leading to gain or loss of DNA binding (38). The extensive phosphorylation of TFs predicted that
the transcriptional effects of the STPKs could be significant. To test this idea, we measured the
STPKs' downstream transcriptional effects by RNA-seq using samples matched to the samples
used for phosphoproteomics. We used a cutoff of at least 4-fold up- or down-regulation
compared to WT and a significance cutoff of p <0.01 to define differentially expressed genes
(DEGs).
284
Overall, 1,155 of 4,030 detected genes met our criteria for DEG, representing ~30% of *Mtb*
genes (Fig. 4B, S4). All STPKs caused changes in gene expression and two distinct kinase
activation patterns were apparent: Several STPKs showed a larger change in gene expression
upon LoF (PknB, G), suggesting a constitutive housekeeping function. In particular PknB elicited
differential expression of 30-fold more genes in the LoF strain than the GoF strain (Fig. 4B, S4).
Another group of STPKs including PknD, F, H and J showed no or little expression changes
upon LoF, but large changes upon GoF (Fig 4B, S4). These data mirror findings from our
phosphoproteomic analysis showing that some STPKs are inactive in culture and likely require a
signal for activation that is absent *in vitro*. Except for PknE and PknF, the number of
differentially regulated phosphosites and the number of DEGs corresponded well (r=0.92) (Fig.
S5A). The range of fold-change varied widely to several hundred-fold (Fig. 4C). Although
proteomic analyses described above suggested no or little artificial phosphorylation of the GoF
mutants, we re-examined this question using transcriptional data. To test if the abundance of
STPKs in the GoF mutants was within physiologic levels, we mined existing transcriptional data
for STPK expression during stress conditions and found that in most cases, STPK expression
levels in the GoF mutants were within the range found in WT bacteria subjected to physiologic
stresses such as hypoxia, oxidative stress, and nitrosative stress (Fig. S5B). The PknK GoF
mutant showed the largest difference from PknK induction under stress in WT, however, the
levels of induction of 8 STPKs in our GoF strains were in fact below those reported in WT under

specific stresses. Non-specific transcriptional effects of the STPKs would also suggest a correlation between the magnitude of STPK expression and the number of DEGs. The number of differentially expressed genes was not correlated with induction of the STPK (r=0.37) (Fig. S5C). To test if the transcriptional effects of STPK perturbation are indeed due to the kinase activity of the STPKs, we measured the transcriptional effects of a kinase-dead PknL mutant (Fig. S6A). While the PknL GoF mutant led to 260 changes in gene expression compared to WT, the kinase dead mutant led to one, showing that the kinase activity is responsible for gene expression changes. Domain prediction showed that in addition to the kinase domain, PknK also has a transcriptional regulator domain of the MalT family. To test if gene expression changes in the PknK GoF strain data reflect effects of the kinase activity or perhaps also effects of the MalT domain, we compared DEGs in strains overexpressing PknK WT and kinase dead mutant. The WT GoF mutant showed changes in 181 genes, the kinase dead mutant in 13, suggesting that <10% of DEGs in the PknK GoF mutant may be due to the activity of the MalT domain (Fig. S6B).

To understand the higher-level organization of the STPKs' transcriptional effects, we visualized the DEGs regulated by STPK LoF and GoF mutants (Fig. S7). While some genes were affected by a single STPK, 40% of genes were affected by several STPKs, some by up to seven. The co-regulation of these genes showed that signaling by multiple *Mtb* STPKs converges on the same cellular pathways and suggests a high degree of redundancy between the STPKs. We observed co-regulation in the same and in the opposite direction. For example, Rv2590 was induced in PknG and PknL LoF mutants while Rv1875 was repressed in PknE and PknK LoF mutants, both cases of redundant kinase function. In contrast, Rv1739c was upregulated in the PknE LoF mutant and downregulated in the PknB LoF mutant. The connectedness of STPK gene expression given as the average number of STPKs that affect expression of one gene was on average 1.75. These data reveal a large, highly coordinated interface between STPK signaling and transcription.

Integration of phosphoproteomic and transcriptional data predicts functional TF phosphosites

The share of functional phosphorylation events in any phosphoproteome is unknown, and their prediction remains a major challenge. Because we determined STPK-TF substrate pairs and global functional readouts for the STPKs by measuring transcriptional changes, our data offered an opportunity to predict function of phosphorylation events on TFs. To this end, we integrated our phosphoproteomic data on STPK-TF kinase-substrate pairs, the matched STPK transcriptional data, and a global transcriptional dataset mapping the transcriptional effects of ~200 *Mtb* TFs (37). A significant overlap between the regulon of an STPK and that of its cognate TF substrate would suggest that the phosphorylation site is functional. To correlate regulons of phosphorylated TFs and those of their cognate STPKs, we determined the genes altered by an STPK and its TF partner and plotted the % overlap of the TFs regulon with the STPK's regulon (Fig. 5). 52 TFs showed significant regulon overlap (p=0.05) with their cognate STPK's regulon. 24 TFs showed >50% of shared regulated genes, and the regulons of 3 TFs (hns, pyrR and Rv0792c) were fully contained in their STPK's regulon. These data predict a set of TFs whose phosphorylation is likely regulatory and leads to downstream gene expression changes. The strength of the predictions increase with higher regulon overlap and lower STPK regulon size (Fig. 5).

O-phosphorylation alters Zur transcription factor activity

To validate the STPK-TF functional predictions and to illustrate how these predictions can be used to assemble signaling pathways, we explored the interaction between PknK and the Zinc uptake repressor Zur (Rv2359), a negative transcriptional regulator of metal homeostasis (39).

The PknK GoF mutant showed significant Zur hyperphosphorylation on Thr67 in our MS data (8.7-fold, p-value=0.001) when compared to WT, consistent with a direct kinase-substrate interaction, and the phosphorylation event was predicted to be functional based on significant regulon overlap in our STPK-TF regulon analysis (Fig. 6). We validated the PknK-Zur interaction using an *in vitro* kinase assay with recombinant PknK and Zur. PknK efficiently phosphorylated Zur *in vitro* (Fig. 6B), but PknG, which did not affect Zur phosphorylation in our phosphoproteomic data, did not. To test a regulatory connection between PknK and Zur, we first compared the previously identified Zur LoF and GoF regulons (37,39) to genes regulated in the PknK GoF mutant and found that >80% of the Zur regulon is also regulated in the PknK GoF mutant, suggesting a function in the same pathway (Fig. 6C). While the Zur GoF mutant strain repressed its regulon consistent with its role as a negative regulator, the PknK GoF mutant strain led to induction of nearly all genes in the Zur regulon. The opposite effects of Zur and PknK GoF predicted that phosphorylation of Zur by PknK derepresses Zur. Structural analysis using a high-confidence model based on the *E. coli* Zur structure bound to DNA showed that Thr67 is adjacent to the DNA-binding helix-loop-helix motif of Zur and faces directly towards the minor groove of target DNA. In its phosphorylated form, Thr67 would likely create charge repulsion with the phosphatedeoxyribose backbone of the DNA, predicting a negative effect of phosphorylation on DNA binding. To test the effect of Zur Thr67 phosphorylation on its ability to bind DNA, we purified recombinant WT, a Thr67 phosphoablative (Thr67Ala) and phosphomimetic (Thr67Asp) mutant and tested DNA-binding activity by electrophoretic mobility shift assays (Fig. 6D). WT Zur efficiently bound to DNA, resulting in a shift of the DNA probe. The phosphoablative mutant did not show any difference to WT in DNA-binding. The phosphomimetic mutant, however, showed decreased DNA-binding, consistent with our prediction. Together, these data define a phosphoregulation pathway in which PknK phosphorylates Thr67 of Zur, resulting in loss of DNA binding, which de-represses the Zur regulon (Fig. 6E).

A combined transcriptional and phosphoproteomic resource for studying kinase function
The matched kinome-wide phosphoproteomic and transcriptomic datasets provide the set of phosphorylation sites and genes that are affected by individual STPKs. An overview of the combined interactions is shown in Fig. 7. Our data assign STPKs to substrates and genes and predict STPK functions. Conversely, these data allow for the identification of candidate STPKs affecting a particular physiologic response if that response is characterized by the expression of a gene set or a change in phosphorylation. In addition to several STPKs for which no experimental functional data is available to date, in particular the GoF data furnishes a large set of phosphosites that were not accessible before. The complete dataset showing all transcriptional and phosphorylation effects of the STPK GoF and LoF mutants are presented in Table S1 (all phosphosites, label-free and TMT), Table S2 (regulated phosphosites), and Table S3 (transcriptional changes). Together, these data show that a large swath of *Mtb* physiology is integrated with STPK signaling, provide a resource for assigning kinases to substrates and regulated genes and thus the foundation for detailed mechanistic understanding of this pervasive *Mtb* phosphosignaling mechanism.

DISCUSSION

The contribution of protein *O*-phosphorylation to bacterial cell signaling remains an open question, and the relatively small numbers of *O*-phosphorylation sites previously reported in bacteria have reinforced the notion that bacterial *O*-phosphorylation systems are simpler and more rudimentary than its eukaryotic counterparts. Here, we challenge this view with the deepest phosphoproteome measured in a bacterium yet, which expands the number of reported

Mtb O-phosphorylation sites by >5-fold. The degree of *O*-phosphorylation in *Mtb*, with >70% of
proteins decorated by at least one phosphosite, is on par with that found in many eukaryotic
organisms (28), and shows a signaling network of a complexity that was thought to be exclusive
to eukaryotes. These findings place *O*-phosphorylation at the center of cellular signaling in the
pandemic human pathogen *Mtb* and argue that the STPKs constitute the backbone of bacterial
phosphosignaling in *Mtb*.
412
To further understand phosphosignaling beyond the general inventory of the phosphoproteome
requires parsing of contributions of individual kinases and their substrates, or the kinase-
substrate relationships. Assigning these kinase-substrate relationships remains a major
experimental bottleneck. Although several substrates, partial pathways, and functions of the *Mtb*
STPKs have previously been identified, they remain unknown for most. Global
phosphoproteomic approaches are arguably best suited to identify kinase-substrate
relationships. These global approaches typically use LoF perturbations and then measure the
resulting changes in the phosphoproteome by MS (40). While the changes resulting from a LoF
perturbation are highly informative, this approach requires a baseline activity of a kinase in the
experimental condition, which was an untested premise for most *Mtb* kinases grown in broth
culture. Given that *Mtb* STPKs are mostly sensor kinases that likely survey the host, it is not
surprising that many are largely inactive in broth culture, where host-specific signals are absent.
Our GoF approach captured many of these previously intractable phosphorylation events and
downstream transcriptional effects of these silent STPKs.
427
Our study vastly expands the overall number of *Mtb O*-phosphorylation sites and candidate
substrates for the individual STPKs, including for PknB that has been the subject of dozens of
studies including one that applied LoF-MS-based approaches that are comparable to ours (11).
The higher number of phosphosites identified in our study is likely a result of high-sensitivity MS
hardware, using label-free and TMT approaches combined with extensive sample fractionation,
and the analysis of GoF mutants. The extensive substrate sets for the STPKs presented here
provide a directory for identifying STPK pathways and functions. One global functional
consequence of STPK signaling in particular was the sweeping effect on gene expression, with
~30% of all *Mtb* genes affected by one or more STPKs. The most direct connection between
STPKs and gene expression is through the phosphorylation of TFs, whose DNA binding activity
can change upon phosphorylation. Here, we uncover extensive phosphorylation of TFs by the
STPKs. A key question for these and in fact most known phosphosites is whether they are
functional. Prediction of functional phosphorylation sites is exceedingly difficult. Our data, by
measuring both STPK-TF pairs and transcriptional effects, allowed for the prediction of
functional phosphorylation sites on TFs and identifies putative signaling pathways from kinase
to gene. Most TF phosphorylation events, however, did not produce significant gene regulatory
changes that could be ascribed to its cognate STPK. The lack of detectable function of these TF
sites may be due to the limited, one-directional perturbation in the TF dataset (37), indirect
effects of STPK and/or TF, functions other than transcription, but likely also reflects a large
degree of phosphorylation with no function. These data provide a rare experimental datapoint
for gauging the share of functional phosphorylation sites within a large protein family.
449
The many STPK-TF connections predicted to be functional and the regulation of Zur by PknK
suggest a general model of STPK-mediated gene regulation in *Mtb* —direct regulation of TFs by
STPKs. The number of these interactions was surprising given the few previous examples of
STPK-TF interactions reported in *Mtb*. Other direct interactions between STPKs and the
transcriptional machinery suggested by our data include the STPK-specific phosphorylation of
sigma factors, RNA polymerase, but also of TCSs. TCSs and STPKs were thought to be largely
insulated from one another, but the extensive phosphorylation of His kinases and response

regulators by the STPKs suggest close integration of the two arms of phosphosignaling. The
complex and highly redundant architecture of the STPK signaling network may be better suited
to translate complex signals into effective and robust responses than the typically simpler and
more linear TCS pathways. The remarkable interconnectivity between signal transduction and
gene regulation underscores how understanding the interplay of these networks will be critical to
dissect how *Mtb* and potentially other bacteria sense, process, and respond to environmental
cues.
464
MS data are probabilistic in nature. We sought to minimize the uncertainty inherent in our
phosphoproteomic data by setting stringent cutoffs for peptide identification, false discovery
rate, and phosphosite localization. One limitation of our study is the analysis of the
phosphoproteome in only one *in vitro* condition, stationary phase. In addition, gene KO can lead
to compensatory effects that can obscure STPK functions. Further, the large effects of the
STPK GoF mutants on phosphorylation and transcription also raise the question of experimental
artifacts. However, we found no evidence for capricious phosphorylation. The induction levels of
STPKs in the GoF mutants were within the range elicited in WT responses to physiologic stress
conditions, and the frequency of phosphorylation did not correlate with STPK expression levels.
In addition, most changes in the GoF strains were in fact changes in the abundance of
phosphosites that were also present in WT. The likelihood of an inadvertent functional impact of
erratic phosphosites appear small: Phosphosignals propagate through evolved protein-protein
interactions that are unlikely to be replicated by artificial phosphorylation events, and the low
occupancy of erratic phosphorylation would further reduce phenotypic consequences. However,
a phosphosite or phenotypic effect that can be attributed to a LoF mutation may ultimately be
more reliable than an effect observed in a GoF mutant, and corresponding effects of LoF and
GoF mutants can further increase the confidence in the detected change.
482
On a systems level, this study shows a central and dominant place of *O*-phosphorylation in the
important human pathogen *Mtb* and a cooperative signaling system of a complexity that is
typically only associated with eukaryotes. These findings revise, at least for *Mtb*, the long-held
view of the TCSs as the canonical bacterial phosphosignaling system. On a molecular level, this
study provides rich detail of this system through a resource with >14,000 unique phosphosites
and >3,700 STPK-specific phosphosite changes and their transcriptional effects. These data lay
a foundation for understanding the cellular functions of these enzymes and provide signaling
context for a large part of *Mtb* physiology.
491
492
493 **FIGURE LEGENDS**
494
Figure 1: A deep *Mtb* phosphoproteome
(A) Quantitation of STPK protein levels in LoF and GoF *Mtb* mutant strains by selected reaction
monitoring. PknB LoF is a knockdown strain, all other LoF strains are knockouts.
(B) Comparison of the number of unique protein *O*-phosphorylation sites and -proteins
previously published and identified in this study.
(C) Number of unique phosphosites identified by TMT and label-free approaches.
(D) Distribution of phosphosite localization Ascore (for TMT data) and MaxQuant probability (for
label-free data). An Ascore of 19 corresponds to 99% probability of correct localization.
(E) Relative share of pSer, pThr, and pTyr sites in *Mtb*, *E. coli*, and human cells.
(F) Number of phosphorylation sites per protein and highly phosphorylated proteins with >20
phosphosites (inset).
(G) Share of phosphorylated essential gene products stratified by number of phosphosites.
Essentiality was based on transposon mutagenesis (30).

508
Figure 2: STPK perturbation defines kinase-substrate interactions
Magnitude and direction of changes in phosphorylation sites in LoF (Rows 1 and 3) and GoF
strains (Rows 2 and 4). Volcano plots show $\log_2$-fold changes in phosphorylation levels by
phosphosite (X-axis) versus the -log p value of the change (y-axis). Horizontal and vertical lines
show significance cutoffs applied to the data for further analysis (>2-fold change in abundance,
p<0.005). Significantly changing phosphosites are shown in red and blue and their numbers are
given in the respective quadrant. The STPKs altered in the respective strain were not included
as they showed the largest changes and distorted the plots.

Figure 3: *Mtb* protein *O*-phosphorylation establishes a network
(A) Effects of STPK perturbation on the abundance of unique phosphosites by STPK compared
to WT. Changes in LoF and GoF strains were combined for each STPK.
(B) Correlation between global protein abundance and phosphosite abundance of all
phosphorylation sites showing a change in STPK LoF and GoF mutants. Quantitative TMT data
of total protein abundance was plotted against the phosphosite abundance for each protein,
showing that most increases in phosphorylation are not concomitant with higher protein
abundance.
(C) Directionality of phosphorylation changes in response to STPK perturbation. The volcano
plots show assignment of direct (hypophosphorylation in LoF, hyperphosphorylation in GoF),
and indirect phosphorylation events with inverted directionality. Table shows putative direct and
indirect phosphosites (phosphoproteins) by STPK.
(D) Changes in Tyr phosphorylation sites upon STPK perturbation. The numbers given above
the STPK mutant strain is from LoF and GoF mutants combined.
(E) An overview of the STPKs' substrates and overlap in substrate phosphorylation. The purple
squares represent the STPKs. The size of the square is proportional to the number of that
STPK's substrates, which is also given in parentheses. The round nodes show the number of
substrates phosphorylated by one or multiple STPKs. The thickness of the edge corresponds to
the number of substrates phosphorylated by the respective STPK.
(F) Phosphorylation of members of the proteasome/Clp protein degradation systems by different
STPKs.
(G) Phosphorylation of RNA polymerase subunits by multiple STPKs.

Figure 4: *O*-phosphorylation has large transcriptional effects
(A) Overview of TF phosphorylation by STPKs. STPKs are shown as purple squares, TFs are
shown as green circles. Edges represent kinase-substrate interactions. No Ascore cutoff was
applied as analysis was on the basis of phosphoproteins, not phosphosites. Table shows
number of TF substrates for each STPK.
(B) Overall number of differentially expressed genes upon STPK perturbation by STPK and by
type of perturbation. Differentially expressed genes are defined as >4-fold change with p<0.01.
(C) The most induced and repressed genes by STPK and type of perturbation.

Figure 5: Prediction of functional TF phosphorylation events
Genes regulated by both TFs and their cognate STPK were plotted as % regulon overlap
against the significance of the overlap. Colors denote different STPKs. Dotted line indicates a p-
value of 0.05. These data are based on protein-level phosphorylation and no localization score
cutoff was applied.

Figure 6: PknK regulates the transcription factor Zur
(A) *In vitro* phosphorylation assay showing specific phosphorylation of recombinant Zur by
PknK, but not PknG.

559 (B) Transcriptional response of Zur LoF, Zur GoF, and PknK GoF mutations on the Zur operon
560 shows overlapping but opposite effects of PknK and Zur GoF, predicting derepression of Zur
561 activity by PknK phosphorylation.
562 (C) Effect of Zur Thr67 phosphoablative and -mimetic mutations on DNA binding activity.
563 (D) Model of the PknK-Zur pathway.
564
Figure 7: Overview of the *Mtb* STPK *O*-phosphoproteome and transcriptional network.
The phosphoproteomic and transcriptional effects of STPK mutation are shown, illustrating the
strongly overlapping nature of the network. The STPKs are shown as round nodes, with their
total number of phosphorylated substrates in parentheses. The outer circle of each STPK node
shows the relative share of substrates observed in the LoF (orange), GoF (green), or both
(down in LoF, up in GoF, yellow). The color of the inner STPK circle indicates whether the
STPK's effect on DEGs on average is induction (red), repression (blue), or a combination
(purple). The inner square nodes represent genes regulated by the STPKs, with the thickness of
edges connecting the STPKs to genes corresponding to the number of genes regulated by an
STPK. Single and multiple refers to the number of STPKs that have the particular effect on the
group of genes (for example, 130 genes are repressed by more than one STPK). Interestingly,
the expression of 61 genes is activated by some STPKs and repressed by others.
577
578
579 **SUPPLEMENTAL FIGURE AND TABLE LEGENDS**
580
581 **Figure S1: Overview of the MS experimental workflow.**
582
Figure S2: Fitness test of STPK mutant strains.
(A) LoF strains were grown to stationary phase, diluted to OD_{600} of 0.25 and regrown to test for
fitness. The PknB LoF knockdown mutant was induced with ATc.
(B) GoF mutants were induced with ATc, grown to stationary phase, diluted to OD_{600} of 0.25,
and regrown without ATc.
Figure S3: Correlation of phosphosites and STPK protein abundance and motif analysis.
(A) The abundance of STPKs in the GoF mutants does not overall correlate with the number of
phosphosites detected in these strains.
(B) Preferred phosphorylation motifs of all STPKs combined and by phosphoresidue. Height of
sequence logo represents frequency of residue at that position.
593
Figure S4: Transcriptional effects of STPK perturbation.
Volcano plots show the DEGs in STPK mutant strains, direction of change, magnitude, and p-
value associated with the change. Horizontal and vertical red lines show significance cutoffs
(>4-fold, p<0.01) used for further analysis of DEGs. The STPKs that were altered in the
respective strain were removed from the data as they showed the largest changes and distorted
the plots.
600
Figure S5: Relationship between STPK induction, phosphosites, and DEGs.
(A) More phosphorylation sites in the GoF mutant strains led to more DEGs. PknJ affects the
expression of a large number of genes through few phosphosites. Red line shows linear
regression if outliers PknE and PknF are excluded.
(B) The induction of STPKs in the GoF strains as determined by RNA-seq was within the range
observed in historical data (41-45) under physiologic stress conditions for most STPKs. PknK
and PknL showed higher induction, most STPKs showed lower induction, suggesting largely
physiologic levels of STPK in the GoF strains.

(C) STPK abundance is not correlated with the number of DEGs in the GoF strains.
610
Figure S6: Transcriptional effects of the PknL and PknK kinase dead mutants
(A) The PknL kinase dead mutant reduces the DEGs compared to WT GoF from 260 to 1. PknL
WT and kinase dead transcripts are highlighted in green, showing induction in both cases.
(B) The PknK kinase dead mutant reduces DEGs compared to WT GoF from 181 to 13,
suggesting a small transcriptional effect of the PknK malT domain in the GoF. PknK WT and
kinase dead transcripts are highlighted in green, showing induction in both cases.
617
618 **Figure S7: STPK signaling is highly integrated with transcription.**
619 STPKs (purple nodes) affect the expression of a large number of genes (number in
620 parentheses). Nodes in the center represent the genes grouped by the number of STPKs they
621 are regulated by. The number in parentheses indicates the number of genes in that group,
622 illustrating the redundancy and/or cooperativity of the network.
623
624 **Table S1: Total and unique phosphosites.**
625
626 **Table S2: The STPK-related *O*-phosphorylation changes.**
627
628 **Table S3: The STPK-related differentially expressed genes.**
629
630
631 **MATERIALS AND METHODS**
632
***Mtb* STPK mutant strain generation**
Loss-of-function (LoF) mutants were constructed by recombineering (22). The recombineering
substrate was generated by amplifying 500 bp upstream and downstream of each STPK's
coding sequence by PCR and cloning them 5' and 3' of a hygromycin-resistance cassette,
respectively. A linear recombineering substrate was amplified by PCR, purified, and transformed
into *Mtb* strain H37Rv carrying pNIT-ETc (46). Successful integration was confirmed using DNA
sequencing. Gain-of-function (GoF) mutants were constructed using the episomal pDTCF
plasmid encoding a tetracycline-inducible promoter, C-terminal FLAG tag, and a hygromycin-
selectable marker. STPKs were amplified from *Mtb* strain H37Rv genomic DNA, cloned into the
pDTCF plasmid 5' of the FLAG tag using Gibson assembly cloning, and confirmed with
sequencing. Plasmids were transformed into *Mtb* and expression was confirmed by anti-FLAG
Western blot. The *pknB* CRISPRi-knockdown strain was constructed using a pJR965 plasmid
encoding a tetracycline-inducible dCas9, a kanamycin-selectable marker, and a tetracycline-
inducible sgRNA targeting the 5' coding region of *pknB*. STPK LoF and GoF were confirmed by
DNA sequencing and selected reaction monitoring and label-free LC-MS/MS.
648
***Mtb* culture**
All strains were grown in Middlebrook 7H9 with (10%) OADC and (0.2%) glycerol at 37°C in
rolling culture. For experiments involving STPK GoF, cultures were grown in 50 µg/ml
hygromycin B. For experiments involving STPK CRISPRi knockdown, cultures were grown in 30
µg/ml kanamycin. WT and STPK LoF strains were grown to stationary phase in triplicate. STPK
GoF and CRISPRi knockdown strains were grown in triplicate to OD_{600} 0.4-0.6, induced with
100 ng/ml anhydrotetracycline, and grown to stationary phase.
656
Lysate production
Mtb cultures were pelleted at 4,000 x g for 5 min at 4°C, washed with phosphate buffered saline,
and resuspended in lysis buffer (8 M Urea; 50 mM Tris pH 8.0; 75 mM NaCl; cOmplete protease

inhibitor cocktail (Sigma 5892791001); 10 mM sodium fluoride (Sigma 919), 1% phosphatase
inhibitor cocktail 2 (Sigma, P 5726), 1% phosphatase inhibitor cocktail 3 (Sigma, P 0044)).
Lysates were generated by bead-beating three times at 6 m/s for 30 sec with cooling on ice
between steps. Samples were centrifuged at 12,000 x g for 1 min, and the supernatant sterilized
by filtration. Protein concentrations were determined by Bradford assay and the lysates stored
at -80°C until further processing.
666
Proteomic and phosphoproteomic sample processing
1 mg of protein from each sample in 1 ml lysis buffer was incubated for 15 min at 4°C. Protein
concentrations were determined by BCA assay (ThermoFisher Scientific). Proteins were
reduced with 5 mM dithiothreitol for 1 h at 37 °C, and subsequently alkylated with 10 mM
iodoacetamide for 45 min at 25°C in the dark. Samples were diluted 4-fold with 50 mM Tris-HCl,
pH 8.0 to decrease a final concentration of urea below 2 M for the digesting with Lys-C (Wako)
at 1:50 enzyme-to-substrate ratio. After 2 h of Lys-C digestion at 25 °C, sequencing grade
modified trypsin (Promega, V5117) at 1:50 enzyme-to-substrate ratio was added to the samples
and samples were incubated at 25°C for 14 h. The reaction was stopped by acidifying the
samples with 100% formic acid (Sigma-Aldrich, St. Louis, MO) to a final concentration of 1%
formic acid. Acidified samples were centrifuged for 15 min at 1,500 g to clear digest from
possible precipitation. Tryptic peptides were desalted on 200-mg tC18 SepPak (Waters,
WAT054925) SPE and concentrated down using Speed-Vac concentrator. Final peptide
concentration was determined via BCA assay. Desalted peptide samples from each sample
were labeled with 10-plex tandem mass tags (TMT, ThermoFisher Scientific) with one of the
TMT channels (131) occupied with a pooled mixture of peptides from all the samples, which
serves as a reference to normalize across different sets of samples. Peptides (400 μg) were
dissolved in 80 μl of 50 mM HEPES, pH 8.5 solution, and mixed with 400 μg of TMT reagent
that was dissolved freshly in 20 μl of anhydrous acetonitrile. To quench the labeling reaction, 12
μl of 5% hydroxylamine was added to the samples before for 15 min at RT incubation. Peptides
labeled by different TMT reagents were then mixed, dried in Speed-Vac concentrator,
reconstituted with 3% acetonitrile, 0.1% formic acid and desalted on tC18 SepPak SPE
columns. 3.5 mg of 10-plex TMT-labeled sample was separated on a reversed phase Agilent
Zorbax 300 Extend-C18 column (250 mm × 4.6 mm column containing 3.5-μm particles) using
Agilent 1200 HPLC System. Solvent A was 4.5 mM ammonium formate, pH 10, 2% acetonitrile
and solvent B was 4.5 mM ammonium formate, pH 10, 90% acetonitrile. The flow rate was 1
ml/min and the injection volume was 900 μL. The LC gradient started with a linear increase of
solvent B to 16% in 6 min, then linearly increased to 40% B in 60 min, 4 min to 44% B, 5 min to
60% B and another 14 of 60% solvent B. A total of 96 fractions were collected into a 96 well
plate throughout the LC gradient. These fractions were concatenated into 6 fractions by
combining 16 fractions into one for a total of 6 fractions. For proteome analysis, 5% of each
concatenated fraction was dried and re-suspended in 2% acetonitrile, 0.1% formic acid to a
peptide concentration of 0.1 mg/ml for LC-MS/MS analysis. The rest of the fractions (95%) were
subjected to immobilized metal affinity chromatography (IMAC) for phosphopeptide enrichment
(47,48).
702
Phosphopeptide enrichment
Fe3+-NTA-agarose beads were freshly prepared using the Ni-NTA Superflow agarose beads
(QIAGEN, #30410) for phosphopeptide enrichment. For each of the 12 fractions (~ 200 ug per
each fraction) and for each label-free sample (100 ug), peptides were reconstituted to 0.5 μg/μl
in IMAC binding/wash buffer (80% acetonitrile, 0.1% trifluoroacetic acid) and incubated with 10
μl of the Fe3+-NTA-agarose beads for 30 min at RT. After incubation, the beads were washed 2
times each with 50 μl of wash buffer and once with 50 μl of 1% formic acid on the stage tip
packed with 2 discs of Empore C18 material (Empore Octadecyl C18, 47 mm; CDS Analytical,

98-0604-0217-3). Phosphopeptides were eluted from the beads on C18 using 70 µl of Elution
Buffer (500 mM potassium phosphate buffer). 50% acetonitrile, 0.1% formic acid was used for
elution of phosphopeptides from the C18 stage tips. Samples were dried using Speed-Vac, and
later reconstitute with 12 µl of 3% acetonitrile, 0.1% formic acid containing 0.01% DDM (n-
Dodecyl-beta-Maltoside) for LC-MS/MS analysis.
716

Global and phosphoproteome analysis
Both TMT and label-free global and phosphopeptide enriched samples were subjected to a
custom high mass accuracy LC-MS/MS system as previously described (11), where the LC
component consisted of automated reversed-phase columns prepared in-house by slurry
packing 3-µm Jupiter C18 (Phenomenex) into 35-cm x 360 µm o.d. x 75 µm i.d fused silica
(Polymicro Technologies Inc.). The MS component consisted of a Q Exactive HF Hybrid
Quadrupole-Orbitrap mass spectrometer (Thermo Scientific) outfitted with a custom
electrospray ionization interface. Electrospray emitters were custom made using 360 µm o.d. x
20 µm i.d. chemically etched fused silica capillary. Analysis of the phosphoproteome samples
applied similar conditions as used in the global proteome sample analysis, except that the spray
voltage was 2.2 kV. All other instrument conditions were set as previously described (49). Raw
spectral data and analysis information is available via the MassIVE database for the public
accession #MSV000088254. LC–MS/MS raw data were converted into dta files using Bioworks
Cluster 3.2 (Thermo Fisher Scientific, Cambridge, MA, USA). The MSGF+ algorithm was used
to search MS/MS spectra against the *Mtb* database (RefSeq H37Rv_uid57777_2014-08-14,
20198 entries). Search parameters included 20 ppm tolerance for precursor ion masses, +2.5
Da and -1.5 Da window on fragment ion mass tolerances, no limit on missed cleavages, partial
tryptic search, no exclusion of contaminants, dynamic oxidation of methionine (15.9949 Da),
static IAA alkylation on cysteine (57.0215 Da), and static TMT modification of lysine and N-
termini (+144.1021 Da) for TMT analyses. The decoy database searching methodology (50,51)
was used to control the false discovery rate at the unique peptide level to <0.01% and
subsequent protein level to <0.1% (52). Quantification for TMT was based upon initially
summing to the peptide (phospho) or protein (global) level the sample specific peptide reporter
ion intensities captured for each channel across all 24 or 12 analytical fractions. Final data for
statistical analysis was the scaling and central tendency normalization of each peptide or protein
summed value across all observations within each TMT10 experiment to adjust for experiment
specific variability. Label-free quantification used MaxQuant (53) generated LFQ values at the
peptide (phospho) and protein (global) levels as previously described (54). All collected protein
and peptide abundance values were subjected to statistical comparison using WT as the basis
for comparison. Primary comparisons were based upon ANOVA analysis at a <0.005 p-value
threshold coupled with a fold change criteria minimum of +/- 2 between abundances.
748

Selected reaction monitoring
SRM was performed on the panel of WT, GoF, and LoF mutants as described above. Crude
heavy peptides labeled with $^{13}C/^{15}N$ on C-terminal lysine and arginine were purchased from
vivitide (Gardner, MA). Trypsin digested samples that had been stored at -80°C until use were
processed as previously described (55). For each sample the digested peptides were diluted to
0.25 µg/µl containing standards at a final concentration of 20 fmol/µl. All the samples were
analyzed with a nanoACQUITY UPLC system (Waters Cooperation, Milford, MA) coupled online
to a TSQ Altis triple quadrupole mass spectrometer (Thermo Fisher Scientific, San Jose, CA).
The LC-SRM platform was configured and utilized as previously described (56). The abundance
of PknE and PknI in wild-type, PknE LOF mutant, and PknI LOF mutant were too low to be
detectable through the standard LC-SRM analysis. An extra PRISM procedure was applied to
these low abundance samples to enrich the targeted peptides for PknE and PknI before LC-
SRM analysis (57). SRM data acquired on the TSQ Altis were analyzed using Skyline-daily

software (version 21.0.9.139) (58). Peak detection and integration were determined based on retention time and the relative SRM peak intensity ratios across multiple transitions between light peptide and heavy peptide standards (59). All the data were manually inspected to ensure correct peak assignment and peak boundaries. The peak area ratios of endogenous light peptides and their heavy isotope-labeled internal standards (i.e., L/H peak area ratios) were then automatically calculated by Skyline-daily, and the average peak area ratios from all the transitions were used for quantitative analysis of the samples. For targets that had more than one surrogate peptide, correlation graphs were plotted to verify a strong correlation and ultimately the peptide that had the most sensitive response was selected for obtaining quantitative values.

RNA Isolation and RNA sequencing

Mtb cultures were pelleted at 4000 x g for 5 min at 4°C, resuspended in Trizol, transferred to a tube containing Lysing Matrix B, and lysed via bead beating using 30 sec shaking at 6 m/s in a homogenizer three times, with cooling on ice. The samples were centrifuged at 21,000 x g for 1 minute, and the supernatant was transferred to a tube containing 300 µl chloroform and Heavy Phase Lock Gel. The tubes were inverted for 2 minutes and centrifuged at 21,000 x g for 5 minutes. RNA in the aqueous phase was precipitated using 300 µl isopropanol and 300 µl high-salt solution (0.8 M Na citrate, 1.2 M NaCl). RNA was purified using a RNeasy kit with one on-column DNase treatment. Total RNA was quantified using a Nanodrop. Messenger RNA was enriched by first depleting ribosomal RNA from samples using the Ribo-Zero rRNA removal (bacteria) magnetic kit (Illumina). The purified mRNA was prepared for Illumina sequencing using the NEBNext Ultra RNA Library Prep Kit for Illumina according to the manufacturer's instructions in combination with AMPure XP reagent for clean-up of adaptor-ligated DNA. Each replicate was barcoded in the DNA library using the NEBNext Multiplex Oligos for Illumina (Dual Index Primers Set 1). 30-40 libraries were multiplexed per sequencing run, and libraries were quantified using the KAPA qPCR quantification kit. The libraries were sequenced at the University of Washington Northwest Genomics Center with the Illumina NextSeq 500 High Output v2 Kit. Read alignment was performed using the custom processing pipeline that uses Bowtie 2 utilities, available at https://github.com/robertdouglasmorrison/DuffyNGS and https://github.com/robertdouglasmorrison/DuffyTools.

Phosphorylation motif analysis

The custom background peptideome was generated by using the *Mtb* H37Rv proteome from UniProt.org (Proteome ID: UP000001584, modification date 12/1/2019). A custom R script was used to partition the proteome into unique, 11 amino acid long peptides. Sequences were further subdivided into lists whose central residue was either serine, threonine, or tyrosine. The foreground peptidome was generated by compiling the differentially modulated phosphosites for each kinase mutant. An R script was used to create a 11 amino acid sequence centered on the modulated phosphosite. Additionally, the same script was used to generate a list of all phosphosites identified in the WT phosphoproteome (basal phosphoproteome). For the motif analysis, four separate analyses were performed for each of the kinase mutants and the basal phosphoproteome: a site-centered or a Ser/Thr/Tyr-centered motif search. Foreground and respective background peptidomes were uploaded to the pLogo web application for analysis using default parameters.

***In vitro* phosphorylation assay**

The kinase domains of PknB-PknL were cloned from *Mtb* genomic DNA into inducible pET28b protein expression plasmid using Gibson assembly cloning and confirmed by DNA sequencing. pET28b plasmids were transformed into *E. coli* BL21 for protein expression. Cultures were grown to 0.4-0.6 OD, induced with 250 µM IPTG, and grown at 20°C for 18-24 hours. Cultures

were pelleted at 4000 x g for 20 minutes, resuspended with lysis buffer (20 mM Tris-HCl pH 7.5,
150 mM NaCl, 20 mM imidazole), and lysed by sonication. Lysates were centrifuged to remove
the insoluble fraction, and purified using affinity chromatography with Ni-NTA. Proteins were
further purified using size exclusion chromatography and stored in size exclusion buffer (20 mM
Tris-HCl, 150 mM NaCl, 5% glycerol, pH 7.4). For the auto-phosphorylation assay, 1 µM STPK
was combined with reaction buffer (50 µM $MnCl_2$, 50 µM $MgCl_2$, 1 mM DTT, 50 mM Tris pH 8.0,
75 mM NaCl) and initiated with 50 µM ATP and 15 µCi ATP [γ-32P]. The reaction was incubated
at 37°C for 30 min. To terminate the reaction, 4x loading buffer with β-mercaptoethanol was
added and incubated at 95°C for 5 minutes. Proteins were separated by SDS-PAGE in MES
buffer and 4-15% Bis-Tris Plus gels. Gels were dried using a gel dryer at 80°C for 3 hours.
Phosphor screens were exposed to the dried gels overnight and then imaged using a Sapphire
biomolecular imager. For Zur phosphorylation assay, 10 µM Rv2359 was combined with 1 µM
STPK in reaction buffer (50 µM $MnCl_2$, 50 µM $MgCl_2$, 1 mM DTT, 50 mM Tris pH 8.0, 75 mM
NaCl), with 50 µM ATP and 15 µCi ATP [γ -32P], and then incubated at 37°C for 30 min.
Proteins were separated by SDS-PAGE and visualized as above.
828
Zur mutagenesis
Zur phosphomimetic (Thr67Asp) and phosphoablative (Thr67Ala) mutants were generated by
QuikChange site-directed mutagenesis. In summary, overlapping primers containing the desired
mutation were used to linearize and introduce the mutation into the pET28b plasmid containing
Zur. PCR products were gel extracted and DpnI-digested, followed by transformation into
competent *E. coli*. Mutagenesis was confirmed by DNA sequencing.
835
Electrophoretic Mobility Shift Assay (EMSA)
Cy5-labeled DNA probe was made by resuspending three oligos (Integrated DNA Technologies)
to 100 µM in dsDNA annealing buffer (10 mM Tris-HCl-pH 7.5, 100 mM NaCl, 1 mM EDTA).
Oligo 1 corresponded to 100-200 bp binding region of Rv2359. Oligo 2 corresponded to the
reverse complement of Oligo 1, as well as a 12-nucleotide sequence at the 3' end to which
Oligo 3 was the reverse complement. Oligo 3 was a 12-mer with Cy5 covalently coupled to the
5' end. The three oligos were combined to a final concentration of 100 µM, vortexed, and
heated to 95°C for 10 min. The reaction was cooled to room temperature over 3-4 hours
protected from light and used for EMSA. For the EMSA, various concentrations of purified
recombinant Rv2359 WT or mutant protein were combined with 2 µM final DNA probe in
reaction buffer (20 mM Tris-HCl pH 8.0, 75 mM NaCl, 1 mM DTT, 50 µM $MgCl_2$, 50 µM $MnCl_2$,
5% glycerol, 1% zinc sulfate, 50 µg/mL bovine serum albumin, 50 µg/mL salmon sperm DNA)
and incubated for 30 min at room temperature, protected from light. After incubation, 4 µl native
sample loading buffer was added to each sample. The whole reaction was loaded on a native
10% Tris-Glycine gel, and gel was maintained on ice and ran at a constant 100 volts for 2 hours.
The gel was rinsed with water and visualized on a Li-cor Odyssey scanner. Bands were
quantified using the Odyssey software.
853
854

855 **ACKNOWLEDGEMENTS**
856
This work was supported by NIH grants R01AI117023, R21AI137571, and R03AI131223 and by
a grant by the American Lung Association to C.G. A.F. was supported by the Interdisciplinary
Program in Bacterial Pathogenesis 5T32AI053396. Portions of this research were supported by
NIH NIGSM GM103493. Some work was performed in the Environmental Molecular Sciences
Laboratory, a U.S. Department of Energy Office of Biological and Environmental Research
national scientific user facility located at Pacific Northwest National Laboratory in Richland,

REFERENCES

1.   Dworkin, J. (2015) Ser/Thr phosphorylation as a regulatory mechanism in bacteria. *Curr*
Opin Microbiol **24**, 47-52
2.   Stock, A. M., Robinson, V. L., and Goudreau, P. N. (2000) Two-component signal
transduction. *Annu Rev Biochem* **69**, 183-215
3.   Kannan, N., Taylor, S. S., Zhai, Y., Venter, J. C., and Manning, G. (2007) Structural and
functional diversity of the microbial kinome. *PLoS Biol* **5**, e17
4.   Krupa, A., and Srinivasan, N. (2005) Diversity in domain architectures of Ser/Thr kinases
and their homologues in prokaryotes. *BMC Genomics* **6**, 129
5.   Leonard, C. J., Aravind, L., and Koonin, E. V. (1998) Novel families of putative protein
kinases in bacteria and archaea: evolution of the "eukaryotic" protein kinase superfamily.
Genome Res **8**, 1038-1047
6.   Stancik, I. A., Sestak, M. S., Ji, B., Axelson-Fisk, M., Franjevic, D., Jers, C., Domazet-
Loso, T., and Mijakovic, I. (2018) Serine/Threonine Protein Kinases from Bacteria,
Archaea and Eukarya Share a Common Evolutionary Origin Deeply Rooted in the Tree
of Life. *J Mol Biol* **430**, 27-32
7.   Kennelly, P. J. (2014) Protein Ser/Thr/Tyr phosphorylation in the Archaea. *J Biol Chem*
289, 9480-9487
8.   Av-Gay, Y., and Everett, M. (2000) The eukaryotic-like Ser/Thr protein kinases of
Mycobacterium tuberculosis. *Trends Microbiol* **8**, 238-244
9.   Cole, S. T., Brosch, R., Parkhill, J., Garnier, T., Churcher, C., Harris, D., Gordon, S. V.,
Eiglmeier, K., Gas, S., Barry, C. E., 3rd, Tekaia, F., Badcock, K., Basham, D., Brown, D.,
Chillingworth, T., Connor, R., Davies, R., Devlin, K., Feltwell, T., Gentles, S., Hamlin, N.,
Holroyd, S., Hornsby, T., Jagels, K., Krogh, A., McLean, J., Moule, S., Murphy, L.,
Oliver, K., Osborne, J., Quail, M. A., Rajandream, M. A., Rogers, J., Rutter, S., Seeger,
K., Skelton, J., Squares, R., Squares, S., Sulston, J. E., Taylor, K., Whitehead, S., and
Barrell, B. G. (1998) Deciphering the biology of Mycobacterium tuberculosis from the
complete genome sequence. *Nature* **393**, 537-544
10.   Carette, X., Platig, J., Young, D. C., Helmel, M., Young, A. T., Wang, Z., Potluri, L. P.,
Moody, C. S., Zeng, J., Prisic, S., Paulson, J. N., Muntel, J., Madduri, A. V. R., Velarde,
J., Mayfield, J. A., Locher, C., Wang, T., Quackenbush, J., Rhee, K. Y., Moody, D. B.,
Steen, H., and Husson, R. N. (2018) Multisystem Analysis of Mycobacterium
tuberculosis Reveals Kinase-Dependent Remodeling of the Pathogen-Environment
Interface. *mBio* **9**
11.   Zeng, J., Platig, J., Cheng, T. Y., Ahmed, S., Skaf, Y., Potluri, L. P., Schwartz, D., Steen,
H., Moody, D. B., and Husson, R. N. (2020) Protein kinases PknA and PknB
independently and coordinately regulate essential Mycobacterium tuberculosis
physiologies and antimicrobial susceptibility. *PLoS Pathog* **16**, e1008452
12.   Molle, V., and Kremer, L. (2010) Division and cell envelope regulation by Ser/Thr
phosphorylation: Mycobacterium shows the way. *Mol Microbiol* **75**, 1064-1077
13.   Kusebauch, U., Ortega, C., Ollodart, A., Rogers, R. S., Sherman, D. R., Moritz, R. L.,
and Grundner, C. (2014) Mycobacterium tuberculosis supports protein tyrosine
phosphorylation. *Proc Natl Acad Sci U S A* **111**, 9265-9270
14.   Prisic, S., Dankwa, S., Schwartz, D., Chou, M. F., Locasale, J. W., Kang, C. M., Bemis,
G., Church, G. M., Steen, H., and Husson, R. N. (2010) Extensive phosphorylation with

overlapping specificity by Mycobacterium tuberculosis serine/threonine protein kinases.
Proc Natl Acad Sci U S A **107**, 7521-7526

15.   Alber, T. (2009) Signaling mechanisms of the Mycobacterium tuberculosis receptor
Ser/Thr protein kinases. *Curr Opin Struct Biol* **19**, 650-657

16.   Sherman, D. R., and Grundner, C. (2014) Agents of change - concepts in
Mycobacterium tuberculosis Ser/Thr/Tyr phosphosignalling. *Mol Microbiol* **94**, 231-241

17.   Breitkreutz, A., Choi, H., Sharom, J. R., Boucher, L., Neduva, V., Larsen, B., Lin, Z. Y.,
Breitkreutz, B. J., Stark, C., Liu, G., Ahn, J., Dewar-Darch, D., Reguly, T., Tang, X.,
Almeida, R., Qin, Z. S., Pawson, T., Gingras, A. C., Nesvizhskii, A. I., and Tyers, M.
(2010) A global protein kinase and phosphatase interaction network in yeast. *Science*
328, 1043-1046

18.   Schneiker, S., Perlova, O., Kaiser, O., Gerth, K., Alici, A., Altmeyer, M. O., Bartels, D.,
Bekel, T., Beyer, S., Bode, E., Bode, H. B., Bolten, C. J., Choudhuri, J. V., Doss, S.,
Elnakady, Y. A., Frank, B., Gaigalat, L., Goesmann, A., Groeger, C., Gross, F., Jelsbak,
L., Jelsbak, L., Kalinowski, J., Kegler, C., Knauber, T., Konietzny, S., Kopp, M., Krause,
L., Krug, D., Linke, B., Mahmud, T., Martinez-Arias, R., McHardy, A. C., Merai, M.,
Meyer, F., Mormann, S., Munoz-Dorado, J., Perez, J., Pradella, S., Rachid, S., Raddatz,
G., Rosenau, F., Ruckert, C., Sasse, F., Scharfe, M., Schuster, S. C., Suen, G., Treuner-
Lange, A., Velicer, G. J., Vorholter, F. J., Weissman, K. J., Welch, R. D., Wenzel, S. C.,
Whitworth, D. E., Wilhelm, S., Wittmann, C., Blocker, H., Puhler, A., and Muller, R.
(2007) Complete genome sequence of the myxobacterium Sorangium cellulosum. *Nat*
Biotechnol **25**, 1281-1289

19.   Baer, C. E., Iavarone, A. T., Alber, T., and Sassetti, C. M. (2014) Biochemical and
spatial coincidence in the provisional Ser/Thr protein kinase interaction network of
Mycobacterium tuberculosis. *J Biol Chem* **289**, 20422-20433

20.   Roskoski, R., Jr. (2021) Properties of FDA-approved small molecule protein kinase
inhibitors: A 2021 update. *Pharmacol Res* **165**, 105463

21.   Prisic, S., and Husson, R. N. (2014) Mycobacterium tuberculosis Serine/Threonine
Protein Kinases. *Microbiol Spectr* **2**

22.   van Kessel, J. C., and Hatfull, G. F. (2007) Recombineering in Mycobacterium
tuberculosis. *Nat Methods* **4**, 147-152

23.   Rock, J. M., Hopkins, F. F., Chavez, A., Diallo, M., Chase, M. R., Gerrick, E. R.,
Pritchard, J. R., Church, G. M., Rubin, E. J., Sassetti, C. M., Schnappinger, D., and
Fortune, S. M. (2017) Programmable transcriptional repression in mycobacteria using an
orthogonal CRISPR interference platform. *Nat Microbiol* **2**, 16274

24.   Soares, N. C., Spat, P., Krug, K., and Macek, B. (2013) Global dynamics of the
Escherichia coli proteome and phosphoproteome during growth in minimal medium. *J*
Proteome Res **12**, 2611-2621

25.   Kelkar, D. S., Kumar, D., Kumar, P., Balakrishnan, L., Muthusamy, B., Yadav, A. K.,
Shrivastava, P., Marimuthu, A., Anand, S., Sundaram, H., Kingsbury, R., Harsha, H. C.,
Nair, B., Prasad, T. S., Chauhan, D. S., Katoch, K., Katoch, V. M., Kumar, P.,
Chaerkady, R., Ramachandran, S., Dash, D., and Pandey, A. (2011) Proteogenomic
analysis of Mycobacterium tuberculosis by high resolution mass spectrometry. *Mol Cell*
Proteomics **10**, M111 011627

26.   Arnvig, K. B., Comas, I., Thomson, N. R., Houghton, J., Boshoff, H. I., Croucher, N. J.,
Rose, G., Perkins, T. T., Parkhill, J., Dougan, G., and Young, D. B. (2011) Sequence-
based analysis uncovers an abundance of non-coding RNA in the total transcriptome of
Mycobacterium tuberculosis. *PLoS Pathog* **7**, e1002342

27.   Hornbeck, P. V., Zhang, B., Murray, B., Kornhauser, J. M., Latham, V., and Skrzypek, E.
(2015) PhosphoSitePlus, 2014: mutations, PTMs and recalibrations. *Nucleic Acids Res*
43, D512-520

28.   Olsen, J. V., Vermeulen, M., Santamaria, A., Kumar, C., Miller, M. L., Jensen, L. J.,
Gnad, F., Cox, J., Jensen, T. S., Nigg, E. A., Brunak, S., and Mann, M. (2010)
Quantitative phosphoproteomics reveals widespread full phosphorylation site occupancy
during mitosis. *Sci Signal* **3**, ra3
29.   Sharma, K., D'Souza, R. C., Tyanova, S., Schaab, C., Wisniewski, J. R., Cox, J., and
Mann, M. (2014) Ultradeep human phosphoproteome reveals a distinct regulatory nature
of Tyr and Ser/Thr-based signaling. *Cell Rep* **8**, 1583-1594
30.   Griffin, J. E., Gawronski, J. D., Dejesus, M. A., Ioerger, T. R., Akerley, B. J., and
Sassetti, C. M. (2011) High-resolution phenotypic profiling defines genes essential for
mycobacterial growth and cholesterol catabolism. *PLoS Pathog* **7**, e1002251
31.   Kanshin, E., Giguere, S., Jing, C., Tyers, M., and Thibault, P. (2017) Machine Learning
of Global Phosphoproteomic Profiles Enables Discrimination of Direct versus Indirect
Kinase Substrates. *Mol Cell Proteomics* **16**, 786-798
32.   O'Shea, J. P., Chou, M. F., Quader, S. A., Ryan, J. K., Church, G. M., and Schwartz, D.
(2013) pLogo: a probabilistic approach to visualizing sequence motifs. *Nat Methods* **10**,
1211-1212
33.   Bradley, D., and Beltrao, P. (2019) Evolution of protein kinase substrate recognition at
the active site. *PLoS Biol* **17**, e3000341
34.   Frades, I., Resjo, S., and Andreasson, E. (2015) Comparison of phosphorylation
patterns across eukaryotes by discriminative N-gram analysis. *BMC Bioinformatics* **16**,
239
35.   Bach, H., Wong, D., and Av-Gay, Y. (2009) Mycobacterium tuberculosis PtkA is a novel
protein tyrosine kinase whose substrate is PtpA. *Biochem J* **420**, 155-160
36.   Bodenmiller, B., Wanka, S., Kraft, C., Urban, J., Campbell, D., Pedrioli, P. G., Gerrits, B.,
Picotti, P., Lam, H., Vitek, O., Brusniak, M. Y., Roschitzki, B., Zhang, C., Shokat, K. M.,
Schlapbach, R., Colman-Lerner, A., Nolan, G. P., Nesvizhskii, A. I., Peter, M., Loewith,
R., von Mering, C., and Aebersold, R. (2010) Phosphoproteomic analysis reveals
interconnected system-wide responses to perturbations of kinases and phosphatases in
yeast. *Sci Signal* **3**, rs4
37.   Rustad, T. R., Minch, K. J., Ma, S., Winkler, J. K., Hobbs, S., Hickey, M., Brabant, W.,
Turkarslan, S., Price, N. D., Baliga, N. S., and Sherman, D. R. (2014) Mapping and
manipulating the Mycobacterium tuberculosis transcriptome using a transcription factor
overexpression-derived regulatory network. *Genome Biol* **15**, 502
38.   Hunter, T., and Karin, M. (1992) The regulation of transcription by phosphorylation. *Cell*
70, 375-387
39.   Maciag, A., Dainese, E., Rodriguez, G. M., Milano, A., Provvedi, R., Pasca, M. R., Smith,
I., Palu, G., Riccardi, G., and Manganelli, R. (2007) Global analysis of the
Mycobacterium tuberculosis Zur (FurB) regulon. *J Bacteriol* **189**, 730-740
40. Baros, S. S., Blackburn, J. M., and Soares, N. C. (2020) Phosphoproteomic Approaches
to Discover Novel Substrates of Mycobacterial Ser/Thr Protein Kinases. *Mol Cell*
Proteomics **19**, 233-244
41. Boshoff, H. I., Myers, T. G., Copp, B. R., McNeil, M. R., Wilson, M. A., and Barry, C. E.,
3rd. (2004) The transcriptional responses of Mycobacterium tuberculosis to inhibitors of
metabolism: novel insights into drug mechanisms of action. *J Biol Chem* **279**, 40174-
40184
42. Honaker, R. W., Leistikow, R. L., Bartek, I. L., and Voskuil, M. I. (2009) Unique roles of
DosT and DosS in DosR regulon induction and Mycobacterium tuberculosis dormancy.
Infect Immun **77**, 3258-3263
43. O'Donnell, G., Poeschl, R., Zimhony, O., Gunaratnam, M., Moreira, J. B., Neidle, S.,
Evangelopoulos, D., Bhakta, S., Malkinson, J. P., Boshoff, H. I., Lenaerts, A., and

Gibbons, S. (2009) Bioactive pyridine-N-oxide disulfides from Allium stipitatum. *J Nat*
Prod **72**, 360-365
44. Raghavan, S., Manzanillo, P., Chan, K., Dovey, C., and Cox, J. S. (2008) Secreted
transcription factor controls Mycobacterium tuberculosis virulence. *Nature* **454**, 717-721
45. Rustad, T. R., Harrell, M. I., Liao, R., and Sherman, D. R. (2008) The enduring hypoxic
response of Mycobacterium tuberculosis. *PLoS One* **3**, e1502
46. van Kessel, J. C., and Hatfull, G. F. (2008) Mycobacterial recombineering. *Methods Mol*
Biol **435**, 203-215
47. Wang, Y., Yang, F., Gritsenko, M. A., Wang, Y., Clauss, T., Liu, T., Shen, Y., Monroe, M.
E., Lopez-Ferrer, D., Reno, T., Moore, R. J., Klemke, R. L., Camp, D. G., 2nd, and
Smith, R. D. (2011) Reversed-phase chromatography with multiple fraction
concatenation strategy for proteome profiling of human MCF10A cells. *Proteomics* **11**,
2019-2026
48. Mertins, P., Tang, L. C., Krug, K., Clark, D. J., Gritsenko, M. A., Chen, L., Clauser, K. R.,
Clauss, T. R., Shah, P., Gillette, M. A., Petyuk, V. A., Thomas, S. N., Mani, D. R., Mundt,
F., Moore, R. J., Hu, Y., Zhao, R., Schnaubelt, M., Keshishian, H., Monroe, M. E.,
Zhang, Z., Udeshi, N. D., Mani, D., Davies, S. R., Townsend, R. R., Chan, D. W., Smith,
R. D., Zhang, H., Liu, T., and Carr, S. A. (2018) Reproducible workflow for multiplexed
deep-scale proteome and phosphoproteome analysis of tumor tissues by liquid
chromatography-mass spectrometry. *Nat Protoc* **13**, 1632-1661
49. Zhang, H., Liu, T., Zhang, Z., Payne, S. H., Zhang, B., McDermott, J. E., Zhou, J. Y.,
Petyuk, V. A., Chen, L., Ray, D., Sun, S., Yang, F., Chen, L., Wang, J., Shah, P., Cha,
S. W., Aiyetan, P., Woo, S., Tian, Y., Gritsenko, M. A., Clauss, T. R., Choi, C., Monroe,
M. E., Thomas, S., Nie, S., Wu, C., Moore, R. J., Yu, K. H., Tabb, D. L., Fenyo, D.,
Bafna, V., Wang, Y., Rodriguez, H., Boja, E. S., Hiltke, T., Rivers, R. C., Sokoll, L., Zhu,
H., Shih, I. M., Cope, L., Pandey, A., Zhang, B., Snyder, M. P., Levine, D. A., Smith, R.
D., Chan, D. W., Rodland, K. D., and Investigators, C. (2016) Integrated Proteogenomic
Characterization of Human High-Grade Serous Ovarian Cancer. *Cell* **166**, 755-765
50. Elias, J. E., and Gygi, S. P. (2007) Target-decoy search strategy for increased
confidence in large-scale protein identifications by mass spectrometry. *Nat Methods* **4**,
207-214
51. Qian, W. J., Liu, T., Monroe, M. E., Strittmatter, E. F., Jacobs, J. M., Kangas, L. J.,
Petritis, K., Camp, D. G., 2nd, and Smith, R. D. (2005) Probability-based evaluation of
peptide and protein identifications from tandem mass spectrometry and SEQUEST
analysis: the human proteome. *J Proteome Res* **4**, 53-62
52. Kim, S., Gupta, N., and Pevzner, P. A. (2008) Spectral probabilities and generating
functions of tandem mass spectra: a strike against decoy databases. *J Proteome Res* **7**,
3354-3363
53. Cox, J., and Mann, M. (2008) MaxQuant enables high peptide identification rates,
individualized p.p.b.-range mass accuracies and proteome-wide protein quantification.
Nat Biotechnol **26**, 1367-1372
54. Clair, G., Piehowski, P. D., Nicola, T., Kitzmiller, J. A., Huang, E. L., Zink, E. M., Sontag,
R. L., Orton, D. J., Moore, R. J., Carson, J. P., Smith, R. D., Whitsett, J. A., Corley, R.
A., Ambalavanan, N., and Ansong, C. (2016) Spatially-Resolved Proteomics: Rapid
Quantitative Analysis of Laser Capture Microdissected Alveolar Tissue Samples. *Sci*
Rep **6**, 39223
55. Shi, T., Sun, X., Gao, Y., Fillmore, T. L., Schepmoes, A. A., Zhao, R., He, J., Moore, R.
J., Kagan, J., Rodland, K. D., Liu, T., Liu, A. Y., Smith, R. D., Tang, K., Camp, D. G.,
2nd, and Qian, W. J. (2013) Targeted quantification of low ng/mL level proteins in human
serum without immunoaffinity depletion. *J Proteome Res* **12**, 3353-3361

56.   Nielson, C. M., Jones, K. S., Chun, R. F., Jacobs, J. M., Wang, Y., Hewison, M., Adams,
1064   J. S., Swanson, C. M., Lee, C. G., Vanderschueren, D., Pauwels, S., Prentice, A., Smith,
1065   R. D., Shi, T., Gao, Y., Schepmoes, A. A., Zmuda, J. M., Lapidus, J., Cauley, J. A.,
1066   Bouillon, R., Schoenmakers, I., Orwoll, E. S., and Osteoporotic Fractures in Men
1067   Research, G. (2016) Free 25-Hydroxyvitamin D: Impact of Vitamin D Binding Protein
1068   Assays on Racial-Genotypic Associations. *J Clin Endocrinol Metab* **101**, 2226-2234
1069   57.   Shi, T., Fillmore, T. L., Sun, X., Zhao, R., Schepmoes, A. A., Hossain, M., Xie, F., Wu,
1070   S., Kim, J. S., Jones, N., Moore, R. J., Pasa-Tolic, L., Kagan, J., Rodland, K. D., Liu, T.,
1071   Tang, K., Camp, D. G., 2nd, Smith, R. D., and Qian, W. J. (2012) Antibody-free, targeted
1072   mass-spectrometric approach for quantification of proteins at low picogram per milliliter
1073   levels in human plasma/serum. *Proc Natl Acad Sci U S A* **109**, 15395-15400
1074   58.   MacLean, B., Tomazela, D. M., Shulman, N., Chambers, M., Finney, G. L., Frewen, B.,
1075   Kern, R., Tabb, D. L., Liebler, D. C., and MacCoss, M. J. (2010) Skyline: an open source
1076   document editor for creating and analyzing targeted proteomics experiments.
1077   *Bioinformatics* **26**, 966-968
1078   59.   He, J., Sun, X., Shi, T., Schepmoes, A. A., Fillmore, T. L., Petyuk, V. A., Xie, F., Zhao,
1079   R., Gritsenko, M. A., Yang, F., Kitabayashi, N., Chae, S. S., Rubin, M. A., Siddiqui, J.,
1080   Wei, J. T., Chinnaiyan, A. M., Qian, W. J., Smith, R. D., Kagan, J., Srivastava, S.,
1081   Rodland, K. D., Liu, T., and Camp, D. G., 2nd. (2014) Antibody-independent targeted
1082   quantification of TMPRSS2-ERG fusion protein products in prostate cancer. *Mol Oncol*
1083   **8**, 1169-1180

1084

A

STPK	Number of TF substrates
PknB	9
PknD	21
PknE	39
PknF	48
PknG	3
PknH	20
PknI	2
PknJ	4
PknK	94
PknL	76

B

Differentially expressed genes (LoF / GoF):
- PknB: 489 / 16
- PknD: 8 / 85
- PknE: 120 / 321
- PknF: 0 / 312
- PknG: 44 / 1
- PknH: 2 / 46
- PknI: 0 / 8
- PknJ: 0 / 103
- PknK: 103 / 181
- PknL: 1 / 260

C

Most induced genes

STPK	Mutant	Gene	Fold-change	p-value
PknH	GoF	mhpE	333	3.84e-43
PknH	GoF	Rv1271c	259	2.43e-55
PknK	GoF	echA13	227	1.28e-85
PknH	GoF	Rv1291c	168	1.89e-33
PknK	GoF	whiB5	164	4.56e-32
PknK	GoF	fadE17	163	2.29e-52
PknK	GoF	Rv1936	109	3.07e-142
PknH	GoF	Rv1290A	68	4.68e-20
PknK	GoF	Rv0077c	61	2.69e-51
PknB	LoF	papA1	59	6.30e-26

Most repressed genes

STPK	Mutant	Gene	Fold-change	p-value
PknL	GoF	nirB	-89	0.00e+00
PknL	GoF	Rv3371	-72	0.00e+00
PknB	LoF	Rv2625c	-58	1.58e-25
PknB	LoF	Rv2628	-55	5.52e-23
PknB	LoF	hrp1	-55	1.90e-13
PknB	LoF	TB31.7	-45	1.44e-17
PknB	LoF	Rv2624c	-38	1.68e-20
PknB	LoF	Rv3129	-38	4.08e-11
PknL	GoF	Rv3131	-33	8.74e-24
PknL	GoF	glbN	-32	0.00e+00

PknG [38]

PknH [204]

PknF [720]

PknI [15]

Repressed (Single) 140

Induced (Single) 536

Induced and Repressed 61

PknE [506]

PknJ [50]

Repressed (Multiple) 130

Induced (Multiple) 288

PknD [282]

PknB [114]

PknL [968]

PknK [869]

Proportion of substrates identified in:

LoF

GoF

Reciprocal

A

B

C

A

	H37Rv+ WT STPK	H37Rv+ Kinase-Dead STPK
Total DEG:	260	1

B

Total DEG:	181	13

STPK-DEG Interaction Network