Modularity in temporal co-occurrence networks at a stopover site reveals strong

co-migration fidelity in African-European migratory landbirds over time

Bruno Bellisario[1], Massimiliano Cardinale[2], Ivan Maggini[3], Leonida Fusani[3,4*], Claudio Carere[5]

[1]Department of Agricultural and Forestry Sciences (DAFNE), University of Tuscia, Viterbo, Italy

[2]Swedish University of Agricultural Sciences, Department of Aquatic Resources, Institute of Marine

Research, Lysekil, Sweden

[3]Ornithology, Konrad-Lorenz Institute of Ethology, University of Veterinary Medicine, Vienna, Austria

[4]Department of Behavioural and Cognitive Biology, University of Vienna, Austria

[5]Department of Ecological and Biological Sciences (DEB), University of Tuscia, Viterbo, Italy

[*]Corresponding author

Author ORCID:

BB 0000-0002-5339-0757
MC 0000-0003-1870-3882
IM 0000-0002-1528-2288
LF 0000-0001-8900-796X
CC 0000-0003-1644-2113
20
21

Abstract

Migratory species are changing the timing of departures from wintering areas and

arrival to breeding sites (i.e., migration phenology), as response to climate change to

exploit maximum food availability at higher latitudes and improve their fitness. Despite

the impact of changing migration phenology on migratory connectivity at population

and community level, the extent to which such individual and species-specific response

affects associations among co-migrating species has been seldom explored. By applying

temporal co-occurrence network models on 15 years of standardized bird ringing data at

a spring stopover site, we show that African-European migratory landbirds tend to

migrate in well-defined groups of species with high temporal overlap. Such 'co-

migration fidelity' is significantly increasing over the years and higher in trans-Saharan

than North-African migrants. Our findings suggest non-random patterns of associations

in co-migrating species, possibly related to the existence of regulatory mechanisms

associated with changing climate conditions and different uses of stopover sites by

migrants, ultimately influencing the global migration economy of landbirds in the

Palearctic-African migration system.

Keywords: stopover; co-occurrence networks; migration phenology; co-migrants

1. Introduction

Migration represents a key life-history event for a large number of species and is typically associated with the temporal and geographic variation in resource availability [1]. Alongside endogenous (e.g., circannual rhythms) and exogenous (e.g., photoperiodic cues) mechanisms [2,3], favourable environmental conditions and food availability *en route* are key elements driving the timing of migration in birds [4–6].

Climate change may pose serious problems to migrants spending different parts of their annual cycle in different parts of the world as it drives the restructuring of important phenological events [7–9]. For instance, arrival dates to breeding grounds of Central-European populations of Pied flycatchers (*Ficedula hypoleuca*) showed no significant synchronization with advances in food-peak dates due to warming temperature, leading to a drastic decline (ca. 90%) in the abundance of populations that arrived after the peak [10]. Overall, differences in the extent of climatic change along migratory routes, as well as at wintering, stopover and breeding areas [11–13], increase the chance of phenological mismatches, influencing the interactions among co-occurring species including their food sources, predators, and competitors.

Within this context, migratory connectivity, defined as the 'geographic linking of individuals and populations between one life cycle stage and another' [14] represents a key factor having far-reaching consequences at multiple ecological scales, from individual fitness to community dynamics and ecosystem functioning. To date, most studies rely on a purely spatial perspective aimed at determining the extent to which different areas of a species' annual range are linked by the movement paths of individuals [14,15], providing useful insights on how climate change influence the ability of such areas in sustaining migratory flows (e.g., 'migration hot spots' [16]). However, an explicit consideration of the temporal dimension [17] is fundamental to

understand how, when and for how long species change their migratory patterns and

what are the consequences at multiple ecological levels.

Advances in the date of departure from wintering areas in response to endogenous

and/or exogenous factors reflecting changing climate conditions have been observed in

several species [18–20]. Populations of origin, sex, age, ecological requirements might

all affect the ability of species to change migration phenology [21–26], leading in some

cases to asynchronous migration patterns. Moreover, phenotypic plasticity and

microevolutionary changes can influence the long-term responses of species to track

changes along migratory routes, leading in this case to differences in the consistency of

migration pattern (i.e., how repeatable migration phenology and synchrony are over

time [17]).

Migrations involve the simultaneous movement of multiple species at once,

connecting separated and diverse communities that may have a strong impact on the

dynamic and stability of ecosystems. Migratory routes of different species often

converge both in space and time [27], resulting in direct and indirect ecological

interactions such as competition for resources, predation and even social interactions.

For instance, studies have shown how parasites transmission, as well as transient

dynamics in lake ecosystems, can be related to migration timing [28,29]. Similarly,

intra- and interspecific competition may be affected by synchronous migrations by

excluding inferior competitors from high-quality foraging areas [30,31], or promoting

coexistence when the joint use of a resource increases its quality and productivity [32].

Migration timing can rely on social interactions, as observed in songbirds migrating

along the coasts of Baltic Sea, which use vocalizations from conspecifics as cues to

assess the suitability of potential stopover habitats [33,34]. However, despite the

ecological importance of interactions between co-migrants, we still lack a proper

understanding on how long-term phenological changes affect the restructuring of avian

assemblages *en route*.

Investigating the causes and the effects of changing migration phenology from a

community point of view entails inherent difficulties associated with the need to

constantly track and monitor animals across oceans and continents [27]. A further

source of uncertainty derives from the need to appropriately model multi-species

temporal movements to identify recurrent patterns in the aggregation of species and

forecast the consequences of climate change on migrations. Within this context, co-

occurrence analysis and co-occurrence networks are valuable tools to elucidate

mechanisms driving species assemblages. Recently, García-Navas et al. (2021) [35]

used null model approaches on 20 years of abundance data to investigate co-occurrence

patterns in bird communities inhabiting habitats with different degree of productivity,

showing the role of social information and microhabitat preferences over interspecific

competition. Furthermore, Montaño-Centellas (2020) [36] used spatial data to build and

explore co-occurrence networks of avian mixed-species flocks across elevation

gradients, identifying changes in interactions as the main driver of species turnover

among Andean flocking birds. Co-occurrence networks can thus help illustrating

complex properties of animal movements with simple graphical metrics [37], although

the actual ecological meaning of revealed links must be critically evaluated (see [38]

and references therein).

By using temporal co-occurrence network models on time series data spanning 15

years of standardized bird ringing activity at the Mediterranean spring stopover area of

Ponza (Italy), we searched for a consistent and repeated order of association over time

to study how long-term changes in the migration timing of African-European migratory

landbirds affect the structuring of associations among co-migrants.

2. Materials and methods

2.1 Sampling site and bird ringing activity

The study was conducted on Ponza (figure 1), a small island in the Tyrrhenian Sea with an extension of ca. 9.87 km^2, located about 50 km off the Western coast of Italy (40°55' N, 12°58' E), where we have been monitoring spring migration through capture and ringing since 2002 (http://www.inanellamentoponza.it). Ponza is located along one of the main Mediterranean migratory routes and attracts a large number of African-European migratory landbirds during spring migration [20]. The bird capture season started in March and ended in May almost every year (exact start and end dates can be found in the supplementary Table S1 in [20]). Mist nets were opened every day except for days with heavy rain or strong winds (>15 knots) and the number of days with reduced effort was less than 1% of the total. Mist-nets were checked hourly from dawn until one hour after dusk, and birds were captured throughout the day with an average of 227 m of mist-nets over the time-period. We kept the net brand and model (Lavorazione Reti Bonardi, Monte Isola, BS, Italy, http://www.vbonardi.it/; 2.4 m height, 16 mm mesh size), as well as the vegetation height constant throughout the entire study period. After being captured, birds were ringed and measured using standard EURING procedures [39]. Here we used the daily abundances of the 31 most abundant species passing through Ponza (table 1), using ringing data over a period of 15 years (2007 to 2021), during which the procedures were fully standardized.

2.2 Network inference

For each year of survey ($n = 15$), we reconstructed the temporal co-occurrence network

by calculating the time-lagged Spearman rank correlations using the daily abundances

of each species with each other in the previous time step (i.e., the lag-1 correlation

[40]). To avoid spurious correlations due to extremely low abundances, we removed

from computation those species for which the total number of individuals in each year

was below the 25th percentile of the entire distribution. We used the packages 'Hmisc'

[41] and 'qvalue' [42] of R [43] to adjust the significance of correlations using the

Bonferroni-Holm procedure. Co-occurrences were considered significant for correlation

coefficients $r > 0.5$ and $p < 0.05$. We used unweighted links, where potential co-

occurrences were considered irrespective of the strength of the correlation.

2.3 Network modularity

We used an eigenvector-based maximizing algorithm (i.e., the Louvain algorithm [44])

to measure modularity (Q), that is, the degree to which a network subdivides in densely

connected groups of nodes (aka modules) with only sparser connections between groups

[45]. From an ecological point of view, modularity is a straightforward structural

property able to detect groups of species having similar responses to the main processes

structuring biodiversity in space and time [46–48]. In our case, the higher the

modularity, the greater will be the tendency of networks to cluster into subgroups of

species characterized by common migration timing (i.e., high temporal co-occurrence).

We considered networks with $Q > 0.3$ as having a strong modular subdivision [45].

The yearly measured modularity values were filtered by means of a simple moving

average (SMA), to reduce background noise in the original time series and emphasize

trends. We used SMA with an automated selection procedure based on the Akaike

information criterion to find the optimal order of the moving average [49]. Modularity

was measured by using the function 'cluster_louvain' [44] in the package 'igraph' [50]

of R, while we used the functions 'sma' and 'sen.slope' in the packages 'smooth' [49]

and 'trend' [51], respectively, to smooth the time series and measure the Mann-Kendall

statistics.

2.4 Co-migration fidelity

We defined 'co-migration fidelity' as the frequency with which two species tended to

co-occur in the same module over the years, measured by using the normalized mutual

information (*nmi* [52]). Basically, this metric assesses whether two classifications (i.e.,

module affiliation in different years) on the same set (i.e., species) well explain one

another, providing a robust metric for comparing networks as it is not affected by the

number and size of modules found in each network. The normalized mutual information

is bounded between 0 and 1, so that if two species were always part of the same module

over the years (i.e., perfect congruence, high fidelity), then *nmi* = 1; conversely, if two

species never co-occurred within the same module over the years (i.e., independent, no

fidelity), then *nmi* = 0. We thus measured all *nmi* values between species, resulting in a

symmetric similarity matrix expressing the pairwise mutual information among species.

We finally used the permutational multivariate analysis of variance using distance

matrices for partitioning the measured normalized mutual information among sources of

variation given by wintering areas and foraging niches. We therefore divided the species

in three main groups (table 1) corresponding to those wintering either north of the

Sahara Desert (North African), in the sub-Saharan belt extended between the Sahara

Desert and the Sudan savannah (Sahel), or in the Afrotropical ecozone (Tropical).

Following Pigot et al. (2020) [53], species were also characterized by considering their

main foraging niches (table 1). The normalized mutual information and permutational

multivariate analysis of variance were measured by using the function 'compare' and

'adonis2' in the 'igraph' and 'vegan' [50,54] packages of R, respectively.

3. Results

A total of 226,533 individuals were ringed over the 15 years of field activity, with an

average ($\pm$ SD) of 15,117 $\pm$ 4,031 per year and a minimum of 8,861 in 2010 and a

maximum of 23,186 in 2015. Three main families accounted for 84% of the total,

Sylviidae (four species, 40%), Muscicapidae (seven species, 29%) and Phylloscopidae

(three species, 15 %). The garden warbler (*Sylvia borin*) and common whitethroat

(*Sylvia communis*) were the most abundant species, while other as the common

chaffinch (*Fringilla coelebs*) and the great reed warbler (*Acrocephalus arundinaceus*)

were less abundant (table 1).

Overall, temporal co-occurrence networks always showed relatively strong modular

structures, with an average Q ($\pm$ SD) of 0.357 $\pm$ 0.053 and ranging from a minimum of

0.288 in 2010 and a maximum of 0.448 in 2018, also showing a significant increase

over the years (Mann-Kendall trend test $z = 3.865$, $p < 0.001$) (figure 2). The pairwise

co-migration fidelity, measured by the normalized mutual information (*nmi*), showed a

well-defined pattern, with species subdividing in (at least) six to seven main clusters

(figure 3a). Interestingly, one main cluster was characterized by species with very high

levels of co-migration fidelity (*S. borin*, *Hippolais icterina*, *Muscicapa striata*, *Oriolus*

oriolus), and always co-occurring in the same module with the same species over the

years, with no temporal overlap with other species (figure 3a). Other, as the black

redstart (*Phoenicurus ochruros*), the song thrush (*Turdus philomelos*), the common

stonechat (*Saxicola torquata*), the European robin (*Erithacus rubecula*) and the

common chiffchaff (*Phylloscopus collybita*) showed more variable *nmi* index, with

partial temporal overlap with other species (figure 3a). Co-migration fidelity

significantly differed between wintering areas ($F = 2.635$, $p = 0.004$), with birds

wintering in North Africa showing on average lower values of *nmi* compared to those

wintering in the sub-Saharan belt and Afrotropical ecozone (figure 3b). We did not find

significant differences with respect to foraging niches ($F = 1.379$, $p = 0.088$, figure 3b).

4. Discussion

By applying network models to a long-term series of captures of spring migratory

passerines at a stopover site, we showed that changes in the timing of migration of

African-European landbirds lead to non-random patterns in the assemblage of co-

occurring migrants. We found that modularity is a common pattern in temporal co-

occurrence networks, with a trend for an increase over the years. This indicates a

general tendency of birds to arrive at the stopover area in well-defined groups of species

characterized by common migration phenology, a tendency likely to be influenced by

changes in climatic conditions and to which species respond by restructuring their

migration patterns in a non-random way.

Spring migration phenology of birds crossing the Mediterranean Sea can be related to

specific ecological factors associated with travel distance and competition for nesting

sites, as well as to the degree of sexual dimorphism [55]. For instance, species wintering

closer to the breeding areas (e.g., North Africa) usually have faster adaptive responses

due to their ability to better track changing conditions at the breeding grounds [56].

Latitudinal differences in wintering sites might thus determine the phenology of spring

migration because species wintering north of Sahara advance their peak date of passage

of more days compared to sub-Saharan ones [20]. Moreover, recent findings have

shown a substantial increase (up to one day per year) of the migration window for

species wintering south of the Sahara Desert, possibly related to more favourable

conditions at wintering areas that allow faster refuelling, earlier departure, and/or longer

residence time [20]. The strong modular subdivision observed over the study period

(with the only exception of 2010) suggests that passerine species moving within the

African-West Eurasian flyway during spring migration always followed a well-defined

order of migration, a tendency likely to be predominant in those species forced to cover

larger distance to reach their breeding areas.

How migratory birds manage to detect and adapt to changing environmental

conditions during their journey to breeding areas remains an open question, possibly

related to whether shifts in migration phenology are driven by microevolutionary

changes or phenotypic plasticity [57]. Phenotypic plasticity may provide faster adaptive

responses, especially for species travelling shorter distance and capable of immediate

response to changing environmental conditions [58]. Although our findings do not

allow for direct evidence, they nevertheless indicate a close relationship between bird's

phenotypic responses to migration and co-migration fidelity. This would explain why,

for instance, North-African migrants showed lower co-migration fidelity than trans-

Saharan ones, a pattern likely to be related to more flexible responses to changing

climate conditions and possibly influencing the repeatability and synchrony of

migration phenology over time.

4.1 Consequences for coexistence

Although co-occurrence patterns cannot be used to infer actual interspecific interactions

(see [38] and references therein), they can reveal potential for interferences among

species, especially when these are forced to share small areas in limited time frames,

such as at migratory bottlenecks [59]. This is the case of stopover areas, where birds

stop to rest, refuel and seek shelter during their migration activity, and where the

decision on where, when and how long to rest is driven by physiological adaptations

associated with migratory behaviour and habitat quality [60–62]. For instance, the

amount of forest cover and productivity relates to the two-step mechanism determining

the spatiotemporal use of stopover areas, *via* identification of high-quality habitat prior

to landing and sustained refuelling rates during stopover [61,63]. Overall, stopover

areas are important to replenish energetic reserves and can influence subsequent flight

ranges [64,65]. However, stopover sites can have functions different from providing

food, and their ecological context (e.g., proximity to ecological barriers, spatial

isolation) and intrinsic characteristics (e.g., diversity and abundance of resources) may

determine the use made by migrants [20,62,66,67].

Most individuals do not use Ponza for refuelling [67], as they might just need to rest

before continuing the last part of their journey [68,69]. Therefore, coexistence at

stopover areas might not be regulated exclusively by avoidance of competition for

resources, but also by the need to reduce the risk of predation or to gain information on

novel habitats and resources [33,34,70]. Almost all species investigated in our study

have a strictly carnivorous diet and feed mainly on insects, with the exception of the

European turtle dove, *Streptopelia turtur* and the Eurasian golden oriole *Oriolus*

oriolus, which are classified mainly as granivores' ground feeding species, but have

nevertheless variable foraging niches [53]. Several studies have shown that species with

similar diets switch to alternative foraging substrates and different predatory behaviour

to avoid competition [53,71]. Our findings show no association between co-migration

fidelity and foraging strategies, suggesting the existence of species-independent

foraging mechanisms promoting coexistence at stopover areas.

However, if refuelling is not the main reason for migrants to land in an area [20],

social (facilitative) interactions could explain the observed differences in species co-

migration fidelity. Social interactions have been observed in migrating songbirds that

use interspecific calls as cues to estimate stopover quality [33]. More recently, Gayk et

al. (2021) [72] showed how new world warblers that migrate in mixed-species flocks

produce flight calls to reduce disorientation during migration or to increase the chance

of finding high quality stopover sites, a pattern even more pronounced in

phylogenetically close species that breed at the same latitude with overlapping

migration timing. Thus, it could be hypothesized that the observed tendency to migrate

in well-defined and consistent subgroups of species with synchronized migration times

is functional to optimize the costs associated with predation risks *en route* and resting at

stopover area, in order to improve their competitive ability for territories at breeding

areas.

5. Conclusions

As animal migrations involve the simultaneous movements of individuals and species in

space and time, the emergence of potential interactions is far from being rare and is

likely to influence the costs and benefits of migration. Long-term phenological changes

have the potential to determine when, how long and with whom each species co-occurs

with other members of the community. Furthermore, the consistency and

synchronization of such changes might influence the global economy of migration. Here

we show that modularity is a common pattern in the multi-species assemblage of

African-European migratory landbirds, meaning that during spring migration birds

follow a well-defined schedule driven by common responses to the effects of climate

change on the timing of migration. Such pattern is likely to increase over time, mainly

affecting species with low phenotypic plasticity and which cover longer distance during

migration, with consequent fragmentation of migration timing. Further studies are

however needed to better understand the consistency of observed patterns and the

consequence for coexistence, for instance by extending the spatiotemporal coverage of

analysis to species' co-occurrence at wintering, stopover and breeding areas, or by

experimental tests of the effects of competitive interactions in the field. Our study

represents a first attempt in modelling multi-species temporal data to gain new insights

in the consequence of long-term changes in migration phenology on the restructuring of

avian assemblages *en route*.

Acknowledgements

We would love to dedicate this work to the loving memory of our friend and colleague
Prof. Dario Angeletti, torn from life too early. We would also like to thank Prof. Hanna
Kokko for the stimulating discussion that greatly improved the drafting of the
manuscript. The authors would also like to thank all the hundreds of volunteers at the
bird ringing station of Ponza (http://www.inanellamentoponza.it), who have made it
possible to constantly collect data over the years. This is publication N. XX of the
Piccole Isole project.

Conflict of Interest

The authors have no conflict of interest to declare.

Authors' contributions

BB conceived the paper together with CC; BB analysed the data and led the writing of
the manuscript; MC prepared and analysed the data. IM and LF contributed to data
collection and to writing of the manuscript. All authors have no perceived conflict of
interest and contributed critically to the writing of the manuscript and gave final
approval for publication.

Funding statement

Data used for this study were collected thanks to funds of the University of Ferrara,
University of Veterinary Medicine, Vienna, Grant Nr. 196451/V40 of the Research
Council of Norway, Grant Nr. P31037-B29 of the Austrian Science Fund (FWF), and a
number of donations and contributions to Centro Italiano per lo Studio e la
Conservazione dell'Ambiente (CISCA).

Data Availability Statement

Data used in this paper are available at Figshare:

Bird ringing data: https://doi.org/10.6084/m9.figshare.20438997.v1

Selected bird traits: https://doi.org/10.6084/m9.figshare.20439027.v1

R code: https://doi.org/10.6084/m9.figshare.20438982.v1.

Aggregated data-only can also be found at https://phaidra.vetmeduni.ac.at/view/o:114.

References

1.	Gauthreaux Jr. S.A. 1982 The ecology and evolution of avian migration systems. In *Avian Biology* (eds. Farner D.S., King J.R., Parkes K.C.), pp. 93-168. Amsterdam, Academic Press. (doi:https://doi.org/10.1016/B978-0-12-249406-2.50011-3)

2.	Gwinner E. 1996 Circadian and circannual programmes in avian migration. *J. Exp Biol* **199**, 39–48. (doi:10.1242/jeb.199.1.39)

3.	Maggini I., Bairlein F. 2010 Endogenous rhythms of seasonal migratory body mass changes and nocturnal restlessness in different populations of Northern Wheatear *Oenanthe oenanthe*. *J. Biol Rhythms* **25**(4), 268–276. (doi:10.1177/0748730410373442)

4.	Hasselquist D., Montràs-Janer T., Tarka M., Hansson B. 2017 Individual consistency of long-distance migration in a songbird: significant repeatability of autumn route, stopovers and wintering sites but not in timing of migration. *J. Avian Biol* **48**(1), 91–102. (doi:10.1111/jav.01292)

5.	Ruthrauff D.R., Tibbitts T.L., Gill R.E. 2019 Flexible timing of annual movements across consistently used sites by Marbled Godwits breeding in Alaska. *The Auk* **136**, 1–11. (doi:10.1093/auk/uky007)

6.	Thorup K., Tøttrup A.P., Willemoes M., Klaassen R.H.G., Strandberg R., Vega M.L., Dasari H.P., Araújo M.B., Wikelski M., Rahbek C. 2017 Resource tracking within and across continents in long-distance bird migrants. *Sci Adv* **3**(1), e1601360. (doi:10.1126/sciadv.1601360)

7.	Cresswell W., McCleery R.H. 2003 How great tits maintain synchronization of their hatch date with food supply in response to long-term variability in temperature. *J Anim Ecol* **72**(2), 356-366. (doi:10.1046/j.1365-2656.2003.00701.x)

8.	Visser M.E., Both C., Lambrechts M.M. 2004 Global Climate Change Leads to Mistimed Avian Reproduction. *Adv Ecol Res* **35**, 89-110. (doi:10.1016/s0065-2504(04)35005-1)

9.	Walther G.-R., Post E., Convey P., Menzel A., Parmesan C., Beebee T.J.C., Fromentin J.-M., Hoegh-Guldberg O., Bairlein F. 2002 Ecological responses to recent climate change. *Nature* **416**(6879), 389-395. (doi:10.1038/416389a)

10.	Both C., Bouwhuis S., Lessells C.M., Visser M.E. 2006 Climate change and population declines in a long-distance migratory bird. *Nature* **441**(7089), 81-83. (doi:10.1038/nature04539)

11.	Tryjanowski P., Kuźniak S., Sparks T. 2002 Earlier arrival of some farmland migrants in western Poland. *Ibis* **144**(1), 62-68. (doi:10.1046/j.0019-1019.2001.00022.x)

12.	Ahola M., Laaksonen T., Sippola K., Eeva T., Rainio K., Lehikoinen E. 2004 Variation in climate warming along the migration route uncouples arrival and breeding dates. *Glob Change Biol* **10**(9), 1610-1617. (doi:10.1111/j.1365-2486.2004.00823.x)

13.	Gordo O., Brotons L., Ferrer X., Comas P. 2005 Do changes in climate patterns in wintering areas affect the timing of the spring arrival of trans-Saharan migrant birds? *Glob Change Biol* **11**(1), 12-21. (doi:10.1111/j.1365-2486.2004.00875.x)

14.	Webster M.S., Marra P.P., Haig S.M., Bensch S., Holmes R.T. 2002 Links between worlds: unraveling migratory connectivity. *Trends Ecol Evol* **17**(2), 76-83. (doi:10.1016/S0169-5347(01)02380-1)

15.	Finch T., Butler S.J., Franco A.M.A., Cresswell W. 2017 Low migratory connectivity is common in long-distance migrant birds. *J Anim Ecol* **86**(3), 662-673. (doi:10.1111/1365-2656.12635)

16.	Bellisario B. 2018 Conserving migration in a changing climate, a case study: The Eurasian spoonbill, *Platalea leucorodia leucorodia*. *Biol Conserv* **217**, 222-231.

417 (doi:10.1016/j.biocon.2017.11.013)

417 17. Bauer S., Lisovski S., Hahn S. 2016 Timing is crucial for consequences of migratory connectivity. *Oikos* **125**(5), 605-612. (doi:10.1111/oik.02706)

418 18. Bairlein F., Winkel W. 2001 Birds and climate change. In *Climate of the 21st Century: Changes and Risks* (eds. Lozan J.L., Graßl H., Hupfer P.), pp. 278-282. Hamburg.

423 19. Lehikoinen E.S.A., Sparks T.H., Zalakevicius M. 2004 Arrival and Departure Dates. *Adv Ecol Res* **35**, 1-31. (doi:10.1016/s0065-2504(04)35001-4)

425 20. Maggini I., Cardinale M., Sundberg J.H., Spina F., Fusani L. 2020 Recent phenological shifts of migratory birds at a Mediterranean spring stopover site: Species wintering in the Sahel advance passage more than tropical winterers. *PLoS ONE* **15**(9), e0239489. (doi:10.1371/journal.pone.0239489)

429 21. Hötker H. 2002 Arrival of Pied Avocets *Recurvirostra avosetta* at the breeding site: Effects of winter quarters and consequences for reproductive success. *Ardea* **90**(3), 379–387.

432 22. Both C. 2010 Flexibility of Timing of Avian Migration to Climate Change Masked by Environmental Constraints En Route. *Curr Biol* **20**(3), 243–248. (doi:https://doi.org/10.1016/j.cub.2009.11.074)

435 23. Emmenegger T., Hahn S., Arlettaz R., Amrhein V., Zehtindjiev P., Bauer S. 2016 Shifts in vegetation phenology along flyways entail varying risks of mistiming in a migratory songbird. *Ecosphere* **7**(6), e01385. (doi:10.1002/ecs2.1385)

438 24. Hedlund J.S.U., Jakobsson S., Kullberg C., Fransson T. 2015 Long-term phenological shifts and intra-specific differences in migratory change in the willow warbler *Phylloscopus trochilus*. *J Avian Biol* **46**(1), 97-106. (doi:10.1111/jav.00484)

441 25. Rubolini D., Møller A.P., Rainio K., Lehikoinen E. 2007 Intraspecific consistency and geographic variability in temporal trends of spring migration phenology among European bird species. *Clim Res* **35**(1-2), 135-146. (doi:10.3354/Cr00720)

444 26. Saino N., Szép T., Romano M., Rubolini D., Spina F., Møller A.P. 2004 Ecological conditions during winter predict arrival date at the breeding quarters in a trans-Saharan migratory bird. *Ecol Lett* **7**(1), 21-25. (doi:10.1046/j.1461-0248.2003.00553.x)

448 27. Cohen E.B., Satterfield D.A. 2020 'Chancing on a spectacle:' co-occurring animal migrations and interspecific interactions. *Ecography* **43**(11), 1657-1671. (doi:10.1111/ecog.04958)

451 28. Brodersen J., Ådahl E., Brönmark C., Hansson L.-A. 2007 Ecosystem effects of partial fish migration in lakes. *Oikos* **117**(1), 40-46. (doi:10.1111/j.2007.0030-1299.16118.x)

454 29. Peacock S.J., Krkošek M., Lewis M.A., Molnár P.K. 2020 A unifying framework for the transient parasite dynamics of migratory hosts. *Proc Natl Acad Sci USA* **117**(20), 10897-10903. (doi:10.1073/pnas.1908777117)

457 30. Eichhorn G., Drent R.H., Stahl J., Leito A., Alerstam T. 2009 Skipping the Baltic: the emergence of a dichotomy of alternative spring migration strategies in Russian barnacle geese. *J Anim Ecol* **78**(1), 63-72. (doi:10.1111/j.1365-2656.2008.01485.x)

461 31. Beauchamp G. 2012 Foraging speed in staging flocks of semipalmated sandpipers: evidence for scramble competition. *Oecologia* **169**(4), 975-980. (doi:10.1007/s00442-012-2269-0)

464 32. Holdo R.M., Holt R.D., Coughenour M.B., Ritchie M.E. 2007 Plant productivity and soil nitrogen as a function of grazing, migration and fire in an African savanna. *J Ecol* **95**(1), 115-128. (doi:10.1111/j.1365-2745.2006.01192.x)

467 33. Mukhin A., Chernetsov N., Kishkinev D. 2008 Acoustic information as a distant cue for habitat recognition by nocturnally migrating passerines during landfall. *Behav*

Ecol **19**(4), 716-723. (doi:10.1093/beheco/arn025)

34. Németh Z., Moore F.R. 2007 Unfamiliar stopover sites and the value of social
information during migration. *J Ornithol* **148**(2), 369-376. (doi:10.1007/s10336-007-
0209-2)

35. García-Navas V., Sattler T., Schmid H., Ozgul A. 2021 Bird species co-
occurrence patterns in an alpine environment supports the stress-gradient hypothesis.
Oikos **130**(11), 1905-1918. (doi:10.1111/oik.08588)

36. Montaño-Centellas F.A. 2020 Interaction networks of avian mixed-species
flocks along elevation in the tropical Andes. *Ecography* **43**(6), 930-942.
(doi:10.1111/ecog.05135)

37. Pasquaretta C., Dubois T., Gomez-Moracho T., Delepoulle V.P., Le Loc'h G.,
Heeb P., Lihoreau M. 2021 Analysis of temporal patterns in animal movement
networks. *Methods Ecol Evol* **12**(1), 101-113. (doi:10.1111/2041-210x.13364)

38. Blanchet F.G., Cazelles K., Gravel D. 2020 Co-occurrence is not evidence of
ecological interactions. *Ecol Lett* **23**(7), 1050-1063. (doi:10.1111/ele.13525)

39. Bairlein F. 1995 *Manual of Field Methods. European-African Songbird*
Migration Network. revised ed. European-African songbird migration network.
Wilhelmshaven, Germany. 32 p.

40. Sander E.L., Wootton J.T., Allesina S. 2017 Ecological Network Inference From
Long-Term Presence-Absence Data. *Sci Rep* **7**(1), 7154. (doi:10.1038/s41598-017-
07009-x)

41. Harrell Jr. F.E., Dupont C. 2021 *Hmisc: Harrell Miscellaneous*. R package
version 4.5-0. (https://hbiostat.org/R/Hmisc/)

42. Storey J.D., Bass A.J., Dabney A., Robinson D. 2020 *qvalue: Q-value*
estimation for false discovery rate control. R package version 2.22.0.
(http://github.com/jdstorey/qvalue)

43. R Core Team. 2020 R: *A language and environment for statistical computing*.
(http://www.R-project.org/)

44. Blondel V.D., Guillaume J.-L., Lambiotte R., Lefebvre E. 2008 Fast unfolding
of communities in large networks. *J Stat Mech Theory Exp* **2008**(10), p10008.
(doi:10.1088/1742-5468/2008/10/P10008)

45. Newman M., Girvan M. 2004 Finding and evaluating community structure in
networks. *Physical Review E* **69**, 026113. (doi:10.1103/PhysRevE.69.026113)

46. Carstensen, D.W., Dalsgaard, B., Svenning, J.-C., Rahbek, C., Fjeldså, J.,
Sutherland, W.J. and Olesen, J.M. 2012 Biogeographical modules and island roles: a
comparison of Wallacea and the West Indies. *J Biogeogr* **39**, 739-749.
(doi:10.1111/j.1365-2699.2011.02628.x)

47. Fortuna M.A., García C., Guimarães Jr. P.R., Bascompte J. 2008 Spatial mating
networks in insect-pollinated plants. *Ecol Lett* **11**, 490–498. (doi:10.1111/j.1461-
0248.2008.01167.x)

48. Bellisario B., Cerfolli F., Nascetti G. 2010 Spatial network structure and
robustness of detritus-based communities in a patchy environment. *Ecol Res* **25**, 813–
821. (doi:10.1007/s11284-010-0711-5)

49. Svetunkov I. 2021 *smooth*: *Forecasting Using State Space Models*. R package
version 3.1.0. (https://github.com/config-i1/smooth)

50. Csárdi G., Nepusz T. 2006 *The igraph software package for complex network*
research, *InterJournal*, *Complex Systems* 1695. (http://igraph.org)

51. Pohlert T. In press. Non-Parametric Trend Tests and Change-Point Detection. ,
18.

52. Danon L., Díaz-Guilera A., Duch J., Arenas A. 2005 Comparing community
structure identification. *J Stat Mech Theory Exp* **2005**(09), P09008-P09008.
(doi:10.1088/1742-5468/2005/09/p09008)

53.   Pigot A.L., Sheard C., Miller E.T., Bregman T.P., Freeman B.G., Roll U.,
Seddon N., Trisos C.H., Weeks B.C., Tobias J.A. 2020 Macroevolutionary convergence
connects morphological form to ecological function in birds. *Nat Ecol Evol* **4**(2), 230-
239. (doi:10.1038/s41559-019-1070-4)
54.   Oksanen J., Blanchet F.G., Friendly M., Kindt R., Legendre P., McGlinn D.,
Minchin P.R., O'Hara R.B., Simpson G.L., Solymos P., et al. 2020 *vegan: Community*
Ecology Package. R package version 2.5-7. (http://CRAN.R-
project.org/package=vegan)
55.   Rubolini D., Spina F., Saino N. 2005 Correlates of timing of spring migration in
birds: a comparative study of trans-Saharan migrants. *Biol J Linn Soc* **85**(2), 199-210.
(doi:10.1111/j.1095-8312.2005.00484.x)
56.   Both C., Visser M.E. 2001 Adjustment to climate change is constrained by
arrival date in a long-distance migrant bird. *Nature* **411**(6835), 296-298.
(doi:10.1038/35077063)
57.   Jonzén N., Linden A., Ergon T., Knudsen E., Vik J.O., Rubolini D., Piacentini
D., Brinch C., Spina F., Karlsson L., et al. 2006 Rapid advance of spring arrival dates in
long-distance migratory birds. *Science* **312**(5782), 1959-1961.
(doi:10.1126/science.1126119)
58.   Usui T., Butchart S.H.M., Phillimore A.B. 2017 Temporal shifts and
temperature sensitivity of avian spring migratory phenology: a phylogenetic meta-
analysis. *J Anim Ecol* **86**(2), 250-261. (doi:10.1111/1365-2656.12612)
59.   Maggini I, Cardinale M, Favaretto A, Voříšek P, Spina F, Maoret F, Ferri A,
Riello S, Fusani L. 2021 Comparing population trend estimates of migratory birds from
breeding censuses and capture data at a spring migration bottleneck. *Ecol Evol* **11**, 967–
977. (doi:10.1002/ece3.7110)
59.   Fusani L., Cardinale M., Carere C., Goymann W. 2009 Stopover decision during
migration: physiological conditions predict nocturnal restlessness in wild passerines.
Biol Lett **5**(3), 302–305. (doi:10.1098/rsbl.2008.0755)
60.   Lafleur J.M., Buler J.J., Moore F.R. 2016 Geographic position and landscape
composition explain regional patterns of migrating landbird distributions during spring
stopover along the northern coast of the Gulf of Mexico. *Landscape Ecol* **31**(8), 1697-
1709. (doi:10.1007/s10980-016-0354-1)
61.   Ferretti A., Maggini I., Fusani L. 2021 How to recover after sea crossing: the
importance of small islands for passerines during spring migration. *Ethol Ecol Evol*
33(3), 307-320. (doi:10.1080/03949370.2021.1886181)
62.   Buler J.J., Moore F.R. 2011 Migrant–habitat relationships during stopover along
an ecological barrier: extrinsic constraints and conservation implications. *J Ornithol*
152, 101-112. (doi:10.1007/s10336-010-0640-7)
63.   Bayly N.J., Gomez C., Hobson K.A. 2013 Energy reserves stored by migrating
Gray-cheeked Thrushes *Catharus minimus* at a spring stopover site in northern
Colombia are sufficient for a long-distance flight to North America. *Ibis* **155**(2), 271-
283. (doi:10.1111/ibi.12029)
64.   Gómez C., Bayly N.J., Norris D.R., Mackenzie S.A., Rosenberg K.V., Taylor
P.D., Hobson K.A., Daniel Cadena C. 2017 Fuel loads acquired at a stopover site
influence the pace of intercontinental migration in a boreal songbird. *Sci Rep* **7**(1), 3405.
(doi:10.1038/s41598-017-03503-4)
65.   Mehlman D.W., Mabey S., E., Ewert D., N., Duncan C., Abel B., Cimprich D.,
Sutter R., D., Woodrey M. 2005 Conserving stopover sites for forest-dwelling
migratory landbirds. *Auk* **122**(4), 1281-1290. (doi:10.1642/0004-
8038(2005)122[1281:CSSFFM]2.0.CO;2)
66.   Ferretti A., McWilliams S.R., Rattenborg N.C., Maggini I., Cardinale M., Fusani
L. 2020 Energy Stores, Oxidative Balance, and Sleep in Migratory Garden Warblers

(*Sylvia borin*) and Whitethroats (*Sylvia communis*) at a Spring Stopover Site. *Integr*
Organ Biol **2**(1), obaa010. (doi:10.1093/iob/obaa010)

67.   Maggini I, Trez M, Cardinale M, Fusani L. 2020 Stopover dynamics of 12
passerine migrant species in a small Mediterranean island during spring migration.
Journal of Ornithology **161**, 793–802. (doi:10.1007/s10336-020-01768-7)

68.   Ferretti A, McWilliams SR, Rattenborg NC, Maggini I, Cardinale M, Fusani L.
2020 Energy Stores, Oxidative Balance, and Sleep in Migratory Garden Warblers
(Sylvia borin) and Whitethroats (Sylvia communis) at a Spring Stopover Site. *Integr*
Organ Biol **2**. (doi:10.1093/iob/obaa010)

69.   Ferretti A, Rattenborg N, Ruf T, Mcwilliams S, Cardinale M, Fusani L. 2019
Sleeping Unsafely Tucked in to Conserve Energy in a Nocturnal Migratory Songbird.
Curr Biol **29**, 2766-2772.e4. (doi:10.1016/j.cub.2019.07.028)

70.   Cooper WE, Pyron RA, Garland T. 2014 Island tameness: living on islands
reduces flight initiation distance. *Proc R Soc B* **281**, 20133019.
(doi:10.1098/rspb.2013.3019)

71.   Petalas C, Lazarus T, Lavoie RA, Elliott KH, Guigueno MF. 2021 Foraging
niche partitioning in sympatric seabird populations. *Sci Rep* **11**, 1–12.
(doi:10.1038/s41598-021-81583-z)

72.   Gayk ZG, Simpson RK, Mennill DJ. 2021 The evolution of wood warbler flight
calls: Species with similar migrations produce acoustically similar calls. *Evolution* **75**,
719–730. (doi:10.1111/evo.14167)

List of tables

Table 1 – List of the 31 most abundant species ringed at the stop over site of Ponza Island during the period 2007-2021.

Species	Wintering area	Foraging niche	Number of sampled individuals		
			Total	Mean (± SD)	Range [min-max]
Acrocephalus arundinaceus	Tropical	Invertivore glean arboreal	1,156	25.2 ± 11.391	[13-58]
Acrocephalus schoenobaenus	Sahel	Invertivore glean arboreal	378	77.067±48.792	[40-236]
Anthus trivialis	Tropical	Invertivore ground	2,197	146.467±44.171	[72-207]
Delichon urbica	Tropical	Invertivore sally air	1,283	85.534±49.393	[21-187]
Erithacus rubecula	North Africa	Invertivore ground	14,948	996.534±877.128	[84-2599]
Ficedula albicollis	Tropical	Invertivore sally air	11,504	121.734±99.349	[5-279]
Ficedula hypoleuca	Tropical	Invertivore sally air	1,826	766.934±271.304	[369-1393]
Fringilla coelebs	Tropical	Invertivore ground	421	28.067±20.537	[1-58]
Hippolais icterina	Tropical	Invertivore glean arboreal	18,850	1256.667±706.695	[325-2922]
Hirundo rustica	Sahel	Invertivore sally air	6,402	426.8±132.102	[208-700]
Jynx torquilla	Sahel	Invertivore ground	501	33.4±15.263	[17-67]
Lanius senator	Tropical	Invertivore ground	621	41.4±14.603	[17-71]
Luscinia megarhynchos	Tropical	Invertivore ground	1,976	131.733±50.941	[56-214]
Merops apiaster	Sahel	Invertivore sally air	757	50.466±26.848	[21-114]
Muscicapa striata	Sahel	Invertivore sally air	9,500	633.333±246.003	[187-1108]
Oenanthe oenanthe	Sahel	Invertivore sally ground	2,222	148.133±59.677	[67-259]

Oriolus oriolus	Tropical	Granivore ground	1,343	89.533±37.557	[39-182]
Phoenicurus ochruros	North Africa	Invertivore sally ground	2,750	183.333±142.749	[6-397]
Phoenicurus phoenicurus	Sahel	Invertivore sally ground	6,728	448.533±255.998	[155-1118]
Phylloscopus collybita	North Africa	Invertivore glean arboreal	5,880	392±288.111	[88-1050]
Phylloscopus sibilatrix	Tropical	Invertivore glean arboreal	15,402	1026.8±285.567	[597-1430]
Phylloscopus trochilus	Sahel	Invertivore glean arboreal	13,466	897.733±315.275	[424-1421]
Saxicola rubetra	Tropical	Invertivore sally ground	13,392	892.8±432.842	[7-1646]
Saxicola torquata	North Africa	Invertivore sally ground	780	52±60.523	[2-184]
Streptopelia turtur	Sahel	Granivore ground	724	48.266±36.526	[18-139]
Sylvia atricapilla	North Africa	Invertivore glean arboreal	13,885	925.666±1738.913	[29-5614]
Sylvia borin	Tropical	Invertivore glean arboreal	35,655	2377±1485.051	[337-5967]
Sylvia cantillans	Sahel	Invertivore glean arboreal	14,117	941.133±659.287	[259-2789]
Sylvia communis	Sahel	Invertivore glean arboreal	25,358	1690.533±1023.444	[40-3589]
Turdus philomelos	North Africa	Invertivore ground	1,985	132.333±146.932	[16-477]
Upupa epops	Sahel	Invertivore ground	526	35.066±18.525	[15-76]

List of figures

Figure 1 - Geographical framework of the study area within the Palearctic-African

migration system. The dotted circle in the box shows the 'Pontine' islands system with

the exact location of Ponza Island. Colours in the map show a synthetic view of the

main African ecoregions corresponding to the supposed wintering areas of species

analysed in this study.

Figure 2 - Trend of modularity (*Q*) in temporal co-occurrence networks. Dashed black

lines indicate the original time series, while solid and dotted lines show the measured

trends and uncertainty in trend slopes, respectively.

Figure 3 - (a) Heatmap showing the species' pairwise co-migration fidelity measured

by the normalized mutual information, *nmi*. Black boxes within the heatmap correspond

to the main cluster subdivision, while coloured boxes below and alongside dendrograms

show to which foraging niche and wintering area, respectively, a species belongs.

Colour codes are the same of (b), which show the boxplots of the *nmi* distribution

between different categories.