1 **Title:** Rapid thalamocortical network switching mediated by cortical synchronization underlies
2 propofol-induced EEG signatures: a biophysical model
3

Authors: Austin E. Soplata[1,2,+], Elie Adam[3,4+], Emery N. Brown[1,3,4,5], Patrick L. Purdon[1],
Michelle M. McCarthy[2,*], Nancy Kopell[2,*]
Affiliations:
1. Department of Anesthesia, Critical Care and Pain Medicine, Massachusetts General
Hospital, Harvard Medical School, Boston, MA, USA
2. Department of Mathematics & Statistics, Boston University, Boston, MA, USA
3. Department of Brain and Cognitive Sciences, Massachusetts Institute of Technology,
Cambridge, MA, USA
4. Picower Institute for Learning and Memory, Massachusetts Institute of Technology,
Cambridge, MA, USA
5. Institute for Medical Engineering and Science, Massachusetts Institute of Technology,
Cambridge, MA, USA
* Co-senior authors
+ Co-first authors

18
Contributions:
Austin E. Soplata: Conceptualization, Data curation, Investigation, Methodology, Software,
Analysis, Validation, Visualization, Writing, Editing. Elie M. Adam: Software, Visualization,
Analysis, Writing, Editing. Emery N. Brown: Funding acquisition, Project administration,
Supervision, Editing. Patrick Purdon: Funding acquisition, Data curation, Project administration,
Supervision, Editing. Michelle M. McCarthy: Conceptualization, Analysis, Writing, Editing.
Nancy Kopell: Conceptualization, Analysis, Funding acquisition, Project administration,
Supervision, Writing, Editing.
27
28 **Running Head:** Synchronization switches between propofol oscillatory states
29

30 **Address for Correspondence:** Austin E. Soplata, email: austin.soplata@gmail.com , address:
31 École Polytechnique Fédérale de Lausanne, Blue Brain Project, Campus Biotech, Chemin des
32 Mines 9, 1202 Geneva, Switzerland.

33
34 **Key Words:** thalamus, cholinergic modulation, slow rhythms, alpha rhythms, synchrony
35

Abstract:
Propofol-mediated unconsciousness elicits strong alpha/low-beta and slow oscillations in the
electroencephalogram (EEG) of patients. As anesthetic dose increases, the EEG signal changes
in ways that give clues to the level of unconsciousness; the network mechanisms of these
changes are only partially understood. Here, we construct a biophysical thalamocortical
network involving brainstem influences that reproduces transitions in dynamics seen in the EEG
involving the evolution of the power and frequency of alpha/low beta and slow rhythm, as well
as their interactions.

Our model suggests propofol engages thalamic spindle and cortical sleep mechanisms to elicit
persistent alpha/low-beta and slow rhythms, respectively. The thalamocortical network fluctuates
between two mutually exclusive states on the timescale of seconds. One state is characterized by

continuous alpha/low-beta frequency spiking in thalamus (C-state), while in the other, thalamic
alpha spiking is interrupted by periods of co-occurring thalamic and cortical silence (I-state). In
the I-state, alpha co-localizes to the peak of the slow; in the C-state, there is a variable
relationship between an alpha/beta rhythm and the slow oscillation. The C-state predominates
near loss of consciousness; with increasing dose, the proportion of time spent in the I-state
increases, recapitulating EEG phenomenology. Cortical synchrony drives the switch to the I-state
by changing the nature of the thalamocortical feedback. Brainstem influence on the strength of
thalamocortical feedback mediates the amount of cortical synchrony. Our model implicates loss
of low-beta, cortical synchrony, and coordinated thalamocortical silent periods as contributing to
the unconscious state.

57
New & Noteworthy:
GABAergic anesthetics induce alpha/low-beta and slow oscillations in the EEG, which interact
in dose-dependent ways. We construct a thalamocortical model to investigate how these
interdependent oscillations change with propofol dose. We find two dynamic states of
thalamocortical coordination, which change on the timescale of seconds and dose-dependently
mirror known changes in EEG. Thalamocortical feedback determines the oscillatory coupling
and power seen in each state, and this is primarily driven by cortical synchrony and brainstem
neuromodulation.
66
Introduction:
Unconsciousness mediated by GABAergic anesthetics, such as propofol and sevoflurane, is char-
acterized by a presence of alpha/low-beta (8-20 Hz) and slow oscillations (0.5-2.0 Hz) in the
electroencephalogram (EEG) (Brown, Purdon, and Van Dort 2011; Lewis et al. 2012; Purdon et
al. 2013; Mukamel et al. 2014; Scheinin et al. 2018; Malekmohammadi et al. 2019; Stephen et al.
2020). As effect site concentration (dose) is increased, the EEG shows less low-beta, lower fre-
quency alpha, less alpha power, and increased slow wave power (Purdon et al. 2013; Mukamel et
al. 2014; Stephen et al. 2020). The alpha (8-14 Hz) and slow oscillations are interdependent and
the level of interdependence was found to indicate the level of unconsciousness (Purdon et al.
2013). (Purdon et al. 2013; Mukamel et al. 2014; Gaskell et al. 2017; Stephen et al. 2020) high-
lighted two states they called "peak-max" and "trough-max". These are two states in which the
slow and alpha rhythms are coupled by phase-amplitude coupling (PAC). In the peak-max, the
amplitude of the alpha rhythm is maximal during the peak of the slow rhythm; in trough-max,
the maximal alpha amplitude appears in the trough of the slow. The trough-max state appears
close to the loss of consciousness (LOC), whereas the peak-max state occurs with deeper levels
of propofol. However, during most of the time spent under propofol, the EEG does not reflect
either of these states. This is shown in Fig. 1.
84

Figure 1. Typical spectral changes seen in the EEG as propofol dose is increased. Human experimental data from a single subject used in (Purdon et al. 2013; Mukamel et al. 2014) (see Methods). (A) Raw EEG traces (top) and EEG filtered at alpha/low-beta, alpha and slow (middle) at three different doses of propofol (bottom) starting near loss of consciousness (LOC) and up to but not including the state of "burst suppression" (Brown, Lydic, and Schiff 2010; Ching et al. 2012). Low-beta power appears predominately near LOC. (B) Spectrogram of the EEG during stepped increases in propofol dose in a healthy volunteer.

Anesthetics act on their molecular targets, and it is then through network mechanisms that they alter brain rhythms (Brown, Lydic, and Schiff 2010; Brown, Purdon, and Van Dort 2011). The network mechanisms underlying the EEG phenomena above remain unclear. Our previous work (Soplata et al. 2017) suggests alpha emerges from thalamus under propofol and can be modulated by an external source of slow oscillation such as that modeled in (Compte et al. 2003; Benita et al. 2012). Additionally, the alpha/slow peak-max increases as dose increases. However, our model was open loop, where cortex provides feedforward input to thalamus without feedback. It is unknown what dynamics are possible under a closed-loop interaction. Yet this interaction is essential during unconsciousness, for instance, to explain the anteriorization of alpha oscillations (Vijayan et al. 2013). In this work, we develop a biophysical model of closed-loop thalamocortical interaction that explains all the dose-dependent EEG phenomena above and shows the importance of thalamocortical feedback. Our Hodgkin-Huxley type model builds on

105 (Soplata et al. 2017; Ching et al. 2010; Compte et al. 2003; Benita et al. 2012) to incorporate
106 influences of cortex, thalamus and brainstem on EEG dynamics.
107
108 In this paper, we use 8-12 Hz as the range for alpha rather than the 8-14 Hz range used in
109 (Purdon et al. 2013; Mukamel et al. 2014). We note that 8-12 Hz is the standard range for alpha
110 in the cognitive literature (Van Diepen, Foxe, and Mazaheri 2019; Foster and Awh 2019). In the
111 current paper, we aim to relate the propofol-induced rhythms to loss of cognitive function; and
112 thus, we make a distinction between alpha (8-12 Hz) and low-beta (13-20 Hz).
113
We show that the dynamics on small space and time scales are highly complex: on each slow
cycle, there is one of two network states, which can change after some indeterminate number of
slow cycles. One of those states is called the "C-state" and is characterized by a continuous ("C")
alpha/low-beta oscillations in thalamus; the other state is called "I-state" in which there is
thalamic spiking at an alpha frequency interrupted ("I") by a time period within a slow
oscillation cycle in which there is no thalamic activity. The I-state is highly related to what has
been called peak-max, since the alpha activity is co-localized with the peak of the slow
oscillation. The interaction of alpha, low-beta, and slow during the C-state is more variable than
in the I-state, and the alpha portion includes what has previously been called "trough-max". We
show that the statistics of these states are dose-dependent, with higher doses of propofol
corresponding to a larger percentage of the I-state. A significant finding of the work is that the
statistics of the two states are strongly influenced by the synchrony of the cortical cells. Thus, the
depth of anesthesia corresponds to the statistics of the I-states and C-states. Unconsciousness is
associated with a prevalence of the I-state and thus a higher degree of cortical synchronization.
Such increased synchronization has been reported experimentally (Gutiérrez et al. 2022). See the
Discussion for more details.
130
Systemic administration of GABAergic anesthetics effects all structures in the brain, and the
influence of these drugs on the brainstem can alter neuromodulatory systems (Moody et al.
2021). Our simulations show that when feedback from thalamus to cortex is potentiated because
of these alterations, the synchronization of cortex is facilitated, and makes the switch to the I-
state more probable. Endogenous noise in the system facilitates switching back to the C-state.
Our work suggests that cortical synchronization due to potentiated thalamic feedback resulting
from neuromodulatory alterations are key in understanding the mechanism of GABAergic
anesthetics mediated unconsciousness. The loss of beta with increasing dose also has
implications for loss of long-distance communication needed for consciousness.
140
141 **Results**
142
Model of propofol acts on cortical, thalamic and brainstem biophysics.
We develop a model consisting of interacting thalamic and cortical circuits (see Methods). Our
goal is to understand, and relate to loss of consciousness, the physiologic and network
mechanisms related to EEG changes as propofol dose increases (Purdon et al. 2013; Mukamel et
al. 2014; Flores et al. 2017; Stephen et al. 2020; Bastos et al. 2021) (Fig. 1). Among the spectral
features that we investigate with our modeling are: (a) increased co-localization of the alpha with
the peak of slow as dose is increased (Fig. 1A), (b) increased amplitude of the slow oscillation
with increasing propofol dose (Fig. 1A), (c) increased low-beta power near loss of consciousness

(LOC) and return of consciousness (ROC) (Fig. 1A,B), and (d) decreased mean frequency of
alpha and decreased alpha power with higher doses (Fig. 1B). We show below that each of these
features reflects properties of the model suggestive of mechanisms of loss of consciousness.
154
We model the cortical circuit using 100 pyramidal cells (PY) and 20 cortical interneurons (IN)
from (Compte et al. 2003; Benita et al. 2012) (Fig. 2A). The pyramidal cells are modeled with
two compartments: the soma (PYso) and the dendrite (PYdr). The thalamic circuit is modeled as
in (Soplata et al. 2017), except with 20 thalamocortical cells (TC) and 20 thalamic reticular
neurons (TRN). Our model currents are conductance-based with Hodgkin-Huxley dynamics.
Details of the currents used in each neuron type, network connectivity, and all other aspects of
the model can be found in the Methods and Appendix.
162
163

164
Figure 2: Propofol enables thalamocortical network slow and alpha oscillations. (A) Schematic of
thalamocortical network model. TC: thalamocortical cells; TRN: thalamic reticular cells; PY: pyramidal cells
(PYdr: dendrite, PYso: soma); IN: cortical interneurons. EEG is modeled as the sum of PYdr voltages. (B)
Rastergram of each cell/compartment in wake-like mode with no sensory input, in which each black line
represents a spike by each cell/compartment. (C) Rastergram of a simulation under propofol conditions. (Left)
Simulation over 30 seconds. (Right) A zoom-in showing alpha/low-beta spiking in thalamus. (D) Rastergram
of a cortex-only simulation showing slow wave generation in the absence of thalamus when the maximal
conductance of the K(Na) current was 0.10 mS/cm^2 representing the level seen during sleep. (E) Variations in
model EEG. (Top) EEG taken from 23 to 25 seconds. Bottom EEG taken from 11 to 13 seconds. (F)
Spectrogram of the model EEG from the simulation shown in (C).
175
We modeled the addition of propofol as five changes from wake-like conditions. The first is an
increase in the maximal conductance and the inhibition time constant of the GABA receptors,
which are known to be produced by propofol (McCarthy, Brown, and Kopell 2008; Ching et al.
2010). The second is a decrease in the maximal conductance of the TC cell H-current (Ying et al.
2006). Both were used in our previous work without feedback connections from thalamus to
cortex (Soplata et al. 2017). The last three effects of propofol account for its action on brainstem
circuits: specifically, propofol potentiates inhibitory GABAergic circuits in various arousal
centers in brainstem (Brown, Purdon, and Van Dort 2011), including cholinergic centers. We

focus on the enhanced inhibitory effects on brainstem cholinergic circuits (but see Discussion for
other types of neuromodulation). We do not model the brainstem circuitry directly but rather
model the effect of decreasing cortical cholinergic tone in the presence of propofol (Kikuchi et
al. 1998; Nemoto et al. 2013; Meuret et al. 2000; Pal and Mashour 2021; Luo et al. 2020). This
brainstem effect on decreased cortical cholinergic tone is modeled by strengthening (1)
intracortical AMPAergic synaptic conductance, (2) TC→PY thalamocortical AMPAergic
synaptic conductances, and (3) PY K(Na)-current maximal conductance (see Methods). As
propofol dose is increased, we increase the strength of the intracortical and thalamocortical
AMPA conductances. All other propofol changes are not further changed with dose. The EEG of
the model is produced by the summation of the voltages in the dendrites of all cortical pyramidal
cells (see Methods).
195
Propofol generates slow waves in model cortex and alpha/low-beta waves in model
thalamus.
The "wake-like" condition without propofol produces a "depolarized relay state" (Destexhe et al.
1996) in thalamus as shown in Fig. 2B. In the depolarized relay state, the thalamus can transmit
incoming signals; however, no such signals are included in the wake-like state and thus thalamus
is silent (see (Ching et al. 2010; Krishnan et al. 2016) for similar models of wake-like thalamus).
This simulation shows infra-slow oscillations in cortical cells (Fig. 2B), which are known to be
present in the awake state (Krishnan, González, and Bazhenov 2018). The slow oscillation in our
model originates from the cortex due to the K(Na) current (Compte et al. 2003) (see Methods).
The firing rates of cells are consistent with known data (Dash, Autio, and Crandall 2022).
206
With propofol, the thalamocortical network produces prominent slow and alpha/low-beta
oscillations that are visible in the raster plots (Fig. 1C). Here we show with a blow-up of Fig. 1C
that the alpha/low-beta oscillation is visible in the thalamic TC cell spiking. The network
dynamics can change in time in a way that will be examined in detail below. We find that the
model cortex alone can generate slow oscillations and thus thalamic involvement is not needed
(Fig. 2D). With propofol, an alpha oscillation arises in thalamus from the potentiation of GABA-
A and the decrease in H-current, as shown in our previous work, which simulates thalamus in the
absence of cortex. A description of the alpha generating mechanism is in (Soplata et al. 2017)in
the section entitled "Propofol induces sustained alpha via changing the balance of
excitation/inhibition." This alpha uses the interaction between TC and TRN and engages
thalamic spindling mechanisms. In our previous work, we did not include thalamocortical
feedback; our current work shows that the thalamic alpha oscillations persist in the presence of
thalamic feedback (Fig. 2C).
220
Our model EEG shows strong slow and alpha/low-beta oscillations (Fig. 2E). The alpha/low-beta
oscillations are more evident (Fig. 2E, top) in the first and last 10 seconds of the simulation in
Fig. 2C than during the middle 10 seconds (Fig. 2E, bottom). A spectrogram of the model EEG
from the simulation in Fig. 2C, shows a continuous slow oscillation and a prominent higher
frequency oscillation that switches between alpha/low-beta (8-20 Hz) and a lower average
frequency alpha (~ 8 Hz) (Fig. 2F).
227
Propofol induces rapid switching between two distinct thalamocortical network states.

With propofol, the network switches between two mutually exclusive network states on a rapid
timescale (seconds) both on the single cell level and the population level (Fig. 2C, 3A-F), each of
which can span over multiple cycles of the slow oscillation. The main difference between these
states is the periodic cessation of spiking in the thalamus in one state but not the other; when this
happens, the cortex and thalamus are simultaneously silent (Fig. 3A, D). The thalamic silence
occurs because the thalamus enters a silent depolarized state, which stops the thalamus from
spiking and thus thalamic input to cortex is lost. We call this network state the "I-state" (short for
"interrupt"). In contrast, in the other network state, there are no thalamic silent periods because
spiking from thalamus in continuous. Thus, we refer to this network state as the "C-state" (short
for "continuous").
239
240

241
Figure 3: Different slow cycles display I-state or C-state during propofol. (A-C) Representative voltage
traces of each cell/compartment (A) during an I-state, (B) across the entire propofol simulation, with I-state
highlighted in orange and C-state highlighted in blue, and (C) during a C-state. (D-F) Rastergram of all spiking
activity (D) during the I-state (E) across the entire simulation, with coupling regimes highlighted, and (F)
during the C-state. (G-I) Model EEG filtered at alpha/low beta, alpha and slow (G) during the I-state, (H)
across the entire simulation and (I) during the C-state.

248
Both states show slow oscillations and either alpha or alpha/low-beta oscillations in their spiking
patterns (Fig. 3D-F) as well as in the model EEG (Fig. 3G-I) (See Methods for definition of
model EEG). Although the alpha and low-beta rhythms are not a cortically generated rhythm,
they appear in the model EEG due to the TC connections onto the pyramidal cell dendrites (Fig.

2A). During the I-state the thalamic fast frequency is predominately alpha (8-12 Hz) (Fig. 3D,G);
during the C-state the fast thalamic frequencies range from 8-20 Hz and thus spans the
alpha/low-beta range (Fig. 3F,I). These two network states correspond to different interactions
between the slow and the alpha or alpha/low-beta in the EEG. In the I-state, there is always high
amplitude alpha associated with the peak (or rising phase) of the slow oscillations (Fig. 3G); the
trough of the slow oscillations corresponds to the simultaneous silent periods in thalamus and
cortex (Fig. 3G). In contrast, the C-state has a variable relationship between alpha/low-beta and
slow (Fig. 3I). The thalamic spiking persists continuously throughout the slow oscillation cycles
with some variation in the relationship of the alpha/low-beta amplitude to the phase of the slow.
In particular, the alpha and low-beta tend to couple at different phases of the slow oscillation
during the C-state. Our simulations suggest that one possible coupling during the C-state is a
transient coupling of alpha to the trough of the slow oscillation as seen on some slow cycles in
Fig. 3I, as well as in experimental literature (Purdon et al. 2013; Mukamel et al. 2014). When
this occurs, the low-beta couples to a different phase of the slow oscillation (Fig. 3I). Note that,
in the C-state, the slow oscillation has lower amplitude than in the I-state (compare Fig. 3G and
Fig. 3I).
269
In summary, the thalamocortical network rapidly switches between two states with propofol. The
I-state is characterized by periods of coordinated thalamic and cortical silence as well as high
cortical synchrony. In contrast, the C-state displays ongoing activity in thalamus and less cortical
synchronization. The EEG during the I-state shows alpha oscillations but lacks low-beta
oscillations, large amplitude slow oscillations, and co-localization of the alpha to the peak of the
slow. In contrast, during the C-state the EEG has alpha and low-beta oscillations, smaller
amplitude slow oscillations and variable alpha/slow coupling with alpha coupling to the trough
being one possibility.
278
Cortical synchrony determines thalamocortical state by modulating network feedback
dynamics.
Cortical synchrony is a critical determinant of thalamocortical state under propofol determining
both the thalamic state (continuous spiking or interrupted spiking) as well as the spectral features
of each state. The spiking in the cortex is noticeably more synchronized during the I-state than in
C-state (Fig. 3D, F). The level of cortical synchrony affects the depolarization level of the
thalamus and thus the thalamic spiking dynamics. Indeed, the less synchronized cortex in the C-
state allows the thalamus to remain hyperpolarized and thus to continue engaging spindle
dynamics (Soplata et al. 2017), while the additional synchronization in cortex during the I-state
depolarizes the thalamus into its silent depolarized state. Thus, cortical synchrony is a critical
determinant of thalamocortical state under propofol.
290
The C-state and I-state have different types of thalamocortical interactions under propofol. In the
C-state, when the thalamus is spiking continuously (due to low cortical synchrony), the thalamus
provides positive feedback to the cortex. In the I-state, the synchronous cortical input stops the
thalamus by putting it its silent depolarized phase, whereas cortical silence promotes thalamic
spiking (due to the hyperpolarization of the thalamus). Thus, the type of feedback in the I-state is
homeostatic: cortical excitation leads to negative feedback from thalamus via silencing thalamus,
whereas reduced cortical firing causes excitatory feedback from thalamus. The switch in
feedback regimes (and thus network state) is abrupt because, as soon as the cortical

synchronization level is sufficient to silence the thalamus, the network switches to homeostatic
feedback.
301
The type of corticothalamic feedback dictates the spectral coupling observed during the different
states. In the C-state, the alpha often occurs near the trough of the slow wave, while the low-beta
occurs more towards the peak of the slow wave. This phenomenon stems from the influence of
cortical spiking on the excitation level of the thalamus: when cortical spiking is low, as in the
trough of the slow wave, the thalamus is more hyperpolarized and thus has a lower frequency of
spiking (alpha spiking). In contrast, when the cortical spiking is higher, as in the peak of the slow
wave, the thalamus is more depolarized and thus the thalamic spiking is at beta. The positive
feedback from thalamus to cortex engages the activity-dependent K(Na)-current with the thala-
mus giving less excitation (alpha) to cortex when the cortical activity is low and more excitation
(low-beta) when the cortical excitation is high. Note that this accounts for increasing frequency
of the K(Na)-mediated slow oscillations between the wake-like state (~ 0.2 Hz) and the C-state
(1 Hz) (Fig. 2B, C). In contrast, when the network is in its homeostatic feedback regime (I-state),
cortical synchrony leads to thalamic silence, which in turn leads to a profound loss of cortical
spiking. This simultaneous loss of thalamic and cortical spiking is reflected in a large slow wave
trough with no alpha coupling. The thalamus responds to the loss of cortical input by hyperpolar-
izing into its spindling regime and again spiking, now at predominately alpha, which depolarizes
the cortex into its active phase. The diminished cortical K(Na), which decreased during the corti-
cal inactive phase, additionally primes the cortex to spike more synchronously in response to the
thalamic input. The more synchronous spiking during the cortical active phase is reflected in the
EEG as a higher amplitude peak in the slow wave. The alpha spiking in the thalamus is coupled
to the peak of the slow wave since this is the only phase at which the thalamus is active.
323
Summarizing the connection between cortical synchronization and thalamic state, we find that
under propofol the thalamocortical network can abruptly switch between two dynamic networks
states governed by the level of synchronization in the cortex. The spectral features of the EEG
during these two states are the results of a switch in thalamic feedback dynamics: positive feed-
back during the C-state and homeostatic feedback during the I-state. In particular, we find that
during the state of positive corticothalamic feedback (C-state), the amplitude and duration of the
slow oscillation is controlled by the kinetics of the activity-dependent K(Na) current in the cor-
tex, whereas during the state of homeostatic corticothalamic feedback (I-state), the amplitude and
duration of the slow oscillation is influenced by the thalamic feedback. The larger amplitude
slow waves during the I-state are a consequence of engaging the homeostatic thalamocortical
feedback, while the low amplitude slow waves during the C-state result primarily from cortical
K(Na) dynamics.
336
To verify the relationship between the cortical synchronization and the thalamocortical state, we
tested whether we could change the thalamocortical state by introducing artificial cortical
synchronization or de-synchronization. We applied 100 milliseconds of either a synchronizing or
desynchronizing input to cortex during a period when the system was in a C-state. Artificial
synchronization of cortex switched the thalamocortical network to an I-state, whereas with
artificial the thalamocortical network remained in the C-state (Fig. 4A-E). These results support
cortical synchronization as a driver of C-state to I-state transitions. When cortical synchronizing
or de-synchronizing inputs were applied to the network when the system was in an I-state, we

found that de-synchronizing inputs induced a transition to the C-state, whereas the network
remained in the I-state in response to synchronizing inputs (Fig. 5). These results highlight
cortical de-synchronization as a key determinant of I-state to C-state transitions, and thus a key
determinant of thalamocortical state under propofol.
349
350

351
Figure 4. Cortical synchronization induces a C-state to I-state transition. (A) Rastergram of a "baseline"
simulation in which most of the time is spent in the C-state. For every 100 millisecond interval expressing the
C-state in the baseline simulation, two additional simulations were run: one simulation where a Synchronizing
Input stimulus (see Methods) was applied to PYdrs during that specific 100 millisecond interval (example is
shown in B), and another simulation where a DE-synchronizing Input stimulus (see Methods) was applied to
PYdrs over that specific 100 millisecond interval (example is shown in C). (B) Rastergram showing an
example simulation where Synchronizing Input was applied during the time 5700-5800 ms, causing the
remainder of the simulation to exhibit I-state behavior. Red line marks the time of the input. (C) Rastergram

showing an example of DE-synchronizing Input for the same time interval as B, but where the simulation
almost completely remains in the C-state. (D) Bar graph showing, for all simulations receiving Synchronizing
Inputs at different times, the proportion of post-stimulus simulation time that exhibited a C-to-I-state transition.
Scoring was done by visual inspection, where each simulation was marked as "All" if >95% of simulation time
after the Synchronizing Input exhibited I-state; analogously, each simulation was otherwise marked as "Most"
= 95%-50%, "Half" = ~50%, "Some" = 50%-10%, or "None" = <10%. The bar chart is the summation of all
simulations receiving a Synchronizing Input and indicates that the artificial synchrony brought by the
Synchronizing Input was very effective at converting C-states to I-states. (E) Similar to (D) but for baseline
simulations receiving DE-synchronizing input, indicated that DE-synchronizing Input was not effective in
eliciting C-state to I-state transitions.
370
371

372
Figure 5. Cortical de-synchronization causes an I-state to C-state transition. This figure is analogous to
Figure 4 but studying I-to-C transitions instead of C-to-I transitions. (A) Rastergram of a "baseline" simulation
in which most of the time is spent in the I-state. For every 100 millisecond interval expressing the I-state in the

baseline simulation, two additional simulations were run: one simulation where a Synchronizing Input stimulus
(see Methods) was applied to PYdrs over that specific 100 millisecond interval (example is shown in B), and
another simulation where a DE-synchronizing Input stimulus (see Methods) was applied to PYdrs over that
specific 100 millisecond interval (example is shown in C). (B) Rastergram showing an example simulation
where Synchronizing Input was applied during time 3200-3300 ms but did not lead to C-state behavior. Red
line marks the time of the input. (C) Rastergram showing an example of DE-synchronizing Input for the same
time interval as B, but where the remainder of the simulation exhibits C-state behavior. (D) Bar graph
showing, for all simulations receiving Synchronizing Inputs at different times, the proportion of post-stimulus
simulation time that exhibited an I-to-C state transition. Scoring was done as in Fig. 4. The bar chart is the
summation of all simulations receiving a Synchronizing Input and indicates that the artificial synchrony
brought by the Synchronizing Input was not effective at converting I-states to C-states. (E) Similar to (D) but
for baseline simulations receiving DE-synchronizing input, indicated that DE-synchronizing Input was very
effective in eliciting I-state to C-state transitions.
389
390 **Propofol potentiates TC feedback via brainstem neuromodulation, facilitating**
391 **synchronization of cortex and a switch to the I-state as dose increases.**
392 In the previous section, we showed that cortical synchronization is a key driver of C-state to I-
393 state transitions in corticothalamic circuits. Here we propose the physiological mechanism by
394 which increased cortical synchronization occurs as propofol dose is increased.
395
We model increasing propofol dose by progressively decreasing ACh neuromodulation on
corticothalamic circuits (see Methods): we increased maximal AMPA synaptic strength between
cortical pyramidal cells ($\bar{g}_{AMPA:PY \rightarrow PY}$) and from TC cells to cortical pyramidal cells
($\bar{g}_{AMPA:TC \rightarrow PY}$), which strengthens feedback from thalamus to cortex. As propofol dose is
increased in our model, the amount of time spent in the C-state progressively decreases and is
replaced by more time spent in the I-state (Table 1). The progression in co-localization of alpha
to the peak of the slow oscillation recapitulates the findings in subjects undergoing propofol-
anesthesia induced unconsciousness.
404
405

gAMPA	Mean ± Std Dev
0.002	83.7% ± 9.81%
0.003	83.26% ± 16.74%
0.004	70.37% ± 25.87%
0.005	44.39% ± 30.94%
0.006	32.03% ± 34.22%
0.007	15.61% ± 23.60%
0.008	6.97% ± 15.61%
0.009	4.15% ± 11.35%
0.01	1.29% ± 3.26%
0.012	0.65% ± 3.27%

406 **Table 1: Proportion of time spent in the C-state when both TC→PY and PY→PY are set to the**
407 **corresponding maximal gAMPA conductance**. The units for gAMPA are mS/cm^2. For each gAMPA value,
408 measurements were taken across approximately 70 independent simulations that were each 30 seconds long.
409
To determine whether the change to more I-states relies primary on TC→PY or PY→PY
synapses, we looked at the incidence of I-states when changing one of these synapses at a time.
We found that the dose-dependent increased time spent in the I-state is primarily due to

increasing the TC→PY AMPA synapse (Table 2). The strengthened feedback leads to more
synchronized cortical cells, which in turn causes the switch to the thalamocortical homeostatic
feedback regime (see previous section) associated with I-state dynamics.
416
417

		TC→PY gAMPA (mS/cm^2)			
		0.004	0.006	0.008	0.01
PY→PY gAMPA (mS/cm^2)	0.004	**58.49%** ± 25.07%	**32.69%** ± 30.00%	**12.79%** ± 14.31%	**4.82%** ± 11.01%
	0.006	**54.53%** ± 32.70%	**21.33%** ± 24.49%	**8.42%** ± 10.33%	**3.15%** ± 8.92%
	0.008	**52.52%** ± 25.46%	**27.92%** ± 27.83%	**6.05%** ± 11.04%	**4.67%** ± 8.56%
	0.01	**43.06%** ± 25.29%	**17.71%** ± 19.18%	**7.07%** ± 13.46%	**2.36%** ± 5.54%

418 **Table 2: Proportion of time spent in the C-state with differential values of TC→PY and PY→PY**
419 **maximal gAMPA conductance.** All values in form of "**mean** ± standard deviation". For each set of gAMPA
420 values, measurements were taken across approximately 40 independent simulations that were each 30 seconds
421 long.
422
That the C-state to I-state switch requires thalamocortical feedback is supported by the finding
that in the absence of TC feedback, the thalamocortical networks remain in the C-state (Fig. 6). It
also shows that the cortical synchronization caused by the K(Na)-production of the slow wave
does not produce sufficient cortical synchrony to induce the C-state to I-state transition. In
contrast, we showed in the last section that the switch from the I-state to the C-state is facilitated
by cortical desynchronization. This can occur physiologically under propofol due to cortical
noise (Fig. 3).
430
431

432
433 **Figure 6. Lack of thalamocortical connections disables transitions from C-state to I-state.** Rastergram of
434 a "high-dose" ($\bar{g}_{AMPA:PY→PY}$ = 0.008 mS/cm^2) propofol simulation where TC→PY AMPA synapses have been
435 set to zero. Simulations without thalamocortical feedback never exhibited an I-state.
436

The increasing time spent in the I-state with higher doses of propofol accounts for several dose-
related EEG findings including: (1) less low-beta because this frequency is predominately in the
C-state, (2) more lower frequency alpha because the alpha in the I-state has a lower average
frequency than the alpha in the C-state (see Fig. 4B), (3) decreased alpha power because alpha is
only present for a short period on the peak of the slow wave in the I-state, (4) loss of trough-max
because trough-max is seen only in the C-state and only occasionally in that state, (5) increased
peak-max due to thalamic alpha spiking occurring only during the peak of the slow wave in the
I-state, (6) increased slow wave amplitude/power due to the I-state having larger amplitude slow
oscillations as a result of increased cortical synchronization and its coordinated cortico-thalamic
silent states.
447
Discussion
Overview and Clinical Implications
The anesthetic propofol produces oscillatory signatures on the electroencephalogram (EEG):
prominent alpha/low-beta oscillations (8-20 Hz), slow oscillations (0.5-2.0 Hz), and increased
co-localization of the alpha to the peak of the slow with increasing effect site concentration
(Mukamel et al. 2014). Here we use computational models to examine the role of brainstem,
thalamus, and cortex in producing these oscillations and shaping the interactions between them.
Specifically, our results help to understand the biophysical origin of the rhythms and the role of
the rhythms in producing loss of consciousness.
457
One surprising finding in our simulations is that the thalamocortical network switches at the
timescale of seconds between two dynamic states characterized by continuous alpha/low-beta in
thalamus (C-state) or by transient interruption of the thalamic alpha (I-state). In the latter, the
alpha is co-localized to the peak of the slow rhythm ("peak-max"), whereas in the former, there
need not be consistent co-localization. What is called "trough-max" in the literature is here a
special case of the C-state seen in the literature at sustained low doses of propofol that are used
in experiments with volunteers; with clinical use, these stages are passed through quickly
(Purdon et al. 2013; Mukamel et al. 2014; Stephen et al. 2020). In the I-state, cortical and
thalamic silent periods are coordinated, which is the reason for the peak-max coupling. We
further show that increases in propofol dose lead to statistically more prevalence of the I-state
versus the C-state. Cortical synchrony is a key driver of the transitions between the I-state and
the C-state: we show that corticothalamic state transitions are mediated by the level of cortical
synchrony. We illustrate this by introducing artificial synchrony and desynchrony in cortical
spiking during ongoing periods of the I-state or the C-state (Figs. 4 and 5). Cortical synchrony
can switch the C-state to the I-state, and desynchrony can cause the I-state to switch to the C-
state. One determinant of cortical synchrony in our network is thalamocortical feedback; this is
strengthened by neuromodulatory changes related to propofol dose, biasing the expression of the
I-state over the C-state as the dose of propofol increases.
476
Our model reproduces several clinical EEG observations as propofol dose increases: less low-
beta power, lower frequency alpha, less alpha power, loss of trough-max coupling, increased
peak-max coupling, and increased slow wave amplitude/power. The model provides physiologi-
cal and network mechanisms for these effects, which give insights into how propofol works to
produce loss of consciousness. The dynamic switch between two types of thalamocortical feed-
back, positive feedback in the C-state and homeostatic feedback in the I-state, determine the

corticothalamic spiking patterns as well as the EEG spectral changes that correlate with each state. As dose is increased, brainstem effects on the thalamocortical feedback enable increased cortical synchrony, and thus, I-states become more prevalent. The increasing proportion of time spent in the I-state with larger doses of propofol in our model accounts for all the experimental spectral changes mentioned above associated with increasing propofol dose. Cortical synchronization has been found experimentally to relate to the level of unconsciousness (Gutiérrez et al. 2022), and thus these EEG measures may indicate the level of propofol-induced unconsciousness.

Our model helps explain why earlier work emphasized the trough-max state (Purdon et al. 2013; Mukamel et al. 2014): they were filtering in the alpha band (8-12 Hz), while our C-state shows non-slow oscillations between 8-20 Hz. As can be seen in Figs. 1 and 3I, the alpha band is sometimes found in the trough of the slow oscillation with beta oscillations at other phases. Lower doses of propofol associated with the time of loss of consciousness (LOC) are also characterized by a larger beta power than seen either at resting wakefulness or at higher doses of propofol (Purdon et al. 2013; Mukamel et al. 2014; Stephen et al. 2020; Krom et al. 2020; Malekmohammadi et al. 2019). Our modeling suggests this beta comes from thalamus in the presence of propofol. The model also suggests why the lower doses of propofol that commonly occur around LOC are associated with a light state of anesthesia (Mukamel et al. 2014): the beta oscillations communicated to the cortex are believed to be the basis of long-distance coordination in cortex (Siegel et al., 2012) and hence, signals from the thalamus can be transferred more widely throughout the cortex. By contrast, modeling has suggested that alpha disrupts such coordination (Weiner et al. 2023; Jones et al. 2000), which is consistent with alpha being the predominant rhythm with deeper levels of propofol anesthesia. Thus, a ratio of low-beta power to alpha power may be a way to monitor the depth of propofol-induced unconsciousness near the onset of LOC.

Relation to prior modeling work
Our previous computational modeling (Soplata et al. 2017) suggested thalamus as the source of propofol-induced alpha (although not low-beta), and showed that dose-dependent coupling between alpha and slow occurred if slow oscillations were imposed on the thalamus from cortex via an "open-loop". In this work, we simulate a "closed-loop" thalamocortical network of Hodgkin-Huxley cells with feedback from both thalamus to cortex and cortex to thalamus. Most of the results we find are a consequence of this feedback. Our cortical model is derived from models that can generate slow oscillations during sleep (Compte et al. 2003; Benita et al. 2012).

While prior modeling demonstrated thalamocortical circuits could produce propofol alpha oscillations (Ching et al. 2010; Vijayan et al. 2013), or propofol alpha and slow (Krishnan et al. 2016), our previous work (Soplata et al. 2017) was the first modeling investigation of which we are aware into the unique alpha-slow PAC dynamics of propofol. The essential results of the previous paper continue to hold in our current model: (1) The alpha is generated by propofol acting on thalamic circuits; (2) Our previous hypothesis that propofol-induced slow oscillations can come from the cortex is supported by the new modeling; (3) Cortical slow and alpha can interact in different ways depending on the hyperpolarization of the thalamus. However, details of the coupling between alpha and slow oscillation are altered from our previous work by the interaction between the thalamus and cortex, which was not explicitly modeled in that prior work.

Some mechanisms have been changed from our prior model and yield a better fit with
experimental data: (1) in the prior thalamus-only model, the maximal thalamic spiking frequency
under propofol was alpha. In this modeling work, the thalamus can spike as high as low-beta (~
20 Hz) (which has been seen in experiments (Ujma et al. 2022; Wang et al. 2022)). (2) in the
previous model, thalamic alpha emerged only during the cortical silent phase of the slow during
low-dose propofol and only during the cortical active phase of the slow during high-dose. In the
current model, in which the hyperpolarization level of the thalamus is influenced by the cortex,
we find that, during the C-state, the thalamus does not hyperpolarize enough to completely stop
thalamic spiking during the cortical active phase; rather, the thalamus produces alpha/low-beta
throughout all phases of the slow oscillation. As a result of this, alpha/low-beta is more
prominent, and thus will show higher power, during low-dose time periods, which the C-state is
more prevalent in (see Fig. 1B) (Purdon et al. 2013; Mukamel et al. 2014; Stephen et al. 2020).
(3) The addition of thalamocortical feedback allows changes in cortical synchronization with
increase in dose: continuous alpha/low-beta in thalamus results from less cortical
synchronization, and hence the slow waves that appear in cortical EEG have a significantly
lower amplitude and greater variability (Fig. 1B,C; Fig. 3 G,I) during the C-state than during the
I-state (Mukamel et al. 2014). (4) Another difference from previous work is that, in the current
model during the I-state, the thalamic silence during the cortical silent period is due to
depolarization from corticothalamic excitation, rather than hyperpolarization as suggested by the
thalamus-only model. This difference is significant in linking cortical synchronization to state
changes: cortical synchronization depolarizes thalamus out of its alpha spiking into a silent state,
changing the dynamics to the I-state. (5) In this paper, we explicitly model the cortex rather than
using it just as an input to the thalamus. Thus, we were able to model the EEG by using the sum
of pyramidal cell dendritic voltages (Kirschstein and Köhling 2009).
553
Other models of alpha oscillations exist in the literature with cortical as well as thalamic sources.
Most of the modeling literature about cortical alpha concerns the awake state (Silva, Amitai, and
Connors 1991; Jones et al. 2000; Bollimunta et al. 2011; Sigala et al. 2014; Lozano-Soldevilla,
ter Huurne, and Oostenveld 2016; Köster and Gruber 2022) (but also see (Vijayan et al. 2013)).
In the context of propofol anesthesia however, the prominent alpha observed in the EEG is likely
to come from the thalamus: alpha oscillations under propofol are globally coherent, whereas
slow oscillations are not (Lewis et al. 2012; Purdon et al. 2013). This suggests cortical alpha is
coming from a non-cortical source such as the thalamus. Our earlier modeling suggests the
thalamus can produce an ongoing alpha in the presence of propofol via potentiated GABA-A
hijacking thalamic spindle mechanisms (Soplata et al. 2017). This is a prediction of both our
previous and current model that remains to be tested experimentally. A recent experimental and
computational study looking at oscillations with another GABAergic anesthetic, isoflurane,
suggests this anesthetic can also generate thalamic alpha oscillations (Jiang et al. 2022).
567
Propofol, slow oscillations, and brainstem-mediated neuromodulation
A notable interpretation of this model is that the anesthetic effects of propofol depend on
neuromodulatory effects as much as on GABA-A inhibition and TC cell H-currents. ACh release
in rodent frontal cortex can be decreased by 70-85% in the presence of propofol (Kikuchi et al.
1998; Nemoto et al. 2013). In the model, lowered ACh affects several cortical currents: (1)
lowered ACh increases the K(Na) current (Schwindt, Spain, and Crill 1989; McCormick 1992)
enabling slow oscillations; (2) lowered cortical ACh increases intracortical AMPAergic synaptic

conductances (Compte et al. 2003; Benita et al. 2012; Krishnan et al. 2016); and (3) cortical
muscarinic receptors respond to lowered ACh by increasing thalamocortical AMPA conductance
(Kruglikov and Rudy 2008; Favero, Varghese, and Castro-Alamancos 2012). The latter is critical
in our model for producing the greater synchrony of the cortex needed to switch to the I-state
more often as dose is increased.
580
Noradrenergic tone is also decreased in the presence of GABAergic anesthetics: propofol and
sevoflurane potentiate inhibitory GABAergic synaptic activity from the pre-optic area of the
hypothalamus onto noradrenergic neurons of the locus coeruleus, which diffusely project to
cortex among other areas (Brown, Pavone, and Naranjo 2018). In the cortex, norepinephrine
(NE), as well as ACh, abolish slow oscillatory activity mediated by cortical Up and Down states
(Favero, Varghese, and Castro-Alamancos 2012). Thus, lowered cortical NE due to propofol
may work collaboratively with ACh in the producing increased slow oscillations in cortex.
Unlike ACh, which non-specifically suppresses PY→PY and TC→PY excitatory (AMPA), NE
selectively suppresses intracortical excitatory inputs (Favero, Varghese, and Castro-Alamancos
2012). Although we find the dominant effect on cortical synchronization with propofol is due to
increasing the thalamocortical AMPA conductance, NE likely contribute to the overall cortical
synchronization by increasing the intracortical AMPA conductance.
593
We note that propofol likely also utilizes known slow mechanisms via its effects on non-
cholinergic brainstem neuromodulatory systems. Propofol affects not just the cholinergic sources
in the basal forebrain, laterodorsal tegmental area, and pedunculopontine tegmental area, but also
inhibits the tuberomammillary nucleus, locus coeruleus, dorsal raphe nucleus, ventral
periaqueductal gray, and lateral hypothalamus (Brown, Lydic, and Schiff 2010; Brown, Purdon,
and Van Dort 2011). These areas respectively provide histamine, norepinephrine, serotonin,
dopamine, and orexin/hypocretin to the cortex (Brown, Lydic, and Schiff 2010; Brown, Purdon,
and Van Dort 2011). Many of these neuromodulators affect various potassium currents that are
critical in known models of slow oscillations, including the K(Na)-current, the NaP-current, and
potassium leak currents (Schwindt, Spain, and Crill 1989; McCormick 1992). These
neuromodulators can also affect both excitatory and inhibitory currents in the cortex, and can
change the relative impact of thalamocortical synapses (McCormick 1992; Favero, Varghese,
and Castro-Alamancos 2012; Kuo and Dringenberg 2008). Additionally, cortical
neuromodulation by NE, as well as ACh, is known abolish SWO and is active during
awake/relay states (Moody et al. 2021). There is still much we do not understand about how all
these neuromodulators work in concert together (Krishnan et al. 2016; Moody et al. 2021).
610
It is known that the cholinesterase inhibitor physostigmine reverses propofol LOC (Meuret et al.
2000); assuming LOC depends on slow oscillations and synchronization of the cortex, our model
suggests that the engagement of ACh-modulated currents by propofol may explain this
experimental result: physostigmine acts to increase ACh levels, which would decrease $\bar{g}_{K(Na)}$ in
our model and stop the K(Na)-dependent slow oscillation produced by propofol. ACh also acts to
weaken thalamocortical connections (Favero, Varghese, and Castro-Alamancos 2012) and
therefore release the cortex from over-synchronization; less synchronized states are associated
with the awake state. Thus, loss of ACh may be a major contributor to propofol-induced loss of
consciousness. The fact that our model requires neuromodulatory changes to produce propofol
oscillations and their coupling suggests that the effects of propofol on the brainstem may be

critical for its oscillatory phenomena, which is supported by active experimental research on
propofol and other anesthetics (Moody et al. 2021; Minert, Yatziv, and Devor 2017; Minert,
Baron, and Devor 2020; Muindi et al. 2016; Vlasov et al. 2021; Pal and Mashour 2021). Since
the transition from C-state to I-state in our model is associated with lowering ACh (without
changing GABAergic effects), we predict that a smaller dose of physostigmine may produce less
I-state and more C-state and thus, may lead to more EEG low-beta, higher frequency alpha,
increased alpha power, less peak-max, and lower slow wave amplitude.
628
Propofol, slow oscillations, and sleep
Our work suggests that propofol utilizes not only thalamic spindling mechanisms (Soplata et al.
2017), but also natural sleep slow mechanisms and changes in neuromodulation to produce its
oscillatory effects. The K(Na)-current is the primary mechanism of slow oscillation generation in
the cortical sleep slow oscillation model we used (Sanchez-Vives, Nowak, and McCormick
2000; Schwindt, Spain, and Crill 1989; Compte et al. 2003). Most other slow oscillation models
rely on a combination of changes to cortical excitatory/inhibitory plasticity and/or the persistent
sodium current (NaP) (Bazhenov et al. 2002; Hill and Tononi, Giulio 2004; V. Crunelli et al.
2011; Sanchez-Vives and McCormick 2000; Timofeev et al. 2000; Krishnan et al. 2016). The
NaP-current has been shown to be functionally coupled to the K(Na)-current (Hage and Salkoff
2012), and therefore the K(Na)-current may contribute to these mechanisms. Some models of
slow UP state initiation, also called DOWN-to-UP transitions, rely on random cortical excitation
(Timofeev et al. 2000), synaptic plasticity changes (Krishnan et al. 2016; Sanchez-Vives and
McCormick 2000), or TC initiation of cortical slow UP states (V. Crunelli et al. 2011). In our
simulations, the DOWN-to-UP transition is not reliant on the above mechanisms but rather
initiates when the hyperpolarizing K(Na)-current in a PY cell has decayed sufficiently to allow
significant PY spiking. In our I-state, however, the slow oscillation is primarily a manifestation
of the change in thalamocortical feedback dynamics. The network feedback switches from a
positive feedback dynamic in the C-state to a homeostatic feedback dynamic in the I-state. The
homeostatic feedback produces wide swings in the amplitude of the slow oscillations, which
differs from low amplitude K(Na)-mediated slow oscillation in the C-state. Future work designed
to differentiate natural sleep slow oscillations versus general anesthetic slow mechanisms will
enable finer-grained experiments into how the loss of consciousness occurs in these two distinct
states.
653
Propofol, memory consolidation, and aging
Our investigation of thalamocortical dynamics under propofol may have implications for
memory and aging. During natural sleep, memory consolidation onto cortical axo-dendrite
connections likely occurs during the nesting of hippocampal ripples during thalamic spindles,
which themselves are nested inside thalamocortical sleep slow oscillations (Penagos, Varela, and
Wilson 2017). Based on our current and previous work (Soplata et al. 2017), propofol alpha and
slow oscillations hijack some of the same mechanisms used during sleep-related memory
consolidation: propofol alpha derives from the thalamic spindling mechanism, and propofol slow
oscillations engage the cortical K(Na) current also engaged by sleep. Proper memory
consolidation requires correct encoding of worthwhile memories during sleep (Stickgold 2005),
but if application of propofol abnormally activates some of the same oscillations in this process,
this may lead to invalid memory consolidation or interfere with synaptic-dendrite networks
involved in storing memory. A recent experiment showed promising results in using propofol to

disrupt reconsolidation of traumatic memories (Galarza Vallejo et al. 2019), which could help
treat Post-Traumatic Stress Disorder patients. Additionally, alpha power and, to a lesser extent,
slow power under propofol may indicate a subject's "brain age" (Purdon et al. 2015). Our
modeling predicts that propofol alpha may predominately arise from the thalamus, and therefore
a decrease in propofol alpha power across age could correlate with brain fitness via losses in the
ability of the thalamus to burst at alpha or enter the spindling regime (Purdon et al. 2015),
myelination retention of thalamocortical afferents (Peters 2002), or the strength of
thalamocortical synapses onto cortical dendrites (Morrison and Baxter 2012).
675
Caveats, Limitations, and Future Modeling
Our model is relatively small considering its significant complexity: there are a total of 160
neurons among multiple types in both cortex and thalamus. Larger models may sometimes
display behavior that is not captured in smaller ones because of the degree of heterogeneity that
is possible. Delineating such behavior is beyond the scope of this work. Also, not included in our
model is complexity associated with the multiple layers of the cortex and the multiple kinds of
nuclei in the thalamus. The use of Hodgkin-Huxley modeling in this study is justified when
examining the effects of anesthetic drugs such as propofol, which work to change the kinetics of
synaptic receptors and the conductances of intrinsic membrane currents that interact to sculpt the
network behavior (Brown, Purdon, and Van Dort 2011). As such, the size and complexity of the
model is driven by the questions asked. We found that simpler models, both in cellular
components and neuron numbers, were not sufficient to replicate the spectrum of propofol-
induced EEG changes explored in this paper and understand their mechanisms. As to further
detail, our philosophy is to look for a minimal model, however complex, that will reproduce the
interactions of the slow and alpha rhythms consistent with experimental results. Regarding the
effects of brainstem modulation, we focused almost entirely on cholinergic modulation. Effects
of other neuromodulators, briefly mentioned, are left for future work. Even the effects of ACh
were not completely investigated; notably, we did not look at the effects of changing K(Na)
strength with successively larger effect site concentrations of propofol.
695
Heterogeneity in TC→PY and PY→PY strength and connectivity across cortical regions and
layers may contribute to diversity in cortical synchronization levels (Redinbaugh et al. 2020;
Malekmohammadi et al. 2019) and therefore patient- or region-specific diversity in time spent in
the I-state versus the C-state. Our simulations indicate that different effect site concentrations of
propofol tend to express different proportions of these two network states on a small spatial
scale. Our results also suggest that, under propofol, different local cortical networks may, on a
fast timescale, switch between the I-state and the C-state even while a regional EEG signal
predominantly shows a single type of dose-dependent PAC. By introducing region-specific
heterogeneity to cortex (e.g., sensory and higher-order) and thalamus (e.g., core and matrix),
future simulations may be able to investigate the significant spatiotemporal changes between
low- and high-dose propofol. "Anteriorization" is a well-known phenomenon in which propofol
administration initially leads to the loss of awake, occipital alpha and an increase in frontal alpha
(Tinker, Sharbrough, and Michenfelder 1977; Cimenser et al. 2011; Vijayan et al. 2013). This
frontal alpha is at its strongest and most persistent state during low-dose propofol, before
spreading to become region-nonspecific during high-dose propofol (Cimenser et al. 2011;
Purdon et al. 2013; Mukamel et al. 2014; Stephen et al. 2020) and decreasing in power with
increasing dose (Gutiérrez et al. 2022). Slow power is also greater during high-dose than low-

dose propofol (Purdon et al. 2013; Mukamel et al. 2014; Mhuircheartaigh et al. 2013; Lee et al.
2017) and may modulate higher frequencies more in frontal regions during high-dose (Stephen et
al. 2020). We note that our current modeling captures many of these results for frontal cortex: (1)
alpha is strongest and most persistent during low-dose propofol; (2), alpha decreases in power
with increasing dose; (3) slow power is greater during high dose than low-dose propofol.
718
In the future, modeling multiple cortices will allow us to probe why holding propofol at a low-
dose results in trough-max coupling that is most prevalent in frontal cortex (Mukamel et al.
2014), why there is stronger frontal slow modulation at higher doses (Stephen et al. 2020), and
why there is increased thalamocortical alpha coherence in this region (Flores et al. 2017).
Modeling multiple cortices will also allow us to explore coherence, phase (Malekmohammadi et
al. 2019), and firing rate (Krom et al. 2020) discrepancies found between frontal and sensory
regions under anesthesia. Understanding how region-specific heterogeneity affects cross-cortical
communication and frontal cortex specifically may help to validate theories of loss of
consciousness, including frontoparietal disconnection (Hudetz and Mashour 2016) and similar
connectivity changes (Banks et al. 2020), brainstem changes to neuromodulation (Brown,
Purdon, and Van Dort 2011), alpha blocking of processing (Palva and Palva 2007), and slow
oscillation control of activity (Gemignani et al. 2015; Stephen et al. 2020).
731
There is strong recent evidence that thalamic simulation can reverse unconsciousness during
low-dose propofol (Bastos et al. 2021). For our model to reproduce this, we would need to
introduce additional mechanisms. One possible mechanism for this interesting phenomenon is
that 30-seconds-long thalamic stimulation may activate group I metabotropic glutamatergic
receptors (mGluRs) on the cortical cells receiving thalamocortical input. Thalamocortical
activation of these receptors can produce depolarizing effects (Sherman 2014) and antagonism of
cortical mGluRs leads to anesthesia-like effects (Suzuki and Larkum 2020). The very strong
thalamocortical excitation elicited by thalamic stimulation may lead to activation of these
mGluRs in the hyperpolarized cortical cells, depolarizing the cortical cells. This cortical
depolarization could lead to a temporary reduction in cortical DOWN states (reducing SWO
power) and an increase in higher power activity. Conversely, since mGluRs also exist at
corticothalamic synapses (V. Crunelli et al. 2011; Sherman 2014), another contributor to the
reversal of sedation could be that thalamic stimulation leads to thalamocortical excitation, which
leads to corticothalamic stimulation that activates mGluRs, temporarily depolarizing the
thalamus.
747
Implications for unconsciousness
Higher doses of propofol corresponding to increased co-localization of alpha and the peak of
slow have been associated with unarousable consciousness, whereas lower doses of propofol
with less peak-max co-localization lead to a state of unconsciousness in which arousability is
possible (Purdon et al. 2013; Mukamel et al. 2014; Scheinin et al. 2018). Our modeling suggests
that the depth of anesthesia is associated with the degree of predominance of the I-state over the
C-state. Since, in our model, cortical synchrony is a key driver of the transition from the C-state
to the I-state and cortical desynchrony causes the opposite transition, our results suggest that
increased cortical synchrony corresponds to deeper levels of unconsciousness. This has also been
noted in the experimental paper (Bharioke et al. 2022) on unconsciousness and synchrony in

layer 5 cortical cells. Such increased synchronization prevents the flexible and fast changes of
coordination needed for normal cognitive processing (Cannon et al. 2014; Buzsáki 2006).
760
761 Our model suggests lower doses of propofol are associated with more time spent in the C-state.
762 This state is associated with low-beta (as well as alpha) oscillations in the thalamus. As discussed
763 above, beta oscillations have been documented to be highly involved in long-distance
764 coordination in the brain (Siegel, Donner, and Engel 2012). Thus, the loss of this beta could
765 contribute to the higher degree of unconsciousness associated with higher doses.
766
One unintuitive finding suggested by our model was that TC neurons may be depolarized into
"relay mode" during I-state and could potentially relay sensory information during this window,
even during deep anesthesia. In our simulations, strong corticothalamic excitation after
synchronized active cortical states increased the membrane potential of TC cells during the I-
state, as shown in Fig. 3A,D. This increase was enough to interrupt the intrinsic alpha bursts of
the thalamus, but if this occurs at the same time as strong sensory input spikes, the TC cells may
be depolarized enough to briefly relay sensory spiking information up to the cortex. Recently,
even in humans under low-dose propofol anesthesia, auditory stimuli resulted in wake-like
cortical neural activity in primary auditory cortex but not higher-order cortex (Krom et al. 2020).
This suggests that some thalamic sensory relay may still occur under propofol anesthesia, even if
changes to cross-cortical communication prevent its higher-order processing. Furthermore, in our
simulations, the I-state may occur during individual slow cycles of both low- and high-dose
propofol, indicating this brief sensory relay may occur at any point during propofol anesthesia
(Malekmohammadi et al. 2019; Krom et al. 2020).
781
782 **Methods**
783
784 *Model Design*
785
Our Hodgkin-Huxley network, illustrated in Figure 2 A, consists of 100 cortical pyramidal
dendritic compartments (PYdr), 100 corresponding cortical pyramidal somatic/axonal
compartments (PYso), 20 cortical interneuron cells (INs), 20 thalamic thalamocortical cells
(TCs), and 20 thalamic reticular neurons (TRNs). All equations and parameters used in the
model are available in both the Appendix and the model code (Soplata 2023b; 2023a). The
thalamic cells are identical to those used in (Soplata et al. 2017) and therefore derived from
(Destexhe et al. 1996; Ching et al. 2010), except that we used a population size of 20 for each
cell class rather than 50 due to memory/RAM limitations. The cortical compartments and cells
are implemented according to their original description in (Compte et al. 2003), except that we
include a simple leak-current in our PYdr compartments; we suspect that the original paper
accidentally neglected to list this current. While there are many cortical slow models to choose
from (Lytton, Destexhe, and Sejnowski 1996; Destexhe et al. 1996; Sanchez-Vives and
McCormick 2000; Timofeev et al. 2000; Bazhenov et al. 2002; Destexhe and Sejnowski 2003;
Hill and Tononi, Giulio 2004; Vincenzo Crunelli and Hughes 2010; V. Crunelli et al. 2011), we
use this particular K(Na)-based sleep slow cortical model (Compte et al. 2003) due to its
simplicity, experimental basis (Sanchez-Vives and McCormick 2000), and effective utilization in
other slow models (Benita et al. 2012; Taxidis et al. 2013).
803

804 *Model Connectivity*
805
All connections are illustrated in Figure 2 A and available in both the Appendix and the model
mechanism code (Soplata 2023a). AMPA connections include from PYso to neighbor-only PYdr
(PYso→PYdr also called PY→PY), from PYso to IN (PY→IN), from TC to TRN (TC→TRN),
from TC to PYdr (TC→PY), from TC to IN (TC→IN), from PYso to TRN (PY→TRN), and
from PYso to TC (PY→TC). Intracortical AMPA connections (PYso→PYdr and PYso→IN)
included synaptic depression. NMDA connections include from PYso to PYdr and from PYso to
IN and include synaptic depression. GABA-A connections include from IN to PYso (IN→PY),
from IN to neighbor-only IN (IN→IN), from TRN to TC (TRN→TC), and from TRN to TRN
(TRN→TRN). GABA-B connections are only from TRN to TC (TC→TRN). Finally, the only
connections that are not chemical synapses are the simple compartmental connections between
each PYdr compartment and its corresponding PYso compartment. Note that we use PY→PY to
refer exclusively to AMPAergic PYso→PYdr connections.
818
For all synapses, each source cell is connected to its "nearest-neighbors" (2*radius+1) target
cells, where the radius is 10 cells. This makes all connectivity ratios, or how many connections
are projecting from each source cell, approximately 1:20. For our intrathalamic connectivity,
since our population sizes are 20, this effectively makes all thalamic connections all-to-all
connected, just like in (Ching et al. 2010; Soplata et al. 2017). For our intracortical connectivity
ratio of 1:20, we based this primarily on the model from which we drew our SWO mechanism,
(Compte et al. 2003) (1:20 with a standard deviation of 5), and subsequent work (Benita et al.
2012). Other models utilized either higher (Traub et al. 2005; Cruikshank et al. 2012) (ranging
from 1:5 to 1:50, with most at 1:10 or 1:20) or lower (Krishnan et al. 2016) (1:10) connectivity
ratios. Since the original model used a connectivity ratio that was comparable to similar models,
and so as not to disturb network dependencies of the SWO mechanism, we elected to use their
same intracortical connectivity ratio. Similarly, for the thalamocortical connectivity ratio of 1:20,
we based our parameters on (Krishnan et al. 2016) (TC:PY ratio of 1:20). This gives us equal
intracortical and thalamocortical connectivity ratios of both 1:20.
833
All synaptic conductances are normalized across the number of incoming connections of a given
synapse type. For PY→PY connections, this is straightforward, since the size of the source and
target cell populations are equal, leading to each PY cell receiving 20 of these connections. The
maximal conductance of each of these incoming connections is computed to be 1/20th of the
total maximal conductance for this synapse type. For connections where the source and
population sizes are not equal, such as TC→PY connections, this is more complex. Each TC cell
has 20 projections to the PY population, but since there are 20 TCs, this equals 400 total
projections to go to 100 PY cells. Ultimately, this results in each PY cell receiving connections
from 4 TC synapses, resulting in each synapse being normalized to 1/4th of the total maximum
conductance. If the total maximum conductances of PY→PY and TC→PY are equal (see next
paragraph), then each connection from a single source TC cell to a single target PY cell will have
a larger maximal synaptic conductance than each single PY to PY connection, but there will be
fewer TC to PY connections.
847
Except for the simulations done for Table 2, the total maximal conductances of our intracortical
PY→PY and thalamocortical TC→PY AMPA synapses were kept equal to each other and

changed simultaneously. We assumed that the ratio between these two maximal synaptic
conductances could be held to be equal due to prior thalamocortical models using the same
(Ching et al. 2010) or similar (Krishnan et al. 2016) (0.020:0.024) ratios. Additionally, by using
equal values for these conductances, we could model the effects of propofol decreasing ACh on
each of these synapse types in identical ways (see Table 1). We could then compare the effects
on the system of treating these conductances heterogeneously (see Table 2) to understand their
relative contribution to the case where they are set identically.
857
The total maximal conductances used for the PY→PY and TC→PY AMPA synapses ranged
from 0.002 mS/cm^2 to 0.012 mS/cm^2. These values change in response to the concentration of
ACh present, but the relationship between concentration and proportion of change in
conductance is not clear, so we were required to explore a range of values; for the relationship of
ACh to these conductances, see the subsection "Propofol effects" below. We initially based our
intracortical and thalamocortical total maximal conductances on the default value used for
PY→PY AMPA total maximal conductance in the original cortical SWO model paper (Compte
et al. 2003): 5.4 nS, which is divided over a pyramidal dendritic surface area of 0.035 mm^2 to
give 0.0154 mS/cm^2. Since this value is meant to correspond to an NREM sleep state which
already exhibits low ACh concentration (Compte et al. 2003), we could use this value to simulate
anesthetized states, but would need to decrease the value to simulate states of lower levels of
anesthesia (see the aforementioned subsection below). We did simulate higher values such as
0.0154 mS/cm^2 as in the original paper, but these were not significantly different than the I-state
shown in the paper, and continued the trend shown in the Tables of a dominance of I-state.
872
873 *Model Inputs*
874
For all simulations, in order to model background activity, excitatory Poisson spiketrains were
input into all PY, TC, and TRN cells. These spiketrains had a firing rate of 40 Hz and were
convolved with an exponential with a decay time of 2 milliseconds. The total maximal
conductance of these virtual synapses were always held equal to that of the total maximal
conductance of PY→PY and TC→PY AMPA synapses, except for the simulations in Table 2,
where the Poisson inputs had conductances of 0.004 mS/cm^2. See the mechanism code (Soplata
2023a) for implementation details.
882
For Figures 4 and 5, the "Synchronizing Input" was applied to all PYdr compartments for a
duration of 100 milliseconds in each simulation and had a constant amplitude of 1.0 uA/cm^2. For
the "DE-synchronizing Input", the input had a duration of 100 milliseconds, but each PYdr
compartment received a different constant amplitude randomly pulled from a uniform
distribution of -1.0 to +1.0 uA/cm^2. In both cases, the amplitudes did not change for the duration
of the stimulus and were 0 outside of the stimulus time.
889
890 *Propofol effects*
891
Similarly to our previous work (Soplata et al. 2017), we model how increasing propofol affects
the thalamus via changing three parameters: decreasing TC cell H-current maximal conductance
($\bar{g}_H$), and potentiating all GABA$_A$ synapses via increasing maximal conductance ($\bar{g}_{GABA_A}$) and
GABA$_A$ decay time constants (τ_{GABA_A}). Propofol may decrease $\bar{g}_H$ (Ying et al. 2006; Cacheaux et

al. 2005), although the magnitude of this change is experimentally unknown (Chen 2005). To
shift from a wake-like state to a propofol-anesthetized state, we decrease $\bar{g}_H$ from 0.04 to 0.005
mS/cm^2, which is in line with previous anesthetic and sleep research using this thalamic model
(Destexhe et al. 1996; Vijayan et al. 2013; Ching et al. 2010; Soplata et al. 2017).
900
For our propofol simulations, we tripled $\bar{g}_{GABA_A}$ and τ_{GABA_A} for all GABA$_A$ synapses (both
cortical and thalamic), since doubling these GABA$_A$ parameters produced only I-state, but
tripling lead to the presence of C-state. We originally based the magnitude of our propofol
GABA$_A$ changes on prior modeling work (McCarthy, Brown, and Kopell 2008). In our previous
paper (Soplata et al. 2017), we found that our thalamus-only network could produce persistent
alpha oscillations if we doubled or tripled these GABA$_A$ parameters. In Figure 4 of (Soplata et al.
2017), we showed that thalamic persistent alpha occurred across a broader range of inputs when
tripling the parameters compared to doubling. In the current paper, for all anesthetic simulation
variations, doubling GABA$_A$ parameters produces little simulation time with persistent thalamic
alpha oscillations. Instead, only by tripling GABA$_A$ parameters do the simulations produce C-
state for a substantial or majority of simulation time. This may be due to effects on lower-dose
behavior caused by additional cortical cell types that we did not include, namely from those of
(McCarthy, Brown, and Kopell 2008).
914
Propofol decreases cortical acetylcholine (ACh) (see Introduction), and we model these
cholinergic changes via increasing intracortical AMPAergic synaptic conductances ($\bar{g}_{AMPA:PY \to PY}$)
(Compte et al. 2003; Benita et al. 2012; Krishnan et al. 2016), TC→PY thalamocortical
AMPAergic synaptic conductances ($\bar{g}_{AMPA:TC \to PY}$) (Kruglikov and Rudy 2008; Favero, Varghese,
and Castro-Alamancos 2012), and K(Na)-current maximal conductance ($\bar{g}_{K(Na)}$) (Compte et al.
2003; Benita et al. 2012). ACh affects thalamocortical afferent synapses in different ways:
decreased nicotinic ACh receptor activation weakens thalamocortical synapses, but decreased
muscarinic ACh receptor activation strengthens them (Kruglikov and Rudy 2008; Favero,
Varghese, and Castro-Alamancos 2012; Gil, Connors, and Amitai 1997; Hsieh, Cruikshank, and
Metherate 2000; Oldford and Castro-Alamancos 2003; Eggermann and Feldmeyer 2009). Based
on the rapid desensitization of nicotinic ACh receptors (Quick and Lester 2002), the slowly-
changing, metabotropic nature of muscarinic receptors, and their similar shifts in natural sleep
(McCormick 1992), we believe that muscarinic receptors could exert a stronger effect than
nicotinic receptors on thalamocortical afferents, therefore increasing $\bar{g}_{AMPA:TC \to PY}$ with increasing
propofol dose.
930
The actual proportion of change that ACh can cause in the intracortical and thalamocortical
synapses is unclear. Different computational models use a range of proportional increases to
$\bar{g}_{AMPA:PY \to PY}$ caused by ACh, including up to +75% (Vijayan and Kopell 2012) or +15% to
+100% (Krishnan et al. 2016). Data on how much ACh may increase $\bar{g}_{AMPA:TC \to PY}$ is much more
scarce, but the increase may be as high as +300% (see Figure 8 F of (Favero, Varghese, and
Castro-Alamancos 2012)). We therefore focused our synaptic changes on a wide range of
decreases from 0.0154 mS/cm^2, the derived value of the low-ACh sleep state used in the cortical
SWO model of (Compte et al. 2003).
939
For the K(Na)-current maximal conductance $\bar{g}_{K(Na)}$, we only used two values: 0.10 mS/cm^2 in
our wake-like simulation (similar to the tonic, wake-like state of Figure 14 of the original paper

(Compte et al. 2003), or 1.33 mS/cm^2 for all anesthetic states. We specifically chose to not make
small changes to $\bar{g}_{K(Na)}$ between anesthetic states because it has impacts on the frequency of the
SWO produced, which the original paper investigates. Keeping the SWO frequency stable
allowed us to more effectively investigate how PAC can arise, and how it can be changed by
covariation of intracortical and thalamocortical synaptic strength.
947
948 *Specific simulation parameters*
949
The changes to parameters for the wake-like simulation from those of the Appendix are as
follows (all units in mS/cm^2): $\bar{g}_{AMPA:TC \to PY}$, $\bar{g}_{AMPA:PY \to PY}$, and all Poisson inputs set to 0.004, PY
$\bar{g}_{K(Na)}$ 0.10, TC $\bar{g}_H$ 0.4, TC $\bar{g}_{K\text{-Leak}}$ 0.001. Additionally for the wake-like simulation, the
following synapses were decreased by 60% compared to their values in the Appendix:
$\bar{g}_{NMDA:PY \to PY}$, $\bar{g}_{NMDA:PY \to IN}$, $\bar{g}_{AMPA:PY \to IN}$, $\bar{g}_{AMPA:TC \to IN}$, $\bar{g}_{AMPA:PY \to TC}$, and $\bar{g}_{AMPA:PY \to TRN}$. Our wake-
like simulation was the only simulation where we included changes to the six synapse types in
the previous sentence; this was done to better align with the wake-like simulations in Figures 14
and 15 of the original paper (Compte et al. 2003). The script used to run the wake-like simulation
is available with the rest of the simulation scripts (Soplata 2023b).
959
960 For anesthetic states, all parameters were as given in the Appendix unless otherwise indicated,
961 such as changes to $\bar{g}_{AMPA:PY \to PY}$ and $\bar{g}_{AMPA:TC \to PY}$ for Tables 1 and 2 and Fig. 6. All parameters are
962 available in the mechanism (Soplata 2023a) and simulation (Soplata 2023b) code.
963
964 *EEG Model*
965
966 We define our simulated EEG as the sum of dendritic voltages of PY cells as these are
967 considered to be the main contributors (Kirschstein and Köhling 2009), bandpass filtered
968 between 0.1-50 Hz. The results are unchanged if we considered PY somatic voltages.
969
970 *Simulations and Reproducibility*
971
972 All of the simulation parameters (Soplata 2023b) and model mechanism code (Soplata 2023a)
973 needed to reproduce the simulations shown in this work are available online on GitHub. All
974 simulations were run using the "dev" branch of the MATLAB simulation toolbox DynaSim
975 (Sherfey et al. 2018) located online. Individual simulations should be reproducible on a modern
976 desktop computer with access to RAM of 32 gigabytes or higher.
977
978 Raw data for Tables 1 and 2 and the bar charts in Figures 4 and 5 were obtained by visual
979 inspection from simulations which are reproducible from the simulation code (Soplata 2023b).
980 The aggregate simulation data is available at Supplemental Data 1.
981
982 *Human Data for Figure 1*
983
984 Human experimental data used in Figure 1 is from a single subject used in (Purdon et al. 2013;
985 Mukamel et al. 2014). Analyses in Fig. 1A follows those in (Purdon et al. 2013; Mukamel et al.
986 2014). EEG traces were band-pass filtered to 0.1-1 Hz, 8-12 Hz and 8-20Hz using a Butterworth

987 filter of order 2. EEG spectrogram in Figure 1B (not previously published) was computed using
988 the multi-tapered method (Babadi and Brown 2014).
989

Acknowledgements:
Current affiliation for Austin Soplata is École Polytechnique Fédérale de Lausanne, Blue Brain
Project, Campus Biotech, Chemin des Mines 9, 1202 Geneva, Switzerland. We thank the
reviewers for their helpful comments on a previous draft of this paper. We would also like to
thank Jason Sherfey, Erik A. Roberts, Emily Stephen, and Caroline Moore-Kochlacs for their
suggestions during the investigation.
996

997 **Grants:**
998 All authors were supported by NIH Grant P01GM118269. Other sources include Guggenheim
999 Fellowship in Applied Mathematics, NIH R01-GM104948, funds from MGH, and NSF DMS-
1000 1042134-5.
1001

Disclosures:
Massachusetts General Hospital has licensed intellectual property for EEG monitoring developed
by Drs. Brown and Purdon to Masimo Corporation. Drs. Purdon and Brown have a financial
interest in PASCALL Systems, Inc., a company developing closed loop physiological
control systems for anesthesiology. Dr. Purdon and Dr. Brown's interests were reviewed and
are managed by Massachusetts General Hospital and Mass General Brigham in accordance with
their conflict-of-interest policies.
1009

1010 **Appendix:**
1011 **Equations for Computational Models (attached as PDF)**
1012

1013 **Supplemental Data 1:**
1014 **Simulation Data and Plots, available at**
1015 **https://doi.org/10.6084/m9.figshare.22228858.v1**
1016

References

Babadi, Behtash, and Emery N. Brown. 2014. "A Review of Multitaper Spectral Analysis."
IEEE Transactions on Biomedical Engineering 61 (5): 1555–64.
https://doi.org/10.1109/TBME.2014.2311996.

Banks, Matthew I., Bryan M. Krause, Christopher M. Endemann, Declan I. Campbell,
Christopher K. Kovach, Mark Eric Dyken, Hiroto Kawasaki, and Kirill V. Nourski. 2020.
"Cortical Functional Connectivity Indexes Arousal State during Sleep and Anesthesia."
NeuroImage 211 (May): 116627. https://doi.org/10.1016/j.neuroimage.2020.116627.

Bastos, André M, Jacob A Donoghue, Scott L Brincat, Meredith Mahnke, Jorge Yanar, Josefina
Correa, Ayan S Waite, et al. 2021. "Neural Effects of Propofol-Induced Unconsciousness
and Its Reversal Using Thalamic Stimulation." Edited by Eunji Cheong and Laura L
Colgin. *ELife* 10 (April): e60824. https://doi.org/10.7554/eLife.60824.

Bazhenov, Maxim, Igor Timofeev, Mircea Steriade, and Terrence J Sejnowski. 2002. "Model of
Thalamocortical Slow-Wave Sleep Oscillations and Transitions to Activated States." *The
Journal of Neuroscience : The Official Journal of the Society for Neuroscience.* 22 (19):
8691–8704. https://doi.org/10.1523/JNEUROSCI.22-19-08691.2002.

Benita, Jose M., Antoni Guillamon, Gustavo Deco, and Maria V. Sanchez-Vives. 2012.
"Synaptic Depression and Slow Oscillatory Activity in a Biophysical Network Model of
the Cerebral Cortex." *Frontiers in Computational Neuroscience* 6.
https://doi.org/10.3389/fncom.2012.00064.

Bharioke, Arjun, Martin Munz, Alexandra Brignall, Georg Kosche, Max Ferdinand Eizinger,
Nicole Ledergerber, Daniel Hillier, et al. 2022. "General Anesthesia Globally
Synchronizes Activity Selectively in Layer 5 Cortical Pyramidal Neurons." *Neuron* 0 (0).
https://doi.org/10.1016/j.neuron.2022.03.032.

Bollimunta, A., J. Mo, C. E. Schroeder, and M. Ding. 2011. "Neuronal Mechanisms and
Attentional Modulation of Corticothalamic Alpha Oscillations." *Journal of Neuroscience*
31 (13): 4935–43. https://doi.org/10.1523/JNEUROSCI.5580-10.2011.

Brown, Emery N., Ralph Lydic, and Nicholas D. Schiff. 2010. "General Anesthesia, Sleep, and
Coma." *The New England Journal of Medicine* 363 (27): 2638–50.
https://doi.org/10.1056/nejmra0808281.

Brown, Emery N., Kara J. Pavone, and Marusa Naranjo. 2018. "Multimodal General Anesthesia:
Theory and Practice." *Anesthesia & Analgesia* 127 (5): 1246–58.
https://doi.org/10.1213/ANE.0000000000003668.

Brown, Emery N., Patrick L. Purdon, and Christa J. Van Dort. 2011. "General Anesthesia and
Altered States of Arousal: A Systems Neuroscience Analysis." *Annual Review of
Neuroscience* 34 (1): 601–28. https://doi.org/10.1146/annurev-neuro-060909-153200.

Buzsáki, György. 2006. *Rhythms of the Brain*. Oxford University Press.
https://doi.org/10.1093/acprof:oso/9780195301069.001.0001.

Cacheaux, L. P., Norbert Topf, Gareth R. Tibbs, Ulrich R. Schaefer, Roberto Levi, Neil L.
Harrison, Geoffrey W. Abbott, and Peter A. Goldstein. 2005. "Impairment of
Hyperpolarization-Activated, Cyclic Nucleotide-Gated Channel Function by the
Intravenous General Anesthetic Propofol." *The Journal of Pharmacology and
Experimental Therapeutics* 315 (2): 517–25. https://doi.org/10.1124/jpet.105.091801.

Cannon, Jonathan, Michelle M. McCarthy, Shane Lee, Jung Lee, Christoph Börgers, Miles A.
Whittington, and Nancy Kopell. 2014. "Neurosystems: Brain Rhythms and Cognitive

Processing." *European Journal of Neuroscience* 39 (5): 705–19.
https://doi.org/10.1111/ejn.12453.

Chen, X. 2005. "HCN Subunit-Specific and CAMP-Modulated Effects of Anesthetics on Neuronal Pacemaker Currents." *Journal of Neuroscience* 25 (24): 5803–14.
https://doi.org/10.1523/JNEUROSCI.1153-05.2005.

Ching, S., A. Cimenser, P. L. Purdon, E. N. Brown, and N. J. Kopell. 2010. "Thalamocortical Model for a Propofol-Induced Alpha-Rhythm Associated with Loss of Consciousness." *Proceedings of the National Academy of Sciences of the United States of America* 107 (52): 22665–70. https://doi.org/10.1073/pnas.1017069108.

Ching, S., P. L. Purdon, S. Vijayan, N. J. Kopell, and E. N. Brown. 2012. "A Neurophysiological-Metabolic Model for Burst Suppression." *Proceedings of the National Academy of Sciences of the United States of America* 109 (8): 3095–3100.
https://doi.org/10.1073/pnas.1121461109.

Cimenser, A., P. L. Purdon, E. T. Pierce, J. L. Walsh, A. F. Salazar-Gomez, P. G. Harrell, C. Tavares-Stoeckel, K. Habeeb, and E. N. Brown. 2011. "Tracking Brain States under General Anesthesia by Using Global Coherence Analysis." *Proceedings of the National Academy of Sciences* 108 (21): 8832–37. https://doi.org/10.1073/pnas.1017041108.

Compte, Albert, Maria V. Sanchez-Vives, David A. McCormick, and Xiao-Jing Wang. 2003. "Cellular and Network Mechanisms of Slow Oscillatory Activity (<1 Hz) and Wave Propagations in a Cortical Network Model." *Journal of Neurophysiology* 89 (5): 2707–25. https://doi.org/10.1152/jn.00845.2002.

Cruikshank, S. J., O. J. Ahmed, T. R. Stevens, S. L. Patrick, A. N. Gonzalez, M. Elmaleh, and B. W. Connors. 2012. "Thalamic Control of Layer 1 Circuits in Prefrontal Cortex." *Journal of Neuroscience* 32 (49): 17813–23. https://doi.org/10.1523/JNEUROSCI.3231-12.2012.

Crunelli, V., A. C. Errington, S. W. Hughes, and T. I. Toth. 2011. "The Thalamic Low-Threshold Ca2+ Potential: A Key Determinant of the Local and Global Dynamics of the Slow (<1 Hz) Sleep Oscillation in Thalamocortical Networks." *Philosophical Transactions of the Royal Society of London. A* 369 (1952): 3820–39.
https://doi.org/10.1098/rsta.2011.0126.

Crunelli, Vincenzo, and Stuart W Hughes. 2010. "The Slow (<1 Hz) Rhythm of Non-REM Sleep: A Dialogue between Three Cardinal Oscillators." *Nature Neuroscience* 13 (1): 9–17. https://doi.org/10.1038/nn.2445.

Dash, Suryadeep, Dawn M. Autio, and Shane R. Crandall. 2022. "State-Dependent Modulation of Activity in Distinct Layer 6 Corticothalamic Neurons in Barrel Cortex of Awake Mice." *The Journal of Neuroscience* 42 (34): 6551–65.
https://doi.org/10.1523/JNEUROSCI.2219-21.2022.

Destexhe, Alain, Thierry Bal, David A. McCormick, and Terrence J. Sejnowski. 1996. "Ionic Mechanisms Underlying Synchronized Oscillations and Propagating Waves in a Model of Ferret Thalamic Slices." *Journal of Neurophysiology* 76 (3): 2049–70.
https://doi.org/10.1152/jn.1996.76.3.2049.

Destexhe, Alain, and T. J. Sejnowski. 2003. "Interactions between Membrane Conductances Underlying Thalamocortical Slow-Wave Oscillations." *Physiological Reviews* 83 (4): 1401–53. https://doi.org/10.1152/physrev.00012.2003.

Eggermann, Emmanuel, and Dirk Feldmeyer. 2009. "Cholinergic Filtering in the Recurrent Excitatory Microcircuit of Cortical Layer 4." *Proceedings of the National Academy of Sciences* 106 (28): 11753–58. https://doi.org/10.1073/pnas.0810062106.

Favero, Morgana, Gladis Varghese, and Manuel A Castro-Alamancos. 2012. "The State of Somatosensory Cortex during Neuromodulation." *Journal of Neurophysiology* 108 (4): 1010–24. https://doi.org/10.1152/jn.00256.2012.

Flores, Francisco J., Katharine E. Hartnack, Amanda B. Fath, Seong-Eun Kim, Matthew A. Wilson, Emery N. Brown, and Patrick L. Purdon. 2017. "Thalamocortical Synchronization during Induction and Emergence from Propofol-Induced Unconsciousness." *Proceedings of the National Academy of Sciences of the United States of America* 114 (32): E6660–68. https://doi.org/10.1073/pnas.1700148114.

Foster, Joshua J, and Edward Awh. 2019. "The Role of Alpha Oscillations in Spatial Attention: Limited Evidence for a Suppression Account." *Current Opinion in Psychology*, Attention & Perception, 29 (October): 34–40. https://doi.org/10.1016/j.copsyc.2018.11.001.

Galarza Vallejo, Ana, Marijn C W Kroes, Enrique Rey, Maria Victoria Acedo, Stephan Moratti, Guillén Fernández, and Bryan A Strange. 2019. "Propofol-Induced Deep Sedation Reduces Emotional Episodic Memory Reconsolidation in Humans." *Science Advances* 5 (3): eaav3801. https://doi.org/10.1126/sciadv.aav3801.

Gaskell, A. L., D. F. Hight, J. Winders, G. Tran, A. Defresne, V. Bonhomme, A. Raz, J. W. Sleigh, and R. D. Sanders. 2017. "Frontal Alpha-Delta EEG Does Not Preclude Volitional Response during Anaesthesia: Prospective Cohort Study of the Isolated Forearm Technique." *British Journal of Anaesthesia*, August. https://doi.org/10.1093/bja/aex170.

Gemignani, Angelo, Danilo Menicucci, Marco Laurino, Andrea Piarulli, Francesca Mastorci, Laura Sebastiani, and Paolo Allegrini. 2015. "Linking Sleep Slow Oscillations with Consciousness Theories: New Vistas on Slow Wave Sleep Unconsciousness." *Archives Italiennes de Biologie* 153 (2–3): 135–43. https://doi.org/10.4449/aib.v153i2-3.4041.

Gil, Ziv, Barry W Connors, and Yael Amitai. 1997. "Differential Regulation of Neocortical Synapses by Neuromodulators and Activity." *Neuron* 19 (3): 679–86. https://doi.org/10.1016/S0896-6273(00)80380-3.

Gutiérrez, Rodrigo, Felipe Maldonado, Jose I. Egaña, and Antonello Penna. 2022. "Electroencephalographic Alpha and Delta Oscillation Dynamics in Response to Increasing Doses of Propofol." *Journal of Neurosurgical Anesthesiology* 34 (1): 79–83. https://doi.org/10.1097/ANA.0000000000000733.

Hage, Travis A, and Lawrence Salkoff. 2012. "Sodium-Activated Potassium Channels Are Functionally Coupled to Persistent Sodium Currents." *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience.* 32 (8): 2714–21. https://doi.org/10.1523/JNEUROSCI.5088-11.2012.

Hill, Sean and Tononi, Giulio. 2004. "Modeling Sleep and Wakefulness in the Thalamocortical System." *Journal of Neurophysiology* 93 (3): 1671–98. https://doi.org/10.1152/jn.00915.2004.

Hsieh, C Y, S J Cruikshank, and R Metherate. 2000. "Differential Modulation of Auditory Thalamocortical and Intracortical Synaptic Transmission by Cholinergic Agonist." *Brain Res.* 880 (1–2): 51–64.

Hudetz, Anthony G., and George A. Mashour. 2016. "Disconnecting Consciousness: Is There a Common Anesthetic End Point?" *Anesthesia & Analgesia* 123 (5): 1228–40. https://doi.org/10.1213/ANE.0000000000001353.

Jiang, Jingyao, Yi Zhao, Jin Liu, Yaoxin Yang, Peng Liang, Han Huang, Yongkang Wu, Yi Kang, Tao Zhu, and Cheng Zhou. 2022. "Signatures of Thalamocortical Alpha

Oscillations and Synchronization With Increased Anesthetic Depths Under Isoflurane." *Frontiers in Pharmacology* 13. https://www.frontiersin.org/articles/10.3389/fphar.2022.887981.

Jones, Stephanie R., David J. Pinto, Tasso J. Kaper, and Nancy Kopell. 2000. "Alpha-Frequency Rhythms Desynchronize over Long Cortical Distances: A Modeling Study." *Journal of Computational Neuroscience* 9 (3): 271–91.

Kikuchi, T, Y Wang, K Sato, and F Okumura. 1998. "In Vivo Effects of Propofol on Acetylcholine Release from the Frontal Cortex, Hippocampus and Striatum Studied by Intracerebral Microdialysis in Freely Moving Rats." *British Journal of Anaesthesia* 80 (5): 644–48. https://doi.org/10.1093/bja/80.5.644.

Kirschstein, Timo, and Rüdiger Köhling. 2009. "What Is the Source of the EEG?" *Clinical EEG and Neuroscience* 40 (3): 146–49. https://doi.org/10.1177/155005940904000305.

Köster, Moritz, and Thomas Gruber. 2022. "Rhythms of Human Attention and Memory: An Embedded Process Perspective." *Frontiers in Human Neuroscience* 16. https://www.frontiersin.org/articles/10.3389/fnhum.2022.905837.

Krishnan, Giri P., Sylvain Chauvette, Isaac Shamie, Sara Soltani, Igor Timofeev, Sydney S. Cash, Eric Halgren, and Maxim Bazhenov. 2016. "Cellular and Neurochemical Basis of Sleep Stages in the Thalamocortical Network." *ELife* 5: e18607. https://doi.org/10.7554/eLife.18607.

Krishnan, Giri P., Oscar C. González, and Maxim Bazhenov. 2018. "Origin of Slow Spontaneous Resting-State Neuronal Fluctuations in Brain Networks." *Proceedings of the National Academy of Sciences* 115 (26): 6858–63. https://doi.org/10.1073/pnas.1715841115.

Krom, Aaron J., Amit Marmelshtein, Hagar Gelbard-Sagiv, Ariel Tankus, Hanna Hayat, Daniel Hayat, Idit Matot, et al. 2020. "Anesthesia-Induced Loss of Consciousness Disrupts Auditory Responses beyond Primary Cortex." *Proceedings of the National Academy of Sciences*, May. https://doi.org/10.1073/pnas.1917251117.

Kruglikov, Illya, and Bernardo Rudy. 2008. "Perisomatic GABA Release and Thalamocortical Integration onto Neocortical Excitatory Cells Are Regulated by Neuromodulators." *Neuron* 58 (6): 911–24. https://doi.org/10.1016/j.neuron.2008.04.024.

Kuo, Min-Ching, and Hans C. Dringenberg. 2008. "Histamine Facilitates in Vivo Thalamocortical Long-Term Potentiation in the Mature Visual Cortex of Anesthetized Rats." *European Journal of Neuroscience* 27 (7): 1731–38. https://doi.org/10.1111/j.1460-9568.2008.06164.x.

Lee, Minji, Robert D. Sanders, Seul-Ki Yeom, Dong-Ok Won, Kwang-Suk Seo, Hyun Jeong Kim, Giulio Tononi, and Seong-Whan Lee. 2017. "Network Properties in Transitions of Consciousness during Propofol-Induced Sedation." *Scientific Reports* 7 (1): 16791. https://doi.org/10.1038/s41598-017-15082-5.

Lewis, Laura D., Veronica S. Weiner, Eran A. Mukamel, Jacob A. Donoghue, Emad N. Eskandar, Joseph R. Madsen, William S. Anderson, et al. 2012. "Rapid Fragmentation of Neuronal Networks at the Onset of Propofol-Induced Unconsciousness." *Proceedings of the National Academy of Sciences of the United States of America* 109 (49): E3377–86. https://doi.org/10.1073/pnas.1210907109.

Lozano-Soldevilla, Diego, Niels ter Huurne, and Robert Oostenveld. 2016. "Neuronal Oscillations with Non-Sinusoidal Morphology Produce Spurious Phase-to-Amplitude Coupling and Directionality." *Frontiers in Computational Neuroscience* 10 (August). https://doi.org/10.3389/fncom.2016.00087.

Luo, Tian-Yuan, Shuang Cai, Zai-Xun Qin, Shao-Cheng Yang, Yue Shu, Cheng-Xi Liu, Yu Zhang, et al. 2020. "Basal Forebrain Cholinergic Activity Modulates Isoflurane and Propofol Anesthesia." *Frontiers in Neuroscience* 14: 1086. https://doi.org/10.3389/fnins.2020.559077.

Lytton, W. W., A. Destexhe, and T. J. Sejnowski. 1996. "Control of Slow Oscillations in the Thalamocortical Neuron: A Computer Model." *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience.* 70 (3): 673–84. https://doi.org/10.1016/S0306-4522(96)83006-5.

Malekmohammadi, Mahsa, Collin M. Price, Andrew E. Hudson, Jasmine A. T. DiCesare, and Nader Pouratian. 2019. "Propofol-Induced Loss of Consciousness Is Associated with a Decrease in Thalamocortical Connectivity in Humans." *Brain*. https://doi.org/10.1093/brain/awz169.

McCarthy, Michelle M., Emery N. Brown, and Nancy Kopell. 2008. "Potential Network Mechanisms Mediating Electroencephalographic Beta Rhythm Changes during Propofol-Induced Paradoxical Excitation." *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience.* 28 (50): 13488–504. https://doi.org/10.1523/JNEUROSCI.3536-08.2008.

McCormick, David A. 1992. "Neurotransmitter Actions in the Thalamus and Cerebral Cortex and Their Role in Neuromodulation of Thalamocortical Activity." *Progress in Neurobiology* 39 (4): 337–88. https://doi.org/10.1016/0301-0082(92)90012-4.

Meuret, P., S. B. Backman, V. Bonhomme, G. Plourde, and P. Fiset. 2000. "Physostigmine Reverses Propofol-Induced Unconsciousness and Attenuation of the Auditory Steady State Response and Bispectral Index in Human Volunteers." *Anesthesiology* 93 (3): 708–17.

Mhuircheartaigh, Róisín Ní, Catherine Warnaby, Richard Rogers, Saad Jbabdi, and Irene Tracey. 2013. "Slow-Wave Activity Saturation and Thalamocortical Isolation during Propofol Anesthesia in Humans." *Science Translational Medicine* 5 (208): 208ra148-208ra148.

Minert, Anne, Mark Baron, and Marshall Devor. 2020. "Reduced Sensitivity to Anesthetic Agents upon Lesioning the Mesopontine Tegmental Anesthesia Area in Rats Depends on Anesthetic Type." *Anesthesiology* 132 (3): 535–50. https://doi.org/10.1097/ALN.0000000000003087.

Minert, Anne, Shai-Lee Yatziv, and Marshall Devor. 2017. "Location of the Mesopontine Neurons Responsible for Maintenance of Anesthetic Loss of Consciousness." *Journal of Neuroscience* 37 (38): 9320–31. https://doi.org/10.1523/JNEUROSCI.0544-17.2017.

Moody, Olivia A., Edlyn R. Zhang, Kathleen F. Vincent, Risako Kato, Eric D. Melonakos, Christa J. Nehs, and Ken Solt. 2021. "The Neural Circuits Underlying General Anesthesia and Sleep." *Anesthesia & Analgesia* 132 (5): 1254–64. https://doi.org/10.1213/ANE.0000000000005361.

Morrison, John H., and Mark G. Baxter. 2012. "The Aging Cortical Synapse: Hallmarks and Implications for Cognitive Decline." *Nature Reviews Neuroscience* 13 (4): 240–50. https://doi.org/10.1038/nrn3200.

Muindi, Fanuel, Jonathan D. Kenny, Norman E. Taylor, Ken Solt, Matthew A. Wilson, Emery N. Brown, and Christa J. Van Dort. 2016. "Electrical Stimulation of the Parabrachial Nucleus Induces Reanimation from Isoflurane General Anesthesia." *Behavioural Brain Research* 306 (June): 20–25. https://doi.org/10.1016/j.bbr.2016.03.021.

Mukamel, E. A., E. Pirondini, B. Babadi, K. F. K. Wong, E. T. Pierce, P. G. Harrell, J. L. Walsh, et al. 2014. "A Transition in Brain State during Propofol-Induced Unconsciousness." *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience.* 34 (3): 839–45. https://doi.org/10.1523/JNEUROSCI.5813-12.2014.

Nemoto, Chiaki, Masahiro Murakawa, Takahiro Hakozaki, Tuyoshi Imaizumi, Tuyoshi Isosu, and Shinju Obara. 2013. "Effects of Dexmedetomidine, Midazolam, and Propofol on Acetylcholine Release in the Rat Cerebral Cortex in Vivo." *Journal of Anesthesia* 27 (5): 771–74. https://doi.org/10.1007/s00540-013-1589-5.

Oldford, E, and M. A Castro-Alamancos. 2003. "Input-Specific Effects of Acetylcholine on Sensory and Intracortical Evoked Responses in the 'Barrel Cortex' in Vivo." *Neuroscience* 117 (3): 769–78. https://doi.org/10.1016/S0306-4522(02)00663-2.

Pal, Dinesh, and George A. Mashour. 2021. "Consciousness, Anesthesia, and Acetylcholine." *Anesthesiology* 134 (4): 515–17. https://doi.org/10.1097/ALN.0000000000003696.

Palva, Satu, and J. Matias Palva. 2007. "New Vistas for α-Frequency Band Oscillations." *Trends in Neurosciences* 30 (4): 150–58. https://doi.org/10.1016/j.tins.2007.02.001.

Penagos, Hector, Carmen Varela, and Matthew A Wilson. 2017. "Oscillations, Neural Computations and Learning during Wake and Sleep." *Current Opinion in Neurobiology* 44 (June): 193–201. https://doi.org/10.1016/j.conb.2017.05.009.

Peters, Alan. 2002. "Structural Changes That Occur during Normal Aging of Primate Cerebral Hemispheres." *Neuroscience and Biobehavioral Reviews* 26 (7): 733–41. https://doi.org/10.1016/S0149-7634(02)00060-X.

Purdon, P. L., K. J. Pavone, O. Akeju, A. C. Smith, A. L. Sampson, J. Lee, D. W. Zhou, K. Solt, and E. N. Brown. 2015. "The Ageing Brain: Age-Dependent Changes in the Electroencephalogram during Propofol and Sevoflurane General Anaesthesia." *British Journal of Anaesthesia* 115 (Suppl 1): i46–57. https://doi.org/10.1093/bja/aev213.

Purdon, P. L., E. T. Pierce, E. A. Mukamel, M. J. Prerau, J. L. Walsh, K. F. K. Wong, A. F. Salazar-Gomez, et al. 2013. "Electroencephalogram Signatures of Loss and Recovery of Consciousness from Propofol." *Proceedings of the National Academy of Sciences of the United States of America* 110 (12): E1142–51. https://doi.org/10.1073/pnas.1221180110.

Quick, Michael W., and Robin A. J. Lester. 2002. "Desensitization of Neuronal Nicotinic Receptors." *Journal of Neurobiology* 53 (4): 457–78. https://doi.org/10.1002/neu.10109.

Redinbaugh, Michelle J., Jessica M. Phillips, Niranjan A. Kambi, Sounak Mohanta, Samantha Andryk, Gaven L. Dooley, Mohsen Afrasiabi, Aeyal Raz, and Yuri B. Saalmann. 2020. "Thalamus Modulates Consciousness via Layer-Specific Control of Cortex." *Neuron* 106 (1): 66-75.e12. https://doi.org/10.1016/j.neuron.2020.01.005.

Sanchez-Vives, Maria V., and David A. McCormick. 2000. "Cellular and Network Mechanisms of Rhythmic Recurrent Activity in Neocortex." *Nature Neuroscience* 3 (10): 1027–34.

Sanchez-Vives, Maria V., Lionel G. Nowak, and David A. McCormick. 2000. "Cellular Mechanisms of Long-Lasting Adaptation in Visual Cortical Neurons In Vitro." *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience.* 20 (11): 4286–99. https://doi.org/10.1523/JNEUROSCI.20-11-04286.2000.

Scheinin, Annalotta, Roosa E. Kallionpää, Duan Li, Minna Kallioinen, Kaike Kaisti, Jaakko Långsjö, Anu Maksimow, et al. 2018. "Differentiating Drug-Related and State-Related Effects of Dexmedetomidine and Propofol on the Electroencephalogram." *Anesthesiology: The Journal of the American Society of Anesthesiologists*, April. https://doi.org/10.1097/ALN.0000000000002192.

Schwindt, P C, W J Spain, and W E Crill. 1989. "Long-Lasting Reduction of Excitability by a Sodium-Dependent Potassium Current in Cat Neocortical Neurons." *Journal of Neurophysiology* 61 (2): 233–44. https://doi.org/10.1152/jn.1989.61.2.233.

Sherfey, Jason S., Austin E. Soplata, Salva Ardid, Erik A. Roberts, David A. Stanley, Benjamin R. Pittman-Polletta, and Nancy J. Kopell. 2018. "DynaSim: A MATLAB Toolbox for Neural Modeling and Simulation." *Frontiers in Neuroinformatics* 12. https://doi.org/10.3389/fninf.2018.00010.

Sherman, S. M. 2014. "The Function of Metabotropic Glutamate Receptors in Thalamus and Cortex." *The Neuroscientist* 20 (2): 136–49. https://doi.org/10.1177/1073858413478490.

Siegel, Markus, Tobias H. Donner, and Andreas K. Engel. 2012. "Spectral Fingerprints of Large-Scale Neuronal Interactions." *Nature Reviews Neuroscience* 13 (2): 121–34. https://doi.org/10.1038/nrn3137.

Sigala, Rodrigo, Sebastian Haufe, Dipanjan Roy, Hubert Dinse, and Petra Ritter. 2014. "The Role of Alpha-Rhythm States in Perceptual Learning: Insights from Experiments and Computational Models." *Frontiers in Computational Neuroscience* 8. https://www.frontiersin.org/articles/10.3389/fncom.2014.00036.

Silva, Laurie R., Yael Amitai, and Barry W. Connors. 1991. "Intrinsic Oscillations of Neocortex Generated by Layer 5 Pyramidal Neurons." *Science* 251 (4992): 432–35. https://doi.org/10.1126/science.1824881.

Soplata, Austin E. 2023a. "DynaSim Mechanism Files for Simulating Thalamically-Extended (Benita et al., 2012) Slow Wave Model." MATLAB. https://doi.org/10.5281/zenodo.7725210.

———. 2023b. "Simulation and Plotting Scripts for (Soplata et al., 2023)." MATLAB. https://doi.org/10.5281/zenodo.7725212.

Soplata, Austin E., Michelle M. McCarthy, Jason Sherfey, Shane Lee, Patrick L. Purdon, Emery N. Brown, and Nancy Kopell. 2017. "Thalamocortical Control of Propofol Phase-Amplitude Coupling." *PLoS Computational Biology* 13 (12): e1005879. https://doi.org/10.1371/journal.pcbi.1005879.

Stephen, Emily P., Gladia C. Hotan, Eric T. Pierce, P. Grace Harrell, John L. Walsh, Emery N. Brown, and Patrick L. Purdon. 2020. "Broadband Slow-Wave Modulation in Posterior and Anterior Cortex Tracks Distinct States of Propofol-Induced Unconsciousness." *Scientific Reports* 10 (1): 13701. https://doi.org/10.1038/s41598-020-68756-y.

Stickgold, Robert. 2005. "Sleep-Dependent Memory Consolidation." *Nature* 437 (7063): 1272–78. https://doi.org/10.1038/nature04286.

Suzuki, Mototaka, and Matthew E. Larkum. 2020. "General Anesthesia Decouples Cortical Pyramidal Neurons." *Cell* 180 (4): 666-676.e13. https://doi.org/10.1016/j.cell.2020.01.024.

Taxidis, Jiannis, Kenji Mizuseki, Robert Mason, and Markus R. Owen. 2013. "Influence of Slow Oscillation on Hippocampal Activity and Ripples through Cortico-Hippocampal Synaptic Interactions, Analyzed by a Cortical-CA3-CA1 Network Model." *Frontiers in Computational Neuroscience* 7. https://doi.org/10.3389/fncom.2013.00003.

Timofeev, I., F. Grenier, M. Bazhenov, T. J. Sejnowski, and M. Steriade. 2000. "Origin of Slow Cortical Oscillations in Deafferented Cortical Slabs." *Cerebral Cortex (New York, N.Y.: 1991)* 10 (12): 1185–99. https://doi.org/10.1093/cercor/10.12.1185.

Tinker, John H., Frank W. Sharbrough, and John D. Michenfelder. 1977. "Anterior Shift of the Dominant EEG Rhythm during Anesthesia in the Java Monkey: Correlation with Anesthetic Potency." *Anesthesiology.*

Traub, Roger D, Diego Contreras, Mark O Cunningham, Hilary Murray, Fiona E N LeBeau, Anita Roopun, Andrea Bibbig, W Bryan Wilent, Michael J Higley, and Miles A Whittington. 2005. "Single-Column Thalamocortical Network Model Exhibiting Gamma Oscillations, Sleep Spindles, and Epileptogenic Bursts." *J. Neurophysiol.* 93 (4): 2194–2232.

Ujma, Péter P., Orsolya Szalárdy, Dániel Fabó, Loránd Erőss, and Róbert Bódizs. 2022. "Thalamic Activity during Scalp Slow Waves in Humans." *NeuroImage*, May, 119325. https://doi.org/10.1016/j.neuroimage.2022.119325.

Van Diepen, Rosanne M., John J Foxe, and Ali Mazaheri. 2019. "The Functional Role of Alpha-Band Activity in Attentional Processing: The Current Zeitgeist and Future Outlook." *Current Opinion in Psychology*, Attention & Perception, 29 (October): 229–38. https://doi.org/10.1016/j.copsyc.2019.03.015.

Vijayan, S., S. Ching, P. L. Purdon, E. N. Brown, and N. J. Kopell. 2013. "Thalamocortical Mechanisms for the Anteriorization of Alpha Rhythms during Propofol-Induced Unconsciousness." *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience.* 33 (27): 11070–75. https://doi.org/10.1523/JNEUROSCI.5670-12.2013.

Vijayan, S., and N. J. Kopell. 2012. "Thalamic Model of Awake Alpha Oscillations and Implications for Stimulus Processing." *Proceedings of the National Academy of Sciences* 109 (45): 18553–58. https://doi.org/10.1073/pnas.1215385109.

Vlasov, Ksenia, JunZhu Pei, Christa J. Nehs, Jennifer A. Guidera, Edlyn R. Zhang, Jonathan D. Kenny, Timothy T. Houle, Gary J. Brenner, Norman E. Taylor, and Ken Solt. 2021. "Activation of GABAergic Neurons in the Rostromedial Tegmental Nucleus and Other Brainstem Regions Promotes Sedation and Facilitates Sevoflurane Anesthesia in Mice." *Anesthesia & Analgesia* 132 (4): e50. https://doi.org/10.1213/ANE.0000000000005387.

Wang, Dengyu, Witold J. Lipski, Alan Bush, Anna Chrabaszcz, Christina A. Dastolfo-Hromack, Michael Dickey, Julie A. Fiez, and R. Mark Richardson. 2022. "Lateralized and Region-Specific Thalamic Processing of Lexical Status during Reading Aloud." *Journal of Neuroscience* 42 (15): 3228–40. https://doi.org/10.1523/JNEUROSCI.1332-21.2022.

Weiner, Veronica S., David W. Zhou, Pegah Kahali, Emily P. Stephen, Robert A. Peterfreund, Linda S. Aglio, Michele D. Szabo, et al. 2023. "Propofol Disrupts Alpha Dynamics in Functionally Distinct Thalamocortical Networks during Loss of Consciousness." *Proceedings of the National Academy of Sciences* 120 (11): e2207831120. https://doi.org/10.1073/pnas.2207831120.

Ying, Shui-Wang, Syed Y. Abbas, Neil L. Harrison, and Peter A. Goldstein. 2006. "Propofol Block of Ih Contributes to the Suppression of Neuronal Excitability and Rhythmic Burst Firing in Thalamocortical Neurons: Propofol Inhibition of Thalamic Ih." *The European Journal of Neuroscience* 23 (2): 465–80. https://doi.org/10.1111/j.1460-9568.2005.04587.x.

Appendix: Equations for Computational Models

Austin Soplata

March 26, 2023

1 General Notes

Units of maximal conductances and currents for all equations below are in $\frac{mS}{cm^2}$ and $\frac{\mu A}{cm^2}$, respectively, based on the original formulation of [Hodgkin and Huxley, 1952]. Capacitance C_m for all cells was $1\frac{\mu F}{cm^2}$. Thalamic circuitry is based on [Destexhe et al., 1993], [Destexhe et al., 1996], and [Ching et al., 2010] and is mostly identical to [Soplata et al., 2017]. Cortical circuitry is based on a from-scratch implementation of [Compte et al., 2003] and [Benita et al., 2012]. Thalamocortical and corticothalamic synaptic equations were derived from the same AMPAergic synapse equations as in [Soplata et al., 2017]. Occasionally, multiplication dots will be used ($\cdot$) to help legibility. All simulations were run for 30 seconds of simulation time (unless otherwise indicated) and solved using Euler integration, with a time resolution (dt) of 0.01 ms.

2 (PYdr) Cortical Pyramidal Dendritic Cell Compartment Equations

2.1 Voltage / Membrane Potential

$$\dot{V}_{PYdr} = \frac{1}{C_m}(I_{app} - I_{HVA} - I_{K(Ca)} - I_{K(Na)} +$$
$$-I_{NaP} - I_{AR} - I_{Leak} - I_{COM} - I_{syn})$$

$$I_{app} = 0.0\frac{\mu A}{cm^2}$$

2.2 (HVA) High-Voltage-Activated Ca Channel

$$I_{HVA} = \bar{g}_{HVA}m_\infty^2(V - E_{HVA})$$

Parameters and Functions:

$$\bar{g}_{HVA} = 0.43\frac{mS}{cm^2}$$
$$E_{HVA} = 120mV$$
$$m_\infty = 1/(1 + exp(\frac{-(V+20)}{9}))$$

2.3 (K(Ca)) Slow Calcium-activated K Channel

$$I_{K(Ca)} = \bar{g}_{K(Ca)}\frac{[Ca_i]}{[Ca_i] + K_D}(V - E_{K(Ca)})$$

Parameters:

$$\bar{g}_{K(Ca)} = 0.57\frac{mS}{cm^2}$$
$$E_{K(Ca)} = -100mV$$
$$K_D = 30\mu M$$
$$\alpha_{Ca} = 0.005 \cdot 1000\frac{\mu M}{\mu A \cdot ms}$$
$$A_{DR} = 0.00035cm^2$$
$$[Ca_i]_{initial} = 0.001\mu M$$

State Variable Equations: $[Ca_i]$

$$[\dot{Ca_i}] = max(-\alpha_{Ca} \cdot A_{DR} \cdot I_{HVA}, 0) - [Ca_i]$$

2.4 (K(Na)) Na-activated K Channel

See the section "Non-synaptic Connection Equations".

2.5 (NaP) Persistent Na Channel

$$I_{NaP} = \bar{g}_{NaP}m_\infty^3(V - E_{NaP})$$

Parameters and Functions:

$$\bar{g}_{NaP} = 0.0686\frac{mS}{cm^2}$$
$$E_{NaP} = 55mV$$
$$m_\infty = 1/(1 + exp(\frac{-(V+55.7)}{7.7}))$$

2.6 (AR) Inwardly Rectifying K Channel

$$I_{AR} = \bar{g}_{AR}h_\infty(V - E_{AR})$$

Parameters and Functions:

$$\bar{g}_{AR} = 0.0257\frac{mS}{cm^2}$$
$$E_{AR} = -100mV$$
$$h_\infty = 1/(1 + exp(\frac{V+75}{4}))$$

2.7 (Leak) Leak Channel

$$I_{Leak} = \bar{g}_{Leak}(V - E_{Leak})$$

Parameters:

$$\bar{g}_{Leak} = 0.005\frac{mS}{cm^2}$$
$$E_{Leak} = -60.95mV$$

3 (PYso) Cortical Pyramidal Axo-somatic Cell Compartment Equations

3.1 Voltage / Membrane Potential

$$\dot{V}_{PYso} = \frac{1}{C_m}(-I_{Na} - I_K - I_A - I_{KS}+$$
$$-I_{Leak} - I_{COM} - I_{syn})$$

3.2 (Na) Na Channel

$$I_{Na} = \bar{g}_{Na}m_\infty^3 h(V - E_{Na})$$

Parameters:

$$\bar{g}_{Na} = 50\frac{mS}{cm^2}$$

$$E_{Na} = 55mV$$

$$\phi = 4$$

State Variable Equations: m, h

$$m_\infty = \frac{\alpha_m}{\alpha_m + \beta_m}$$
$$\alpha_m = \frac{0.1(V+33)}{(1 - exp(\frac{-(V+33)}{10}))}$$
$$\beta_m = 4exp(\frac{-(V+53.7)}{12})$$
$$\dot{h} = \phi \cdot (\alpha_h \cdot (1 - h)) - \beta_h \cdot h$$
$$\alpha_h = 0.07 \cdot exp(\frac{-(V+50)}{10})$$
$$\beta_h = 1/(1 + exp(\frac{-(V+20)}{10}))$$

3.3 (K) K Channel

$$I_K = \bar{g}_K n^4 (V - E_K)$$

Parameters:

$$\bar{g}_K = 10.5\frac{mS}{cm^2}$$

$$E_K = -100mV$$

$$\phi = 4$$

State Variable Equations: n

$$\dot{n} = \phi \cdot (\alpha_n \cdot (1 - n)) - \beta_n \cdot n$$
$$\alpha_n = \frac{0.01(V+34)}{(1 - exp(\frac{-(V+34)}{10}))}$$
$$\beta_n = 0.125exp(\frac{-(V+44)}{25})$$

3.4 (A) Fast A-type K Channel

$$I_A = \bar{g}_A m_\infty^3 h(V - E_A)$$

Parameters:

$$\bar{g}_A = 1\frac{mS}{cm^2}$$

$$E_A = -100mV$$

$$\phi = 1$$

$$\tau_h = 15ms$$

State Variable Equations: h

$$\dot{h} = \phi \cdot \frac{h_\infty - h}{\tau_h}$$
$$\alpha_h = 1/(1 + exp(\frac{-(V+50)}{20}))$$
$$\beta_h = 1/(1 + exp(\frac{(V+80)}{6}))$$

3.5 (KS) Non-inactivating K Channel

$$I_{KS} = \bar{g}_{KS}m^3(V - E_{KS})$$

Parameters:

$$\bar{g}_{KS} = 0.576\frac{mS}{cm^2}$$

$$E_{KS} = -100mV$$

$$\phi = 1$$

State Variable Equations: m

$$\dot{m} = \phi \cdot \frac{m_\infty - m}{\tau_m}$$
$$m_\infty = 1/(1 + exp(\frac{-(V+34)}{6.5}))$$
$$\tau_m = 8/\left(exp(\frac{-(V+55)}{30}) + exp(\frac{(V+55)}{30})\right)$$

3.6 (Leak) Leak Channel

$$I_{Leak} = \bar{g}_{Leak}(V - E_{Leak})$$

Parameters:

$$\bar{g}_{Leak} = 0.0667\frac{mS}{cm^2}$$

$$E_{Leak} = -60.95mV$$

4 (IN) Cortical Interneuron Cell Equations

4.1 Voltage / Membrane Potential

$$\dot{V}_{IN} = \frac{1}{C_m}(-I_{Na} - I_K - I_{Leak} - I_{syn})$$

4.2 (Na) Na Channel

$$I_{Na} = \bar{g}_{Na}m_\infty^3 h(V - E_{Na})$$

5.2 (Na) Na Channel

$$I_{Na} = \bar{g}_{Na} m^3 h (V_T - E_{Na})$$

Parameters:

$$V_T = V + 40mV$$

$$\bar{g}_{Na} = 90 \frac{mS}{cm^2}$$

$$E_{Na} = 50mV$$

State Variable Equations: m, h

$$\dot{m} = \alpha_m \cdot (1 - m) - \beta_m \cdot m$$

$$\alpha_m = \frac{0.32 \cdot (13 - V_T)}{exp(\frac{13 - V_T}{4}) - 1}$$

$$\beta_m = \frac{0.28 \cdot (V_T - 40)}{exp(\frac{V_T - 40}{5}) - 1}$$

$$\dot{h} = \alpha_h \cdot (1 - h) - \beta_h \cdot h$$

$$\alpha_h = 0.128 \cdot exp(\frac{17 - V_T}{18})$$

$$\beta_h = \frac{4}{exp(\frac{40 - V_T}{5}) + 1}$$

Parameters:

$$\bar{g}_{Na} = 35 \frac{mS}{cm^2}$$

$$E_{Na} = 55mV$$

State Variable Equations: m, h

$$m_\infty = \frac{\alpha_m}{\alpha_m + \beta_m}$$

$$\alpha_m = \frac{0.5(V+35)}{(1 - exp(\frac{-(V+35)}{10}))}$$

$$\beta_m = 20 exp(\frac{-(V+60)}{18})$$

$$\dot{h} = (\alpha_h \cdot (1 - h)) - \beta_h \cdot h$$

$$\alpha_h = 0.35 \cdot exp(\frac{-(V+58)}{20})$$

$$\beta_h = 5/(1 + exp(\frac{-(V+28)}{10}))$$

4.3 (K) K Channel

$$I_K = \bar{g}_K n^4 (V - E_K)$$

Parameters:

$$\bar{g}_K = 9 \frac{mS}{cm^2}$$

$$E_K = -90mV$$

State Variable Equations: n

$$\dot{n} = (\alpha_n \cdot (1 - n)) - \beta_n \cdot n$$

$$\alpha_n = \frac{0.05(V+34)}{(1 - exp(\frac{-(V+34)}{10}))}$$

$$\beta_n = 0.625 exp(\frac{-(V+44)}{80})$$

5.3 (K) K Channel

$$I_K = \bar{g}_K n^4 (V_T - E_K)$$

Parameters:

$$V_T = V + 25mV$$

$$\bar{g}_K = 10 \frac{mS}{cm^2}$$

$$E_K = -100mV$$

State Variable Equations: n

$$\dot{n} = \alpha_n \cdot (1 - n) - \beta_n \cdot n$$

$$\alpha_n = \frac{0.032(15 - V_T)}{exp(\frac{15 - V_T}{5}) - 1}$$

$$\beta_n = 0.5 exp(\frac{10 - V_T}{40})$$

4.4 (Leak) Leak Channel

$$I_{Leak} = \bar{g}_{Leak}(V - E_{Leak})$$

Parameters:

$$\bar{g}_{Leak} = 0.1025 \frac{mS}{cm^2}$$

$$E_{Leak} = -63.8mV$$

5.4 (T) T-type Calcium Channel

$$I_T = \bar{g}_T m_\infty^2 h (V_T - E_T)$$

Parameters:

$$V_T = V + 2mV$$

$$\bar{g}_T = 2 \frac{mS}{cm^2}$$

$$E_T = 1000 \frac{8.31441 * 309.15}{2 * 96486} ln \frac{2}{[Ca]_i} mV$$

$$\phi_h = 3.73$$

$$A = \frac{10}{2 \cdot 96489} \frac{mM \cdot cm^2}{ms \cdot \mu A}$$

$$[Ca_i]_{initial} = 0.000001mM$$

5 (TC) Thalamocortical Cell Equations

5.1 Voltage / Membrane Potential

$$\dot{V}_{TC} = \frac{1}{C_m}(-I_{Na} - I_K - I_T - I_H +$$

$$- I_{Leak} - I_{KLeak} - I_{syn})$$

State Variable Equations: m

$$m_\infty = \frac{1}{1+exp(\frac{-(V_T+57)}{6.2})}$$

State Variable Equations: h

$$\dot{h} = \frac{h_\infty - h}{\tau_h}$$

$$h_\infty = \frac{1}{1+exp(\frac{V_T+81}{4})}$$

$$\tau_h = \left(30.8 + \frac{211.4+exp(\frac{V_T+113.2}{5})}{1+exp(\frac{V_T+84}{3.2})}\right)/\phi_h$$

State Variable Equations: $[Ca]_i$

$$[\dot{Ca}]_i = max\left(\frac{-10 \cdot I_T}{2 \cdot 96489}, 0\right) + \frac{0.00024 - [Ca]_i}{5}$$

5.5 (H) Hyperpolarization-activated Cation Channel

$$I_H = \bar{g}_H(O_U + 2O_L)(V - E_H)$$

Note: This is the more complex [Destexhe et al., 1996] formulation of the H-current, not that of [Destexhe et al., 1993]. O_U is the proportion of unlocked channels, O_L is the proportion of locked-open channels, and P_1 is the proportion of utilized substrate for channel locking.

Parameters:

$\bar{g}_H = 0.005\frac{mS}{cm^2}$ (depending on dose)

$E_H = -40mV$

$k_2 = 0.0004$

$k_4 = 0.001$

$[Ca]_{crit} = 0.002$ (mM of Calcium)

$n_{Ca} = 4$ (number binding sites of Ca on protein)

$n_{exp} = 1$ (number binding sites on H-channel)

$\bar{g}_H$ changes depending on the simulation under investigation. See Propofol Effects section for more detail.

State Variable Equations: O_U, P_1, O_L

$$\dot{O}_U = \alpha_h(1 - O_U - O_L) - \beta_h O_U$$

$$\dot{P}_1 = k_2(\frac{[Ca_i]}{[Ca_i]_{crit}})^{n_{Ca}}(1 - P_1) - k_2 P_1$$

- $[Ca]_i$ is a state variable determined by the T-current.

$$\dot{O}_L = k_4(\frac{P_1}{0.007})^{n_{exp}} O_U - k_4 O_L$$

$$\alpha_h = \frac{h_\infty}{\tau_s}$$

$$\beta_h = \frac{1 - h_\infty}{\tau_s}$$

$$h_\infty = 1/(1 + exp(\frac{V+75}{5.5}))$$

$$\tau_s = 20 + \frac{1000}{exp(\frac{V+71.5}{14.2}) + exp(\frac{-(V+89)}{11.6})}$$

5.6 (Leak) Leak Current

$$I_{Leak} = \bar{g}_{Leak}(V - E_{Leak})$$

Parameters:

$\bar{g}_{Leak} = 0.01\frac{mS}{cm^2}$

$E_{Leak} = -70mV$

5.7 (KLeak) K Leak Current

$$I_{KLeak} = \bar{g}_{KLeak}(V - E_{KLeak})$$

Parameters:

$\bar{g}_{KLeak} = 0.0172\frac{mS}{cm^2}$

$E_{KLeak} = -100mV$

6 (TRN) Thalamic Reticular Nucleus Cell Equations

6.1 Voltage / Membrane Potential

$$\dot{V}_{TRN} = \frac{1}{C_m}(-I_{Na} - I_K - I_T - I_{Leak}+$$
$$-I_{KLeak} - I_{syn})$$

6.2 (Na) Na Channel

$$I_{Na} = \bar{g}_{Na}m^3 h(V_T - E_{Na})$$

Parameters:

$V_T = V + 55mV$

$\bar{g}_{Na} = 200\frac{mS}{cm^2}$

$E_{Na} = 50mV$

State Variable Equations: m, h

$$\dot{m} = \alpha_m \cdot (1 - m) - \beta_m \cdot m$$

$$\alpha_m = \frac{0.32 \cdot (13 - V_T)}{exp(\frac{13 - V_T}{4}) - 1}$$

$$\beta_m = \frac{0.28 \cdot (V_T - 40)}{exp(\frac{V_T - 40}{5}) - 1}$$

$$\dot{h} = \alpha_h \cdot (1 - h) - \beta_h \cdot h$$

$$\alpha_h = 0.128 \cdot exp(\frac{17 - V_T}{18})$$

$$\beta_h = \frac{4}{exp(\frac{40 - V_T}{5}) + 1}$$

6.3 (K) K Channel

$$I_K = \bar{g}_K n^4(V_T - E_K)$$

$$V_T = V + 55mV$$

$$\bar{g}_K = 20\frac{mS}{cm^2}$$

$$E_K = -100mV$$

$$\dot{n} = \alpha_n \cdot (1 - n) - \beta_n \cdot n$$

$$\alpha_n = \frac{0.032(15 - V_T)}{exp(\frac{15 - V_T}{5}) - 1}$$

$$\beta_n = 0.5exp(\frac{10 - V_T}{40})$$

6.4 (T) T-type Calcium Channel

$$I_T = \bar{g}_T m^2 h(V_T - E_T)$$

Parameters:

$$V_T = V + 4mV$$

$$\bar{g}_T = 3\frac{mS}{cm^2}$$

$$E_T = 120mV$$

$$\phi_m = 6.81$$

$$\phi_h = 3.73$$

State Variable Equations: m

$$\dot{m} = \frac{m_\infty - m}{\tau_m}$$

$$m_\infty = 1/(1 + exp(\frac{-(V_T + 50)}{7.4}))$$

$$\tau_m = \frac{1}{\phi_m}\left(3 + \frac{1}{exp(\frac{V_T + 25}{10}) + exp(\frac{-(V_T + 100)}{15})}\right)$$

State Variable Equations: h

$$\dot{h} = \frac{h_\infty - h}{\tau_h}$$

$$h_\infty = \frac{1}{1 + exp(\frac{V_T + 78}{5})}$$

$$\tau_h = \frac{1}{\phi_h}\left(85 + \frac{1}{exp(\frac{V_T + 46}{4}) + exp(\frac{-(V_T + 405)}{50})}\right)$$

6.5 (Leak) Leak Currents

$$I_{Leak} = \bar{g}_{Leak}(V - E_{Leak})$$

Parameters:

$$\bar{g}_{Leak} = 0.05\frac{mS}{cm^2}$$

$$E_{Leak} = -90mV$$

7 Non-synaptic Connection Equations

7.1 Direct Compartmental Connections

Each PYdr compartment has a special connection to and from a single PYso compartment called I_{COM}, meant to simulate voltage fluctuations between the axo-soma and dendrite of a single cell. Note that these conductances are not exact inverses of each other due to the difference in size between the two compartments. These two currents are calculated using the following:

$$I_{COM|PYdr\to PYso} = \bar{g}_{COM|PYdr\to PYso}(V_{PYdr} - V_{PYso})$$

$$\bar{g}_{COM|PYdr\to PYso} = 11.667\frac{mS}{cm^2}$$

$$I_{COM|PYso\to PYdr} = \bar{g}_{COM|PYso\to PYdr}(V_{PYso} - V_{PYdr})$$

$$\bar{g}_{COM|PYso\to PYdr} = 5\frac{mS}{cm^2}$$

7.2 (K(Na)) Intercompartmental Na-activated K Channel

Additionally, there is a special and very important current, $I_{K(Na)}$, which requires information on the state of the Na present in directly-connected PYso and PYdr compartments. This changes depending on ACh dose. Practically speaking, this can be modeled somewhat easily by programming it as if it is a synaptic current. The equations follow:

$$I_{K(Na)} = \bar{g}_{K(Na)}\frac{0.37}{1 + (\frac{38.7}{[Na_i]})^{3.5}}(V_{PYso} - E_{K(Na)})$$

$$\bar{g}_{K(Na)} = 1.33\frac{mS}{cm^2}$$

$$E_{K(Na)} = -100mV$$

$$\alpha = 0.01 \cdot 1000\frac{mM}{\mu A \cdot ms}$$

$$R_{pump} = 0.018\frac{mM}{ms}$$

$$[Na_{eq}] = 9.5mM$$

$$[\dot{Na_i}] = -\alpha(0.00015 \cdot I_{Na|PYso} + 0.00035 \cdot I_{NaP|PYdr}) +$$

$$-R_{pump}(\frac{[Na_i]^3}{[Na_i]^3 + 15^3} - \frac{[Na_{eq}]^3}{[Na_{eq}]^3 + 15^3})$$

8 Synaptic Equations

8.1 Channel Equations

The electrical current equations for each class of synapse ($AMPA$, $AMPA_D$, $NMDA$, $GABA_A$, and $GABA_B$) were the same within their class, with the sole exception of different connections having different maximal conductances. All conductances of individual synapses were divided by a normalizing factor of how many connections each target cell received. Note that $AMPA_D$ is referred to as $AMPA$ in the main manuscript; its different name is only relevant here for explaining that it contains depression mechanisms in the implementation.

<u>*AMPA*:</u>

$I_{AMPA} = -\bar{g}_{AMPA} \cdot s_{AMPA} \cdot (V_{target} - E_{AMPA})$

$E_{AMPA} = 1mV$

$\tau_{AMPA} = 2ms$

$\dot{s}_{AMPA} = 5(1 + tanh\frac{V_{source}}{4})(1 - s_{AMPA}) - \frac{s_{AMPA}}{\tau_{AMPA}}$

<u>*AMPA_D*:</u>

AMPA synapse, with synaptic depression

$I_{AMPA_D} = -\bar{g}_{AMPA_D} \cdot R \cdot s_{AMPA_D} \cdot (V_{target} - E_{AMPA})$

$E_{AMPA} = 0mV$

$\alpha = 3.48$

$\tau_s = 2ms$

$\tau_R = 400ms$

$DF = \text{Depression Factor} = 0.9$

$SD = \text{Spike-detection function} = \text{if (time}$

since last spike$) < 2 \cdot dt$, where dt is

the time resolution of the ODE solver

$\dot{s}_{AMPA_D} = \alpha/(1 + exp\frac{-(V_{source}-20)}{2}) - \frac{s_{AMPA_D}}{\tau_s}$

$\dot{R} = \frac{1-R}{\tau_R} + SD \cdot (\frac{-(1-R)}{\tau_R} + \frac{1}{dt}(DF \cdot R - R))$

<u>*NMDA*:</u>

$I_{NMDA} = -\bar{g}_{NMDA} \cdot R \cdot s_{NMDA} \cdot (V_{target} - E_{NMDA})$

$E_{AMPA} = 0mV$

$\tau_{AMPA} = 2ms$

$\alpha_s = 0.5$

$\tau_s = 100ms$

$\alpha_x = 3.48$

$\tau_x = 2ms$

$\tau_R = 400ms$

$DF = \text{Depression Factor} = 0.9$

$SD = \text{Spike-detection function} = \text{if (time}$

since last spike$) < 2 \cdot dt$, where dt is

the time resolution of the ODE solver

$\dot{s}_{NMDA} = \alpha_s \cdot x_{NMDA} \cdot (1 - s_{NMDA}) - \frac{s_{NMDA}}{\tau_s}$

$\dot{x}_{NMDA} = \alpha_x/(1 + exp(\frac{-(V_{source}-20)}{2})) - \frac{x_{NMDA}}{\tau_x}$

$\dot{R} = \frac{1-R}{\tau_R} + SD \cdot (\frac{-(1-R)}{\tau_R} + \frac{1}{dt}(DF \cdot R - R))$

<u>*GABA_A*:</u>

$I_{GABA_A} = -PM \cdot \bar{g}_{GABA_A} \cdot s_{GABA_A} \cdot (V_{target} - E_{GABA_A})$

$E_{GABA_A} = -80mV$

$\tau_{GABA_A} = PM \cdot 5ms$

$PM = 1$ propofol multiplier, see Propofol Effects section.

Note that PM multiplies both $\bar{g}_{GABA_A}$ and τ_{GABA_A}.

$\dot{s}_{GABA_A} = 2(1 + tanh\frac{V_{source}}{4})(1 - s_{GABA_A}) - \frac{s_{GABA_A}}{PM \cdot \tau_{GABA_A}}$

<u>*GABA_B*:</u>

$I_{GABA_B} = -\bar{g}_{GABA_B} \cdot \frac{s_{GABA_B}^4}{s_{GABA_B}^4 + 100} \cdot (V_{target} - E_{GABA_A})$

$E_{GABA_B} = -95mV$

$\dot{s}_{GABA_B} = 0.18 \cdot r_{GABA_B} - 0.034 \cdot s_{GABA_B}$

$\dot{r}_{GABA_B} = 0.5(2(1 + tanh\frac{V_{source}}{4}))(1 - r_{GABA_B}) +$

$-0.0012 \cdot r_{GABA_B}$

8.2 Synaptic connectivity and conductances

See the Methods section for general description of connectivity. For the "nearest neighbors" connection algorithm, see the code file "models/netconNearestNeighbors.m". Additionally, incoming synapse maximal conductances are divided by a Normalization Factor (NF). This NF ensures that the total maximal conductance for a particular synaptic type into a target cell is balanced across the number of incoming cell synapses. NF is defined as:

$$NF = min(\frac{1 + 2 \cdot radius}{N_{target}/N_{source}}, N_{source})$$

Where N_{source} and N_{target} are the number of cells in the source and target populations, respectively.

The following table lists all synaptic connection radii and synaptic maximal conductances (before NF has been applied). $PY so \to PY dr$ synaptic connections only connect to $2 \cdot radius$ target cells so as not to synapse onto their corresponding compartment. All maximal conductances are giving in units of $\frac{mS}{cm^2}$. For all synapses modulated by propofol, see the following section Propofol Effects.

Connection	Synapse Class	$\bar{g}(\frac{mS}{cm^2})$	radius
$PYso \rightarrow PYdr$	$AMPA_D$	0.005	10
	$NMDA$	0.00257	10
$PYso \rightarrow IN$	$AMPA_D$	1.0	10
	$NMDA$	0.0025	10
$IN \rightarrow PYso$	$GABA_A$	0.1	10
$IN \rightarrow IN$	$GABA_A$	0.000825	10
$PYso \rightarrow TC$	$AMPA$	0.4	10
$PYso \rightarrow TRN$	$AMPA$	0.2	10
$TC \rightarrow PYdr$	$AMPA$	0.005	10
$TC \rightarrow IN$	$AMPA$	0.1	10
$TC \rightarrow TRN$	$AMPA$	0.4	10 (all-to-all)
$TRN \rightarrow TC$	$GABA_A$	0.1	10 (all-to-all)
$TRN \rightarrow TC$	$GABA_B$	0.001	10 (all-to-all)
$TRN \rightarrow TRN$	$GABA_A$	0.1	10 (all-to-all)

9 Propofol Effects

For the "wakelike" non-propofol state, see the Methods. For the anesthetic simulations, the following parameters were set. Note that PM or "Propofol Multiplier" multiplies both $\bar{g}_{GABA_A}$ and τ_{GABA_A}.

$$IN \rightarrow PYso: \quad PM = 3$$
$$IN \rightarrow IN: \quad PM = 3$$
$$TRN \rightarrow TC: \quad PM = 3$$
$$TRN \rightarrow TRN: \quad PM = 3$$
$$TC: \quad \bar{g}_H = 0.005\frac{mS}{cm^2}$$
$$PYdr: \quad \bar{g}_{K(Na)} = 1.33\frac{mS}{cm^2}$$

10 Reproducibility and Code

All simulations were run using the MATLAB-based DynaSim software package [Sherfey et al., 2018] on the "dev" branch, using MATLAB 2021a. The individual mechanism files for use with DynaSim are available online [Soplata, 2023a], and so are all the runscripts needed to reproduce simulations of the paper [Soplata, 2023b].

References

Jose M. Benita, Antoni Guillamon, Gustavo Deco, and Maria V. Sanchez-Vives. Synaptic depression and slow oscillatory activity in a biophysical network model of the cerebral cortex. *Frontiers in Computational Neuroscience*, 6, 2012. ISSN 1662-5188. doi: 10.3389/fncom.2012.00064. URL http://journal.frontiersin.org/article/10.3389/fncom.2012.00064/abstract.

S. Ching, A. Cimenser, P. L. Purdon, E. N. Brown, and N. J. Kopell. Thalamocortical model for a propofol-induced alpha-rhythm associated with loss of consciousness. *Proceedings of the National Academy of Sciences of the United States of America*, 107(52):22665–22670, December 2010. ISSN 0027-8424, 1091-6490. doi: 10.1073/pnas.1017069108. URL http://www.pnas.org/cgi/doi/10.1073/pnas.1017069108.

Albert Compte, Maria V. Sanchez-Vives, David A. McCormick, and Xiao-Jing Wang. Cellular and Network Mechanisms of Slow Oscillatory Activity (<1 Hz) and Wave Propagations in a Cortical Network Model. *Journal of Neurophysiology*, 89(5):2707–2725, May 2003. ISSN 0022-3077. doi: 10.1152/jn.00845.2002. URL https://www.physiology.org/doi/full/10.1152/jn.00845.2002.

A Destexhe, A Babloyantz, and T J Sejnowski. Ionic mechanisms for intrinsic slow oscillations in thalamic relay neurons. *Biophysical Journal*, 65(4):1538–1552, October 1993. ISSN 0006-3495. doi: 10.1016/S0006-3495(93)81190-1. URL http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1225880/.

Alain Destexhe, Thierry Bal, David A. McCormick, and Terrence J. Sejnowski. Ionic mechanisms underlying synchronized oscillations and propagating waves in a model of ferret thalamic slices. *Journal of Neurophysiology*, 76(3):2049–2070, 1996. doi: 10.1152/jn.1996.76.3.2049.

Alan L. Hodgkin and Andrew F. Huxley. A quantitative description of membrane current and its application to conduction and excitation in nerve. *The Journal of physiology*, 117(4):500, 1952. URL http://www.ncbi.nlm.nih.gov/pmc/articles/pmc1392413/.

Jason S. Sherfey, Austin E. Soplata, Salva Ardid, Erik A. Roberts, David A. Stanley, Benjamin R. Pittman-Polletta, and Nancy J. Kopell. DynaSim: A MATLAB Toolbox for Neural Modeling and Simulation. *Frontiers in Neuroinformatics*, 12, 2018. ISSN 1662-5196. doi: 10.3389/fninf.2018.00010. URL https://www.frontiersin.org/articles/10.3389/fninf.2018.00010/full.

Austin E. Soplata. DynaSim mechanism files for simulating thalamically-extended (Benita et al., 2012) Slow Wave model, March 2023a. URL https://doi.org/10.5281/zenodo.7725210.

Austin E. Soplata. Simulation and plotting scripts for (Soplata et al., 2023), March 2023b. URL https://doi.org/10.5281/zenodo.7725212.

Austin E. Soplata, Michelle M. McCarthy, Jason Sherfey, Shane Lee, Patrick L. Purdon, Emery N. Brown, and Nancy Kopell. Thalamocortical control of propofol phase-amplitude coupling. *PLoS Computational Biology*, 13(12):e1005879, December 2017. ISSN

1553-7358. doi: 10.1371/journal.pcbi.1005879. URL http://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.1005879.