Dependence of diffusion in *Escherichia coli* cytoplasm on protein size, environmental conditions and cell growth

N. Bellotto[1], J. Agudo-Canalejo[2,†], R. Colin[1,†], R. Golestanian[2,3,+], G. Malengo[1,+], V. Sourjik[1,+]

[1]Max Planck Institute for Terrestrial Microbiology and LOEWE Center for Synthetic Microbiology (SYNMIKRO), Karl-von-Frisch-Strasse 16, D-35043 Marburg, Germany

[2]Max Planck Institute for Dynamics and Self-Organization, Am Faßberg 17, D-37077 Göttingen, Germany

[3]Rudolf Peierls Centre for Theoretical Physics, University of Oxford, Oxford OX1 3PU, United Kingdom

[†]These authors equally contributed to this work

[+]Corresponding authors: ramin.golestanian@ds.mpg.de, gabriele.malengo@synmikro.mpi-marburg.mpg.de, victor.sourjik@synmikro.mpi-marburg.mpg.de

Abstract

Inside prokaryotic cells, passive translational diffusion typically limits the rates with which cytoplasmic proteins can reach their locations and interact with partners. Diffusion is thus fundamental to most cellular processes, but the understanding of protein mobility in the highly crowded and non-homogeneous environment of a bacterial cell is still limited. Here we investigated the mobility of a large set of proteins in the cytoplasm of *Escherichia coli*, by employing fluorescence correlation spectroscopy (FCS) combined with simulations and theoretical modeling. We conclude that cytoplasmic protein mobility could be well described by Brownian diffusion in the confined geometry of the bacterial cell and at high viscosity imposed by macromolecular crowding. The size dependence of protein diffusion is well consistent with the Stokes-Einstein relation for small fusion proteins when taking into account their shape, but larger proteins diffuse slower than expected from this relation. Pronounced subdiffusion and hindered mobility is observed for proteins with extensive interactions within the cytoplasm. Finally, we show that cytoplasmic viscosity has a temperature dependence comparable to that of water, and that protein mobility is increased by biosynthetic activity and cell growth.

Introduction

Diffusion of molecules is important for the function of any cellular system, setting the upper limit for mobility of proteins and other (macro)molecules and for the rates of many biochemical reactions that rely on random encounters between molecules (Schavemaker, Boersma, and Poolman 2018). Although the fundamental physics of diffusion in dilute aqueous solutions is well understood and mathematically described (Einstein 1906; Langevin 1908; Perrin 1909), diffusion in a cellular environment may be quite different (Schavemaker, Boersma, and Poolman 2018; Mika and Poolman 2011). The concentration of macromolecules in the bacterial cytoplasm, primarily proteins but also ribonucleic acids (RNAs), is extremely high. For *Escherichia coli*, it is around 300 mg/mL, which corresponds to a volume fraction of 25-30% (Cayley et al. 1991; Zimmerman and Trach 1991). Such macromolecular crowding could hinder free diffusion and influence kinetics of protein association and of gene expression (Klumpp et al. 2013; Tabaka et al. 2014; van den Berg, Boersma, and Poolman 2017). The effects of crowding on protein diffusion have been demonstrated both *in vitro* and *in vivo* (Dix and Verkman 2008; Rivas and Minton 2016). Compared to water, the diffusion of a free green fluorescent protein (GFP) was reported to be three to four times slower in the eukaryotic cytoplasm (Swaminathan, Hoang, and Verkman 1997) and up to ten times slower in the bacterial cytoplasm (Elowitz Michael et al. 1999; Nenninger, Mastroianni, and Mullineaux 2010; Mika et al. 2010; Kumar, Mommer, and Sourjik 2010).

In addition to the high density of macromolecules, the diversity in the size and chemical properties of the solutes makes the cytoplasmic environment highly inhomogeneous (Luby-Phelps 1999; Spitzer and Poolman 2013). How much the diffusion of a particular molecule is affected by macromolecular crowding might thus depend on the size (Muramatsu and Minton 1988) and shape of the molecule (Balbo et al. 2013) as well as on the nature of the crowders (Banks and Fradin

2005; Goins, Sanabria, and Waxham 2008). The effects of crowding observed in living cells appear

to be even more complex, varying not only with the diffusing particle but also with the

physiological state of the cell (Parry et al. 2014; Joyner et al. 2016) and the local cellular

environment (Konopka et al. 2006; Persson, Ambati, and Brandman 2020). Moreover, non-trivial

effects on diffusion arise due to reversible assembly and disassembly of the diffusing protein

complexes (Agudo-Canalejo, Illien, and Golestanian 2020), and possibly also due to the active

enhancement of enzyme diffusion by catalytic reactions (Golestanian 2015; Agudo-Canalejo et

al. 2018; Zhang and Hess 2019).

The dependence of the diffusion coefficient (D) of a protein in the cytoplasm on its size might thus

not necessarily follow the Stokes-Einstein (also called Stokes-Einstein-Sutherland-

Smoluchowski) relation that is valid in dilute solutions, $D \propto T/(\eta R)$ (Einstein 1906), where T is

absolute temperature in Kelvin, η is the viscosity of the medium and R is the hydrodynamic radius

of the particle. For globular proteins, R is given by the radius of gyration (Tyn and Gusek 1990)

and depends on the molecular mass (MM) as $R \propto MM^{\beta}$, where the exponent β would be 1/3 for

perfectly compact and globular proteins but is in practice within the range of 0.35-0.43 for typical

proteins, reflecting the fractal nature of the spatial distribution of protein mass (Smilgies and Folta-

Stogniew 2015; Enright and Leitner 2005). Several studies of protein diffusion in the cytoplasm

of *E. coli* have yielded different dependencies on the molecular mass, with an average $\beta \sim 0.7$

estimated based on the data pooled from multiple studies (Nenninger, Mastroianni, and Mullineaux

2010; Kumar, Mommer, and Sourjik 2010; Mika and Poolman 2011; Kalwarczyk, Tabaka, and

Holyst 2012), and thus substantially steeper than predicted by the Stokes-Einstein relation. Similar

exponent of ~0.7 was observed for a limited set of differently-sized proteins (Mika et al. 2010;

Stracy et al. 2021). However, neither of these studies took explicitly into account the non-

globularity of the used fluorescent constructs, where two or more proteins are typically connected

by flexible linkers. For such multidomain proteins, shape fluctuations and hydrodynamic

interactions between the different domains can have a sizeable effect on the effective diffusion

coefficient of the whole protein (Agudo-Canalejo and Golestanian 2020).

Besides macromolecular crowding, the translational diffusion of cytoplasmic proteins is also

influenced by intracellular structures, such as cytoskeletal filaments (Sabri et al. 2020), and by

(transient) binding to other macromolecules (Guigas and Weiss 2008; Saxton 2007). Both these

factors can not only reduce protein mobility but also lead to the anomalous subdiffusive behavior,

where the mean square displacement (MSD) of diffusing particles does not scale linearly with

time, as for Brownian diffusion in dilute solutions, but rather follows MSD $\propto t^{\alpha}$ with the anomalous

diffusion exponent α being <1 (Saxton 1996; Etoc et al. 2018). Subdiffusion is commonly observed

in eukaryotes, particularly at longer spatial scales, primarily due to the obstruction by the

cytoskeletal filaments (Di Rienzo et al. 2014; Sabri et al. 2020). Mobility of mRNA molecules in

the bacterial cytoplasm is also subdiffusive (Golding and Cox 2004; Lampo et al. 2017). In

contrast, the diffusion of several tested cytoplasmic bacterial proteins was apparently Brownian

(Bakshi, Bratton, and Weisshaar 2011; English et al. 2011).

Even for the same protein, e.g. GFP or its spectral variants, estimates of the diffusion coefficient

in the cytoplasm obtained in different studies vary widely (Schavemaker, Boersma, and Poolman

2018), which could be in part due to differences in methodologies. Most early studies in bacteria

relied on fluorescence recovery after photobleaching (FRAP), where diffusion is quantified from

the recovery of fluorescence in a region of the cell bleached by a high intensity laser (Loren et al.

2015). These measurements provided values of diffusion coefficient for GFP ranging from 3 μm^{2}s$^-$

^{1} to 14 μm^{2}s^{-1} (Elowitz Michael et al. 1999; Mullineaux et al. 2006; Konopka et al. 2009; Kumar,

Mommer, and Sourjik 2010; Mika et al. 2010; Nenninger, Mastroianni, and Mullineaux 2010;

Schavemaker, Smigiel, and Poolman 2017). More recently, single-particle tracking (SPT), where

diffusion is measured by following the trajectories of single fluorescent molecules over time

(Kapanidis, Uphoff, and Stracy 2018), became increasingly used. Lastly, diffusion can also be

studied *in vivo* using fluorescence correlation spectroscopy (FCS) (Cluzel, Surette, and Leibler

2000), which measures the time required by a fluorescent molecule to cross the observation volume

of a confocal microscope (Elson 2011). SPT and FCS measure protein mobility locally within the

cell, with FCS having also a significantly better temporal resolution than FRAP and SPT. Both

methods provided higher but still varying values of D_{GFP}, from 8 $\mu m^2 s^{-1}$ up to 18 $\mu m^2 s^{-1}$ (Meacci

et al. 2006; English et al. 2011; Sanamrad et al. 2014; Diepold et al. 2017; Rocha et al. 2019).

Protein mobility also depends on the environmental and cellular conditions that affect the structure

of the cytoplasm (Schavemaker, Boersma, and Poolman 2018). Previously studied examples

include effects of high osmolarity that increases macromolecular crowding and creates diffusion

barriers (Konopka et al. 2006; Konopka et al. 2009; Liu et al. 2019) and of the energy-dependent

fluidization of the cytoplasm to enable diffusion of large particles (Parry et al. 2014). Furthermore,

the surface charge of cytoplasmic proteins has been shown to have a dramatic effect on their

mobility (Schavemaker, Smigiel, and Poolman 2017).

Variations between values of diffusion coefficients observed even for the same model organism in

different studies, each investigating only a limited number of protein probes, using different

strains, growth conditions and measurement techniques, hampered drawing general conclusions

about the effective viscosity of bacterial cytoplasm and its dependence on the protein size. Here,

we address these limitations by systematically analyzing the mobility of a large number of

differently-sized cytoplasmic fluorescent protein constructs under standardized conditions by FCS.

We further combined experiments with Brownian dynamics simulations and theoretical modeling

of diffusion to correct for effects of confined cell geometry. For the majority of studied constructs,

we observe a uniform dependence of the diffusion coefficient on the protein size, with a

pronounced upper limit on diffusion at a given molecular mass. When corrected for the

confinement due to the bacterial cell geometry, the diffusion of these constructs was nearly

Brownian. Moreover, part of the deviation of the mass-dependence of their diffusion coefficients

from the Stokes-Einstein relation might be explained by the specific shape of the fusion proteins.

An aberrantly slow and anomalous diffusion of several protein constructs was apparently due to

their strong interactions with other cellular proteins and protein complexes, and disruption of these

interactions restored a Brownian diffusion close to the upper limit expected for their mass. We

further investigated the dependence of protein mobility on environmental temperature, on

exposure to antibiotics and on cellular growth, and we tested the mobility of proteins that are non-

native to *E. coli*. Finally, we show that under the same experimental conditions FCS and FRAP

measurements yield similar values of diffusion coefficients, suggesting that, in contrast to

eukaryotic cells (Di Rienzo et al. 2014), no pronounced dependence of protein mobility on spatial

scale could be observed in the bacterial cytoplasm.

141   Results

142   Dependence of cytoplasmic protein mobility on molecular mass measured by

143   FCS

For our analysis of cytoplasmic protein mobility, we generated a plasmid-encoded library of 31

cytoplasmic proteins (Figure 1 – supplementary table 1 and supplementary table 2) of *E. coli* fused

to superfolder GFP (sfGFP) (Pédelacq et al. 2006). We selected monomeric proteins with known

and roughly globular protein structure, to avoid effects on mobility due to multimerization or

irregular shape of proteins. These proteins belong to different cellular pathways, but DNA-binding

proteins were excluded. The expected size and stability of each construct was verified by gel

electrophoresis and immunoblotting (Figure 1 – figure supplement 1). Only one of the constructs,

ThpR-sfGFP, showed >20% degradation to free sfGFP, and it was therefore excluded from further

analyses. This was also the sole construct with an atypically high isoelectric point (pI), and all

remaining constructs have pI ranging from 5.1 to 6.2, as common for cytoplasmic proteins

(Schwartz, Ting, and King 2001). We further imaged the distribution of fusion proteins in the

cytoplasm. Except for RihA-sfGFP and NagD-sfGFP that were subsequently excluded, all other

constructs showed uniform localization (Figure 1A).

The mobility of the remaining 28 fusion constructs and of free sfGFP was investigated in living *E.*

coli cells by FCS (see Materials and Methods and Appendix 1). In order to facilitate the

measurements, cells were treated with the division inhibitor cephalexin for 45 min to induce

moderate elongation, yielding final cell length of ~ 5 μm (Figure 1A). During an FCS

measurement, the laser focus was positioned close to the polar region in the cell cytoplasm, and

the fluorescence intensity in the confocal volume was measured over time (Figure 1 – figure

supplement 2). For each individual cell, six subsequent acquisitions of 20 s each were performed

at the same position. The autocorrelation function (ACF) of the fluorescence intensity fluctuations

was independently calculated for each time interval and fitted to extract the mobility parameters

of the fluorescent proteins. Although we initially considered both the Brownian diffusion and the

anomalous diffusion models, the latter model proved to be considerably better in fitting the

experimental data (Figure 1 – figure supplement 3). The anomalous diffusion model was therefore

used to determine the diffusion (or residence) time (τ_D) of a fluorescent molecule in the confocal

volume and the anomalous diffusion exponent α for all ACFs (Figure 1B and Figure 1 –

supplementary table 1 and figure supplement 2). The averaged values of τ_D and α for each

individual cell were then calculated from these six individual acquisitions (Figure 1C and Figure

1 – figure supplement 4).

Despite their substantial intercellular variability, thus obtained mean values of the diffusion time

were clearly different between protein constructs (Figure 1C and Figure 1 – supplementary table

1. We next plotted the mean values of $1/\tau_D$, which reflect protein mobility, against the molecular

mass of protein constructs (Figure 1D). This dependence revealed a clear trend, where mobility of

more than half of the constructs decreased uniformly with their molecular mass, while some

exhibited much lower mobility than the other constructs of similar mass. In contrast, the anomalous

diffusion exponent α showed no apparent dependence on the protein size, ranging from 0.8 to 0.86

for most of the constructs (Figure 1E). Notably, the few protein constructs with α of ~ 0.8 or lower

were also among the ones with low mobility (Figure 1D and 1E, colored symbols).

Figure 1. Dependence of protein mobility in bacterial cytoplasm on molecular mass and cellular interactions.

(A) Examples of fluorescence microscopy images of *E. coli* cells expressing either sfGFP or the indicated sfGFP-tagged cytoplasmic proteins. Scale bar is 2 μm. (B) Representative autocorrelation functions (ACFs) measured by FCS for the indicated protein constructs. Data were fitted using the anomalous diffusion model (solid lines). All ACF curves were normalized to their respective maximal values to facilitate comparison. (C) Diffusion times (τ_D) measured for the indicated protein constructs. Each dot in the box plot represents the value for one individual cell, averaged over six consecutive acquisitions (Figure 1 – figure supplement 2) *** $p<0.0001$ in a two-tailed heteroscedastistic *t*-test. Exact p-values for all significance analyses performed in this study could be found in Appendix 4. (D, E). Dependence of protein mobility ($1/\tau_D$) (D) and apparent anomaly of diffusion (α) (E) on molecular mass. Each symbol represents

the average value for all individual cells that have been measured for that particular construct and the error bars

represent the standard error of the mean. Individual values are shown in Figure 1 – figure supplement 4 and the

numbers of measured cells are shown in Appendix 5. Protein constructs with most aberrant mobility are highlighted

in color and labeled. For fusions to ClpS, Map and DnaK, the values of $1/\tau_D$ and α for both original constructs

(diamonds) and constructs where mutations were introduced to disrupt interactions (circles) are shown. For Map, two

alternative amino acid substitutions that disrupt its interaction with the ribosome are shown (see Figure 1 – figure

supplement 6). (F-H) Cartoons illustrating the cellular interactions that could affect mobility of ClpS (F), Map (G),

and DnaK (H). ClpS engages with the ClpAP protease and with substrates, cartoon adapted from (Roman-Hernandez

et al. 2011). Map interacts with the actively translating ribosomes, cartoon adapted from (Sandikci et al. 2013). DnaK

interacts with unfolded client protein. Amino acidic residues that were mutated to disrupt interactions are highlighted

(see text for details).

Figure 1 – source data 1. Original values used for plots in Figure 1C-E.

Macromolecular interactions reduce protein mobility

We reasoned that the main group of constructs that exhibit mobility close to the apparent mass-

dependent upper limit represents proteins whose diffusion is only limited by macromolecular

crowding, and that the aberrantly low $1/\tau_D$ and α of other constructs might be due to their specific

interactions with other cellular proteins or protein complexes. Indeed, for three of these proteins

(ClpS, Map, DnaK) such interactions are well characterized and can be specifically disrupted. ClpS

is the adaptor protein that delivers degradation substrates to the protease ClpAP (Roman-

Hernandez et al. 2011). The substrate-binding site of ClpS is constituted by three amino acid

residues (D35, D36, H66) that interact with the N-terminal degron of target proteins (Figure 1F).

If these residues are mutated into alanine, substrate binding *in vitro* is substantially reduced

(Roman-Hernandez et al. 2011; Humbard et al. 2013). Additionally, ClpS directly docks to the

hexameric ClpA. Consistently, we observed that while stability of the mutant construct

ClpS$^{D35A\_D36A\_H66A}$-sfGFP was not affected (Figure 1 – figure supplement 5), its mobility in a

$\Delta clpA$ strain became significantly higher and less anomalous, with both $1/\tau_D$ and α reaching levels

similar to those of other proteins of similar mass (Figure 1D and 1E and Figure 1 – figure

supplement 6).

Similar results were obtained for the other two constructs. Map is the methionine aminopeptidase

that cleaves the N-terminal methionine from nascent polypeptide chains (Solbiati et al. 1999). Map

interacts with the negatively charged backbone of ribosomes through four positively charged

lysine residues (K211, 218, 224, 226) located in a loop (Figure 1G). If these residues are mutated

into alanine, the *in vitro* affinity of Map for the ribosomes is reduced (Sandikci et al. 2013). The

mobility of Map-sfGFP was indeed much increased by alanine substitutions at all four lysine sites

(Figure 1D and 1E and Figure 1 – figure supplement 5 and figure supplement 6). Interestingly,

charge inversion of lysines to glutamic acid did not further increase Map-sfGFP mobility as was

expected based on *in vitro* experiments (Sandikci et al. 2013).

DnaK is the major bacterial chaperone that binds to short hydrophobic polypeptide sequences,

which become exposed during protein synthesis, membrane translocation or protein unfolding

(Genevaux, Georgopoulos, and Kelley 2007). DnaK accommodates its substrate peptides inside a

hydrophobic pocket (Figure 1H). The substitution of the valine residue 436 with bulkier

phenylalanine creates steric hindrance that markedly decreases substrate binding to DnaK *in vitro*

(Mayer, Rüdiger, and Bukau 2000), and both the $1/\tau_D$ and α of DnaK^{V436F}-sfGFP were significantly

higher than for the correspondent wild-type construct (Figure 1D and 1E and Figure 1 – figure

supplement 5 and figure supplement 6). Nevertheless, in this case the $1/\tau_D$ did not reach the levels

of other proteins of similar molecular mass, which is likely explained by multiple interactions of

DnaK with other components of the cellular protein quality control machinery besides its binding

to substrates (Kumar and Sourjik 2012).

Cytoplasmic diffusion rate is conserved between bacterial species

We then investigated whether sfGFP fusions to non-native proteins, originating from other

bacteria, may show diffusion different from that of their *E. coli* counterparts. The existence of an

organism-dependent "quinary" code of unspecific, short living interactions has been recently

proposed in order to explain reduced mobility of heterologous human proteins in *E. coli* cytoplasm

(Mu et al. 2017). Thus, we investigated the mobility of proteins from other Gram-negative

proteobacteria *Yersinia enterocolitica*, *Vibrio cholerae*, *Caulobacter crescentus*, *Myxococcus*

xanthus and from the Gram-positive bacterium *Bacillus subtilis* that are homologous to several

freely-diffusing *E. coli* protein constructs (Figure 1 – figure supplement 7). Within this set of

constructs, we observed no significant differences of $1/\tau_D$ values from their *E. coli* counterparts,

except for AcnA from *M. xanthus* (Figure 1 – figure supplement 7A, C). Such lower mobility

might be a sign of multimerization of *M. xanthus* AcnA, although cellular distribution of this

construct was uniform. In contrast, most of these constructs showed slightly but significantly

increased anomaly of diffusion (Figure 1 – figure supplement 7B, D), which might reflect

increased "stickiness" of non-native proteins in *E. coli* cytoplasm.

Apparent anomaly of diffusion could be largely explained by confinement

When FCS measurements are performed in a confined space with dimensions comparable to those

of the observation volume, such confinement may affect the apparent mobility of fluorescent

molecules (Gennerich and Schild 2000; Jiang et al. 2020). To investigate the effect of confinement

on our FCS measurements, we performed Brownian dynamics simulations of FCS experiments

with particles undergoing three-dimensional, purely Brownian diffusion inside a bacterial-cell like

volume (Figure 2A *Inset*; see Materials and Methods). For the values of cell diameter commonly

observed under our growth conditions, 0.8-0.9 μm, and over a wide range of particle diffusion

coefficients, simulated autocorrelation functions could be indeed successfully fitted with the

anomalous diffusion model, yielding an anomalous diffusion exponent of around 0.8-0.9 (Figure

2A and 2B). This made us hypothesize that the relatively small apparent deviation from Brownian

diffusion in the fit, common to most constructs, may primarily reflect a confinement-induced effect

rather than proper subdiffusion. In order to test this, we grew *E. coli* in presence of the inhibitor

A22, that is known to affect the biosynthesis of bacterial cell wall and lead to increased cell width

(Ouzounov et al. 2016). Indeed, the cell diameter (Figure 2C) and the anomalous diffusion

exponent of sfGFP (Figure 2D) both significantly increased upon such A22 treatment. We

conclude that the values of α above 0.8, as observed for most protein constructs, reflect Brownian

diffusion with confinement rather than truly anomalous diffusion.

We thus derived an Ornstein-Uhlenbeck (OU) model for fitting FCS data, where the confinement

of Brownian diffusing fluorescent particles within the width of the cell is approximated by trapping

in a harmonic potential of the same width (Appendix 2). The two models fit the ACF of the

Brownian dynamic simulations comparably well and better than the model of unconfined

Brownian diffusion (Figure 2A and Figure 2 - figure supplement 1), with the OU model having

one less free parameter than the anomalous diffusion model. The OU model directly estimates the

ansatz diffusion coefficient with $\pm$ 5% accuracy for the typical cell widths observed in our

experiments (Figure 2 – figure supplement 2).

We then applied the OU model to re-fit the ACF data for all faster diffusing constructs and to

estimate their Brownian diffusion coefficients. The OU proved accurate in fitting the experimental

data, comparably to the anomalous diffusion model (Figure 2 – figure supplement 3). The

dependence of D on molecular mass for this set of constructs was scaling as $(MM)^{-\beta}$ with $\beta = 0.56$

$\pm\ 0.05$ (Figure 2E), less steep compared to the previous estimates (Kumar, Mommer, and Sourjik

2010; Mika et al. 2010; Stracy et al. 2021) but still steeper than expected from the Stokes-Einstein

relation, even when assuming $\beta = 0.4$ for not perfectly globular proteins (Figure 2E *Inset*) (Enright

and Leitner 2005; Smilgies and Folta-Stogniew 2015). In order to elucidate whether part of this

residual deviation may be accounted for by the specific shape of fusion constructs, where sfGFP

is fused to the differently-sized target proteins by a short flexible linker, we further applied a

previously derived model describing diffusion of such linked proteins (Appendix 3) (Agudo-

Canalejo and Golestanian 2020). The dependence of D on molecular mass predicted by this linked-

protein model seems indeed to better recapitulate our experimental data, particularly for smaller

protein fusions (Figure 2E *Inset*). Nevertheless, this model still clearly overestimates D for larger

protein fusions, suggesting that at least in this size range the dependence of D on molecular mass

in bacterial cytoplasm deviates from the Stokes-Einstein relation.

Figure 2. Protein diffusion in bacterial cytoplasm corrected for confinement. (A) Representative ACFs of simulated fluorescence intensity fluctuations. Simulations were performed in a confined geometry of a cell with indicated length L and diameter d, and dimensions of the measurement volume ω_0 and z_0, representing an experimental FCS measurement (*Inset*; see Materials and Methods) for two different values of the ansatz diffusion coefficient. Solid lines are fits by the models of unconfined Brownian diffusion, anomalous diffusion and by the Ornstein-Uhlenbeck (OU) model of Brownian diffusion under confinement, as indicated. (B) The exponent α extracted from the fit of the anomalous diffusion model to the ACFs data that were simulated at different values of the cell diameter. Corresponding values of the diffusion coefficient are shown in Figure 2 – figure supplement 2. (C,D) *E. coli* cells

grown for ~ 4 hours either in absence or in presence of 1 µg/mL of A22, show A22-dependent increase in the measured

cell diameter (C) and higher values of the exponent α extracted from the fit to the ACF measurements (D). For

consistency with other experiments, the cultures were also treated with cephalexin for 45 min. *** p<0.0001; * p<0.05

in a two-tailed heteroscedastistic *t*-test. (E) Dependence of the diffusion coefficient calculated from fitting the

experimental ACFs with the OU model of confined diffusion. Only the subset of apparently freely diffusing constructs

from Figure 1D has been analyzed with the OU model (see also Figure 1 – supplementary table 1). Each circle

represents the average value for all individual cells that have been measured for that particular construct, and the error

bars represent the standard error of the mean. *Inset* shows the fit of the mass dependence with an inverse power law

(solid blue line, exponent $\beta = 0.56 \pm 0.05$), and predictions of the Stokes-Einstein relation (black dashed line) and of

the model describing diffusion of two linked globular proteins (solid yellow line), both with exponent $\beta = 0.4$.

Figure 2 – source data 1. Original values used for plots in Figure 2B-E.

Protein diffusion coefficients measured using FRAP or FCS are consistent

Since many previous measurements of protein diffusion in bacteria were performed using FRAP,

we aimed to directly compare the results of FRAP and FCS measurements for a set of constructs

of different mass. Importantly, we used the same growth conditions and microscopy sample

preparation protocols as for the FCS experiments. The cells were photobleached in a region close

to the pole, similar to the position that was used for the FCS experiment. The recovery of

fluorescence was then followed for 11 seconds with the time resolution of 18 ms (Figure 3A). The

diffusion coefficients were computed from the time course of recovery with the plugin for ImageJ,

simFRAP (Blumenthal et al. 2015), which utilizes a simulation-based approach (Figure 3B). We

observed very good correlation between both values of diffusion coefficients, although for most

constructs the diffusion coefficients determined by FRAP were 5 to 30% lower than those obtained

from the FCS data (Figure 3C and Figure 1 – supplementary table 1).

Figure 3. Comparison between protein diffusion coefficients measured by FCS and FRAP. (A) Examples of FRAP measurements for two different constructs, sfGFP and AcnA-sfGFP. A 3x3 pixels area close to one cell pole (red circle) was photobleached with a high intensity laser illumination for 48 ms and the recovery of fluorescence in the bleached area was monitored for 11 s with the time resolution of 18 ms. (B) Representative curves of fluorescence recovery in FRAP experiments and their fitting using simFRAP. The experimental data (colored dots) are used by the simFRAP algorithm to simulate the underlying diffusional process (colored lines). The simulation is then used to compute the diffusion coefficient. The simulation proceeds until the recovery curve reaches a plateau, therefore it is interrupted at a different time for each curve. (C) Correlation between the diffusion coefficients measured in FCS experiments (D_{FCS}, fitting with the OU model; data from Figure 2E) and in FRAP experiment (D_{FRAP}, fitting with simFRAP). Error bars represent the standard error of the mean.

Figure 3 – source data 1. Original values used for plot in Figure 3C.

Effects of temperature, osmolarity and antibiotics on protein mobility

Having established the assay of protein mobility, we selected several differently-sized constructs with high mobility to characterize the impact of environmental and cellular perturbations that can be experienced by bacterial cells. We first confirmed the previously well characterized (Konopka et al. 2006; Konopka et al. 2009; Mika et al. 2010; Liu et al. 2019) decrease in protein mobility in bacterial cytoplasm upon osmotic upshift. Exposing *E. coli* cells to increased ionic strength by addition of 100 mM NaCl indeed decreased mobility of sfGFP by 30% (Figure 4 – figure supplement 1), which is comparable to previously measured values (Konopka et al. 2009).

Next, we studied the effect of temperature on protein mobility in bacterial cytoplasm. According to the Stokes-Einstein equation, the diffusion of a particle directly depends on the system's temperature in Kelvin and on the viscosity of the fluid, which itself changes with temperature. In the biologically relevant range, the temperature sensitivity of diffusion is primarily determined by the temperature dependence of water viscosity. The measured increase in mobility of sfGFP and two other constructs, by approximately 20-25% between 25°C and 35°C (Figure 4A), agrees well with the expected temperature-dependent decrease in water viscosity over 10°C (Huber et al. 2009). Importantly, growth temperature of *E. coli* culture had no apparent effect on protein mobility (Figure 4 – figure supplement 2), suggesting that – at least in the tested temperature range – *E. coli* lacks the growth-temperature dependent regulation of cytoplasmic viscosity that has been recently reported in the budding yeast (Persson, Ambati, and Brandman 2020).

Antibiotics that inhibit transcription (e.g., rifampicin) or translation (e.g., chloramphenicol) are well known to affect the spatial organization of bacterial chromosomes (Bakshi et al. 2014), but

their effects on the diffusive properties of the cytoplasm are less established. We observed that

protein mobility significantly increased upon inhibition of RNA transcription by rifampicin

treatment, roughly independently of protein size (Figure 4B and Figure 4 – figure supplement 3).

In contrast, inhibition of protein translation by chloramphenicol slightly reduced mobility.

Figure 4. Effects of temperature and antibiotics on protein mobility. (A) The effect of environmental temperature

on mobility of indicated protein constructs. As for other experiments, *E. coli* culture was grown at 37°C. (B) The

effects of antibiotics treatment on mobility of indicated protein constructs. Chloramphenicol (Cam; 200 µg/mL),

rifampicin (Rif; 200 µg/mL, in 0.1% v/v DMSO), or DMSO control (0.1% v/v) were added to growing *E. coli* culture

60 minutes prior to harvesting. *** p<0.0001; ** p<0.001; * p<0.05; NS: no statistically significant difference in a

two-tailed heteroscedastistic *t*-test.

Figure 4 – source data 1. Original values used for plots in Figure 4.

Effects of cell growth on protein mobility

In order to further investigate whether protein mobility might be affected by cell growth, we

compared cells that were incubated at 35°C on agarose pads containing either only M9 salts or M9

salts plus glucose and casamino acids. The observed mobility was indeed significantly higher in

the presence of nutrients, and this increase was similar (~ 40%) for the two tested differently-sized

constructs (Figure 5).

Figure 5. Effect of nutrient availability and growth on protein mobility. The mobility of indicated protein

constructs was measured in cells incubated at 35°C on agarose pads containing either only M9 salts or M9 salts

together with 20 mM glucose and 0.2% casamino acids (Glu + CA). Where indicated, chloramphenicol (Cam) was

also added to the agarose pads. *** p<0.0001; ** p<0.001; * p<0.05 in a two-tailed heteroscedastistic t-test.

Figure 5 – source data 1. Original values used for plots in Figure 5.

To distinguish the respective contributions of metabolic activity, and of biosynthesis and resulting

cell growth, we incubated cells in presence of both nutrients and chloramphenicol on the agarose

pad. Similar to our previous experiments where chloramphenicol was added to the batch culture,

its addition had no or little effect on protein mobility in absence of nutrients. In contrast, protein

mobility in presence of nutrients was strongly affected by chloramphenicol treatment. Thus, the

enhanced protein mobility in presence of nutrients largely depends on active protein production

and cell growth. Nevertheless, even chloramphenicol-treated cells exhibited a moderate increase

in protein mobility in presence of nutrients, indicating that the metabolic activity might also

contribute to the overall effect.

Discussion

Bacteria rely on translational diffusion to deliver proteins and other macromolecules to their

cellular destinations, including their reaction partners, and the diffusional properties of bacterial

cytoplasm are therefore fundamental to the understanding of bacterial cell biology. Consequently,

a number of studies have investigated protein mobility in bacteria, all showing strong effects of

macromolecular crowding in the bacterial cytoplasm on diffusion (Konopka et al. 2006;

Mullineaux et al. 2006; Konopka et al. 2009; Kumar, Mommer, and Sourjik 2010; Mika et al.

2010; Nenninger, Mastroianni, and Mullineaux 2010). Nevertheless, the relatively small number

of proteins investigated in each of these previous studies, and the differences between strains,

growth conditions and, particularly, between methodologies, limited general conclusions about

protein mobility, even in the most-studied environment of *E. coli* cytoplasm. For example,

combining data from different studies to determine the relation between size of a protein and its

cytoplasmic diffusion coefficient yielded only uncertain estimates (Mika and Poolman 2011;

Schavemaker, Boersma, and Poolman 2018). Such variability between different studies might be

further compounded by potentially profound effects on diffusion of size-independent protein

properties such as surface charge (Schavemaker, Smigiel, and Poolman 2017). Similarly, it

remains unclear whether a typical protein in the cytoplasm exhibits Brownian diffusion, as has

been shown in few examples (Bakshi, Bratton, and Weisshaar 2011; English et al. 2011), or rather

a subdiffusive behavior as common for proteins in eukaryotic cells (Di Rienzo et al. 2014; Sabri

et al. 2020) and for bacterial mRNA-protein complexes (Lampo et al. 2017; Golding and Cox

2004), and what might be the possible causes of anomalous diffusion in bacteria.

Here, we addressed these questions by systematically investigating the diffusive behavior of a

large set of fluorescent protein fusions to differently-sized cytoplasmic proteins of *E. coli*. We

demonstrate that, despite their diverse cellular functions, the majority of studied proteins exhibit a

rather uniform relation between their molecular mass and cytoplasmic mobility, with a clear upper

bound on protein mobility at a given molecular mass. Furthermore, the apparent weak anomaly of

diffusion observed in the FCS data analysis, with $\alpha \sim$ 0.85-0.9 for most proteins, could be

accounted for by confinement of the otherwise purely Brownian diffusing particles in the small

volume of a bacterial cell. Similar conclusions have been drawn by previous SPT studies for

several individual proteins (Bakshi, Bratton, and Weisshaar 2011; English et al. 2011). This

interpretation was directly confirmed by performing FCS measurements in *E. coli* with an

increased cell diameter, which indeed yielded $\alpha \sim$ 0.95.

We therefore used a model of purely Brownian diffusion under confinement (Ornstein-Uhlenbeck

model) to determine diffusion coefficients by directly fitting the autocorrelation functions of our

FCS measurements. The obtained overall dependence of diffusion coefficients on the molecular

mass of the fusion protein showed the exponent β = 0.56, steeper than predicted by the Stokes-

Einstein relation, with β = 0.33 for fully compact proteins or β = 0.4 for the more realistic case

where proteins are assumed to be not entirely compact (Enright and Leitner 2005; Smilgies and

Folta-Stogniew 2015). Nevertheless, at least for smaller constructs, the observed dependence of

the diffusion coefficient on the molecular mass could be well reproduced when the specific shape

of fusion constructs, where two roughly globular proteins are fused by a short linker, was taken

into account along with their imperfect globularity (Agudo-Canalejo and Golestanian 2020). In

contrast, the diffusion of larger proteins was still slower than calculated by this model.

Our analysis thus suggests that, despite the high crowdedness of the bacterial cytoplasm, the

diffusion of typical cytoplasmic proteins in bacteria is mostly Brownian and can be well described

by treating the cytoplasm as a viscous fluid, with only a moderate dependence of the effective

viscosity on the size of diffusing proteins. Given the diffusion coefficient determined in our study

for free sfGFP, ~14 $\mu m^2 s^{-1}$, for small proteins this effective viscosity of bacterial cytoplasm is only

approximately six times higher than in dilute solution (Potma et al. 2001). This diffusion

coefficient for GFP is substantially larger than the values reported in the early studies that used

FRAP (Elowitz Michael et al. 1999; Konopka et al. 2006; Mullineaux et al. 2006; Konopka et al.

2009; Kumar, Mommer, and Sourjik 2010; Mika et al. 2010; Nenninger, Mastroianni, and

Mullineaux 2010), although it is consistent with other FCS and SPT studies (Meacci et al. 2006;

English et al. 2011; Sanamrad et al. 2014; Diepold et al. 2017; Rocha et al. 2019). These

differences are apparently due to the limitations of early FRAP analyses that generally

underestimated protein mobility, rather than due to different spatial and temporal scales assessed

by the two techniques, since our direct comparison between FCS and FRAP measurements yielded

similar values of diffusion coefficients. Indeed, a more recent FRAP study also reported higher

diffusion coefficients for GFP (Schavemaker, Smigiel, and Poolman 2017).

Several proteins in our set showed much lower mobility than expected from their size, and in some

cases also clearly subdiffusive behavior. For three selected examples, such deviation could be

explained by specific association of these proteins with other proteins or multiprotein complexes.

This is consistent with theoretical studies suggesting that binding of diffusing molecules to

crowders can lead to subdiffusion (Saxton 2007; Guigas and Weiss 2008). A similar explanation

is likely to hold for the other aberrantly slow proteins.

Unspecific transient interactions might also explain the slightly subdiffusive behavior of sfGFP

fusions to proteins from other bacteria in *E. coli* cytoplasm. However, this anomaly was weak and

there was overall only little difference between the mobility of these non-native proteins and their

similarly-sized *E. coli* homologues, in contrast to pronounced differences observed between

bacterial and mammalian proteins (Mu et al. 2017). Thus, there is apparently little organism-

specific adaptation of freely diffusing proteins to their "bacterial host", with a possible exception

of bacteria with very different pH or ionic strength of the cytoplasm (Schavemaker, Smigiel, and

Poolman 2017), which might facilitate horizontal gene transfer among bacteria.

We further probed how the effective viscous properties of bacterial cytoplasm changed under

different physicochemical perturbations, using a subset of proteins that showed fast diffusion for

their molecular mass as "reporters" of effective viscosity. Consistent with the importance of

macromolecular crowding and in agreement with previous results (Konopka et al. 2009), protein

mobility decreased upon osmotic upshift that increases cytoplasmic crowding. In contrast, the

effective cytoplasmic viscosity significantly (~20%) decreases upon treatment with rifampicin that

inhibits transcription and thereby reduces the overall macromolecular crowding. This observation

is consistent with recent SPT measurements (Wlodarski et al. 2017; Rotter et al. 2021), and it

agrees well with the relative contribution of RNA to the macromolecular composition of an *E. coli*

cell (Cayley et al. 1991).

Despite multiple effects of environmental temperature on cellular processes, temperature

dependence of the cytoplasmic viscosity in the tested range was similar to that of water and

consistent with the Stokes-Einstein relation, decreasing by 20-30% for a temperature increase of

10°C (Huber et al. 2009). We observed no dependence of the effective cytoplasmic viscosity on

growth temperature, in contrast to the homeostatic adaptation of bacterial membrane fluidity

(Sinensky 1974) and of bacterial signaling (Oleksiuk et al. 2011; Almblad et al. 2021) to the growth

temperature. Since growth-temperature dependent adaptation of the cytosolic viscosity was

recently reported for budding yeast (Persson, Ambati, and Brandman 2020), it is surprising that

such compensation apparently does not exist in *E. coli*. One possible explanation for this difference

might be a broader range of growth temperatures for budding yeast *Saccharomyces cerevisiae*

compared to *E. coli* and a stronger temperature effect on protein diffusion in the yeast cytosol. Of

note, here we did not explore protein diffusion in thermally stressed *E. coli* cells, which might have

more profound effects on the properties of bacterial cytoplasm as recently shown for *Listeria*

monocytogenes (Tran et al. 2021).

Finally, we observed that protein mobility was significantly higher in actively growing cells. This

"fluidizing" effect of growth seems to be primarily due to the biosynthetic processes, likely protein

translation, or to cell growth itself, whereas the effect of metabolic activity in presence of nutrients

was weaker although still significant. Thus, the observed phenomenon may be different from

previously characterized energy-dependent fluidization of the bacterial cytoplasm that enables

mobility of large multiprotein complexes but apparently does not affect free GFP (Montero Llopis

et al. 2012; Parry et al. 2014). The interplay between these energy-, metabolism- and growth-

dependent effects on diffusional properties of bacterial cytoplasm remains to be investigated.

Summarizing, here we present a comprehensive study of the diffusional properties of proteins in

the cytoplasm of *E. coli*, combining FCS measurements with theoretical modeling of diffusion in

a confined geometry of a bacterial cell. This analysis enabled us to distinguish a large set of

proteins with essentially Brownian diffusion, which could be used to characterize the effective

cytoplasmic viscosity over a range of molecular masses and its dependence on macromolecular

crowding, protein interactions, environmental temperature and cell growth.

Materials and methods

Bacterial strains, plasmids and media

All strains and plasmids used in this study are listed in Figure 1 – supplementary table 2. All

experiments were performed in the *Escherichia coli* strain W3110 (Serra et al. 2013). Genes of

interest were amplified by PCR using Q5 polymerase (New England Biosciences) and cloned

along with sfGFP using Gibson assembly (Gibson et al. 2009) into pTrc99A vector (Amann, Ochs,

and Abel 1988), under control of the *trc* promoter inducible by isopropyl ß-D-1-

thiogalactopyranoside (IPTG). In all cases sfGFP was fused at the C-terminus of the protein of

interest with a GGGGS linker. The stability of the fusion constructs was verified by gel

electrophoresis and immunoblotting using an anti-GFP primary antibody (JL-8 monoclonal,

Takara). Point mutations were introduced by site-directed mutagenesis (New England

Biosciences). *ΔclpA* was generated by transferring the kanamycin resistance cassette from the

corresponding mutant in the Keio collection (Baba et al. 2006) by P1 transduction. The cassette

was further removed by FLP recombinase carried on the temperature-sensitive plasmid pCP20

(Cherepanov and Wackernagel 1995).

E. coli cultures were grown in M9 minimal medium (48 mM Na_2HPO_4, 22 mM KH_2PO_4, 8.4 mM

NaCl, 18.6 mM NH_4Cl) supplemented with 0.2% casamino acids, 0.4% glucose and 100 µg/mL

ampicillin for selection. The overnight cultures were diluted to $OD_{600} = 0.035$ and grown for 3.5

hours at 37°C and 200 rpm shaking. Cultures were treated for additional 45 minutes, under the

same temperature and shaking conditions, with 10 µg/mL cephalexin and with 0 - 15 µM IPTG

(Figure 1 – supplementary table 1), to induce expression of the fluorescent protein constructs.

Where indicated, cultures were further incubated with 200 µg/mL rifampicin, 200 µg/mL

chloramphenicol or DMSO as a mock treatment for 1 hour under the same temperature and shaking

conditions.

FCS data acquisition

Cells were harvested by centrifugation at 7000g for 3 minutes and washed 3 times in tethering

buffer (10 mM K_2HPO_4, 10 mM KH_2PO_4, 1 µM methionine, 10 mM sodium lactate, buffered with

NaOH to pH 7). 2.5 µL of bacterial cells were then spread on a small 1% agarose pad prepared in

tethering buffer salts (10 mM K_2HPO_4, 10 mM KH_2PO_4 buffered with NaOH to pH 7)**,** unless

differently stated. Imaging was performed on Ibidi 2-well µ-Slides (#1.5H, 170 µm ± 5 µm). After

the 45 minutes treatment with cephalexin, length of most bacterial cells was in a range of 4-8 µm.

For consistency, only cells of 5 ± 1 µm in length were selected for FCS experiments.

FCS measurements were performed on a LSM 880 confocal laser scanning microscope (Carl Zeiss

Microscopy) using a C-Apochromat 40×/1.2 water immersion objective selected for FCS. sfGFP

was excited with a 488 nm Argon laser (25 mW) and fluorescence emission was collected from

490 to 580 nm. Each sample was equilibrated for at least 20 minutes at 25°C (or 35°C when

specified), on the stage of the microscope and measurements were taken at the same temperature.

FCS measurements were acquired within 60 minutes from the sample preparation. The pinhole

was aligned on a daily basis, by maximizing the fluorescence intensity count rate of an Alexa488

(Invitrogen) solution (35 nM) in phosphate-buffered saline (PBS; 137 mM NaCl, 2.7 mM KCl, 8

mM Na_2HPO_4, 1.8 mM KH_2PO_4, pH 7.4). The coverslip collar adjustment ring of the water

immersion objective was also adjusted daily, maximizing the fluorescence intensity signal and the

brightness of Alexa 488. The laser power was adjusted in order to obtain molecular brightness (i.e.

photon counts per second per molecule, cpsm) of 10 kcpsm for Alexa 488, using the ZEN software

(Carl Zeiss Microscopy). The brightness of Alexa 488 was used as a daily reference to ensure

constant laser power and adjusting it using the software-provided laser power percentage whenever

necessary (range over the entire set of measurements was $0.11 - 0.18\%$). Before each measurement

session, we acquired three sequential FCS measurements of Alexa488 in PBS, to verify the

reproducibility of the confocal volume shape and size. The ratio between axial and lateral beam

waist $S = \frac{z_0}{\omega_0} = 8.0 \pm 0.2$ was obtained from a Brownian fit of the Alexa 488 autocorrelation

curves using the ZEN software. For the lateral beam waist, we obtained $\omega_0 = 0.186 \pm 0.001$ μm,

calculated from the diffusion time $\tau_D = 20.9 \pm 0.11$ μs obtained from the Brownian fit, being

$$D = \frac{\omega_0^2}{4\tau_D} \qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad (1)$$

and being $D_{Alexa488} = 414$ μm^2/s at 25°C (Petrov et al. 2006).

For the FCS measurements *in vivo*, the laser was positioned at the center of the short length axis

and typically 0.8-1 μm from one of the cell poles along the long axis. The positioning was chosen

to maintain distance from the cell membranes and to keep the confocal volume possibly away from

the bacterial chromosome. For each cell, six sequential fluorescence intensity acquisitions of 20

seconds each were performed on the same spot (Figure 1 – figure supplement 2). The laser power

used for measurements *in vivo* was fixed to a value about 7 times lower than for Alexa488 in PBS,

in order to reduce photobleaching. Confocal images of the selected cell were routinely acquired

before and after the FCS measurement to verify focal (*z*) and positioning (*xy*) stability (see

Appendix 1 for additional information on the FCS measurements).

FCS data analysis

Due to the small size of bacterial cells, fluorescence intensity traces are affected by photobleaching

(Appendix 1). The effect of photobleaching on autocorrelation curves was corrected by detrending

the long-time fluorescence decrease of each of the six fluorescence intensity traces using an ImageJ

plugin (Jay Unruh, https://research.stowers.org/imagejplugins/index.html, Stowers Institute for

Medical Research, USA). The plugin calculates the autocorrelation function (ACF) from each

fluorescence intensity trace, correcting it for the photobleaching effect by approximating the

decreasing fluorescence intensity trend with a multi-segment line (the number of segments was

fixed to two). We obtained almost identical ACFs correcting for photobleaching effects by local

averaging (Appendix 1 - figure 3) using the FCS-dedicated software package Fluctuation Analyzer

(Wachsmuth et al. 2015). In both cases, ACFs were calculated starting at 2 μs, since at times

shorter than 2 μsec, ACFs can be significantly affected by the GaAsp photomultipliers

afterpulsing.

For each FCS measurement, we fitted all the six ACFs, calculated using the multi-segment

detrending method, with a three-dimensional anomalous diffusion model that includes one

diffusive component and one blinking component due to the protonation-deprotonation of the

chromophore of sfGFP, according to the Equation (2):

$$G(\tau) = G_\infty + \frac{1}{N} \left(\frac{1 - F_P + F_P e^{-\frac{\tau}{\tau_P}}}{1 - F_P} \right) \frac{1}{1 + \left(\frac{\tau}{\tau_D}\right)^\alpha \sqrt{1 + \frac{1}{S^2}\left(\frac{\tau}{\tau_D}\right)^\alpha}} \qquad (2)$$

where N is the average number of particles in the confocal volume, F_P is the fraction of particles

in the non-fluorescent state, τ_P is the protonation-deprotonation lifetime at pH 7.5, $S = \frac{z_0}{\omega_0}$, the

aspect ratio of the confocal volume with z_0 and ω_0 being the axial and lateral beam waists, τ_D is the

diffusion time in the confocal volume, α is the anomalous diffusion exponent, and G_∞ is the offset

of the autocorrelation function. The protonation-deprotonation lifetime (τ_P) for sfGFP was fixed

to 25 µs according to FCS measurements for sfGFP in PBS at pH 7.5 (Cotlet et al. 2006). The

aspect ratio of the confocal volume was fixed to $S = 8$ in the fittings to be consistent with the

experimental calibration (see above). All other parameters were left free. For each FCS

measurement, we calculated the average diffusion time τ_D and the average anomalous diffusion

exponent α based on the autocorrelation curves of the six sequential fluorescence intensity traces.

Importantly, no significant trend in τ_D or α was apparent when comparing the six sequential

autocorrelation functions acquired for a given bacterial cell (Appendix 1 - figure 4). Fitting to the

anomalous diffusion model was performed using the Levenberg-Marquardt algorithm in the FCS

analysis-dedicated software QuickFit 3.0 developed by Jan Wolfgang Krieger and Jörg Langowski

(Deutsches Krebsforschungszentrum, Heidelberg, https://github.com/jkriege2/QuickFit3). Identical

results were obtained when fitting the data with OriginPro.

Alternatively, the autocorrelation functions were fitted by the Ornstein-Uhlenbeck model

(Appendix 2) according to Equation (3):

$G(\tau) = G_\infty + \frac{1}{N}\left(\frac{1-F_P+F_Pe^{-\frac{\tau}{\tau_P}}}{1-F_P}\right)\left(1 + 2\frac{\sigma^2}{\omega_0^2}\frac{1-e^{-\frac{1}{2}\frac{\omega_0^2}{\sigma^2}\frac{\tau}{\tau_D}}}{1+\frac{1}{8}\frac{\omega_0^2}{\sigma^2}}\right)^{-\frac{1}{2}}\left(1+\frac{\tau}{\tau_D}\right)^{-\frac{1}{2}}\left(1 + 2\frac{\sigma^2}{S^2\omega_0^2}\frac{1-e^{-\frac{1}{2}\frac{\omega_0^2}{\sigma^2}\frac{\tau}{\tau_D}}}{1+\frac{1}{8}\frac{S^2\omega_0^2}{\sigma^2}}\right)^{-\frac{1}{2}}$

(3)

where S and τ_P were fixed to the same values mentioned for equation (2), ω_0 was fixed to 0.19 and

σ was fixed to $d/2$=0.42 μm, being d the typical diameter of an *E. coli* cell (see Ornstein-Uhlenbeck

model validation paragraph). Fitting to the Ornstein-Uhlenbeck model was performed with

OriginPro.

FRAP data acquisition and analysis

Cells for FRAP experiments were grown and prepared for imaging following the same protocol as

for the FCS measurements. Due to the higher sensitivity of FCS at low fluorophore concentrations,

several fusion constructs required higher induction by IPTG (Figure 1 – supplementary table 1) to

obtain fluorescence intensity suitable for FRAP. The same LSM 880 confocal microscope,

including objective and light path was used for FRAP as for the FCS measurements. The bacterial

cell was imaged at 40×40 pixels with 30× zoom (pixel size 0.177 μm) with a pixel dwell time of

3.15 μs. First, 15 pre-bleaching frames were acquired at 2% laser power, subsequently the

photobleaching was performed on a 3×3 pixels area on one cell pole with 100% laser power for a

total of 48 ms and 584 post-bleaching frames were acquired to monitor the fluorescence recovery.

We observed that the mobile fraction for all constructs was >0.9. FRAP measurements were

analyzed using simFRAP (Blumenthal et al. 2015), an ImageJ plugin based on a simulation

approach implemented in a fast algorithm, which bypasses the need of using analytical models to

interpolate the data. The simFRAP algorithm simulates two-dimensional random walks in each

pixel, using the first image acquired after bleaching to define initial and boundary conditions, and

it resolves numerically the diffusion equation by iterative simulation. The frame time and pixel

size were fixed respectively to 0.018 s and 0.177 μm, and the target cell and the bleached region

were defined as ImageJ ROIs (regions of interest). Of note, we used the target cell itself as a

reference to compensate for the gradual bleaching during the measurement, as done previously

(Kumar, Mommer, and Sourjik 2010). This enabled us to achieve the highest possible temporal

resolution, by reducing the acquisition area to a single *E. coli* cell. The FRAP derived diffusion

coefficient D_{FRAP} was directly obtained as output of the plugin.

Brownian dynamics simulations

We performed Brownian dynamics simulations of uncorrelated point particles under confinement.

The $N = 50$ fluorescent particles performed a random walk with steps taken from a Gaussian

distribution of width $\sqrt{2D\Delta t}$, with D the free diffusion coefficient and $\Delta t = 10^{-6}$ s the simulation

step. Confinement was imposed by redrawing the random steps that moved out of the confinement

volume. Imposing elastic reflections on the walls yielded identical results. The confinement

volume was assumed to be a cylinder of diameter d and length (*L-d*) closed at both ends by

hemispheric caps of diameter d, idealizing the shape of *E. coli*. The cell length was fixed to $L = 5$

μm. The diameter was varied in the range $d = [0.7, 1]$ μm. The collected fluorescence intensity

was computed at each time step assuming a Gaussian intensity profile of the laser beam, $I(t) =$

$\sum_{i=1}^{N} I_G(\boldsymbol{r}_i(t) - \boldsymbol{r}_0)$ with $\boldsymbol{r}_i(t)$ the position of particle i, $\boldsymbol{r}_0$ the center of the confocal volume and

$I_G(r = (x, y, z)) = \exp\left(-2\left(\frac{x^2 + y^2}{\omega_0^2} + \frac{z^2}{z_0^2}\right)\right)$, with $\omega_0 = 200$ nm and $z_0 = 800$ nm the lateral and

axial widths of the confocal volume. The normalized intensity autocorrelation $C(dt) =$

$\frac{\langle I(t+dt)I(t)\rangle}{\langle I(t+dt)\rangle\langle I(t)\rangle} - 1$ is computed for logarithmically spaced lag times dt, to reflect experimental

practices. The center of the confocal volume was chosen in the center of the cell along the y and

z axes and 1 μm away from the edge of the cell along the longitudinal x axis of the cell, similarly

to experimental conditions. The intensity autocorrelation function was finally multiplied by an

exponential decay, $\left(1 + 0.1 * \exp - \frac{dt}{\tau_H}\right)/1.1$ with $\tau_H = 25$ μs, to mimic the blinking component

due to the protonation-deprotonation process of sfGFP, before fitting with the different models of

diffusion. The code used for this simulation is available in GitHub

(https://github.com/croelmiyn/Simulation_FCS_in_Bacteria) and via

DOI: 10.5281/zenodo.5940484.

Validation of fitting by the Ornstein-Uhlenbeck model

We first estimated the relation between the width σ of the potential well and the diameter d of the

bacteria by fitting the ACF of the Brownian simulations with the OU model, fixing all parameters

except σ to their ansatz values. The best fit was obtained for $\sigma \simeq d/2$ over the whole range of

tested parameters. To mimic the fit procedure of experimental data and evaluate the accuracy of

the diffusion coefficient estimation by the OU model (Fig. 2C), we then fixed $\sigma = d/2$, and ω_0

and z_0 to their ansatz values, since they are measured independently in experiments, whereas the

diffusion coefficient, number of particles N in the confocal volume, fraction of triplet excitation

and background noise were taken as free parameters.

Acknowledgements

We thank Lotte Søgaard-Andersen, Martin Thanbichler, Knut Drescher and Andreas Diepold for

providing bacterial genomic DNA. This work was supported by the Max Planck Society.

Competing Interests

The authors declare no competing interests.

References

Agudo-Canalejo, Jaime, Tunrayo Adeleke-Larodo, Pierre Illien, and Ramin Golestanian. 2018.
'Enhanced Diffusion and Chemotaxis at the Nanoscale', *Accounts of Chemical Research*,
51: 2365-72.

Agudo-Canalejo, Jaime, and Ramin Golestanian. 2020. 'Diffusion and steady state distributions
of flexible chemotactic enzymes', *The European Physical Journal Special Topics*, 229:
2791-806.

Agudo-Canalejo, Jaime, Pierre Illien, and Ramin Golestanian. 2020. 'Cooperatively enhanced
reactivity and "stabilitaxis" of dissociating oligomeric proteins', *Proceedings of the
National Academy of Sciences*, 117: 11894.

Almblad, Henrik, Trevor E. Randall, Fanny Liu, Katherine Leblanc, Ryan A. Groves, Weerayuth
Kittichotirat, Geoffrey L. Winsor, Nicolas Fournier, Emily Au, Julie Groizeleau,
Jacquelyn D. Rich, Yuefei Lou, Elise Granton, Laura K. Jennings, Larissa A. Singletary,
Tara M. L. Winstone, Nathan M. Good, Roger E. Bumgarner, Michael F. Hynes, Manu
Singh, Maria Silvina Stietz, Fiona S. L. Brinkman, Ayush Kumar, Ann Karen Cornelia

Brassinga, Matthew R. Parsek, Boo Shan Tseng, Ian A. Lewis, Bryan G. Yipp, Justin L. MacCallum, and Joe Jonathan Harrison. 2021. 'Bacterial cyclic diguanylate signaling networks sense temperature', *Nature Communications*, 12: 1986.

Amann, E., B. Ochs, and K. J. Abel. 1988. 'Tightly regulated tac promoter vectors useful for the expression of unfused and fused proteins in *Escherichia coli*', *Gene*, 69: 301-15.

Baba, T., T. Ara, M. Hasegawa, Y. Takai, Y. Okumura, M. Baba, K. A. Datsenko, M. Tomita, B. L. Wanner, and H. Mori. 2006. 'Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection', *Molecular Systems Biology*, 2: 2006 0008.

Bakshi, Somenath, Benjamin P Bratton, and James C Weisshaar. 2011. 'Subdiffraction-Limit Study of Kaede Diffusion and Spatial Distribution in Live *Escherichia coli*', *Biophysical Journal*, 101: 2535-44.

Bakshi, Somenath, Heejun Choi, Jagannath Mondal, and James C. Weisshaar. 2014. 'Time-dependent effects of transcription- and translation-halting drugs on the spatial distributions of the *Escherichia coli* chromosome and ribosomes', *Molecular Microbiology*, 94: 871-87.

Balbo, Jessica, Paolo Mereghetti, Dirk-Peter Herten, and Rebecca C. Wade. 2013. 'The shape of protein crowders is a major determinant of protein diffusion', *Biophysical Journal*, 104: 1576-84.

Banks, Daniel S., and Cécile Fradin. 2005. 'Anomalous Diffusion of Proteins Due to Molecular Crowding', *Biophysical Journal*, 89: 2960-71.

Blumenthal, D., L. Goldstien, M. Edidin, and L. A. Gheber. 2015. 'Universal Approach to FRAP Analysis of Arbitrary Bleaching Patterns', *Scientific Reports*, 5: 11655.

Cayley, Scott, Barbara A. Lewis, Harry J. Guttman, and M. Thomas Record. 1991. 'Characterization of the cytoplasm of *Escherichia coli* K-12 as a function of external osmolarity: Implications for protein-DNA interactions in vivo', *Journal of Molecular Biology*, 222: 281-300.

Cherepanov, Peter P., and Wilfried Wackernagel. 1995. 'Gene disruption in *Escherichia coli*:
TcR and KmR cassettes with the option of Flp-catalyzed excision of the antibiotic-
resistance determinant', *Gene*, 158: 9-14.

Cluzel, Philippe, Michael Surette, and Stanislas Leibler. 2000. 'An Ultrasensitive Bacterial
Motor Revealed by Monitoring Signaling Proteins in Single Cells', *Science*, 287: 1652-
55.

Cotlet, M., P. M. Goodwin, G. S. Waldo, and J. H. Werner. 2006. 'A comparison of the
fluorescence dynamics of single molecules of a green fluorescent protein: one- versus
two-photon excitation', *Chemphyschem*, 7: 250-60.

Di Rienzo, Carmine, Vincenzo Piazza, Enrico Gratton, Fabio Beltram, and Francesco Cardarelli.
2014. 'Probing short-range protein Brownian motion in the cytoplasm of living cells',
Nature Communications, 5: 5891.

Diepold, A., E. Sezgin, M. Huseyin, T. Mortimer, C. Eggeling, and J. P. Armitage. 2017. 'A
dynamic and adaptive network of cytosolic interactions governs protein export by the
T3SS injectisome', *Nature Communications*, 8: 15940.

Dix, James A., and A. S. Verkman. 2008. 'Crowding Effects on Diffusion in Solutions and Cells',
Annual Review of Biophysics, 37: 247-63.

Einstein, Albert 1906. 'On the theory of the Brownian movement', *Annalen der Physik*, 19: 371-
81.

Elowitz Michael, B., G. Surette Michael, Pierre-Etienne Wolf, B. Stock Jeffry, and Stanislas
Leibler. 1999. 'Protein Mobility in the Cytoplasm of *Escherichia coli*', *Journal of
Bacteriology*, 181: 197-203.

Elson, E. L. 2011. 'Fluorescence correlation spectroscopy: past, present, future', *Biophysical
Journal*, 101: 2855-70.

Enderlein, Jörg. 2012. 'Polymer Dynamics, Fluorescence Correlation Spectroscopy, and the
Limits of Optical Resolution', *Physical Review Letters*, 108: 108101.

Enderlein, Jörg, Ingo Gregor, Digambara Patra, Thomas Dertinger, and U. Benjamin Kaupp. 2005. 'Performance of Fluorescence Correlation Spectroscopy for Measuring Diffusion and Concentration', *Chemphyschem*, 6: 2324-36.

English, Brian P., Vasili Hauryliuk, Arash Sanamrad, Stoyan Tankov, Nynke H. Dekker, and Johan Elf. 2011. 'Single-molecule investigations of the stringent response machinery in living bacterial cells', *Proceedings of the National Academy of Sciences of the United States of America*, 108: E365-E73.

Enright, Matthew B., and David M. Leitner. 2005. 'Mass fractal dimension and the compactness of proteins', *Physical Review E*, 71: 011912.

Etoc, Fred, Elie Balloul, Chiara Vicario, Davide Normanno, Domenik Liße, Assa Sittner, Jacob Piehler, Maxime Dahan, and Mathieu Coppey. 2018. 'Non-specific interactions govern cytosolic diffusion of nanosized objects in mammalian cells', *Nature Materials*, 17: 740-46.

Genevaux, P., C. Georgopoulos, and W. L. Kelley. 2007. 'The Hsp70 chaperone machines of *Escherichia coli*: a paradigm for the repartition of chaperone functions', *Molecular Microbiology*, 66: 840-57.

Gennerich, Arne, and Detlev Schild. 2000. 'Fluorescence Correlation Spectroscopy in Small Cytosolic Compartments Depends Critically on the Diffusion Model Used', *Biophysical Journal*, 79: 3294-306.

Gibson, D. G., L. Young, R. Y. Chuang, J. C. Venter, C. A. Hutchison, 3rd, and H. O. Smith. 2009. 'Enzymatic assembly of DNA molecules up to several hundred kilobases', *Nature Methods*, 6: 343-5.

Goins, Aron B., Hugo Sanabria, and M. Neal Waxham. 2008. 'Macromolecular crowding and size effects on probe microviscosity', *Biophysical Journal*, 95: 5362-73.

Golding, Ido, and Edward C. Cox. 2004. 'RNA dynamics in live *Escherichia coli* cells', *Proceedings of the National Academy of Sciences of the United States of America*, 101: 11310.

Golestanian, Ramin. 2015. 'Enhanced Diffusion of Enzymes that Catalyze Exothermic
Reactions', *Physical Review Letters*, 115: 108102.

Guigas, Gernot, and Matthias Weiss. 2008. 'Sampling the Cell with Anomalous Diffusion—The
Discovery of Slowness', *Biophysical Journal*, 94: 90-94.

Huber, M. L., R. A. Perkins, A. Laesecke, D. G. Friend, J. V. Sengers, M. J. Assael, I. N.
Metaxa, E. Vogel, R. Mareš, and K. Miyagawa. 2009. 'New International Formulation for
the Viscosity of H2O', *Journal of Physical and Chemical Reference Data*, 38: 101-25.

Humbard, M. A., S. Surkov, G. M. De Donatis, L. M. Jenkins, and M. R. Maurizi. 2013. 'The N-
degradome of *Escherichia coli*: limited proteolysis in vivo generates a large pool of
proteins bearing N-degrons', *The Journal of Biological Chemistry*, 288: 28913-24.

Jiang, Yanfei, Bingxian Xu, Artem Melnykov, Guy M. Genin, and Elliot L. Elson. 2020.
'Fluorescence Correlation Spectroscopy and Photon Counting Histograms in Finite,
Bounded Domains', *Biophysical Journal*, 119: 265-73.

Joyner, Ryan P., Jeffrey H. Tang, Jonne Helenius, Elisa Dultz, Christiane Brune, Liam J. Holt,
Sebastien Huet, Daniel J. Müller, and Karsten Weis. 2016. 'A glucose-starvation response
regulates the diffusion of macromolecules', *Elife*, 5: e09376.

Kalwarczyk, Tomasz, Marcin Tabaka, and Robert Holyst. 2012. 'Biologistics--diffusion
coefficients for complete proteome of *Escherichia coli*', *Bioinformatics (Oxford,*
England), 28: 2971-78.

Kapanidis, A. N., S. Uphoff, and M. Stracy. 2018. 'Understanding Protein Mobility in Bacteria
by Tracking Single Molecules', *Journal of Molecular Biology*, 430: 4443-55.

Klumpp, Stefan, Matthew Scott, Steen Pedersen, and Terence Hwa. 2013. 'Molecular crowding
limits translation and cell growth', *Proceedings of the National Academy of Sciences*,
110: 16754.

Konopka, M. C., I. A. Shkel, S. Cayley, M. T. Record, and J. C. Weisshaar. 2006. 'Crowding and
confinement effects on protein diffusion in vivo', *Journal of Bacteriology*, 188: 6115-23.

Konopka, M. C., K. A. Sochacki, B. P. Bratton, I. A. Shkel, M. T. Record, and J. C. Weisshaar.
811   2009. 'Cytoplasmic protein mobility in osmotically stressed *Escherichia coli*', *Journal of*
812   *Bacteriology*, 191: 231-7.

813   Kumar, M., M. S. Mommer, and V. Sourjik. 2010. 'Mobility of cytoplasmic, membrane, and
814   DNA-binding proteins in *Escherichia coli*', *Biophysical Journal*, 98: 552-9.

815   Kumar, Mohit, and Victor Sourjik. 2012. 'Physical map and dynamics of the chaperone network
816   in *Escherichia coli*', *Molecular Microbiology*, 84: 736-47.

817   Lampo, Thomas J., Stella Stylianidou, Mikael P. Backlund, Paul A. Wiggins, and Andrew J.
818   Spakowitz. 2017. 'Cytoplasmic RNA-Protein Particles Exhibit Non-Gaussian
819   Subdiffusive Behavior', *Biophysical Journal*, 112: 532-42.

820   Langevin, Paul. 1908. 'Sur la théorie du mouvement brownien', *Comptes-rendus de l'Académie*
821   *des sciences*, 146: 530-33.

822   Liu, B., Z. Hasrat, B. Poolman, and A. J. Boersma. 2019. 'Decreased Effective Macromolecular
823   Crowding in *Escherichia coli* Adapted to Hyperosmotic Stress', *Journal of Bacteriology*,
824   201.

825   Loren, N., J. Hagman, J. K. Jonasson, H. Deschout, D. Bernin, F. Cella-Zanacchi, A. Diaspro, J.
826   G. McNally, M. Ameloot, N. Smisdom, M. Nyden, A. M. Hermansson, M. Rudemo, and
827   K. Braeckmans. 2015. 'Fluorescence recovery after photobleaching in material and life
828   sciences: putting theory into practice', *Quarterly Reviews of Biophysics*, 48: 323-87.

829   Luby-Phelps, Katherine. 1999. 'Cytoarchitecture and Physical Properties of Cytoplasm: Volume,
830   Viscosity, Diffusion, Intracellular Surface Area.' in Harry Walter, Donald E. Brooks and
831   Paul A. Srere (eds.), *International Review of Cytology* (Academic Press).

832   Macháň, Radek, Yong Hwee Foo, and Thorsten Wohland. 2016. 'On the Equivalence of FCS and
833   FRAP: Simultaneous Lipid Membrane Measurements', *Biophysical Journal*, 111: 152-61.

834   Mayer, Matthias P., Stefan Rüdiger, and Bernd Bukau. 2000. 'Molecular Basis for Interactions of
835   the DnaK Chaperone with Substrates', *Biological Chemistry*, 381: 877-85.

Meacci, G., J. Ries, E. Fischer-Friedrich, N. Kahya, P. Schwille, and K. Kruse. 2006. 'Mobility
of Min-proteins in *Escherichia coli* measured by fluorescence correlation spectroscopy',
Physical Biology, 3: 255-63.

Mika, J. T., G. van den Bogaart, L. Veenhoff, V. Krasnikov, and B. Poolman. 2010. 'Molecular
sieving properties of the cytoplasm of *Escherichia coli* and consequences of osmotic
stress', *Molecular Microbiology*, 77: 200-7.

Mika, Jacek T., and Bert Poolman. 2011. 'Macromolecule diffusion and confinement in
prokaryotic cells', *Current Opinion in Biotechnology*, 22: 117-26.

Montero Llopis, Paula, Oleksii Sliusarenko, Jennifer Heinritz, and Christine Jacobs-Wagner.
2012. '*In vivo* biochemistry in bacterial cells using FRAP: insight into the translation
cycle', *Biophysical Journal*, 103: 1848-59.

Mu, Xin, Seongil Choi, Lisa Lang, David Mowray, Nikolay V. Dokholyan, Jens Danielsson, and
Mikael Oliveberg. 2017. 'Physicochemical code for quinary protein interactions in
Escherichia coli', *Proceedings of the National Academy of Sciences*, 114: E4556.

Mullineaux, C. W., A. Nenninger, N. Ray, and C. Robinson. 2006. 'Diffusion of green
fluorescent protein in three cell environments in *Escherichia coli*', *Journal of
Bacteriology*, 188: 3442-8.

Muramatsu, N., and A. P. Minton. 1988. 'Tracer diffusion of globular proteins in concentrated
protein solutions', *Proceedings of the National Academy of Sciences of the United States
of America*, 85: 2984-88.

Nenninger, A., G. Mastroianni, and C. W. Mullineaux. 2010. 'Size dependence of protein
diffusion in the cytoplasm of *Escherichia coli*', *Journal of Bacteriology*, 192: 4535-40.

Oleksiuk, O., V. Jakovljevic, N. Vladimirov, R. Carvalho, E. Paster, W. S. Ryu, Y. Meir, N. S.
Wingreen, M. Kollmann, and V. Sourjik. 2011. 'Thermal robustness of signaling in
bacterial chemotaxis', *Cell*, 145: 312-21.

Ouzounov, N., J. P. Nguyen, B. P. Bratton, D. Jacobowitz, Z. Gitai, and J. W. Shaevitz. 2016.
'MreB Orientation Correlates with Cell Diameter in *Escherichia coli*', *Biophysical*
Journal, 111: 1035-43.

Parry, B. R., I. V. Surovtsev, M. T. Cabeen, C. S. O'Hern, E. R. Dufresne, and C. Jacobs-
Wagner. 2014. 'The bacterial cytoplasm has glass-like properties and is fluidized by
metabolic activity', *Cell*, 156: 183-94.

Pédelacq, Jean-Denis, Stéphanie Cabantous, Timothy Tran, Thomas C. Terwilliger, and Geoffrey
S. Waldo. 2006. 'Engineering and characterization of a superfolder green fluorescent
protein', *Nature Biotechnology*, 24: 79-88.

Perrin, Jean. 1909. 'Mouvement brownien et réalité moléculaire', *Annales de chimie et de*
physique.

Persson, Laura B., Vardhaan S. Ambati, and Onn Brandman. 2020. 'Cellular Control of Viscosity
Counters Changes in Temperature and Energy Availability', *Cell*, 183: 1572-85.e16.

Petrov, E. P., T. Ohrt, R. G. Winkler, and P. Schwille. 2006. 'Diffusion and segmental dynamics
of double-stranded DNA', *Physical Review Letters*, 97: 258101.

Potma, Eric O., Wim P. de Boeij, Leonard Bosgraaf, Jeroen Roelofs, Peter J. M. van Haastert,
and Douwe A. Wiersma. 2001. 'Reduced Protein Diffusion Rate by Cytoskeleton in
Vegetative and Polarized Dictyostelium Cells', *Biophysical Journal*, 81: 2010-19.

Rivas, Germán, and Allen P. Minton. 2016. 'Macromolecular Crowding *In Vitro*, *In Vivo*, and In
Between', *Trends in Biochemical Sciences*, 41: 970-81.

Rocha, J., J. Corbitt, T. Yan, C. Richardson, and A. Gahlmann. 2019. 'Resolving Cytosolic
Diffusive States in Bacteria by Single-Molecule Tracking', *Biophysical Journal*, 116:
1970-83.

Roman-Hernandez, G., J. Y. Hou, R. A. Grant, R. T. Sauer, and T. A. Baker. 2011. 'The ClpS
adaptor mediates staged delivery of N-end rule substrates to the AAA+ ClpAP protease',
Molecular Cell, 43: 217-28.

Rotter, D. A. O., C. Heger, L. M. Oviedo-Bocanegra, and P. L. Graumann. 2021. 'Transcription-
dependent confined diffusion of enzymes within subcellular spaces of the bacterial
cytoplasm', *BMC Biology*, 19: 183.

Sabri, Adal, Xinran Xu, Diego Krapf, and Matthias Weiss. 2020. 'Elucidating the Origin of
Heterogeneous Anomalous Diffusion in the Cytoplasm of Mammalian Cells', *Physical*
Review Letters, 125: 058101.

Sanamrad, A., F. Persson, E. G. Lundius, D. Fange, A. H. Gynna, and J. Elf. 2014. 'Single-
particle tracking reveals that free ribosomal subunits are not excluded from the
Escherichia coli nucleoid', *Proceedings of the National Academy of Sciences of the*
United States of America, 111: 11413-8.

Sandikci, A., F. Gloge, M. Martinez, M. P. Mayer, R. Wade, B. Bukau, and G. Kramer. 2013.
'Dynamic enzyme docking to the ribosome coordinates N-terminal processing with
polypeptide folding', *Nature Structural & Molecular Biology*, 20: 843-50.

Saxton, M. J. 1996. 'Anomalous diffusion due to binding: a Monte Carlo study', *Biophysical*
Journal, 70: 1250-62.

Saxton, Michael J. 2007. 'A Biological Interpretation of Transient Anomalous Subdiffusion. I.
Qualitative Model', *Biophysical Journal*, 92: 1178-91.

Schavemaker, P. E., W. M. Smigiel, and B. Poolman. 2017. 'Ribosome surface properties may
impose limits on the nature of the cytoplasmic proteome', *Elife*, 6.

Schavemaker, Paul E., Arnold J. Boersma, and Bert Poolman. 2018. 'How Important Is Protein
Diffusion in Prokaryotes?', *Frontiers in Molecular Biosciences*, 5: 93.

Schwartz, Russell, Claire S. Ting, and Jonathan King. 2001. 'Whole Proteome pI Values
Correlate with Subcellular Localizations of Proteins for Organisms within the Three
Domains of Life', *Genome Research*, 11: 703-09.

Serra, D. O., A. M. Richter, G. Klauck, F. Mika, and R. Hengge. 2013. 'Microanatomy at cellular
resolution and spatial order of physiological differentiation in a bacterial biofilm', *mBio*,
4: e00103-13.

Sinensky, M. 1974. 'Homeoviscous adaptation--a homeostatic process that regulates the viscosity
of membrane lipids in *Escherichia coli*', *Proceedings of the National Academy of*
Sciences of the United States of America, 71: 522-25.

Smilgies, Detlef- M., and Ewa Folta-Stogniew. 2015. 'Molecular weight-gyration radius relation
of globular proteins: a comparison of light scattering, small-angle X-ray scattering and
structure-based data', *Journal of applied crystallography*, 48: 1604-06.

Solbiati, José, Anne Chapman-Smith, Judith L. Miller, Charles G. Miller, and John E. Cronan.
1999. 'Processing of the N termini of nascent polypeptide chains requires deformylation
prior to methionine removal', *Journal of Molecular Biology*, 290: 607-14.

Spitzer, Jan, and Bert Poolman. 2013. 'How crowded is the prokaryotic cytoplasm?', *FEBS*
Letters, 587: 2094-98.

Stasevich, Timothy J., Florian Mueller, Ariel Michelman-Ribeiro, Tilman Rosales, Jay R.
Knutson, and James G. McNally. 2010. 'Cross-validating FRAP and FCS to quantify the
impact of photobleaching on in vivo binding estimates', *Biophysical Journal*, 99: 3093-
101.

Stracy, M., J. Schweizer, D. J. Sherratt, A. N. Kapanidis, S. Uphoff, and C. Lesterlin. 2021.
'Transient non-specific DNA binding dominates the target search of bacterial DNA-
binding proteins', *Molecular Cell*, 81: 1499-514 e6.

Swaminathan, R., C. P. Hoang, and A. S. Verkman. 1997. 'Photobleaching recovery and
anisotropy decay of green fluorescent protein GFP-S65T in solution and cells:
cytoplasmic viscosity probed by green fluorescent protein translational and rotational
diffusion', *Biophysical Journal*, 72: 1900-07.

Tabaka, Marcin, Tomasz Kalwarczyk, Jedrzej Szymanski, Sen Hou, and Robert Holyst. 2014.
'The effect of macromolecular crowding on mobility of biomolecules, association
kinetics, and gene expression in living cells', *Frontiers in Physics*, 2.

Tran, Buu Minh, Haritha Prabha, Aditya Iyer, Conor O'Byrne, Tjakko Abee, and Bert Poolman.
2021. 'Measurement of Protein Mobility in *Listeria monocytogenes* Reveals a Unique
Tolerance to Osmotic Stress and Temperature Dependence of Diffusion', *Frontiers in*
Microbiology, 12.

Tyn, Myo T., and Todd W. Gusek. 1990. 'Prediction of diffusion coefficients of proteins',
Biotechnology and Bioengineering, 35: 327-38.

van den Berg, Jonas, Arnold J. Boersma, and Bert Poolman. 2017. 'Microorganisms maintain
crowding homeostasis', *Nature Reviews Microbiology*, 15: 309-18.

Wachsmuth, M., C. Conrad, J. Bulkescher, B. Koch, R. Mahen, M. Isokane, R. Pepperkok, and J.
Ellenberg. 2015. 'High-throughput fluorescence correlation spectroscopy enables analysis
of proteome dynamics in living cells', *Nature Biotechnology*, 33: 384-9.

Wlodarski, Michal, Bianca Raciti, Jurij Kotar, Marco Cosentino Lagomarsino, Gillian M. Fraser,
and Pietro Cicuta. 2017. 'Both genome and cytosol dynamics change in *E. coli* challenged
with sublethal rifampicin', *Physical Biology*, 14: 015005.

Zhang, Yifei, and Henry Hess. 2019. 'Enhanced Diffusion of Catalytically Active Enzymes', *ACS*
Central Science, 5: 939-48.

Zimmerman, Steven B., and Stefan O. Trach. 1991. 'Estimation of macromolecule concentrations
and excluded volume effects for the cytoplasm of *Escherichia coli*', *Journal of Molecular*
Biology, 222: 599-620.

Supplementary files

This article contains the following supplementary files:

Figure 1 – supplementary table 1

Figure 1 – supplementary table 2

Figure 1 – figure supplement 1

Figure 1 – figure supplement 2

Figure 1 – figure supplement 3

Figure 1 – figure supplement 4

Figure 1 – figure supplement 5

Figure 1 – figure supplement 6

Figure 1 – figure supplement 7

Figure 2 – figure supplement 1

Figure 2 – figure supplement 2

Figure 2 – figure supplement 3

Figure 4 – figure supplement 1

Figure 4 – figure supplement 2

Figure 4 – figure supplement 3

Appendix 1: Notes on the acquisition and analysis protocols for FCS measurements in bacterial

cells

Appendix 2: Ornstein-Uhlenbeck model for confinement effect in FCS measurements

Appendix 3: Effective diffusion coefficient of two linked proteins

Appendix 4: Exact p-values for each figure

Appendix 5: Numerosity of the constructs and conditions for each experiment

Source data: Source data files and raw immunoblot images.

Supplementary figures

984 **Figure 1 – supplementary table 1. sfGFP-tagged library of cytoplasmic proteins**

Protein name	Molecular mass of sfGFP-construct	Biological function	IPTG concentration for FCS (for FRAP)	Number of cells (FCS)	τ_D (μs; avg. ± SE)	α (Avg. ± SE)	Diffusion coefficient, FCS (μm²/s, Avg. ± SE)	Number of cells (FRAP)	Diffusion coefficient, FRAP (μm²/s, Avg. ± SE)
sfGFP	26.9	-	5 μM (15 μM)	42	580 ± 15	0.86 ± 0.01	14.4 ± 0.4	11	11.3 ± 1.3
YggX	39.2	Probable Fe (2+)-trafficking protein	5 μM (5 μM)	8	611 ± 19	0.85 ± 0.01	12.9 ± 0.4	10	9.4 ± 1.6
ClpS	39.2	ATP-dependent Clp protease adapter protein	0 μM	11	1054 ± 33	0.75 ± 0.01			
FolK	45.1	2-amino-4-hydroxy-6-hydroxymethyldihydropteridine pyrophosphokinase	0 μM	8	734 ± 24	0.87 ± 0.01	11.6 ± 0.4		
Crr	45.2	Component of glucose-specific phosphotransferase enzyme IIA	0 μM	14	1065 ± 36	0.87 ± 0.01			
UbiC	45.7	Chorismate pyruvate-lyase	15 μM	14	1140 ± 58	0.87 ± 0.01			
ThpR	46.9	RNA 2'3'-cyclic phosphodiesterase			Discarded due to instability of sfGFP-construct				
CoaE	49.6	Dephospho-CoA kinase	0 μM	11	854 ± 47	0.87 ± 0.01	9.8 ± 0.6		
Adk	50.6	Adenylate kinase	5 μM (15 μM)	23	802 ± 26	0.88 ± 0.00	10.6 ± 0.4	16	9.8 ± 1.5
Cmk	51.7	Cytidylate kinase	5 μM	16	1163 ± 58	0.87 ± 0.01			
NagD	54.1	Ribonucleotide monophosphatase			Discarded due to non-uniform protein localization				
KdsB	54.6	3-deoxy-manno-octulosonate cytidylyltransferase	0 μM	11	1659 ± 70	0.84 ± 0.01			
Map	56.3	Methionine aminopeptidase	0 μM	20	1830 ± 78	0.81 ± 0.01			
MmuM	60.4	Homocysteine S-methyltransferase	5 μM	14	2241 ± 138	0.73 ± 0.01			
RihA	60.8	Pyrimidine-specific ribonucleoside hydrolase			Discarded due to non-uniform protein localization				
PanE	60.8	2-dehydropantoate 2-reductase	0 μM (5 μM)	18	1059 ± 26	0.85 ± 0.01	7.8 ± 0.2	11	5.2 ± 0.6
SolA	67.9	N-methyl-L-tryptophan oxidase	0 μM	7	795 ± 31	0.82 ± 0.01	9.9 ± 0.5		
Pgk	68.1	Phosphoglycerate kinase	0 μM	16	991 ± 41	0.90 ± 0.01	8.6 ± 0.3		
EntC	69.9	Isochorismate synthase	15 μM	15	1777 ± 119	0.82 ± 0.01	8.7 ± 0.7		
AroA	73.1	3-phosphoshikimate 1-carboxyvinyltransferase	0 μM	9	995 ± 69	0.86 ± 0.01	8.7 ± 0.7		
ThrC	74.1	Threonine synthase	0 μM	14	908 ± 28	0.87 ± 0.01	9.1 ± 0.3		
MurF	74.4	UDP-N-acetylmuramoyl-tripeptide--D-alanyl-D-alanine ligase	0 μM	7	1008 ± 76	0.85 ± 0.02	8.3 ± 0.7		
DsdA	74.9	D-serine dehydratase	0 μM	14	1017 ± 53	0.89 ± 0.01	8.4 ± 0.4	10	7.8 ± 0.7
HemN	79.7	Oxygen-independent coproporphyrinogen III oxidase	0 μM	13	1262 ± 54	0.86 ± 0.01	6.7 ± 0.4		
PrpD	80.9	2-methylcitrate dehydratase	0 μM	12	1866 ± 140	0.84 ± 0.01			
DnaK	96.0	Molecular chaperone	5 μM	10	2296 ± 78	0.76 ± 0.01			
MalZ	96.0	Maltodextrin glucosidase	0 μM	9	3725 ± 229	0.77 ± 0.01			
GlcB	107.5	Malate synthase G	5 μM (15 μM)	16	1315 ± 45	0.86 ± 0.01	6.4 ± 0.2	10	6.7 ± 1.1
MetE	111.7	5-methyltetrahydropteroyltriglutamate--homocysteine methyltransferase	0 μM	8	1137 ± 53	0.87 ± 0.01	7.4 ± 0.3		
LeuS	124.2	Leucine--tRNA ligase	0 μM	14	1637 ± 75	0.86 ± 0.01	5.1 ± 0.2		
AcnA	124.7	Aconitate hydratase A	5 μM (15 μM)	19	1415 ± 56	0.86 ± 0.01	6.1 ± 0.2	10	4.3 ± 0.4
MetH	163.0	Methionine synthase	0 μM (5 μM)	9	1402 ± 45	0.81 ± 0.01	5.8 ± 0.1	15	4.0 ± 0.5

Figure 1 – supplementary table 2. Strains and plasmids used in this study

Strain or plasmid	Relevant genotype or phenotype	Reference or source
	Strains	
E. coli W3110	W3110 derivative with functional RpoS [rpoS396(Am)]	(Serra Diego et al.)
NB63	W3110 Δ*clpA*	This work
	Plasmids	
pTrc99A	Amp^r; expression vector; pBR ori; *trc* promoter, IPTG inducible	(Amann, Ochs, and Abel 1988)
pCP20	Amp^r, Cam^r; *flp*	(Cherepanov and Wackernagel 1995)
pNB1	Amp^r; *sfGFP* in pTrc99A	This work
pNB3	Amp^r; *Adk-sfGFP* in pTrc99A	This work
pNB4	Amp^r; *CoaE-sfGFP* in pTrc99A	This work
pNB5	Amp^r; *Cmk-sfGFP* in pTrc99A	This work
pNB6	Amp^r; *Pgk-sfGFP* in pTrc99A	This work
pNB7	Amp^r; *MmuM-sfGFP* in pTrc99A	This work
pNB8	Amp^r; *PrpD-sfGFP* in pTrc99A	This work
pNB9	Amp^r; *DsdA-sfGFP* in pTrc99A	This work
pNB11	Amp^r; *GlcB-sfGFP* in pTrc99A	This work
pNB13	Amp^r; *HemN-sfGFP* in pTrc99A	This work
pNB14	Amp^r; *Map^{WT}-sfGFP* in pTrc99A	This work
pNB15	Amp^r; *ThrC-sfGFP* in pTrc99A	This work
pNB16	Amp^r; *MalZ-sfGFP* in pTrc99A	This work
pNB17	Amp^r; *EntC-sfGFP* in pTrc99A	This work
pNB18	Amp^r; *ThpR-sfGFP* in pTrc99A	This work
pNB19	Amp^r; *AroA-sfGFP* in pTrc99A	This work
pNB20	Amp^r; *ClpS^{WT}-sfGFP* in pTrc99A	This work
pNB21	Amp^r; *Crr-sfGFP* in pTrc99A	This work
pNB22	Amp^r; *KdsB-sfGFP* in pTrc99A	This work
pNB23	Amp^r; *LeuS-sfGFP* in pTrc99A	This work
pNB24	Amp^r; *MurF-sfGFP* in pTrc99A	This work
pNB25	Amp^r; *NagD-sfGFP* in pTrc99A	This work
pNB26	Amp^r; *RihA-sfGFP* in pTrc99A	This work
pNB27	Amp^r; *SolA-sfGFP* in pTrc99A	This work
pNB28	Amp^r; *UbiC-sfGFP* in pTrc99A	This work
pNB29	Amp^r; *PanE-sfGFP* in pTrc99A	This work
pNB30	Amp^r; *FolK-sfGFP* in pTrc99A	This work
pNB39	Amp^r; *AcnA-sfGFP* in pTrc99A	This work
pNB40	Amp^r; *MetE-sfGFP* in pTrc99A	This work
pNB42	Amp^r; *MetH-sfGFP* in pTrc99A	This work
pNB44	Amp^r; *YggX-sfGFP* in pTrc99A	This work
pNB45	Amp^r; *Adk$^{C.c.}$-sfGFP* in pTrc99A	This work
pNB46	Amp^r; *Adk$^{V.c.}$-sfGFP* in pTrc99A	This work
pNB47	Amp^r; *Adk$^{M.x.}$-sfGFP* in pTrc99A	This work
pNB48	Amp^r; *AcnA$^{M.x.}$-sfGFP* in pTrc99A	This work
pNB49	Amp^r; *AcnA$^{V.c.}$-sfGFP* in pTrc99A	This work
pNB51	Amp^r; *Adk$^{B.s.}$-sfGFP* in pTrc99A	This work
pNB52	Amp^r; *AcnA$^{B.s.}$-sfGFP* in pTrc99A	This work
pNB54	Amp^r; *Adk$^{Y.e.}$-sfGFP* in pTrc99A	This work
pNB56	Amp^r; *Pgk$^{C.c.}$-sfGFP* in pTrc99A	This work
pNB58	Amp^r; *Pgk$^{V.c.}$-sfGFP* in pTrc99A	This work
pNB59	Amp^r; *Pgk$^{M.x.}$-sfGFP* in pTrc99A	This work
pNB60	Amp^r; *AcnA$^{Y.e.}$-sfGFP* in pTrc99A	This work
pNB61	Amp^r; *DnaK^{WT}-sfGFP* in pTrc99A	This work
pNB62	Amp^r; *Map$^{K211E\_K218E\_K224E\_K226E}$-sfGFP* in pTrc99A	This work
pNB63	Amp^r; *Map$^{K211A\_K218A\_K224A\_K226A}$-sfGFP* in pTrc99A	This work
pNB64	Amp^r; *DnaK^{V436F}-sfGFP* in pTrc99A	This work
pNB66	Amp^r; *ClpS$^{D35A\_D36A\_H66A}$-sfGFP* in pTrc99A	This work

 Figure 1 – figure supplement 1. Expression analysis for all *E. coli* protein constructs made in this study.

990 Expression of proteins was analyzed by SDS-PAGE and immunoblotting using a primary antibody specific for GFP.

991 All fusion proteins displayed a dominant band corresponding to the expected molecular mass of the full-length fusion

992 (Figure 1 – supplementary table 1).

993 **Figure 1 – figure supplement 1 – source data 1-10.** Raw images of immunoblots used for Figure 1 – figure

994 supplement 1.

Measure	τ_D (µs)	α
R1	536	0.79
R2	535	0.81
R3	517	0.84
R4	595	0.88
R5	531	0.81
R6	605	0.90
Avg. (±SD)	553 (±34)	0.84 (±0.04)

Figure 1 – figure supplement 2. Workflow of a typical FCS experiment. The focus of the confocal microscope is positioned near one pole in the cell of interest, and six subsequent acquisitions of the fluorescence intensity of 20 seconds each are performed. The example for sfGFP shows only traces for R1, R3 and R5 measurements (different colors). The autocorrelation functions (ACFs) are then calculated independently for each acquisition. The values for τ_D and α are extracted independently from the fit to each ACF using the anomalous diffusion model (solid lines) and then averaged to obtain the τ_D and α for each individual cell.

1002

1003 **Figure 1 – figure supplement 3. Comparison between fits of the experimental data with Brownian and**

1004 **anomalous diffusion models.** The experimental data (here the example for the R3 measurement from Figure 1 –

1005 figure supplement 2) were fitted by the model of free Brownian diffusion and by the anomalous diffusion model as

1006 indicated by different colors (upper panel), with the corresponding values of residuals (lower panel).

1007

Figure 1 – figure supplement 4. Individual measurements of τ_D and α for all *E. coli* protein constructs included

in the analysis of mass dependence. Each dot in the box plot represents the values of τ_D (A) and α (B) for one

individual cell. The averages values calculated from these datasets are shown in Figure 1.

Figure 1 – figure supplement 4 – source data 1. Original values used for plots in Figure 1 – figure supplement 4.

1012

Figure 1 – figure supplement 5. Expression analysis for the mutants with impaired interactions. The expression

of indicated point mutants of ClpS, DnaK and Map was analyzed by SDS-PAGE and immunoblotting using a primary

antibody specific for GFP. All mutants displayed a dominant band corresponding to the expected molecular mass of

the full-length fusion, and comparable to that of the wild-type counterpart. ClpS$^{D35A\_D36A\_H66A}$-sfGFP was measured

in the same $\Delta clpA$ background as subsequently used for the FCS experiments.

Figure 1 – figure supplement 5 – source data 1-2. Raw images of immunoblots used for Figure 1 – figure supplement

5.

1020

1021 **Figure 1 – figure supplement 6. Mobility of ClpS, Map and DnaK and of indicated mutants with disrupted**

1022 **protein interactions.** Each dot in the box plot represents the values of τ_D (A) and α (B) for one individual cell. ClpS

1023 mutant was measured in *ΔclpA* background. *** p<0.0001; * p<0.05; NS: no statistically significant difference in a

1024 two-tailed heteroscedastistic *t*-test.

1025 **Figure 1 – figure supplement 6 – source data 1.** Original values used for plots in Figure 1 – figure supplement 6.

1026

Figure 1 – figure supplement 7. Mobility of homologous proteins from other bacterial species in *E. coli*. The

mobility (A) and anomalous diffusion exponent (B) of sfGFP fusions to homologues of Adk, Pgk and AcnA from

indicated bacterial species (*E.c. = Escherichia coli; Y.e. = Yersinia enterocolitica; V.c. = Vibrio cholerae; C.c. =

Caulobacter crescentus; M.x. = Myxococcus xanthus; B.s. = Bacillus subtilis) compared with that of their counterpart

from *E. coli*. *** $p<0.0001$; ** $p<0.001$; * $p<0.05$ in a two-tailed heteroscedastistic *t*-test. When not indicated, no

statistically significant difference is observed. (C, D) Mass dependence of diffusion parameters of these sfGFP-

constructs. Each symbol represents the average value for all individual cells that have been measured for that construct

and the error bars represent the standard error of the mean.

Figure 1 – figure supplement 7 – source data 1. Original values used for plots in Figure 1 – figure supplement 7.

1036

1037 **Figure 2 – figure supplement 1. Residuals of fitting the simulated ACFs with different models.** Simulated

1038 fluorescence intensity ACF data were fitted by the models of anomalous and Brownian diffusion, as well as by the

1039 OU model of Brownian diffusion under confinement, as indicated.

1040

Figure 2 – figure supplement 2. Diffusion coefficients fitted from simulation data. Diffusion coefficients

computed ($D = \omega_0^2/4\tau_D$) from the diffusion times extracted from the fit of the Brownian simulation data with (A) the

anomalous diffusion model or (B) the Ornstein-Uhlenbeck (OU) model of Brownian diffusion under confinement at

various value of the ansatz D and of the cell diameter d, normalized by the ansatz D. The grey areas represent ± 5%

accuracy.

Figure 2 – figure supplement 2 – source data 1. Original values used for plots in Figure 2 – figure supplement 2.

1047

1048 **Figure 2 – figure supplement 3. Comparison between fits of the experimental data with confined diffusion and**

1049 **anomalous diffusion models.** The experimental data (here the example for the R3 measurement from Figure 1 –

1050 figure supplement 2) were fitted by the model of confined diffusion and by the anomalous diffusion model as indicated

1051 by different colors (upper panel), yielding comparable values of residuals (lower panel).

1052

1053 **Figure 4 – figure supplement 1. Effect of osmotic upshift on protein mobility.** The effect of osmotic upshift on

1054 sfGFP mobility, measured in cells that were either resuspended in tethering buffer (ionic strength of 105 mM) or in

1055 the same buffer but supplemented with additional 100 mM NaCl (total ionic strength of 305 mM). *** p<0.0001 in a

1056 two-tailed heteroscedastistic *t*-test.

1057 **Figure 4 – figure supplement 1 – source data 1.** Original values used for plots in Figure 4 – figure supplement 1.

1058

Figure 4 – figure supplement 2. Effect of growth and measurement temperature on protein diffusion. Mobility

for sfGFP in cells grown either at 25°C or at 37°C and measured either at 25°C or at 35°C, as in Figure 4A. ***

$p<0.0001$; ** $p<0.001$; NS: no statistically significant difference in a two-tailed heteroscedastistic *t*-test.

Figure 4 – figure supplement 2 – source data 1. Original values used for plots in Figure 4 – figure supplement 2.

1063

1064 **Figure 4 – figure supplement 3. Mass dependence of the effect of antibiotics on protein diffusion.** Data from

1065 Figure 4B were plotted as a function of molecular mass for indicated protein constructs and treatments. Error bars

1066 represent the standard error of the mean.

1067 **Figure 4 – figure supplement 3 – source data 1.** Original values used for plots in Figure 4 – figure supplement 3.

Appendix 1

Notes on the acquisition and analysis protocols for FCS measurements in bacterial cells

Due to the limited size of bacterial cells, FCS measurements require precise positioning of the confocal volume in the bacterial cytoplasm and minimization of the photobleaching-induced effects. In order to ensure that, before fitting an autocorrelation function, we verified the stability of the lateral (xy) positioning of the observation volume by visually analyzing for lateral drifts in confocal images acquired immediately before and after the FCS acquisition. This was done by annotating the xy position in the pre-aquisition image and verifying that the positioning did not change in the post-aquisition image after 120 seconds. Measurements showing xy drift were excluded from the analysis (Appendix 1 - figure 1). Furthermore, the focal stability of the sample was increased by thermal equilibration on the microscope stage before measurements.

Appendix 1 - figure 1. Typical examples of presence or absence of lateral focal drift during FCS measurements. Substantial lateral drift could be observed for <10% of experiments (upper images), whereas most measurement showed no perceptible lateral drift (lower images).

Long-term photobleaching due to the progressive decrease of the total number of fluorescent

proteins during FCS experiments (Appendix 1 - figure 2) is unavoidable due to the small volume

of *E. coli* cells, and it requires correction to avoid artifacts. We observed that almost identical ACFs

were obtained when correcting for the photobleaching using either multi-segment detrending (Jay

Unruh, https://research.stowers.org/imagejplugins/index.html, Stowers Institute for Medical

Research, USA) or a local averaging approach (Wachsmuth et al. 2015) (Appendix 1 - figure 3).

Appendix 1 - figure 2. Typical traces of fluorescence intensity during FCS measurements. Examples of

fluorescence intensity traces for indicated protein fusions. The vertical red dashed lines separate sequential

fluorescence intensity acquisitions on the same cell.

Appendix 1 - figure 3. Results of detrending with multi-segments and local averaging approaches. Comparison

of experimental ACFs corrected using either multi-segments or local averaging approaches (as indicated) for sfGFP

and Adk-sfGFP and different data acquisition segments (R1 vs R6).

We also confirmed that there was no systematic trend in the fitted values of τ_D and α with the time

of the fluorescence trace acquisition (Appendix 1 - figure 4). An additional process that could

potentially affect autocorrelation functions is short-term photobleaching of the fluorophore in the

confocal volume, also known as cryptic photobleaching, which can artificially accelerate the

decrease of the autocorrelation function and lead to an underestimation of the protein residence

time (Macháň, Foo, and Wohland 2016). This process is different from long-term photobleaching,

which is caused by the continuous illumination in the entire illumination light cone. However, the

effect of cryptic photobleaching was shown to be typically <5%, even for proteins that diffuse 10

to 100 times slower and have higher bleaching rates than our constructs (Stasevich et al. 2010;

Wachsmuth et al. 2015).

Appendix 1 - figure 4. Values of τ_D or α for the six sequential ACFs. Values were determined by fitting the anomalous diffusion model to experimental ACFs for the six sequential time segments per individual cell expressing sfGFP (A) or MetH-sfGFP (B).

48   Appendix 2

49   Ornstein-Uhlenbeck model for confinement effect in FCS measurements

50   We aim to derive the autocorrelation function for an FCS experiment in which the fluorescent

51   particles are confined by a (possibly anisotropic) harmonic potential centered at $(x, y, z) = (0,0,0)$,

52   i.e.

$$53 \qquad V(x, y, z) = \frac{k_x x^2}{2} + \frac{k_y y^2}{2} + \frac{k_z z^2}{2}$$

where k_i represents the stiffness of the potential in each dimension, and thus the extent σ_i of

confinement along that dimension given by $\sigma_i^2 \equiv k_{\mathrm{B}} T / k_i$.

We can treat each dimension independently, using x without loss of generality in what follows.

Diffusion in a harmonic potential is described by the Ornstein-Uhlenbeck process. The

corresponding Green's function $P(x, t | x_0)$, representing the probability of finding a particle at

position x at time t given that it was at position x_0 at time $t = 0$, is

$$60 \qquad P(x, t | x_0) = \frac{1}{\sqrt{2\pi\sigma_x^2 \left(1 - e^{-\frac{2Dt}{\sigma_x^2}}\right)}} \exp\left[-\frac{1}{2\sigma_x^2} \frac{\left(x - x_0 e^{-\frac{Dt}{\sigma_x^2}}\right)^2}{1 - e^{-\frac{2Dt}{\sigma_x^2}}}\right]$$

61   at long times, $t \to \infty$, we recover the stationary state given by the Boltzmann distribution

62   corresponding to the harmonic trap

$$63 \qquad P_{\mathrm{st}}(x) = \frac{1}{\sqrt{2\pi\sigma_x^2}} \exp\left[-\frac{x^2}{2\sigma_x^2}\right].$$

The autocorrelation function (ACF) $G(t)$ of an FCS measurement is given as the multiple integral

over the product of the probability to detect a photon from a molecule at some initial position x_0,

the probability density that it diffuses from this position to a final position x within time t (given

by Green's function), and the probability to detect a photon from a molecule at this final position

(Enderlein et al. 2005; Enderlein 2012). Note that the probability of detection of a molecule will

necessarily be proportional to the intensity of the laser beam, which we can assume Gaussian and

also centered at $(x, y, z) = (0,0,0)$, with the usual form

71
$$I(x, y, z) = I_0 \exp\left(-\frac{2x^2}{\omega_0^2}\right) \exp\left(-\frac{2y^2}{\omega_0^2}\right) \exp\left(-\frac{2z^2}{S^2\omega_0^2}\right) \equiv I_0 I_x(x) I_y(y) I_z(z)$$

72   where ω_0 is the width of the (circular) laser beam along the x and y directions, and S is a

73   dimensionless factor accounting for the anisotropy along the z direction, i.e. the axial direction of

74   the beam.

75   Ignoring constant normalization factors and baselines, the time dependent part of the

76   autocorrelation function is then given by $G(t) = G_x(t) G_y(t) G_z(t)$ with

77
$$G_x(t) = \int dx \int dx_0 I_x(x) P(x, t|x_0) I_x(x_0) P_{st}(x_0)$$

78   which can be directly integrated to give, after normalizing so that $G_x(t = 0) = 1$, the expression

79
$$G_x(t) = \left[1 + 2\frac{\sigma_x^2}{\omega_0^2}\frac{1 - e^{-\frac{2Dt}{\sigma_x^2}}}{1 + \frac{1}{8}\frac{\omega_0^2}{\sigma_x^2}}\right]^{-\frac{1}{2}} . \quad (A2\text{-}1)$$

80   Note that, in the limit of no confinement, $\sigma_x^2 \to \infty$, this equation reduces to the well-known ACF

81   for unconfined diffusion

82
$$G_x(t) = \left[1 + \frac{4Dt}{\omega_0^2}\right]^{-\frac{1}{2}} = \left[1 + \frac{t}{\tau_D}\right]^{-\frac{1}{2}}$$

83 where we have defined the diffusion time $\tau_D \equiv \omega_0^2/(4D)$. With this definition, the ACF <u>in</u> Eq.

84 (A2-1) can be rewritten as

85
$$G_x(t) = \left[1 + 2\frac{\sigma_x^2}{\omega_0^2}\frac{1 - e^{-\frac{1}{2}\frac{\omega_0^2}{\sigma_x^2}\frac{t}{\tau_D}}}{1 + \frac{1}{8}\frac{\omega_0^2}{\sigma_x^2}}\right]^{-\frac{1}{2}}.$$

86 The full three-dimensional ACF is then, in general,

87
$$G(t) = \left[1 + 2\frac{\sigma_x^2}{\omega_0^2}\frac{1 - e^{-\frac{1}{2}\frac{\omega_0^2}{\sigma_x^2}\frac{t}{\tau_D}}}{1 + \frac{1}{8}\frac{\omega_0^2}{\sigma_x^2}}\right]^{-\frac{1}{2}}\left[1 + 2\frac{\sigma_y^2}{\omega_0^2}\frac{1 - e^{-\frac{1}{2}\frac{\omega_0^2}{\sigma_y^2}\frac{t}{\tau_D}}}{1 + \frac{1}{8}\frac{\omega_0^2}{\sigma_y^2}}\right]^{-\frac{1}{2}}\left[1 + 2\frac{\sigma_z^2}{S^2\omega_0^2}\frac{1 - e^{-\frac{1}{2}\frac{\omega_0^2}{\sigma_z^2}\frac{t}{\tau_D}}}{1 + \frac{1}{8}\frac{S^2\omega_0^2}{\sigma_z^2}}\right]^{-\frac{1}{2}}.$$

88 The cylindrical geometry of a bacterium can be approximated by an infinite cylinder along the y

89 direction, so that $\sigma_x = \sigma_z = \sigma$ and $\sigma_y \to \infty$, resulting in the ACF

90
$$G(t) = \left[1 + 2\frac{\sigma^2}{\omega_0^2}\frac{1 - e^{-\frac{1}{2}\frac{\omega_0^2}{\sigma^2}\frac{t}{\tau_D}}}{1 + \frac{1}{8}\frac{\omega_0^2}{\sigma^2}}\right]^{-\frac{1}{2}}\left[1 + \frac{t}{\tau_D}\right]^{-\frac{1}{2}}\left[1 + 2\frac{\sigma^2}{S^2\omega_0^2}\frac{1 - e^{-\frac{1}{2}\frac{\omega_0^2}{\sigma^2}\frac{t}{\tau_D}}}{1 + \frac{1}{8}\frac{S^2\omega_0^2}{\sigma^2}}\right]^{-\frac{1}{2}}. \quad (A2\text{-}2)$$

91 Eq. (A2-2), with the added baseline and multiplicative correction accounting for particles in the

92 non-fluorescent state, corresponds to Eq. (3) in the main text.

94 Appendix 3

95 Effective diffusion coefficient of two linked proteins

96 In previous work, we studied the diffusion of two spherical objects with radii a_1 and a_2, joined

97 together by a flexible linker (Agudo-Canalejo and Golestanian 2020). In the limit of a rigid linker

98 of length ℓ, the effective diffusion coefficient of the composite object goes as:

99
$$D \simeq \frac{k_B T}{6\pi\eta a_1} \frac{a_1}{(a_1+a_2)} \left[1 + 2\frac{a_1 a_2}{(a_1+a_2)(a_1+a_2+\ell)} - \frac{9}{8}\frac{a_1 a_2 (a_1-a_2)^2}{(a_1+a_2)^2 (a_1+a_2+\ell)^2} \right] \quad \text{(A3-1)}$$

plus higher order correction terms of order $O\left(\frac{a_i^3}{(a_1+a_2+\ell)^3}\right)$.

We can then consider what is the effective diffusion coefficient of two proteins that are linked to

each other. For that, we first need to connect the molecular mass to the effective radius of the

protein. If we identify subunit 1 with GFP, and subunit 2 with the protein attached to it, and we call

M_{GFP} the molecular mass of GFP and M_{tot} the total molecular mass (sum of GFP and the protein),

we expect relations of the form

106
$$a_1 = C M_{GFP}^\beta$$

107
$$a_2 = C(M_{tot} - M_{GFP})^\beta$$

where C is a proportionality constant assumed to be typical for all proteins (which is of order 1,

with values reported in the literature of about 0.65 (Smilgies and Folta-Stogniew 2015), when the

mass is in kDa and the radius is in nm), and β is the scaling exponent introduced in the main text.

For the linker which is made of 6 amino-acids, we may use the typical conversion factor 0.35

nm/amino-acid to estimate $\ell \approx 2$ nm.

Plugging these expressions for a_1 and a_2 into Eq. (A3-1) above, one obtains an expression for the

diffusion coefficient D as a function of M_{tot} that depends only on three parameters (since the

molecular mass of GFP is known): (i) the diffusion coefficient of GFP $\frac{k_B T}{6\pi\eta a_1}$, (ii) the exponent β,

and (iii) the rescaled linker length ℓ/C.

Appendix 4

Exact p-values for all significance analyses

Figure 1C

Testing pair	P-value
sfGFP vs Adk-sfGFP	0.000000010
Adk-sfGFP vs AcnA-sfGFP	0.00000000044

Figure 1 – figure supplement 6A

Testing pair	P-value
ClpS^{WT}-sfGFP vs ClpS$^{D35A\_D36A\_H66A}$-sfGFP	0.000094
Map^{WT}-sfGFP vs Map$^{Lys\rightarrow Ala}$-sfGFP	0.0000012
Map^{WT}-sfGFP vs Map$^{Lys\rightarrow Glu}$-sfGFP	0.000000016
Map$^{Lys\rightarrow Ala}$-sfGFP vs Map$^{Lys\rightarrow Glu}$-sfGFP	0.10
DnaK^{WT}-sfGFP vs DnaK^{V436F}-sfGFP	0.00023

Figure 1 – figure supplement 6B

Testing pair	P-value
ClpS^{WT}-sfGFP vs ClpS$^{D35A\_D36A\_H66A}$-sfGFP	0.000014
Map^{WT}-sfGFP vs Map$^{Lys\rightarrow Ala}$-sfGFP	0.0092
Map^{WT}-sfGFP vs Map$^{Lys\rightarrow Glu}$-sfGFP	0.065
Map$^{Lys\rightarrow Ala}$-sfGFP vs Map$^{Lys\rightarrow Glu}$-sfGFP	0.37
DnaK^{WT}-sfGFP vs DnaK^{V436F}-sfGFP	0.00000019

Figure 1 – figure supplement 7A

Testing pair	P-value
Adk$^{E.c.}$-sfGFP vs Adk$^{Y.e.}$-sfGFP	0.17
Adk$^{E.c.}$-sfGFP vs Adk$^{V.c.}$-sfGFP	0.056
Adk$^{E.c.}$-sfGFP vs Adk$^{C.c.}$-sfGFP	0.93
Adk$^{E.c.}$-sfGFP vs Adk$^{M.x.}$-sfGFP	0.21
Adk$^{E.c.}$-sfGFP vs Adk$^{B.s.}$-sfGFP	0.23
Pgk$^{E.c.}$-sfGFP vs Pgk$^{V.c.}$-sfGFP	0.75
Pgk$^{E.c.}$-sfGFP vs Pgk$^{C.c.}$-sfGFP	0.26
Pgk$^{E.c.}$-sfGFP vs Pgk$^{M.x.}$-sfGFP	0.33
AcnA$^{E.c.}$-sfGFP vs AcnA$^{Y.e.}$-sfGFP	0.093
AcnA$^{E.c.}$-sfGFP vs AcnA$^{V.c.}$-sfGFP	0.084
AcnA$^{E.c.}$-sfGFP vs AcnA$^{M.x.}$-sfGFP	0.0000000023
AcnA$^{E.c.}$-sfGFP vs AcnA$^{B.s.}$-sfGFP	0.069

129 **Figure 1 – figure supplement 7B**

Testing pair	P-value
Adk$^{E.c.}$-sfGFP vs Adk$^{Y.e.}$-sfGFP	0.000060
Adk$^{E.c.}$-sfGFP vs Adk$^{V.c.}$-sfGFP	0.18
Adk$^{E.c.}$-sfGFP vs Adk$^{C.c.}$-sfGFP	0.042
Adk$^{E.c.}$-sfGFP vs Adk$^{M.x.}$-sfGFP	0.070
Adk$^{E.c.}$-sfGFP vs Adk$^{B.s.}$-sfGFP	0.0029
Pgk$^{E.c.}$-sfGFP vs Pgk$^{V.c.}$-sfGFP	0.11
Pgk$^{E.c.}$-sfGFP vs Pgk$^{C.c.}$-sfGFP	0.0082
Pgk$^{E.c.}$-sfGFP vs Pgk$^{M.x.}$-sfGFP	0.0087
AcnA$^{E.c.}$-sfGFP vs AcnA$^{Y.e.}$-sfGFP	0.035
AcnA$^{E.c.}$-sfGFP vs AcnA$^{V.c.}$-sfGFP	0.16
AcnA$^{E.c.}$-sfGFP vs AcnA$^{M.x.}$-sfGFP	0.083
AcnA$^{E.c.}$-sfGFP vs AcnA$^{B.s.}$-sfGFP	0.00024

131 **Figure 2C**

Testing pair	P-value
Untreated vs A22 treatment	0.00000000000032

133 **Figure 2D**

Testing pair	P-value
Untreated vs A22 treatment	0.0022

135 **Figure 4A**

Testing pair	P-value
sfGFP 25°C vs 35°C	0.000040
Adk-sfGFP 25°C vs 35°C	0.0000025
AcnA-sfGFP 25°C vs 35°C	0.00077

137 **Figure 4B**

Testing pair	P-value
sfGFP Untreated vs Chloramphenicol	0.61
sfGFP DMSO vs Rifampicin	0.012
Adk-sfGFP Untreated vs Chloramphenicol	0.12
Adk-sfGFP DMSO vs Rifampicin	0.00048
Pgk-sfGFP Untreated vs Chloramphenicol	0.17
Pgk-sfGFP DMSO vs Rifampicin	0.0000012
AcnA-sfGFP Untreated vs Chloramphenicol	0.000011
AcnA-sfGFP DMSO vs Rifampicin	0.0085

141   **Figure 4 – figure supplement 1**

Testing pair	P-value
sfGFP ionic strength 105 mM vs 305 mM	0.000027

143   **Figure 4 – figure supplement 2**

Testing pair	P-value
sfGFP Grown 25°C, measured 25°C vs 35°C	0.0020
sfGFP Measured 25°C, grown 25°C vs 37°C	0.060
sfGFP Measured 35°C, grown 25°C vs 37°C	0.98
sfGFP Grown 37°C, measured 25°C vs 35°C	0.000040

145   **Figure 5**

Testing pair	P-value
sfGFP M9 salts vs M9 salts, Cam	0.82
sfGFP M9 salts vs M9 salts, Glu+CA	0.000023
sfGFP M9 salts, Cam vs M9 salts, Cam, Glu+CA	0.019
sfGFP M9 salts, Glu+CA vs M9 salts, Cam, Glu+CA	0.019
AcnA-sfGFP M9 salts vs M9 salts, Cam	0.37
AcnA-sfGFP M9 salts vs M9 salts, Glu+CA	0.0073
AcnA-sfGFP M9 salts, Cam vs M9 salts, Cam, Glu+CA	0.0035
AcnA-sfGFP M9 salts, Glu+CA vs M9 salts, Cam, Glu+CA	0.014

Appendix 5

Numerosity of the constructs and conditions for each experiment

Table 1. Figure 1 and Figure 1 – figure supplement 4 and 6

Construct	*Numerosity (n)*	*Construct*	*Numerosity (n)*
sfGFP	42	Pgk-sfGFP	16
YggX-sfGFP	8	EntC-sfGFP	15
ClpS^{WT}-sfGFP	11	AroA-sfGFP	9
ClpS$^{D35A\_D36A\_H66A}$-sfGFP	10	ThrC-sfGFP	14
FolK-sfGFP	8	MurF-sfGFP	7
Crr-sfGFP	14	DsdA-sfGFP	14
UbiC-sfGFP	14	HemN-sfGFP	13
CoaE-sfGFP	11	PrpD-sfGFP	12
Adk-sfGFP	23	DnaK^{WT}-sfGFP	10
Cmk-sfGFP	16	DnaK^{V436F}-sfGFP	10
KdsB-sfGFP	22	MalZ-sfGFP	9
Map^{WT}-sfGFP	20	GlcB-sfGFP	16
Map$^{Lys\rightarrow Glu}$-sfGFP	10	MetE-sfGFP	8
Map$^{Lys\rightarrow Ala}$-sfGFP	12	LeuS-sfGFP	14
MmuM-sfGFP	14	AcnA-sfGFP	19
PanE-sfGFP	18	MetH-sfGFP	9
SolA-sfGFP	7		

Table 2. Figure 1 – figure supplement 7

Construct	*Numerosity (n)*	*Construct*	*Numerosity (n)*
Adk$^{E.c.}$-sfGFP	Same as table 1	Pgk$^{C.c.}$-sfGFP	5
Adk$^{Y.e.}$-sfGFP	5	Pgk$^{M.x.}$-sfGFP	11
Adk$^{V.c.}$-sfGFP	10	AcnA$^{E.c.}$-sfGFP	Same as table 1
Adk$^{C.c.}$-sfGFP	5	AcnA$^{Y.e.}$-sfGFP	5
Adk$^{M.x.}$-sfGFP	11	AcnA$^{V.c.}$-sfGFP	10
Adk$^{B.s.}$-sfGFP	10	AcnA$^{M.x.}$-sfGFP	10
Pgk$^{E.c.}$-sfGFP	Same as table 1	AcnA$^{B.s.}$-sfGFP	10
Pgk$^{V.c.}$-sfGFP	10		

Table 3. Figure 2C and 2D

Condition	*Numerosity (n)*
Untreated	10
A22-treated	12

165 **Table 4. Figure 3**

Construct	Numerosity (n)	Construct	Numerosity (n)
sfGFP	11	DsdA-sfGFP	10
YggX-sfGFP	10	GlcB-sfGFP	10
Adk-sfGFP	16	AcnA-sfGFP	10
PanE-sfGFP	11	MetH-sfGFP	15

167 **Table 5. Figure 4A**

Construct and condition	Numerosity (n)	Construct and condition	Numerosity (n)
sfGFP, 25°C	See table 1	AcnA-sfGFP, 25°C	See table 1
sfGFP, 35°C	14	AcnA-sfGFP, 35°C	18
Adk-sfGFP, 25°C	See table 1		
Adk-sfGFP, 35°C	21		

169 **Table 5. Figure 4B**

Construct and condition	Numerosity (n)	Construct and condition	Numerosity (n)
sfGFP, untreated	See table 1	Pgk-sfGFP, untreated	See table 1
sfGFP, chloramphenicol	10	Pgk-sfGFP, chloramphenicol	10
sfGFP, DMSO	15	Pgk-sfGFP, DMSO	10
sfGFP, rifampicin	15	Pgk-sfGFP, rifampicin	10
Adk-sfGFP, untreated	See table 1	AcnA-sfGFP, untreated	See table 1
Adk-sfGFP, chloramphenicol	10	AcnA-sfGFP, chloramphenicol	10
Adk-sfGFP, DMSO	10	AcnA-sfGFP, DMSO	10
Adk-sfGFP, rifampicin	10	AcnA-sfGFP, rifampicin	10

171 **Table 6. Figure 4 – figure supplement 1**

Construct and condition	Numerosity (n)
sfGFP, 105 mM ionic strength	See table 1
sfGFP, 305 mM ionic strength	11

173 **Table 6. Figure 4 – figure supplement 2**

Construct and condition	Numerosity (n)
sfGFP, grown 25°C, measured 25°C	10
sfGFP, grown 25°C, measured 35°C	10
sfGFP, grown 37°C, measured 25°C	See table 1
sfGFP, grown 37°C, measured 35°C	See table 5

179 **Table 7. Figure 5**

Construct and condition	Numerosity (n)	Construct and condition	Numerosity (n)
sfGFP, M9 salts	10	AcnA-sfGFP, M9 salts	5
sfGFP, M9 salts, Cam	13	AcnA-sfGFP, M9 salts, Cam	8
sfGFP, M9 salts, Glu+CA	15	AcnA-sfGFP, M9 salts, Glu+CA	10
sfGFP, M9 salts, Cam, Glu+CA	13	AcnA-sfGFP, M9 salts, Cam, Glu+CA	10