# Watching the release of a photopharmacological drug from tubulin using time-resolved serial crystallography

Maximilian Wranik[1], Tobias Weinert[1], Chavdar Slavov[2], Tiziana Masini[3], Antonia Furrer[1], Natacha Gaillard[1], Dario Gioia[3], Marco Ferrarotti[3], Daniel James[1], Hannah Glover[1], Melissa Carrillo[1], Demet Kekilli[1], Robin Stipp[1], Petr Skopintsev[1], Steffen Brünle[1], Tobias Mühlethaler[1], John Beale[1], Dardan Gashi[4], Karol Nass[4], Dmitry Ozerov[5], Philip J.M. Johnson[4], Claudio Cirelli[4], Camila Bacellar[4], Markus Braun[2], Meitian Wang[4], Florian Dworkowski[4], Chris Milne[4], Andrea Cavalli[3,6], Josef Wachtveitl[2], Michel O. Steinmetz[1,7,*], Jörg Standfuss[1,*]

[1] Division of Biology and Chemistry, Paul Scherrer Institut, 5232 Villigen, Switzerland.
[2] Institute of Physical and Theoretical Chemistry, Goethe University, Frankfurt am Main, Germany.
[3] Computational & Chemical Biology, Istituto Italiano di Tecnologia, 16163 Genova, Italy.
[4] Photon Science Division, Paul Scherrer Institut, 5232 Villigen, Switzerland.
[5] Scientific Computing, Theory and Data, Paul Scherrer Institut, 5232 Villigen, Switzerland.
[6] Department of Pharmacy and Biotechnology, University of Bologna, 40126 Bologna, Italy.
[7] Biozentrum, University of Basel, 4056 Basel, Switzerland.

*Correspondence to: michel.steinmetz@psi.ch and joerg.standfuss@psi.ch

## Abstract

The binding and release of ligands from their protein targets is central to fundamental biological processes as well as to drug discovery. Photopharmacology introduces chemical triggers that allow the changing of ligand affinities and thus biological activity by light. Insight into the molecular mechanisms of photopharmacology is largely missing because the relevant transitions during the light-triggered reaction cannot be resolved by conventional structural biology. Using time-resolved serial crystallography at a synchrotron and X-ray free-electron laser, we captured the release of the anti-cancer compound azo-combretastatin A4 and the resulting conformational changes in tubulin. Nine structural snapshots from 1 ns to 100 ms complemented by simulations show how *cis*-to-*trans* isomerization of the azobenzene bond leads to a switch in ligand affinity, opening of an exit channel, and collapse of the binding pocket upon ligand release. The resulting global backbone rearrangements are related to the action mechanism of microtubule-destabilizing drugs.

## Introduction

The dynamic interaction of small molecule ligands with their target proteins is the basis for numerous physiological processes in living cells. To catalyze chemical reactions, enzymes have to bind their substrates and release the products. Cellular signaling is coordinated by the binding of ligands that activate their cognate receptors in the cytoplasm or the cellular membrane. Small molecule ligands that act as drugs to selectively intercede in these physiological processes provide the most common strategy for medical intervention.

Photopharmacology introduces photochemical triggers into small molecule ligands to be able to change binding affinities and manipulate complex biological processes with the help of light[1]. The most widely used photochemical affinity switches are based on the azobenzene scaffold that allows for high quantum efficiencies, fast trigger rates, and reversible photoisomerization upon illumination[2]. A prominent example is an azobenzene derivative of combretastatin A4 (azo-CA4)[3], a photopharmacological compound that binds the colchicine-site of the αβ-tubulin heterodimer (hereafter called tubulin)[4]. Tubulin is the key building block of the microtubule cytoskeleton in all eukaryotic cells and amongst others essential for cell division[5]. It is thus a common target for medically relevant agents including the first line anti-cancer drug paclitaxel (Taxol). While colchicine modulates multiple pro- and anti-inflammatory pathways associated with gouty arthritis[6], combretastatin A4 (CA4) is a strong cell growth inhibitor[7] that has been tested in clinical trials against different types of tumors[8, 9]. Interestingly, the phase III drug sabizabulin, that decreases mortality in intense-care COVID-19 patients by 55%[10], is also a colchicine-site ligand, sharing one binding moiety with combretastatins. The diverse applications of these small molecule drugs highlight the importance of the colchicine-site as a hotspot to influence the various cellular processes related to microtubule structure and function. A better understanding of how the dynamic interplay between a ligand and the colchicine-site is related to the conformational plasticity of tubulin may help further developments of this class of anti-tubulin agents for medical applications.

Besides its applications in photopharmacology[1] and material sciences[11], the azobenzene scaffold is also ideally suited for time-resolved pump-probe spectroscopy to understand protein motion[12]. Among the latest incarnations of the pump-probe technique is time-resolved serial crystallography to resolve molecular snapshots of proteins that can be assembled into stop-motion movies of how proteins function[13, 14]. The method has been most successful when studying photoactive proteins that are easy to trigger with high temporal precision; however, these represent less than 0.5% of all known proteins[15]. In order to reach the full potential of

time-resolved crystallography, new ways have to be developed to trigger protein-ligand

interaction and protein dynamics in non-photoactive protein targets.

To this end, we show here how to utilize photopharmacological compounds to assess protein-

ligand interaction dynamics and the resulting protein response over time. We first obtained the

structure of azo-CA4 in complex with tubulin, before following its photoflash-induced release

at the Swiss Light Source (SLS) and the Swiss X-ray Free Electron Laser (SwissFEL). We

resolved the structural evolution at up to 1.7 Å spatial resolution in nine logarithmically-timed

structural snapshots ranging from 1 ns to 100 ms. Our molecular snapshots show the formation

and dissolution of a metastable, intermediate ligand-binding pose, and the accompanying

reorganizations of the most frequently targeted drug-binding site in tubulin[16]. We anticipate

that similar photoswitchable compounds developed to control kinases, channels, G protein-

coupled receptors, and other pharmaceutically relevant protein targets will be important future

tools to study protein-ligand interaction dynamics in non-photoactive protein targets at high

spatial resolution and without compromising time resolution.

**Results**

**Adapting photopharmacology for time-resolved serial crystallography**

Time-resolved crystallography is ever more sample efficient (Supplementary Figure 1) and the

technology is now available at a growing number of synchrotrons and X-ray free-electron lasers

(XFELs)[17,18]. With increased availability of beamtime at these advanced X-ray sources, future

challenges will shift towards enabling dynamic studies on a wide range of relevant biological

targets. Photopharmacology could be an important part of the solution as the field already

developed a variety of photosensitive compounds that can be utilized as efficient triggers to

activate diverse soluble and membrane proteins of medical interest (for some examples, see

Supplementary Figure 2).

In the last few years, tubulin-targeting photoswitches based on different scaffolds have been

synthesized, for some of which we also obtained structures[19, 20]. To demonstrate the use of a

photochemical affinity switch for time-resolved crystallography, we chose azo-CA4 as a

prominent photopharmacological compound[21] with minimal modifications to the parent

compound CA4. Replacing the central C=C stilbene bond found in CA4 with an N=N

azobenzene bond in azo-CA4 results in a shift of the absorption maximum out of the ultraviolet

range and in an increased photoreactivity. After the modification, illumination with blue or

green light allows for reversible switching of the molecule between the high-affinity *cis* and

low-affinity *trans* conformations, respectively[3]. This effect allows perturbing the division of

cancer cells using a localized pulse of monochromatic light[3]. However, how azobenzene

isomerization changes binding affinity, how this induces structural adaptations within the

binding pocket, and how this then affects the global conformational plasticity of tubulin that is

needed during microtubule formation is largely unknown.

The idea behind our experiment was to use time-resolved crystallography to follow how the

ligand relaxes after photoactivation, allowing it to be observed bound to the protein in its out-

of-equilibrium, low affinity state, and eventually observe ligand unbinding and the resulting

changes in the protein (Figure 1). As a first step, we synthesized azo-CA4 and characterized its

tubulin-binding properties under crystallization conditions using the TD1 crystal system

composed of one αβ-tubulin heterodimer and one DARPin D1 molecule[22, 23] (Supplementary

Figure 3). Based on these results, we designed time-resolved crystallography experiments to

reveal the structural changes upon ligand isomerization and release.

Overall, we collected 1'139'945 indexable diffraction patterns with a resolution of up to 2.1 Å

when using serial synchrotron crystallography at the SLS[24, 25] and 1.7 Å when using serial

femtosecond crystallography at the SwissFEL[26] (Supplementary Table 1). The XFEL data are

divided into "dark" images collected without light exposure, images from crystals without

ligand, and pump-probe data taken at logarithmically spaced time delays of 1 ns, 10 ns, 100 ns,

1 μs, 10 μs, 100 μs, 1 ms, and 10 ms after illumination with flashes of laser light.

This untargeted approach relies on both synchrotron and XFEL setups to cover nine orders of

magnitude in time, which increases accessibility and remains an option even for cases where

initial spectroscopic data is scarce. We have characterized the ultrafast regime and initial

photochemical reaction using transient absorption spectroscopy (Supplementary Figure 3);

however, here we focus on the temporal domain starting from the nanoseconds where the

biochemically most relevant conformational changes in the protein are to be expected[18].

Difference electron density maps ($F_{obs}$(light)-$F_{obs}$(dark)) calculated from the time-resolved data

are of high quality (Figure 1D, Supplementary Figure 4, Supplementary Movie 1) and can be

qualitatively followed by Pearson correlation analysis to highlight transitions between

structural intermediates[27] (Figure 1B). We observed three major blocks of elevated correlation

compatible with a nonequilibrium photoproduct population that relaxes via several

intermediates into the ligand-free state of tubulin. Following the build-up of negative density

around azo-CA4 (Figure 1C) suggests that ligand release starts after the 1 ms delay at the end

of the second block observed in the Pearson correlation analysis. Using structural refinements

against extrapolated data from the light-activated fraction of the structure factors[28, 29], we were

able to generate atomic models for predominant intermediate states (Figure 1D, Supplementary

Table 2). In addition, we used conventional refinement approaches to obtain room temperature

crystal structures of tubulin in its *cis*-azo-CA4 bound and ligand-free states.

Together, these data provide a comprehensive overview of the structural changes occurring

after *cis-trans* isomerization and the following release of the azo-CA4 compound from its

tubulin binding pocket. A molecular movie obtained from morphing between these states

(Supplementary Movie 2) summarizes the critical molecular rearrangements that are described

in the following.

**Formation of a metastable binding pose after azo-CA4 photoisomerization**

The conformation of a ligand is generally presumed to precisely fit the architecture of its

binding pocket. The photoinduced release of azo-CA4 provides a system to study what happens

when a ligand deviates from its primary binding pose (Figure 2A). Before light-activation, the

*cis* isomer of azo-CA4 is bound to the colchicine site at the interface between the $\alpha$ and $\beta$

subunits of tubulin (Figure 1A). Interactions with the protein are primarily hydrophobic in

nature, except for hydrogen bonds to the backbone carbonyl of $\alpha$Thr179 and the backbone

amide of $\alpha$Val181, respectively (Figure 2B). Further interactions are mediated by a water

molecule interacting directly with the azobenzene bond of azo-CA4. Overall, the binding pose

of the ligand is nearly identical to that of the parent compound CA4 that contains a stilbene

instead of the azobenzene group connecting the A and B rings of the ligand[4].

Transient absorption spectroscopy shows how illumination of azo-CA4 induces *cis-trans*

isomerization around the azobenzene bond, which is completed in the late picosecond range

following by its relaxation into the nanoseconds (Supplementary Figure 3). Our structures show

how the compound transitions from the partially disordered conformation observed in the 1 ns

time delay into a relaxed *trans*-azo-CA4 state observed at 100 ns after light activation. At this

point in time, the molecule is stretched by 2.2 Å, which relocates its A ring by 2.6 Å towards

the $\beta$T7 loop known to be involved in ligand unbinding[4, 23] (Figure 2C). Stretching along the

azobenzene bond changes the interaction pattern with the protein and the binding pocket. The

water-mediated interactions are broken and relocation of the two ring moieties of azo-CA4

reduces their hydrophobic stacking interactions with the protein.

To bridge the gap between structural effects and binding energy we relied on molecular

dynamics (MD) simulations[30]. We aimed at estimating the relative binding free energy

difference between the *cis* and *trans* isomers ($\Delta\Delta G_{cis-trans}$) using the computational protocol

utilized for the tubulin-combretastatin A4 complex[4]. As expected, the *trans* isomer is the most

stable form in both the bound and unbound simulations with $\Delta G$ values of $\Delta G_{cis-trans-bound} = -14$

kcal/mol and $\Delta G_{cis\text{-}trans\text{-}unbound}$ = -19 kcal/mol, respectively. The resulting relative binding free

energy difference of $\Delta\Delta G_{cis\text{-}trans}$ = 5 kcal/mol clearly favors the *cis*-isomer stabilized interactions

with tubulin (free energy profiles are given in Supplementary Figure 5). Such a free energy

difference translates into an increased affinity of ~4300 times based on the equilibrium

probability ratio and Boltzmann statistics. This change is in good agreement with radioligand

scintillation proximity assays that indicated an $EC_{50}$ value of ~30 μM for illuminated azo-CA4

(representing a mixture of cis-trans isomers) but no significant competitive binding of the

relaxed *trans*-azo-CA4 to tubulin[3]. The low affinity of *trans*-CA4 also explains why we were

unable to crystallize this isomer in complex with tubulin using classical crystallography

methods but instead needed a time-resolved experiment to resolve the metastable binding pose

before azo-CA4 release.

**Binding pocket changes upon azo-CA4 release**

The colchicine site undergoes pronounced conformational changes between the ligand-bound

and -unbound states that are incompatible with a simple "lock-and-key" model of protein-ligand

interactions[4, 31, 32, 33]. In our time-resolved data, the relevant temporal regime starts at 100 ns,

when the A ring of the ligand has been repositioned into close contacts with residue βAla248

of the βT7 loop (Figure 3A). Initial conformational changes of the βT7 loop follow on the

microsecond scale with further rearrangements of the loop, including a displacement of

βAsn247 and βLeu246 occurring in the millisecond time range. The binding pocket initially

expands starting from a volume of 446 $\text{Å}^3$ before illumination (*cis*-azo-CA4) to 525 $\text{Å}^3$ at 100

ns (relaxed *trans*-azoCA4), and then collapses from 363 $\text{Å}^3$ at 1 ms (after βT7 loop movement)

down to 71 $\text{Å}^3$ at 10 ms when the compound has left the binding pocket. Based on visual

inspection of difference maps and plotting electron density changes on the ligand over time, we

have modeled the release of azo-CA4 between the 1 and 10 ms time delays (Figure 1C).

However, residual ligand can still be observed at low occupancy in the 10 ms data and is only

fully released at about 100 ms, where the structure is essentially identical to the one determined

in the absence of azo-CA4 (Supplementary Figure 6). The long release time is consistent with

a stochastic nature of the process where unbinding events of individual molecules occur over

several milliseconds in time once they are initiated on faster time scales.

The binding pocket is not freely accessible to the solvent but our structural analysis suggests

the opening of an exit channel between 100 μs and 1 ms (Figure 3B). Reorganization of the

βT7 loop, particularly through a rotation of the βLeu246 side chain, results into an 8 Å long

channel forming a possible exit site close to the A ring of the ligand (Figure 3C). The diameter

of the channel is 4-6 Å, which seems small given the 7 Å diameter for the larger A ring of azo-

CA4. As the unbinding takes place over many milliseconds throughout the crystal, it seems

likely that the channel transiently widens in short-lived intermediates that did not sufficiently

accumulate to be observed. To study this possibility, we have used MD simulations (Figure 3D)

starting from the 1 ms delay where the opening of the channel is most prominent. Almost 50 μs

of simulations (Supplementary Datasets 1 and 2) suggests that the azo-CA4 approaches a

position close to the proposed exit site of the protein, where only minimal changes would be

sufficient to let the ligand diffuse out of its binding pocket. Once the ligand detaches our

crystallographic data again shows how the βT7 loop flips back and packs against β-tubulin

residues that were previously engaged in azo-CA4 binding.

Together, these data are consistent with the flexible βT7 loop to act as a "molecular gate" that

opens and closes the colchicine site and, depending on its current conformation, directly

competes with ligand binding[32, 33]. Notably, all colchicine-site ligands must displace the βT7

loop upon binding[16, 34] as the pocket is contracted fully in ligand-free tubulin structures. Along

with the dynamic nature of the pocket at room temperature[24], this observation suggests

conformational selection as the predominant mechanism of how microtubule-destabilizing

agents interact with the colchicine-site.

**Global backbone rearrangements**

Tubulin displays substantial structural plasticity that is key to microtubule dynamics and thus

its cellular function. This includes the global conformation from a "curved" free tubulin to a

"straight" microtubule-lattice incorporated tubulin conformation[35]. This reorganization of the

tubulin dimer is accompanied by a contraction of the colchicine site[32, 33]. Ligand binding

prevents pocket contraction and thus stabilizes the curved conformation[16] as the principal action

mechanism for many tubulin-targeting anti-cancer drugs[16, 32, 33, 35].

To illustrate how global backbone rearrangements develop during the release of azo-CA4, we

superimposed our time-resolved structures and visualized the rearrangements of Cα backbone

atoms as individual vectors. We observed several motions that correspond well with our initial

Pearson-correlation analysis of electron density changes, and which coincide in time with the

major events in the ligand-binding pocket (Figure 4). Global structural changes are already

present in the nanosecond time range, where we observed coordinated inward motions of both

the α- and β-tubulin subunits (Figure 4A). These changes correlated well to the time range

where the A ring of azo-CA4 shifts to a new place completing the *cis*-to-*trans* relaxation process

within the binding pocket. Interestingly, some of these movements are reverted in the

microsecond time range, where the βT7 loop starts moving (Figure 4B). At this stage of the

reaction, global movements in the α and β subunits are directed predominately away from the

colchicine-site and may cause opening of ligand release pathway at 1 ms. In the later

millisecond range, where the βT7 loop finally collapses into the binding pocket, backbone

movements are primarily located in the α-tubulin subunit and bring the structure back to its

ligand-free state (Figure 4C).

To analyze whether the global backbone rearrangements could have an impact on the overall

conformation of microtubules, we analysed the curvature of an oligomer constructed from three

longitudinally, head-to-tail aligned tubulin dimers, as previously described[36]. Indeed, the

curvature is reduced significantly towards the direction of the straight protofilament

conformation found in microtubules once azo-CA4 is released. Our time-resolved structures

thus suggest that the global tubulin backbone movements, as well as overall changes in tubulin

conformational curvature evolve in par with structural alterations of the ligand that is bound to

the colchicine site.

**Discussion**

In the field of photopharmacology ligand design is done largely "blind" in the sense that little

is known about the molecular mechanism of how these modifications will alter dynamics and

activity of the target protein. In this context, it was reassuring to see that the interaction pattern

of the parent compound CA4 and the photoactive azo-CA4 remain largely the same. Further

knowledge of both the initial binding mode and the metastable binding pose after illumination

provides the molecular basis to understand differences in binding affinity between the *cis* and

*trans* forms of the compound. Together with molecular information on how the pocket adapts

during ligand release, this may provide innovative angles to improve the potency of azo-CA4

and perhaps other photopharmaceutical compounds.

In the wider context of general protein-ligand interaction dynamics, our work provides

molecular information beyond the classical "key and lock" model. The pharmaceutical industry

relies on such information as an integral part of a rational approach to discover new drugs and

understand their mechanisms of action. The colchicine-site is the most frequently targeted drug-

binding site in tubulin[16] with broad relevance to combat gout[6], cancer[7, 8, 9], and COVID-19[10].

Following the relaxation and release of the same ligand within different isoforms of tubulin[23]

or characterizing ligands specific for tubulin variants of parasitic origin[37] may offer new

avenues to rationally design innovative anti-tubulin ligands.

Finally, our study demonstrates how compounds developed for photopharmacology can be used

for time-resolved crystallography. While promising results using photo-caged compounds[38] and

rapid mixing have been shown[39, 40], these approaches will not allow observations on the fast

time scales relevant for protein-ligand interactions. It is furthermore difficult to predict how

ligands diffuse within a crystal, as this depends on crystal shape, size, packing and solvent

content among other parameters [41]. Starting a time-resolved experiment with a prebound ligand

is an elegant solution to the problem, even though it would be interesting to study to what extent

ligand binding and unbinding follow the same principles.

The underlying X-ray technology is now widely available, with many new endstations at $4^{th}$

generation  synchrotrons  and  XFELs  purpose  build  for  time-resolved  pump-probe

measurements[17, 18]. The rapidly expanding field of photopharmacology delivers growing

numbers of photoswitches to, for example, modulate the activities of TRP channels, kinases, G

protein-coupled receptors, or to tackle antibiotic resistance[1, 11, 12, 42, 43, 44]. Together these

developments promise more direct experimental insights into the dynamic nature of protein-

ligand interactions in a wide range of relevant protein targets.

**Methods**

**Purification of tubulin-DARPin D1 complex**

Bovine brain tubulin was purchased from the Centro de Investigaciones Biológicas

(Microtubule Stabilizing Agents Group), CSIC, Madrid, Spain. DARPin D1 was expressed and

purified according to previous descriptions[22]. Detailed protocols for obtaining high-resolution

tubulin crystals can be found here[45].

Briefly, the pET-derived vector PSTCm9 includes the gene for the artificial DARPin D1 as a

N-terminal fusion with thioredoxin followed by a hexahistidine-tag and a thrombin cleavage

site. For expression the DARPin D1 carrying plasmid was transfected into competent E.coli

BL21 (DE3) cells and the protein expressed in LB medium in 2 L Erlenmeyer flask. The cells

were harvested by centrifugation at 5000 g and stored at 80 degrees Celsius.

The cell walls were lysed on ice using an ultrasonic cell disrupter applying alterations of 2 s on

and off pulses with 30% amplitude for 22 minutes. The lysate was centrifuged at 20.000 g at 4

degrees Celsius for 40 minutes, and the supernatant was collected. Filtered supernatant was

applied to a Ni-NTA column on a fast protein liquid chromatography system (FPLC). Eluted

protein fractions were analyzed by SDS-Page, pooled together and the hexahistidine-tag

cleaved using thrombin. After cleavage, the solution was applied to a Ni-NTA column on a fast

protein liquid chromatography system and the flow through collected. Suitable fractions were

identified by characteristic UV absorbance at 280 nm, concentrated and applied on a Superdex

75 16/60 column for size-exclusion chromatography. Respective fractions were pooled

together, concentrated to a final DARPin D1 concentration of ~25 mg/mL using a 3 kDa cutoff

concentrator, flash-frozen in liquid nitrogen in 50 μL aliquots, and stored at -80 degrees Celsius.

The concentration was determined in mg/mL using a Nanodrop- 1000 instrument against the

size exclusion buffer using an extinction coefficient of 0.416 $Lg^{-1} cm^{-1}$.

To extract the active and non-aggregated tubulin from the delivered sample stocks, a sequence

of solution-based tubulin polymerization and depolymerization steps was performed. On ice,

the lyophilized tubulin is dissolved and diluted to 10 mg/mL with a polymerization buffer

containing 50 mM 2-(*N*-morpholino)ethanesulfonic acid (MES-KOH) pH 6.8, 0.5 mM ethylene

glycol-bis(β-aminoethyl ether)-N,N,N′,N′-tetraacetic acid (EGTA), 0.4 mM Guanosine-5'-

triphosphate (GTP), 6 mM $MgCl_2$, and 33% glycerol. The lyophilized pellet was dissolved over

~30 minutes by regular shaking and pipette resuspension on ice. Afterward, the clear solution

was ultracentrifuged at 150.000 g at 4 degree Celsius for 10 minutes. The supernatant was

carefully extracted, transferred to pre-warmed tubes, and incubated for 25 minutes in a water-

filled heating block at 37 degree Celsius. This procedure induces tubulin polymerization

causing the solution to turn viscous and turbid. The solution was then ultracentrifuged at

300.000 g at 30 degree Celsius for 15 minutes. The supernatant was carefully discarded, and

the tubulin pellet was poured over with depolymerization buffer containing 80 mM Pipes-KOH

pH 6.8, 0.5 mM EGTA, 2 mM Guanosine-5'-diphosphate (GDP), 1 mM $MgCl_2$, and 2 mM

$CaCl_2$. After 15 minutes of incubation, the pellet was resuspended with a pipette, and the

solution was incubated again on ice for additional 15 minutes. This step was repeated until the

pellet was dissolved entirely. The depolymerized tubulin solution was diluted with

depolymerization buffer to a concentration of 9.8 mg/mL and stored at -80 degree Celsius.

**Tubulin crystallization**

For crystallization, the tubulin-DARPin D1 (TD1) complex was formed by mixing the

respective components in a 1:1.1 molar ratio. The TD1 complex was crystallized using

EasyXtal 15-well plates by the hanging drop vapor diffusion method (drop size 2 μL, drop ratio

1:1, 8 drops per well) at a concentration of 9.8 mg $mL^{-1}$ at 20°C with a precipitant solution

containing 21% (w/v) polyethylene glycol (PEG) 3000, 0.2 M ammonium sulfate and 0.1 M

bis-tris methane, pH 5.5. All drops were subsequently hair-seeded with crystalline material

obtained in previous crystallization trials to increase homogeneity and density of crystals. After

48 hours, crystals were washed off the plates with precipitant solution, collected in 0.6 ml tubes

and vortexed for a few seconds. This procedure induces a batch-crystallization within the tubes

and a sedimented crystal pellet after 24 hours of incubation at 20°C. Initially crystals grew as

long needles that could be broken into smaller fragments of approximately 20x20x5 μm$^3$ during

the sample preparation described below.

**Azo-CA4 synthesis**

The synthetic photoswitch azo-CA4 was prepared following the experimental procedure

described in detail by Borowiak *et al* [3]. First, catechol (4.44 g, 40.3 mmol) was reacted with

tosylchloride (9.53 g, 50 mmol) in pyridine (10 mL) for 2 hours at room temperature. The

reaction was quenched with aq HCl, and the organic phases extracted with dichloromethane,

dried on MgSO$_4$, filtered, concentrated and purified by column chromatography (gradient of

cyclohexane/EtOAc, with EtOAc from 10% to 50%), resulting in the mono-tosylated product

(7.5 g, 28.4 mmol) in 70% yield. Mono-tosylcatechol was then reacted with commercial 3,4,5-

trimethoxyanline via a diazo coupling with isopentyl nitrite. The aniline (297 mg, 1.62 mmol)

was dissolved in methanol (8.1 mL) and concentrated HCl (0.4 mL), and the mixture was cooled

in an ice bath. A solution of isopentyl nitrite (0.22 mL, 1.65 mmol) in methanol (0.97 mL) was

added dropwise and the reaction stirred for 30 minutes in the cold. A cold solution of the mono-

tosylcathecol (450 mg, 1.7 mmol) in methanol (3.2 mL) and NaOH (2.0 M, 2.9 mL) was

prepared, and to it was added the solution of the diazonium dropwise over 1 minute. After 5

hours stirring in the cold, the pH was adjusted to 7 with phosphate buffer, dichloromethane was

added, and the aqueous phase was extracted with dichloromethane. The combined organic

layers were washed with water and brine, dried on MgSO$_4$, filtered, concentrated and purified

by column chromatography (gradient of cyclohexane/EtOAc, with EtOAc from 20% to 50%),

affording the para-phenolic azobenzene (492 mg, 1.07 mmol) in 66% yield. The resulting

diazocompound (480 mg, 1.05 mmol) was then methylated in acetone (9.6 mL) with K$_2$CO$_3$

(431 mg, 3.12 mmol) and MeI (0,13 mL, 2.12 mmol) at room temperature for 16 hours. The

crude was concentrated and purified by column chromatography (cyclohexane/EtOAc, with

EtOAc from 10% to 50%) affording the methylated diazocompound (491 mg, 1,04 mmol) in

99% yield. The resulting compound (415 mg, 0.88 mmol) was then treated in basic media (KOH

in MeOH, reflux for 1 hour) to remove the tosyl group. The reaction mixture was concentrated,

dissolved in EtOAc and aq KH$_2$PO$_4$ (10%), the organic phases extracted with EtOAc,

combined, washed with water and brine and dried on MgSO$_4$. Purification by column

chromatography (cyclohexane/EtOAc, with EtOAc from 20% to 50%) delivered azo-CA4 (220

mg, 0.69 mmol) with 78% yield. Comparison to published NMR spectra[3] confirmed successful

synthesis (1H NMR (400 MHz, DMSO-d6) δ 9.47 (s, 1H), 7.46 (dd, J = 8.5, 2.4 Hz, 1H), 7.32 (d, J = 2.4 Hz, 1H), 7.18 (s, 2H), 7.12 (d, J = 8.6 Hz, 1H), 3.88 (s, 6H), 3.87 (s, 3H), 3.75 (s, 3H)).

**Sample preparation for serial crystallography**

Crystal pellets were resuspended and the contents of two 0.6 mL tubes were combined and collected into 2 mL tubes. After centrifugation in a tabletop centrifuge (1000 g, 15 seconds), a concentrated crystal pellet of 45-60 μL volume was obtained. The pellets were extracted, transferred into PCR tubes and mixed with the *trans*-azo-CA4 to a final concentration of 1 mM. To switch the compound into the binding *cis* conformation, the tubes were then illuminated for at least 4 hours at 385 nm before incorporation into hydroxyethylcellulose hydrogel (Sigma-Aldrich)[46]. Hydrogels containing 22% (w/v) hydroxyethylcellulose were prepared in syringes and left to cure for 5 days. Illuminated crystals were transferred into a Hamilton syringe and mixed into hydrogel matrix at a ratio of 4:7 using a three-way-coupler[47] before loading into the high viscosity injector.

**Experimental setup and XFEL data collection**

Time-resolved serial crystallographic data of light-induced azo-CA4 from tubulin were collected within 2 days of beamtime in September 2020 at the Alvra experimental station of SwissFEL. The X-ray source provided pulses with a photon energy of 12.1 keV and a pulse energy between 300 μJ and 550 μJ at a repetition rate of 100 Hz. To reduce X-ray scattering (background level), the sample chamber was pumped down to a pressure of 150 mbar and then refilled with helium to 500 mbar. This process was continuously repeated during the measurement. Data were collected using a Jungfrau 16M operated in 4M mode. Hit rate determination and initial data processing during the experiment were done with the online data analysis pipeline described by Nass et al.[48].

TD1 crystals homogeneously incorporated into hydroxyethylcellulose were loaded into a high viscosity injector connected to an HPLC pump[49]. The system extruded the crystal loaded hydrogel matrix into the pump–probe interaction point through a 75 μm capillary at a flow rate of 5.3 μl per minute. In the interaction point, the probing X-ray pulses intersected with a circularly polarized pump beam originating from an optical parametric amplifier producing laser pulses with 350 fs duration ($1/e^2$), 475 nm wavelength and 10 μJ total energy in a focal spot of 65 μm ($1/e^2$) diameter; corresponding to a maximal laser fluence of 301 mJ/cm$^2$ and laser power density of 861 GW/cm$^2$. In order to follow the reaction of light-induced azo-CA4

ligand release from tubulin between the pump laser and the probing, XFEL pulses were chosen

at $\Delta t$ = 1 ns, 10 ns, 100 ns, 1 μs, 10 μs, 100 μs, 1 ms and 10 ms.

**Serial synchrotron crystallography**

Serial synchrotron data were collected using a previously described setup[25]. Crystals embedded

into hydroxyethylcellulose were injected in the intersecting X-ray and laser beam paths by a

high viscosity injector[49] with a 75 μm diameter nozzle.

The jet speed was set to 0.5 mm/ms and data frames were recorded using an Eiger16M detector

with a frame rate of 100 Hz. The beam size was set to 20 x 5 μm$^2$ to cover the central part of

the hydroxyethylcellulose column on one axis and extrusion by one frame on the other. In this

way we ensured the majority of crystals were exposed, whilst minimizing similarities in

consecutive frames for more reliable statistics. A photon flux of around 1.5 x 10$^{12}$ photons/s

resulted in a radiation dose of about 5 kGy per recorded detector frame when calculated with

RADDOSE-3D[50]. Depending on their size and orientation while traversing the beam this

corresponds to between 5 and maximally 100 kGy per crystal. Illumination was done using

light at 445 nm from a laser diode with a peak optical power of 5 mW (Roithner Lasertechnik).

Light intensity at the interaction region was approximately 2.5 mW in a spot size of about 100

x 50 mm$^2$. The illumination time was set to approximately 100 ms by adjusting the distance

between the laser and X-ray interaction regions.

**Data processing**

Indexing, integrating and merging of obtained data was performed using Crystfel version

0.9.1[51, 52]. In detail, the peakfinder9 algorithm was used for peak detection, xgandalf[53] was used

to index obtained data; and peaks were integrated using --ring-radius=2,3,6 ; partialator options

--model=unity, --iterations=1.0, --push-res=1.5 were used to merge selected patterns. A general

resolution cutoff of 1.7 Å was chosen (see Supplementary Table 1) and refinements for

collected dark-state and apo-state data were carried out to the full resolution range. Structural

refinements of time-resolved data were carried out to 2.2 Å resolution due to data degradation

when extrapolating data (see Supplementary Table 1). Serial synchrotron crystallography data

obtained at SLS showed stronger anisotropy than the data obtained at the SwissFEL. The data

were therefore corrected for anisotropy using the STARANISO server[54].

**Difference density maps calculation**

Calculations of $F_{obs}$(light) − $F_{obs}$(dark) difference maps were performed using PHENIX

v1.19.2[55]. In detail, the multi-scaling option was used excluding amplitudes smaller than 2 σ

and exclusively considering the resolution range between 6 Å and 1.7 Å. For all calculated

$F_{obs}$(light) − $F_{obs}$(dark) difference maps, the phases of the refined dark state were used.

$F_{calc}$(light) − $F_{calc}$(dark) difference maps were computed to the same resolution using the CCP4

suite v8.0[56].

**Pearson correlation of difference maps**

To compute Pearson correlations independent of individual dataset multiplicities we reduced

the number of patterns to match the time delay where we collected the least data (55,065

patterns) and calculated difference maps as stated above. These difference maps where then

used to calculate the Pearson correlation of integrated difference electron densities[27] around all

protein atoms and around the ligand and all residues lying within a 3 Å sphere around the ligand.

Water molecules and hydrogen atoms were excluded in the selection. The density maps were

integrated within a 2 Å radius around the atoms and densities below 1.5 σ were ignored. The

resulting Pearson correlation values where normalized to the largest correlation, displayed on

a gray scale and values smaller than 0.3 were omitted from the figures for clarity.

**Integration of negative density around the ligand**

To plot the release of azo-CA4 over time we integrated the negative density around the ligand

in same way as the Pearson correlation data for the whole protein (1.5 A radius, 3.0 sigma

cutoff). The integrated negative density was normalized: the highest value (apo) was set to 1

the lowest negative ligand density (1 ms) was set to 0. Since apo data and time-resolved data

from the XFEL experiment are not fully comparable due to a roughly 4 fold difference in

occupancy. The negative density of the apo state was downscaled to lower occupancy by

assuming a roughly 4-fold occupancy increase is similar to a 4-fold increase of the number of

observations, resulting in an estimated scaling factor of 0.5 (assuming signal to noise improves

by $\sqrt{2}$ for doubling the number of observations).

**Data extrapolation**

Calculation of extrapolated data was performed using the method described by Pandey et al.[29].

Briefly, the linear approximation was used: $F_{extra} = [100/A \times (F_{obs}(light) − F_{obs}(dark)] + F_{calc}$,

with A representing the activation level in percent, Fcalc representing amplitudes of the high-

resolution dark state model and $F_{extra}$ representing the extrapolated structure factor amplitude.

The activation level A, defined as the percentage of dark state molecules being activated after

pump laser pulse exposure, was determined as described[29]. Briefly, extrapolated data were

calculated using a decreasing activation level A. Then the negative electron density in $2F_{extra}$-

$F_{calc}$ maps calculated using the dark model phases, was integrated around nitrogen atoms of the

ligand which have strong negative difference density peaks on them in each of the light

activated structures and hence leave their position. The negative density was then plotted

against the activation level. The intersection of the linear parts of the resulting graph (negative

density is zero for not enough dark state subtracted and starts to linearly increase as dark is over

subtracted) was taken as the activation level. The activation levels determined in this way

fluctuated between 18-27% but since all data were measured under the same conditions, we

chose the average of 22% to refine the structures. As a further control we determined the

activation level using the re-appearance of dark state features in $2F_{extra}$-$F_{calc}$ maps[57, 58], which

confirmed our initial assignment.

**Structure determination and refinement**

The structure of the dark-state was solved by molecular replacement with 5NQT as a search

model. Structural refinements of the model were done using PHENIX v1.19.2[55] with iterative

cycles of manual adjustments made in Coot v0.9.8.1[59]. As a final step the ligand was refined

using Buster v2.10.4[60], employing the -qm option to use a quantum mechanics target for the

ligand rather than restraints which may be a source of errors, especially for molecules which

are not at their energy minimum. The structure of the apo-state was directly refined using the

final dark-state structure as a starting point.

Before the actual refinement and model building of the obtained time-resolved data, the

negative amplitudes resulting from the extrapolation procedure were removed. The models

were manually adjusted to best fit observed difference map features as well as extrapolated

maps using the final dark-state model as a starting point.

Before the actual refinement and model building of the obtained time-resolved data, the

negative amplitudes resulting from the extrapolation procedure were removed. The models

were manually adjusted to best match observed difference map features, to produce similar

$F_{calc}$(light)-$F_{calc}$(dark) maps and to fit extrapolated maps using the final dark model as a starting

point. To match the observed difference density features using $F_{calc}$(1 ms)-$F_{calc}$(dark) maps we

chose to exclude Leu246 from XYZ refinement after manual adjustment for the 1 ms structure.

Given the activation level of 22 percent, we choose to refine the predominant structural species

by comparing the models to previous and consecutive structures.

**Binding pocket and protein backbone analysis**

Protein interactions in the binding pocket were identified with the protein-ligand interaction profiler[61]. Changes in the volume of the colchicine-site over time were determined using PyVOL[62] with the parameters ($\beta$–tubulin, minimum radius = 1.65; maximum radius = 3.8; minimum volume = 50, standard partitioning).

To visualize global protein backbone movements, alternative conformers were removed and structures truncated to poly-Ala using Phenix PDB tools[55]. Respective structural displacements of secondary structural elements were indicated using the PyMOL Python script Modevectors (by Sean M. Law, University of Michigan). Displacements greater than 0.5 Å were classified as significant (cutoff=0.5). For visualization purposes, only the displacement vectors of every second (skip=2) C$\alpha$-backbone atom (atom=CA) were shown and the vectors amplified by a factor of 6 (factor=6). Additional settings involve the vector representation (head=2.0, tail=0.85, head_length=3.0, cut=0).

To find possible release pathways during reorganization of the binding pocket we used the program Caver 3.03[63] using the following settings: minimum probe radius = 1.9 Å, shell depth = 4 Å, and shell radius = 5 Å, azo-CA4 position as starting coordinate.

**Time-resolved spectroscopy measurements**

The time-resolved transient absorption data were recorded using a home-built pump–probe setup, as described in detail previously[64]. In short, the fundamental laser pulses (1 mJ, 775 nm, 130 fs, 1 kHz) were provided by a Ti:Sa amplifier (Clark, MXR-CPA-iSeries). The pump pulses at 438 nm were generated using a home-built two stage NOPA (noncollinear optical parametric amplifier) and sum frequency mixing. The pulses were pre-compressed in a prism compressor located between the two NOPA stages. White light continuum pulses (300-750 nm) were generated by focusing a fraction of the laser fundamental beam into a $CaF_2$-crystal (5 mm). The continuum pulses were split into probe and reference beams. The reference beam was guided directly into a spectrograph, while the probe beam was focused at the sample position then collected and directed into a second spectrograph. The spectrographs (AMKO Multimode) contained gratings with 600 grooves/mm blazed at 300 nm and a photodiode array. The instrument response function (IRF) of ~100 fs in the experiments was estimated from the pump probe cross correlation. Anisotropic contributions to the measurements were avoided by performing the experiments at the magic angle condition (54.7° pump-probe polarization difference). The samples were held in a fused silica cuvette with an optical path length of 1 mm

which was constantly moved in the plane perpendicular to the direction of the probe pulse
propagation to avoid accumulation of photoproducts. To keep the azo-CA4 samples (bound and
free) in the *cis* isomer state, the cuvette was continuously illuminated with a high-power LED
at 385 nm. Global analysis of the experimental data was performed using OPTIMUS
(www.optimusfit.org)[64] to recover the lifetimes describing the *cis* → *trans* photoisomerization
kinetics. With the help of target analysis, a sequential kinetic scheme was fitted to the
experimental data to obtain the times of the highest population of each kinetic state, which were
helpful in determining the optimal time points for the XFEL experiments.
557
**Computational Methods**
**System preparation**
The starting models for simulations were built from the respective XFEL-based time-resolved
structures. We removed the DARPin protein and reconstructed the βM-loop (res 272-288) using
the one in chain D of PDB 4I4T as a template by employing the Prime[65] module of the
Schrödinger 2020-4 suite. Analogously, $Mg^{2+}$ ion in chain B (β-tubulin) was added by
superimposition with the one of chain D of 4I4T structure. Calcium ions were removed from
the model.
The final model consisted of an αβ-tubulin heterodimer in complex with azo-CA4, GTP, and
GDP molecules bound to the α- and β-tubulin subunits, respectively, as well as their associated
$Mg^{2+}$ ions and their coordinating water molecules. The resulting protein structure possessed
437 out of 451 residues of α-tubulin (UniProtKB ID P81947) and 431 out of 445 residues of β-
tubulin (UniProtKB ID Q6B856). Missing residues belonging to the intrinsically disordered C-
terminal tails of α- and β-tubulin were not modeled and C-termini were capped with N-methyl
amide (NME) groups. Residue protonation states were evaluated at pH 7.0 using the Protein
Preparation Wizard tool[66] implemented in the Schrödinger 2020-4 suite[67]. The αβ-tubulin
heterodimer structure was solvated with the TIP3P-model[68] for water molecules in a truncated
octahedron box using 12 Å as the minimum distance between the protein and the box edges.
The system was neutralized by adding Na+ ions resulting in a total of about 122k atoms. The
atomistic force field Amber-ff14SB[69] was used for all simulations. Parameters for $Mg^{2+}$ ions
and the GTP and GDP molecules were developed by Allner et al.[70] and Meagher et al.[71],
respectively.
Concerning the ligand, partial atomic charges were obtained with the Restrained Electrostatic
Potential (RESP) method on a DFT-optimized trans conformation at the B3LYP/6-31g* level.
The geometric optimization was carried out with Terachem v1.94. The General Amber

ForceField (GAFF) was used, with custom parameters for the three main dihedral angles: 1-1a-
1b-1', 6-1-1a-1b, 1a-1b-1'-6'.

To obtain such optimized parameters, we fitted the GAFF energy curves associated with a rigid
torsion of the dihedral angles to the corresponding DFT energy curves. For each dihedral, we
proceeded as follows: we started out with the geometrically optimized *trans* structure. We then
performed a clockwise rigid rotation with 15° steps followed by an analogous counter-
clockwise rigid rotation. For each rotation step we performed a DFT geometric optimization
restraining the rotated dihedral, and we evaluated the energy. We retained the minimum energy
among the clockwise and anticlockwise rotation cycles to overcome hysteresis. The three
resulting DFT energy profiles consisting of 24 evaluation points were used as a target for the
fitting procedure. DFT calculations were performed with Terachem v1.94, at the B3LYP/6-
31g* level. The fitting procedure was carried out separately for the three energy profiles with
mdgx, part of AmberTools21. The αβ-tubulin heterodimer system was assembled with the
LEaP tool implemented in the AmberTools 21 software package[67].

The designed models for QM calculations consisted of protein residues α(99-101, 178-181),
β(237-242, 248-259, 314-321, 349-354, 378-380), with the respective N- and C- termini capped
with acetyl (ACE) and NME groups, the azo-CA4 molecule, and its two closest water
molecules. They resulted in 743 atoms.

**Molecular dynamics simulations**

Plain molecular dynamics simulations starting from the tubulin-trans-azo-CA4-tubulin
complex 1 ms structure were performed using the Amber20 engine[67]. The system, after a
preliminary stage of energy minimization (1000 steps of steepest descent and 9000 of conjugate
gradient), was heated to the target temperature of 300 K with a 250 ps ramp using an integration
time-step of 1 ps and the weak-coupling algorithm, followed by another 500 ps of simulation
in the NVT ensemble after switching to an integration time-step of 2 ps. Subsequently, in order
to the equilibrate box dimensions, the system was simulated for 1 ns in the NPT ensemble using
the Berendsen barostat. For these initial stages, heavy atoms were harmonically restrained with
a force constant of 5 kcal/mol/Å$^2$. Finally, a last stage of 5 ns in the NVT ensemble, using the
Langevin thermostat with a collision frequency g of 2.0 ps-1. Bonds involving hydrogen atoms
were restrained, and a short-range, non-bonded cut-off of 8 Å was applied, whereas long-range
electrostatics were treated with the particle mesh Ewald (PME) method[72]. Periodic boundary
conditions (PBC) were applied.

After the equilibration stage, a cumulative 30 μs-long MD simulation was conducted with an

integration time step of 2 fs in the NVT ensemble at a target temperature of 300 K.

In order to evaluate the $\Delta\Delta G_{cis\text{-}trans}$, well-tempered metadynamics (WTMD) simulations[73] were

performed with GROMACS 2020.2[74] and PLUMED 2.6.1[75]. We performed WTMD enhancing

the sampling of the two dihedral angles 1-1a-1b-1' and 6-1-1a-1b for both the bound and the

unbound state. The bias factor, the deposition rate, the initial hill height, and the hill width were

set to 45, 0.5 ps−1, 2.5 kJ·mol−1, and 0.3 rad, respectively.

The WTMD simulation of the bound state was carefully designed to sample conformations

compatible with the XFEL-based time-resolved structures. A positional harmonic restraint with

K=100 kJ/mol nm$^2$ was introduced on the backbone of the protein together with the supervision

of the hydrogen bond to the backbone carbonyl of αThr179. With this regard, the simulation

was stopped and restarted with regenerated velocities whenever the H-bond was broken. The

resulting trajectory of 400 ns was reweighted in order to reconstruct the free energy profile.

Block analysis with a block size of 50 ns was performed to assess convergence and to plot 95%

confidence interval error bands. We performed WTMD enhancing the sampling of the two

dihedral angles 1-1a-1b-1' and 6-1-1a-1b. The bias factor, the deposition rate, the initial hill

height, and the hill width were set to 45, 0.5 ps−1, 2.5 kJ·mol−1 740, and 0.3 rad, respectively.

The WTMD simulation of the unbound state was performed with the same parameters used for

the bound state. In this case, we simulated a single ligand molecule in a solvated cubic box

using 10 Å as the minimum distance between the ligand and the box edges. After standard

minimization, thermalization and equilibration phases WTMD was carried out for 400 ns. The

resulting trajectory was reweighted in order to reconstruct the free energy profile. Block

analysis with block size of 50 ns was performed to assess convergence and to plot 95%

confidence interval error bands.

**Data availability**

The pdb entry 4I4T [http://doi.org/10.2210/pdb4I4T/pdb] (tubulin-RB-TTL complex) was used

as starting point to prepare the QM simulations. For molecular replacement we used pdb entry

5NQT [http://doi.org/10.2210/pdb5NQT/pdb] (tubulin-darpin complex). We used the pbd entry

5SYE [http://doi.org/10.2210/pdb5SYE/pdb] (taxol-stabilized microtubule) to represent the

tubulin conformation found in microtubules. Coordinates and structure factors of the presented

serial femtosecond crystallographic structures have been deposited in the PDB database under

accession code 7YYQ [http://doi.org/10.2210/pdb7YYQ/pdb] (dark) and 7YZ3

[http://doi.org/10.2210/pdb7YZ3/pdb] (apo). The serial synchrotron data have been deposited

under 7YZ6 [http://doi.org/10.2210/pdb7YZ6/pdb] (dark) and 7YZ5

[http://doi.org/10.2210/pdb7YZ5/pdb] (100 ms). Models refined against extrapolated data,

extrapolated structure factors and light data used for data extrapolation have been deposited

under accession codes 7YYV [http://doi.org/10.2210/pdb7YYV/pdb] (1 ns), 7YYW

[http://doi.org/10.2210/pdb7YYW/pdb] (10 ns), 7YYX

[http://doi.org/10.2210/pdb7YYX/pdb] (100 ns), 7YYY

[http://doi.org/10.2210/pdb7YYY/pdb] (1 μs), 7YYZ [http://doi.org/10.2210/pdb7YYZ/pdb]

(10 μs), 7YZ0 [http://doi.org/10.2210/pdb7YZ0/pdb] (100 μs), 7YZ1

[http://doi.org/10.2210/pdb7YZ1/pdb] (1 ms) and 7YZ2 [http://doi.org/10.2210/pdb7YZ2/pdb]

(10 ms). The start and end coordinates of the molecular dynamics simulations are provided as

Source Data file.

**References:**

1. Hull K, Morstein J, Trauner D. In Vivo Photopharmacology. *Chem Rev* **118**, 10710-
10747 (2018).
665

666 2. Beharry AA, Woolley GA. Azobenzene photoswitches for biomolecules. *Chem Soc*
667 *Rev* **40**, 4422-4437 (2011).
668

669 3. Borowiak M, *et al.* Photoswitchable Inhibitors of Microtubule Dynamics Optically
670 Control Mitosis and Cell Death. *Cell* **162**, 403-411 (2015).
671

672 4. Gaspari R, Prota AE, Bargsten K, Cavalli A, Steinmetz MO. Structural Basis of cis-
673 and trans-Combretastatin Binding to Tubulin. *Chem-Us* **2**, 102-113 (2017).
674

675 5. Brouhard GJ, Rice LM. Microtubule dynamics: an interplay of biochemistry and
676 mechanics. *Nat Rev Mol Cell Biol* **19**, 451-463 (2018).
677

678 6. Dalbeth N, Lauterio TJ, Wolfe HR. Mechanism of action of colchicine in the
679 treatment of gout. *Clin Ther* **36**, 1465-1479 (2014).
680

681 7. Pettit GR, Singh SB, Hamel E, Lin CM, Alberts DS, Garcia-Kendall D. Isolation and
682 structure of the strong cell growth and tubulin inhibitor combretastatin A-4.
683 *Experientia* **45**, 209-211 (1989).
684

685 8. Grisham R, Ky B, Tewari KS, Chaplin DJ, Walker J. Clinical trial experience with
686 CA4P anticancer therapy: focus on efficacy, cardiovascular adverse events, and
687 hypertension management. *Gynecol Oncol Res Pract* **5**, 1 (2018).
688

689 9. Baytas SN. Recent Advances in Combretastatin A-4 Inspired Inhibitors of Tubulin
690 Polymerization: An Update. *Curr Med Chem* **29**, 3557-3585 (2022).
691

692 10. Barnette KG, *et al.* Oral Sabizabulin for High-Risk, Hospitalized Adults with Covid-
693 19: Interim Analysis. *NEJM Evidence* **0**, EVIDoa2200145 (2022).
694

695    11.    Jerca FA, Jerca VV, Hoogenboom R. Advances and opportunities in the exciting
696            world of azobenzenes. *Nature Reviews Chemistry* **6**, 51-69 (2022).
697

698    12.    Bozovic O, Jankovic B, Hamm P. Using azobenzene photocontrol to set proteins in
699            motion. *Nature Reviews Chemistry* **6**, 112-124 (2022).
700

701    13.    Neutze R, Moffat K. Time-resolved structural studies at synchrotrons and X-ray free
702            electron lasers: opportunities and challenges. *Curr Opin Struc Biol* **22**, 651-659
703            (2012).
704

705    14.    Tenboer J*, et al.* Time-resolved serial crystallography captures high-resolution
706            intermediates of photoactive yellow protein. *Science* **346**, 1242-1246 (2014).
707

708    15.    Monteiro DCF, Amoah E, Rogers C, Pearson AR. Using photocaging for fast time-
709            resolved structural biology studies. *Acta Crystallographica Section D* **77**, (2021).
710

711    16.    Steinmetz MO, Prota AE. Microtubule-Targeting Agents: Strategies To Hijack the
712            Cytoskeleton. *Trends Cell Biol* **28**, 776-792 (2018).
713

714    17.    Branden G, Neutze R. Advances and challenges in time-resolved macromolecular
715            crystallography. *Science* **373**, 980-+ (2021).
716

717    18.    Orville AM. Recent results in time resolved serial femtosecond crystallography at
718            XFELs. *Curr Opin Struc Biol* **65**, 193-208 (2020).
719

720    19.    Gao L*, et al.* A Robust, GFP-Orthogonal Photoswitchable Inhibitor Scaffold Extends
721            Optical Control over the Microtubule Cytoskeleton. *Cell Chem Biol* **28**, 228-241 e226
722            (2021).
723

724    20.    Gao L*, et al.* In Vivo Photocontrol of Microtubule Dynamics and Integrity, Migration
725            and Mitosis, by the Potent GFP-Imaging-Compatible Photoswitchable Reagents
726            SBTubA4P and SBTub2M. *J Am Chem Soc* **144**, 5614-5628 (2022).
727

728    21.    Kirchner S, Pianowski Z. Photopharmacology of Antimitotic Agents. *Int J Mol Sci* **23**,
729            (2022).
730

731    22.    Pecqueur L*, et al.* A designed ankyrin repeat protein selected to bind to tubulin caps
732            the microtubule plus end. *Proc Natl Acad Sci U S A* **109**, 12011-12016 (2012).
733

734    23.    La Sala G*, et al.* Structure, Thermodynamics, and Kinetics of Plinabulin Binding to
735            Two Tubulin Isotypes. *Chem-Us* **5**, 2969-2986 (2019).
736

737    24.    Weinert T*, et al.* Serial millisecond crystallography for routine room-temperature
738            structure determination at synchrotrons. *Nat Commun* **8**, 542 (2017).
739

740    25.    Weinert T*, et al.* Proton uptake mechanism in bacteriorhodopsin captured by serial
741            synchrotron crystallography. *Science* **365**, 61-65 (2019).
742

743    26.    Abela R*, et al.* Perspective: Opportunities for ultrafast science at SwissFEL. *Struct*
744            *Dyn* **4**, 061602 (2017).
745

746 27.  Wickstrand C, *et al.* A tool for visualizing protein motions in time-resolved
747      crystallography. *Struct Dyn* **7**, 024701 (2020).
748

749 28.  Skopintsev P, *et al.* Femtosecond-to-millisecond structural changes in a light-driven
750      sodium pump. *Nature*, (2020).
751

752 29.  Pandey S, *et al.* Time-resolved serial femtosecond crystallography at the European
753      XFEL. *Nature Methods* **17**, 73-+ (2020).
754

755 30.  Decherchi S, Cavalli A. Thermodynamics and Kinetics of Drug-Target Binding by
756      Molecular Simulation. *Chem Rev* **120**, 12788-12833 (2020).
757

758 31.  Prota AE, *et al.* The Novel Microtubule-Destabilizing Drug BAL27862 Binds to the
759      Colchicine Site of Tubulin with Distinct Effects on Microtubule Organization. *Journal*
760      *of Molecular Biology* **426**, 1848-1860 (2014).
761

762 32.  Dorleans A, Gigant B, Ravelli RBG, Mailliet P, Mikol V, Knossow M. Variations in
763      the colchicine-binding domain provide insight into the structural switch of tubulin. *P*
764      *Natl Acad Sci USA* **106**, 13775-13779 (2009).
765

766 33.  Ravelli RBG, *et al.* Insight into tubulin regulation from a complex with colchicine and
767      a stathmin-like domain. *Nature* **428**, 198-202 (2004).
768

769 34.  Muhlethaler T, Gioia D, Prota AE, Sharpe ME, Cavalli A, Steinmetz MO.
770      Comprehensive Analysis of Binding Sites in Tubulin. *Angew Chem Int Ed Engl* **60**,
771      13331-13342 (2021).
772

773 35.  Knossow M, Campanacci V, Khodja LA, Gigant B. The Mechanism of Tubulin
774      Assembly into Microtubules: Insights from Structural Studies. *iScience* **23**, 101511
775      (2020).
776

777 36.  Brouhard GJ, Rice LM. The contribution of alphabeta-tubulin curvature to
778      microtubule dynamics. *J Cell Biol* **207**, 323-334 (2014).
779

780 37.  Gaillard N, *et al.* Inhibiting parasite proliferation using a rationally designed anti-
781      tubulin agent. *Embo Mol Med* **13**, (2021).
782

783 38.  Mehrabi P, *et al.* Time-resolved crystallography reveals allosteric communication
784      aligned with molecular breathing. *Science* **365**, 1167-1170 (2019).
785

786 39.  Pandey S, *et al.* Observation of substrate diffusion and ligand binding in enzyme
787      crystals using high-repetition-rate mix-and-inject serial crystallography. *Iucrj* **8**, 878-
788      895 (2021).
789

790 40.  Butryn A, *et al.* An on-demand, drop-on-drop method for studying enzyme catalysis
791      by serial crystallography. *Nat Commun* **12**, 4461 (2021).
792

793 41.  Schmidt M. Reaction Initiation in Enzyme Crystals by Diffusion of Substrate. *Crystals*
794      **10**, (2020).
795

796 42. Ricart-Ortega M, Font J, Llebaria A. GPCR photopharmacology. *Mol Cell Endocrinol*
797 **488**, 36-51 (2019).
798
799 43. Velema WA, Szymanski W, Feringa BL. Photopharmacology: beyond proof of
800 principle. *J Am Chem Soc* **136**, 2178-2191 (2014).
801
802 44. Leippe P, Frank JA. Designing azobenzene-based tools for controlling
803 neurotransmission. *Curr Opin Struct Biol* **57**, 23-30 (2019).
804
805 45. Muhlethaler T*, et al.* Crystallization Systems for the High-Resolution Structural
806 Analysis of Tubulin-Ligand Complexes. *Methods Mol Biol* **2430**, 349-374 (2022).
807
808 46. Sugahara M*, et al.* Hydroxyethyl cellulose matrix applied to serial crystallography. *Sci*
809 *Rep* **7**, 703 (2017).
810
811 47. James D*, et al.* Improving High Viscosity Extrusion of Microcrystals for Time-
812 resolved Serial Femtosecond Crystallography at X-ray Lasers. *J Vis Exp*, (2019).
813
814 48. Nass K*, et al.* Pink-beam serial femtosecond crystallography for accurate structure-
815 factor determination at an X-ray free-electron laser. *IUCrJ* **8**, (2021).
816
817 49. Weierstall U*, et al.* Lipidic cubic phase injector facilitates membrane protein serial
818 femtosecond crystallography. *Nat Commun* **5**, 3309 (2014).
819
820 50. Zeldin OB, Gerstel M, Garman EF. RADDOSE-3D: time- and space-resolved
821 modelling of dose in macromolecular crystallography. *Journal of Applied*
822 *Crystallography* **46**, 1225-1230 (2013).
823
824 51. White TA*, et al.* CrystFEL: a software suite for snapshot serial crystallography.
825 *Journal of Applied Crystallography* **45**, 335-341 (2012).
826
827 52. White TA*, et al.* Recent developments in CrystFEL. *J Appl Crystallogr* **49**, 680-689
828 (2016).
829
830 53. Gevorkov Y*, et al.* XGANDALF - extended gradient descent algorithm for lattice
831 finding. *Acta Crystallogr A Found Adv* **75**, 694-704 (2019).
832
833 54. Tickle IJ, Flensburg, C., Keller, P., Paciorek, W., Sharff, A., Vonrhein, C., Bricogne,
834 G. . STARANISO.). Cambridge, United Kingdom: Global Phasing Ltd. (2018).
835
836 55. Adams PD*, et al.* PHENIX: building new software for automated crystallographic
837 structure determination. *Acta Crystallogr D Biol Crystallogr* **58**, 1948-1954 (2002).
838
839 56. Winn MD*, et al.* Overview of the CCP4 suite and current developments. *Acta*
840 *Crystallogr D Biol Crystallogr* **67**, 235-242 (2011).
841
842 57. Gorel A, Schlichting I, Barends TRM. Discerning best practices in XFEL-based
843 biological crystallography - standards for nonstandard experiments. *Iucrj* **8**, 532-543
844 (2021).
845

58.    Nogly P*, et al.* Retinal isomerization in bacteriorhodopsin captured by a femtosecond
x-ray laser. *Science* **361**, (2018).

59.    Emsley P, Cowtan K. Coot: model-building tools for molecular graphics. *Acta*
*Crystallogr D Biol Crystallogr* **60**, 2126-2132 (2004).

60.    Bricogne G BE, Brandl M, Flensburg C, Keller P, Paciorek W, Roversi P, Sharff A,
Smart O, Vonrhein C, Womack T BUSTER version 2.10.4.). Cambridge, United
Kingdom: Global Phasing Ltd. (2021).

61.    Salentin S, Schreiber S, Haupt VJ, Adasme MF, Schroeder M. PLIP: fully automated
protein-ligand interaction profiler. *Nucleic Acids Res* **43**, W443-447 (2015).

62.    Smith RHB, Dar AC, Schlessinger A. PyVOL: a PyMOL plugin for visualization,
comparison, and volume calculation of drug-binding sites. *bioRxiv*, 816702 (2019).

63.    Chovancova E*, et al.* CAVER 3.0: a tool for the analysis of transport pathways in
dynamic protein structures. *PLoS Comput Biol* **8**, e1002708 (2012).

64.    Slavov C*, et al.* The ultrafast reactions in the photochromic cycle of water-soluble
fulgimide photoswitches. *Phys Chem Chem Phys* **18**, 10289-10296 (2016).

65.    Jacobson MP, Friesner RA, Xiang Z, Honig B. On the role of the crystal environment
in determining protein side-chain conformations. *J Mol Biol* **320**, 597-608 (2002).

66.    Sastry GM, Adzhigirey M, Day T, Annabhimoju R, Sherman W. Protein and ligand
preparation: parameters, protocols, and influence on virtual screening enrichments. *J*
*Comput Aided Mol Des* **27**, 221-234 (2013).

67.    D.A. Case HMA, K. Belfon, I.Y. Ben-Shalom, S.R. Brozell, D.S. Cerutti, T.E.
Cheatham, III, G.A. Cisneros, V.W.D. Cruzeiro, T.A. Darden, R.E. Duke, G.
Giambasu, M.K. Gilson, H. Gohlke, A.W. Goetz, R. Harris, S. Izadi, S.A. Izmailov,
C. Jin, K. Kasavajhala, M.C. Kaymak, E. King, A. Kovalenko, T. Kurtzman, T.S. Lee,
S. LeGrand, P. Li, C. Lin, J. Liu, T. Luchko, R. Luo, M. Machado, V. Man, M.
Manathunga, K.M. Merz, Y. Miao, O. Mikhailovskii, G. Monard, H. Nguyen, K.A.
O'Hearn, A. Onufriev, F. Pan, S. Pantano, R. Qi, A. Rahnamoun, D.R. Roe, A.
Roitberg, C. Sagui, S. Schott-Verdugo, J. Shen, C.L. Simmerling, N.R. Skrynnikov, J.
Smith, J. Swails, R.C. Walker, J. Wang, H. Wei, R.M. Wolf, X. Wu, Y. Xue, D.M.
York, S. Zhao, and P.A. Kollman. Amber 21. In: *University of California, San*
*Francisco*) (2021).

68.    Jorgensen WL, Chandrasekhar J, Madura JD, Impey RW, Klein ML. Comparison of
Simple Potential Functions for Simulating Liquid Water. *J Chem Phys* **79**, 926-935
(1983).

69.    Maier JA, Martinez C, Kasavajhala K, Wickstrom L, Hauser KE, Simmerling C.
ff14SB: Improving the Accuracy of Protein Side Chain and Backbone Parameters
from ff99SB. *J Chem Theory Comput* **11**, 3696-3713 (2015).

70.    Allner O, Nilsson L, Villa A. Magnesium Ion-Water Coordination and Exchange in
Biomolecular Simulations. *J Chem Theory Comput* **8**, 1493-1502 (2012).

71.     Meagher KL, Redman LT, Carlson HA. Development of polyphosphate parameters
899           for use with the AMBER force field. *J Comput Chem* **24**, 1016-1025 (2003).

901   72.     Salomon-Ferrer R, Gotz AW, Poole D, Le Grand S, Walker RC. Routine Microsecond
902           Molecular Dynamics Simulations with AMBER on GPUs. 2. Explicit Solvent Particle
903           Mesh Ewald. *Journal of Chemical Theory and Computation* **9**, 3878-3888 (2013).

905   73.     Barducci A, Bussi G, Parrinello M. Well-tempered metadynamics: A smoothly
906           converging and tunable free-energy method. *Phys Rev Lett* **100**, (2008).

908   74.     Van der Spoel D, Lindahl E, Hess B, Groenhof G, Mark AE, Berendsen HJC.
909           GROMACS: Fast, flexible, and free. *J Comput Chem* **26**, 1701-1718 (2005).

911   75.     Tribello GA, Bonomi M, Branduardi D, Camilloni C, Bussi G. PLUMED 2: New
912           feathers for an old bird. *Comput Phys Commun* **185**, 604-613 (2014).

913
914

**Acknowledgments**

We are grateful for the excellent support from the PSI Crystallization Facility and the

Macromolecular Crystallography group during growing and testing of crystals at the Swiss

Light Source. The Biomolecular Structure and Mechanism Program of the Life Science Zürich

Graduate School is acknowledged for their academic framework for our graduate students. This

project was founded by the following agencies: The Swiss National Science Foundation project

grants 31003A_179351 (to J.S.), to 310030_197674 (to T.W.), 310030_192566 (to M.O.S.)

and the NCCR:MUST (to C.M. and J.S.), the Swiss Nanoscience Institute SNI #1904 (to M.C.),

Deutsche Forschungsgemeinschaft, Project WA 1850/4-3 (C.S. and J.W.).

**Author contributions**

The project was coordinated and led by J.S. following input on time-resolved serial

crystallographic data acquisition and evaluation from T.W., computer simulations from A.C.,

and time-resolved spectroscopy from J.W. and tubulin structural biology from M.O.S.. The

TD1 protein complex was purified and crystallized by M.Wr. with the help from A.F., N.G.,

and other members of the team during the large-scale preparation phase. The azo-CA4

compound was synthesized by T.Ma.. Crystal injection was optimized by M.Wr., A.F., and

D.J.. The lipidic cubic phase injector was operated and aligned during the beamtime by A.F.

D.Ga., and D.J.. The endstation including the laser system were aligned and operated by

P.J.M.J., D.O., K.N., C.C., J.B., C.B. and C.M. Sample preparation during the beamtime was

done by M.Wr., A.F., M.C., R.S., D.K. and H.G.. Data processing during the beamtime was

initially done using the online data analysis pipeline established by K.N. and D.O. followed by

manual optimization by S.B. and P.S. with instructions from T.W.. The synchrotron setup was

aligned and optimized by F.D., D.J. and W.Wa..

Final structures were refined by M.Wr. and T.W. and interpreted together with J.S.. Quantum

chemical calculations were done by M.F. and D.Gi. under supervision of A.C.. The time-

resolved spectroscopic experiments were done by N.G., T.W., and C.S. and interpreted together

with M.B. and J.W.. The manuscript was written by T.W., M.Wr., M.O.S. and J.S. with further

suggestions from the other authors.

**Competing interest declaration**

The authors declare that they have no competing interests.

953
954

**Figure 1: Tracking ligand release from tubulin using a photochemical affinity switch.** (**A**) The cis-azo-CA4 (yellow sticks) binds to the colchicine site that is formed by secondary structural elements stemming from both the α (blue) and β (cyan) subunits of the αβ-tubulin heterodimer. Illumination with a flash of a laser light triggers the release of *trans*-azo-CA4 (colored in orange, diffusion indicated by including several molecules, cylindrical arrows designate conformational changes of the protein) from its binding pocket. (**B**) Difference electron density maps ($F_{obs}$(light)-$F_{obs}$(dark)) from time-resolved XFEL data were integrated and subjected to Pearson correlation analysis. Three areas of elevated correlation can be identified. The box in the nanoseconds (**1**) corresponds to the relaxed conformation of the ligand observed as a metastable binding pose. Correlations between maps in the microseconds (**2**) are less homogeneous and stretch from 1 μs to opening of the exit pathway at 1 ms. The final snapshots in the milliseconds (**3**) corresponds to the predominantly unliganded protein. (**C**) Plot showing negative density integrated around azo-CA4 with a strong increase after 1 ms indicating release of the ligand. (**D**) Time-resolved structural snapshots of azo-CA4 release. The time arrow depicts the investigated time regime. The panels show from left to right the isomorphous difference maps obtained at 100 ns with changes centered on the ligand, 100 μs with changes centered on the binding pocket, and 100 ms with conformational changes propagating throughout the protein. All panels show isomorphous difference maps in red (negative) and green (positive) at 3 σ. The structure in the given time range (colored in orange) is compared to that of the previous time range (colored in grey). A movie showing the electron changes within the binding pocket over all nine time delays is available as supplementary material.

**Figure 2: Photoisomerization leads to the formation of a metastable binding state. (A)** The two chemical structures of azo-CA4 show the wavelength dependent conformational change between the *cis* (top) and *trans* (bottom) positions. Central to the switch is the N=N azo bond that allows to switch azo-CA4 reversibly and with high efficiency between its high and low affinity stereoisomers, respectively. **(B)** Before laser induced isomerization, *cis*-azo-CA4 (yellow sticks) is bound in the colchicine-site located between the α (blue) and β (cyan) subunits close to the βT7 loop (green) of β-tubulin. Hydrophobic interactions (green lines) between tubulin residues and the A and B rings of *cis*-azo-CA4 as well as hydrophilic interactions (blue dashed lines) towards the ring substituents anchor the ligand within the colchicine- site. **(C)** Within nanoseconds after illumination, *trans*-azo-CA4 (orange sticks) has relaxed into a metastable binding pose with altered interaction network and reduced affinity. For clarity only interacting residues are shown.

**A Binding pocket adaptations over time**

**B Molecular gating of ligand release**

**C Pathway identification**    **D Molecular dynamics simulation**

**Figure 3: Molecular changes in the βT7 gating loop allow ligand release.** (**A**) Within
nanoseconds, the light induced isomerization and relaxation repositions the A ring of azo-CA4
closer to the βT7 loop (green), which acts like a lid on the colchicine-site. Further
reorganizations (black arrows) in the microseconds opens a channel between the βH8 helix and
the βT7 loop. In the millisecond range, the βT7 loop is folded back and packs against residues
of the empty binding pocket. (**B**) Opening of a release pathway through relocation of the βT7
loop in the 1 ms structure is illustrated by the position of βLeu246 (green sticks and surface) at
the indicated time delays. (**C**) The release pathway (red spheres) plotted onto the 1 ms structure.
(**D**) The experimentally deduced unbinding pathway of azo-CA4 is confirmed by MD
simulations. The small red spheres depict the centers of masses of ligand atoms, plotted every
50 frames along the simulation.
1011

**Figure 4: Correlation between colchicine-site dynamics and global tubulin backbone rearrangements**. (**A**) Relaxation of azo-CA4: dark to 100 ns. (**B**) Movements of βT7 loop: 100 ns to 1 ms. (**C**) Collapse of the colchicine-site: 1 ms to 100 ms. The three rows depict conformational changes at different spatial levels: Binding pocket in the upper row, tubulin backbone in the middle row, and longitudinally aligned tubulin dimers in the lower row. Starting structures are shown in grey. Backbone movements within tubulin (middle row) are represented by modevectors between consecutive time delays. Every second Cα-backbone atom in the main secondary structural elements with a displacement of at least 0.5 Å is shown. The N-terminal nucleotide-binding domains, the C-terminal domains, and the intermediate domains for both α and β tubulin are shown in blue, red, and green, respectively. The release pathway (red spheres) was added to illustrate the directional movements of the tubulin subunits. The global conformational changes (lower row) are visualized by three longitudinally aligned tubulin dimers ("Straight" tubulin dimers as found in the main shaft of a microtubule are shown in surface representation (dark gray, α-tubulin; light grey, β-tubulin; PDB ID 5SYE); "Curved" tubulin dimers (α-tubulin in blue and β-tubulin in cyan) at the indicated time delay and in relation to the previous state are shown in black). Deviations demonstrate the conformational plasticity of tubulin during azo-CA4 relaxation in its binding pocket and βT7-loop movements of β-tubulin, as well as a directed curvature adjustment after azo-CA4 release. Arrows indicate the main directions of movements in the indicated time delays.