Representational dynamics of memories for real-life events

Olivier Jeunehomme[1]*, Rebekka Heinen[2]*, David Stawarczyk[1,2], Nikolai Axmacher[2], and

Arnaud D'Argembeau[1,3]

[1] Psychology and Neuroscience of Cognition, Department of Psychology, University of Liège,

Belgium

[2] Department of Neuropsychology, Institute of Cognitive Neuroscience, Faculty of Psychology,

Ruhr University Bochum, Germany

[3] GIGA-CRC In Vivo Imaging, University of Liège, Belgium

* These authors contributed equally to this work

Correspondence: o.jeunehomme@uliege.be (O.J.), rebekka.heinen@rub.de (R.H.),

nikolai.axmacher@rub.de (N.A.), a.dargembeau@uliege.be (A.D.)

Acknowledgments: This work was supported by the University of Liège (Fonds spéciaux

– Crédits sectoriels no. 9893). O. Jeunehomme and A. D'Argembeau are supported by the Fund

for Scientific Research (F.R.S.-FNRS), Belgium.

Competing interests: The authors have declared no competing interest.

Abstract

The continuous flow of experience that characterizes real-life events is not recorded as such in episodic memory. The unfolding of events is represented as a succession of moments of prior experience (referred to as "experience units") that are separated by temporal discontinuities, such that events are temporally compressed when remembered. To unravel the neural basis of this temporal structure of episodic memories, we recorded a series of real-life events using wearable camera technology and then used fMRI to investigate brain activity during the temporal unfolding of these events in memory. We found that, compared to the representation of static scenes in memory, dynamically unfolding memory representations were associated with greater activation of posterior brain regions that support episodic recollection (i.e., the precuneus, posterior cingulate cortex, retrosplenial cortex, and angular gyrus). Strikingly, by analyzing the autocorrelation of brain activity patterns at successive time points throughout the retrieval period, we found that these brain regions showed higher temporal dynamics when remembering the unfolding of events compared to static scenes. Furthermore, these effects were more pronounced when recalling events that included a higher density of experience units (i.e., actions compared to spatial displacements). These results reveal the key role of the posterior medial network in representing the dynamic unfolding of event segments that constitute episodic memories.

Keywords:

Episodic memory, fMRI, Temporal structure, Event segmentation, Scene construction.

INTRODUCTION

Decades of cognitive, neuropsychological, and neuroimaging research on episodic memory have deepened our understanding of the neurocognitive processes supporting the ability to remember past events (Conway, 2009; Moscovitch et al., 2016; Rugg & Vilberg, 2013; Schacter et al., 1998; Tulving, 2002). However, most studies have focused on memory for static stimuli (e.g., lists of words or pictures), so important questions remain about how the dynamic unfolding of events is represented in episodic memory. People make sense of the continuous stream of experience by segmenting it into meaningful units (i.e., events and sub-events), with this segmentation process playing a key role in the temporal structure of memory representations (Clewett & Davachi, 2017; Radvansky & Zacks, 2017; Zacks, 2020). Memories are composed of a succession of event segments (i.e., moments or slices of prior experience) that are organized in chronological order (Anderson & Conway, 1993; Radvansky et al., 2005) and usually include temporal discontinuities in the unfolding of events: Parts of the continuous flow of information that characterized the previous experience are omitted during event recall (Folville et al., 2020; Jeunehomme et al., 2018; Jeunehomme & D'Argembeau, 2020). This structure of memories allows one to mentally represent events in a temporally compressed form, such that it usually takes less time to remember an event than it took for this event to occur in the past (Arnold et al., 2016; Bonasia et al., 2016; Chen et al., 2017; Jeunehomme & D'Argembeau, 2019; Michelmann et al., 2019). To date, however, the neural basis of this temporal structure of memory representations remains unclear.

Memory for real-life events is supported by a specific set of brain regions that includes the medial temporal lobes (e.g., hippocampus, parahippocampal gyrus), cortical midline structures (medial prefrontal cortex, precuneus, posterior cingulate cortex, and retrosplenial cortex), and lateral fronto-temporo-parietal areas (Benoit & Schacter, 2015; Cabeza & St Jacques, 2007; Kim, 2012; McDermott et al., 2009; Svoboda et al., 2006). Within this core

memory network, the hippocampus and medial parietal cortex have been associated with the

process of "scene construction" (i.e., the generation, maintenance, and visualization of

complex spatial scenes), which is considered one of the key components of past event recall

(Hassabis & Maguire, 2007). Supporting this view, it has been shown that episodic memory

retrieval and the imagination of fictitious experiences are associated with overlapping patterns

of brain activation, notably in the hippocampus, parahippocampal gyrus, and retrosplenial

cortex (Hassabis et al., 2007). However, most studies on scene construction required

participants to mentally represent static scenes (e.g., visualizing an old library) (Zeidman et

al., 2015), whereas, as noted above, remembering past events involves a dynamic succession

of moments of prior experience.

A recent fMRI study has provided important insight into the neural mechanisms

underlying the dynamic unfolding of events in episodic memory (Baldassano et al., 2017).

During the initial encoding of events from a movie, event segmentation processes were

evidenced at multiple time scales throughout the cortical hierarchy: While early sensory

regions represented relatively short-lived segments of experience, the posteromedial cortex

and angular gyrus supported more extended episode representations that corresponded closely

to event boundaries identified by human observers. Event boundaries in these higher-level

areas triggered the hippocampus to encode the current situation model into episodic memory,

which was later reinstated in the posteromedial cortex and angular gyrus during memory

retrieval (see also Ben-Yakov & Dudai, 2011; Ben-Yakov & Henson, 2018; DuBrow &

Davachi, 2016).

While these results present an important first step towards understanding the temporal

dynamics of events during their initial experience, the dynamic unfolding of event segments

during recollection remains to be elucidated. As noted above, episodic memories are not

literal reproductions of past events, but instead consist of a succession of event segments that

typically includes temporal discontinuities. The density of the segments of experience that

constitute memories, and therefore the amount of temporal discontinuities within memories,

can vary substantially across events, leading to more or less compressed representations

(Jeunehomme & D'Argembeau, 2019). Here, we aimed to unravel the neural basis of this

dynamic unfolding of the units of experience that constitute episodic memories.

We first sought to examine brain regions that are involved in remembering dynamic

and temporally extended events as compared to static scenes. Memories were prompted using

a paradigm that capitalizes on wearable camera technology to investigate the remembrance of

real-life experiences (Jeunehomme et al., 2018; Nielson et al., 2015; Stawarczyk et al., 2018).

Specifically, participants performed a 40-min walk on a university campus during which they

experienced a series of daily life events, while the content and timing of these experiences

were recorded using a wearable camera (Chow & Rissman, 2017) (Figure 1A). Immediately

after the walk, participants took part in an fMRI session in which they had to remember

events that were cued by pictures that had been taken during their walk (Figure 1B). By

contrasting these episodic memories to a control condition involving the mental representation

of static scenes, we identified the brain regions that specifically underlie the recall of

temporally unfolding events. We focused on the recall of real-life events rather than movies

because the former involve a sense of agency that is absent from passive video viewing but

may play an important role in structuring memory representations, given that a key function

of episodic memory is to remember actions and goal-relevant information (Conway, 2009;

Zimmer et al., 2007).

Another aim of the current study was to compare the neural correlates of episodic

memories that are associated with different event compression rates. Previous studies have

shown that the density of the units of experience that constitute memories, and therefore the

temporal compression of events, is not constant but varies with the content of remembered

events. In particular, memories for actions contain a higher density of experience units than

memories for spatial displacements (Jeunehomme et al., 2018; Jeunehomme & D'Argembeau,

2019, 2020). Indeed, spatial displacements involve a relatively simple structure that is

primarily organized in terms of changes in direction (i.e., turns), whereas actions typically

have a more complex and fine-grained structure in which event segments correspond to

changes in goals and sub-goals (Tversky et al., 2004). This difference in the density of event

segments should be reflected in the dynamics of neural activity when remembering the

unfolding of events.

Our main hypothesis is that the posterior medial network, which is thought to support

the representation of event models (Ranganath & Ritchey, 2012; Ritchey & Cooper, 2020;

Stawarczyk et al., 2021), plays a key role in representing the temporal structure of memories.

Indeed, regions such as the retrosplenial cortex, posterior cingulate cortex, precuneus, and

angular gyrus might contribute to the integration of successive changes in experience when

remembering the unfolding of events, and might also maintain and elaborate integrated event

representations (Ritchey & Cooper, 2020). To test these hypotheses, we not only examined

brain activity over the entire duration of the retrieval period using classical univariate

analyses, but also investigated to what extent multivariate patterns of brain activity change

across time during the unfolding of events in memory. To the extent that memories for actions

contain a higher density of experience units, we expected that they would be associated with

higher temporal dynamics (i.e., lower autocorrelations) of brain activity patterns at successive

time points during the retrieval period.

144
Figure 1: Experimental procedures. (A) Overview of the walk on the campus of the University of Liège.
Locations in which participants performed an action are indicated by blue circles and the paths taken to go to
these locations are indicated by yellow lines. For each event, an example of picture taken by the wearable
camera that was used as cue for the episodic memory task is provided. (B) Trial structure in the episodic memory
conditions (top panel) and in the control condition (bottom panel). Participants were first presented with a cue
specifying the experimental condition. In the episodic memory conditions, a picture taken during the walk of the
participant (corresponding to the beginning of an action or a spatial displacement) was then presented on the
screen. As soon as they identified the moment of the walk depicted by the picture, participants were asked to

remember the unfolding of the event, starting at the precise moment they identified on the picture. In the control
condition (i.e., scene condition), participants were asked to imagine the visual scene depicted by the picture
(which was not part of their walk). For each condition, participants were asked to close their eyes and to press a
response key once they started to perform the task; they did the task with eyes closed until the presentation of a
sound signal (which was presented 12 seconds after their button press). Finally, each trial ended with the
presentation of a Likert scale on which participants rated the temporal dynamics of their mental representation
(i.e., static vs. dynamic). In total, 20 trials were presented for each condition. FMRI analyses focused on the time
interval between the key press and the end of the 12 second period, thus corresponding to the phase of mental
replay.

RESULTS

Behavioral results

Participants successfully identified the presented pictures for nearly all trials ($M = 98.24\%$,

95% CI [98.11%, 98.36%]). Verbal descriptions of episodic memories showed that

participants re-experienced a succession of experience units when recalling the events (see

Materials and Methods). Memories of actions contained more experience units than memories

of spatial displacements, $t = 7.29$, $p < .001$, $\xi = .85$ (Table 1). To assess the density of recalled

experience units, we computed the number of reported units per minute of the actual event

duration (see Materials and Methods). In line with previous studies(Jeunehomme et al., 2018;

Jeunehomme & D'Argembeau, 2019, 2020), the density of experience units was higher when

recalling actions than spatial displacements, $t = 6.77$, $p < .001$, $\xi = .88$. As shown in Table 1,

all memory representations were judged as clear, easy to recall/construct, and associated with

a subjective feeling of re-experiencing the events.

Table 1. Behavioral data for the episodic memory (action, spatial displacement) and control (scene) conditions

	Action	Spatial displacement	Scene
Number of experience units	8.51 [7.51, 9.55]	4.93 [4.08, 5.78]	-
Number of experience units / min	12.42 [10.28, 15.62]	5.02 [4.05, 6.56]	-
Clarity	5.88 [5.48, 6.32]	4.68 [4.31, 5.19]	5.34 [4.80, 5.82]
Easiness	6.09 [5.60, 6.57]	4.93 [4.55, 5.42]	5.32 [5.02, 5.74]
Re-experiencing	5.71 [5.18, 6.20]	4.94 [4.57; 5.43]	-

Note: Data represent 20% trimmed means and their 95% confidence intervals. The score for clarity, easiness and re-experiencing ratings made during the post-scan debriefing ranged from 1 = '*not at all*' to 7 = '*extremely*'.

fMRI results

Episodic memories versus scene representations

We first conducted univariate analyses to identify brain regions within the autobiographical memory network (see Materials and Methods) that showed higher activity when recalling the unfolding of events compared to static scene representations (see Figure 2A, Table 2). Results showed that the recall of actions (action > scene) was associated with increased activation in bilateral parietal areas, including the precuneus, posterior cingulate, retrosplenial cortex, and angular gyrus. Moreover, we found higher activity in the superior/middle frontal gyrus, the posterior middle temporal gyrus, middle occipital cortex, and cerebellum. When looking at memories of spatial displacements (spatial displacement > scene), similar activations were observed in medial parietal areas (i.e., precuneus, posterior cingulate, and retrosplenial cortex). Additional clusters of activation were detected in the right

200   angular gyrus/middle occipital cortex, and bilaterally in the middle frontal gyrus. Figure 2A

201   also illustrates overlapping activations associated with each contrast, which show that the two

202   types of episodic memories were associated with common activation in medial parietal areas,

203   including the precuneus, posterior cingulate cortex, and retrosplenial cortex.

Table 2. Brain regions showing increased activity for episodic memories as compared to scene representations

Brain regions	Side	MNI coordinates			Cluster Size	t
		X	Y	Z		
Actions > scenes						
Precuneus/PCC/RSC	R	12	-62	36	2111	11.46
	R	8	-54	44		9.93
	R	8	-38	26		8.35
	L	-12	-60	20		8.79
	R	12	-62	28		10.99
Middle cingulate gyrus	L	-2	-32	40	28	6.71
Middle/superior frontal gyrus	L	-40	10	54	84	6.40
	R	30	14	58	35	5.85
Middle frontal gyrus	L	-24	30	48	172	4.73
AG	R	44	-68	36	165	5.66
	L	-58	-56	26	21	4.40
MTG/AG	L	-48	-54	20	66	5.64
Middle OC	L	-40	-72	38	64	5.38
MTG	R	62	-52	10	20	5.18
Cerebellum	L	-6	-54	-40	28	4.18
Spatial displacements > Scenes						
Precuneus/PCC/RSC	L	-10	-58	18	1983	11.03
	R	12	-52	20		8.64
	R	4	-44	12		7.75
	R	8	-64	44		7.48
AG/Middle OC	R	42	-70	34	92	537
Middle frontal gyrus	R	28	16	54	31	4.91
	L	-34	6	60	40	4.72
	L	-38	8	54		4.40
	L	-22	26	42	23	4.17

205   **Note:** PCC = Posterior cingulate cortex; RSC = Retrosplenial cortex; OC = Occipital cortex; AG = Angular gyrus;

206   MTG = Middle temporal gyrus.

Figure 2. Brain regions activated when recalling the unfolding of events (univariate analyses). (A) Regions that were more activated when recalling actions and spatial displacements compared to the scene construction condition. (B) Regions showing differential activity when recalling actions and spatial displacements. Activations are displayed at $p < .001$ (uncorrected) within the autobiographical network mask.

Remembering actions versus spatial displacements

Next, we compared the two episodic memory conditions (actions vs. spatial displacements;

see Figure 2B, Table 3). The retrieval of actions was associated with higher activity in the

precuneus, left middle frontal gyrus, right middle temporal gyrus, middle cingulate cortex,

and angular gyrus compared to the retrieval of spatial displacements. The reverse contrast

(spatial displacements > actions) revealed increased activations in the retrosplenial cortex, the

right parahippocampal gyrus, and the right middle occipital gyrus.

Table 3. Brain regions showing differential activity for actions vs spatial displacements

Brain regions	Side	MNI coordinates			Cluster Size	*t*
		X	*Y*	*Z*		
Action > Spatial displacement						
Middle frontal gyrus	L	-32	26	36	34	6.25
Precuneus	L	-2	-62	38	131	5.51
MTG	R	64	-48	6	22	5.45
Middle cingulate gyrus	R	2	-26	38	21	5.39
AG	L	-58	-54	30	24	4.82
Spatial displacement > Action						
RSC	L	-12	-52	12	483	9.44
	R	12	-46	6	363	7.84
PHG/FG	R	20	-34	-10	170	7.68
Middle occipital gyrus	R	42	-74	26	21	4.28

Note: RSC = Retrosplenial cortex; MTG = Middle temporal gyrus; PHG = Parahippocampal gyrus; FG = Fusiform

gyrus; AG = Angular gyrus.

Distinct brain activity patterns across recall conditions

We next examined whether distributed patterns of brain activity in the autobiographical

memory network reflected the content of memories – i.e., whether these patterns differed

systematically between experimental conditions. Using representational similarity analysis,

we compared all recall trials of the three experimental conditions (actions, spatial

displacements, scenes) with trials of the same condition as compared to trials of other

conditions. A repeated measures ANOVA revealed a significant main effect of within versus

between similarities ($F(1,20) = 34.58$, $p = <0.0001$, Figure 3A). Post-hoc contrasts showed

significantly higher within condition correlations compared to between condition correlations

for all three conditions (Figure 3B). Trials within the action condition were more similar to

one another compared to trials of other conditions (within: $M = 0.43$, $SD = 0.13$; between: M

$= 0.41$, $SD = 0.13$; $t(20) = 3.13$, $p_{Bonferroni} = 0.015$), and the same pattern was found for spatial

displacement (within: $M = 0.44$, $SD = 0.15$; between: $M = 0.41$, $SD = 0.13$; $t(20) = 3.13$,

$p_{Bonferroni} = 0.00044$) and for the control condition (within: $M = 0.43$, $SD = 0.15$; between: $M =$

0.39 , $SD = 0.15$; $t(20) = 3.12$, $p_{Bonferroni} = 0.015$). These results show that voxel patterns

indeed reflect the remembered contents and can thus be used to analyze the temporal

dynamics of episodic memories, which we did in the next step.

Figure 3. Content-specific memory representations in the autobiographical memory network. (A)
Distributed patterns of neural activity are more similar when recalling events from the same category than events
from different categories. (B) This higher within- than between-condition similarity is observed for all
experimental conditions.

Higher temporal dynamics of brain activity patterns during recall of high-density events

We next computed temporal autocorrelations between successive patterns of brain activity as

a measure of the temporal dynamics of memories (see Figure 4A). A searchlight analysis

revealed higher temporal dynamics (i.e., lower autocorrelations) during the recall of episodic

memories compared to the representation of static scenes. Compared to the control condition,

the recall of actions was associated with higher temporal dynamics in the precuneus, posterior

cingulate cortex, angular gyrus, and lingual gyrus ($t(20)= 10.8$; $p_{corr} < 0.05$, FWE corrected

using TFCE, Figure 4B/C). This extended cluster also included the middle frontal gyrus,

frontal pole, middle temporal gyrus, and precentral gyrus. Compared to the control condition,

the recall of spatial displacements was associated with higher temporal dynamics in the

posterior cingulate cortex, cuneus, angular gyrus, and lateral occipital cortex ($t(20)= 5.48$; p_{corr}

< 0.05, FWE corrected using TFCE, Figure 6B/C).

Next, we compared the temporal dynamics of the two types of episodic memories and

found higher temporal dynamics for the recall of actions than spatial displacements in the

lateral occipital cortex, precuneus, posterior cingulate, cuneus, and angular gyrus ($t(20) =$

7.06; $p_{corr} < 0.05$, FWE corrected using TFCE, Figure 4B/C). None of the opposite contrasts

showed any significant differences.

Figure 4. Temporal dynamics of patterns of brain activity during the recall period. (A) Within each searchlight sphere (first volume blue circle), we correlated 10 consecutive volumes (starting with the fifth volume after the recall onset to account for the delayed hemodynamic response function), resulting in a 10x10 spearman rho correlation matrix. From this matrix we extracted the mean of the first off-diagonal (i.e., correlations of patterns between all pairs of temporally adjacent volumes). This procedure was applied to all trials of all conditions and for all sphere locations, resulting in three brain maps (one for each condition) indicating the averaged autocorrelation as an (inverse) measure of the temporal dynamics of episodic recall. (B) The searchlight analysis revealed significantly higher temporal dynamics for the action condition compared to the control and spatial displacements conditions. Spatial displacements showed higher temporal dynamics compared to the control condition. (C) The conjunction of higher temporal dynamics for actions compared to both the spatial displacements and control conditions is shown in purple (The conjunction of spatial displacements vs. control and of action vs. control is not shown as it overlaps completely with the action vs. control contrast.).

In addition to this whole-brain analysis, we conducted a second searchlight analysis within the autobiographical network mask that we used for the univariate analyses (Figure 5). This analysis revealed significantly higher temporal dynamics during the recall of actions compared to the control task within the precuneus and posterior cingulate (cluster 1: $t(20) =$

12.3; $p_{corr} < 0.05$, FWE corrected using TFCE), middle temporal gyrus (cluster 2: $t(20) =$

5.52; $p_{corr} < 0.05$, FWE corrected using TFCE), superior frontal gyrus (cluster 3: $t(20) = 4.48$;

$p_{corr} < 0.05$, FWE corrected using TFCE), lateral occipital cortex and angular gyrus (cluster 4:

$t(20) = 4.15$; $p_{corr} < 0.05$, FWE corrected using TFCE). We also found higher temporal

dynamics for the action condition compared to the recall of spatial displacements within one

cluster including the precuneus, lingual gyrus, cuneus, angular gyrus, and the lateral occipital

cortex ($t(20)= 8.75$; $p_{corr} < 0.05$, FWE corrected using TFCE). The contrast between the

spatial displacements and control conditions did not yield significant differences within the

autobiographical network mask.

Figure 5. Searchlight analysis of temporal dynamics within the autobiographical network mask.
(A). This analysis revealed significantly higher temporal dynamics for the action condition compared to both the
control and the spatial displacement condition. (B) Conjunction of higher temporal dynamics for actions compared
to spatial displacements and compared to the control condition shown in green
313

DISCUSSION

The aim of our study was to better understand the neural mechanisms underlying the

dynamic unfolding of events within episodic memory representations. Compared to mental

representations of static scenes, memories for dynamic events elicited greater activation in

medial parietal areas (including the precuneus, posterior cingulate cortex, and retrosplenial

cortex) and in the angular gyrus. Furthermore, when investigating the autocorrelation of

patterns of brain activity throughout the retrieval period, we found that these posterior brain

regions showed greater temporal dynamics when remembering dynamic events compared to

static scenes. These effects were more pronounced for memories that included a higher

density of experience units (i.e., memories for actions compared to spatial displacements).

Memories for real-life events depend on an extended fronto-temporo-parietal network,

but the exact functions of specific regions within this network are not fully understood

(Benoit & Schacter, 2015; Cabeza & St Jacques, 2007; Kim, 2012; McDermott et al., 2009;

Svoboda et al., 2006). Medial parietal areas, including the precuneus, posterior cingulate, and

retrosplenial cortex, are central areas of this network(Svoboda et al., 2006) that share

reciprocal connections (Cavanna & Trimble, 2006) and have been referred to as the posterior

medial episodic system (Baldassano et al., 2017; Ranganath & Ritchey, 2012; Ritchey &

Cooper, 2020). A first contribution of the present study is to show that this posterior network

is more active when recalling the dynamic unfolding of events compared to mentally

representing static scenes. Remembering the unfolding of events involves the retrieval of

moments or slices of prior experience (each moment itself consisting of a bounded set of

various event features) that need to be sequenced and integrated into a stable and coherent

representation (Conway, 2009; D'Argembeau et al., 2021). The posterior medial episodic

network is considered to play a central role in the elaboration and use of event models (i.e.,

models that specify the spatial, temporal, and causal relationships between entities within a

particular context) (Ranganath & Ritchey, 2012; Ritchey & Cooper, 2020; Stawarczyk et al.,

2021), and may thus contribute to organize and integrate the sequence of experience units that

represent the unfolding of events during episodic recall.

A second important aim of this study was to compare the neural correlates of episodic

memories that are composed of different densities of experience units. Consistent with

previous findings (Jeunehomme et al., 2018; Jeunehomme & D'Argembeau, 2019, 2020), our

behavioral results showed that the density of experience units was higher when remembering

actions compared to spatial displacements. Actions typically have a more complex and fine-

grained segmental structure than spatial displacements (Tversky et al., 2004), which is

manifest when mentally representing the unfolding of prior experience during episodic recall.

At the neural level, we found that some regions of the posterior medial episodic network,

including the precuneus, were more active when remembering actions compared to spatial

displacements. Furthermore, and most importantly, we observed higher temporal dynamics

within patterns of activity in the precuneus, as well as the posterior cingulate and occipital

cortex when remembering the course of action events than when remembering spatial

displacements or mentally representing static scenes. The precuneus has been associated with

central aspects of autobiographical memory retrieval, including visuo-spatial imagery, first-

person perspective taking, and an experience of agency (Cavanna & Trimble, 2006).

Interestingly, recent findings suggest that the precuneus plays a causal role in the dynamics of

memory retrieval (Hebscher et al., 2020), and may notably contribute to the temporal

structure and order of moments of past events (Aly et al., 2018; Kwok et al., 2012). The

higher temporal dynamics of activity patterns in the precuneus and other posterior regions that

we observed suggest that these regions are sensitive to changes in recalled content during the

retrieval period and may thus contribute to implementing and organizing transitions between

experience units when representing the temporal unfolding of past events. This interpretation

fits with previous research showing that patterns of activity in posterior medial regions are

less similar across than within segments of a video episode (Baldassano et al., 2017).

We also found that the angular gyrus was more active and showed greater temporal

dynamics when remembering actions compared to spatial displacements. Several studies have

established relationships between activity in the angular gyrus, especially left-lateralized, and

the amount of recollected information and strength of recollection (Gilmore et al., 2015;

Rissman et al., 2016; Vilberg & Rugg, 2007; Wagner et al., 2005). Although the exact role of

this region has been a matter of considerable debate (Cabeza et al., 2012; Seghier, 2012), a

recent unifying framework has proposed that the angular gyrus supports the integration and

representation of the multimodal contextual details that constitute a mnemonic representation

and promote vivid re-experiencing (Ramanan et al., 2018). The present findings suggest that

this integration process contributes to representing the dynamic unfolding of experience units

within episodic memory representations.

Compared to the recall of actions, the recall of spatial displacements was associated

with increased activity in the retrosplenial cortex and parahippocampal gyrus. These regions

are well known for their role in spatial navigation, and notably respond to relevant spatial

elements of the environment such as landmarks (Epstein et al., 2017). Interestingly, the

parahippocampal gyrus coordinates observed in the present study overlap with those reported

in a previous study on the role of the parahippocampal place area on scene perception (Park &

Chun, 2009). Evidence suggests that the parahippocampal place area and the retrosplenial

cortex might play complementary functions in building a coherent and detailed representation

of a scene across several viewpoints (Epstein et al., 2003; Park & Chun, 2009). Specifically, it

has been proposed that when viewpoint changes occur continuously in a scene, the

parahippocampal place area is sensitive to changes in physical details across each view,

thereby representing scenes as a succession of individual representations of each view rather

than a unique and integrated representation of multiple views. On the other hand, the

retrosplenial cortex links the current and previous views of the same scene, creating an

integrated representation of this scene over changes in visual perspective (Park & Chun,

2009). In the current study, it is therefore possible that the parahippocampal gyrus and the

retrosplenial cortex were more active when remembering the unfolding of spatial

displacements compared to actions because the former implied more important updating and

integration of successive views of the surrounding environment when mentally travelling

from one particular moment of past experience to the next.

The critical role of the hippocampus in the retrieval of episodic memories is now well

established (Moscovitch et al., 2016). This region is considered to play an important role in

binding the various elements that constitute memories, including information about the

sequence of events (Bellmund et al., 2020; Clewett et al., 2019; Lee et al., 2020). Therefore,

one could have expected that, in the present study, the hippocampus would be differentially

activated when remembering events with different densities of experience units, but this is not

what we found. We have no ready explanation for this null result, but it is worth noting that in

previous work on memory for dynamic events, hippocampal activity was observed during

encoding but not during event recall (Baldassano et al., 2017). One possible explanation could

be that hippocampal activity during retrieval is transient in nature (Vilberg & Rugg, 2012) and

thus was not captured by our analysis. Alternatively, the episodic segments supported by the

hippocampus may be substantially longer (~1 minute) (Baldassano et al., 2017) than the

relatively short recall periods that we analyzed (12 seconds). Future studies should analyze

the contribution of the hippocampus to the temporal dynamics of memories in more detail.

From a methodological perspective, it would also be interesting in future studies to

compare our method of analysis of the temporal dynamics of brain activity patterns with other

methods that detect state boundaries in activity patterns using hidden Markov models (HMM)

(Baldassano et al., 2017) or greedy state boundary search (GSBS) (Geerligs et al., 2021).

HMM and GSBS were not applicable to our data for several reasons. First, these methods

have been applied to fMRI recordings ranging from 8 minutes (Geerligs et al., 2021) up to 45

minutes (Baldassano et al., 2017), whereas recall trials in our study were limited to 12

seconds. State boundaries might not be detectable within such short trials, as Geerligs et al.

(2021) found that the algorithms start to struggle with short state durations. Second, HMM

and GSNS methods are applied to group data, detecting state boundaries across participants.

The real-life events experienced by our participants were not completely comparable across

participants, contrary to the standardized movies used in the aforementioned studies. In future

studies, applying state detection algorithms on the recall of real-life events while taking these

issues into account would be a promising next step for further characterizing the

representational dynamics of memories.

Here, we investigated the temporal dynamics of brain activity as a function of the

density of recalled experience units by comparing memories for two types of events that have

been repeatedly shown to differ in terms of event compression rates (i.e., actions vs spatial

displacements) (Jeunehomme et al., 2018; Jeunehomme & D'Argembeau, 2019). However, it

should be noted that, in addition to these overall differences between event types,

compression rates can be voluntarily modulated according to retrieval goals (Bellmund et al.,

2020). For example, when people scan their memory for a particular piece of information,

they skip more event segments and thus compression rates are higher than when they try to

mentally relive the events (Michelmann et al., 2019, 2021). An interesting avenue for future

research would be to use the method we developed to analyze the temporal dynamics of brain

activity patterns to study the neural basis of this flexible modulation of event compression

during episodic retrieval.

In sum, the present study provides first insights on the neural correlates of the dynamic

succession of the experience units that constitute memories for real-life events. Our results

showed that remembering the unfolding of events activates posterior brain structures

(including the precuneus, posterior cingulate cortex, retrosplenial cortex, and angular gyrus)

that support event models and may contribute to the representation and integration of episodic

details to form coherent representations of the succession of moments of prior experience.

Furthermore, an important finding of the current study is that changes in patterns of activity

within these regions reflect differences in the density of recalled experience units, suggesting

that the posterior medial network plays an important role in the temporal dynamics of event

recall and may notably contribute to implement transitions between the units of experience

that represent the unfolding of events.

MATERIALS AND METHODS

Participants

Twenty-one right-handed young students (9 women, mean age = 22.29 years, SD = 2.12, range = 18-27 years) from the University of Liège took part in this study. All participants were highly familiar with the university campus where the pre-scan walk took place. None of the participants reported a current use of psychoactive medication or a history of a neurological or psychiatric disorder. Five additional participants were tested but were excluded from the analyses due to noncompliance with the task, either during the pre-scan walk (one participant) or in the scanner (four participants). All participants gave their written informed consent to take part in the study, which was approved by the ethics committee of the Medical School of the University of Liège and was performed in accordance with the ethical standards described in the Declaration of Helsinki (1964).

Materials and procedure

Pre-scan session

Immediately before scanning, participants performed a series of daily-life activities at different locations on the campus of the University of Liège while wearing an Autographer (OMG Life Ltd., Kidlington, UK) tied around their neck. The Autographer is a small wearable camera that automatically and silently takes a continuous series of pictures from the first-person perspective through a 136° fish-eye lens, depending on several electronic sensors (e.g., temperature, accelerometer, magnetometer, etc.). This device has been used in recent neuroimaging and behavioral studies investigating episodic memories for real-life events (Chow & Rissman, 2017). To record the entire set of events as completely as possible, the Autographer was set to the fastest capture rate (approximatively 10 pictures per minute).

The entire set of events was designed to simulate activities that college students would

perform in their daily life. It consisted of two kinds of events: goal-directed actions at specific

locations, and spatial displacements that did not involve any particular action other than

walking from a place to another. These two kinds of events correspond respectively to what

Tversky et al. (2004) referred to as "events by hands" (i.e., interactions with objects; e.g.,

buying a snack) and "events by feet" (i.e., changes of location; e.g., going from the laboratory

building to the bus stop). We initially planned to examine a third kind of events that consisted

in waiting at specific locations without doing any particular action (e.g., waiting at the bus

stop); this condition was included as we assumed it would involve fewer perceptual changes

than the two previous types of events. However, preliminary behavioral analyses revealed that

the variance in recalled elements (i.e., information about persons, objects, thoughts,

perceptual details, and spatial details) was much higher in memories for waiting moments

than in memories for actions and spatial displacements, showing that the former condition did

not involve a homogenous set of events (the waiting moments took place in public spaces,

such as a bus stop, where many different events could occur while participants were waiting).

Thus, we do not report analyses about waiting moments (but note that these moments were

modeled in the design matrix; see below) as this heterogeneity in recalled contents make the

results difficult to interpret.

Participants were instructed to conduct the following series of successive actions:

First, they left the laboratory, went to the bus stop, and stayed there as if they were waiting for

the bus. After having waited approximately two minutes, they went to the campus shop to buy

a snack of their choice. Next, they went to the elementary school near the University campus

and stayed in front of the school for approximately two minutes as if they were waiting for a

kid. After that, participants went to the Speech and Language Therapy Service to post a letter

in a particular mailbox. Having posted the letter, they came back to the laboratory building to

print a document and to wait in the laboratory building hall (again for approximately two

minutes) as if they were waiting for an appointment with a professor. Then, they went to a

cafeteria to buy a drink of their choice, after which they chose a table inside the cafeteria to

wait as if they were waiting for a friend. Next, participants went to a classroom and waited

outside as if they were waiting for the beginning of a course. Finally, they were instructed to

return to the laboratory and answered an email sent by the experimenter during the walk.

Before starting the walk, participants were given the letter that they should post, as

well as 5 euros for purchasing the snack and the drink. They were told that they could take the

route they wanted to go from each location to the next, but not to perform any additional

actions during spatial displacements or waiting moments (e.g., using their smartphones,

listening to music, or counting down time). In addition, participants were asked to avoid

obstructing the lens of the camera and to behave as naturally as possible. Finally, they were

informed that pictures taken during their walk would be presented during a subsequent fMRI

session, but no mention was made of the memory task. Instead, we used a cover story in

which participants were told that they would have to perform aesthetic judgments on the

pictures in the scanner.

FMRI Session

Immediately after the walk on campus, participants took part in an fMRI session that

involved the recall of events experienced during the walk on campus, as well as a control

condition. The entire task comprised 80 trials (20 trials in each condition). The order of trial

presentation was randomized with the constraint that two subsequent trials of the same

condition were not presented in direct succession or with an interval of more than five trials.

Episodic memory tasks. In the episodic memory conditions (i.e., action, spatial

displacement, and waiting conditions), participants were asked to remember the unfolding of

actions, spatial displacements, or waiting periods. Each trial started with a fixation cross (with

a variable duration ranging between one and three seconds) followed by the presentation of a

written experimental instruction (for one second), specifying the task that participants should

perform on the trial (i.e., remembering the unfolding of an event or remembering a particular

moment of waiting). Then, a picture taken during the pre-scan walk of the participant was

presented, which corresponded to the beginning of either a particular goal-directed action, a

spatial displacement, or a waiting moment. Experimental instructions specified that each

picture would be displayed several times during the fMRI session. In total, 20 trials were

presented in each of the episodic memory conditions (five pictures that were each repeated

four times).

For each picture, participants were first asked to identify the corresponding moment of

the walk (without starting to remember the corresponding event); all pictures were presented

to the participants before the scan session to ensure that they would be able to easily identify

the corresponding moment of their walk during the fMRI session. As soon as they identified

the moment depicted, participants were instructed to close their eyes and then to press a

response key to indicate that they began their mental replay (their button press was confirmed

by a sound signal). Depending on the condition, they were asked to mentally replay either the

unfolding of the event (for actions and spatial displacements) or the waiting period, in as

much detail as possible, starting at the precise moment they identified on the picture.

Participants were told that their mental replay could be of anything that they encountered

during the walk (e.g., people, objects, actions, thoughts, etc.). Moreover, in the unfolding

conditions, it was specified that the task was not to proceed as far as possible in the recall of

the walk or action, but rather to be as precise and detailed as possible. In the waiting

condition, participants were told that they should only remember what they experienced

during the moments in which they waited at a particular location (without remembering the

unfolding of what happened after these waiting moments). The mental replay phase lasted 12

seconds after which a sound signal prompted participants to stop their mental replay and to

open their eyes.

Following the mental replay phase, a one-second fixation cross was shown,

immediately followed by the presentation of a four-point Likert scale assessing the extent to

which participants' mental replay was static (meaning that it was focused on a particular

moment of past experience) or dynamic (meaning that it involved the unfolding of several

moments of past experience; see Figure 1B). Responses were self-paced with a maximum of

five seconds to respond.

Control task. In the control task (scene condition), participants had to form mental

images of static scenes. The structure and timing of the control condition was identical to the

episodic memory conditions: Each trial started with a fixation cross, followed by written

instructions. Then a picture of a scene was presented in the middle of the screen. Each picture

was selected to represent a specific location on the campus of the University of Liège that

participants did not visit during their walk. In total, five different pictures were presented for

four times each, resulting in 20 trials. As in the episodic memory conditions, each picture was

also presented to the participants before the scan. After having examined the picture,

participants were asked to close their eyes and then to press the response key to indicate that

they began forming a mental representation of the scene (a sound signal confirmed their

button press). They were instructed to maintain their mental representation of the scene in as

much detail as possible until the presentation of a second sound signal (i.e., after 12 seconds).

It was specified that their task was not to retrieve personal memories (e.g., past events

experienced at the presented location or past experiences that happened at similar locations)

but only to mentally represent the scene that was depicted on the picture. The control

condition trials ended with the presentation of a four-point Likert scale assessing to what

extent each trial involved a static representation or the unfolding of several moments (as in the

episodic memory conditions).

Post-scan debriefing

Immediately after the fMRI session, participants were shown the 20 pictures presented

during the scanning session (i.e., 5 pictures per condition) and, for each picture, they were

asked to describe the content of the mental representation they formed in the scanner (i.e.,

their memories or scene representations). Verbal reports were recorded using a digital audio

recorder. Furthermore, in the episodic memory conditions, each verbal report was followed by

a review of the pictures that had been taken during the walk and participants were instructed

to select the picture that best corresponded to their mental representation of the last moment

they remembered while mentally replaying the unfolding of the event in the scanner (i.e., the

moment they remembered when hearing the second sound signal). These tasks allowed us to

determine (a) whether the participants correctly performed all conditions in the scanner and

(b) to estimate the density of experience units recalled (see Behavioral data analyses).

Finally, participants were asked to rate the overall clarity of the memory or scene they

formed in response to each picture (from 1 = *not clear at all*, to 7 = *extremely clear*), as well

as the easiness of remembering/imagining (from 1 = *not easy at all*, to 7 = *extremely easy*).

Furthermore, for trials of the episodic memory conditions, participants were asked to estimate

their overall feeling of re-experiencing the unfolding of events (from 1 = *not at all*, to 7 =

completely).

Behavioral data analyses

Verbal descriptions of the memories reported during the post-scan interview consisted

of a succession of distinct moments of past experience (referred to as experience units; see

(Jeunehomme et al., 2018; Jeunehomme & D'Argembeau, 2019, 2020). For example, a

typical verbal report would start by "I got out of the building" (first experience unit), "then I

walked along the parking" (second experience unit), "then, I saw a man I know on the

opposite side of the road" (third experience unit), and so on. The segmentation of verbal

reports into distinct experience units was performed by the first author based on indications of

transitions or temporal discontinuities between reported moments of experience (i.e., verbal

clues such as "then", "next", and "after that" or moments of silence); most of the time,

transitions between experience units involved significant changes in actions, environmental

elements, and/or thoughts (e.g., "I saw a sheet of paper on the ground just in front of the door

of the cafeteria," followed by "Once inside of the cafeteria, I was happy to see that there were

few students queuing"). Although in most cases this segmentation procedure was obvious, we

nevertheless assessed its reliability by asking a second trained rater to independently segment

a random selection of 20% of verbal reports. The Intraclass Correlation Coefficient (ICC)

computed on the number of experience units that were identified within verbal reports showed

an excellent agreement between the two raters (ICC = 0.91).

In addition to computing the number of experience units reported by participants, we

estimated the density of recalled experience units in the two episodic memory conditions by

using temporal information associated with the pictures acquired during the walk. More

specifically, for each trial, we estimated the actual event duration by calculating the amount of

time between the beginning of each event and the end of the event. The beginning

corresponded to the picture presented on the computer screen during each fMRI trial. The end

of the event corresponded to the picture selected by participants as the last moment of the

experience they remembered in the scanner. Then, we estimated the density of recalled

experience units as the number of experience units reported per minute of the actual event

duration (Jeunehomme et al., 2018; Jeunehomme & D'Argembeau, 2019).

Behavioral data were analyzed using robust statistical methods (i.e., Yuen's *t* test)

(Field & Wilcox, 2017) because the assumptions underlying classical inferential methods

(normality and homoscedasticity) were violated for our main measures of interest (i.e., the

number of reported experience units per minute of the actual event duration). These methods

perform well in terms of type I error control and statistical power, even when the normality

and homoscedasticity assumptions are violated (Erceg-Hurn & Mirosevich, 2008). All

descriptive statistics refer to the 20% trimmed means and their 95% confidence intervals.

fMRI data acquisition

All MRI data were acquired on a whole-body 3T scanner (Magnetom Prisma, Siemens

Medical Solutions, Erlangen, Germany) operated with a 20-channel receiver head coil.

Multislice T2*-weighted functional images were acquired using a gradient-echo-planar

imaging (EPI) sequence (TR = 1170 ms, TE = 30 ms, FA 90°, matrix size $72 \times 72 \times 36$, voxel

size $3 \times 3 \times 3$ mm³). Thirty-six 3 mm thick transverse slices (FOV 216×216 mm²) were

acquired, with a 25% interslice gap, covering the whole brain. Around 1400 functional

volumes were obtained for the entire task and we dropped the five initial volumes to avoid T1

saturation effect. After image acquisitions, we generated field maps using a double echo

gradient-recalled sequence (TR = 634 ms, TE = 10 and 12.46 ms, FoV = 192×192 mm², $64 \times$

64 matrix, 40 transverse slices with 3 mm thickness and 25% gap, flip angle = 90°, bandwidth

= 260 Hz/pixel) that we used to correct echo-planar images for geometric distortion due to

field inhomogeneities. Finally, we acquired a structural MR scan [T1-weighted 3D

magnetization-prepared rapid acquisition gradient echo (MP-RAGE) sequence, TR = 1900

ms, TE = 2.19 ms, FOV 256 × 240 mm², matrix size 256 × 240 × 224, voxel size 1 × 1 × 1

mm]. All experimental stimuli were presented on a screen positioned at the rear of the

scanner, which the subject could see through a mirror mounted on the head coil.

fMRI data preprocessing

Image preprocessing was performed using the SPM 12 software (Wellcome

Department of Imaging Neuroscience, http//www.fil.ion.ucl.ac.uk/spm) implemented in

MATLAB R2017a. EPI time series were corrected for motion and distortion artefacts using

Realign and Unwarp (Andersson et al., 2001) with the Fieldmap Toolbox (Hutton et al.,

2002). The mean realigned EPI image was coregistered with the structural T1 image, and the

coregistration parameters were applied to the realigned EPI time series. The T1 image was

segmented into gray matter, white matter, and cerebrospinal fluid, using the unified

segmentation approach (Ashburner & Friston, 2005), and we used the normalization

parameters obtained from the segmentation procedure to normalize the functional images to

MNI space (2×2×2 mm voxels). Finally, we performed a spatial smoothing of functional

images using an 8 mm FWHM Gaussian kernel.

For the representational similarity analyses, volumes coregistered to the first scan of

each subject were used to perform the searchlight in the native space of each subject. To

exclude variance explained by motion, regressors from motion correction were used as

predictors in a GLM for the voxel time series in every voxel. The resulting residuals were

then used as the final input for the analyses. To exclude that the searchlight results are

influenced by condition differences in motion inside the MRI we analyzed the motion

parameters resulting from fMRI preprocessing. As the assumptions for an analysis of variance

(ANOVA) were violated we evaluated motion effects with a non-parametric permutation test

using ez-Perm (Lawrence, 2016) in R (R Core Team, 2021). No significant differences of the

motion regressor between the three conditions were found.

fMRI data analyses

Univariate analyses

Univariate analyses were performed using SPM 12. For each participant, BOLD

responses were modeled at each voxel, using a general linear model (GLM). The

memory/scene representation phase of each trial was modeled separately for each condition

(i.e., action, spatial displacement, waiting, and scene representation) as epoch-related

responses (beginning at the key press and ending at the second sound signal after 12 seconds)

and convolved with the canonical hemodynamic response function to create the regressors of

interest. Trials for which participants did not press the response button to indicate that they

identified the picture and trials for which ratings indicated that they did not perform the

requested task were modelled in a separate regressor of no interest. This corresponds to trials

for which they provided a rating of 1 or 2 in the two episodic memory conditions (indicating

that they mentally represented only one single moment of past experience), or a rating of 3 or

4 in the scene condition (indicating that they mentally represented the unfolding of several

moments of past experience). Experimental phases corresponding to the presentation of the

pictures and the presentation of the Likert scales were also modelled as epoch-related

responses with two regressors of no interest (i.e., a single regressor for each phase across all

conditions). Finally, we also modelled the two beeps presented respectively at the beginning

and the end of each task as an event-related response, with a single regressor across all

conditions. Finally, the design matrix included the realignment parameters to account for any

residual movement-related effect. The canonical hemodynamic response function was used

and a high-pass filter was implemented using a cutoff period of 128 seconds to remove low-

frequency drifts from the time series.

A series of linear contrasts was computed to identify the brain regions involved in the

processing of episodic versus scene representations, and in the retrieval of actions versus

spatial displacements. More specifically, we first investigated brain regions showing higher

activity during each episodic memory condition compared to scene representation (i.e., Action

> Scene; Spatial displacement > Scene). Next, we examined brain regions showing

differential activity during the retrieval of the goal-directed actions and spatial displacements

(Action > Spatial displacement, and vice versa).

The contrasts of interest were first computed for each participant and were then

entered into random effects one-sample *t*-tests. For all analyses, statistical parametric maps

(SPMs) were thresholded at $p < .001$, uncorrected, and statistical inferences were corrected

for multiple comparisons at the voxel level ($p < .05$, FWE-corrected) using a small volume

correction (SVC) within a priori regions of interest. These a priori regions of interest

corresponded to brain areas that have been previously related to autobiographical memory, as

determined by a meta-analysis of 84 studies performed with Neurosynth

(www.neurosynth.org) (Yarkoni et al., 2011). Only clusters with a size of k > 15 voxels are

reported.

Representational Similarity Analysis

In addition to the univariate analyses, we conducted a representational similarity

analysis, a well-established method to study neural representational structures in brain activity

patterns (Kriegeskorte & Kievit, 2013). First, we used this method to assess whether the

different conditions (i.e., mental representations of Actions, Spatial Displacements, and

Scenes) showed distinct patterns of voxel activity during the recall period. This would

substantiate further analysis of the temporal dynamics of multi-voxel pattern activity (see

below). We computed a single-trial GLM for each subject with one regressor for each recall

period of 12 seconds, resulting in 60 beta weights for each subject. Within the

autobiographical network mask, we tested whether trials within one condition were more

similar to each other compared to trials of other conditions. We calculated the cross-

correlation between all trials using spearman rank correlations, resulting in a 60 x 60 matrix.

Correlation coefficients between all pairs of trials in each condition were used to build the

mean within-condition correlation coefficient. Correlations of each trial with itself were

excluded before computing the mean. To generate between-condition correlation coefficients,

trials of one condition were correlated to trials of the other conditions. Differences for within

and between condition similarities were analyzed using a repeated measures ANOVA

implemented in the ez-package (Lawrence, 2016) in R (R Core Team, 2021).

Analysis of temporal dynamics

Next, we investigated the temporal dynamics of patterns of brain activity during the

memory/scene construction period by computed correlations between consecutive MRI

volumes during the recall phase. We expected dynamic and high-density events to show

higher differences in brain activity patterns across time, as indicated by smaller volume to

volume correlations. The analysis was limited to grey matter voxels by applying a subject

specific grey matter mask segmented using Freesurfer (version 6) (Fischl, 2012). We

constructed a sphere with a radius of 6 voxels which was centered at each voxel. If the sphere

consisted of fewer than 20 voxels no further analysis was performed at this location. At each

valid sphere location, 10 consecutive volumes from the recall period, starting with the fifth

volume after the recall onset to account for the delay of the hrf, were used to compute a 10x10

cross-correlation matrix (Figure 4A). To analyze consecutive changes in voxel patterns, the

first off-diagonal was used to extract Spearman's rank correlations of each volume with the

next (i.e., volume 5 was correlated with volume 6; volume 6 with volume 7; etc.). Then, we

computed the mean over the first off-diagonal for the resulting 9 volume-to-volume

correlations. This was done for each trial in each condition. In a final step, the mean over all

fisher-z-transformed volume-to-volume correlations in all trials per condition was saved at the

current sphere location. Thus, three correlation coefficient maps, one per condition, were

created for each subject. For group level analysis, the subject-wise correlation coefficient

maps were normalized to the FSL standard 2mm 152 MNI template using a non-linear

registration in FSL (version 5.0.9, http://fsl.fmrib.ox.ac.uk/fsl/fslwiki/).

On the group level, the correlation coefficient maps were used to identify regions with

consistent differences in correlations between the conditions. Using *randomise* in FSL

(Winkler et al., 2014), a non-parametric version of the paired one-sample t-test for each

condition pairing was conducted. *Randomise* analysis was performed using 10,000 random

sign-flips and threshold-free cluster enhancement (TFCE) (Smith & Nichols, 2009) was used

to identify significant clusters. TFCE has been shown to be more sensitive than clusterwise

thresholding methods by taking both voxelwise signal strength and extent of neighboring

voxels into account without relying on an absolute threshold (Smith & Nichols, 2009). All

results reported are FWE corrected based on TFCE and thresholded at $p < 0.05$ to control for

multiple comparisons. We used the Harvard-Oxford Cortical Structural atlas (Desikan et al.,

2006) for anatomical labels.

References

Aly, M., Chen, J., Turk-Browne, N. B., & Hasson, U. (2018). Learning Naturalistic Temporal Structure in the Posterior Medial Network. *Journal of Cognitive Neuroscience*, *30*(9), 1345–1365. https://doi.org/10.1162/jocn_a_01308

Anderson, S. J., & Conway, M. A. (1993). Investigating the structure of autobiographical memories. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, *19*(5), 1178–1196. https://doi.org/10.1037/0278-7393.19.5.1178

Andersson, J. L., Hutton, C., Ashburner, J., Turner, R., & Friston, K. (2001). Modeling geometric deformations in EPI time series. *Neuroimage*, *13*(5), 903–919.

Arnold, A. E. G. F., Iaria, G., & Ekstrom, A. D. (2016). Mental simulation of routes during navigation involves adaptive temporal compression. *Cognition*, *157*, 14–23. https://doi.org/10.1016/j.cognition.2016.08.009

Ashburner, J., & Friston, K. J. (2005). Unified segmentation. *Neuroimage*, *26*(3), 839–851.

Baldassano, C., Chen, J., Zadbood, A., Pillow, J. W., Hasson, U., & Norman, K. A. (2017). Discovering Event Structure in Continuous Narrative Perception and Memory. *Neuron*, *95*(3), 709-721.e5. https://doi.org/10.1016/j.neuron.2017.06.041

Bellmund, J. L. S., Polti, I., & Doeller, C. F. (2020). Sequence memory in the hippocampal-entorhinal region. *Journal of Cognitive Neuroscience*, *32*, 2056–2070.

Benoit, R. G., & Schacter, D. L. (2015). Specifying the core network supporting episodic simulation and episodic memory by activation likelihood estimation. *Neuropsychologia*, *75*, 450–457.

Ben-Yakov, A., & Dudai, Y. (2011). Constructing Realistic Engrams: Poststimulus Activity of Hippocampus and Dorsal Striatum Predicts Subsequent Episodic Memory. *Journal of Neuroscience*, *31*(24), 9032–9042. https://doi.org/10.1523/JNEUROSCI.0702-11.2011

Ben-Yakov, A., & Henson, R. N. (2018). The Hippocampal Film Editor: Sensitivity and

Specificity to Event Boundaries in Continuous Experience. *The Journal of*

Neuroscience: The Official Journal of the Society for Neuroscience, *38*(47), 10057–

10068. https://doi.org/10.1523/JNEUROSCI.0524-18.2018

Bonasia, K., Blommesteyn, J., & Moscovitch, M. (2016). Memory and navigation:

Compression of space varies with route length and turns. *Hippocampus*, *26*(1), 9–12.

https://doi.org/10.1002/hipo.22539

Cabeza, R., Ciaramelli, E., & Moscovitch, M. (2012). Cognitive contributions of the ventral

parietal cortex: An integrative theoretical account. *Trends Cogn Sci*, *16*(6), 338–352.

Cabeza, R., & St Jacques, P. (2007). Functional neuroimaging of autobiographical memory.

Trends Cogn Sci., *11*(5), 219–227.

Cavanna, A. E., & Trimble, M. R. (2006). The precuneus: A review of its functional anatomy

and behavioural correlates. *Brain*, *129*(3), 564–583.

https://doi.org/10.1093/brain/awl004

Chen, J., Leong, Y. C., Honey, C. J., Yong, C. H., Norman, K. A., & Hasson, U. (2017).

Shared memories reveal shared structure in neural activity across individuals. *Nature*

Neuroscience, *20*(1), 115–125. https://doi.org/10.1038/nn.4450

Chow, T. E., & Rissman, J. (2017). Neurocognitive mechanisms of real-world

autobiographical memory retrieval: Insights from studies using wearable camera

technology. *Annals of the New York Academy of Sciences*, *1396*(1), 202–221.

https://doi.org/10.1111/nyas.13353

Clewett, D., & Davachi, L. (2017). The ebb and flow of experience determines the temporal

structure of memory. *Current Opinion in Behavioral Sciences*, *17*, 186–193.

https://doi.org/10.1016/j.cobeha.2017.08.013

820 Clewett, D., DuBrow, S., & Davachi, L. (2019). Transcending time in the brain: How event

821   memories are constructed from experience. *Hippocampus*, *29*(3), 162–183.

822   https://doi.org/10.1002/hipo.23074

823 Conway, M. A. (2009). Episodic memories. *Neuropsychologia*, *47*(11), 2305–2313.

824 D'Argembeau, A., Jeunehomme, O., & Stawarczyk, D. (2021). Slices of the past: How events

825   are temporally compressed in episodic memory. *Memory*, *0*(0), 1–6.

826   https://doi.org/10.1080/09658211.2021.1896737

827 Desikan, R. S., Ségonne, F., Fischl, B., Quinn, B. T., Dickerson, B. C., Blacker, D., Buckner,

828   R. L., Dale, A. M., Maguire, R. P., Hyman, B. T., Albert, M. S., & Killiany, R. J.

829   (2006). An automated labeling system for subdividing the human cerebral cortex on

830   MRI scans into gyral based regions of interest. *NeuroImage*, *31*(3), 968–980.

831   https://doi.org/10.1016/j.neuroimage.2006.01.021

832 DuBrow, S., & Davachi, L. (2016). Temporal binding within and across events. *Neurobiology*

833   *of Learning and Memory*, *134*, 107–114. https://doi.org/10.1016/j.nlm.2016.07.011

834 Epstein, R., Graham, K. S., & Downing, P. E. (2003). Viewpoint-specific scene

835   representations in human parahippocampal cortex. *Neuron*, *37*(5), 865–876.

836   https://doi.org/10.1016/s0896-6273(03)00117-x

837 Epstein, R., Patai, E. Z., Julian, J. B., & Spiers, H. J. (2017). The cognitive map in humans:

838   Spatial navigation and beyond. *Nature Neuroscience*, *20*(11), 1504–1513.

839   https://doi.org/10.1038/nn.4656

840 Erceg-Hurn, D. M., & Mirosevich, V. M. (2008). Modern robust statistical methods: An easy

841   way to maximize the accuracy and power of your research. *The American*

842   *Psychologist*, *63*(7), 591–601. https://doi.org/10.1037/0003-066X.63.7.591

Field, A. P., & Wilcox, R. R. (2017). Robust statistical methods: A primer for clinical

psychology and experimental psychopathology researchers. *Behaviour Research and*

Therapy, *98*, 19–38. https://doi.org/10.1016/j.brat.2017.05.013

Fischl, B. (2012). FreeSurfer. *NeuroImage*, *62*(2), 774–781.

https://doi.org/10.1016/j.neuroimage.2012.01.021

Folville, A., Jeunehomme, O., Bastin, C., & D'Argembeau, A. (2020). The impact of age on

the temporal compression of daily life events in episodic memory. *Psychology and*

Aging, *35*(4), 484–496. https://doi.org/10.1037/pag0000456

Geerligs, L., van Gerven, M., & Güçlü, U. (2021). Detecting neural state transitions

underlying event segmentation. *NeuroImage*, *236*, 118085.

https://doi.org/10.1016/j.neuroimage.2021.118085

Gilmore, A. W., Nelson, S. M., & McDermott, K. B. (2015). A parietal memory network

revealed by multiple MRI methods. *Trends in Cognitive Sciences*, *19*(9), 534–543.

https://doi.org/10.1016/j.tics.2015.07.004

Hassabis, D., Kumaran, D., & Maguire, E. A. (2007). Using imagination to understand the

neural basis of episodic memory. *J Neurosci*, *27*(52), 14365–14374.

Hassabis, D., & Maguire, E. A. (2007). Deconstructing episodic memory with construction.

Trends Cogn Sci., *11*(7), 299–306.

Hebscher, M., Ibrahim, C., & Gilboa, A. (2020). Precuneus stimulation alters the neural

dynamics of autobiographical memory retrieval. *NeuroImage*, *210*, 116575.

https://doi.org/10.1016/j.neuroimage.2020.116575

Hutton, C., Bork, A., Josephs, O., Deichmann, R., Ashburner, J., & Turner, R. (2002). Image

distortion correction in fMRI: A quantitative evaluation. *Neuroimage*, *16*(1), 217–240.

Jeunehomme, O., & D'Argembeau, A. (2019). The time to remember: Temporal compression

and duration judgements in memory for real-life events. *Quarterly Journal of*

Experimental Psychology, *72*(4), 930–942.

https://doi.org/10.1177/1747021818773082

Jeunehomme, O., & D'Argembeau, A. (2020). *Event segmentation and the temporal*

compression of experience in episodic memory. *84*, 481–490.

Jeunehomme, O., Folville, A., Stawarczyk, D., Van der Linden, M., & D'Argembeau, A.

(2018). Temporal compression in episodic memory for real-life events. *Memory*,

26(6), 759–770. https://doi.org/10.1080/09658211.2017.1406120

Kim, H. (2012). A dual-subsystem model of the brain's default network: Self-referential

processing, memory retrieval processes, and autobiographical memory retrieval.

Neuroimage, *61*(4), 966–977.

Kriegeskorte, N., & Kievit, R. A. (2013). Representational geometry: Integrating cognition,

computation, and the brain. *Trends in Cognitive Sciences*, *17*(8), 401–412.

https://doi.org/10.1016/j.tics.2013.06.007

Kwok, S. C., Shallice, T., & Macaluso, E. (2012). Functional anatomy of temporal

organisation and domain-specificity of episodic memory retrieval. *Neuropsychologia*,

50(12), 2943–2955. https://doi.org/10.1016/j.neuropsychologia.2012.07.025

Lawrence, M. A. (2016). *ez: Easy analysis and visualization of factorial experiments*.

Lee, H., Bellana, B., & Chen, J. (2020). What can narratives tell us about the neural bases of

human memory? *Current Opinion in Behavioral Sciences*, *32*, 111–119.

https://doi.org/10.1016/j.cobeha.2020.02.007

McDermott, K. B., Szpunar, K. K., & Christ, S. E. (2009). Laboratory-based and

autobiographical retrieval tasks differ substantially in their neural substrates.

Neuropsychologia, *47*(11), 2290–2298.

Michelmann, S., Hasson, U., & Norman, K. (2021). *Event boundaries are steppingstones for*

memory retrieval. PsyArXiv. https://doi.org/10.31234/osf.io/k8j94

Michelmann, S., Staresina, B. P., Bowman, H., & Hanslmayr, S. (2019). Speed of time-

compressed forward replay flexibly changes in human episodic memory. *Nature*

Human Behaviour, *3*(2), 143–154. https://doi.org/10.1038/s41562-018-0491-4

Moscovitch, M., Cabeza, R., Winocur, G., & Nadel, L. (2016). Episodic Memory and

Beyond: The Hippocampus and Neocortex in Transformation. *Annual Review of*

Psychology, *67*, 105–134. https://doi.org/10.1146/annurev-psych-113011-143733

Nielson, D. M., Smith, T. A., Sreekumar, V., Dennis, S., & Sederberg, P. B. (2015). Human

hippocampus represents space and time during retrieval of real-world memories.

Proceedings of the National Academy of Sciences, *112*(35), 11078–11083.

https://doi.org/10.1073/pnas.1507104112

Park, S., & Chun, M. M. (2009). Different roles of the parahippocampal place area (PPA) and

retrosplenial cortex (RSC) in panoramic scene perception. *NeuroImage*, *47*(4), 1747–

1756. https://doi.org/10.1016/j.neuroimage.2009.04.058

R Core Team. (2021). *R: A language and environment for statistical computing*. R

Foundation for Statistical Computing.

Radvansky, G. A., Copeland, D. E., & Zwaan, R. A. (2005). A novel study: Investigating the

structure of narrative and autobiographical memories. *Memory (Hove, England)*,

13(8), 796–814. https://doi.org/10.1080/09658210444000412

Radvansky, G. A., & Zacks, J. M. (2017). Event Boundaries in Memory and Cognition.

Current Opinion in Behavioral Sciences, *17*, 133–140.

https://doi.org/10.1016/j.cobeha.2017.08.006

Ramanan, S., Piguet, O., & Irish, M. (2018). Rethinking the Role of the Angular Gyrus in

Remembering the Past and Imagining the Future: The Contextual Integration Model.

The Neuroscientist: A Review Journal Bringing Neurobiology, Neurology and

Psychiatry, *24*(4), 342–352. https://doi.org/10.1177/1073858417735514

Ranganath, C., & Ritchey, M. (2012). Two cortical systems for memory-guided behaviour.

Nat Rev Neurosci, *13*(10), 713–726.

Rissman, J., Chow, T. E., Reggente, N., & Wagner, A. D. (2016). Decoding fMRI signatures

of real-world autobiographical memory retrieval. *Journal of Cognitive Neuroscience*,

28, 1–17. https://doi.org/10.1162/jocn_a_00920

Ritchey, M., & Cooper, R. A. (2020). Deconstructing the Posterior Medial Episodic Network.

Trends in Cognitive Sciences, *24*(6), 451–465.

https://doi.org/10.1016/j.tics.2020.03.006

Rugg, M. D., & Vilberg, K. L. (2013). Brain networks underlying episodic memory retrieval.

Current Opinion in Neurobiology, *23*(2), 255–260.

https://doi.org/10.1016/j.conb.2012.11.005

Schacter, D. L., Norman, K. A., & Koutstaal, W. (1998). The cognitive neuroscience of

constructive memory. *Annual Review of Psychology*, *49*, 289–318.

https://doi.org/10.1146/annurev.psych.49.1.289

Seghier, M. L. (2012). The Angular Gyrus: Multiple Functions and Multiple Subdivisions.

Neuroscientist.

Smith, S. M., & Nichols, T. E. (2009). Threshold-free cluster enhancement: Addressing

problems of smoothing, threshold dependence and localisation in cluster inference.

NeuroImage, *44*(1), 83–98. https://doi.org/10.1016/j.neuroimage.2008.03.061

Stawarczyk, D., Bezdek, M. A., & Zacks, J. M. (2021). Event Representations and Predictive

Processing: The Role of the Midline Default Network Core. *Topics in Cognitive*

Science, *13*(1), 164–186. https://doi.org/10.1111/tops.12450

Stawarczyk, D., Jeunehomme, O., & D'Argembeau, A. (2018). Differential Contributions of

Default and Dorsal Attention Networks to Remembering Thoughts and External

Stimuli From Real-Life Events. *Cerebral Cortex (New York, N.Y.: 1991)*, *28*(11), 4023–4035. https://doi.org/10.1093/cercor/bhx270

Svoboda, E., McKinnon, M. C., & Levine, B. (2006). The functional neuroanatomy of autobiographical memory: A meta-analysis. *Neuropsychologia*, *44*(12), 2189–2208.

Tulving, E. (2002). Episodic memory: From mind to brain. *Annu.Rev.Psychol.*, *53*, 1–25.

Tversky, B., Zacks, J., & Lee, P. (2004). Events by Hands and Feet. *Spatial Cognition and Computation*, *4*, 5–14. https://doi.org/10.1207/s15427633scc0401_2

Vilberg, K. L., & Rugg, M. D. (2007). Dissociation of the neural correlates of recognition memory according to familiarity, recollection, and amount of recollected information. *Neuropsychologia*, *45*(10), 2216–2225. https://doi.org/10.1016/j.neuropsychologia.2007.02.027

Vilberg, K. L., & Rugg, M. D. (2012). The neural correlates of recollection: Transient versus sustained FMRI effects. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *32*(45), 15679–15687. https://doi.org/10.1523/JNEUROSCI.3065-12.2012

Wagner, A. D., Shannon, B. J., Kahn, I., & Buckner, R. L. (2005). Parietal lobe contributions to episodic memory retrieval. *Trends Cogn Sci*, *9*(9), 445–453.

Winkler, A. M., Ridgway, G. R., Webster, M. A., Smith, S. M., & Nichols, T. E. (2014). Permutation inference for the general linear model. *NeuroImage*, *92*, 381–397. https://doi.org/10.1016/j.neuroimage.2014.01.060

Yarkoni, T., Poldrack, R. A., Nichols, T. E., Van Essen, D. C., & Wager, T. D. (2011). Large-scale automated synthesis of human functional neuroimaging data. *Nature Methods*, *8*(8), 665–670. https://doi.org/10.1038/nmeth.1635

Zacks, J. M. (2020). Event Perception and Memory. *Annual Review of Psychology*, *71*(1), 165–191. https://doi.org/10.1146/annurev-psych-010419-051101

967   Zeidman, P., Mullally, S. L., & Maguire, E. A. (2015). Constructing, Perceiving, and

968   Maintaining Scenes: Hippocampal Activity and Connectivity. *Cerebral Cortex*,

969   *25*(10), 3836–3855. https://doi.org/10.1093/cercor/bhu266

970   Zimmer, H. D., Helstrup, T., & Nilsson, L.-G. (2007). Action events in everyday life and their

971   remembering. In *Everyday memory* (pp. 57–91). Psychology Press.

972