The influence of leaf anatomical traits on photosynthetic in coffee leaves in Yunnan of China

Ziwei Xiao[1*], Guanrun Ma[1]*, Xuehui Bai[1], Jinhong Li[1], Mingzhu Zhao[1], Linlin Su[1], Hongying Ma[2], Hua Zhou[1]

1. *Dehong Tropical Agriculture Research Institute of Yunnan, Ruili, Yunnan 678600, China*

2. Ruining No.1 Nationality Middie Schooi

*Correspondence:xiaozw86@126.com or marui1956@126.com

Abstract

Leaf photosynthesis is largely determined by anatomical features. This work aimed to reveal the quantitative effects of *Coffea arabica* leaf anatomical structure on photosynthesis. Pearson's correlation and path analysis was used to explore the correlation between leaf structure and photosynthesis. The comprehensive evaluation value of the correlation between leaf anatomical traits and photosynthetic parameters is calculated by using the Criteria Importance Though Intercrieria Correlation (CRITIC) method to obtain the objective weight of each photosynthetic parameter. Most of the leaf's anatomical traits significantly ($P<0.05$) to highly significantly ($P<0.01$) correlated with photosynthetic parameters, suggesting that anatomical traits greatly affected photosynthetic capacity in *Coffea arabica* leaf. Likewise, path coefficient analysis revealed that all the leaf anatomical traits influenced photosynthetic capacity directly and indirectly. The results of the comprehensive evaluation showed that leaf thickness (LT) and stomatal density (d) are the anatomical characteristics most closely related to photosynthesis. These results may help pinpoint anatomical features required to gain higher photosynthesis in coffee leaves.

Keywords: *Coffea arabica* L., leaf anatomy, photosynthesis, correlation and path coefficient, CRITIC method, comprehensive evaluation

Introduction

Coffee (*Coffea arabica*) considered an important tropical crop is one of the most traded agricultural products in the world. Under suboptimal growing conditions, shade trees, which could reduce the stress of coffee by ameliorating adverse climatic conditions, have been recommended (Beer *et al.*, 2001; Franck and Vaast, 2009a). Yunnan province, located in Southwest China, is a predominantly coffee-growing region (ICO, 2015). Although the mountainous landscape and mild climate in this region are well-suited to coffee production, many areas are suboptimal for coffee cultivation (ICO, 2015; Ma *et al.*, 2019). Thus, agroforestry systems have been recommended for these areas, especially at altitudes below 1000m (Ma *et al.*, 2019). However, the physiological trait characteristic of shade-adapted coffee is the lower photosynthetic capacity when compared to C_3 woody plants grown in tropical climates (Ayalew, 2018; DaMatta, 2004). Leaves have evolved as photosynthetic organs in land plants (Oguchi *et al.*, 2018) so the leaf of coffee is the main organ of photosynthesis.

Because leaves are specialized organs that enable plants to intercept light and uptake CO_2 through the stomata for photosynthesis (Lambers and Oliveira, 2019), light and CO_2 availability in the mesophyll of plant leaves largely determine the net carbon

assimilation rate (A) of leaves (Franck and Vaast, 2009b). To analyze the response of photosynthesis to irradiance, the light response curves (LRCs) were used to describe the relationship between photosynthesis and light intensity (Herrmann *et al.*, 2020). It is an important method to elucidate the response mechanism of photosynthesis and evaluate photosynthetic efficiency by fitting of light-response model (Lambers and Oliveira, 2019). Often, LRCs are described mathematically using the empirical model. The operation state of plant photosynthetic apparatus and photosynthetic parameters, such as the information of the maximum photosynthetic capacity (P_{nmax}), intrinsic and apparent quantum yield (IQY and AQY), light compensation and saturation point (LCP and LSP), and dark respiration rate (R_D), can be obtained from the LRCs. In addition, mechanistic information about the key biochemical parameters of carbon assimilation can be estimated by fitting *A/C*i curves with the mechanistic Farquhar-von Caemmerer-Berry (FvCB) biochemical model described by Farquhar et al. (1980) (Farquhar *et al.*, 1980; Sun *et al.*, 2014; Wang *et al.*, 2017). This model has been widely used to simulate CO_2 assimilation and applied for decades (Farquhar *et al.*, 1980; GU *et al.*, 2010). It elegantly explains the physiological properties of photosynthesis of intact leaves (Lambers and Oliveira, 2019), and the parameters reflecting biochemical properties of leaves were easier to estimate from this model with gas exchange measurements (Wang *et al.*, 2017).

The structural organization of plant leaves could facilitate or sometimes constrain the photochemical reactions of the photosynthetic process (Boom *et al.*; Oguchi *et al.*, 2018). Therefore, the leave structural changes may affect photosynthetic efficiency (Akhkha *et al.*, 2001; Herrmann *et al.*, 2020). However, although a variety of ecological studies have revealed a correlation between leaf structural parameters and photosynthetic performance (Bolhar-Nordenkampf and Draxler, 1993; James and Bell, 2001; Vogelman *et al.*, 1996), there were contradictory and not universal differences between foliage structural and functional variables in the previous work (Niinemets, 1999). To understand the effects of leaf structure on each photosynthetic performance parameter, it is necessary to analyze the relationship between leaf structure and leaf photosynthesis and comprehensively evaluate the importance of each component in leaf photosynthesis. Because the evaluation of the relationship between leaf structure and leaf photosynthesis could provide ecophysiological evidence for understanding the structural properties of leaves how to influence photosynthetic performance. Even though a number of studies analyze the effects of leaf structure on photosynthetic performance (Bolhar-Nordenkampf and Draxler, 1993; Oguchi *et al.*, 2018), there are few works synthetically analyzing the correlation between leaf structural parameters and photosynthetic performance (Boom *et al.*). In particular, a few works were related to coffee. Therefore, the aims of this study were to explore the relationship between leaf structural parameters and photosynthetic performance, clarify the importance of the correlation between leaf structural parameters and photosynthetic performance by using a comprehensive evaluation method. Gaining these results may provide useful information for future coffee breeding and plantation management.

2 Materials and methods

2.1 Study site characterization and Plant material

The experiment was conducted in the shade coffee plantations of Dehong Tropical Agriculture Research Institute of Yunnan (DTARI), Ruili, southwest China. The plantations are located at 890 meters of altitude, at 24.025 south latitude and 97.855 west longitude, with an average temperature of 22.2 ℃ and precipitation of 1260 mm in the studies conducted years (2017-2018). The soil in the plantation is characterized as acidic lateritic red soil.

The coffee trees of the cultivar are 'Catimor', a hybrid derived from the cross between two cultivars: Caturra (*Coffea arabica* L.) and Timor (*C. arabica* × *C. canephora* Pierre ex Froehner). The coffee trees were approximately five years old and spaced 1×2 m. Shade trees (*Dimocarpus longan* Lour.) planted at 12×4 m, East to West rows-oriented.

A completely randomized design was the following: six coffee plants per treatment combination and used as replicates. All the sampling and measurement leaves used the third or fourth completely expanded leaf pair from the apex of the plagiotropic (lateral) branches located in the upper third of the plant. Nine leaves in different directions on each tree were selected to measure the chlorophyll content values and leaf gas exchange parameters.

2.2 Gas exchange measurements

The leaf gas exchange parameters (net CO_2 assimilation rate (A), stomatal conductance to water vapor (g_s), transpiration rate (E), internal CO_2 concentration (C_i), vapor pressure deficit (VPD), and leaf water use efficiency (WUE=A.E^{-1})) and environmental indicators(leaf temperature (TL), relative humidity (RH) and ambient temperature (TA)) were determined simultaneously by using a portable photosynthesis system CIRAS 3 (PP Systems, USA). Leaf water use efficiency (WUE), representing the units assimilated CO_2 per unit water lost through transpiration, is following the user manual version 1.06. All measurements were taken on sunny days from 09:00 to 11:30 and 15:00 to 18:00. The reference CO2 concentration was set at 390 ppm using a buffer bottle, the temperature at 27 ∘C, flow rate at 300 $\mu mol.s^{-1}$, relative humidity at 50 ± 10%, and irradiance at 800 $\mu mol\ m^{-2}.s^{-1}$. The leaf of photosynthetic light response (A/PPFD) and internal CO_2 response (A/C_i) curves were constructed by using leaf gas exchange. To collect data for building A/PPFD curves, net photosynthesis was measured at 0, 50, 100, 150, 200, 250, 300, 350, 400, 500, 600, 700, 800, 1000, and 1200 $\mu mol\ m^{-2}.s^{-1}$ (PAR) with a cuvette environment that maintained leaf temperature at 27 ℃, reference CO2 concentration at 400 ppm. After the leaf tissue was clamped in the leaf chamber, the gas exchange rates usually stabilized within approximately 3–4 min. Data to construct A/C_i curves were collected from the same sample leaves immediately following A/PPFD measurements. P_n was measured at CO_2 concentrations of 400, 300, 200, 100, 150, 250, 350, 450, 550, 650, 750, 850, 950, 1050, 1250, 1450 and 1650 $\mu mol.m^{-2}.s^{-1}$ at photosynthesis saturating irradiance 800 $\mu mol\ m^{-2}.s^{-1}$，which light intensity is sufficiently high to saturate the photosynthetic machinery without causing photoinhibition. A total of 41 A/PPFD and 42 A/C_i curves were generated, respectively.

2.3 Measurement of leaf anatomical traits

Following gas exchange measurement, five sample leaves of each plant were

harvested, sealed in polyethylene bags, and immediately brought back to the laboratory. Two segments (3 mm × 5 mm) were cut from each leaf halfway between the leaf apex and base. Then, the material was dehydrated on sucrose solutions, followed by the inclusion of Gum Arabic aqueous solution. Leaf transverse sections (7–10 μm) thickness cut using a rotary microtome (Leica CM3050S, Germany) and stained with toluidine blue. Surface sections of the leaves were taken manually and stained with safranine aqueous solution (0.1%). For each leaf, five transversal sections and surface sections were selected systematically to measure the leaf thickness (LT), upper cuticle thickness (UCT), upper epidermal thickness (UET), palisade tissue thickness (PT), spongy tissue thickness (ST), lower epidermal thickness (LET), lower cuticle thickness (LCT), stomatal density (d), and guard cell length (l) and width (w), using a microscope binocular with 10 × and 40 × objectives (Leica DM5000 B). Subsequently, five photographs of microscope fields per section were taken with a photomicroscope equipped with a Leica DFC500 camera (Leica Microsystems, Rijswijk, The Netherlands). Images were analyzed with Leica LAS X software.

2.4 Calculations and statistical analysis

A/PPFD curves were fitted by applying a modified rectangular hyperbola model using Photosynthetic Calculation Software 4.1.1(Ye, 2007; Ye, 2010; Ye and Yu, 2008). The equation of the modified rectangular hyperbola model is described by:

$$P_n(I) = \alpha \frac{1 - \beta I}{1 + \gamma I} I - R_d \qquad (1)$$

where P_n is the net photosynthetic rate ($\mu mol \cdot CO_2 \ m^{-2} \ s^{-1}$), I is the photosynthetic photon flux density ($\mu mol \cdot m^{-2} \ s^{-1}$), I_c is compensation irradiance, β and γ are the coefficients which are independent of I, and α is the absolute value of slope between $I = 0$ and $I = I_c$.

For $I = 0$, the quantum yield at this point is defined as intrinsic quantum yield (IQY), which is given by:

$$IQY = \alpha(1 + (\gamma + \beta)I_c) \qquad (2)$$

The apparent quantum yield (AQY) ignored the Kok effect is obtained by:

$$AQY = \alpha \frac{1 + (\gamma - \beta)I_c - \gamma \beta I_c^2}{(1 + \gamma I_c)^2} \qquad (3)$$

For $I = 0$, the rate of dark respiration (R_d) is:

$$R_d = -P_n(I_0) = -\alpha I_c \qquad (4)$$

The saturation irradiance I_{sat} is obtained by:

$$I_{sat} = \frac{\sqrt{(\beta + \gamma)/\beta} - 1}{\gamma} \qquad (5)$$

The maximum photosynthetic rate P_{nmax} is given by:

$$P_{nmax} = \alpha \left(\frac{\sqrt{\beta + \gamma} - \sqrt{\beta}}{\gamma} \right)^2 - R_d \qquad (6)$$

A/Ci curves were fit by uploading the datasets to the website (http://Leafweb.ornl.gov) to estimate the parameters. In this study, A/C_i curves were at saturating light levels. Thus, the rate of electron transport (J) was to assume the maximum rate of electron

transport (J_{max}).

The total stomach pore area (TSP) was calculated as $d \times 1/4\pi \times l \times w$ (7), and by

assuming that the stomatal pore was an ellipse(Batos *et al.*, 2010; James and Bell,

2001; Patricia *et al.*, 2014). Where d is the stomatal frequency calculated by dividing

the stomatal count by the area of the field of view, l is the guard cell length, and w is

the guard cell width.

The relationships of photosynthetic capacity with leaf anatomical traits were analyzed

by Pearson's correlation. The possible direct and indirect effects of each leaf

anatomical trait on the photosynthetic parameter were further identified by path

analysis(Zhang *et al.*, 2005). For path analysis, direct and indirect effects were

obtained from partial least squares (PLS) regression of leaf anatomical traits on

photosynthetic parameter and Pearson's correlations between leaf anatomical

traits(Guohua *et al.*, 2017). To combine effect leaf anatomical traits on photosynthetic

parameters, a multiple criteria decision analysis (MCDA) has been utilized in this

study. The objective weight of each photosynthetic parameter (w_j) is obtained by

using the Criteria Importance Though Intercrieria Correlation (CRITIC) method,

which was a correlation method using a standard deviation of column and correlation

coefficients of all paired columns(Diakoulaki *et al.*, 1995). The comprehensive

evaluation of the effects of leaf anatomical traits on photosynthetic parameters was

calculated by using the following equation: $Q_i = \sum_{j=i}^{n}(w_j \times r_{ij})$(8), where r_{ij} was

the correlation coefficient between leaf anatomical trait i and photosynthetic

parameter j. All statistical analyses were conducted with IBM SPSS Statistics 24.

Results

Results showed the photosynthetic light-response curve (Fig.1) fits very well with

the modified rectangular hyperbola model ($R^2=0.945 \pm 0.006$) described by Ye (2007)

(Ye, 2007). The P_n increased with I below the I_{sat}. When I was above I_{sat}, the P_n

decreased as I increased, which means the photoinhibition phenomenon. The

calculated AQY would be 0.028 ± 0.003 if the Kok effect were ignored. This value was

lower than α (0.031 ± 0.003) and the IQY (0.033 ± 0.003). The maximum P_n values

calculated by modified rectangular hyperbola model at I_{sat} of 621.762 ± 37.358

$\mu mol(photon)\ m^{-2}s^{-1}$ was $5.131 \pm 0.478\ \mu mol(CO_2)m^{-2}s^{-1}$ for coffee leaf. The values of

I_c and R_D were $60.238 \pm 11.332 \mu mol(photon)m^{-2}s^{-1}$ and $0.930 \pm 0.098 \mu mol(CO_2)m^{-2}s^{-1}$,

respectively.

Fig.1 Photosynthetic light-response curve of coffee plants. Data shown in the figure indicate mean ± stand error.

IQY–intrinsic quantum yield; AQY–apparent quantum yield; α–the absolute value of slope between $I = 0$ and $I = I_c$;

P_{nmax}–maximum photosynthetic rate [$\mu mol(CO_2)$ m^{-2} s^{-1}]; I_{sat} – light saturation point [μmol (photon) m^{-2} s^{-1}]; I_c

–light compensation point [μmol (photon) m^{-2} s^{-1}]; R_D– dark respiration [$\mu mol(CO_2)$ m^{-2} s^{-1}].

Based on the FvCB model structure, the A/C_i curve was fitted with the interactive

website (http://www.leafweb.ornl.gov)(Gu and Sun, 2014). Figure 2 shows the

response of A to C_i for coffee leaves. The parameters of values for V_{cmax}, J_{max}, T_p, R_d,

Γ^*, g_i, and K_{co} were 61.749±5.394, 80.721±17.963, 5.316±0.576, 3.225±0.441,

3.309±0.333, 1.101±0.287, 103.096±11.721, respectively. The transition between the

Rubisco- and RuBP regeneration-limited states (C_{i_CJ}) occurs around 30 Pa

intercellular CO_2 and the TPU-limited state (C_{i_JP}) is around 90 Pa (Fig 2.).

Fig 2. A/C_i curves for coffee plants. Data shown in the figure indicate mean ± stand error. C_i–Intercellular CO_2

concentration (μmol mol^{-1}); V_{cmax}–Maximal Rubisco carboxylation rate (μmol m^{-2} s^{-1}); J_{max}–Maximal electron

transport rate (μmol m^{-2} s^{-1}); T_p–Rate of triose phosphate export from the chloroplast (μmol m^{-2} s^{-1}); R_d–Day

respiration (μmol m^{-2} s^{-1}); Γ^*–CO_2 compensation point in the absence of dark respiration (Pa); g_i– Internal

(mesophyll) conductance to CO_2 transport (μmol m^{-2} s^{-1} Pa^{-1}); K_{oc} –A composite parameter (Pa): $K_{oc} =$

$K_c \left(1 + {}^O/_{K_o}\right)$.

Interrelationships with correlation analysis between photosynthetic capacity and

leaf anatomical traits were presented in Table 1. There were many significant

differences between photosynthetic capacity and leaf anatomical traits. The UCT

correlated negatively with IQY, AQY, α, I_{sat}, and g_i ($p < 0.05$). The UET appeared to be positively related to AQY, α, and P_{max} ($p < 0.05$), and negatively related to I_c, R_d, C_i, and VPD ($p < 0.05$) (Table 1). The relationship between PT and IQY, AQY and α was significantly negative ($p < 0.05$). The LET had a significantly negative effect on the I_c was significantly negative ($p < 0.05$). The LCT was important to I_{sat}, and the Pearson correlation coefficient was significantly negative ($p < 0.05$). The guard cell width (w) significantly affected I_c and R_d. The guard cell length (l) correlated negatively with IQY, AQY, α, and I_{sat}, and positively with J_{max}. A strongly positive ($p < 0.01$) correlation relationship between stomatal density (d) and J_{max} was found in this study. Based on Pearson correlation analysis, however, LT and ST had relatively little effect on the photosynthetic capacity.

Path analysis was performed to reveal the direct and indirect effects of leaf anatomical traits on photosynthetic capacity. The results indicating the effects were shown in Fig 3. Although correlation analysis suggested that there were no significant correlations between LT and photosynthetic capacity variables ($p > 0.05$), its direct effects on IQY (0.488), α (0.429), R_d (0.390), and AQY (0.381) were relatively high. In addition, there were high negative direct effects of LT on I_{sat} (-0.431) and Γ^* (-0.379). The negative direct effect of UCT on IQY (-0.756), α (-0.720), and AQY (-0.679) was found to be higher than the other photosynthetic capacity variables. UET has a high negative direct effect on R_d (-0.785), TPU (-0.520), and a high positive direct effect on E (0.563), g_s (0.451), and P_{max} (0.448). PT exhibited an impressive negative direct effect on TPU (-0.518), AQY (-0.410), α (-0.410), and IQY (-0.401), while a high positive direct effect was shown between PT and E (0.406). The leaves of ST presented a high positive direct effect on I_{sat} (0.583) and WUE (0.578). In addition, a similar high negative direct effect was also observed between ST and VPD (-0.588). LET presented high positive direct effects on WUE (0.514), V_{cmax} (0.483), and TPU (0.391). However, a relatively high negative direct effect was observed between LET and R_D (-0.362), VPD (-0.359), and C_i (-0.321). There was a direct effect with a relatively high negative magnitude of LCT on V_{cmax} (-0.478), I_c (-0.416), J_{max} (-0.371), and VPD (-0.342), while the LCT has a higher positive direct effect on the Γ^* (0.335) than photosynthetic capacity variables. The magnitude of negative direct effect guard cell width (w) on I_{sat} (-0.461) was observed. Also, a large relatively negative direct effect of guard cell length (l) on I_{sat} (-0.422) and g_i (-0.364) was found. However, l also observed high positive direct effects on C_i (0.369). The stomatal density (d) exhibited large negative direct effects on g_i (-0.379) and high positive direct effects on J_{max} (0.350) was found. The direct effect of TSP on the photosynthetic capacity variable was low, and the value range was from -0.179 to 0.185.

Table 1. The correlation between leaf anatomical traits and photosynthetic parameter

Parameters	LT	UCT	UET	PT	ST	LET	LCT	w	l	d	TSP
IQY	-0.044	-0.336	0.296	-0.343	-0.010	0.000	-0.232	0.175	-0.315	0.232	0.103
	ns	*	ns	*	ns	ns	ns	ns	*	ns	ns
AQY	-0.024	-0.322	0.386*	-0.394	0.012	0.039	-0.224	0.197	-0.358	0.206	0.068

	ns	*	*		ns	ns	ns	ns	*	ns	ns
α	-0.033	-0.332	0.351*	-0.375	0.005	0.022	-0.230	0.189	-0.342	0.217	0.082
	ns	*	*		ns	ns	ns	ns	*	ns	ns
P_{max}	0.178	-0.108	0.324*	-0.036	0.103	0.100	-0.065	-0.037	-0.233	0.143	-0.003
	ns	ns		ns	ns	ns	ns	ns	ns	ns	ns
I_{sat}	-0.057	-0.359	0.157	-0.278	0.189	-0.202	-0.377	-0.227	-0.460	0.031	-0.313
	ns	*	ns	ns	ns	ns	*	ns	**	ns	*
I_c	-0.064	-0.143	-0.324	0.164	-0.008	-0.336	-0.174	-0.323	-0.071	0.017	-0.150
	ns	ns	*	ns	ns	*	ns	*		ns	ns
R_D	0.025	-0.234	-0.195	0.204	0.057	-0.190	-0.150	-0.243	-0.005	0.156	0.036
	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns
R_d	0.097	0.161	-0.477	0.101	0.003	-0.129	0.071	-0.354	0.133	-0.043	-0.114
	ns	ns	**	ns	ns	ns	ns	*	ns	ns	ns
V_{cmax}	-0.021	-0.097	0.009	0.106	0.188	0.019	-0.161	0.172	-0.003	0.139	0.200
	ns	ns	ns	ns	ns	ns	ns	ns	ns	*	ns
J_{max}	-0.086	-0.169	-0.104	-0.047	-0.035	-0.188	-0.183	-0.003	-0.212	0.403*	0.246
	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns
TPU	-0.033	-0.145	-0.209	0.012	0.045	-0.008	0.015	-0.162	0.045	0.245	0.183
	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns
Γ^*	-0.210	0.181	-0.086	-0.159	-0.239	0.228	0.119	-0.163	0.104	-0.285	-0.276
	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns
g_i	0.207	-0.433	0.311	-0.092	0.371	-0.071	-0.420	-0.006	-0.301	-0.149	-0.301
	ns	*	ns	ns	ns	ns	*	ns	ns	ns	ns
C_i	-0.066	0.108	-0.396	0.257	-0.125	-0.164	0.187	-0.053	0.311*	-0.080	0.077
	ns	ns	**	ns	ns	ns	ns	ns		ns	ns
g_s	0.140	-0.111	0.278	0.073	0.044	0.146	0.053	0.005	-0.085	0.174	0.116
	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns
VPD	-0.023	0.071	-0.372	0.281	-0.143	-0.230	0.106	-0.063	0.221	0.052	0.148
	ns	ns	*	ns	ns	ns	ns	ns	ns	ns	ns
A	0.157	-0.079	0.346*	0.007	0.049	0.203	0.047	0.048	-0.117	0.149	0.094
	ns	ns		ns	ns	ns	ns	ns	ns	ns	ns
E	0.196	-0.057	0.224	0.191	0.031	0.068	0.106	0.002	0.003	0.164	0.158
	ns	ns	*	ns	ns	ns	ns	ns	ns	ns	ns
WUE	0.049	-0.115	0.350*	-0.230	0.184	0.226	-0.142	0.041	-0.216	0.074	-0.038
	ns	ns		ns	ns	ns	ns	ns	ns	ns	ns

LT–leaf thickness, UCT–upper cuticle thickness, UET–upper epidermal thickness, PT–palisade tissue thickness, ST–spongy tissue thickness, LET–lower epidermal thickness, LCT–lower cuticle thickness, w–guard cell width, l–guard cell length, d–stomatal density, TSP–total stomach pore area. Ns = non-significant. *, ** significant significance difference at 5 % and 1 % probability levels, respectively.

The estimates of each indirect effect of leaf anatomical trait on photosynthetic capacity via other leaf anatomical traits were presented in Fig 3(b). According to this figure, the values of the indirect effect of LT and ST on photosynthetic capacity variables through other anatomical traits were lower than 0.300. UCT revealed a

negative indirect effect on IQY (-0.335), I_{sat} (-0.343), α (-0.326), g_i (-0.319), and AQY (-0.313) through other anatomical traits. UET indirectly exerted relatively high positive effects on AQY (0.357), A (0.345), P_{max} (0.325), and α (0.322) via other anatomical traits. Conversely, UET exhibited relatively high negative indirect effects on R_d (-0.479) and VPD (-0.367) via other anatomical traits. The indirect effects of PT on g_i via other anatomical traits were negative and high (-0.329). There were indirect effects with a high negative magnitude of LCT on I_{sat} and g_i via other anatomical traits(-0.359 and -0.340, respectively). The R_d and I_c were influenced indirectly with a relatively high negative magnitude by w via other anatomical traits (-0.352 and -0.307, respectively). There was a relatively high positive indirect effect of l on C_i (0.314) and negative on I_{sat} (-0.450), AQY (-0.320) and α (-0.302) through other anatomical traits. In addition, a highly positive indirect effect of d on J_{max} (0.399) was observed. The TSP presented a relatively high negative indirect effect on I_{sat} (-0.312) and g_i (-0.344) through other anatomical traits.

(a)

(b)

To identify the combined effect of leaf anatomical traits on photosynthetic parameters, a comprehensive evaluation was analyzed. Based on the evaluation values (Q_i), the ranking of the comprehensive scores of relationships between photosynthetic capacity and leaf anatomical traits was shown in Table 2. The highest comprehensive score of direct effect was obtained by the LT followed by LET, d, TSP, ST, L, LCT, UET, PT, W, and UCT. For the comprehensive score of indirect effect on of leaf anatomical traits photosynthetic capacity, the d held the highest comprehensive score, and ST was the second followed by LT, UET, TSP, PT, LET, W, LCT, L, and UCT. Similarly, for Pearson's correlation of comprehensive evaluation analysis, the results showed that the first comprehensive score was d, and the second was also ST. Then, the ranking order after the third was UET, LT, TSP, LET W, PT, LCT, L, and UCT.

Table 2. Comprehensive ordering of the effects of leaf anatomical traits on photosynthetic capacity with CRITIC method.

Anatomical traits	Direct effects		Indirect effects		Pearson's correlation	
	Comprehensive score	Ranking	Comprehensive score	Ranking	Comprehensive score	Ranking
LT	0.123	1	0.009	3	0.007	4
UCT	-0.168	11	-0.125	10	-0.137	11
UET	-0.034	8	0.002	4	0.027	3
PT	-0.056	9	-0.021	5	-0.049	8
ST	0.022	5	0.032	2	0.033	2
LET	0.098	2	-0.049	6	-0.042	6
LCT	-0.020	7	-0.079	8	-0.102	9
W	-0.062	10	-0.060	7	-0.045	7
L	-0.008	6	-0.085	9	-0.104	10
d	0.048	3	0.081	1	0.076	1
TSP	0.039	4	0.002	4	-0.007	5

LT–leaf thickness, UCT–upper cuticle thickness, UET–upper epidermal thickness, LET–lower epidermal thickness, PT–palisade tissue thickness, ST–spongy tissue thickness, LCT–lower cuticle thickness, w–guard cell width, l–guard cell length, d –stomatal density, TSP–total stomach pore area.

Discussion

Several studies involving gas exchange measurements of coffee leaves have been performed. These studies have pointed out that the photosynthetic capacity of coffee leaves varied with a variation environment (Ayalew, 2018; DaMatta, 2004; DaMatta and Ramalho, 2006; Kumar and Tieszen, 1980). For single coffee leaves, the saturating irradiance is between 300 and 700 mmol photons m^{-2} s^{-1}, with shade leaves showing a lower value than lower sun leaves (DaMatta, 2004; MATTA). This study found that the saturating irradiance of sun leaves was about 620 mmol photons m^{-2} s^{-1}, which was consistent with the result of the above mention. Quantum yields revealed the relationship between a given light-dependent product and the number of absorbed photons (Taiz and Zeige, 2006). Quantum yield is 0 when none of the light energy is used in photosynthesis. Quantum yield is 1 when all the absorbed light is used. Based

on the results of YE (2007), the quantum yield decreased with I increasing (Ye, 2007).
Thus, the IQY was higher than AQY. In the past, there were several pertinent points
about maximal photosynthetic rates of coffee leaves. For example, Kumar and
Tieszen (1980) and Cannell (1985) have pointed out that the maximal photosynthetic
rates of sun leaves of coffee are around 7 mmol CO_2 m^{-2} s^{-1} and 8.8 mmol CO_2 m^{-2} s^{-1},
respectively (Cannell, 1985; Kumar and Tieszen, 1980). However, Bote et al. (2018)
reported that the maximal photosynthetic rates of sun leaves of coffee were lower than
7 mmol CO_2 m^{-2} s^{-1} with a larger N supply (Bote *et al.*, 2018). In this study, the
maximal photosynthetic rates were lower than observations of Kumar and Tieszen
(1980) and Cannell (1985) and consistent with the result of Bote et al. (2018). For
coffee leaves, light saturation would lead to an excess of electron transport capacity
(Martins *et al.*, 2014). Therefore, higher J_{max} was observed under light saturation. In
addition, the values of V_{cmax} and J_{max} were larger than Martins et al. (2014) reported
result.
The correlation analysis is a great utility method for understanding the relationships
between variables. In this study, significant correlations between photosynthetic
capacity and leaf anatomical traits were observed, indicating that leaf anatomical traits
could affect photosynthetic capacity. Despite the correlation coefficients could
provide a good way to understand the relationships between variables, the cause and
effect relationships between variables couldn't be estimated (Gonzalez-Paleo and
Ravetta, 2018). Path analysis, which investigates the cause and effect relationship, can
provide a detailed knowledge of positive and negative correlations between traits
(Gonzalez-Paleo and Ravetta, 2018). The result of the path analysis was shown in Fig
3, indicating that there were high-level direct and indirect effects of leaf anatomical
traits on photosynthetic capacity. This may explain the importance of leaf structure in
photosynthesis. In order to quantitatively evaluate the effect of each structure of leaf
on photosynthesis, the comprehensive evaluation values are calculated by using the
formula (8) introduced in the methods. The ordering comprehensive score showed
that leaf thickness (LT) and stomatal density (d) were the most important leaf
anatomical traits for coffee leaf photosynthesis. For Arabica coffee, the photosynthetic
is greatly limited by CO_2 diffusion (DaMatta *et al.*, 2007; Franck and Vaast, 2009b).
The CO_2 diffusion from the air to the leaf mesophyll is mainly modulated by the
stomatal conductance (g_s), which is associated with stomatal density (d) (Lambers and
Oliveira, 2019; Rodrigues *et al.*, 2018). In fact, higher stomatal density will increase
g_s by increasing the same total pore area (Rodrigues *et al.*, 2018). This may partly
explain why the stomatal density (d) had higher comprehensive evaluation values in
the effect of coffee leaf photosynthesis.
The leaf thickness strongly positively correlated with the fraction of intercellular air
space in the leaves (Niinemets, 1999). Given that the CO_2 diffusion in the gaseous
phase is about three orders of magnitude larger than in the liquid (Oguchi *et al.*,
2018)(108). Thus, the diffusive resistance should not necessarily increase
considerably with increasing leaf thickness (Smith *et al.*, 1997). Moreover, due to the
chloroplasts adhering to the inner surface of exposed mesophyll cell walls, thicker
leaves provide space for more chloroplasts per unit leaf area (Lambers and Oliveira,

2019) and possess more chloroplast surface area, which results in a substantial
increase in carbon dioxide absorption because of a larger surface area for diffusion
(Smith *et al.*, 1997). Furthermore, thicker leaves may have higher total contents of
Rubisco and leaf N per unit area (Niinemets and Sack, 2006). In this study, thick
leaves have the highest most important direct effects on photosynthetic capacity as
compared with other anatomical traits by using the CRITIC method. In addition to the
result of path analysis, the relatively high direct effects of thick leaves (TL) on IQY, α,
R_d, and AQY were also observed.

The structure of the leaf cuticle consists of epicuticular wax, birefringent wax
embedded in cutin, cutin matrix, and pectic substance (Lee and Graham, 1986). The
epicuticular wax can decrease light by scattering (Bote *et al.*, 2018). The most cutin
matrix in the cuticle is heterogeneous, and the areas of embedded waxes exhibit
birefringence in polarized light (Lee and Graham, 1986). This heterogeneous structure
of the leaf cuticle may affect epidermal focusing, allowing actinic light to penetrate
deeper into the mesophyll (Bote *et al.*, 2018). These findings most likely explained
the significant relationship between cuticle thickness and photosynthetic gas exchange
parameters. In addition, the cuticle thickness would increase resistance to CO_2
diffusion (Miranda *et al.*, 1981), and the negative relationships between UCT and
LCT and photosynthetic parameters were also observed in this study.

The epidermal cell usually has a large transparent central vacuole and lacks mature
chloroplasts, thus the light can directly pass from these cells into the mesophyll (Bote
et al., 2018). However, these epidermal cells have various shapes, such as
plano-convex, spherical, conical, or some other convex shape (Jackson *et al.*, 2012).
The curved outermost epidermal cell wall can affect the focal point within the leaf
(Bote *et al.*, 2018). However, the result of Brodersen and Vogelmann (2007) showed
that there was no relative between the convexity of epidermal cells and the
absorptance of diffuse light (Brodersen and Vogelmann, 2007). Although two
contradictory hypotheses have been proposed with regard to the impacts of epidermal
cells on photosynthesis, this study examined epidermal lens effects on many
photosynthetic parameters and elucidated cells property plays an important role in
photosynthetic capacity by using the CRITIC method.

When intrinsic quantum yields are high, light capture and utilization are most
efficient(Herrmann *et al.*, 2020). However, structural changes of a leaf may affect
light absorption (Herrmann *et al.*, 2020). For example, thicker mesophyll would
enhance light absorptance in leaves, due to greater internal light scattering (Lambers
and Oliveira, 2019). Because photon irradiance absorptance varied with variation
chlorophyll concentration of leaf (Lambers and Oliveira, 2019), the quantum yield
would be affected by chlorophyll concentration. Most chlorophylls were contained in
the palisade cells. This could explain the significant relationship between quantum
yield and PT. In addition, measurements of light gradients in leaves showed that
palisade tissue (PT) facilitates the penetration of direct sunlight (Vogelman *et al.*,
1996). The result of this study also indicated that a large relatively negative direct
effect of PT on IQY, AQY, and J_{max} were observed.

The concentration of CO_2 in the mesophyll is affected by air temperature, and each

degree rise in temperature above 24 ℃ results in a 20 ppm rise in CO_2 concentration
(Nunes *et al.*, 1968). However, Khairi and Hall's (1976) studies of citrus
photosynthesis showed that mesophyll conductance to CO_2 decreased with
temperature increase when the temperature increased from 22 ℃ to around 40 ℃
(Khairi and Hall, 1976). In the analysis of effects of leaf anatomical traits on
photosynthetic capacity, the spongy tissue thickness (ST) had larger indirect effects
and Pearson's correlation comprehensive score, suggesting that spongy tissue
thickness (ST) plays an important role in the photosynthetic process. In addition,
Martins et al. (2014) showed that mesophyll thickness and stomatal attributes related
to water flux and gas exchange per leaf area (Martins *et al.*, 2014), which is consistent
with the result of this study. On the other hand, the irregular shaped spongy cells
alternating with air spaces increases light absorption due to the optical path length
through a leaf increase (Oguchi *et al.*, 2018)(105), and these absorbed light will be
advanced for photosynthesis (Brodersen and Vogelmann, 2007; Cannell, 1985).
Although there was little relationship between spongy tissue thickness (ST) and
photosynthetic parameters, the high direct spongy tissue thickness (ST) effects upon
I_{sat}, WUE, and VPD was observed.

Leaf structure plays a key role in determining the conductance of CO_2 through the
leaf because changes in leaf structure will change gas diffusion within the leaves
(Evans *et al.*, 1994). Moreover, the CO_2 diffusion conductance is also influenced by
the density and arrangement of the stomata (Oguchi *et al.*, 2018). For the coffee leaf,
the net carbon assimilation rate (A) greatly is governed by diffusive conductances of
CO_2, thus the photosynthesis of the coffee leaf was largely limited by a diffusive
factor(Araujo *et al.*, 2008). Traditionally, stomata have been considered as one of the
major roles in controlling CO_2 diffusive (Araujo *et al.*, 2008) and were a key
parameter in the photosynthesis of *C. arabica*(Ayalew, 2018). Among the various
stomatal characters, stomatal density plays a major role in gas exchange and
photosynthesis (Pompelli *et al.*, 2010). In this study, it is shown that stomatal density
had the highest comprehensive scores compared with other anatomical traits in
indirect effects and Pearson's correlation comprehensive evaluation. In addition,
structural modification of the stomata is one of the key factors affecting stomatal
conductance (Taiz and Zeige, 2006). The result of this study also showed that the
guard cell length (*l*) significantly correlated positively with C_i.

For light absorption, although the result of Martins et al. (2014) showed that the
guard cell length did not change in response to light treatments (Martins *et al.*, 2014),
there was a significant correlation with IQY and AQY. The coordination between
increased initiation of stomata cells and expansion of epidermal cells was observed in
coffee leaves, implying that there was an optimization of the trade-off between
transpirational costs and CO_2 assimilation (Martins *et al.*, 2014). In addition, stomatal
size and density largely controlled the leaf's diffusive conductance to CO_2, and the
balance between the CO_2 required for photosynthesis and the level of water
availability was also determined by stomatal traits (Rodrigues *et al.*, 2018). Thus,
lower epidermal stomatal density had a significant correlation with photosynthetic
parameters or direct effects and indirect effects on photosynthetic parameters.

Moreover, these traits also had larger comprehensive scores in a comprehensive evaluation. This means that stomatal density was a key parameter in the regulation photosynthesis of *C. arabica*.

Conclusion

Many leaf anatomical traits significantly affected photosynthetic parameters. This result provided holistic relationships between leaf anatomical traits and photosynthetic parameters in coffee. It is tempting to suggest that these relationships might to some extent explain the photosynthetic behaviour of coffee (*Coffea arabica* L.). The relationships between leaf anatomical traits and photosynthetic parameters emphasize the importance of leaf anatomy in determining photosynthesis at the tissue level. Moreover, these correlations between leaf anatomical traits and photosynthetic parameters were useful to explore the impact of different anatomical features on photosynthetic efficiency and accordingly provide some information to design leaf anatomy for enhanced photosynthetic efficiency. To a certain extent, path analysis and comprehensive evaluation could indicate interrelationship between photosynthetic capacity and leaf anatomical traits. However, the causal interrelationship between photosynthetic parameters and leaf anatomical traits is less clear. In spite of this, the result of the study also suggests that improvements of the photosynthetic performance of coffee through selecting thinker leaf and higher stomatal density that might support higher photosynthetic capacity could be a useful strategy to facilitate the selection of promising genotypes with enhanced coffee growth and production.

Author contributions

Z.X. and G.M. conceptualization and research design; X.H. and J.L. project administration;M.Z.and L.S. investigation and fieldwork, Z.X. analysed the data and wrote the manuscript; H.M. and H.Z. resources; all theauthors read and edited the manuscript, and approved the final version.

Funding

This research was funded by the Basic Research Project of Dehong Tropical Agriculture Research Institute of Yunnan (DTARI-JJ2020-02) and DR155 Coffee Variety Industrialization Technology Integration and Application Demonstration (202003AD150006), and National Tropical Plants Germplasm Resource Center.

References

Akhkha A, Reid I, Clarke DD, Dominy P. 2001. Photosynthetic light response curves determined with the leaf oxygen electrode: minimisation of errors and significance of the convexity term. *Planta* **214**, 135-141.

Araujo WL, Dias PC, Moraes GABK, Celin EF, Cunha RL, Barros RS, Damatta FM. 2008. Limitations to photosynthesis in coffee leaves from different canopy positions. *Plant Physiol Biochem* **46**, 884-890.

Ayalew B. 2018. Impact of shade on morpho-physiological characteristics of coffee plants, their pests and diseases: A review. *African Journal of Agricultural Research* **13**, 2016-2024.

Batos, Vilotic, Orlovic, Miljkovic. 2010. Inter and intra-population variation of leaf stomatal traits of quercus robur l. In northern serbia. *ARCH BIOL SCI* **2010,62(4)**, 1125-1136.

Beer JW, Muschler RG, Somarriba E, Kass D. 2001. Shade management in coffee and cacao plantations- a review. *Journal of Experimental Botany* **52**, 2057-2065.

Bolhar-Nordenkampf H, Draxler G. 1993. Functional leaf anatomy. *Photosynthesis and production in a changing environment*: Springer, 91-112.

Boom, Tholen, Zhu XG. Opinion: prospects for improving photosynthesis by altering leaf anatomy.

Bote AD, Zana Z, Ocho FL, Vos J. 2018. Analysis of coffee (Coffea arabica L.) performance in relation to radiation level and rate of nitrogen supply II. Uptake and distribution of nitrogen, leaf photosynthesis and first bean yields. *European journal of Agronomy* **92**, 107-114.

Brodersen CR, Vogelmann TC. 2007. Do epidermal lens cells facilitate the absorptance of diffuse light? *American Journal of Botany* **94**, 1061-1066.

Cannell M. 1985. Physiology of the coffee crop. *Coffee*, 108-134.

DaMatta FM. 2004. Ecophysiological constraints on the production of shaded and unshaded coffee: a review. *Field Crops Research* **86**, 99-114.

DaMatta FM, Ramalho JDC. 2006. Impacts of drought and temperature stress on coffee physiology and production: a review. *Brazilian Journal of Plant Physiology* **18**, 55-81.

DaMatta FM, Ronchi CP, Maestri M, Barros RS. 2007. Ecophysiology of coffee growth and production. *Braz. J. Plant Physiol* **19**, 485-510.

Diakoulaki D, Mavrotas G, Papayannakis L. 1995. Determining objective weights in multiple criteria problems: The critic method. *Computers & Operations Research* **22**, 763-770.

Evans JR, Caemmerer S, Setchell BA, Hudson GS. 1994. The relationship between CO2 transfer conductance and leaf anatomy in transgenic tobacco with a reduced content of Rubisco. *Functional Plant Biology* **21**, 475-495.

Farquhar GD, Caemmerer SV, Berry JA. 1980. A biochemical model of photosynthetic CO2 assimilation in leaves of C3 species. *Planta* **149**, 78-90.

Franck N, Vaast P. 2009a. Limitation of coffee leaf photosynthesis by stomatal conductance and light availability under different shade levels. *Trees structure and function*.

Franck Ns, Vaast P. 2009b. Limitation of coffee leaf photosynthesis by stomatal conductance and light availability under different shade levels. *Trees* **23**, 761-769.

Gonzalez-Paleo L, Ravetta DA. 2018. Relationship between biochemical and photosynthetic traits with Asian soybean rust. *Photosynthetica* **90**, 3925-3940.

GU L, Pallardy GS, TU K, Law BE, Wullschleger SD. 2010. Reliable estimation of biochemical parameters from C-3 leaf photosynthesis-intercellular carbon dioxide response curves. *Plant, Cell & Environment* **33**, 1852-1874.

Gu L, Sun Y. 2014. Artefactual responses of mesophyll conductance to CO2 and irradiance estimated with the variable J and online isotope discrimination methods. *Plant Cell & Environment* **37**, 1231-1249.

Guohua W, Hu Z, Yuesong W. 2017. Implementation of Partial Least Squares Regression in SPSS Software. *Statistics & Decision*, 67-71.

Herrmann HA, Schwartz JM, Johnson GN. 2020. From empirical to theoretical models of light response curves - linking photosynthetic and metabolic acclimation. *Photosynthesis Research* **145**, 5-14.

ICO. 2015. Coffee in China. In: Organization IC, ed. *International Coffee Council*. Milan, Italy: International Coffee Organization.

Jackson D, Skillman J, Vandermeer J. 2012. Indirect biological control of the coffee leaf rust, Hemileia vastatrix, by the entomogenous fungus Lecanicillium lecanii in a complex coffee agroecosystem. *Biological Control* **61**, 89-97.

James S, Bell D. 2001. Leaf morphological and anatomical characteristics of heteroblastic Eucalyptus
globulus ssp. globulus (Myrtaceae). *Australian Journal of Botany* **49**, 259-269.

Khairi MMA, Hall AE. 1976. Comparative Studies of Net Photosynthesis and Transpiration of Some
Citrus Species and Relatives. *Physiologia plantarum* **36**, 35-39.

Kumar D, Tieszen LL. 1980. Photosynthesis in Coffea arabica. I. Effects of light and temperature.
Experimental Agriculture **16**, 13-19.

Lambers H, Oliveira RS. 2019. *Plant physiological ecology*. Switzerland: Springer.

Lee DW, Graham R. 1986. Leaf optical properties of rainforest sun and extreme shade plants.
American Journal of Botany **73**, 1100-1108.

Ma GR, Liu HQ, Tian SM, Bai XH, Zhao MZ, Li JH, Xiao ZW. 2019. Soil nutrient status in coffee
plantation of Yunnan and the main factors related with quality of green coffee beans. *Journal of Plant*
Nutrition and Fertilizers **25**, 154-161.

Martins SC, Galmés J, Cavatte PC, Pereira LF, Ventrella MC, DaMatta FM. 2014. Understanding
the low photosynthetic rates of sun and shade coffee leaves: bridging the gap on the relative roles of
hydraulic, diffusive and biochemical constraints to photosynthesis. *PloS one* **9**, e95571.

MATTA FD. Ecophysiological constraints on the production of shaded and unshaded coffee: a review.
Field Crops Research.

Miranda V, Baker NR, Long SP. 1981. Anatomical variation along the length of the *Zea mays* leaf in
relation to photosynthesis. *New Phytologist* **1981**, 595-605.

Niinemets Ü. 1999. Research review. Components of leaf dry mass per area – thickness and density –
alter leaf photosynthetic capacity in reverse directions in woody plants. *New Phytologist* **144**, 35-47.

Niinemets Ü, Sack L. 2006. Structural determinants of leaf light-harvesting capacity and
photosynthetic potentials. *Progress in Botany* **67**.

Nunes M, Bierhuizen J, Ploegman C. 1968. Studies on productivity of coffee: I. Effect of light,
temperature and CO2 concentration on photosynthesis of Coffea arabica. *Acta botanica neerlandica* **17**,
93-102.

Oguchi R, Onoda Y, Terashima I, Tholen D. 2018. Leaf anatomy and function. In: III WWA,
Terashima I, eds. *The leaf: a platform for performing photosynthesis*. Switzerland: Springer, 97-139.

Patricia BL, Laclau JP, Beri C, Mietton L, Muniz MRA, Arenque BC, De Cassia Piccolo M,
Lionel JM, Bouillet JP, Nouvellon Y. 2014. Photosynthetic and anatomical responses of E ucalyptus
grandis leaves to potassium and sodium supply in a field experiment. *Plant, Cell & Environment* **37**,
70-81.

Pompelli M, Martins S, Celin E, Ventrella M, DaMatta F. 2010. What is the influence of ordinary
epidermal cells and stomata on the leaf plasticity of coffee plants grown under full-sun and shady
conditions? *Brazilian Journal of Biology* **70**, 1083-1088.

Rodrigues WP, Silva JR, Ferreira LS, Machado Filho JA, Figueiredo FA, Ferraz TM, Bernado
WP, Bezerra LB, de Abreu DP, Cespom L, Ramalho JC, Campostrini E. 2018. Stomatal and
photochemical limitations of photosynthesis in coffee (Coffea spp.) plants subjected to elevated
temperatures. *Crop and Pasture Science* **69**, 317-325.

Smith WK, Vogelmann TC, DeLucia EH, Bell DT, Shepherd KA. 1997. Leaf form and
photosynthesis. *BioScience* **47**, 785-793.

Sun Y, Gu L, Dickinson RE, Pallardy SG, Baker J, Cao Y, DaMatta FM, Dong X, Ellsworth D,
Goethem DV, Jensen AM, Law BE, Loos R, Martins SCV, Norby RJ, Warren J, Weston D,
Winter K. 2014. Asymmetrical effects of mesophyll conductance on fundamental photosynthetic

parameters and their relationships estimated from leaf gas exchange measurements. *Plant, Cell & Environment* **37**, 978-994.

Taiz L, Zeige E. 2006. *Plant Physiology*. Sunderland Massachusetts Sinauer Associates, Inc.

Vogelman TC, Nishio JN, Smith WK. 1996. Leaves and light capture: light propagation and gradients of carbon fixation within leaves. *Trends in plant science* **1**, 65-70.

Wang Q, Chun A, J., Fleisher D, Reddy V, Timlin D, Resop J. 2017. Parameter Estimation of the Farquhar—von Caemmerer—Berry Biochemical Model from Photosynthetic Carbon Dioxide Response Curves. *Sustainability* **9**, 1228.

Ye Z-P. 2007. A new model for relationship between irradiance and the rate of photosynthesis in Oryza sativa. *Photosynthetica* **45**, 637-640.

Ye Z-P. 2010. A review on modeling of responses of photosynthesis to light and CO_2. *Chinese Journal of Plant Ecology* **34**, 727-740.

Ye Z-P, Yu Q. 2008. Comparison of new and several classical models of photosynthesis in response to irradiance. *Journal of Plant Ecology (Chinese Version)*, 149-154.

Zhang H, Schroder JL, Fuhrman JK, Basta NT, Storm DE, Payton ME. 2005. Path and Multiple Regression Analyses of Phosphorus Sorption Capacity. *Soil science society of America journal* **69**, 96-106.