Mechanisms of frustrated phagocytic spreading of human neutrophils on antibody-coated surfaces

Emmet Francis, Hugh Xiao, Lay Heng Teng, Volkmar Heinrich[*]

Department of Biomedical Engineering, University of California Davis, Davis, CA 95616, USA

*Author for correspondence (vheinrich@ucdavis.edu)

Running Title: Cell spreading in phagocytosis

Keywords: Motility, Cytoskeleton, Phagocytosis, Spreading, Adhesion

Summary

The detailed analysis of immune-cell spreading on antibody-coated surfaces establishes that active cytoskeletal protrusion rather than passive substrate adhesion drives phagocytic spreading.

Abstract

Complex motions of immune cells are an integral part of diapedesis, chemotaxis, phagocytosis, and other vital processes. To better understand how immune cells execute such motions, we present a detailed analysis of phagocytic spreading of human neutrophils on flat surfaces functionalized with different densities of immunoglobulin G (IgG) antibodies. We visualize the cell-substrate contact region at high resolution and without labels using reflection interference contrast microscopy (RICM) and quantify how the area, shape, and position of the contact region evolves over time. We find that the likelihood of the cell commitment to spreading strongly depends on the surface density of IgG, but the rate at which the substrate-contact area of spreading cells increases does not. Validated by a theoretical companion study, our results resolve controversial notions about the mechanisms controlling cell spreading, establishing that active forces generated by the cytoskeleton rather than cell-substrate adhesion primarily drive cellular protrusion. Adhesion, on the other hand, aids phagocytic spreading by regulating the cell commitment to spreading, the maximum cell-substrate contact area, and the directional movement of the contact region.

Introduction

Innate immune cells, in particular white blood cells, stand out among animal cells due to their deformability and extraordinary motility. Their mechanical flexibility enables them to carry out vital immune functions, such as chemotaxis toward sites of infection, and phagocytosis of foreign particles. These and other processes critically depend on the cells' ability to spread onto surfaces, for example, during firm arrest of white blood cells at the endothelium, or during the engulfment of target particles.

Phagocytic spreading is distinct from other forms of cell spreading in that it engages specific cell-surface receptors known to trigger phagocytosis (Cougoule et al., 2004; Flannagan et al., 2012; Lim et al., 2017). In many physiological situations, phagocytosis is initiated by opsonic ligands such as immunoglobulin G (IgG) antibodies and complement fragments. When bound to a pathogen surface, these ligands are recognized by the immune cells' Fcγ receptors (FcγRs) and complement receptors, respectively (Bournazos et al., 2016; Dustin, 2016). Ligation of these receptors activates receptor-specific intracellular signaling (Flannagan et al., 2012; Futosi et al., 2013; Rosales and Uribe-Querol, 2017; Uribe-Querol and Rosales, 2020) and ultimately leads to progressive spreading of the phagocyte over the target surface. In conventional phagocytosis, cell spreading concludes when the phagocytic cup closes to create a phagosome compartment. However, in some cases known as frustrated phagocytosis, the target is too large to be completely engulfed by a single phagocyte. During these encounters, immune cells strive to maximize the area of contact with the target surface, resulting in thinly spread, lamella-shaped cells that partially cover the target. Such frustrated phagocytic spreading has been observed, for instance, in interactions of human neutrophils with spherules of the fungal pathogen *Coccidioides posadasii* (Lee et al., 2015) as well as with large beads (Herant et al., 2005). The extent of spreading in these cases seems to be limited by the maximum apparent cell-surface area that a phagocyte can generate. For example, the maximum surface area observed during phagocytosis by human neutrophils amounts to about 300% of the surface area of the spherical resting shape of these cells (Hallett and Dewitt, 2007; Herant et al., 2005; Lee et al., 2015; Simon and Schmid-Schonbein, 1988), while for macrophages this plateau value is as high as 600% of the respective resting area (Cannon and Swanson, 1992; Lam et al., 2009).

The targets encountered by immune cells during frustrated phagocytosis are characterized by very low surface curvatures. Although curvature-dependent variations in phagocytic behavior have been reported (Cannon and Swanson, 1992; Champion and Mitragotri, 2006; Pacheco et al., 2013; Paul et al., 2013; Simon and Schmid-Schonbein, 1988), they do not appear to reflect qualitative changes in the cellular response but are likely due to the same response program coping with different "input" sizes and shapes of target particles (Herant et al., 2006; van Zon et al., 2009). Thus, it is reasonable to assume that the same preprogrammed cellular response also controls phagocytic spreading of immune cells on surfaces with zero curvature, i.e., flat substrates. Considering the high resolution of imaging methods that inspect cell-substrate interactions directly at the bottom of an experiment chamber, the study of frustrated phagocytic spreading of immune cells on such planar surfaces is a powerful tool to examine fundamental mechanisms of phagocytosis (Barger et al., 2019; Jaumouille et al., 2019; Kovari et al., 2016; Ostrowski et al., 2019).

Here, we present an in-depth quantitative analysis of human neutrophil spreading on IgG-coated coverslips. Our main objective is to address the lacking consensus between two different hypotheses that have been put forward to explain the mechanistic origin of phagocytic spreading, denoted as "Brownian zipper" and "protrusive zipper" mechanisms (Heinrich, 2015). Both hypotheses agree that fresh contact between the phagocyte membrane and target surface results in a zipper-like adhesive attachment that is essentially irreversible. However, they differ in their assumptions about the primary driving force of cellular protrusion that produces fresh cell-target contact in the first place. The Brownian zipper hypothesis views the phagocyte as a passive object and postulates that strong adhesion alone is responsible for pulling the cell membrane onto the target surface. In other words, this hypothesis treats phagocytic spreading as a wetting phenomenon, akin to a water droplet spreading on a hydrophilic surface. In contrast, the protrusive zipper hypothesis assumes that the phagocyte's cytoskeleton actively generates protrusive force to push the front of the cell outwards.

We investigate which of these hypotheses captures the main driving force of phagocytic spreading using well-controlled frustrated phagocytosis experiments. Our strategy is based on the following reasoning. If the Brownian zipper hypothesis of adhesion-driven spreading holds true, then both the rate as well as the extent of spreading should strongly depend on the density

of adhesive binding sites on the substrate. In contrast, because the protrusive zipper hypothesis

assumes that the spreading speed is primarily set by the rate of cytoskeletal protrusion, this

mechanism would predict a weaker dependence of the spreading speed on the ligand density. To

expose which type of behavior human neutrophils exhibit, we allow the cells to settle onto

coverslips pre-coated with different densities of rabbit IgG and analyze their subsequent

spreading dynamics. Visualization of the cells with reflection interference contrast microscopy

(RICM) allows us to image the cell-substrate contact region with exceptional contrast without

the use of labels. In addition to the spreading speed and maximum contact area, this analysis

enables us to further characterize the effects of IgG density on phagocytic spreading by

quantifying parameters such as the spreading probability, contour roundness, and centroid

motion. In an accompanying theoretical study, we leverage these results against computational

models of cell-spreading mechanics (Francis and Heinrich, 2022). Overall, we find that,

although the strength of adhesion can modulate phagocytic spreading, it is the cellular protrusion

that sets the rate of spreading, in support of the protrusive zipper hypothesis.

Results

Comparison with bead standards provides density of IgG on glass coverslips

We prepared surfaces displaying the Fc domains of IgG by saturating BSA-coated glass

coverslips with anti-BSA antibodies (Fig. 1). To adjust the surface density of human-reactive

IgG, we used incubation buffers containing different ratios of polyclonal rabbit IgG and

monoclonal mouse IgG-1. The total concentration of IgG was the same in all buffers. Because

mouse IgG-1 is hardly recognized by human Fcγ receptors (Lubeck et al., 1985;

Shashidharamurthy, 2010; Warmerdam et al., 1990), it served to dilute the amount of human-

reactive (rabbit) IgG decorating the coverslips. This approach appeared to produce consistently

functionalized substrates for cell-spreading experiments. However, because the rabbit and

mouse antibodies may have different affinities for BSA, the ratio of the two types of IgG

deposited on the surface is not necessarily the same as in the incubation buffer.

To quantify the actual amount of deposited rabbit IgG, we labeled IgG-coated coverslips with

fluorescent secondary antibody against rabbit IgG (Fig. 1A) and compared the measured

fluorescence intensity to a suitable standard. This standard consisted of batches of microspheres

pre-functionalized with known numbers of antibody-binding sites (QSC Kit; Bangs Laboratories,

Fishers, IN) that we saturated with the same secondary antibody as used to label the coverslips

(Fig. 1B). We analyzed Z-stacks of confocal images of these beads to determine the intensity of

the fluorescent layer at the underside of each bead local to the center of the bead image

(Materials and methods; Fig. S1). This analysis provided a calibration curve relating the

fluorescence intensity of an essentially flat layer of secondary antibody to the known surface

density of rabbit IgG (Fig. 1B). The calibration curve then allowed us to convert measurements

of the mean fluorescence intensity of confocal images of labeled coverslips (Fig. 1A) to the

density of rabbit IgG deposited on these coverslips. We note that the fluorescence intensity of

the most densely coated coverslips lay outside the intensity range of the bead standards; in this

case, our analysis involved an extrapolation of the calibration curve.

These measurements confirmed that our method to functionalize coverslips resulted in a

reproducible, broad spectrum of controlled rabbit-IgG densities. They also revealed that the

relationship between the rabbit-IgG concentration in the incubation buffer—denoted as the

"nominal" rabbit IgG concentration throughout this paper—and the actual surface density of

rabbit IgG on coverslips was nonlinear (Fig. 1C), indicating that the rabbit and mouse IgGs

indeed had different affinities for BSA. Fitting a mathematical model of competitive binding to

these data allowed us to estimate that the effective affinity of the used polyclonal rabbit antibody

was about 0.4 times that of the monoclonal mouse IgG. The measured surface densities of rabbit

IgG ranged from below our detection limit (at 0.1% nominal rabbit-IgG concentration where the

surface density was indistinguishable from our negative control) to a maximum density of

~25,000 rabbit-IgG molecules per μm^2 (at 100% nominal rabbit-IgG concentration). Assuming a

uniform distribution of bound IgG molecules, the maximum surface density corresponded to an

area of roughly 40 nm^2 occupied by each rabbit IgG, consistent with a dense, essentially

contiguous layer of IgG coating the coverslips.

Cells are more likely to spread on higher densities of IgG

We used an IgG-coated coverslip as the bottom of our assembled experiment chamber (Fig.

2A,B). A plastic insert with funnel-shaped vertical thru holes served as the chamber's top (Fig.

2A). The chamber was filled with Hanks' balanced salt solution supplemented with 2% human

serum albumin and placed on an inverted research microscope set up for epi-illumination. This

design allowed us to introduce small volumes of a suspension of isolated human neutrophils at

non-overlapping locations through the chamber ceiling. Each deposited cell population

dispersed to some extent by diffusion while sinking to the chamber bottom. The cells settled

onto the functionalized coverslip within ~5 minutes, usually without piling into clusters.

We imaged the cells using reflection interference contrast microscopy (RICM, Fig. 2C,D,

Movie S1). The interference patterns produced by RICM markedly enhanced the visibility of

contact regions between the primary light-reflecting interface—the top surface of the coverslip—

and the underside of reflective objects resting on this surface. Regions of direct cell-substrate

contact generally appeared as clearly outlined, distinctive dark patches. It is important to bear in

mind though that light-reflecting structures inside a cell, such as granules, can produce brighter

spots within the dark contact region that could be misinterpreted as areas where part of the cell

has detached from the substrate. Close inspection of the interference patterns revealed that in our

experiments, contiguous contact regions did not appear to contain local areas of cell-substrate

separation.

Contact with the substrate caused a fraction of the cells to start spreading either immediately

or after some lag time. Despite the mixed response of the cells in a given population—reflecting

cell-to-cell variation typical for the behavior of primary human immune cells—we found that the

overall proportion of spreading to non-spreading cells strongly depended on the density of rabbit

IgG coating the coverslip. We quantified the cells' commitment to spreading on a given surface

in terms of the "spreading probability", i.e., the ratio between the number of spreading cells and

the total number of observed cells. We consistently evaluated this ratio 30 minutes after

depositing the cells, defining "spreading cells" as those cells whose cell-substrate contact area

had reached at least $100 \, \mu m^2$ at this time.

The measured spreading probabilities ranged from 0.16 on the lowest density of surface-

bound rabbit IgG to about 0.85 on the highest density (Fig. 3). The result for the lowest density

was not significantly different from the spreading probability on substrates presenting pure

mouse IgG-1 (negative control). While this result may be attributed in part to spurious

recognition of mouse IgG-1 by human neutrophils, it also highlights that neutrophil populations

almost always contain cells that can be activated by non-specific interactions with otherwise

non-stimulatory target surfaces. On the other hand, almost all cells spread on substrates coated

with high-density, 100% rabbit IgG, confirming that antibody-coated targets generally draw a

vigorous response by human neutrophils.

We limit our remaining analysis to spreading neutrophils. Thus, when interpreting the

following results, it is important to bear in mind that they were obtained with selected

subpopulations of cells, and that the fraction of cells making up each subpopulation depended on

the cell response to a particular surface density of rabbit IgG.

The cell-substrate contact region of spreading cells is essentially radially symmetric

After settling on the bottom coverslip, individual neutrophils typically formed one or two initial

attachment spots with the IgG-coated substrate. In cases were more than one spot was visible,

the growing spots quickly merged into a single, contiguous region of cell-substrate contact.

Subsequently, the contact region of spreading neutrophils generally had a roughly circular shape

that expanded in a radially symmetric manner, as expected for phagocytic spreading (Fig. 4). It

typically took neutrophils only about 2 minutes from the start of spreading to form contact

footprints with diameters as large as 20 μm.

After the contact area reached an apparent plateau, neutrophils exhibited a spectrum of

different behaviors. Some remained in place but continued to modify the cell-substrate contact

region, gradually expanding or contracting it, or altering its shape while leaving its net area

essentially unchanged. Other cells appeared to adopt a more migratory phenotype, attempting to

crawl away from the original location of spreading. Common to most post-spreading behaviors

was the formation of trailing membrane extrusions at retracting cell regions. These structures

connected the main cell body to focal cell-substrate adhesion sites from which the moving cell

often was unable to detach (Fig. 4). We have not attempted to analyze this highly variable post-

spreading behavior beyond a coarse empirical classification (Fig. 4), demonstrating that cells

spreading on higher densities of IgG were less likely to try to leave their original spot of

spreading (Fig. 4C).

Density of IgG does not affect spreading speed but weakly correlates with extent of spreading

Our quantitative comparison of neutrophils spreading on different densities of IgG is based on

the analysis of image sequences recorded at intervals of 2 s. We semi-automatically traced the

outline of the cell footprint in each image and stored the resulting polygons of cell peripheries

(Fig. 5A). Our primary quantity of interest was the area circumscribed by these polygons, i.e.,

the cell-substrate contact area. Plots of the contact area as a function of time revealed a

comparatively slow, brief initial spreading phase, followed by a roughly linear expansion of the

contact footprint, and eventually, a gradual approach to a plateau of maximum contact area (Fig.

5A-C).

We define the speed of spreading as the value of the steepest slope of a given contact-area-

versus-time graph. We used sigmoidal or linear fits to such curves to determine the spreading

speeds in suitable experiments. Remarkably, the average spreading speeds, measured over a

1000-fold range of nominal IgG concentration, were all close to 3 μm^2/s, exhibiting no

significant differences on different IgG densities (Fig. 5D).

The similarity in neutrophil behavior during the rapid-spreading phase also was evident in

example curves measured for cells spreading on 0.1% versus 100% rabbit IgG (Fig. 5B). On the

other hand, the collection of graphs exposed an apparent difference between the maximum

contact areas occupied by cells spreading on these two IgG densities. Most cells spreading on

100% IgG reached contact areas in the range of 200-300 μm^2, whereas on the substrates coated

with the lowest IgG density (0.1%), the maximum contact area tended to lie in the range of 150-

250 μm^2. The statistical analysis of all measurements confirmed that the mean maximum contact

areas on surfaces coated with 10% or 100% IgG were indeed significantly larger than those on

0.1% IgG (Fig. 5E).

Recalling that originally spherical cells (with diameter D_0) must increase their apparent

surface area when spreading on a substrate, it is instructive to convert the measured area of cell-

substrate contact footprints into estimates of the overall cell-surface area (A) required to

accommodate such deformations. Our conversion assumes that the cell volume remains constant

during spreading, and that the cell-substrate contact region is circular, denoting its diameter by

D_c. Furthermore, we approximate the shape of the upper, free surface of the spread cell as a

spherical cap, which is the geometry with the smallest possible surface area for the given volume

and a circular contact footprint. For this simplified geometry, one can calculate the ratio

between the final area A and the initial cell-surface area $A_0 = \pi D_0^2$ using the following

prescription

$$\frac{A}{A_0} = \frac{1}{4}\left(\frac{x_c^4}{q} + q\right) \qquad \text{where} \qquad x_c \equiv \frac{D_c}{D_0} \quad \text{and} \quad q \equiv \left(\sqrt{x_c^6 + 16} - 4\right)^{2/3} \qquad (1)$$

The diameter of most non-adherent, resting human neutrophils lies in the range of 8.5-9 µm.

With the choice of $D_0 = 8.75$ µm, Eq. (1) predicts that a typical neutrophil will need to expand

its surface area to ~210% of its initial area in order to form a contact footprint whose diameter is

twice as large as the cell's resting diameter (where $x_c = 2$ and $D_c = 17.5$ µm).

The average of the 10 largest cell-substrate contact areas measured in our experiments is 359

µm^2, corresponding to a typical footprint diameter of $D_c = 21.4$ µm. Thus, we estimate that the

most-spread neutrophils have expanded their apparent surface area to about 303% of their resting

area during our frustrated-phagocytosis experiments, in agreement with previous findings

(Hallett and Dewitt, 2007; Herant et al., 2005; Lee et al., 2015; Simon and Schmid-Schonbein,

1988).

Higher IgG density results in more uniform, concentric spreading

The results of the previous section provide compelling support for the protrusive-zipper

hypothesis. A detailed analysis of additional features of the cell behavior is likely to produce

further insight into the fundamental mechanisms of phagocytic spreading. We next examined

how the IgG density affects the roundness and type of motion of the cell footprint during

spreading.

We assessed the roundness of the cell-substrate contact region in terms of the ratio of the

radii of two circles defined by the outline of this region, i.e., the largest inscribed circle and the

smallest circumscribed circle (Fig. 6A). The value of this roundness measure equals 1 for

circular contours and is smaller otherwise, with lower values corresponding to less round

contours. For each spreading cell, we measured the roundness of individual contours within the

phase of fastest contact-area growth (where the spreading speed was greater than 0.4 µm^2/s),

then applied a moving average filter with a window size of 8 s and finally determined the

maximum of these averages. This approach yielded roundness values in the range of roughly
0.6-0.9 that exhibited no significant differences between surfaces coated with different densities
of IgG (Fig. 6B). Thus, the cell contours generally were characterized by a moderate roundness
that was independent of the IgG density, indicating that phagocytic spreading proceeded in a
more or less uniform fashion on all tested surfaces.

The roundness measure does not capture unambiguous information about possible lateral
displacements of the cell footprint. To quantify the latter, we analyzed the positions of the
centroid of the cell-substrate contact region of spreading cells in suitable video images. We
defined the centroid displacement Δr as the Euclidean distance by which the centroid position
moved during a given time interval Δt (Fig. 6C). We used these measurements to estimate the
instantaneous velocity of the centroid motion in terms of the ratio $\Delta r/\Delta t$ evaluated for
successively recorded centroid positions. For each spreading cell, we then averaged the
instantaneous values over the active spreading phase. Most of the resulting average centroid
speeds were in the range of 0.01-0.06 µm/s and exhibited no significant differences between
substrates coated with different IgG densities (Fig. 6D).

In addition to evaluating the centroid speed, we also used the measured centroid positions to
assess the type of motion of the cell footprint. Now choosing Δt-values in the range from 4 s to
60 s, we averaged the squared centroid displacements of a given cell over all pairs of video
frames whose recording times differed by Δt. We then calculated the averages of these mean
square displacements as a function of Δt for the whole population of cells spreading on a given
substrate (Fig. 6E). Power-law fits of the form $y = ax^b$ to plots of these mean square
displacements as a function of Δt allow for a rough classification of the type of centroid motion.
An exponent of $b = 1$ of such a fit implies a random motion with Gaussian-distributed
individual displacements, as is typical, for example, for diffusion. If the exponent approaches 2,
the motion generally is interpreted as more migratory. The exponent of the power-law fit to the
measured mean square centroid displacements was 1.57 on the highest density of IgG, whereas
on the lowest IgG density it was 1.86, suggesting that lateral displacements of cells spreading on
the highest IgG density were more random, i.e., the cells were less likely to transiently explore
the substrate in a preferred direction.

The lack of a clear correlation between IgG density and centroid speed (Fig. 6D) contrasted with the observed differences in centroid displacement (Fig. 6E), suggesting that on some substrates, moving cells changed direction more frequently than on others. We verified this assumption by assessing the directional persistence of centroid displacements in terms of the decay of the time-dependent autocorrelation function of the tangents of the centroid trajectory (Gorelik and Gautreau, 2014). The resulting "directional persistence time" (Materials and methods) exposes for how long a cell footprint continues to move in roughly the same direction. The measured persistence times generally were small and exhibited a large spread; however, they were significantly shorter (~4.4 s on average) on coverslips coated with the highest density of IgG than on lower IgG densities (~7.5-8.3 s) (Fig. 6F). The averages of the closely related "directional persistence distances" (product of centroid speed and persistence time) traversed in a preferred direction were small, falling in the overall range of 0.16-0.36 µm, and exhibiting a significant difference between substrates coated with nominal IgG concentrations of 100% and 1% (data not shown). In summary, transient lateral polarization and directional movement of the cells, while generally unlikely, indeed met with stronger resistance on higher densities of IgG.

This analysis confirmed that the observed type of spreading motion was specific to the phagocytic ligand IgG. Overall, cells spreading on IgG tended to expand the contact region in a uniform, concentric manner without extending substantial exploratory protrusions in preferred directions. As shown earlier (Fig. 4), this behavior may change once the area of the cell-substrate contact region has reached an apparent plateau.

Computer simulations reinforce mechanistic insights gained from experiments

The comparison of experimental results with theoretical predictions is an essential approach to test mechanistic hypotheses and decide whether observed correlations are coincidental or causative. In a companion study, we have translated different hypotheses about the driving force of phagocytic spreading into mathematical models and examined the mechanical ingredients that such models require to reproduce the cellular behavior (Francis and Heinrich, 2022). This section provides a broadly accessible summary of our comparison between the measured dynamics of the growth of the cell-substrate contact area and theoretical predictions based on the Brownian zipper and the protrusive zipper models, respectively.

Details of our mathematical models are provided in the companion paper (Francis and

Heinrich, 2022). In short, we consider an immune cell as an axisymmetric body with uniform

surface tension that is filled with a highly viscous, incompressible fluid (Tran-Son-Tay et al.,

1991; Yeung and Evans, 1989). The model incorporates adhesive interactions between the cell

surface and a flat, rigid substrate in terms of a short-range attractive force that acts to pull the cell

onto the substrate in an irreversible manner. This basic version of the model captures the

essential assumptions of the Brownian zipper hypothesis.

To introduce active cellular protrusion into this framework, we postulate that fresh contact

between the substrate and the cell surface results in a transient, outward-pushing force local to

the leading edge of the cell-substrate contact region. We do not attempt to model the

microscopic origin of this protrusive force, which would have to account for receptor-induced

signaling, spatiotemporal redistribution of messenger molecules, translation of biochemical cues

into mechanical stresses, etc. Instead, our semi-empirical approach lumps these highly complex

processes into the rationale that receptor activation due to fresh cell-substrate contact ultimately

causes the protrusive force. We also assume that the magnitude of protrusion decays as a

function of the time elapsed after the most recent receptor-ligand binding event. One objective

of our theoretical study is to establish what distribution and magnitude of mechanical stresses

can account for the experimental results, and whether the predicted stresses are physically

realistic and biologically plausible. The model version that includes the protrusive force

represents the protrusive zipper hypothesis.

As reasoned earlier, adhesion-driven, passive spreading, which forms the core of the

Brownian zipper hypothesis, should result in a stronger correlation between the IgG density and

the spreading speed than active, primarily protrusion-driven spreading. To verify this common-

sense expectation and place it on sound quantitative grounds, we simulated the dependence of

the spreading speed on the IgG density using both versions of our mathematical model (Fig. 7).

The simulations indeed revealed a dramatic difference in predicted cell-spreading behavior

between these two models. For realistically chosen parameter values, the Brownian zipper

model predicts a strong dependence of the cell spreading speed on the density of IgG binding

sites, and the spreading speeds on low IgG densities are predicted to be much slower than

observed experimentally. In contrast, the spreading speed predicted by the protrusive zipper

model hardly changes over a 1000-fold range of nominal IgG concentrations and is similar to

experimental results. Comparison of the predictions of Fig. 7D with the measurements shown in

Fig. 5C,D clearly supports the conclusion that the protrusive zipper hypothesis successfully

captures the basic mechanisms underlying phagocytic spreading, whereas the Brownian zipper

hypothesis does not.

The dependences of the maximum contact area on the IgG density (Fig. 7D) reveal another

difference between the two model versions. Recall that our experiments exposed a significant

increase of the maximum contact area at higher IgG densities (Fig. 5E). However, this increase

is much smaller than the large differences between the plateau values of the graphs predicted by

the Brownian zipper model (Fig. 7D). On the other hand, the predictions of the protrusive zipper

model again provide a notably better match with our experimental observations (Fig. 7D).

Discussion

The rapid execution of complex cellular motions is one of the most amazing feats of immune

cells, and critical to phagocytosis, cell migration, and other vital processes. However, precise

descriptions of such motions and, more importantly, understanding of the underlying

mechanisms, remain scarce. The quantitative analysis of frustrated phagocytosis presented in

this paper is part of a larger research area that focuses on the biomechanics of cells spreading on

surfaces, comprising both experimental work (Dobereiner et al., 2004; Dubin-Thaler et al., 2004;

Dubin-Thaler et al., 2008; Hategan et al., 2004; Janmey et al., 2021; Reinhart-King et al., 2005;

Wolfenson et al., 2014) as well as theoretical studies (Chamaraux et al., 2005; Cuvelier et al.,

2007; DiNapoli et al., 2021; Etienne and Duperray, 2011; Fardin et al., 2010; Frisch and

Thoumine, 2002; Odenthal et al., 2013; Xiong et al., 2010). Our integrative

experimental/theoretical approach expands most previous analyses by quantifying how the ligand

density affects the cells' spreading behavior, and to what extent adhesive versus protrusive forces

contribute to the cell motion. In conjunction with our theoretical companion paper, this approach

has allowed us to resolve controversial mechanistic notions about the spreading behavior of

immune cells.

Our experimental strategy has been to expose human neutrophils to target surfaces that

presented a range spanning three orders of magnitude in density of deposited IgG. The strong

dependence of the cells' spreading probability on the IgG density confirms that the ligation of

Fcγ receptors is the original cause of spreading. It also suggests that a threshold in cell

stimulation needs to be reached to induce spreading. The most likely type of stimulating trigger

appears to be a minimum number of engaged receptors. Alternatively, the primary stimulus

could be the spacing between engaged receptors, in which case their total number might play a

lesser role. Our current setup does not allow us to distinguish between these two alternatives. It

also is important to bear in mind that the threshold level that triggers spreading can vary

considerably from cell to cell, suggesting that it is affected by other factors such as the degree of

quiescence of individual neutrophils.

The absence of a significant correlation between the IgG surface density and the spreading

speed of the cells is perhaps the most consequential result of our experiments. It provides a

counterexample to the notion that spreading might be driven by strong cell-substrate adhesion,

thus discrediting the Brownian zipper hypothesis in the case of neutrophils and similar cells. In

contrast, the protrusive zipper hypothesis intuitively provides a better explanation of our results.

The strongest support for the validity of the latter hypothesis comes from the agreement between

computer simulations based on the protrusive zipper model and our experiments performed on

different substrates. Our finding agrees well with reports that during isotropic fibroblast

spreading, the speed of spreading depended on protrusive stress generated by the actin

cytoskeleton (Dubin-Thaler et al., 2008; Fardin et al., 2010; Xiong et al., 2010).

It is worth noting that the spreading speed of macrophage-like cells of the J774 murine cell

line on IgG-coated surfaces also was found to be largely independent of the IgG density (Kovari

et al., 2016). Thus, our conclusion that cell spreading is primarily driven by an active protrusive

force likely holds for motile immune cells in general. It also agrees with a previously reported

"all-or-nothing" signaling response during IgG-mediated phagocytosis by macrophages (Zhang

et al., 2010), as well as with other studies in which macrophages readily consumed particles

coated with low densities of IgG (Ben M'Barek et al., 2015; Pacheco et al., 2013). On the other

hand, it conflicts with computer simulations that neglected the role of cytoskeleton-driven

protrusion and predicted that phagocytosis should proceed most quickly at an optimal ligand

density (Richards and Endres, 2014).

Our finding that the spreading probability, but not the spreading speed, depends on the IgG density reveals that beyond the stimulus threshold that causes a cell to commit to spreading, further strengthening of the stimulus appears to have little effect on the mechanical cell response. In other words, the spreading speed becomes decoupled from the stimulus strength at some point. This raises the interesting question at which step in the sequence of processes leading from receptor ligation to cell movement the decoupling occurs. Does receptor-induced signaling reach a limiting level beyond which the activation of additional receptors is ignored? Is there a bottleneck built into the redistribution of messenger molecules? Or are the cell's resources to generate additional protrusive force exhausted? Answers to questions like these, although beyond the scope of this study, are an essential part of a comprehensive understanding of vital immune-cell functions.

Our analyses of the geometry and type of motion of the contact footprint, as well as the post-spreading behavior of neutrophils, round out this study, allowing us to paint a detailed picture of the events following cell contact with an IgG-coated surface. After reaching the surface, an initially quiescent cell may undergo detectable Brownian motion and/or be dragged along the surface by convection until cell-substrate adhesion causes its arrest. Cell arrest is usually faster—often immediate—on surfaces coated with higher densities of IgG, suggesting that it is mediated by specific bonds between immobilized Fc domains and Fcγ receptors of the cell. It is worth bearing in mind though that neutrophils can arrest even on negative control surfaces through nonspecific interactions.

Ligation of Fcγ receptors ultimately leads to cell activation and spreading. Many intermediate mechanistic details of this response remain uncertain, but our results strongly suggest that an active protrusive force generated by the cell local to the periphery of the contact footprint is the main driver of spreading. There is ample qualitative evidence supporting the cytoskeleton's key role in the generation of this force through local actin polymerization and cross-linking, both generally (Footer et al., 2007; Kovar and Pollard, 2004; Parekh et al., 2005) as well as in the specific context of phagocytosis (Allen and Aderem, 1996; Freeman and Grinstein, 2014; Jaumouille et al., 2019; Ostrowski et al., 2019). For example, F-actin accumulation at the front of the phagocytic cup has been shown to be correlated with force production during macrophage phagocytosis (Nelsen et al., 2020). Novel techniques to measure

phagocytic forces with high resolution hold great promise for illuminating the detailed

mechanisms linking actin remodeling to force generation (Vorselen et al., 2021; Vorselen et al.,

2020). It seems plausible to ascribe the local character of this cytoskeletal remodeling to

localized signaling triggered by fresh cell-substrate contact.

Although cell-substrate adhesion is not the primary cause of spreading, it still plays a critical

role by securing and stabilizing fresh contact regions, thus aiding further cell motion parallel to

the surface. A subtle detail in the cell's control of its motion is the coordination of structural

linkages between the actin cytoskeleton and the plasma membrane. Weakening of such linkages

at the protruding front is likely to facilitate local forward displacements of the cell surface. A

molecular example of such behavior is the calcium-dependent disruption of FERM-mediated

linkages by calpain (Dewitt and Hallett, 2002; Dewitt and Hallett, 2020; Roberts and Hallett,

2019). In contrast, in regions of established cell-substrate contact, strengthening of membrane-

cytoskeleton linkages prevents the main cell body from being lifted away from the surface, while

also providing bracing support to the protrusive motion along the surface (Barger et al., 2019;

Herant et al., 2011; Jaumouille et al., 2019; Jaumouille and Waterman, 2020). Several molecular

players that either link integrins to the cytoskeleton (e.g., talin, paxillin) or directly link the

membrane to the cytoskeleton (e.g., myosin I, ezrin) have recently been shown to colocalize with

the neutrophil or macrophage phagocytic cup and facilitate directional force generation (Barger

et al., 2019; Jaumouille et al., 2019; Ostrowski et al., 2019; Roberts et al., 2020b).

Once the cell commits to spreading, it generally expands its contact footprint in an isotropic,

essentially concentric fashion, resulting in a more or less round contact region throughout the

spreading phase. This behavior is fundamentally different from migratory cell motions like

chemotaxis, which require sustained cell polarization not only in terms of morphology and

mechanics, but also intracellular signaling. It seems logical to attribute this difference to the

nature of the stimulus, which is uniform in one case and localized in the other. However,

especially our substrates coated with low densities of IgG are unlikely to present perfectly

uniform IgG layers. Yet even on these substrates, the cell roundness was unaffected by the IgG

density, providing further support to the "all-or-nothing" notion of phagocytic spreading. This

result also makes sense biologically, considering that a phagocyte should be able to engulf

pathogens that are not uniformly coated with antibodies. Finally, our conclusion that phagocytic

spreading tends to proceed in a symmetric fashion is particularly impactful for theoretical

investigations, as it reinforces the validity of predictions made by mathematical models of

phagocytosis that assume an axisymmetric cell-target configuration.

That said, we did observe larger centroid displacements and a significantly higher directional

persistence of cells spreading on lower densities of IgG, in agreement with observations of

neutrophils spreading on different densities of fibronectin or BSA (Henry et al., 2014) as well as

other cell types, for instance, endothelial cells that spread more unevenly on lower densities of

Arg-Gly-Asp (RGD) (Reinhart-King et al., 2005). This behavior appears to resemble

haptokinesis, albeit on a very small scale considering that the overall cell displacements rarely

exceeded 1 μm.

Our picture of phagocytic spreading would be incomplete without considering its limits.

What determines the maximum extent of spreading? Key to answering this question is the

mechanical resistance to expansion of the apparent cell surface area, i.e., the cortical tension.

This ever-present tension maintains the spherical shape of quiescent immune cells, and it rises as

the cell surface area increases during shape changes (Herant et al., 2005; Lee et al., 2011; Lee et

al., 2015; Roberts et al., 2020a). Geometrical arguments similar to those leading to Eq. (1) show

that the apparent cell surface area grows monotonically as the cell-substrate contact region

increases. Thus, the protrusive force generated by the cytoskeleton meets with increasing

mechanical resistance during cell spreading. The magnitude of the outward pushing force is not

unlimited, and neither is the amount of material that can be recruited to the cell surface to enable

its further growth. Either one of these closely related constraints will stall spreading.

Remarkably, our estimate of the maximum apparent cell surface area produced during frustrated

phagocytic spreading agrees well with values measured previously in various types of

experiments (Hallett and Dewitt, 2007; Herant et al., 2005; Lee et al., 2015; Simon and Schmid-

Schonbein, 1988).

Even before the cell-substrate contact region reaches its maximum extent, the rising

resistance due to the cortical tension slows spreading. A higher substrate density of adhesive

IgG ligands is expected to aid further spreading at this stage, because the distance over which the

protruding cell membrane needs to be displaced before reaching the next adhesion site is smaller.

The observed dependence of the maximum extent of spreading on the IgG density (Fig. 5E)

confirms that this is indeed the case. Yet, this dependence is much less pronounced than

predicted by the Brownian zipper model. This finding underlines the role of adhesion as an

important facilitator of spreading, regulating the cell's commitment to spreading, the maximum

contact area, and the directional movement of the contact region. However, the results of this

and other studies leave little doubt that active cellular protrusion rather than adhesion is the

primary driving force of phagocytic spreading.

Materials and methods

Surface preparation

After partial assembly of the 3D-printed chamber (Fig. 2A), the surface of the glass coverslip

serving as chamber bottom was incubated with 1% (10 mg/ml) bovine serum albumin (BSA;

VWR, Radnor, PA) in phosphate buffered saline (PBS; Bio-Rad, Hercules, CA) for 1 hour at

room temperature, then rinsed with PBS. The surface was then incubated with a mixture of

polyclonal rabbit anti-BSA IgG (Cat# A11133; Invitrogen, Waltham, MA) and monoclonal

mouse anti-BSA IgG-1 (Cat# MA182941, Invitrogen, Waltham, MA) for another hour at room

temperature. All mixtures contained a total of 150 µg/ml IgG, which was sufficient to saturate

the surface, but the ratio of rabbit to mouse IgG concentration was varied to produce solutions

containing 0%, 0.1%, 1%, 10%, or 100% rabbit IgG. After a final rinse with Hanks' balanced

salt solution (HBSS with Ca^{2+} and Mg^{2+}; Thermo Fisher Scientific, Waltham, MA), the chamber

was loaded with HBSS containing 2% human serum albumin (HSA; MP Biomedicals, Irvine,

CA) to block any uncoated regions.

To quantify the actual density of rabbit IgG on coated glass surfaces, coverslips of each batch

were incubated with 30 µg/ml anti-human IgG Fc (cross-reacts with rabbit) conjugated to Alexa

Fluor-488 (Cat# 409321; BioLegend, San Diego, CA) for 45 minutes at room temperature in the

dark, then rinsed with PBS before imaging. Beads from the Quantum Simply Cellular Kit (QSC;

Bangs Laboratories, Fishers, IN) were incubated with the same secondary antibody at 6.25

µg/ml, again for 45 minutes at room temperature in the dark, then washed 3 times in PBS with

0.01% Tween and deposited onto plain glass coverslips for imaging.

Quantification of IgG surface density from fluorescence microscopy

As outlined in the main text (Fig. 1), we determined the density of deposited rabbit IgG by

comparing the fluorescence intensities of labeled coverslips to a calibration curve created using a

set of QSC beads with known numbers of antibody-binding sites. All fluorescence images were

taken on a spinning disc confocal microscope (488 nm laser, 100x oil objective, NA = 1.4). Due

to the large spread of fluorescence intensities of the inspected surfaces, we used two different

exposure times $t_e = 1$ s and $t_e = 2$ s and adjusted the originally measured intensities I to a

corrected value $\bar{I}$ using

528
$$\bar{I} = \frac{I - I_0}{t_e} \tag{2}$$

where I_0 is the background intensity, i.e., the intensity at the substrate surface in the absence of

fluorophore. To minimize photobleaching, we first preadjusted the microscope focus at a

suitable location, then moved the microscope stage to a neighboring field of view and acquired a

Z-stack of images of this unbleached region.

In fluorescence measurements of coverslips, we chose a region of interest of at least 10,000

pixels and determined the most frequent pixel intensity in this region for each image using kernel

density estimation. After plotting the most frequent intensity values of all images of a given

stack as a function of Z-height and fitting a cubic polynomial to these data, we calculated the

maximum of the polynomial fit and identified its function value as the fluorescence intensity of

the plane containing the fluorophore.

A more elaborate procedure was needed to measure the pertinent fluorescence intensity of

labeled QSC beads. We reasoned that the fluorescence at the underside of a bead resting at the

chamber bottom is most closely relatable to the fluorescence at the top surface of labeled

coverslips. Accordingly, we measured for each selected bead the fluorescence intensity at the

center of all bead images in a given Z-stack and then again used polynomial fits to determine the

interpolated maximum value of these intensities, which we identified as the fluorescence

intensity corresponding to the known density of fluorophore coating the bead. Details of this

procedure are illustrated in Fig. S1. Having quantified the relationship between the surface

density of rabbit IgG and the fluorescence intensity in this manner, we were able to establish the

number of rabbit IgG molecules per μm^2 of the surface of functionalized coverslips for each of

the used IgG incubation mixtures, i.e., for each nominal concentration of rabbit IgG (Fig. 1).

Cell isolation

The protocol for this study was approved by the Institutional Review Board of the University of

California, Davis and all donors provided written informed consent. Human neutrophils were

isolated from whole blood of healthy donors by immunomagnetic negative selection using the

EasySep Direct Human Neutrophil Isolation Kit (STEMCELL Technologies, Vancouver,

Canada). Isolation began from an initial blood volume of 0.5-1 ml, and was carried out

according to the protocol described by the manufacturer. Immediately following isolation, the

cells were resuspended at $5\text{-}10\times10^6$ cells per ml in HBSS without Ca^{2+} and Mg^{2+} (Thermo Fisher

Scientific) to ensure their quiescence prior to stimulation by contact with IgG.

Contact area analysis

The cell-substrate contact area was quantified using customized MATLAB software based on

code graciously provided by Daniel Kovari (Kovari et al., 2016). In brief, the RICM image was

thresholded on both intensity and variance to construct an initial binary image by detecting

regions where the local variance was greater than σ^2_{min} or the intensity was less than I_{max} (both

thresholds could be adjusted on an image-by-image basis). The binary image was then

convolved with a Gaussian kernel to smooth the outline of the cell footprint. Holes in the

footprint were filled to define the total contact area. In a few cases where the automatic

recognition algorithm performed poorly (especially during early spreading), the cell outlines

were traced manually.

The spreading speed was quantified using sigmoidal fits to contact-area-versus-time graphs

whenever possible. When necessary, a linear fit was used over a smaller range of the contact

area data. The maximum contact area was determined as the average over the plateau region of

the curve. To calculate the roundness of the cell-substrate contact region, we determined the

largest inscribed and the smallest circumscribed circle of the cell footprint using open-source

MATLAB functions (D'Errico, 2017; Warrier, 2017) (Fig. 6A) and took the ratio the radii of

these two circles.

Analysis of centroid directional persistence

We restricted our analysis of the directionality of the motion of the geometric centroid of the

substrate contact region of each cell to the phase of fastest contact-area growth, where the

spreading speed was greater than 0.4 μm^2/s. To assess for how long the centroid motion

typically maintained a preferred direction, we evaluated the directional autocorrelation function

(Gorelik and Gautreau, 2014). First, for each pair of successive video frames (recorded at times

t_i and t_{i+1}, respectively), we determined the angle $\theta(t_i)$ between the *x*-axis and the line of

centroid displacement using

$$\theta(t_i) = \arctan \frac{y_{i+1} - y_i}{x_{i+1} - x_i} \tag{3}$$

585 Then, after identifying all pairs of angles $\theta(t_i)$ and $\theta(t_j)$ along a given centroid trajectory for

586 which the time gap $t_j - t_i$ equaled a given value Δt, we calculated the autocorrelation function

587 AC(Δt) defined by

$$AC(\Delta t) = \left\langle \cos\left[\theta(t_j) - \theta(t_i) \right] \right\rangle = \left\langle \cos\left[\theta(t_i + \Delta t) - \theta(t_i) \right] \right\rangle \tag{4}$$

where the angular brackets $\langle \ldots \rangle$ are used to denote the average over all such pairs. Persistent

times are typically defined as the time constants of exponentially decaying autocorrelation

functions; however, because of a limited number of data points in this analysis, the

autocorrelation functions of the centroids of individual cells often did not lend themselves to

exponential fits. Instead, recalling that the persistence time of an exponential decay equals the

area under the autocorrelation curve, we here defined the directional persistence time by

$$\tau_{pers} = \int_{0\,\text{s}}^{16\,\text{s}} AC(\Delta t)\,d(\Delta t) \tag{5}$$

Computational modeling

Details of our computer model, including all equations and parameter values, are presented in

our companion paper (Francis and Heinrich, 2022). In short, we represent the cell interior as a

highly viscous Newtonian fluid and assume mechanical equilibrium. This allows us to solve a

perturbed form of the Stokes equations at each time step using the finite element method,

yielding the flow profile inside the cell. In order to complete this calculation, the relevant

stresses are computed for a given shape of the cell, comprising *(i)* adhesion stress, assuming an

attractive potential acting between the cell membrane and the IgG-coated surface, *(ii)* protrusion

stress, assuming this stress acts normal to the cell membrane and decays exponentially along the

membrane, and *(iii)* cortical stress, assuming a uniform cortical tension whose magnitude

depends on the amount of surface area expansion as previously quantified for human neutrophils

(Herant et al., 2005). All calculations are carried out in MATLAB; our code is available at

https://github.com/emmetfrancis/phagocyticSpreading.

Statistics

All tests for significance were conducted using one-way ANOVA followed by Tukey's post-hoc

test in Origin software. Values more than 1.5 interquartile ranges above the upper quartile or

below the lower quartile were excluded as outliers prior to ANOVA.

Acknowledgements

We thank Dr Soichiro Yamada for his help with confocal microscopy, Dr Daniel Kovari for
providing MATLAB code for the analysis of the cell-substrate contact regions, and Dr Robert
Guy for many helpful conversations regarding our computational model of cell spreading.

Competing interests

The authors declare no competing or financial interests.

Funding

This work was supported by grant R01 GM098060 from the National Institutes of Health, USA,
National Science Foundation Graduate Research Fellowship, award number 1650042, and a
graduate student research award from the ARCS Foundation.

References

Allen, L. A. and Aderem, A. (1996). Molecular definition of distinct cytoskeletal structures

involved in complement- and Fc receptor-mediated phagocytosis in macrophages. *J Exp Med* **184**, 627-

37.

Barger, S. R., Reilly, N. S., Shutova, M. S., Li, Q., Maiuri, P., Heddleston, J. M., Mooseker,

M. S., Flavell, R. A., Svitkina, T., Oakes, P. W. et al. (2019). Membrane-cytoskeletal crosstalk

mediated by myosin-I regulates adhesion turnover during phagocytosis. *Nat Commun* **10**, 1249.

Ben M'Barek, K., Molino, D., Quignard, S., Plamont, M. A., Chen, Y., Chavrier, P. and

Fattaccioli, J. (2015). Phagocytosis of immunoglobulin-coated emulsion droplets. *Biomaterials* **51**, 270-

277.

Bournazos, S., Wang, T. T. and Ravetch, J. V. (2016). The Role and Function of Fcgamma

Receptors on Myeloid Cells. *Microbiol Spectr* **4**.

Cannon, G. J. and Swanson, J. A. (1992). The macrophage capacity for phagocytosis. *J Cell Sci*

101 (Pt 4), 907-13.

Chamaraux, F., Fache, S., Bruckert, F. and Fourcade, B. (2005). Kinetics of cell spreading.

Phys Rev Lett **94**, 158102.

Champion, J. A. and Mitragotri, S. (2006). Role of target geometry in phagocytosis. *Proc Natl*

Acad Sci U S A **103**, 4930-4.

Cougoule, C., Wiedemann, A., Lim, J. and Caron, E. (2004). Phagocytosis, an alternative

model system for the study of cell adhesion. *Semin Cell Dev Biol* **15**, 679-89.

Cuvelier, D., Thery, M., Chu, Y. S., Dufour, S., Thiery, J. P., Bornens, M., Nassoy, P. and

Mahadevan, L. (2007). The universal dynamics of cell spreading. *Curr Biol* **17**, 694-9.

D'Errico, J. (2017). A suite of minimal bounding objects. MATLAB Central File Exchange.

Dewitt, S. and Hallett, M. B. (2002). Cytosolic free Ca(2+) changes and calpain activation are

required for beta integrin-accelerated phagocytosis by human neutrophils. *J Cell Biol* **159**, 181-189.

Dewitt, S. and Hallett, M. B. (2020). Calpain Activation by Ca(2+) and Its Role in

Phagocytosis. *Adv Exp Med Biol* **1246**, 129-151.

DiNapoli, K. T., Robinson, D. N. and Iglesias, P. A. (2021). A mesoscale mechanical model of

cellular interactions. *Biophys J* **120**, 4905-4917.

Dobereiner, H. G., Dubin-Thaler, B., Giannone, G., Xenias, H. S. and Sheetz, M. P. (2004).

Dynamic phase transitions in cell spreading. *Phys Rev Lett* **93**, 108105.

Dubin-Thaler, B. J., Giannone, G., Dobereiner, H. G. and Sheetz, M. P. (2004). Nanometer
analysis of cell spreading on matrix-coated surfaces reveals two distinct cell states and STEPs. *Biophys J*
86, 1794-806.

Dubin-Thaler, B. J., Hofman, J. M., Cai, Y., Xenias, H., Spielman, I., Shneidman, A. V.,
David, L. A., Dobereiner, H. G., Wiggins, C. H. and Sheetz, M. P. (2008). Quantification of cell edge
velocities and traction forces reveals distinct motility modules during cell spreading. *PLoS One* **3**, e3735.

Dustin, M. L. (2016). Complement Receptors in Myeloid Cell Adhesion and Phagocytosis.
Microbiol Spectr **4**.

Etienne, J. and Duperray, A. (2011). Initial dynamics of cell spreading are governed by
dissipation in the actin cortex. *Biophys J* **101**, 611-21.

Fardin, M. A., Rossier, O. M., Rangamani, P., Avigan, P. D., Gauthier, N. C., Vonnegut, W.,
Mathur, A., Hone, J., Iyengar, R. and Sheetz, M. P. (2010). Cell spreading as a hydrodynamic process.
Soft Matter **6**, 4788-4799.

Flannagan, R. S., Jaumouille, V. and Grinstein, S. (2012). The cell biology of phagocytosis.
Annu Rev Pathol **7**, 61-98.

Footer, M. J., Kerssemakers, J. W. J., Theriot, J. A. and Dogterom, M. (2007). Direct
measurement of force generation by actin filament polymerization using an optical trap. *Proceedings of*
the National Academy of Sciences of the United States of America **104**, 2181-2186.

Francis, E. A. and Heinrich, V. (2022). Integrative experimental/computational approach
establishes active cellular protrusion as the primary driving force of phagocytic spreading by immune
cells. *(submitted)*.

Freeman, S. A. and Grinstein, S. (2014). Phagocytosis: receptors, signal integration, and the
cytoskeleton. *Immunol Rev* **262**, 193-215.

Frisch, T. and Thoumine, O. (2002). Predicting the kinetics of cell spreading. *J Biomech* **35**,
1137-41.

Futosi, K., Fodor, S. and Mocsai, A. (2013). Neutrophil cell surface receptors and their
intracellular signal transduction pathways. *Int Immunopharmacol* **17**, 638-50.

Gorelik, R. and Gautreau, A. (2014). Quantitative and unbiased analysis of directional
persistence in cell migration. *Nat Protoc* **9**, 1931-43.

Hallett, M. B. and Dewitt, S. (2007). Ironing out the wrinkles of neutrophil phagocytosis.
Trends Cell Biol **17**, 209-14.

Hategan, A., Sengupta, K., Kahn, S., Sackmann, E. and Discher, D. E. (2004). Topographical
pattern dynamics in passive adhesion of cell membranes. *Biophys J* **87**, 3547-60.

Heinrich, V. (2015). Controlled One-on-One Encounters between Immune Cells and Microbes

Reveal Mechanisms of Phagocytosis. *Biophys J* **109**, 469-76.

Henry, S. J., Crocker, J. C. and Hammer, D. A. (2014). Ligand density elicits a phenotypic

switch in human neutrophils. *Integr Biol (Camb)* **6**, 348-56.

Herant, M., Heinrich, V. and Dembo, M. (2005). Mechanics of neutrophil phagocytosis:

behavior of the cortical tension. *J Cell Sci* **118**, 1789-97.

Herant, M., Heinrich, V. and Dembo, M. (2006). Mechanics of neutrophil phagocytosis:

experiments and quantitative models. *J Cell Sci* **119**, 1903-13.

Herant, M., Lee, C. Y., Dembo, M. and Heinrich, V. (2011). Protrusive push versus

enveloping embrace: computational model of phagocytosis predicts key regulatory role of cytoskeletal

membrane anchors. *PLoS Comput Biol* **7**, e1001068.

Janmey, P. A., Hinz, B. and McCulloch, C. A. (2021). Physics and Physiology of Cell

Spreading in Two and Three Dimensions. *Physiology (Bethesda)* **36**, 382-391.

Jaumouille, V., Cartagena-Rivera, A. X. and Waterman, C. M. (2019). Coupling of beta2

integrins to actin by a mechanosensitive molecular clutch drives complement receptor-mediated

phagocytosis. *Nat Cell Biol* **21**, 1357-1369.

Jaumouille, V. and Waterman, C. M. (2020). Physical Constraints and Forces Involved in

Phagocytosis. *Front Immunol* **11**, 1097.

Kovar, D. R. and Pollard, T. D. (2004). Insertional assembly of actin filament barbed ends in

association with formins produces piconewton forces. *Proc Natl Acad Sci U S A* **101**, 14725-30.

Kovari, D. T., Wei, W., Chang, P., Toro, J. S., Beach, R. F., Chambers, D., Porter, K., Koo,

D. and Curtis, J. E. (2016). Frustrated Phagocytic Spreading of J774A-1 Macrophages Ends in Myosin

II-Dependent Contraction. *Biophys J* **111**, 2698-2710.

Lam, J., Herant, M., Dembo, M. and Heinrich, V. (2009). Baseline mechanical

characterization of J774 macrophages. *Biophys J* **96**, 248-54.

Lee, C. Y., Herant, M. and Heinrich, V. (2011). Target-specific mechanics of phagocytosis:

protrusive neutrophil response to zymosan differs from the uptake of antibody-tagged pathogens. *J Cell*

Sci **124**, 1106-14.

Lee, C. Y., Thompson, G. R., 3rd, Hastey, C. J., Hodge, G. C., Lunetta, J. M., Pappagianis,

D. and Heinrich, V. (2015). Coccidioides Endospores and Spherules Draw Strong Chemotactic,

Adhesive, and Phagocytic Responses by Individual Human Neutrophils. *PLoS One* **10**, e0129522.

Lim, J. J., Grinstein, S. and Roth, Z. (2017). Diversity and Versatility of Phagocytosis: Roles

in Innate Immunity, Tissue Remodeling, and Homeostasis. *Front Cell Infect Microbiol* **7**, 191.

Lubeck, M. D., Steplewski, Z., Baglia, F., Klein, M. H., Dorrington, K. J. and Koprowski, H.
(1985). The interaction of murine IgG subclass proteins with human monocyte Fc receptors. *J Immunol*
135, 1299-304.

Nelsen, E., Hobson, C. M., Kern, M. E., Hsiao, J. P., O'Brien Iii, E. T., Watanabe, T.,
Condon, B. M., Boyce, M., Grinstein, S., Hahn, K. M. et al. (2020). Combined Atomic Force
Microscope and Volumetric Light Sheet System for Correlative Force and Fluorescence Mechanobiology
Studies. *Sci Rep* **10**, 8133.

Odenthal, T., Smeets, B., Van Liedekerke, P., Tijskens, E., Van Oosterwyck, H. and Ramon,
H. (2013). Analysis of initial cell spreading using mechanistic contact formulations for a deformable cell
model. *PLoS Comput Biol* **9**, e1003267.

Ostrowski, P. P., Freeman, S. A., Fairn, G. and Grinstein, S. (2019). Dynamic Podosome-
Like Structures in Nascent Phagosomes Are Coordinated by Phosphoinositides. *Dev Cell* **50**, 397-410 e3.

Pacheco, P., White, D. and Sulchek, T. (2013). Effects of microparticle size and Fc density on
macrophage phagocytosis. *PLoS One* **8**, e60989.

Parekh, S. H., Chaudhuri, O., Theriot, J. A. and Fletcher, D. A. (2005). Loading history
determines the velocity of actin-network growth. *Nat Cell Biol* **7**, 1219-23.

Paul, D., Achouri, S., Yoon, Y. Z., Herre, J., Bryant, C. E. and Cicuta, P. (2013).
Phagocytosis dynamics depends on target shape. *Biophys J* **105**, 1143-50.

Reinhart-King, C. A., Dembo, M. and Hammer, D. A. (2005). The dynamics and mechanics of
endothelial cell spreading. *Biophys J* **89**, 676-89.

Richards, D. M. and Endres, R. G. (2014). The mechanism of phagocytosis: two stages of
engulfment. *Biophys J* **107**, 1542-53.

Roberts, R. E., Dewitt, S. and Hallett, M. B. (2020a). Membrane Tension and the Role of Ezrin
During Phagocytosis. *Adv Exp Med Biol* **1246**, 83-102.

Roberts, R. E. and Hallett, M. B. (2019). Neutrophil Cell Shape Change: Mechanism and
Signalling during Cell Spreading and Phagocytosis. *Int J Mol Sci* **20**.

Roberts, R. E., Martin, M., Marion, S., Elumalai, G. L., Lewis, K. and Hallett, M. B.
(2020b). Ca(2+)-activated cleavage of ezrin visualised dynamically in living myeloid cells during cell
surface area expansion. *J Cell Sci* **133**.

Rosales, C. and Uribe-Querol, E. (2017). Phagocytosis: A Fundamental Process in Immunity.
Biomed Res Int **2017**, 9042851.

Shashidharamurthy, R. B., E.; Patel, J.; Kaur, R.; Meganathan, J.; Selvaraj, P. (2010).
Analysis of cross-species IgG binding to human and mouse Fcgamma receptors (FcγRs). *The Journal of*
Immunology **184**.

Simon, S. I. and Schmid-Schonbein, G. W. (1988). Biophysical aspects of microsphere

engulfment by human neutrophils. *Biophys J* **53**, 163-73.

Tran-Son-Tay, R., Needham, D., Yeung, A. and Hochmuth, R. M. (1991). Time-dependent

recovery of passive neutrophils after large deformation. *Biophys J* **60**, 856-66.

Uribe-Querol, E. and Rosales, C. (2020). Phagocytosis: Our Current Understanding of a

Universal Biological Process. *Front Immunol* **11**, 1066.

van Zon, J. S., Tzircotis, G., Caron, E. and Howard, M. (2009). A mechanical bottleneck

explains the variation in cup growth during FcgammaR phagocytosis. *Mol Syst Biol* **5**, 298.

Vorselen, D., Barger, S. R., Wang, Y., Cai, W., Theriot, J. A., Gauthier, N. C. and Krendel,

M. (2021). Phagocytic 'teeth' and myosin-II 'jaw' power target constriction during phagocytosis. *Elife* **10**.

Vorselen, D., Wang, Y., de Jesus, M. M., Shah, P. K., Footer, M. J., Huse, M., Cai, W. and

Theriot, J. A. (2020). Microparticle traction force microscopy reveals subcellular force exertion patterns

in immune cell-target interactions. *Nat Commun* **11**, 20.

Warmerdam, P. A., van de Winkel, J. G., Gosselin, E. J. and Capel, P. J. (1990). Molecular

basis for a polymorphism of human Fc gamma receptor II (CD32). *J Exp Med* **172**, 19-25.

Warrier, R. (2017). max_inscribed_circle. MATLAB Central File Exchange.

Wolfenson, H., Iskratsch, T. and Sheetz, M. P. (2014). Early events in cell spreading as a

model for quantitative analysis of biomechanical events. *Biophys J* **107**, 2508-14.

Xiong, Y., Rangamani, P., Fardin, M. A., Lipshtat, A., Dubin-Thaler, B., Rossier, O.,

Sheetz, M. P. and Iyengar, R. (2010). Mechanisms controlling cell size and shape during isotropic cell

spreading. *Biophys J* **98**, 2136-46.

Yeung, A. and Evans, E. (1989). Cortical shell-liquid core model for passive flow of liquid-like

spherical cells into micropipets. *Biophys J* **56**, 139-49.

Zhang, Y., Hoppe, A. D. and Swanson, J. A. (2010). Coordination of Fc receptor signaling

regulates cellular commitment to phagocytosis. *Proc Natl Acad Sci U S A* **107**, 19332-7.

 Figure legends

Fig. 1. Preparation of substrates with controlled rabbit-IgG surface density. (**A**) BSA-
coated glass coverslips that had been incubated with various ratios of rabbit and mouse anti-BSA
antibodies were labeled with green-fluorescent secondary antibodies and then imaged on a
confocal microscope. (**B**) Standard beads with known numbers of antibody-binding sites were
saturated with the same fluorescent secondary antibody as used in (A). (**C**) The fluorescence
intensity at the underside of the coated beads was measured with the same confocal-microscope
settings as used in (A). The results, along with a straight-line fit (*red line*), are presented as a
function of the known density of antibody-binding sites in a log-log plot. Error bars denote the
geometric standard deviation. Examples of confocal bead images are included. (**D**) The
calibration curve of (C) was used to convert the measured fluorescence intensities of labeled
coverslips into the rabbit-IgG density. The geometric mean values (with error bars denoting
geometric standard deviation) of the IgG density are plotted as a function of the concentration of
rabbit IgG used in the incubation buffer. The result of a linear fit (*dashed line*) to the portion of
the data obtained at IgG concentrations of 1% or higher is included.

Fig. 2. Illustration of a frustrated phagocytic spreading experiment. (**A**) Parts and
assembly of the experiment chamber. (**B**) After settling onto the functionalized coverslip,
initially passive, round cells recognize the deposited antibodies and spread along the surface.
(**C**) Reflection interference contrast microscopy allows us to visualize the cell-substrate contact
area label-free at high resolution. The combination of polarizing beamsplitter and quarter-wave
plate (QWP) ensures that the light components reflected off the coverslip top surface and the cell
are the dominant contributions to the recorded interference pattern. (**D**) Example snapshots
from a series of recorded video images of the dark cell-substrate contact area (see also Movie
S1). Timestamps are included. The scale bar denotes 10 μm.

Fig. 3. Dependence of the spreading probability on the surface density of IgG. (**A**) At low
surface densities of rabbit IgG, the majority of deposited cells do not spread and appear as bright,
out-of-focus spots in our RICM images (*top panel*). In contrast, almost all cells spread on
surfaces coated with the highest density of rabbit IgG (*bottom panel*). Scale bars denote 50 μm.
(**B**) The spreading probability depends strongly on the nominal concentration of rabbit IgG,

confirming that the observed cell response is IgG-specific. Spearman's rank correlation

coefficient ρ is reported with the associated p-value for the null hypothesis $\rho = 0$.

Fig. 4. Overview of the human neutrophil response to IgG-coated surfaces. (**A**) The first

four snapshots of each video sequence illustrate the common morphology of the cell footprint

observed during the outward-spreading phase of almost all cells (see also Movie S1). In

contrast, three qualitatively distinct types of post-spreading cell behavior were observed, as

illustrated in the last two snapshots of each video sequence. Time stamps are included. Scale

bars denote 10 μm. (**B**) Contours of the cell footprint of the first four images of the respective

video sequences of (A) demonstrate the roughly symmetric cell morphology during spreading.

Scale bars denote 10 μm. (**C**) The table reports the fractions of cells that exhibit one of the three

types of post-spreading behavior for each of the tested IgG concentrations. On low densities of

IgG, the majority of cells attempted to migrate after reaching a maximum cell-substrate contact

area. In contrast, on high IgG densities the majority of cells remained in place and exhibited

little further shape changes.

Fig. 5. Quantitative analysis of the cell-substrate contact area of spreading cells. (**A**) The

contact area is determined from a polygonal trace of the circumference of each cell footprint

(*yellow polygons* in the examples included at the top; scale bar denotes 10 μm). A suitable fit to

the time-dependent contact area yields the (maximum) spreading speed. The plateau value of the

contact-area curves provides the maximum contact area. (**B**) Representative examples of

contact-area-versus-time curves measured on the lowest (*red*) and highest (*green*) IgG densities

illustrate the variability of the dynamical behavior of the cell footprint. (**C**) Average curves of

all suitable area-versus-time measurements obtained for each tested rabbit-IgG density expose a

largely conserved spreading speed. On the other hand, the maximum contact area appears to

reach greater values on high IgG densities. Error bars denote standard errors. (**D**) The summary

of all spreading speeds measured over three orders of magnitude of the IgG concentration

confirms that the spreading speed is independent of the surface density of IgG. (**E**) The

summary of maximum contact areas reveals a small but significant increase of the contact area at

high densities of IgG. Error bars in (D) and (E) denote standard deviation.

Fig. 6. Roundness and type of motion of the footprints of spreading cells. (**A**) The

roundness of the cell-substrate contact region was assessed in terms of the ratio between the radii

of the largest inscribed (*red*) and the smallest circumscribed (*blue*) circles, as illustrated with two

examples. (**B**) The summary of all roundness measurements reveals no significant differences

on surfaces coated with different IgG densities. (**C**) Illustration of our measurement of the

displacement Δr of the centroid of the cell footprint during the time interval Δt. (**D**) The

summary of all centroid-speed measurements reveals no significant differences on surfaces

coated with different IgG densities. (**E**) The mean square displacements of the centroid as a

function of Δt (averaged over all suitable measurements obtained for each tested rabbit-IgG

density; error bars denote standard errors) allow us to quantify the degree of randomness of the

centroid motion. The included solid lines show power-law fits of the form $y°=°ax^b$ to the data.

(**F**) The summary of all measured directional persistence times reveals that on the highest

density of IgG, the cell footprint tended to move in roughly the same direction for significantly

shorter times than on lower IgG densities. For more details see the text.

Fig. 7. Summary of pertinent predictions by two different mechanistic models of cell

spreading. (**A**) The models incorporate attractive interactions between the cell surface and the

substrate via short-range adhesion stress (*left panel*). The protrusive-zipper model postulates that

the cell interior generates protrusion stress local to the leading edge of the spreading cell in a

manner that depends on fresh contact between the cell and substrate (*right panel*). (**B**) A

snapshot of a computer simulation of a spreading model cell illustrates the motion of the viscous

cell interior according to the Stokes equations for creeping flow. (**C**) Time series of cross-

sectional shapes of spreading model cells illustrate morphological differences between the

predictions of the two considered models. (**D**) The Brownian-zipper model predicts that both

the cell-spreading speed as well as the maximum contact area exhibit a strong dependence on the

surface density of IgG coating the substrate. The final shapes are predicted to be spherical caps

in this case. These predictions fail to reproduce experimental observations such as presented in

Fig. 5. (**E**) In agreement with our experimental results (Fig. 5C,D), the spreading speeds

predicted by the protrusive-zipper model are essentially independent of the IgG density, and the

IgG dependence of the maximum contact area is much weaker than in (D).

Figure 1

Figure 2

Figure 3

Figure 4

A Monotonic symmetric spreading   Attempted migration   Contraction after spreading

E x a m p l e s

B Contours (during spreading only)

C

Fraction of cells exhibiting one of the observed behaviors [%]

% IgG	Monotonic symmetric spreading	Attempted migration	Contraction after spreading	N
0.1	5.9	76.5	17.6	17
1	0.0	100.0	0.0	7
10	24.0	68.0	8.0	25
100	66.7	33.3	0.0	21

Figure 5

Figure 6

Figure 7