Urocortin-3 neurons in the perifornical area mediate the impact of chronic stress on female infant-directed behavior

Brenda Abdelmesih[1], Robyn Anderson[1], Ilaria Carta[1], Anita E. Autry[1,2*]

1.   Dominick P. Purpura Department of Neuroscience, Albert Einstein College of Medicine, Bronx, NY
2.   Department of Psychiatry and Behavioral Sciences, Albert Einstein College of Medicine, Bronx, NY

*Correspondence should be addressed to Anita E. Autry, anita.autry@einsteinmed.edu

Abbreviated title: Ucn3 neurons mediate impact of stress on female parenting

Abstract

Infant avoidance and aggression are promoted by activation of the Urocortin-3 expressing neurons of the perifornical area of hypothalamus (PeFA^{Ucn3}) in male and female mice. PeFA^{Ucn3} neurons have been implicated in stress, and stress is known to reduce maternal behavior. We asked how chronic restraint stress (CRS) affects infant-directed behavior in virgin and lactating females and what role PeFA^{Ucn3} neurons play in this process. Here we show that infant-directed behavior increases activity in the PeFA^{Ucn3} neurons in virgin and lactating females. Chemogenetic inhibition of PeFA^{Ucn3} neurons facilitates pup retrieval in virgin females. CRS reduces pup retrieval in virgin females and increases activity of PeFA^{Ucn3} neurons but does not affect maternal behavior in mothers. Inhibition of PeFA^{Ucn3} neurons blocks stress-induced deficits in pup-directed behavior in virgin females. Together, these data illustrate the critical role for PeFA^{Ucn3} neuronal activity in mediating the impact of chronic stress on female infant-directed behavior.

Significance statement

While a large body of research has studied the impact of maternal stress on offspring, few studies have focused on the neural circuitry underlying reduced maternal behavior in stressed mothers. In this study, we examine the neural substrates involved in reduced infant-directed behavior caused by chronic stress. We find that perifornical area neurons expressing the neuropeptide urocortin-3 are critical mediators of the impact of stress on infant-directed behavior in females.

Introduction

Many decades of research have focused on the neurobiology of maternal behavior, revealing common mechanisms and pathways involved in infant caregiving behavior across a variety of species (Numan & Insel, 2003; Numan, 2020). Studies have converged on the critical role of the medial preoptic area of hypothalamus in orchestrating the behavioral responses of mothers to their young in frogs, fish, birds, and rodents (Numan, 1974; Slawski & Buntin, 1995; Fischer *et al.*, 2019; Maruska *et al.*, 2020). Recently, it has been appreciated that these mechanisms may also be involved in paternal behaviors as well, suggesting a core circuitry that exists in both sexes to promote caregiving (O'Connell *et al.*, 2012; Wu *et al.*, 2014; Kohl *et al.*, 2018). Moreover, it has been well-documented that neural plasticity mechanisms underlie the facilitation of infant care behavior, including alloparental care towards unrelated young, particularly in females (Numan & Insel, 2003).

In the absence of caregiving behavior, it is possible to observe neglect or even aggression toward infants by adults. Studies have identified circuit nodes in the brain, including the medial and posterior amygdala, the bed nucleus of stria terminalis, and the perifornical area of hypothalamus, that modulate expression of infant-directed neglect and aggression (Tsuneoka *et al.*, 2015; Chen *et al.*, 2019; Sato *et al.*, 2020; Autry *et al.*, 2021). We wondered if this anti-parental circuitry may be active in neglectful animals including virgin females or stressed virgin and lactating females.

Clinical research clearly shows that stress is a critical risk factor for postpartum mental illnesses including postpartum depression or anxiety which affect up to 25% of women and 10% of men annually in the United States (Paulson & Bazemore, 2010; Wisner *et al.*, 2013). However, there are few preclinical studies that examine the neurobiology underlying reduced parent-infant bonding or associated symptoms in animal models (Nephew & Bridges, 2011; Zoubovsky *et al.*, 2020; Rosinger *et al.*, 2021). Extant research has focused on the impact of maternal stress as a model of early life stress either pre- or postnatally on behavior outcomes in

offspring, often with profound behavioral, physiological, and neurobiological impacts on young

raised by stressed mothers (Cameron *et al.*, 2005; Wang *et al.*, 2011; Singh-Taylor *et al.*, 2015;

Delpech *et al.*, 2016; Feifel *et al.*, 2017; Kronman *et al.*, 2021; Rincon-Cortes & Grace, 2021).

During lactation, females are hyporesponsive to acute stress due to hormonal changes that

impact the Hypothalamic-Pituitary-Adrenal (HPA) axis regulation (Walker *et al.*, 2001; Brunton *et

al.*, 2008). HPA axis hypo-responsivity is thought to be protective of anxiety-related behavior

and adult-pup interactions in lactating females (Miller *et al.*, 2011; Medina *et al.*, 2021).

However, chronic stress has been documented to have a long-lasting impact on the regulation

of the HPA axis, leading to reduced parenting, and how the underlying neurobiology is affected

remains poorly understood (Carini *et al.*, 2013; Murgatroyd & Nephew, 2013; Murgatroyd *et al.*,

2015).

Ucn3 is a member of the corticotropin releasing factor (CRF) family of stress hormones

and has the highest endogenous binding affinity for CRF receptor 2 (CRFR2). Previous studies

of this group of neurons suggest that they are sensitive to stress and adrenalectomy (loss of

stress hormones) (Jamieson *et al.*, 2006). Overexpression of Ucn3 in the brain leads to

increased anxiety- and depression-related behaviors and results in a blunted HPA response to

stress (Neufeld-Cohen *et al.*, 2012). Furthermore, overexpression of Ucn3 specifically in the

PeFA is associated with enhanced anxiety-like behaviors in mice (Kuperman *et al.*, 2010).

Social discrimination abilities are altered in a sex-specific manner in total Ucn3 knockout mice

(Deussing *et al.*, 2010). Taken together, these studies suggest that PeFA Ucn3 cells mediate

stress-induced behavioral changes.

Thus, we hypothesized that chronic stress would negatively affect infant-directed

behavior in females and that this disruption is dependent on activation of perifornical area

urocortin-3 expressing neurons. We set out to determine if anti-parental circuit components,

specifically the urocortin-3 positive neurons in the perifornical area of hypothalamus were more

active in naïve or stressed females, and if we could recover parental behavior by blocking

activation of this anti-parental circuit node. We find that increased parental behavior is

accompanied by decreased activity in perifornical area urocortin-3 expressing neurons and

blocking activity in these cells enhances parental behavior in naïve females. Chronic stress

reduces alloparental behavior in naïve females and this stress-induced behavioral effect is

occluded by inhibition of perifornical area urocortin-3 cells. In stressed lactating females,

parental behavior is preserved and perifornical area urocortin-3 cells are less activated in stress.

Together, these data reveal a critical role for perifornical urocortin-3 neurons in the expression

of alloparental behavior in female mice under both normal and pathological conditions.

Materials and Methods

Animals

Mice were maintained on a 12h:12h dark light cycle (10:30am-10:30 pm dark phase)

with access to food and water ad libitum. All experiments were performed in accordance with

NIH guidelines and approved by the Albert Einstein College of Medicine Institutional Animal

Care and Use Committee (IACUC; protocol 20180110; 20180111; 00001386).

C57BL/6J sexually naïve female and pregnant female (E14) mice were ordered from

Jackson Laboratories (Bar Harbor, ME) aged at 6-8 weeks. Ucn3::Cre BAC transgenic line

(STOCK Tg(Ucn3-cre) KF43Gsat/Mmcd 032078-UCD; obtained from laboratory of Catherine

Dulac, Harvard University) were genotyped at weaning (3 weeks of age) and used in

experiments at age 2-5 months. Animals received from Jackson Laboratories habituated to our

facility for 7 days prior to behavioral testing.

Corticosterone Measure

Trunk blood samples were taken at the time of sacrifice and blood serum was isolated

from blood samples by centrifugation. A high-sensitivity corticosterone (CORT) enzyme

immunoassay (EIA) was used and analyzed according to manufacturer's instructions

(Immunodiagnostic Systems Ltd, Fountain Hills, AZ, USA) as previously described (Autry *et al.*,

2009). Briefly, percent binding (B/Bo%) of each calibrator, control and sample was calculated by

dividing the mean absorbance over the mean absorbance for '0' calibrator and multiplied by

100. A calibration curve was used to plot B/Bo% on the ordinate against concentration of

corticosterone. A 4pl curve fit was applied.

<u>Chronic restraint stress model</u>

Virgin and lactating female mice were used. Lactating females were restrained starting

from approximately postpartum (PP) day 2. Both stressed and unstressed mice were brought to

a test room under dim red light during their dark cycle. All animals were weighed, and females

were either placed back into their home cage or placed into a 50 mL conical tube for one hour.

Humidity and temperature in the test room was recorded each day. On the last day of stress,

females remained in the test room for 1-2 hours before being exposed to a foreign-born pup

(see Parental Behavior).

Animals injected with AAV1/DIO-hM4Di (see chemogenetics) recovered from surgery at

least 1 week before the start of restraint. For the stressed virgin female group, mice were

excluded from the stressed group (n=4 females) based on open field test behavior that was

indistinguishable from control. For the groups that received AAV1/DIO-hM4Di injections,

females were excluded if they did not show adequate recombination of the DREADD construct

(n=2, virgin females Figure 2; n=0 stressed virgin females Figure 6).

<u>Intruder stress model</u>

Lactating females were stressed starting from approximately postpartum (PP) day 2.

Both stressed and unstressed mice were brought to a test room under dim red light during their

dark cycle. All animals were weighed, and females were placed back into their home cage and

either placed back on the housing rack or had an intact adult male intruder (C57, ~2 months old)

introduced into their home cage for 10 minutes as described in previous studies (Carini *et al.*,

2013; Murgatroyd *et al.*, 2016). In the event the male was too aggressive toward pups, the

intruder stress period was curtailed. Humidity and temperature in the test room was recorded

each day. On the last day of stress, females remained in the test room for 1-2 hours before

being exposed to a foreign-born pup (see Parental Behavior).

Behavior assays

Mice were individually housed for at least 1 week prior to testing. Experiments were

conducted during the dark phase under dim red light. Tests were recorded by Fly Capture

cameras (Point Grey, Richmond, BC, Canada) and behaviors were scored by an observer blind

to experimental condition using Observer XT13 Software or Ethovision XT 13 (Noldus

Information Technology, Leesburg, VA, USA). Animals were tested for a single behavior per

session with at least 24 hours between sessions.

Parental behavior

Parental behavior tests were conducted in the mouse's home cage as previously

described (Wu *et al.*, 2014). Mice were habituated to the testing environment for 10 minutes.

One to two C57BL6/J pups 1-4 days old were presented in the cage in the opposite corner to

the nest. Test sessions started either at pup introduction or pup approach (female first touches

the pup with its snout) and lasted for 10- 15 minutes. If the mouse became aggressive by biting

and wounding the pup, the session was immediately halted, and the pup was euthanized. The

following behaviors were quantified: latency to retrieve, pup investigation (sniffing, close contact

with snout), grooming (handling with forepaws and licking), nest building, time spent in the nest,

crouching, latency to attack (latency to bite and wound), aggression (roughly handling,

aggressively grooming, aggressive carrying with no retrieval), and tail rattling. A 'parenting

behavior' index was calculated as the sum of duration of grooming, nest building, time spent in

the nest, and crouching.

Open field

Mice were assessed for activity in a 45cm x 45 cm open field at 40 lux for 5 min as

previously described (Autry *et al.*, 2009). Center was considered 15 cm x 15 cm and borders

were 5 cm around the perimeter of the box. Time and frequency in center and borders as well

as distance and velocity were calculated using Ethovision XT13.

Behavioral ethograms were made in Matlab using custom code.

Fluorescence in situ hybridization

Fluorescence in situ hybridization (FISH) was performed as recommended by ACD Bio

(Newark, CA, USA) using V1 RNAscope reagents. Briefly, fresh brain tissue was collected from

animals housed in their home cages or 35 min after the start of the behavior tests for immediate

early gene (*Fos)* studies. Brains were embedded in OCT (Tissue-Tek) and frozen with dry ice.

25µm cryosections were used for mRNA in situ. Adjacent sections from each brain

were collected over replicate slides to stain with multiple probes. Protease 3 was used to digest

tissue. Fos (Cat No. 316921), Ucn3 (Cat No. 464861), and Crh (Cat No. 316091) probes were

used as per manufacturer's instructions. Slides were mounted using Prolong Gold with DAPI.

Zeiss Axioscan was used to image DAPI, Alexa 488, Atto-550, and Alexa 647 at 20X

magnification.

Immunostaining and histology

To visualize c-Fos protein in combination with AAV-hM4Di, perfused tissue was sliced

on a freezing microtome at 30 µm, and every third section throughout the PeFA was stained.

Sections were rinsed with 0.1% PBS with Triton (PBST), blocked with 5% donkey serum diluted

in PBST (blocking solution) for 1 hour at room temperature. Primary antibody chicken anti-

mCherry (Millipore AB3566481) and rabbit anti c-Fos (Cell Signaling 2250S) were diluted at

1:1000 in blocking solution and sections were incubated overnight at 4°C. After rinsing with

PBST, secondary anti-chicken-A594 (Sigma CF594) and anti-rabbit-A647 (Life Technologies

A31573) were applied at 1:200 and 1:1000 dilutions, respectively, in blocking solution and

incubated overnight at 4°C. Sections were rinsed in PBS, mounted to Superfrost Plus slides,

coverslipped with Prolong Gold containing DAPI, and imaged on the Zeiss Axioscan as

described previously.

 Chemogenetics

Ucn3::Cre virgin female mice (or Cre negative littermates as controls) 8-20 weeks old were

used for these experiments. We stereotaxically injected ~225 nL of conditional inhibitory

designer receptor exclusively activated by designer drug (DREADD) virus bilaterally into the

PeFA (AP -0.6mm, ±ML 0.3mm, DV -4.2mm). For the naïve virgin female DREADD

experiment, we used a custom prep from UNC Vector Core (AAV1-hSyn-DIO-HM4D(Gi)-

mCherry; Chapel Hill, North Carolina, USA) and for the stressed virgin female DREADD

experiment we used a custom prep from Vector Builder (AAV1-hSyn-FLEX-HM4D(Gi)-mCherry;

Chicago, Illinois, USA). Animals used for stress study recovered from surgery for 1 week before

the start of restraint and around two-three weeks before behavioral testing.

Cre-positive and Cre-negative females were administered intraperitoneally (i.p) with

either 1x PBS (vehicle) or 0.3mg/kg clozapine-n-oxide (CNO) dissolved in 1x PBS and

habituated to the testing environment for two-three hours prior to pup assay. Females were

presented one to two C57BL6/J pups in the corner of their home-cage opposite the nest and

parental behaviors were recorded for 10-15 minutes.

Four control animals used in the stress study were administered either CNO or saline i.p

and 2 hours later were exposed to a pup. Animals were then perfused 90 minutes later to stain

for c-Fos protein expression. (See Immunostaining and histology)

Data analysis and Statistics

Data was analyzed by Graphpad Prism 9.0 or Matlab scripts. For colocalization

experiments, we used Fisher's exact test to compare the total number of fos+/marker+ positive

cells to the total number of fos-/marker+ positive cell populations across all mice and expressed

the data as percentages from each individual mouse. Pup retrieval percentages are analyzed

by Kolmogorov-Smirnoff (2 groups) or Friedman test (3 groups). For experiments comparing

one manipulation to control (i.e., stress or neuronal inactivation), we used t-test for parameters

with normally distributed data and Mann-Whitney test for non-normally distributed data. To

compare one manipulation and control across several sessions, we used one-way repeated

measures ANOVA tests followed by post-hoc correction. In experiments with comparison of two

manipulations in several sessions (stress and neuronal inactivation), we used two-way repeated

measures ANOVA followed by post-hoc correction. P values reported as follows: <0.05 *, **

P<0.01, *** P<0.001, **** P<0.0001. All data are expressed as mean ± SEM.

<u>Image analysis</u>

Images were exported from Zen Blue software and cells were manually counted for

colocalization using FIJI Cell Counter. Graphpad Prism 9 was used to plot graphs and perform

statistics.

<u>Fiber Photometry</u>

Ucn3-cre animals were injected with a cocktail of 150 nL of AAV-syn-jGCaMP7f-WPRE

(Addgene 104488-AAV9) and 225 nL of AAV-hSyn-DIO-hM4D(Gi)-mCherry into PeFA (ML:

0.3mm; AP -0.6mm; DV 4.2mm). In the same surgery, a 200μM fiber optic cannula was

implanted (ML: 0.3mm; AP -0.6mm; DV 4.2mm). Animals recovered for at least 3 weeks before

behavioral experiments. Animals were brought up to the test room in dim red light and injected

with either vehicle (session 1) or 0.3mg/kg CNO (session 2) i.p. Two hours later, a fiber optic

patch cable (Doric) was attached to the cannula and adjusted to attachment for 10 minutes

before recording. Using a multi-channel fiber photometry system (Neurophotometrics LTD), a

470 nm LED and 415 nm LED (isosbestic control) alternatively illuminated at 60μW via a 20X

objective and fluorescence emission was collected using a CMOS camera sensor. After 1-2

minutes of recording, animals underwent 6 tail suspensions for approximately 5 seconds per

suspension. Data were acquired using the open-source software Bonsai.

Photometry data was analyzed using custom MATLAB code. To correct for

photobleaching and motion artifact, we used normalization similarly described by Hvratin et al

2020 (Hrvatin *et al.*, 2020). In short, the isosbestic signal was fit with a biexponential and then

linearly scaled to fit signal emitted by GCaMP. GCaMP signal was then divided by the scaled fit

for ΔF/F. Tail suspension events were aligned to normalized photometric signal and peri-events

were taken from 5 sec before tail suspension ("pre") to 5 sec after ("post"). The pre-event

baseline was used to calculate the z score $\frac{\chi_i - \chi_{baseline}}{s_{baseline}}$. The mean ΔF/F of each pre- and post- tail

suspension event was taken and averaged across animals per group (vehicle vs CNO) and

compared using a paired t test. Area under curve was calculated with the mean ΔF/F of each

pre- and post- tail suspension event using the MATLAB built-in function "trapz". Standard error

mean is plotted with the average z-score.

<u>Code availability</u>

Custom Matlab code for ethogram generation and analysis of photometry data is

available upon request.

Results

To identify the activation levels of Urocortin-3 in the rostral perifornical area of the

hypothalamus, we exposed C57 virgin females as well as lactating females (postpartum day 2)

to either a foreign pup (P0-P4) or ~25 mg of fresh bedding (control) in their home cage. Animals

were subsequently sacrificed 30 minutes after exposure (Figure 1A). To control for number of

pups as well and foreign pup discrimination (Ostermeyer & Elwood, 1983; Mogi *et al.*, 2017), we

utilized two groups of lactating females that either had their litter removed 10 minutes prior to

foreign pup introduction or kept their litter (Figure 1A). Visualization of immediate early gene Fos

and Ucn3 in PeFA revealed increases in PeFA[Ucn3] cell activation in virgin females exposed to a

pup compared to controls (Figure 1B, C; Supplemental Figure 1-1). However, in lactating

females, Fos levels in PeFA[Ucn3] decrease with pup exposure if litter has been removed but

increase if litter is present. These results suggest that PeFA[Ucn3] cells respond to pup exposure

similarly in virgin females and lactating females that keep their litter, while we observe opposite

impact on PeFA[Ucn3] cell activity in mothers when her litter is removed.

Because virgin females are not as parental as lactating females (Lonstein & De Vries,

2000; Kuroda *et al.*, 2011; Marlin *et al.*, 2015; Carcea *et al.*, 2021) and PeFA[Ucn3] neurons are

activated by infanticide in females (Autry *et al.*, 2021), we wanted to test if suppression of

PeFA[Ucn3] neuronal activity could enhance alloparental behavior in virgin females. To

accomplish this, we used a conditional viral strategy to express the inhibitory designer receptor

exclusively activated by designer drug (DREADD hM4Di) in Ucn3::Cre positive and Cre

negative animals in the PeFA of virgin females naïve to pups. Two-three weeks after viral

injection, both groups of animals were administered CNO (0.3 mg/kg intraperitoneally) and 2-3

hours later, exposed to two pups for fifteen minutes in their home cage (Figure 2A;

Supplemental Figure 2-1). We confirmed viral recombination to include females in subsequent

behavioral analyses (Figure 2B). Cre+ females retrieved more pups in a shorter amount of time

relative to Cre- females (Figure 2C & D). However, there was no difference in latency to retrieve

the 2[nd] pup (Figure 2F). Furthermore, Cre+ animals spent more time in the nest with pups and

started nest-building earlier compared to Cre- females (Figure 2K &N). While other behaviors

were not improved (Figure 2 G-J, L, M,O-P), suppression of PeFA[Ucn3] neurons improved certain

aspects of alloparenting behavior (Figure 2Q), particularly pup retrieval and time spent in the

nest with the pups.

To understand the impact of chronic stress on alloparental behavior in virgin females, we

employed a chronic restraint stress paradigm in which females were placed into a 50 mL Falcon

conical tube for 1 hour a day for 20 days (Figure 3A). Stressed females weighed significantly

less than control females (Figure 3B). On day 19, females were tested for exploratory behavior

in a 5-minute open field task (Figure 3C-F). Stressed females spent less time in the center of

the field compared to control females (Figure 3C). On the last day of stress, day 20, females

were exposed to a newborn pup (P0-4) for 15 minutes and their alloparental behavior was

recorded and analyzed (Figure 3G-R). Only 2 out of 9, or 22%, of stressed females retrieved

pups compared to control females (5 of 9, or 55% retrieved) (Figure 3G). Other than retrieval

latency, virgin stressed females did not show any significant changes in other measures of

alloparental behavior. Altogether, we find that chronic restraint stress significantly reduces pup

retrieval in virgin females.

Next, to understand the impact of chronic stress on maternal behavior in lactating

females, we utilized the same chronic restraint paradigm in females from postpartum day 2-18

(Figure 4A), before weaning age for pups. Like stressed virgin females, stressed lactating

females weighed significantly less than control females (Figure 4B), indicating that chronic

restraint induced physiological changes. On day 16 of chronic restraint, females were tested for

anxiety-related behavior in a 5-minute open field task (Figure 4C-F). Surprisingly, stressed

females spent more time in the center of the field and less time in the borders (Figure 4C, D).

Stressed females also showed an increase in velocity and distance traveled relative to control

females (Figure 4E, F). On the last day of stress, day 17, females had their litters removed and

10 minutes later we introduced a foreign-born pup to their home cage (Figure 4G-R). All females

retrieved pups before the end of the 10 minutes session (Figure 4G, H) and there was no

difference in latency to retrieve between groups. Stressed mothers showed similar levels of

parenting toward pups as control mothers. We also attempted to use an intruder stress

paradigm that has previously been reported to impact parental behavior in lactating females

(Carini *et al.*, 2013; Murgatroyd *et al.*, 2016). We did not see any weight changes or parenting

measures (Supplemental Figure 4-1).

Next, we investigated molecular and physiological impacts of chronic restraint stress in

virgin or lactating females.   In situ hybridization revealed increases in PeFA^{Ucn3}/Fos

colocalization in stressed virgin females (Figure 5A, B; Supplemental Figure 5-1). In control

virgin females, percentage of PeFA^{Ucn3}/Fos colocalization was negatively correlated with time

spent parenting, indicating that activation of PeFA^{Ucn3} may reduce alloparental behaviors (Figure

5E). Because activation of corticotropin releasing factor (CRF) cells in the paraventricular

hypothalamus (PVH) is postulated to disrupt maternal behavior and is critical for physiological

stress responses (Herman & Tasker, 2016; Klampfl & Bosch, 2019), we also quantified

PVH^{CRF}/Fos colocalization (Figure 5C,F). We found that PVH^{CRH}/Fos levels were significantly

reduced in both stressed virgin females and lactating females, and like PeFA^{Ucn3}, PVH^{CRF}

neuronal activation is negatively correlated with parental behaviors (Figure 5F). Chronic

restraint stress did not affect circulating CORT levels in virgin females (Figure 5D). In mothers,

chronic stress led to a decrease in PeFA^{Ucn3}/Fos levels compared to control lactating females,

opposite to the effect we observed in virgin females (Figure 5G, H). Like virgin females,

however, PVH^{CRF} cell activation was significantly decreased in chronically stressed lactating

females (Figure 5G, I; Supplemental Figure 5-1), consistent with previous literature (Girotti *et*

al., 2006; Radley & Sawchenko, 2015; Matovic *et al.*, 2020). We observed a similar negative

trend for correlation of PeFA^{Ucn3} neuronal activation and parental behaviors in control lactating

females that we observed in virgin females, but the trend for PVH^{CRF} cell activation is positively

correlated in lactating females (Figure 5 K, L). We observed that CORT levels were significantly

decreased in stressed lactating females (Figure 5J), suggesting adaptive habituation to the

repeated stress. In our intruder stress experiment, we did not observe molecular changes in

PeFA^{Ucn3} activation or in circulating corticosterone levels, consistent with no changes in weight

or parental behavior (Supplemental Figure 5-2). Altogether, chronic restraint stress induces

differential PeFA^{Ucn3} activation patterns in virgin and lactating females in response to pups,

while chronic stress reduces PVH^{CRF} neuronal activation in both virgins and mothers.

Because our chronic restraint stress paradigm dampened alloparental behavior in virgin

females and increased PeFA^{Ucn3} neuronal activation, we sought to ameliorate deficits in

parenting by inhibiting PeFA^{Ucn3} neurons during pup exposure. To accomplish this, we injected

virgin Ucn3::Cre positive and negative females with AAV1-eF1a-DIO-hM4Di-mCherry and then

started restraint stress 1 week after recovery from surgery. After 16 days of chronic restraint

stress, we injected either vehicle or CNO on 2 consecutive days after the last day of stress.

After several days we then employed randomized CNO/Vehicle open field trials. On day 32, we

performed an additional pup exposure with vehicle treatment (Figure 6A). Stressed females had

a significant difference in weight compared to control females (Figure 6B). During open field,

CNO administration did not induce changes in exploratory behaviors in either group (Figure 6C-

F). In the pup exposure assay, both stressed and control virgin females had improved

cumulative retrieval with CNO treatment, which we did not observe in the Cre negative group

(Figure 6 G, H; Supplemental Figure 6-1). Strikingly, stressed females treated with CNO

displayed improved latency to retrieve, time spent crouching, time spent in nest and overall time

spent parenting which did not occur in unstressed controls or Cre negative controls, or in the

final vehicle session (Figure 6I, L, M, N; Supplemental Figure 6-1). No significant changes were

observed in pup grooming (Figure 6K). Interestingly, pup investigation was significantly

decreased in both groups which may be due to increased familiarity (Figure 6J) (Bielsky *et al.*,

2005; Richter *et al.*, 2005; Moy *et al.*, 2008). CNO administration did not improve any of the

parenting measures in stressed Cre negative females (Supplemental Figure 6-1).   We

confirmed that CNO administration reduced activity in the PeFA^{Ucn3} neurons using both

histology and fiber photometry recording (Supplemental Figure 6-2). Altogether, inhibition of

PeFA^{Ucn3} neuronal activity leads to enhancement in parenting behaviors in stressed virgin

females (Figure 6O, P).

Discussion

Previously, we have observed that virgin females showing alloparental behavior toward

pups have a low-level of activation of PeFA^{Ucn3} neurons while activating these cells highly with

chemogenetic or optogenetic methods leads to infant-directed neglect and aggression toward

pups (Autry *et al.*, 2021). In the present study, we further explored the role of urocortin-3

neurons of the perifornical area during female alloparental and maternal behavior. We aimed to

examine the intersection between the role of these neurons in infant-directed behavior and their

putative role in the hypothalamic-pituitary-adrenal axis. We hypothesized that PeFA^{Ucn3} neurons

become more active with stress and increased PeFA^{Ucn3} cell activity would lead to deficits in

pup-directed behavior. Therefore, we studied activation levels of PeFA^{Ucn3} neurons in virgin

females and mothers exposed to pups with and without stress and studied the effect of

PeFA^{Ucn3} neuron inhibition on alloparental behavior under non-stressed and stressed

conditions.

We find that around 20% of PeFA^{Ucn3} neurons are active during infant-directed behavior

in virgin females, replicating our previous findings (Autry *et al.*, 2021). In addition, we replicated

our finding that mothers have a lower level of PeFA^{Ucn3} neuronal activation during pup exposure

compared to virgin females. However, we noticed that the controls for our mother group, in

which we typically remove the litter, had a high baseline of PeFA^{Ucn3} neuron activity compared

to virgin females exposed to bedding. We therefore added a group of mothers who did not have

their litters removed. This experiment revealed that our control bedding exposure and

experimental pup exposure conditions impact PeFA^{Ucn3} neural activity differentially in mothers

depending on whether the mother's litter is present. These results indicate that PeFA^{Ucn3}

neurons may be sensitive to social contexts; these neurons appear to have high baseline

activity in mothers with their litters removed during a control bedding exposure and this activity

level plummets with introduction of foreign pups. In the future, it will be important to tease apart

whether this heightened PeFA^{Ucn3} neural activity after litter removal may be related to an

aversive or stressed state, or possibly a social motivation set point that is altered after a female

gives birth. Indeed, previous studies have shown that maternal separation can impact both a

mother and their offspring's behavior in measures related to anxiety, social behavior, and

cognition (Lemaire *et al.*, 2000; Weinstock, 2001; Chapillon *et al.*, 2002).

We next assessed the impact of inhibiting this low-level of activity during infant-directed

behavior. We found that there was a subtle but significant effect of inhibition PeFA^{Ucn3} neurons

on pup-retrieval latency in virgin females. We previously observed that activation of the

excitatory PeFA^{Ucn3} neuron projections to the ventromedial hypothalamus or lateral septum

mediate infant avoidance and neglect. Our current results suggest that inhibition of PeFA^{Ucn3}

neurons in virgin females may lead to decreased activity in these target areas responsible for

negative pup directed behavior, allowing for faster infant retrieval. While the behavioral impact

of this manipulation is relatively minor, it is in line with the low-level of activation we observe at

the cellular level.

In a parallel set of experiments, we tested the impact of chronic stress on infant-directed

behavior and PeFA^{Ucn3} neuronal activity in virgin females and mothers. We started by testing a

chronic restraint stress paradigm in virgin females. With this paradigm, we observed significant

weight loss in females exposed to daily restraint compared to unstressed females. Prior to

testing infant-directed behavior, we tested open-field behavior to ensure that the stress

paradigm had a behavioral impact after two weeks of chronic restraint, and indeed we observed

a reduction in exploration time of the center of the arena in stressed females. Thus, we

continued with the infant-directed behavior assay and observed that females with chronic

restraint showed significantly less pup retrieval compared to unstressed females, with 2 out of 9

stressed females displaying infant-directed aggression behavior. When we examined the

physiological effects of stress, we found that circulating CORT levels were not impacted though

PVH^{CRF} neuron activation was significantly reduced in females subjected to stress, suggestive

of HPA axis habituation to the repeated stress. However, we did find enhanced PeFA^{Ucn3}

neuron activity in chronically stressed females. Intriguingly, we found a strong negative

correlation between PeFA^{Ucn3} neuronal activity levels and overall alloparenting behavior in

unstressed females, and this correlation was lost in stressed females. Together, these data

support our hypothesis that stress increases activity of PeFA^{Ucn3} neurons, and heightened

activity in PeFA^{Ucn3} neurons negatively impacts infant-directed behavior.

We therefore proceeded with the chronic restraint paradigm in lactating females. We

used a similar timeline for testing, with daily weighing and an open field test prior to pup

exposure. We observed significant weight loss in lactating females exposed to stress, however

our open field test revealed that females with chronic restraint stress displayed increased

exploration of the center of the open field relative to control females. This behavior may be a

sign of hypervigilance and may be explained by an increase in distance traveled as well as

velocity (Cabib *et al.*, 1988; Sequeira-Cordero *et al.*, 2019; Rudolph *et al.*, 2020). In our

subsequent infant-directed behavior assessment, we found no differences in maternal behavior

between unstressed and chronically restrained mothers. In our physiological measures, we

observed that CORT levels were decreased in mothers with chronic restraint stress with a

decrease in both CRF and Ucn3 neuronal activation. The decrease in CORT and CRF neuronal

activation are indicative of HPA axis habituation to the repeated stress. Indeed, previous

studies have illustrated a reduction in CRF/Fos colabeling or electrophysiological properties of

CRF neurons in the PVH with repeated restraint stress in rodents (Bonaz & Rivest, 1998;

Matovic *et al.*, 2020). We interpret the decrease in Ucn3 neuronal activity as a protective

mechanism, preserving maternal behavior under stressful conditions. Indeed, previous studies

show that lactating mothers display changes in HPA axis responsivity to stress (Johnstone *et*

al., 2000; Douglas *et al.*, 2003; Klampfl & Bosch, 2019), and reduction in PeFA^{Ucn3} neuronal

activity may contribute to behavioral adaptations to stress.

To overcome the habituation to repeated restraint we observed in mothers, we

attempted to perform chronic social stress in lactating females (Supplemental Figure 4-1).

However, the stress did not result in weight changes or impact maternal behavior. In the future,

we hope to identify a stress paradigm for lactating females that impacts maternal behavior.

Furthermore, we aim to study the role of PeFA^{Ucn3} neurons in HPA axis hypo-responsivity in

mothers to uncover their potential role in maternal behavior preservation.

We plotted female infant-directed behaviors as ethograms to gain broader insight into

how stress affects this complex interaction. Surprisingly, we found that virgin females display

longer bouts of infant-directed behaviors during the pup exposure assay relative to mothers,

whose behavioral motifs appear to be more sporadic from one behavior to the next (Zoubovsky

et al., 2020). We suspect that this difference is at least in part due to the removal of the litter

during habituation for mothers that is not required for testing virgin females. However, this

result does imply that, at least in terms of studying the neural architecture of infant-directed

behavior, (1) we can collect a rich dataset from virgin females and (2) that we must be diligent in

considering the conditions under which we test behavior in mothers in the laboratory given that

litter removal may have a significant impact on some experimental parameters (Lonstein, 2005;

Smith & Lonstein, 2008; Miller *et al.*, 2011).

Finally, we tested if we could rescue stress-induced deficits in alloparental behavior by

inhibiting PeFA^{Ucn3} neurons. We confirmed that our chronic restraint stress paradigm led to

weight reductions and proceeded with our pup exposure assay. We found that chemogenetic

inhibition of PeFA^{Ucn3} neurons did lead to improved alloparental behavior on several measures

including latency to retrieve, time in nest, and crouching. We designed the experiment to

observe the effect of stress alone on the first day with vehicle administration followed by

inhibition of PeFA^{Ucn3} neurons by CNO treatment on the second day, with a final vehicle test

after drug washout. We settled on this design based on our previous observation that blocking

PeFA^{Ucn3} neuronal activity optogenetically led to prolonged improvement of infant-directed

behavior. We included two control groups, the nonstressed Cre positive group, and the

stressed Cre negative group to control for the effects of repeated testing. Our data illustrate that

inhibiting PeFA^{Ucn3} neurons in stressed females leads to more substantial effects on alloparental

behavior compared to either control group.

Overall, we find that PeFA^{Ucn3} neuronal activity is higher in females showing lower levels

of positive infant-directed behavior, a trend that can be observed regardless of physiological

status. Chronic stress leads to reduced alloparental behavior accompanied by higher numbers

of active PeFA^{Ucn3} neurons in virgin females. Blocking activity in PeFA^{Ucn3} neurons rescues

infant-directed behavioral deficits in virgin females. Together with previous studies, our results

suggest the important role for the level of PeFA^{Ucn3} neuronal activity in the expression of pro-

and anti-parental behavior (Supplemental Figure 6-3). These results support the critical role for

PeFA^{Ucn3} neurons in the neural circuitry controlling female parental behaviors and the sensitivity

of these behaviors to stress.

Author Contributions

478 B.A. and A.E.A. designed and performed experiments, analyzed, and plotted data, and

479 interpreted data and wrote the paper. R.A. supported animal experiments and analyzed data.

480 I.C. analyzed and plotted data.

Acknowledgements and funding

A.E.A. was supported by a NARSAD Young Investigator Award and a Pathway to

Independence Award (NIH R00HD085188). B.A. was supported by a diversity supplement to

A.E.A.'s NIH award (R00HD085188-S1) and a Tishman Scholarship. We thank Catherine

Dulac for intellectual input and project support at Harvard University. We thank Stacey Sullivan

for assistance with transfer of data and mice from Harvard University to Albert Einstein College

of Medicine. We thank Krysten Garcia for histology assistance. We thank Giovanni Podda for

guidance on the photometry analysis. We thank Kostantin Dobrenis, Vladimir Mudragel, and

Mariah Marrero for guidance on Axioscan usage. We thank Kevin Fisher for assistance with

image export and analysis. We thank all the members of the Autry lab for input on manuscript

preparation.

References

Autry, A.E., Adachi, M., Cheng, P. & Monteggia, L.M. (2009) Gender-specific impact of brain-
derived neurotrophic factor signaling on stress-induced depression-like behavior. *Biol
Psychiatry*, **66**, 84-90.

498
499 Autry, A.E., Wu, Z., Kapoor, V., Kohl, J., Bambah-Mukku, D., Rubinstein, N.D., Marin-
Rodriguez, B., Carta, I., Sedwick, V., Tang, M. & Dulac, C. (2021) Urocortin-3 neurons in
the mouse perifornical area promote infant-directed neglect and aggression. *Elife*, **10**.

502
503 Bielsky, I.F., Hu, S.B., Ren, X., Terwilliger, E.F. & Young, L.J. (2005) The V1a vasopressin
receptor is necessary and sufficient for normal social recognition: a gene replacement
study. *Neuron*, **47**, 503-513.

506
507 Bonaz, B. & Rivest, S. (1998) Effect of a chronic stress on CRF neuronal activity and expression
of its type 1 receptor in the rat brain. *Am J Physiol*, **275**, R1438-1449.

509
510 Brunton, P.J., Russell, J.A. & Douglas, A.J. (2008) Adaptive responses of the maternal
hypothalamic-pituitary-adrenal axis during pregnancy and lactation. *J Neuroendocrinol*,
20, 764-776.

513
514 Cabib, S., Kempf, E., Schleef, C., Mele, A. & Puglisi-Allegra, S. (1988) Different effects of acute
and chronic stress on two dopamine-mediated behaviors in the mouse. *Physiol Behav*,
43, 223-227.

517
518 Cameron, N.M., Champagne, F.A., Parent, C., Fish, E.W., Ozaki-Kuroda, K. & Meaney, M.J.
(2005) The programming of individual differences in defensive responses and
reproductive strategies in the rat through variations in maternal care. *Neurosci Biobehav
Rev*, **29**, 843-865.

522
523 Carcea, I., Caraballo, N.L., Marlin, B.J., Ooyama, R., Riceberg, J.S., Mendoza Navarro, J.M.,
Opendak, M., Diaz, V.E., Schuster, L., Alvarado Torres, M.I., Lethin, H., Ramos, D.,
Minder, J., Mendoza, S.L., Bair-Marshall, C.J., Samadjopoulos, G.H., Hidema, S.,
Falkner, A., Lin, D., Mar, A., Wadghiri, Y.Z., Nishimori, K., Kikusui, T., Mogi, K., Sullivan,
R.M. & Froemke, R.C. (2021) Oxytocin neurons enable social transmission of maternal
behaviour. *Nature*, **596**, 553-557.

529
530 Carini, L.M., Murgatroyd, C.A. & Nephew, B.C. (2013) Using chronic social stress to model
postpartum depression in lactating rodents. *J Vis Exp*, e50324.

532
533 Chapillon, P., Patin, V., Roy, V., Vincent, A. & Caston, J. (2002) Effects of pre- and postnatal
stimulation on developmental, emotional, and cognitive aspects in rodents: a review.
Dev Psychobiol, **41**, 373-387.

536
537 Chen, P.B., Hu, R.K., Wu, Y.E., Pan, L., Huang, S., Micevych, P.E. & Hong, W. (2019) Sexually
538   Dimorphic Control of Parenting Behavior by the Medial Amygdala. *Cell*, **176**, 1206-1221
539   e1218.

540
541 Delpech, J.C., Wei, L., Hao, J., Yu, X., Madore, C., Butovsky, O. & Kaffman, A. (2016) Early life
542   stress perturbs the maturation of microglia in the developing hippocampus. *Brain Behav*
543   *Immun*, **57**, 79-93.

544
Deussing, J.M., Breu, J., Kuhne, C., Kallnik, M., Bunck, M., Glasl, L., Yen, Y.C., Schmidt, M.V.,
Zurmuhlen, R., Vogl, A.M., Gailus-Durner, V., Fuchs, H., Holter, S.M., Wotjak, C.T.,
Landgraf, R., de Angelis, M.H., Holsboer, F. & Wurst, W. (2010) Urocortin 3 modulates
social discrimination abilities via corticotropin-releasing hormone receptor type 2. *The*
Journal of neuroscience : the official journal of the Society for Neuroscience, **30**, 9103-
9116.

551
552 Douglas, A.J., Brunton, P.J., Bosch, O.J., Russell, J.A. & Neumann, I.D. (2003) Neuroendocrine
553   responses to stress in mice: hyporesponsiveness in pregnancy and parturition.
554   *Endocrinology*, **144**, 5268-5276.

555
556 Feifel, A.J., Shair, H.N. & Schmauss, C. (2017) Lasting effects of early life stress in mice:
557   interaction of maternal environment and infant genes. *Genes Brain Behav*, **16**, 768-780.

558
559 Fischer, E.K., Roland, A.B., Moskowitz, N.A., Tapia, E.E., Summers, K., Coloma, L.A. &
560   O'Connell, L.A. (2019) The neural basis of tadpole transport in poison frogs. *Proc Biol*
561   *Sci*, **286**, 20191084.

562
563 Girotti, M., Pace, T.W., Gaylord, R.I., Rubin, B.A., Herman, J.P. & Spencer, R.L. (2006)
564   Habituation to repeated restraint stress is associated with lack of stress-induced c-fos
565   expression in primary sensory processing areas of the rat brain. *Neuroscience*, **138**,
566   1067-1081.

567
568 Herman, J.P. & Tasker, J.G. (2016) Paraventricular Hypothalamic Mechanisms of Chronic
569   Stress Adaptation. *Front Endocrinol (Lausanne)*, **7**, 137.

570
571 Hrvatin, S., Sun, S., Wilcox, O.F., Yao, H., Lavin-Peter, A.J., Cicconet, M., Assad, E.G., Palmer,
572   M.E., Aronson, S., Banks, A.S., Griffith, E.C. & Greenberg, M.E. (2020) Neurons that
573   regulate mouse torpor. *Nature*, **583**, 115-121.

574
Jamieson, P.M., Li, C., Kukura, C., Vaughan, J. & Vale, W. (2006) Urocortin 3 modulates the
neuroendocrine stress response and is regulated in rat amygdala and hypothalamus by
stress and glucocorticoids. *Endocrinology*, **147**, 4578-4588.

Johnstone, H.A., Wigger, A., Douglas, A.J., Neumann, I.D., Landgraf, R., Seckl, J.R. & Russell,
J.A. (2000) Attenuation of hypothalamic-pituitary-adrenal axis stress responses in late
pregnancy: changes in feedforward and feedback mechanisms. *J Neuroendocrinol*, **12**,
811-822.

583
584   Klampfl, S.M. & Bosch, O.J. (2019) Mom doesn't care: When increased brain CRF system
585   activity leads to maternal neglect in rodents. *Front Neuroendocrinol*, **53**, 100735.

586
587   Kohl, J., Babayan, B.M., Rubinstein, N.D., Autry, A.E., Marin-Rodriguez, B., Kapoor, V.,
588   Miyamishi, K., Zweifel, L.S., Luo, L., Uchida, N. & Dulac, C. (2018) Functional circuit
589   architecture underlying parental behaviour. *Nature*, **556**, 326-331.

590
591   Kronman, H., Torres-Berrio, A., Sidoli, S., Issler, O., Godino, A., Ramakrishnan, A., Mews, P.,
592   Lardner, C.K., Parise, E.M., Walker, D.M., van der Zee, Y.Y., Browne, C.J., Boyce, B.F.,
593   Neve, R., Garcia, B.A., Shen, L., Pena, C.J. & Nestler, E.J. (2021) Long-term behavioral
594   and cell-type-specific molecular effects of early life stress are mediated by H3K79me2
595   dynamics in medium spiny neurons. *Nat Neurosci*, **24**, 667-676.

596
597   Kuperman, Y., Issler, O., Regev, L., Musseri, I., Navon, I., Neufeld-Cohen, A., Gil, S. & Chen, A.
598   (2010) Perifornical Urocortin-3 mediates the link between stress-induced anxiety and
599   energy homeostasis. *Proceedings of the National Academy of Sciences of the United
600   States of America*, **107**, 8393-8398.

601
602   Kuroda, K.O., Tachikawa, K., Yoshida, S., Tsuneoka, Y. & Numan, M. (2011) Neuromolecular
603   basis of parental behavior in laboratory mice and rats: with special emphasis on
604   technical issues of using mouse genetics. *Prog Neuropsychopharmacol Biol Psychiatry*,
605   **35**, 1205-1231.

606
607   Lemaire, V., Koehl, M., Le Moal, M. & Abrous, D.N. (2000) Prenatal stress produces learning
608   deficits associated with an inhibition of neurogenesis in the hippocampus. *Proc Natl
609   Acad Sci U S A*, **97**, 11032-11037.

610
611   Lonstein, J.S. (2005) Reduced anxiety in postpartum rats requires recent physical interactions
612   with pups, but is independent of suckling and peripheral sources of hormones. *Horm
613   Behav*, **47**, 241-255.

614
615   Lonstein, J.S. & De Vries, G.J. (2000) Sex differences in the parental behavior of rodents.
616   *Neurosci Biobehav Rev*, **24**, 669-686.

617
618   Marlin, B.J., Mitre, M., D'Amour J, A., Chao, M.V. & Froemke, R.C. (2015) Oxytocin enables
619   maternal behaviour by balancing cortical inhibition. *Nature*, **520**, 499-504.

Maruska, K.P., Butler, J.M., Field, K.E., Forester, C. & Augustus, A. (2020) Neural Activation Patterns Associated with Maternal Mouthbrooding and Energetic State in an African Cichlid Fish. *Neuroscience*, **446**, 199-212.

Matovic, S., Ichiyama, A., Igarashi, H., Salter, E.W., Sunstrum, J.K., Wang, X.F., Henry, M., Kuebler, E.S., Vernoux, N., Martinez-Trujillo, J., Tremblay, M.E. & Inoue, W. (2020) Neuronal hypertrophy dampens neuronal intrinsic excitability and stress responsiveness during chronic stress. *J Physiol*, **598**, 2757-2773.

Medina, J., De Guzman, R.M. & Workman, J.L. (2021) Lactation is not required for maintaining maternal care and active coping responses in chronically stressed postpartum rats: Interactions between nursing demand and chronic variable stress. *Horm Behav*, **136**, 105035.

Miller, S.M., Piasecki, C.C. & Lonstein, J.S. (2011) Use of the light-dark box to compare the anxiety-related behavior of virgin and postpartum female rats. *Pharmacol Biochem Behav*, **100**, 130-137.

Mogi, K., Takakuda, A., Tsukamoto, C., Ooyama, R., Okabe, S., Koshida, N., Nagasawa, M. & Kikusui, T. (2017) Mutual mother-infant recognition in mice: The role of pup ultrasonic vocalizations. *Behav Brain Res*, **325**, 138-146.

Moy, S.S., Nadler, J.J., Young, N.B., Nonneman, R.J., Segall, S.K., Andrade, G.M., Crawley, J.N. & Magnuson, T.R. (2008) Social approach and repetitive behavior in eleven inbred mouse strains. *Behav Brain Res*, **191**, 118-129.

Murgatroyd, C.A., Hicks-Nelson, A., Fink, A., Beamer, G., Gurel, K., Elnady, F., Pittet, F. & Nephew, B.C. (2016) Effects of Chronic Social Stress and Maternal Intranasal Oxytocin and Vasopressin on Offspring Interferon-gamma and Behavior. *Front Endocrinol (Lausanne)*, **7**, 155.

Murgatroyd, C.A. & Nephew, B.C. (2013) Effects of early life social stress on maternal behavior and neuroendocrinology. *Psychoneuroendocrinology*, **38**, 219-228.

Murgatroyd, C.A., Pena, C.J., Podda, G., Nestler, E.J. & Nephew, B.C. (2015) Early life social stress induced changes in depression and anxiety associated neural pathways which are correlated with impaired maternal care. *Neuropeptides*, **52**, 103-111.

Nephew, B.C. & Bridges, R.S. (2011) Effects of chronic social stress during lactation on maternal behavior and growth in rats. *Stress*, **14**, 677-684.

Neufeld-Cohen, A., Kelly, P.A., Paul, E.D., Carter, R.N., Skinner, E., Olverman, H.J., Vaughan, J.M., Issler, O., Kuperman, Y., Lowry, C.A., Vale, W.W., Seckl, J.R., Chen, A. &

Jamieson, P.M. (2012) Chronic activation of corticotropin-releasing factor type 2 receptors reveals a key role for 5-HT1A receptor responsiveness in mediating behavioral and serotonergic responses to stressful challenge. *Biological psychiatry*, **72**, 437-447.

Numan, M. (1974) Medial preoptic area and maternal behavior in the female rat. *J Comp Physiol Psychol*, **87**, 746-759.

Numan, M. (2020) *The parental brain : mechanisms, development, and evolution.* Oxford University Press, New York.

Numan, M. & Insel, T.R. (2003) *The neurobiology of parental behavior.* Springer-Verlag, New York.

O'Connell, L.A., Matthews, B.J. & Hofmann, H.A. (2012) Isotocin regulates paternal care in a monogamous cichlid fish. *Horm Behav*, **61**, 725-733.

Ostermeyer, M.C. & Elwood, R.W. (1983) Pup recognition in Mus musculus: parental discrimination between their own and alien young. *Dev Psychobiol*, **16**, 75-82.

Paulson, J.F. & Bazemore, S.D. (2010) Prenatal and postpartum depression in fathers and its association with maternal depression: a meta-analysis. *JAMA*, **303**, 1961-1969.

Radley, J.J. & Sawchenko, P.E. (2015) Evidence for involvement of a limbic paraventricular hypothalamic inhibitory network in hypothalamic-pituitary-adrenal axis adaptations to repeated stress. *J Comp Neurol*, **523**, 2769-2787.

Richter, K., Wolf, G. & Engelmann, M. (2005) Social recognition memory requires two stages of protein synthesis in mice. *Learn Mem*, **12**, 407-413.

Rincon-Cortes, M. & Grace, A.A. (2021) Postpartum scarcity-adversity disrupts maternal behavior and induces a hypodopaminergic state in the rat dam and adult female offspring. *Neuropsychopharmacology*.

Rosinger, Z.J., Mayer, H.S., Geyfen, J.I., Orser, M.K. & Stolzenberg, D.S. (2021) Ethologically relevant repeated acute social stress induces maternal neglect in the lactating female mouse. *Dev Psychobiol*, **63**, e22173.

Rudolph, S., Guo, C., Pashkovski, S.L., Osorno, T., Gillis, W.F., Krauss, J.M., Nyitrai, H., Flaquer, I., El-Rifai, M., Datta, S.R. & Regehr, W.G. (2020) Cerebellum-Specific Deletion of the GABAA Receptor delta Subunit Leads to Sex-Specific Disruption of Behavior. *Cell Rep*, **33**, 108338.

Sato, K., Hamasaki, Y., Fukui, K., Ito, K., Miyamichi, K., Minami, M. & Amano, T. (2020) Amygdalohippocampal Area Neurons That Project to the Preoptic Area Mediate Infant-Directed Attack in Male Mice. *J Neurosci*, **40**, 3981-3994.

Sequeira-Cordero, A., Salas-Bastos, A., Fornaguera, J. & Brenes, J.C. (2019) Behavioural characterisation of chronic unpredictable stress based on ethologically relevant paradigms in rats. *Sci Rep*, **9**, 17403.

Singh-Taylor, A., Korosi, A., Molet, J., Gunn, B.G. & Baram, T.Z. (2015) Synaptic rewiring of stress-sensitive neurons by early-life experience: a mechanism for resilience? *Neurobiol Stress*, **1**, 109-115.

Slawski, B.A. & Buntin, J.D. (1995) Preoptic area lesions disrupt prolactin-induced parental feeding behavior in ring doves. *Horm Behav*, **29**, 248-266.

Smith, C.D. & Lonstein, J.S. (2008) Contact with infants modulates anxiety-generated c-fos activity in the brains of postpartum rats. *Behavioural brain research*, **190**, 193-200.

Tsuneoka, Y., Tokita, K., Yoshihara, C., Amano, T., Esposito, G., Huang, A.J., Yu, L.M., Odaka, Y., Shinozuka, K., McHugh, T.J. & Kuroda, K.O. (2015) Distinct preoptic-BST nuclei dissociate paternal and infanticidal behavior in mice. *EMBO J*, **34**, 2652-2670.

Walker, C.D., Tilders, F.J. & Burlet, A. (2001) Increased colocalization of corticotropin-releasing factor and arginine vasopressin in paraventricular neurones of the hypothalamus in lactating rats: evidence from immunotargeted lesions and immunohistochemistry. *J Neuroendocrinol*, **13**, 74-85.

Wang, X.D., Rammes, G., Kraev, I., Wolf, M., Liebl, C., Scharf, S.H., Rice, C.J., Wurst, W., Holsboer, F., Deussing, J.M., Baram, T.Z., Stewart, M.G., Muller, M.B. & Schmidt, M.V. (2011) Forebrain CRF(1) modulates early-life stress-programmed cognitive deficits. *J Neurosci*, **31**, 13625-13634.

Weinstock, M. (2001) Alterations induced by gestational stress in brain morphology and behaviour of the offspring. *Prog Neurobiol*, **65**, 427-451.

Wisner, K.L., Sit, D.K., McShea, M.C., Rizzo, D.M., Zoretich, R.A., Hughes, C.L., Eng, H.F., Luther, J.F., Wisniewski, S.R., Costantino, M.L., Confer, A.L., Moses-Kolko, E.L., Famy, C.S. & Hanusa, B.H. (2013) Onset timing, thoughts of self-harm, and diagnoses in postpartum women with screen-positive depression findings. *JAMA Psychiatry*, **70**, 490-498.

Wu, Z., Autry, A.E., Bergan, J.F., Watabe-Uchida, M. & Dulac, C.G. (2014) Galanin neurons in
the medial preoptic area govern parental behaviour. *Nature*, **509**, 325-330.

Zoubovsky, S.P., Hoseus, S., Tumukuntala, S., Schulkin, J.O., Williams, M.T., Vorhees, C.V. &
Muglia, L.J. (2020) Chronic psychosocial stress during pregnancy affects maternal
behavior and neuroendocrine function and modulates hypothalamic CRH and nuclear
steroid receptor expression. *Transl Psychiatry*, **10**, 6.

Figures and legends

Ucn3 Fos DAPI

Figure 1. PeFA Urocortin-3 neuronal activation levels in response to foreign pups

depends on physiological context. (A) Schematic of behavioral paradigm. C57 virgin females

were exposed to a newborn pup and sacrificed 30 minutes after pup exposure or addition of

fresh bedding into home cage (control). Lactating females either had a litter removed or litter

intact and exposed to a foreign-born pup or fresh bedding. **(B)** Rostral perifornical area cells

containing *Urocortin-3* and *Fos* RNA were counted for colocalization in each group. (scale bar

100 µm**) (C)** Quantification of percentage of Ucn3+ cells colocalized with Fos across groups.

Fisher exact test reveals that pup exposure increases PeFA^{Ucn3} activation in virgin females

(Control n=605 N=5; Pup exposed n=655 N=8; ****p=<0.0001), reduced PeFA^{Ucn3} activation in

lactating females with litter removed (Control n=622 N=7; pup exposed n=828 N=8;

**p=0.0044), and increased activation in lactating females that did not have litter removal

compared to control bedding exposure (Control n=816 N=6; pup exposed n=988 N=8

****p<0.0001).

Supplemental Figure 1-1. Colocalization of PeFA Urocortin-3 neurons with Fos in

response to foreign pups. Representative images from Figure 1 with Ucn3 and Fos channels

separated (scale bar 100 µm).

Figure 2. Inhibition of PeFA[Ucn3] neurons enhances alloparental behavior in virgin female

mice. (A) Schematic of viral injection strategy and behavior timeline (n=10 Cre- females; n=8

Cre+ females). **(B)** Representative image of mCherry reporter expression in Ucn3::Cre+

female injected with inhibitory DREADD virus (magenta: mCherry; blue: DAPI; scale bar 100

µm). **(C)** Percentage of pups retrieved by Cre+ females is significantly increased compared to

Cre- females (Kolmogorov-Smirnov test p=0.0048). **(D)** Latency to retrieve pups is significantly

faster in Cre+ females compared to Cre- females (Two-tailed Mann-Whitney test; p=0.0033), as

well as **(E)** latency to retrieve the first pup (Two-tailed Mann-Whiney test; p=0.0259), but there

was no difference between groups in **(F)** latency to retrieve the second pup. **(G-H)** Latency to

approach pups was not significantly different, but time spent investigating pups trended lower in

Cre+ animals (Unpaired t test; p=0.0854). **(I-J)** We observed no significant difference in latency

to pup groom or in time spent pup grooming in Cre+ animals compared to Cre- females. **(K)**

Time spent in nest with pups significantly increased in Cre+ animals (Unpaired t test; p=0.0289).

(L-M) Latency to crouch trended lower in Cre+ animals (Two-tailed Mann-Whiney test;

p=0.1011), but we did not observe a significant difference in time spent crouching. **(N-O)**

Latency to nest build was significantly faster in Cre+ animals (Two-tailed Mann-Whiney test;

p=0.0152) but time spent nest building was not significantly different between groups. **(P)**

Cumulative time spent parenting was unchanged between groups **(Q)** Representative behavior

trace of a Cre- animal (left) and a Cre+ animal (right) during pup assay (time 0 is when pup was

added to home cage).

mCherry DAPI

800 **Supplemental Figure 2-1. Inhibitory DREADD expression in virgin female mice.**

801 Representative image of AAV-mediated hM4di DREADD expression in the perifornical area

802 (PeFA) of Ucn::Cre- (left) or Ucn3::Cre+ (right) females. Third ventricle indicated by outline

803 (scale bar 100 μm).

Figure 3. Chronic restraint stress dampens alloparental behavior in virgin female mice.

(A) Schematic of timeline using restraint stress paradigm and behavioral testing in virgin

females (n=9 control; n=9 stress). **(B)** Weights taken from each group during the period of

chronic restraint stress. Stressed females have significant difference in weight compared to

controls (Two-way repeated measures ANOVA; main effect of interaction of stress x time

$F_{(10,220)}$=22.92 p<0.0001; main effect of time $F_{(10, 220)}$ = 20.35; Bonferroni's multiple comparisons

test) **(C-F)** Time spent in center of open field was significantly reduced for stressed females

compared to controls (C) (unpaired t test; p=0.0289) but no other parameter in open field was

changed. **(G)** Chronic restraint stress significantly decreased cumulative pup retrieval in females

(Kolmogorov-Smirnov test p=0.0059).**(H-Q)** Chronic restraint stress did not significantly change

other parenting measures such as retrieval latency, or time spent pup grooming, time in nest,

crouching, and nest building. **(R)** Representative behavior trace for a control female (left) and

stressed female (right) during pup assay. (*p=0.05; **p=0.01; ***p=0.001; ****p=0.0001).

Figure 4. Chronic restraint stress does not induce changes in parental behavior in

lactating females. (A) Schematic of timeline using restraint paradigm and behavioral testing in

lactating females (Control n=12; Stressed n=13). **(B)** Weights taken from each group during the

period of chronic restraint stress from postpartum day 1 to 18. Stressed females have a

significant difference in weight compared to controls (Two-way repeated measures ANOVA;

main effect of interaction of stress x time $F_{(17,368)}=8.286$ p<0.0001; main effect of time

$F_{(7.579,174.3)}=75.28$ p<0.0001; Sidak's multiple comparisons). **(C-F)** Time spent in center of open

field was significantly increased for stressed females compared to controls (unpaired t test;

p=0.0459) and time spent in borders decreased (unpaired t test; p=0.0459) accompanied by

increased distance traveled (unpaired t test; p=0.0089) and velocity (unpaired t test; p=0.0089)

(G-Q) Chronic restraint stress did not significantly change any parenting measures as retrieval,

pup grooming, time in nest, crouching, and nest building compared to females that did not

receive stress. **(R)** Representative behavior trace for a control female (left) and stressed female

(right) during pup assay.

Supplemental Figure 4-1. Intruder stress in lactating females does not impact weight or parental behavior. (A) Experimental paradigm for intruder stress in lactating females. **(B)** Weight was not affected by stress condition. **(C)** Percentage of pups retrieved was not significantly different between control and stressed females. **(D)** Latency to retrieve **(E)** Pup investigation **(F)** Latency to groom **(G)** Grooming duration **(H)** Time in the nest **(I)** Latency to

841   crouch **(J)** Crouching duration **(K)** Latency to nest build **(L)** Nest building duration and **(M)**

842   Parental behaviors were unchanged between stress and control mothers.

Figure 5. Chronic restraint leads to contrasting molecular changes between virgin and lactating females in parenting. (A) Rostral perifornical area cells containing *Ucn3, CRF,* and *Fos* RNA were counted for colocalization in each virgin female group. (scale bar 100 μm). **(B)**

Colocalization of *Ucn3* and *Fos* in perifornical area reveals increased of activation of PeFA^{Ucn3}

neurons in in stressed virgin females compared to control virgin females (Fisher exact test:

Control n=490 N=9; Stressed n=472 N=9; p=0.038). **(C)** Colocalization of *CRF* and *Fos* in

paraventricular nucleus of the hypothalamus reveals decreased activation of PVH^{CRF} neurons in

stressed virgin females compared to control virgin females (Fisher exact test: Control n=1305

N=9; Stressed n=1272 N=9; p<0.0001). **(D)** Serum corticosterone levels were measured using

ELISA in virgin females. Chronic restraint stress did not contribute to altered corticosterone in

virgin females. **(E)** Plotting time spent parenting against PeFA^{Ucn3} activation levels shows

marked negative correlation between parenting and PeFA^{Ucn3} neuron activation in controls but

this relationship is abolished in stressed virgin females (Control R^2=0.51 Stressed R^2=0.13;

difference in slope F= 10.73 DFn=1, DFd=14; p=0.0055). **(F)** PVH^{CRF} activation levels in

negatively correlated with time spent parenting in control virgin females but this relationship is

indiscernible in stressed virgin females (Control R^2=0.68 Stressed R^2=0.01). **(G)** Rostral

perifornical area cells containing *Ucn3, CRF,* and *Fos* RNA were counted for colocalization in

each lactating female group (scale bar 100 µm). **(H)** PeFA^{Ucn3} neuron activation levels are

reduced in stressed lactating females (Fisher exact test: Control n=338 N=8; Stressed n=360

N=6; p=0.0005). **(I)** Activation levels of CRF cells in the paraventricular nucleus of the

hypothalamus are significantly decreased in stressed lactating females (Control n=919 N=6;

Stressed n=2268 N=5; p<0.0001). **(J)** Chronic restraint stress significantly reduced

corticosterone in lactating females (unpaired t test; p=0.0198). **(K)** Linear regression analysis

reveals a trend towards negative correlation between time spent parenting and PeFA^{Ucn3}

activation levels in control lactating females but not in the stressed group (Control R^2=0.27

Stressed R^2=0.01). **(L)** PVH^{CRF} activation is positively correlated with time spent parenting

(Control R^2=0.45 Stressed R^2=0.39).

Supplemental Figure 5-1. Colocalization of Ucn3 or CRF and Fos in virgin or lactating

females exposed to pups in control or chronic restraint stress conditions. Representative

images from Figure 5 with Ucn3 or CRF and Fos channels separated (Ucn3 or CRF green

channel; Fos magenta channel; scale bar 100 µm).

Supplemental Figure 5-2. Intruder stress in lactating females does not affect PeFA^{Ucn3}

neuronal activation levels or circulating corticosterone. (A) Percentage of Fos+ PeFA^{Ucn3}

neurons is not significantly different between mothers with or without intruder stress. **(B)**

Corticosterone levels are indistinguishable between mothers with or without intruder stress. **(C)**

Linear regression analysis reveals a trend towards negative correlation between time spent

parenting and PeFA^{Ucn3} activation levels in control lactating females but not in mothers with

intruder stress (Control R^2=0.4347 Stressed R^2=0.0052).

Figure 6. PeFA^Ucn3 inhibition ameliorates parenting deficits in stressed virgin females. (A)

Schematic of viral injection strategy and behavior timeline (n=7 stressed Cre+ females; n=7

control Cre+ females). **(B)** Stressed females had a significant difference in weight compared to control females (2-way repeated measures ANOVA, main effect of stress x time p<0.0001). **(C-F)** Open field results show no difference in time spent in center, border, distance moved, or velocity in control or stressed mice with or without CNO treatment. **(G)** Cumulative pup retrieval in control animals improves significantly with CNO injection (Friedman's test; p=0.0013). **(H)** Cumulative pup retrieval in stressed animals improve with CNO administration (Friedman's test; p=0.0003). **(I)** CNO treatment decreases latency to retrieve in stressed females (Two-way repeated measures ANOVA, main effect time $F_{(2,24)}$=8.978 p<0.01; Sidak's multiple comparisons test) but not in control animals. **(J)** Pup investigation dramatically reduces in both animal groups with CNO (Two-way repeated measures ANOVA, main effect of time $F_{(2,24)}$=82.86 p<0.0001; Sidak's multiple comparisons test). **(K)** Pup grooming is unchanged with CNO treatment. **(L)** Time spent crouching increases with CNO administration in stressed females (Two-way repeated measures ANOVA, main interaction effect of stress and drug $F_{(2,24)}$=3.506 p<0.0462); Tukey's multiple comparisons). **(M)** Stressed females spent significantly more time in the nest with CNO administration (Two-way repeated measures ANOVA, main effect of time $F_{(2,24)}$=9.399 p=0.001; Sidak's multiple comparison's test). **(N)** Cumulative time spent parenting increases with CNO treatment in stressed females (Two-way ANOVA, main effect of time $F_{(2,24)}$=3.942 p=0.0331; Sidak's multiple comparisons). **(O-P)** Representative behavior traces showing induction of more parenting behaviors with CNO in stressed females. (Significant post-hoc comparisons noted as follows: *p=0.05; **p=0.01; ***p=0.001; ****p=0.0001).

Supplemental Figure 6-1. CNO treatment does not impact parenting behavior in

Ucn3::Cre- stressed virgin females. (A) Cumulative pup retrieval in control animals improves

significantly with CNO injection (Friedman's test; p=0.0002 followed by Dunn's posthoc

comparisons). **(B)** CNO reduces latency to retrieve pups in stressed Ucn3::Cre+ females but not

in stressed Cre- females (Two-way repeated measures ANOVA main effect of drug treatment

$F_{(2,20)}$=9.667 p<0.0012, Sidak multiple comparisons post-hoc effect significant for Cre+ stressed

group vehicle versus CNO p<0.0224 and Cre- stressed group vehicle vs. vehicle 2 p<0.0036).

(C) Pup investigation is impacted in the stressed Cre+ group but not the stressed Cre- group

(Two-way repeated measures ANOVA main effect of drug treatment $F_{(2,20)}$=10.94 p<0.0006,

Sidak multiple comparisons post-hoc effect significant for Cre+ stressed group vehicle versus

CNO p<0.0010 and vehicle versus vehicle 2 p<0.0009). **(D)** Pup grooming is unaffected. **(E)**

CNO treatment increases crouching significantly in Ucn3::Cre positive females (Two-way

repeated measures ANOVA main effect of drug treatment $F_{(2,20)}$=4.898, p<0.0186, Tukey

multiple comparisons post-hoc effect significant for Cre+ stress group vehicle versus CNO

p<0.05) **(F)** as well as duration in nest (Two-way repeated measures ANOVA main effect of

drug treatment $F_{(2,20)}$=8.098, p<0.0037, Sidak multiple comparisons post-hoc effect significant

for Cre+ stress group vehicle versus CNO p<0.05) **(G)** and cumulative time spent parenting

(Two-way repeated measures ANOVA main effect of drug treatment $F_{(2,20)}$=7.728 p<0.0033,

Sidak multiple comparisons post-hoc effect significant for Cre+ stress group vehicle versus CNO

(p<0.05).

Supplemental Figure 6-2. Confirmation of CNO reduction of PeFA^{Ucn3} on AAV-mediated

hM4di in Ucn3::Cre females. (A) Representative image of changes in PeFA^{Ucn3}/c-Fos levels in

AAV-hM4di DREADD injected Ucn3::Cre females when CNO is injected intraperitoneally

compared to vehicle after 30 minutes of pup interaction (scale bar 100 μm). Arrow indicates

Ucn3/c-Fos colocalization. **(B)** Quantification of DREADD expressing neurons co-expressing c-

Fos is significantly reduced in Ucn3::Cre females injected with CNO. (Fisher exact test: Vehicle

n=180 N=2; Stressed n=101 N=2; p<0.0001). **(C)** Schematic of experimental design to record

photometric signals from the PeFA with expression of inhibitory DREADD hM4Di construct in

the PeFA^{Ucn3} neurons. **(D)** Schematic of tail suspension session design during photometry

recording. **(E)** Z-score of the normalized fluorescence signal (ΔF/F) for each individual trial

during the 10s peri-event window centered around the onset of tail suspension. Left: vehicle;

right: CNO treatment (n=6 trials, N= 4 female mice) **(F)** Average Z-score of ΔF/F averaged

across the 6 trials for each of the four females (dark line represents the mean and the shaded

area represents the SEM). **(G)** Area under the curve calculated on the average ΔF/F signal for

the tail suspension events (n=6 trials, N=4 female mice). AUC of the 5s segment following the

suspension is significantly reduced in the CNO trial compared to vehicle treatment (Paired t-test

p<0.0281, N=4 female mice, average of 6 trials for each). **(H)** Percent change of ΔF/F relative

to baseline reveals significant reduction of fluorescence evoked by tail suspension with CNO

treatment relative to vehicle treatment (Paired t-test p<0.0025, N=4 female mice, average of 6

trials for each).

Supplemental Figure 6-3. Model for PeFA^{Ucn3} neuronal activation in pup-directed

behavior across sexes and physiological states. (Left) Pro-parental behavior such as

grooming, retrieving, and crouching over pups in the nest is accompanied by low-level PeFA^{Ucn3}

neuronal activation (`25%). Mice in this category include alloparental virgin females, unstressed

mothers, and stress-resistant mothers, as demonstrated in this study and Autry et al. 2021. We

predict that this level of activation of PeFA^{Ucn3} neurons may also be observed in other categories

of mice that display pro-parental behavior including alloparental virgin males, unstressed

fathers, and stress-resistant fathers. **(Right)** On the other hand, anti-parental behavior such

infant-directed aggression or neglect is accompanied by increased activation of PeFA^{Ucn3}

neurons (~55%). Mice in this category include stressed virgin females and infanticidal males as

demonstrated in this study and Autry et al., 2021. We predict that this level of activation of

PeFA^{Ucn3} neurons may also be observed in other categories of mice that display anti-parental

behavior including stress-susceptible mothers, stressed virgin males, and stress-susceptible

fathers. Model created with Biorender.com.