Competition contributes to both warm and cool range edges

Shengman Lyu[1]*, Jake M. Alexander[1]

[1] Institute of Integrative Biology, ETH Zürich, 8092 Zürich, Switzerland

* Correspondence: shengman.lyu@usys.ethz.ch

ORCID

Shengman Lyu https://orcid.org/0000-0003-0221-5315

Jake M. Alexander https://orcid.org/0000-0003-2226-7913

Abstract

Competition plays an important role in shaping species' spatial distributions. However, it

remains unclear where and how competition regulates species' range limits. In an

experiment with plants originating from low and high elevations and conducted across an

elevation gradient in the Swiss Alps, we found that the ability of both lowland and highland

species to persist under competition was reduced beyond versus within their elevation

ranges, suggesting that competition helps set both lower and upper elevation range limits of

these species. Furthermore, the reduced ability of pairs of lowland or highland species to

coexist beyond their range edges was mainly driven by diminishing niche differences;

changes in both niche differences and relative fitness differences drove weakening

competitive dominance of lowland over highland species with increasing elevation. These

results highlight the need to account for competitive interactions and investigate underlying

coexistence mechanisms to understand current and future species distributions.

Keywords

Introduction

Climate and biotic interactions, in particular competition, are key factors constraining species' geographic ranges [1, 2, 3]. According to niche theory, a focal species is able to overcome abiotic and biotic constraints on population growth to maintain persistent populations within, but not beyond, its range edge [4]. This hypothesis has been widely supported by empirical studies. For instance, transplant experiments have often found declines in fitness when species are transplanted beyond their range edges, with greater declines in fitness when individuals are transplanted into sites where other species are present (see ref. 5 for a review). Another longstanding hypothesis regarding species' range limits is that competition is especially important for setting warm range edges, such as those at low elevation or latitude [1, 2, 6, 7]. However, while some transplant studies have found patterns consistent with this hypothesis [5], other studies indicate that competition is also important for setting cool range edges such as those at high elevation or latitude [8, 9, 10]. Thus, although the general importance of competition in shaping distributions is well established, how this varies across environmental gradients, as well as the underlying processes involved, remain unclear.

We can use tools and insights from community ecology theory to better understand when competition contributes to setting range limits [11, 12]. From the perspective of coexistence theory, within its range a population of a focal species is predicted to be able to grow from low density when other species are at their equilibrium abundance (i.e., invasion population growth rate, $\ln(\lambda_{invasion}) > 0$) and thus persist with other community members. When beyond its range limit, the same species is predicted to be unable to maintain persistent populations (i.e., $\ln(\lambda_{invasion}) < 0$), resulting in competitive exclusion [11], even if the species would be able to persist in the absence of other competitors (i.e., intrinsic population growth rate, $\ln(\lambda_{intrinsic}) > 0$). In other words, the ability of a species to coexist with its neighbours is expected to be reduced towards its range edge.

A key advantage of conceiving range limits as a coexistence problem is that we can begin to unpack the processes through which competitors influence species' range limits [12]. Modern coexistence theory predicts that coexistence is determined by two sets of differences between competing species [13]. Differences in competitive ability (i.e., relative fitness differences) reflect both intrinsic differences in species' demographic performance and/or their sensitivity to competition, and drive competitive exclusion [14]; species' differences that reduce the intensity of interspecific relative to intraspecific competition (i.e., niche differences) promote stable coexistence [15]. These can result for example from resource partitioning or specialist natural enemies [15]. Accordingly, competition could shape range

limits through two pathways. Firstly, a focal species may be competitively eliminated outside

of its range because it encounters other species with which it has greater niche overlap

compared to species within its range. For example, the replacement of closely related,

ecologically similar species across elevation gradients has been explained by interspecific

competition for shared habitats [16]. Similarly, changing environmental conditions towards

range edges can also reduce niche differences between currently co-coexisting species (i.e.,

decreased niche differences toward the range edge, see ref. 17 for a similar example),

increasing the likelihood of competitive exclusion.

The second pathway through which competition could contribute to range limits is via

changes to relative fitness differences. A focal species may coexist stably with other species

within its range but be competitively excluded beyond its range edge because of reduced

competitive ability. On the one hand, reduced competitive ability could arise because the

focal species has a reduced ability to tolerate environmental conditions outside of its range

in the absence of other competitors (i.e., a diminished $\lambda_{intrinsic}$). On the other hand, the focal

species may become more sensitive to competition, or its competitors exert stronger effects

beyond versus within its range [15, 18]. Taking an elevation gradient as an example, a high

elevation plant could be restricted to higher elevations by competition from lower elevation

species that are better competitors for shared resources, like light [19], under a warmer climate

[20, 21]; but even competition with co-occurring high elevation species might generate a range

limit, if at least some of those species become relatively stronger competitors under warmer

conditions at lower elevation (see refs. 22 and 23 for similar examples).

Here, we evaluate the hypothesis that species experience a reduced ability to coexist with

neighbours beyond versus within their range edges, and examine the contributions of niche

and relative fitness differences to changing coexistence across an elevation gradient. To

address this, we conducted a field experiment interacting seven low and seven high

elevation species with limited range overlap (hereafter lowland and highland species) in

three sites across an elevation gradient. The low (890 m a.s.l.) site was located within the

current elevation range of lowland species and beyond the lower range edge of highland

species, while the high site (1900 m) was located within the elevation range of highland

species and beyond the upper range edge of lowland species, and the middle site (1400 m)

was located near the shared range edges of both groups. In each site, we parameterized

integral projection models (IPMs) to estimate population growth rates (λ) both in the absence

(i.e., intrinsic population growth rates) and presence (i.e., invasion population growth rates)

of neighbour species. Our approach enabled us to quantify coexistence determinants (i.e.,

niche differences and relative fitness differences) and predict the outcomes of competition

between species pairs. We ask two main questions: (1) How does the ability of lowland and

highland species to persist with competitors change across an elevation gradient? (2) To

what extent are any changes in the strength of coexistence driven by changes in niche

differences and changes in relative fitness differences? Our study shows that both lowland

and highland species display a reduced ability to persist with competitors beyond their range

edges, suggesting that competition helps set range limits both at low (i.e., the lower range

edges of highland species) and high (i.e., the upper ranges edges of lowland species)

elevations. Furthermore, we show that competition can influence species' elevation ranges

through changes to both niche differences and relative fitness differences, but that their

relative importance varies depending on the origin of the competitors. These results

therefore highlight the importance of considering competition and its underlying mechanisms

to understand species' distributions across environmental gradients, and ultimately to

forecast range dynamics under changing environments.

Results

Declines in intrinsic and invasion population growth rates towards range edges

Lowland and highland species displayed distinct responses to elevation in the absence of

competition (Fig. 1a). The intrinsic population growth rates of both lowland and highland

species were projected to decline beyond elevational range edges (test based on the mean

of 500 bootstraps; elevation × species origin: $F_{1, 41}$ = 7.062, P = 0.008; Supplementary Table

7; median and 95% CI of elevation × species origin interaction across bootstrap replicates:

0.0011, 0.0008–0.0013). Interestingly, most species were projected to be able to persist in

the absence of neighbours across the whole elevation gradient, regardless of their elevation

origin (i.e., ln($\lambda_{intrinsic}$) > 0; Supplementary Fig. 6).

In the presence of competition, the projected population growth rates displayed greater

declines beyond the range edges of both lowland and highland species (Fig. 1b; test based

on the mean of 500 bootstraps; elevation × species origin: $F_{1, 300}$ = 21.215, P < 0.0001;

Supplementary Table 7; median and 95% CI of elevation × species origin interaction across

bootstrap replicates: 0.0008, 0.0005 – 0.0011). When competing against other species,

highland species were on average predicted to be able to persist at the high site, that is

within their range, but not beyond their lower range edges at the low elevation site. In

contrast, the projected ability of lowland species to persist under competition decreased with

increasing elevation, though most of them were predicted to be able to persist in all three

sites (Fig. 1b; Supplementary Fig. 6).

Changing outcomes of competition across the elevation gradient

The predicted ability of species to coexist changed across the elevation gradient, with effects that depended on competitor identity (Fig. 2). For sympatric pairs (i.e., lowland-lowland and highland-highland pairs), coexistence was more prevalent within than outside of the current ranges. On average across all bootstrap replicates, 63% of all eight and 67% of all five highland-highland pairs were predicted to coexist at the high and middle site versus only 40% at the low site (Fig. 2c). Predictions of coexistence of lowland-lowland pairs were more common and displayed only few changes across the sites, with 78% of all 10 lowland-lowland pairs predicted to coexist at the low and high sites and 50% at the middle site (Fig. 2a). Allopatric pairs (i.e., lowland-highland pairs) were more likely to coexist at the middle elevation near shared range edges, with 58% of all 13 pairs predicted to coexist at the middle site and 46% and 44% at the low and high sites, respectively (Fig. 2b).

These findings based on tallies of coexistence outcomes were supported by an analysis of a metric measuring the strength of coexistence (i.e., the coexistence metric, see Methods). These showed that both lowland-lowland and highland-highland pairs displayed weakened coexistence (i.e., decreasing coexistence metric) beyond their elevation ranges, while lowland-highland pairs displayed the greatest ability to coexistence at the middle elevation (Fig. 2d-f; test based on the mean of 500 bootstraps: elevation × competitor identity: $F_{2,\,90}$ = 11.011, P = 0.004; Supplementary Table 8). The interaction was mainly driven by the opposing responses of lowland and highland pairs, although this contrast was not significantly different from zero after error propagation (individual tests for each bootstrap replicate; median and 95% CI: -0.0006, -0.0017 – 0.00004).

Niche differences and relative fitness differences jointly mediate changing coexistence

Changes in the magnitude of both niche and relative fitness differences mediated variation in coexistence across the elevation gradient (Fig. 3). For sympatric pairs, our results showed that both lowland and highland species displayed greater niche differences (i.e., smaller niche overlap) in sites that lay within their range (Fig. 3a&c; test based on the mean of 500 bootstraps; elevation × competitor identity: $F_{2,\,90}$ = 11.603, P = 0.003; Supplementary Table 8), although this result was not significant after error propagation (individual tests for each bootstrap replicate; median and 95% CI of the contrast between lowland and highland pairs: 0.0005, -0.00005 – 0.0012). In addition, reduced relative fitness differences towards high elevation also contributed to strengthened coexistence of highland species within their range (Fig. 3f; test based on the mean of 500 bootstraps; $F_{1,17}$ = 5.179, P = 0.023), although this

result was not significant after error propagation (individual tests for each bootstrap replicate;
median and 95% CI: -0.0002, -0.0006 – 0.0001). No significant trend on relative fitness
differences was seen for lowland-lowland pairs ($F_{1,29}$ = 0.192, P = 0.661) or when pooling all
pairs together and taking absolute values (i.e., absolute differences in competitive ability
irrespective of competitor origin; test based on the mean of 500 bootstraps, $F_{1,90}$ = 1.406, P
= 0.236; Supplementary Table 8).

For parapatric pairs, lowland species were predicted to be competitively dominant over
highland species in general (i.e., ln(fitness difference) < 0), but the competitive dominance
was predicted to decrease towards high elevations, where some highland species were
predicted to be dominant (Fig. 3e; test based on the mean of 500 bootstraps, $F_{1,44}$ = 3.379, P
= 0.046). However, this result was not significant after error propagation (individual tests for
each bootstrap replicate; median and 95% CI of the slope: 0.0001, -0.0002 – 0.0005). Niche
differences between lowland and highland species tended to be greatest at middle elevation
(Fig. 3b), contributing to the greatest ability to coexist of at the middle site in conjunction with
reduced relative fitness differences (Fig. 2e).

Discussion

Although the role of competition in shaping species' spatial distribution is well acknowledged
[2, 3], how this role varies across environmental gradients, and by which processes (e.g.,
through changes in niche and relative fitness differences) competition regulates species'
range limits, remain unclear [6, 12]. Here we found that the ability of species to persist in the
presence of competition was weakened beyond versus within their elevational ranges,
supporting the role of competition in shaping species' elevation distribution limits. However,
in contrast to the conventional view that competition is more important in abiotically benign
environments, such as at low elevation or latitude (e.g., as predicted by the stress-gradient
hypothesis [6, 7, 24]), our results indicate that competition can also be important towards high
elevation range limits. There are at least two aspects that distinguish our study from
previous work that can partly help explain this discrepancy. Firstly, we focused on the
population-level outcomes of competition rather than the individual-level strength of
competition, as most previous empirical studies have done [5, 25]. Both theoretical and
empirical work demonstrate that competition can be equally important for population
dynamics in harsh and benign environments [10] and its effects cannot be completely
understood by focusing solely on the strength of competition [9, 26]. For example, harsh
environmental conditions could reduce population growth rates directly, making species less
tolerant of interspecific competition and competitive exclusion correspondingly more likely to
occur, despite any reduced intensity or prevalence of interspecific competition itself [9, 27, 28].

Secondly, we grew the same set of species in each experimental site, which allowed us to
isolate changes in the intensity of competition from turnover in community composition
across the environmental gradient [6]. Though this to some extent divorces our study from the
effects of the complex interactions that play-out in natural multispecies communities (see ref.
29 and below), it shows that both changing intensity of interactions and changing identity of
species contribute to the changing competition experienced by a focal species towards its
range edge, and we obtained further insight into these processes through analyses of
changes in niche and fitness differences across elevation.

Although effects of niche and relative fitness differences cannot be understood in isolation [13],
for our species, changes in niche differences appeared to be especially important for
mediating changing coexistence between species across the elevation gradient. Sympatric
species (i.e., lowland-lowland and highland-highland pairs) possessed the greater niche
differences within versus outside of their elevational range, while parapatric species (i.e.,
lowland-highland pairs) displayed the greatest niche differences at the middle elevation site
near to their shared range margins. So far, there have been few a priori predictions made
about how niche differences should change across environmental gradients. Nonetheless,
our finding of greater niche differences between sympatric species within than beyond their
range is consistent with a recent study showing reduced niche differences under drought
conditions due to greater positive effects of drought on the magnitude of interspecific relative
to intraspecific competition [30]. We can only speculate on the biological processes driving the
distinct responses of niche differences without further information (e.g., from functional traits
[31]). However, one possible explanation could be temporal overlap in phenology. Specifically,
the flowering phenology of lowland and highland species overlapped most at the high and
low elevation sites, respectively, while the overlap between lowland and highland species
was smallest at the middle elevation site (Supplementary Fig. 7). Thus, the phenological
overlap between species was associated with the magnitude of niche differences, with
greater phenology overlap leading to smaller niche differences and vice versa [32, 33].

Lowland species were in general competitively superior over highland species, but this
superiority decreased significantly with increasing elevation, and the competitive dominance
of some species pairs was switched in favour of highland plants at the high elevation sites
(i.e., ln(fitness difference) > 0 in Fig. 3e). Nonetheless, in most cases the competitive
advantage of highland species at the high site was not great enough to exclude the lowland
species; possible reasons for this are discussed below. The competitive dominance of
lowland species is consistent with general expectations [6] (but see ref. 20), as well as
empirical evidence showing that lowland species can exert stronger competitive effects on

alpine focal plants than other alpine neighbours [21]. Greater competitive superiority of species

within their home range could result either from their higher intrinsic population growth, in

particular for highland species, or from the fact that species are more resistant to

competition within versus beyond their range [18, 32], though we cannot isolate the relative

contribution of these two processes with the data at hand. Similarly, the increased relative

fitness differences of highland species towards low elevation could result from amplified

differences in intrinsic population growth or competitive responses [30]. Taken together, our

results demonstrate that the range limits of our focal species were influenced by competitors

from both within and outside of their range (i.e., sympatric and parapatric species) and

mediated through both niche differences and relative fitness differences. These results

speak to the need to broadly investigate how environmental conditions impact niche and

relative fitness differences and how they jointly structure communities and species'

distributions across environmental gradients [12].

Although we found significant effects of elevation on the strength of coexistence (i.e., the

coexistence metric), we observed relatively few cases in which coexistence outcomes were

qualitatively altered across the elevation gradient. For example, most lowland species could

still maintain relatively large and positive invasion growth rates, and thus were predicted to

persist, at the high site. The high site is above the current range limit of all lowland species

in this region, and so it is surprising that the climatic and biotic environment appears to be

suitable for them. This might be because while average climate conditions at high elevation

support the growth of our lowland plants, climate extremes do not [34]; if extreme conditions

did not occur during the experiment, then climate limitation would not be observed. Another

possible explanation for lowland species' persistence at high elevation is disequilibrium

between their distributions and current climate. It has been widely reported that the rate of

species' upwards migration across elevation gradients lags behind the pace of climatic

warming [35, 36, 37]. For example, the observed average upward shifts of 183 plant species over

the past 50 years in the European Alps was 30 m [38], while isoclines in the same region have

shifted upwards by 310-355 m [35]. Disequilibrium in the distributions of our lowland species

might be due to their limited dispersal capacities and the often low rate of germination and

establishment of perennial species [36]. Added to this, the growing season temperatures over

the course of our experiment generally exceeded the long-term average

(https://www.meteoswiss.admin.ch/). Thus, the high site could already be within the

potential upper limits of some of the lowland species included in our study. But even if this

were the case, the relationship shown in Fig. 1 implies that lowland species ranges will be

limited by competitors before they are limited by climate.

Manipulative experiments such as ours will tend to underestimate the prevalence of stable

coexistence when they focus on pairwise competition and use homogeneous environmental

conditions over a relatively short temporal scale (e.g., ref. 14). Firstly, such experiments

might miss potentially important stabilizing effects that only emerge from interaction

networks in which any given pair of species is not necessarily able to coexist (e.g., higher-

order interactions [29, 39]). Secondly, standardized, short term experiments might not be able to

capture temporal and spatial variation that could be important for species coexistence (e.g.,

temporal and spatial storage effects [40]). In our experiment, we used the same soil in the

three sites to isolate effects of changing climate from changing soil conditions across the

elevation gradient, which might have reduced opportunities for coexistence driven by

interactions belowground (e.g., plant-soil feedbacks [41]). This failure to capture all possible

coexistence mechanisms might partly explain the lack of stable coexistence between

currently co-occurring species observed in our experiment. For example, we found that only

63% of highland species pairs were predicted to be able to coexist at the high site; small-

scale environmental heterogeneity has been suggested to be especially important for

maintaining the high diversity of alpine plants [42]. Thirdly, due to the short duration the

experiment (four years), our results might be contingent on specific weather conditions over

the course of the experiment, which may also explain the lack of coexistence of highland

species within their range. Lastly, violations of the assumption that monocultures are at

equilibrium abundance could affect our predictions of coexistence. In our experiment, 32%

(11 of 34) of monocultures were predicted to meet this assumption, and those that did not

(10 and 13 monocultures were predicted to be above and below equilibrium, respectively)

were evenly distributed across the sites and so are unlikely to affect the general patterns we

observed (Supplementary Fig. 5).

Our results challenge the conventional view predicting a diminished role for competition in

structuring species' distributions and ecological communities under "harsh" environments at

high elevation or latitude, whilst highlighting the importance of considering species'

interactions to understand and project future species distributions [27, 28]. More generally, our

finding that both lowland and highland species experienced weakened coexistence towards

their range edges (i.e., the upper ranges of lowland species and the lower ranges of

highland species, respectively) suggests that competition might be important for setting both

species' lower and upper elevation limits [10]. An important role for competition at high

elevation could help explain the limited ability of some species to shift their distribution limits

to higher elevation to track climate warming [35, 36]. However, our results suggest that

continuous warming may facilitate the upwards shifts of lowland species and the local

extinction of alpine species [43] by removing the competitive advantage that alpine species

possess under cooler climate. In addition, our results suggest that climatic warming may
destabilize alpine communities by reducing niche differences and amplifying fitness
differences between highland species.

Although modern coexistence theory provides a useful framework for understanding and
predicting species coexistence across environmental gradients, niche differences and
relative fitness differences are only phenomenological syntheses of biological processes
underling species interactions. Future work should seek to uncover the mechanisms through
which competition influences species' distributions along environmental gradients by
manipulating specific abiotic and biotic factors and gathering data on species' physiology,
morphology or life-history strategies (i.e., functional traits [31]). Future studies could also look
at whether competition sets range limits across entire species ranges, including upper and
lower edges of the same species, and how complex interaction networks in multispecies
communities contribute to setting range limits. Such studies would help to provide more
mechanistic insights into the role of species interactions in regulating species' distributions
and bolster our ability to more accurately forecast the impacts of environmental change on
species' distributions and community dynamics.

Materials and Methods

Study area and species

We selected three sites across an elevation gradient in the western Swiss Alps (Bex, Canton de Vaud), situated at 890, 1400 and 1900 m above sea level (hereafter, the low, middle, and high sites; Supplementary Fig. 1). The three sites span a temperature gradient ranging from 2.5 to 9.6 °C (mean annual temperature from 1981 to 2015 [44]; Supplementary Table 1). With increasing elevation, soil moisture increased, and growing season length was shortened by a longer snow-covered period, as measured from July 2019 to June 2020 (Supplementary Fig. 2). All sites were established on south facing and shallow slopes in pasture and fenced to exclude livestock.

We included 14 herbaceous focal species that frequently occur in this region, half of which originated from low elevation (hereafter lowland species) and half from high elevation (highland species, Supplementary Table 2). Lowland species had upper range limits (defined as the 90th percentile of their elevation distribution) below 1500 m (with the exception of *Plantago lanceolata*, with a 90th percentile of 1657 m), while highland species had lower range limits (defined as the 10th percentile of their elevation distribution) above 1500 m, based on a dataset of 550 vegetation plots from the study area [45]. These species consisted of 12 perennial and two biennial species, which are the dominant life histories in this region. Species were selected to include a range of functional types (7 forbs, 4 grasses, 3 legumes) and functional traits (based on plant height, specific leaf area and seed mass). Seeds were obtained from regional suppliers given the large quantities that were needed to establish the experiment (Supplementary Table 2).

Field competition experiment

We designed a field experiment to study the effects of elevation on population growth rates and competitive outcomes by growing focal plants either without competition or competing with a background monoculture of the same or another species (Supplementary Fig. 1). In spring 2017, we established 18 plots (1.6 m x 1 m, 0.2 m deep) at each of the three field sites, lined with wire mesh to exclude rodents (except at the high site) and with weed-suppressing fabric on the sides to prevent roots growing in from outside. To control for soil effects, the beds were then filled with a silt loam soil that originated from a nutrient poor meadow at 1000 m a.s.l. within the study area. Four plots were maintained as bare soil plots (non-competition plots). The other 14 plots received 9 g m^{-2} of viable seeds of each species, which allowed the establishment of a monoculture of relatively high density (competition

plots). We then periodically weeded the plots to maintain monocultures over the course of

the experiment. All species except for two (*Arnica montana* and *Daucus carota*) successfully

established monocultures, of which 11 species (including six lowland species and five

highland species) were fully established by autumn 2017. We then resowed the other plots,

which subsequently established either in spring 2018 (*Poa trivialis* and *Poa alpina* in the low

site and *Bromus erectus* in the middle site) or autumn 2018 (*Aster alpinus, P. trivialis* and *P.*

alpina in the middle site and *Sesleria caerulea* in the low and high sites). Species that failed

to establish were included only as focal species for the calculation of invasion population

growth rates (i.e., the density was low for *A. montana* and *D. carota* in all sites, *Trifolium*

badium in the low site and *S. caerulea* in the middle site, probably due to high mortality rates

caused by drought).

We first raised seedlings of each species in a greenhouse for six weeks on standard

compost and then transplanted them into the field (Supplementary Fig. 1). To test for

responses to elevation in the absence of competition, focal plants were transplanted into

non-competition plots at 25 cm apart in autumn 2017 (n = 9 per species and site). To test for

effects of competition, we transplanted focal individuals into established plots with 14 cm

spacing (n = 9 per focal species, competitor and site). Focal plants that died within two

weeks of transplanting (ca. 5%) were replaced, assuming mortality was caused by transplant

shock. Note that we transplanted focal plants into plots only when the background

monocultures were fully established. In 2018 and 2019, we replaced dead focal individuals

in spring and autumn (ca. 10% each time). The full design included 56 unique interspecific

pairs in each site accounting for 61% of all 14 × 13 = 91 possible pairwise combinations.

These pairs were selected to evenly sample differences in functional trait space based on a

pilot analysis using plant height, specific leaf area and seed mass obtained from the LEDA

dataset [46]. Each focal species competed against four lowland and four highland species,

yielding 14 lowland-lowland and highland-highland pairs and 28 lowland-highland pairs.

Across all three sites, this design resulted in N = 3780 individuals in total ([56 interspecific

pairs × 2 + 14 intraspecific pairs + 14 non-competition] × 9 individuals × 3 sites).

Demographic data

We followed each focal individual between 2017 and 2020 to monitor individual-based

demographic performance (i.e., vital rates; Supplementary Fig. 4). Survival was monitored

twice a year at the beginning and the end of the growing season. Towards the end of the

growing season in each year (August-September), we measured all individuals to record

plant size, whether they flowered, and to estimate seed production on flowering individuals.

To estimate focal plant size, we measured size-related morphological traits on all focal

individuals at each census (i.e., the number and/or length of flowering stalks, leaves or
ramets, depending on species) and estimated dry aboveground biomass using regression
models fitted using collected plant samples (mean R^2 = 0.871; Supplementary Table 3;
Supplementary Methods). To estimate seed production, we counted the number and
measured the size of fruits on reproductive individuals; we then estimated the number of
seeds produced by each individual using regression models fitted using intact fruits of each
species collected at the early fruiting stage on background plants (mean R^2 = 0.806;
Supplementary Table 4; Supplementary Methods). We conducted a separate experiment to
estimate germination and recruitment of each species in each site (Supplementary
Methods).

Population modelling

To estimate population growth rates (λ), we built integral projection models to incorporate
multiple vital rates across the life cycle [47] (see Supplementary Table 5 for model structure
and parameters). Separate IPMs were built to estimate intrinsic growth rates using plants
growing in the absence of competition (in non-competition plots) and invasion growth rates
using plants growing within the background monocultures (in competition plots), under the
assumption that monocultures were at equilibrium (see Supplementary Fig. 5 for a test of
this assumption) and that focal individuals did not interact with each other but only with the
background species. We used plant size (i.e., estimated dry aboveground biomass, log
scale) as a continuous state variable and fitted linear models to estimate vital rate
parameters by combining multiple-year demographic data over three censuses (i.e., 2017-
2018, 2018-2019 and 2019-2020; see Supplementary Methods for consideration of more
complex models). We modeled the probability of survival, flowering, and seedling
establishment using generalized linear models with a binomial distribution, modeled growth
and seed production using general linear models and described the offspring size
distribution using Gaussian probability density functions. We modeled seed germination,
seedling establishment and the seedling size distribution as size-independent functions,
assuming they are unaffected by maternal size (Supplementary Fig. 4; Supplementary Table
5). For each vital rate of each species, we selected the best-fitted vital rate model by
comparing all nested models of the full models using the Akaike information criterion
corrected for small samples (AICc), which allowed us to avoid overfitting models and to
borrow strength across competitor species and sites in cases where full models were
outperformed by reduced models (Supplementary Methods; Supplementary Table 6;
Supplementary Data 1).

We calculated population growth rates (λ) as the dominant eigenvalue of the IPMs, which

represents the discrete per-capita growth rate (i.e., $N_{t+1} = \lambda N_t$) [47]. To evaluate the

uncertainty around λ, we performed parametric bootstraps for size-dependent vital rates

(i.e., survival, growth, flowering, and fecundity). Specifically, we resampled the parameters

of each vital rate model using multivariate normal distributions based on their means and

covariance matrices [48]. We then fitted all IPMs and estimated λ for each of the 500 bootstrap

replicates (Supplementary Data 2).

Estimation of niche differences, relative fitness differences and coexistence outcomes

We quantified niche and relative fitness differences and predicted coexistence outcomes

following the method of Carroll et al. [49]. This method is based on species' sensitivity to

competition defined as the proportional reduction of the growth rate of a focal species *i* when

invading a population of a competitor species *j* that is at its single-species equilibrium, and is

mathematically equivalent to one minus the response ratio:

$$S_{ij} = 1 - \frac{\ln(\lambda_{ij})}{\ln(\lambda_i)}$$

where λ_{ij} denotes the invasion growth rate of focal species *i* and λ_i is its intrinsic growth

rate. The natural logarithm of discrete population growth rates λ estimated from IPMs are

equivalent to per capita growth rate in continuous population growth models [50], and this

transformation makes sensitivities compatible with the coexistence analysis described

below. Sensitivity is > 0 for antagonistic interactions, with higher values equating to stronger

competition, while facilitative interactions lead to negative sensitivities.

For a pair of species, modern coexistence theory predicts that niche differences (ND)

promote coexistence by reducing the intensity of interspecific competition experienced by

both species. Therefore, a pair of species with a large niche difference should display small

mean sensitivities to competition from each other. Consequently, niche differences can be

calculated as one minus the geometric mean of the two sensitivities (i.e., niche overlap). In

contrast, relative fitness differences (RFD) quantify the degree of asymmetry in species'

competitive abilities. Therefore, a pair of species with a large fitness difference should

display large differences in their sensitivities to competition from each other, as quantified as

the geometric standard deviation of sensitivities [49]:

$$ND = 1 - \sqrt{S_{ij}S_{ji}}$$

$$RFD = \sqrt{S_{ji}/S_{ij}}$$

There are three possible outcomes of competition between a given pair of species: stable
coexistence, a priority effect, and competitive exclusion. These can be quantified based on
either invasion criteria or the relative magnitude of niche differences versus relative fitness
differences [15, 51]. Stable coexistence is only possible when both species are able to invade
each other's equilibrium populations; this condition is met when ND > 0 and $RFD < \frac{1}{1-ND}$ [49],
which is equivalent to $\frac{1}{RFD(1-ND)} > 1$, with greater values indicating more stable coexistence
and providing a metric for the strength of coexistence (i.e., coexistence metric [26]). When
neither species can invade when rare, then priority effects occur, meaning that whichever
species is initially established within a community has an advantage and excludes the other.
This could happen when a species pair has a small niche difference and a small relative
fitness difference, that is $ND < 0$ and $RFD < \frac{1}{1-ND}$ [51]. Otherwise, if only one species can
invade, then it is predicted to competitively exclude the inferior competitor; this occurs when
relative fitness differences are big enough to override the stabilizing effect of niche
differences, that is when $RFD > \frac{1}{1-ND}$. We quantified competitive outcomes and coexistence
metrics for each of the 500 bootstrap replicates of the dataset (Supplementary Data 3).

Note that we excluded facilitative interactions that were present in 13% of all pairs because
the equations for niche differences and relative fitness differences are not compatible with
negative values of sensitivity. We quantified the coexistence determinants of species pairs in
cases where either one or both of the species were predicted to be unable to persist in the
absence of neighbours (i.e., ln($\lambda_{intrinsic}$) < 0; 8% of all pairs). This was because the
quantification of niche and fitness differences were based on the competitive impacts on
population growth (i.e., sensitivity to competition) and independent of the actual estimates of
population growth rates. A further analyses after excluding these pairs produced very similar
results that did not change our conclusions (data not shown).

Statistical analysis

We tested the effect of elevation on species' intrinsic and invasion growth rates, and whether
these effects differed between lowland and highland species, by fitting mixed effects models
with elevation (continuous), species origin (i.e., lowland or highland) and their interaction as
fixed effects and focal species identity as a random effect. To test whether coexistence was
strengthened at high versus low elevation, we fitted a mixed-effects model of the

coexistence metric as explained by elevation and competitor identity (i.e., the three types of pairs: lowland-lowland, lowland-highland and highland-highland pairs), with species-pair as a random effect. We fitted similar models for niche and relative fitness differences (absolute values) to gain insights into whether changes in coexistence were driven predominantly by variation in niche or relative fitness differences. Additionally, we tested the effect of elevation on relative fitness differences for each type of pairs separately and used the actual values of lowland-highland pairs since we were interested not only in the magnitude of relative fitness differences but also the direction of effects (i.e., whether lowland or highland species were dominant competitors). The response variables in these models were log-transformed to meet model assumptions. Note that we therefore had to use niche overlap (1-ND) for the log-transformation instead of niche differences, because niche differences included negative values. Firstly, we performed these analyses based on the mean of the 500 bootstrap replicates for a given focal species/background competitor combination and used likelihood ratio tests to determine the significance of fixed effects. To account for the uncertainty around estimated λ, we also performed individual tests for each bootstrap replicate of the dataset and determined the significance of fixed terms based on whether the 95% confidence interval of estimated coefficients, or contrasts of interest, included zero. All population modelling and statistical analysis were conducted in R version 4.0.3 [52].

Data availability

A complete comparison of vital rate models can be found in Supplementary Data 1. Estimated population growth rates of all integral projection models can be found in Supplementary Data 2. Estimates of competitive outcomes and coexistence coefficients can be found in Supplementary Data 3. Supplementary Data 2&3 were used to produce Figs. 1-3. These data are available on Figshare: https://figshare.com/s/cb9e034633f47739466b

Code availability

All R scripts used for modelling and statistical analyses are available on GitHub: https://github.com/ShengmanLyu/competition-contributes-to-range-limits

Acknowledgements

We thank Loïc Liberati, Tim Murray, Jessica Joaquim, Cuicui Zhou, Yan Hess, Kai-Hsiu Chen, Megan Stamp and other members of the Plant Ecology group at ETH Zürich for help with field work. We thank the Commune de Bex and Jean-Louis Putallaz for access to the field sites. We also thank Antoine Guisan and the ECOSPAT group at the University of Lausanne for providing plant community and climatic data. We thank Jonathan Levine and

the Plant Ecology group for their feedback on an earlier version of the manuscript. S.L.

acknowledges the Chinese Scholarship Council for financial support (No.201706100184).

J.M.A. received funding from the European Union's Horizon 2020 research and innovation

programme under grant agreement No. 678841.

Author contributions

J.M.A. designed the experiment. S.L. and J.M.A. collected data. S.L. conducted the

modelling and data analysis. S.L. wrote the manuscript with input from J.M.A.

Competing interests

The authors declare no competing interests.

References

532   1.   Darwin C. *On the Origin of species*. London: British Museum Natural History
533   Cambridge: Cambridge University Press (1859).
534

535   2.   MacArthur RH. *Geographical ecology : patterns in the distribution of species*. Harper & Row
536   (1972).
537

538   3.   Wisz MS*, et al.* The role of biotic interactions in shaping distributions and realised
539   assemblages of species: implications for species distribution modelling. *Biological Reviews*
540   **88**, 15-30 (2013).
541

542   4.   Holt RD. Bringing the Hutchinsonian niche into the 21st century: Ecological and evolutionary
543   perspectives. *Proceedings of the National Academy of Sciences* **106**, 19659-19665 (2009).
544

545   5.   Hargreaves AL, Samis KE, Eckert CG, Associate Editor: Oswald JS, Editor: Judith LB. Are
546   Species Range Limits Simply Niche Limits Writ Large? A Review of Transplant Experiments
547   beyond the Range. *The American Naturalist* **183**, 157-173 (2014).
548

549   6.   Louthan AM, Doak DF, Angert AL. Where and When do Species Interactions Set Range
550   Limits? *Trends in Ecology & Evolution* **30**, 780-792 (2015).
551

552   7.   Paquette A, Hargreaves AL. Biotic interactions are more often important at species' warm
553   versus cool range edges. *Ecology Letters* **n/a**, (2021).
554

555   8.   Liang E*, et al.* Species interactions slow warming-induced upward shifts of treelines on the
556   Tibetan Plateau. *Proceedings of the National Academy of Sciences* **113**, 4380-4385 (2016).
557

558   9.   Chesson P, Huntly N. The Roles of Harsh and Fluctuating Conditions in the Dynamics of
559   Ecological Communities. *The American Naturalist* **150**, 519-553 (1997).
560

561   10.   Chan S-F, Shih W-K, Chang A-Y, Shen S-F, Chen I-C. Contrasting forms of competition set
562   elevational range limits of species. *Ecology Letters* **22**, 1668-1679 (2019).
563

564   11.   Godsoe W, Jankowski J, Holt RD, Gravel D. Integrating Biogeography with Contemporary
565   Niche Theory. *Trends in Ecology & Evolution* **32**, 488-499 (2017).
566

567   12.   Alexander JM, Diez JM, Usinowicz J, Hart SP. Species' Distributions as a Coexistence
568   Problem: A Response to Godsoe et al. *Trends in Ecology & Evolution* **33**, 144-145 (2018).
569

570   13.   Barabás G, D'Andrea R, Stump SM. Chesson's coexistence theory. *Ecological Monographs*
571   **88**, 277-303 (2018).
572

573   14.   Levine JM, HilleRisLambers J. The importance of niches for the maintenance of species
574   diversity. *Nature* **461**, 254-257 (2009).
575

576   15.   Chesson P. Mechanisms of Maintenance of Species Diversity. *Annual Review of Ecology and*
577   *Systematics* **31**, 343-366 (2000).
578

579   16.   Diamond JM. Distributional Ecology of New Guinea Birds: Recent ecological and
580   biogeographical theories can be tested on the bird communities of New Guinea. *Science* **179**,
581   759-769 (1973).
582

583   17.   Harpole WS, Tilman D. Grassland species loss resulting from reduced niche dimension.
584   *Nature* **446**, 791-793 (2007).
585

586   18.   Hart SP, Freckleton RP, Levine JM. How to quantify competitive ability. *Journal of Ecology*
587   **106**, 1902-1909 (2018).
588

589   19.   Vittoz P, Randin C, Dutoit A, Bonnet F, Hegg O. Low impact of climate change on subalpine
590   grasslands in the Swiss Northern Alps. *Global Change Biology* **15**, 209-220 (2009).
591

592   20.   Freeman BG. Lower elevation animal species do not tend to be better competitors than their
593   higher elevation relatives. *Global Ecology and Biogeography* **29**, 171-181 (2020).
594

595   21.   Alexander JM, Diez JM, Levine JM. Novel competitors shape species' responses to climate
596   change. *Nature* **525**, 515-518 (2015).
597

598   22.   Lax S, Abreu CI, Gore J. Higher temperatures generically favour slower-growing bacterial
599   species in multispecies communities. *Nature Ecology & Evolution* **4**, 560-567 (2020).
600

601   23.   Davis AJ, Jenkinson LS, Lawton JH, Shorrocks B, Wood S. Making mistakes when predicting
602   shifts in species range in response to global warming. *Nature* **391**, 783-786 (1998).
603

604   24.   Bertness MD, Callaway R. Positive interactions in communities. *Trends in Ecology &*
605   *Evolution* **9**, 191-193 (1994).
606

607   25.   He Q, Bertness MD, Altieri AH. Global shifts towards positive species interactions with
608   increasing environmental stress. *Ecol Lett* **16**, 695-706 (2013).
609

610   26.   Germain RM, Mayfield MM, Gilbert B. The "filtering" metaphor revisited: competition and
611   environment jointly structure invasibility and coexistence. *Biology Letters* **14**, 20180460
612   (2018).
613

614   27.   Violle C, Pu Z, Jiang L. Experimental demonstration of the importance of competition under
615   disturbance. *Proceedings of the National Academy of Sciences* **107**, 12925-12929 (2010).
616

617   28.   Hart SP, Marshall DJ. Environmental stress, facilitation, competition, and coexistence.
618   *Ecology* **94**, 2719-2731 (2013).
619

620   29.   Levine JM, Bascompte J, Adler PB, Allesina S. Beyond pairwise mechanisms of species
621   coexistence in complex communities. *Nature* **546**, 56 (2017).

30. Matías L, Godoy O, Gómez-Aparicio L, Pérez-Ramos IM. An experimental extreme drought reduces the likelihood of species to coexist despite increasing intransitivity in competitive networks. *Journal of Ecology* **106**, 826-837 (2018).

31. Perez-Ramos IM, Matias L, Gomez Aparicio L, Godoy O. Functional traits and phenotypic plasticity modulate species coexistence across contrasting environments. *bioRxiv*, 539619 (2019).

32. Godoy O, Levine JM. Phenology effects on invasion success: insights from coupling field experiments to coexistence theory. *Ecology* **95**, 726-736 (2014).

33. Alexander JM, Levine JM. Earlier phenology of a nonnative plant increases impacts on native competitors. *Proceedings of the National Academy of Sciences* **116**, 6199-6204 (2019).

34. Zimmermann NE*, et al.* Climatic extremes improve predictions of spatial patterns of tree species. *Proceedings of the National Academy of Sciences* **106**, 19723-19728 (2009).

35. Vitasse Y*, et al.* Phenological and elevational shifts of plants, animals and fungi under climate change in the European Alps. *Biological Reviews* **n/a**, (2021).

36. Alexander JM*, et al.* Lags in the response of mountain plant communities to climate change. *Global Change Biology* **24**, 563-579 (2018).

37. Rumpf SB*, et al.* Extinction debts and colonization credits of non-forest plants in the European Alps. *Nature Communications* **10**, 4293 (2019).

38. Rumpf SB*, et al.* Range dynamics of mountain plants decrease with elevation. *Proceedings of the National Academy of Sciences* **115**, 1848-1853 (2018).

39. Angulo MT, Kelley A, Montejano L, Song C, Saavedra S. Coexistence holes characterize the assembly and disassembly of multispecies systems. *Nature Ecology & Evolution*, (2021).

40. Cáceres CE. Temporal variation, dormancy, and coexistence: A field test of the storage effect. *Proceedings of the National Academy of Sciences* **94**, 9171-9175 (1997).

41. Ke P-J, Wan J. Effects of soil microbes on plant competition: a perspective from modern coexistence theory. *Ecological Monographs* **n/a**, e01391 (2019).

42. Körner C. *Alpine plant life : functional plant ecology of high mountain ecosystems*, Softcover reprint of the hardcover 2nd ed. 2003 edn. Springer (2003).

43. Nomoto HA, Alexander JM. Drivers of local extinction risk in alpine plants under warming climate. *Ecology Letters* **24**, 1157-1166 (2021).

667   44.   Scherrer D, Vitasse Y, Guisan A, Wohlgemuth T, Lischke H. Competition and demography
668   rather than dispersal limitation slow down upward shifts of trees' upper elevation limits in the
669   Alps. *Journal of Ecology* **108**, 2416-2430 (2020).
670

671   45.   Randin CF*, et al.* Climate change and plant distribution: local models predict high-elevation
672   persistence. *Global Change Biology* **15**, 1557-1569 (2009).
673

674   46.   Kleyer M*, et al.* The LEDA Traitbase: a database of life-history traits of the Northwest
675   European flora. *Journal of Ecology* **96**, 1266-1274 (2008).
676

677   47.   Ellner SP, Childs DZ, Rees M. *Data-Driven Modelling of Structured Populations : A Practical
678   Guide to the Integral Projection Model*. Cham : Springer (2016).
679

680   48.   Reed PB*, et al.* Climate manipulations differentially affect plant population dynamics within
681   versus beyond northern range limits. *Journal of Ecology* **109**, 664-675 (2021).
682

683   49.   Carroll IT, Cardinale BJ, Nisbet RM. Niche and fitness differences relate the maintenance of
684   diversity to ecosystem function. **92**, 1157-1165 (2011).
685

686   50.   Chesson P. Multispecies Competition in Variable Environments. *Theoretical Population
687   Biology* **45**, 227-276 (1994).
688

689   51.   Grainger TN, Letten AD, Gilbert B, Fukami T. Applying modern coexistence theory to priority
690   effects. *Proceedings of the National Academy of Sciences* **116**, 6205-6210 (2019).
691

692   52.   R Core Team. R: A language and environment for statistical computing.). R Foundation for
693   Statistical Computing (2020).
694
695

Figure 1. Population growth rates across the elevation gradient. Effects of elevation on
intrinsic (**a**, in the absence of neighbours) and invasion (**b**, in the presence of neighbours)
population growth rate of lowland (orange) and highland (blue) focal species. Populations
are predicted to persist when $ln(\lambda) > 0$ (dashed horizontal line). The points show average (±
SE) population growth rates across species. Solid lines show significant relationships (test
based on the mean of 500 bootstrap replicates; $P < 0.05$). Orange and blue shading
indicates the 95% quantiles of fitted relationships across 500 bootstrap replicates of the
dataset.

Figure 2. Competition outcomes across the elevation gradient. The estimated competitive outcomes (**a-c**) and strength of coexistence (i.e., coexistence metric, log-transformed; **d-f**) of lowland-lowland (orange), lowland-highland (green), and highland-highland (blue) species pairs in the three sites. In **a-c**, grey bars represent stable coexistence while white bars represent competitive exclusion and priority effects, and error bars indicate mean and 95% confidence intervals based on 500 bootstrap replicates. In **d-f**, solid lines indicate a significant interaction of elevation x competitor identity (test based on the mean of 500 bootstraps; $P < 0.05$), and the points are averages (± SE) across species pairs. The shading indicates the 95% quantiles of fitted relationships across 500 bootstrap replicates of the dataset.

Figure 3. Niche differences and relative fitness differences across the elevation

gradient. Estimates of the niche differences (expressed as niche overlap, i.e. 1 – niche

differences; see Methods, **a-c**) and relative fitness differences (**d-f**) of lowland-lowland

(orange), lowland-highland (green) and highland-highland (blue) species pairs. Solid lines

indicate the significant interactions of elevation x competitor identity for niche differences

and significant effects of elevation on relative fitness differences for each type of pair

separately (test based on the mean of 500 bootstrap replicates; $P < 0.05$) while dotted lines

indicate non-significant relationships ($P > 0.05$). The points are averages (± SE) across

species pairs. The shading indicates the 95% quantiles of fitted relationships across 500

bootstrap replicates of the dataset. In **d-f**, absolute values of fitness differences (log-

transformed) are shown for lowland-lowland and highland-highland pairs, while actual values

are shown for lowland-highland pairs, with negative values indicating lowland species are

dominant competitors.